1
2 Transcriptional regulation of the synthesis and secretion of farnesol in the fungus *Candida albicans:*
3 Examination of the Homann transcription regulator knockout collection

11 Daniel J. Gutzmann*, Jaxon J. Kramer, Brigid M. Toomey, Cory H. T. Boone, Audrey L. Atkin* & Kenneth
12 W. Nickerson*
18 School of Biological Sciences
19 University of Nebraska
20 Lincoln, NE USA
27 *For correspondence. E-mail : knickerson1@unl.edu, aatkin@unl.edu
28 daniel.gutzmann@huskers.unl.edu
29 Tel (402) 472-2253; Fax (402) 472-8722.

Keywords

• *Candida albicans*

• Farnesol

• Transcription factor mutant

Abstract

Candida albicans is an efficient colonizer of human gastrointestinal tracts and skin and is an

opportunistic pathogen. *C. albicans* exhibits morphological plasticity and the ability to switch between

yeast and filamentous morphologies is associated with virulence. One regulator of this switch is the

quorum sensing molecule farnesol which is produced by *C. albicans* throughout growth. However, the

synthesis, secretion, regulation, and turnover of farnesol is not fully understood. To address this, we

used our improved farnesol assay to screen a transcription regulator knockout library of 164 mutants for

farnesol accumulation in whole cultures, pellets, and supernatants. All mutants produced farnesol and

they averaged 9.2X more farnesol in the pellet than the supernatant. Nineteen mutants had significant

differences with ten mutants producing more farnesol than their SN152$^+$ parent while nine produced

less. Seven mutants exhibited greater secretion of farnesol while two exhibited less. We examined the

time course for farnesol accumulation in six mutants with the greatest accumulation differences and

found that those differences persisted throughout growth and they were not time dependent.

Significantly, two high accumulating mutants did not exhibit the decay in farnesol levels during

stationary phase characteristic of wild type *C. albicans*, suggesting that a farnesol

modification/degradation mechanism is absent in these mutants. Identifying these transcriptional

regulators provides new insight into farnesol's physiological functions regarding cell cycle progression,

white-opaque switching, yeast-mycelial dimorphism, and response to cellular stress.

Introduction

Candida albicans is an opportunistic pathogen that is present in most human gastrointestinal tracts and

is an efficient colonizer of mucosal surfaces (Neville et al. 2015). A weakened immune system (HIV,

chemotherapy, organ transplantation) or loss of competing flora (antibiotic treatment) allows for *C.*

albicans to colonize and invade host tissues leading to candidiasis (Pfaller and Diekema 2007). Virulence

of *C. albicans* is strongly linked to its ability to switch between yeast and filamentous morphologies

(Romano 1966; Kadosh 2019) as mutants that have lost the ability to switch are avirulent (Lo et al. 1997;

Saville et al. 2003). Dimorphism is also important for adaptation to different host environments (Noble

et al. 2017; Alves et al. 2020) and evading host immune responses (Gow et al. 2012). One regulator of

dimorphic switching is the quorum sensing molecule (QSM) farnesol (Hornby et al. 2001). Farnesol is

produced by *C. albicans* cells continuously throughout growth (Boone et al. 2022), except in

anaerobically grown cells (Dumitru et al. 2004) and opaque cells (Dumitru et al. 2007) where farnesol

production is much lower. However, despite numerous publications on farnesol's role as a QSM and

virulence factor, the actual mechanisms regulating its synthesis and secretion remain unclear.

As an approach to identifying these genes and mechanisms, we tested the hypothesis that

farnesol synthesis and secretion are regulated by the transcriptional networks involved in

morphogenesis and virulence. The transcription regulator knockout library generated by Homann and

colleagues has 165 different transcription regulators deleted, in most cases with two independently

derived knockouts (Homann et al. 2009). Thus, we grew each of the library strains in liquid culture and

then assayed them for total, extracellular, and intracellular farnesol by the improved methods of Boone

et al. (2022) . Key features of this assay include prevention of analyte loss by avoiding filtration and

minimizing evaporation, while incorporating simultaneous cell lysis and analyte extraction by ethyl

acetate. The assay enables comparison of whole culture values with the sum of their cell pellets and

supernatants. Our results suggest farnesol accumulation is integrated with transcriptional regulators of

cell cycle progression, white-opaque switching, yeast-mycelial dimorphism, and responses to cell stress.

Our results provide new insights into the physiological role of farnesol and understanding of the

regulatory mechanisms underlying its production.

Materials and Methods

Strains and media

The homozygous transcription regulator deletion mutants were obtained from the transcription

regulator knockout library provided by Dr. Alexander Johnson's lab (Homann et al. 2009) available

through the Fungal Genomics Stock Center (Manhattan, Kansas, USA; McCluskey et al. 2010). The strains

were maintained at 30°C on plates with YPD (1% yeast extract, 2% peptone, 2% glucose). The

transcription regulator knockout mutants were constructed from *C. albicans* SN152, an auxotroph for

arginine, histidine, and leucine. Each of the transcription regulator (TR) mutants is auxotrophic for

arginine. A wild-type control strain was created by reintroduction of a single allele of HIS1 and LEU2 into

the parent strain (Homann et al. 2009). Throughout this paper, we will refer to this control strain simply

as SN152$^+$. All assays with these mutants include the paired wild-type strain located in the H12 well of

the X2 and Y2 independently derived deletion collections.

Screening the transcription regulator deletion mutants for differences in accumulation farnesol

secretion

The transcription regulator mutant collection was transferred by a 48-pin replicator to YPD

plates and grown at 30°C for 48 hours. Mutants were then transferred to 3 mL YPD liquid medium and

grown for 16 hours at 30°C with rotary agitation at 250 RPM. This culture was used as the inoculum

(1:100) for 75 mL YPD in 250 mL flasks which were incubated 24 hours at 30°C, 250 RPM. Fifty mL of this

culture was used for dry weight determination and 20 mL (2 X 10 mL) was used for determination of

farnesol production as described by Boone et al. (2022). In this method, 10 mL was used to measure

whole culture values while the other was centrifuged to obtain the pellet and supernatant values. Due

to the large number of mutants in the collection, the mutants were assayed in batches of 10-20 mutants

with the SN152$^+$ control assayed in triplicate as part of each batch. Mutants with farnesol accumulation

of greater than 1 log$_2$ fold above or below the mean value for SN152$^+$ were chosen for follow-up

analysis. In follow-up analyses, E,E-farnesol accumulation was measured in at least three independent

experiments using both the X and Y independently-derived TR deletion mutants (n=6).

Temporal dynamics of farnesol accumulation

To examine the time course of farnesol production, we assayed the farnesol accumulation of the

four highest and two lowest accumulating mutants at several cell densities throughout growth. Mutants

were inoculated into to 6 mL YPD liquid medium and grown for 16 hours at 30°C with rotary agitation at

250 RPM. This culture was used as the inoculum (1:100) for replicate flasks for each time point with 75

mL YPD per mutant strain. Farnesol measurements were taken 12, 18, 24, 36, 48, and 80 hours post

inoculation as describe by Boone et al. 2022. Two independent time courses were performed for the

mutant strains and three independent time courses were performed for the SN152$^+$ control. Farnesol

accumulation values were normalized to the 50 mL dry weight at the indicated time point. To quantify

differences in farnesol accumulation across time, area under the curve (AUC) analyses were performed

in GraphPad Prism.

Statistical analysis

Statistical analyses were performed using Microsoft Excel (Version 16.61, Microsoft Office, Las

Vegas, NV, USA) and GraphPad Prism Software (Version 9.5.0, San Diego, CA, USA). All biological data

are represented as mean $\pm$ SD of at least 3 biological replicates for both the X and Y independent

mutants unless otherwise stated. Mutants with significant differences between the X and Y mutant were

excluded and considered a false positive. Normality and homogeneity of variance were assessed by the

D' Agostino-Pearson omnibus (K2) test and Brown-Forsythe test, respectively. Differences between

groups that were normally distributed and homoscedastic (equal variances) were assessed by One-way

ANOVA with Dunnett's multiple comparisons test. Differences between groups that were not normally

distributed or heteroscedastic were assessed by the Kruskal-Wallis test with Dunn's multiple

comparisons tests. Both analyses were performed in GraphPad Prism. Differences were considered

significant at $p < 0.05$ (*$p < 0.05$, **$p < 0.01$, ***$p < 0.001$).

Results

General design and reproducibility of the screen for farnesol regulators

Homann et al. (2009) screened their collection of 165 independent transcription regulator mutants

for possible differences in 55 growth conditions. It was a monumental piece of work. However, for

practical reasons they limited themselves to phenotypes discernible via colonies on agar plates. We

have now extended the list of phenotypes to include accumulation of both intracellular and extracellular

farnesol as determined by our improved GC assay (Boone et al, 2022). For the initial screen, 164 mutants

were screened in 12 batches. One mutant $\Delta\Delta hfl1$ [orf19.3063] was not screened as it exhibited a severe

growth defect on YPD. Each batch included duplicate or triplicate SN152[+] controls, for a total of 198

cultures and 594 GC-FID runs. Farnesol accumulation values were normalized on a per cell basis using

the 50 mL dry weight value of the culture assayed. In this initial screen, the paired WT controls were run

a total of 34 times and their whole culture mean E,E-farnesol accumulation value was 5.29 ± 1.12 ng/μL/

50 mL dry weight (8.27 ± 2.8 μM).

For the rescreen, 27 strains were selected because their farnesol values differed from the

average values for SN152[+] by a $\log_2$ fold change of > 1.0. To these we added 3 strains ($\Delta\Delta upc2, \Delta\Delta tac1,$

and $\Delta\Delta csr1$) because prior reports from the literature indicated they were likely to have altered farnesol

accumulations. These 30 strains were then rescreened five more times, two from the X plate and three

from the Y plate (total n =6) in 10 batches including the SN152[+] parent in biological duplicate or

triplicate, requiring 180 flasks and 540 GC measurements. Thus the rescreen data set examines 30

transcriptional regulator mutants (n=6), reporting the means ± SD for normalized farnesol accumulation

as detected in whole cultures, pellets, and supernatants as well as the supernatant/pellet ratio.

Composite data sets for the initial screen (File S1), rescreen (File S2) and farnesol growth curves (File S3)

are available at Figshare. We now describe several highlights from these data sets.

Mutants altered in farnesol accumulation

Our initial screen of 164 strains, all from the X plate, was done with an n = 1 comparing whole

culture values versus the sum of the cell pellet and supernatant values. The 164 TR mutants were

assayed in 12 batches which always included a biological duplicate or triplicate of the SN152[+] parent.

The whole culture accumulation and supernatant/pellet ratios are shown in Fig. 1. The distribution of

the 34 values for SN152[+] (red) is shown in Fig. 1 where they can be seen in comparison with the

distribution of the 164 TR mutants (black). As expected (Boone et al. 2022), the whole culture (W) and

pellet (P) + supernatant (S) values were always very close to one another, with the average WPS relative

error values (W-(P+S)/W) being 0.07 ± 0.06 for E,E-farnesol.

175   The vast majority of the 164 TR mutants were unchanged from the SN152[+] replicates (Fig. 1).

We then identified all mutants which were > 1 $\log_2$ above or below the mean values, as indicated by the

4 dotted lines in Fig. 1, and designated them as mutants of interest to be rescreened with greater

accuracy. There were 27 mutants with higher or lower farnesol accumulation. Six mutants (*ΔΔcwt1,*

ΔΔrim101, ΔΔcrz2, ΔΔzcf1,ΔΔtea1, ΔΔrme1) were designated as false positives because their rescreened

values from the X and Y plates differed significantly or their rescreen values differed significantly from

the initial screen. Thus, we identified ten mutants with elevated farnesol (Fig. 2A), nine with reduced

farnesol (Fig. 2B), seven with elevated supernatant/pellet ratios (Fig. 2C), and two with lowered farnesol

supernatant/pellet ratios (Fig. 2C). Each of these mutants is described in greater detail in Table 1.

Of the 19 mutants with statistically significant differences in farnesol accumulation, only *ΔΔnrg1*

had been identified in prior literature (Kebaara et al. 2008). Ten mutants (*ΔΔswi4, ΔΔahr1, ΔΔnrg1,*

ΔΔrap1, ΔΔpho4, ΔΔtye7, ΔΔaaf1, ΔΔash1, ΔΔrtg1, ΔΔssn6) had significantly increased farnesol

accumulation either in the whole culture or supernatant fraction compared to the SN152[+] parent strain

(Table 1). The two highest accumulators (*ΔΔswi4* and *ΔΔahr1)* have known roles in cell cycle progression

(Hussein et al. 2011; Sellam et al. 2019). Additionally, two farnesol over accumulating mutants (*ΔΔtye7*

and *ΔΔrtg1*) have roles in carbohydrate metabolism (Askew et al. 2011; Moreno-Velásquez et al. 2020),

four (*ΔΔahr1, ΔΔnrg1, ΔΔash1, ΔΔssn6*) have a role in dimorphism (Fu et al. 1998; Murad et al. 2001;

Inglis and Johnson 2002; García-Sánchez et al. 2005), and two (*ΔΔahr1, ΔΔssn6)* have roles in

white/opaque switching (Zordan et al. 2007; Wang et al. 2011; Hernday et al. 2016). *ΔΔwor2* appeared

to have a modest increase in accumulation but this did not reach the significance threshold (Table 1, Fig.

2A). We note that another regulator of white/opaque switch, Czf1, has role in coordinating switching

with the response to farnesol (Langford et al. 2013).

Nine mutants (*ΔΔorf19.6874, ΔΔmsn4, ΔΔcph2, ΔΔfcr1, ΔΔsko1, ΔΔgzf3, ΔΔcas5, ΔΔcup9,*

ΔΔtec1) had significantly decreased farnesol accumulation either in the whole culture or supernatant

fraction compared to the SN152[+] parent strain (Table 1). Three of these low farnesol accumulating

mutants (*ΔΔmsn4, ΔΔsko1, ΔΔcas5*) have known roles in stress response to cell wall damage and

oxidative stress(Nicholls et al. 2004; Bruno et al. 2006; Rauceo et al. 2008; Xie et al. 2017) and one

mutant (*ΔΔcup9*) has a known role in farnesol response by mediating Nrg1 degradation in the absence

of farnesol (Lu et al. 2014).

The three mutants chosen because of literature precedent had elevated farnesol accumulation

but did not reach statistical significance (*ΔΔcsr1, ΔΔupc2, ΔΔtac1*) (Table 1). Ganguly et al. (2011)

reported decreased accumulation of farnesol by the *ΔΔcsr1* mutant. Notably, the assay conditions used

in our screen (YPD, 30°C) differ significantly from those used in prior studies of *ΔΔcsr1* which employed

biofilm forming conditions at 37°C in SPIDER medium (Ganguly et al. 2011). Thus, the dynamics of

farnesol accumulation are likely influenced not only by temperature, but also by growth medium (Boone

et al. 2022).

Mutants altered in the farnesol supernatant/pellet ratio

The supernatant/pellet (S/P) ratios for farnesol were determined for each mutant (n=1) in

comparison to those for their SN152$^+$ control (Fig. 1). The value for SN152$^+$ (n= 22 batches) was 0.12 $\pm$

0.02 for farnesol. As we had observed previously (Boone et al. 2022), farnesol was predominantly

retained in the cell pellet. The S/P ratios were all tightly clustered from 0.07-0.14 except for four

mutants (*ΔΔmsn4,ΔΔfcr1, ΔΔgzf3, and ΔΔorf19.6874*) which exhibited less secretion (0.05-0.08) and

seven (*ΔΔswi4, ΔΔpho4, ΔΔcas5, ΔΔhap5, ΔΔssn6, ΔΔtye7, and ΔΔrap1*) which exhibited greater

secretion (0.19-0.30) (Fig. 2C). Inexplicably, *ΔΔzcf22* was not among the rescreened mutants; it should

likely be included among the mutants which exhibited less secretion.

Two mutants, *ΔΔhap5* and *ΔΔcas5* are of special interest in that they exhibit greater secretion

without being accompanied by greater total farnesol production (Fig. 1). Thus, they target secretion

specifically. Mutants such as *ΔΔswi4, ΔΔahr1, and ΔΔrap1* with much greater farnesol production (Fig.

2A, Table 1) could exhibit a higher supernatant to pellet ratio because internal and external farnesol are

in equilibrium or there is a need to keep internal farnesol levels below a critical threshold value. In both

cases, the greater secretion would be an indirect consequence of greater production.

Temporal dynamics of farnesol accumulation

The initial screen and rescreen for farnesol accumulation were done at a single time point 24

hours post inoculation. To verify that farnesol differences we observed were not time dependent, we

extended the assay to 12, 18, 24, 36, 48, and 80 hours post inoculation for four over accumulating

(*ΔΔswi4, ΔΔahr1, ΔΔrap1, ΔΔnrg1*) and two under accumulating (*ΔΔmsn4, ΔΔ19.6874*) mutants (Figs. 3

and 4, respectively), with n=2 for each mutant and n=3 for the SN152+ control. Growth of the six mutant

strains paralleled the growth of SN152[+] (Figs. 3A, 4A). As expected, the four over accumulating mutants

had elevated farnesol at all time points in both the whole culture (Fig. 3B) and supernatant fractions

(Fig. 3D) while the two under accumulating mutants had decreased farnesol in both the whole culture

(Fig. 4B) and supernatant fractions (Fig. 4D). To compare total farnesol accumulations, area under the

curve analyses (AUC) were performed. The four over accumulating mutants *ΔΔswi4, ΔΔrap1, ΔΔahr1,*

and *ΔΔnrg1* had significantly higher AUC in the whole culture (Fig. 3C) and supernatant (Fig. 3E) than the

SN152[+] parent while *ΔΔorf19.6874* and *ΔΔmsn4* had significantly lower supernatant AUC (Fig. 4E). Of

great interest, two of the over accumulating mutants (*ΔΔswi4, ΔΔrap1)* still had significant farnesol

remaining at 80 hours (Figs. 3B, 3D). They did not exhibit the decay in farnesol levels during stationary

phase characteristic of wild type *C. albicans* (Boone et al. 2022) as well as SN152[+] and the other two over

accumulating mutants tested (Fig. 3B), suggesting that a farnesol modification or degradation

mechanism is absent from these two mutants.

Opaque phenotype of *ΔΔtup1*

One surprising result of the screen (Fig. 1) was that *ΔΔtup1* did not appear to over accumulate

farnesol in these assay conditions (Table 1). This result was surprising because we had previously

reported that the *ΔΔtup1* mutant produced 17 times more farnesol than its parent (Kebaara et al. 2008).

A possible explanation for this discrepancy is that *C. albicans* can undergo phenotypic switching

between two heritable states: white and opaque, each of which is normally stable for thousands of cell

divisions. Switching between the two cell types is reversible and occurs without any chromosomal

rearrangements or sequence changes. Critically, Tup1 is a key repressor of the opaque state and

Alkafeef et al. (2018) have shown that loss of *TUP1* is sufficient to induce the opaque phase, even in a

MTL a/α background (Alkafeef et al. 2018). The pressure of *ΔΔtup1* to spontaneously convert to the

opaque phase is strong (Alkafeef et al. 2018) and we already know that wild type opaque cells usually

produce far less farnesol than do white cells (Dumitru et al. 2007). Given this, we investigated whether

the *ΔΔtup1* mutant in the Homann collection is in the opaque phase under the assay conditions

performed in the screen (YPD, 30°*C*)*.* It is important to note that *ΔΔtup1* cells are locked in the

filamentous morphology (Kebaara et al. 2008), making visual determination between white and opaque

cells more difficult. When plated on YPD medium supplemented with phloxine B and incubated at 30°C,

the *ΔΔtup1* colonies were pink while the SN152[+] colonies plated under the same conditions were white

(Fig. 5A). The pink colonies for *ΔΔtup1* are consistant with the cells being in the opaque phase. This view

was supported by microscopic examination of the *ΔΔtup1* cells which showed numerous short but

elongated cells consistent with the opaque state (Fig. 5B). The typical budding yeast cells of SN152[+] are

included for comparison (Fig. 5B). We are currently investigating the conditions that influence the

farnesol production by white and opaque *ΔΔtup1* cells.

Discussion

We examined 164 independent transcriptional regulator knockout mutants for their

accumulation and secretion of farnesol. This project was undertaken to define the genes needed for

farnesol's synthesis, secretion, and regulation. For logistical reasons, the initial screen and rescreen

cultures were grown at 30°C in liquid YPD medium and sampled after 24 hours when the cells were in

early stationary phase. The 24 hour time point was chosen based on our prior measurements with wild

type *C. albicans* SC5314 over 72 hours of culture which showed that farnesol accumulation generally

followed increased cell mass, peaked in early stationary phase, and then declined sharply after ~ 30

hours (Boone et al. 2022). Thus, the 24 hour time point was intended to cover the peak of farnesol

accumulation. Similarly, our choice of YPD as a growth medium was to insure that all of the TR mutants

grew well by 24hrs and indeed 164 out of 165 had done so.

With the advantage of now having reliable measures for farnesol presence in whole cultures,

cell pellets, and supernatants, we can approach the genetic basis for their synthesis, secretion,

regulation, and turnover. As expected, most of the TR mutants were unchanged for these parameters

(Fig. 1) while a few potentially useful mutants gave significantly higher and lower levels of farnesol.

Significantly, no mutants were entirely devoid of farnesol; its levels often varied but they never dropped

to zero. A tentative conclusion from this observation is that farnesol has an essential physiological

function over and above acting as a quorum sensing molecule and virulence factor or that it is the

product of a redundant biosynthetic pathway. In examing mutants with significant differences in

farnesol production, several patterns in the transcriptional networks emerge that may highlight

farnesol's physiological functions. These networks include: 1) white-opaque switching, 2) yeast-mycelia

dimorphism, 3) response to cell stress, and 4) cell cycle progression. These networks integrate

environmental signals to influence the morphology of C. *albicans*. Firstly, three of the transcription

regulators studied, Ahr1, Ssn6, and Tup1, have crucial roles in white-opaque switching (Hernday et al.

2013; Hernday et al. 2016; Alkafeef et al. 2018). Given that farnesol is toxic to opaque cells (Dumitru et

al. 2007), it is intriguing that the *AHR1* deletion mutant over accumulates farnesol while also promoting

a higher frequency of white-opaque switching (Hernday et al. 2013). Additionally, it is unclear whether

Ssn6 and Ahr1 participate in the response to farnesol, but there is ample precedent given that another

transcriptional regulator of the white-opaque switch, Czf1, is involved in farnesol response. Secondly, in

addition to Nrg1 and Tup1 (Kebaara et al. 2008), four more transcriptional regulators of yeast-mycelia

dimorphism have been demonstrated in this study to influence farnesol accumulation including Cph2,

Ash1, Ssn6, and Tec1. Thirdly, *C. albicans* is especially resistant to farnesol compared to other fungi

(Semighini et al. 2006; Brasch et al. 2013; Wang et al. 2014), but the mechanism of this tolerance is

elusive. One possibility concerns the protective effect of using longer chain ubiquinones (UQ). *C.*

albicans and *Candida dubliniensis* use UQ9 rather than UQ7 as in other *Candida* species or UQ6 as in *S.*

cerevisiae (Pathirana et al. 2020). Another possibility could be due to differences in their response to

cellular stress. Given that Msn4 and Cas5 have roles in stress response (Nicholls et al. 2004; Xie et al.

2017) and influence farnesol accumulation and secretion (Fig. 2), it is likely they may also contribute to

farnesol tolerance. Fourthly, cell cycle arrest can also induce filamentous growth of *C. albicans,* either

through depletion of cyclins such as Cln3, Clb2, or Clb3 (Bachewich et al. 2005; Bensen et al. 2005),

depletion of polo-like kinase Cdc35 (Bachewich et al. 2003), or through treatment with cell cycle

inhibitors like hydroxyurea (Bachewich et al. 2005; Chen et al. 2018). Interestingly, expression of the

G1/S cyclins *PCL2*, *CLN3*, and *HGC1* are farnesol regulated (Enjalbert and Whiteway 2005). Given that

PLC2 and *CLN3* are Swi4 regulated (Hussein et al. 2011), how these cyclins influence farnesol

accumulation is an important question for subsequent studies.

There are many questions which can now be approached via the genetic information gleaned

from this screen. Among them are: 1/ How is carbon flow through the farnesyl pyrophosphate (FPP)

branch point regulated? 2/ Is farnesol made directly from FPP by an appropriate pyrophosphatase (e.g.

Dpp1p, Dpp2p, or Dpp3p) or are there other more easily regulated and/or safer biosynthetic pathways?

A pathway suggested from *Arabidopsis* invokes the necessary turnover of farnesylated proteins, where

proteolytic degradation of those proteins releases farnesylated cysteine which can be cleaved by a ligase

to release farnesaldehyde which is then converted to farnesol by a farnesol dehydrogenase (Bhandari et

al. 2010). 3/ Is the disappearance of farnesol later in stationary phase due strictly to evaporation of a

volatile molecule or to enzymatic conversion of farnesol to, for instance, 2,3-dihydrofarnesol (Brasch et

al. 2013; Costa et al. 2020). Significantly, regarding its ability to block germ tube formation and hyphal

development, 2,3-dihydrofarnesol is inactive; with only 0.34% of the activity of farnesol (Shchepin et al.

2003). This question has now been partially resolved by extending the *ΔΔswi4* and *ΔΔrap1* mutant

analyses to 80 hours post inoculation (Fig. 3B). While wild type SC5314 (Boone et al. 2022), SN152[+], and

most of the mutants examined had little farnesol remaining by 80 hours, *ΔΔswi4* and *ΔΔrap1* differed in

that they maintained high levels of farnesol. Clearly, something other than evaporation is occurring.

There is ample precedent for farnesol modifying enzymes in fungi. For instance, trans-trans farnesol can

be converted to a cis-trans farnesol by *Helminthosporium sativum* (Imai and Marumo 1974).

Presumably, the corresponding enzymes for *C. albicans* are not made by *Δswi4* and *ΔΔrap1,* and we are

currently using these genetic clues to follow the metabolic disappearance of farnesol during stationary

phase.

Author contributions:

DJG: conceptualization, methodology, validation, investigation, writing — original draft, writing —

review and editing, visualization, formal analysis, data curation, project administration; JJK: validation,

investigation, writing — review and editing; BMT: validation, investigation, writing — review and editing;

CHTB: methodology, validation, writing – review and editing; ALA: conceptualization, methodology,

resources, supervision, writing — review and editing, project administration; KWN: conceptualization,

methodology, resources, supervision, writing — original draft, writing — review and editing, project

administration, funding acquisition.

Data availability:

The datasets generated during and/ or analyzed during the current study are available at FigShare. This

includes the initial screen (File S1), rescreen dataset (File S2), and farnesol growth curves (File S3).

Acknowledgments:

We are grateful to the Fungal Genetics Stock Center (FGSC) for providing strains, to Prof. Rebecca

Roston and Zach Shomo for use and assistance with the gas chromatography system, and to Prof.

Haoping Liu for pointing out the possible role of opaque *ΔΔtup1.*

Compliance with ethical standards:

Funding: This work was supported by Ann L. Kelsall and the Farnesol and *Candida albicans* Research

Fund, University of Nebraska Foundation. This research did not receive any specific grant from funding

agencies in the public, commercial, or not-for-profit sectors.

Conflict of interest: Daniel J. Gutzmann declares that he has no conflict of interest. Jaxon J. Kramer

declares that he has no conflict of interest. Brigid Toomey declares that she has no conflict of interest.

Cory H.T. Boone declares that he has no conflict of interest. Audrey L. Atkin declares that she has no

conflict of interest. Kenneth W. Nickerson declares that he has no conflict of interest.

Ethical approval: This article does not contain any studies with human participants or animals

performed by any of the authors.

References

Alkafeef SS, Yu C, Huang L, Liu H. 2018. Wor1 establishes opaque cell fate through inhibition of the
general co-repressor Tup1 in *Candida albicans*. PLoS Genet. 14(1):e1007176.
doi:10.1371/journal.pgen.1007176.

Alves R, Barata-Antunes C, Casal M, Brown AJP, Van Dijck P, Paiva S. 2020. Adapting to survive: How
Candida overcomes host-imposed constraints during human colonization. PLoS Pathog. 16(5):e1008478.
doi:10.1371/journal.ppat.1008478.

Askew C, Sellam A, Epp E, Mallick J, Hogues H, Mullick A, Nantel A, Whiteway M. 2011. The zinc cluster
transcription factor Ahr1p directs Mcm1p regulation of *Candida albicans* adhesion. Mol Microbiol.
79(4):940–953. doi:10.1111/j.1365-2958.2010.07504.x.

Bachewich C, Nantel A, Whiteway M. 2005. Cell cycle arrest during S or M phase generates polarized
growth via distinct signals in *Candida albicans*. Mol Microbiol. 57(4):942–959. doi:10.1111/j.1365-
2958.2005.04727.x.

Bachewich C, Thomas DY, Whiteway M. 2003. Depletion of a polo-like kinase in *Candida albicans*
activates cyclase-dependent hyphal-like growth. Mol Biol Cell. 14(5):2163–2180. doi:10.1091/mbc.02-
05-0076.

Bensen ES, Clemente-Blanco A, Finley KR, Correa-Bordes J, Berman J. 2005. The mitotic cyclins Clb2p and
Clb4p affect morphogenesis in *Candida albicans*. Mol Biol Cell. 16(7):3387–3400. doi:10.1091/mbc.e04-
12-1081.

Bhandari J, Fitzpatrick AH, Crowell DN. 2010. Identification of a Novel Abscisic Acid-Regulated Farnesol
Dehydrogenase from *Arabidopsis*. Plant Physiology. 154(3):1116–1127. doi:10.1104/pp.110.157784.

Boone CHT, Gutzmann DJ, Kramer JJ, Atkin AL, Nickerson KW. 2022. Quantitative assay for farnesol and
the aromatic fusel alcohols from the fungus *Candida albicans*. Appl Microbiol Biotechnol.
doi:10.1007/s00253-022-12165-w.

Brasch J, Horter F, Fritsch D, Beck-Jendroschek V, Tröger A, Francke W. 2013 Aug. Acyclic sesquiterpenes
released by *Candida albicans* inhibit growth of dermatophytes. Med Mycol.:1–10.
doi:10.3109/13693786.2013.814174.

Bruno VM, Kalachikov S, Subaran R, Nobile CJ, Kyratsous C, Mitchell AP. 2006. Control of the *C. albicans*
cell wall damage response by transcriptional regulator Cas5. PLoS Pathog. 2(3):e21.
doi:10.1371/journal.ppat.0020021.

Chen C, Zeng G, Wang Y. 2018. G1 and S phase arrest in Candida albicans induces filamentous growth via
distinct mechanisms. Mol Microbiol. 110(2):191–203. doi:10.1111/mmi.14097.

Costa CP, Bezerra AR, Almeida A, Rocha SM. 2020. *Candida* Species (Volatile) Metabotyping through
Advanced Comprehensive Two-Dimensional Gas Chromatography. Microorganisms. 8(12):1911.
doi:10.3390/microorganisms8121911.

Dumitru R, Hornby JM, Nickerson KW. 2004. Defined anaerobic growth medium for studying *Candida*
albicans basic biology and resistance to eight antifungal drugs. Antimicrob Agents Chemother.
48(7):2350–2354. doi:10.1128/AAC.48.7.2350-2354.2004.

Dumitru R, Navarathna DHMLP, Semighini CP, Elowsky CG, Dumitru RV, Dignard D, Whiteway M, Atkin
AL, Nickerson KW. 2007. In vivo and in vitro anaerobic mating in *Candida albicans*. Eukaryot Cell.
6(3):465–472. doi:10.1128/EC.00316-06.

Enjalbert B, Whiteway M. 2005. Release from quorum-sensing molecules triggers hyphal formation
during *Candida albicans* resumption of growth. Eukaryot Cell. 4(7):1203–1210. doi:10.1128/EC.4.7.1203-
1210.2005.

Fu Y, Filler SG, Spellberg BJ, Fonzi W, Ibrahim AS, Kanbe T, Ghannoum MA, Edwards JE. 1998. Cloning
and characterization of CAD1/AAF1, a gene from *Candida albicans* that induces adherence to endothelial
cells after expression in *Saccharomyces cerevisiae*. Infect Immun. 66(5):2078–2084.
doi:10.1128/IAI.66.5.2078-2084.1998.

Ganguly S, Bishop AC, Xu W, Ghosh S, Nickerson KW, Lanni F, Patton-Vogt J, Mitchell AP. 2011. Zap1
control of cell-cell signaling in *Candida albicans* biofilms. Eukaryot Cell. 10(11):1448–1454.
doi:10.1128/EC.05196-11.

García-Sánchez S, Mavor AL, Russell CL, Argimon S, Dennison P, Enjalbert B, Brown AJP. 2005. Global
roles of Ssn6 in Tup1- and Nrg1-dependent gene regulation in the fungal pathogen, *Candida albicans*.
Mol Biol Cell. 16(6):2913–2925. doi:10.1091/mbc.e05-01-0071.

Gow NAR, van de Veerdonk FL, Brown AJP, Netea MG. 2012. C*andida albicans* morphogenesis and host
defence: discriminating invasion from colonization. Nat Rev Microbiol. 10(2):112–122.
doi:10.1038/nrmicro2711.

Hernday AD, Lohse MB, Fordyce PM, Nobile CJ, DeRisi JL, Johnson AD. 2013. Structure of the
transcriptional network controlling white-opaque switching in *Candida albicans*. Mol Microbiol.
90(1):22–35. doi:10.1111/mmi.12329.

Hernday AD, Lohse MB, Nobile CJ, Noiman L, Laksana CN, Johnson AD. 2016. Ssn6 Defines a New Level of
Regulation of White-Opaque Switching in *Candida albicans* and Is Required For the Stochasticity of the
Switch. mBio. 7(1):e01565-01515. doi:10.1128/mBio.01565-15.

Homann OR, Dea J, Noble SM, Johnson AD. 2009. A phenotypic profile of the *Candida albicans*
regulatory network. PLoS Genet. 5(12):e1000783. doi:10.1371/journal.pgen.1000783.

Hornby JM, Jensen EC, Lisec AD, Tasto JJ, Jahnke B, Shoemaker R, Dussault P, Nickerson KW. 2001.
Quorum sensing in the dimorphic fungus *Candida albicans* is mediated by farnesol. Appl Environ
Microbiol. 67(7):2982–2992. doi:10.1128/AEM.67.7.2982-2992.2001.

Hussein B, Huang H, Glory A, Osmani A, Kaminskyj S, Nantel A, Bachewich C. 2011. G1/S transcription
factor orthologues Swi4p and Swi6p are important but not essential for cell proliferation and influence
hyphal development in the fungal pathogen *Candida albicans*. Eukaryot Cell. 10(3):384–397.
doi:10.1128/EC.00278-10.

Imai K, Marumo S. 1974. Stereochemistry of c-1 hydrogen exchange during a Δ2, 3 trans-cis
isomerization of farnesol by *Helminthosporium sativum*. Tetrahedron Letters. 15(49–50):4401–4404.

Inglis DO, Johnson AD. 2002. Ash1 protein, an asymmetrically localized transcriptional regulator,
controls filamentous growth and virulence of *Candida albicans*. Mol Cell Biol. 22(24):8669–8680.
doi:10.1128/MCB.22.24.8669-8680.2002.

Kadosh D. 2019. Regulatory mechanisms controlling morphology and pathogenesis in Candida albicans.
Curr Opin Microbiol. 52:27–34. doi:10.1016/j.mib.2019.04.005.

Kebaara BW, Langford ML, Navarathna DHMLP, Dumitru R, Nickerson KW, Atkin AL. 2008. *Candida*
albicans Tup1 is involved in farnesol-mediated inhibition of filamentous-growth induction. Eukaryot Cell.
7(6):980–987. doi:10.1128/EC.00357-07.

Langford ML, Hargarten JC, Patefield KD, Marta E, Blankenship JR, Fanning S, Nickerson KW, Atkin AL.
2013. *Candida albicans* Czf1 and Efg1 coordinate the response to farnesol during quorum sensing, white-
opaque thermal dimorphism, and cell death. Eukaryot Cell. 12(9):1281–1292. doi:10.1128/EC.00311-12.

Lo H-J, Köhler JR, DiDomenico B, Loebenberg D, Cacciapuoti A, Fink GR. 1997. Nonfilamentous *C.*
albicans Mutants Are Avirulent. Cell. 90(5):939–949. doi:10.1016/S0092-8674(00)80358-X.

Lu Y, Su C, Unoje O, Liu H. 2014. Quorum sensing controls hyphal initiation in *Candida albicans* through
Ubr1-mediated protein degradation. Proc Natl Acad Sci U S A. 111(5):1975–1980.
doi:10.1073/pnas.1318690111.

McCluskey K, Wiest A, Plamann M. 2010. The Fungal Genetics Stock Center: a repository for 50 years of
fungal genetics research. J Biosci. 35(1):119–126. doi:10.1007/s12038-010-0014-6.

Moreno-Velásquez SD, Tint SH, Del Olmo Toledo V, Torsin S, De S, Pérez JC. 2020. The Regulatory
Proteins Rtg1/3 Govern Sphingolipid Homeostasis in the Human-Associated Yeast *Candida albicans*. Cell
Rep. 30(3):620-629.e6. doi:10.1016/j.celrep.2019.12.022.

Murad AM, Leng P, Straffon M, Wishart J, Macaskill S, MacCallum D, Schnell N, Talibi D, Marechal D,
Tekaia F, et al. 2001. NRG1 represses yeast-hypha morphogenesis and hypha-specific gene expression in
Candida albicans. EMBO J. 20(17):4742–4752. doi:10.1093/emboj/20.17.4742.

Neville BA, d'Enfert C, Bougnoux M-E. 2015. *Candida albicans* commensalism in the gastrointestinal
tract. FEMS Yeast Research. 15(7):fov081. doi:10.1093/femsyr/fov081.

Nicholls S, Straffon M, Enjalbert B, Nantel A, Macaskill S, Whiteway M, Brown AJP. 2004. Msn2- and
Msn4-like transcription factors play no obvious roles in the stress responses of the fungal pathogen
Candida albicans. Eukaryot Cell. 3(5):1111–1123. doi:10.1128/EC.3.5.1111-1123.2004.

Noble SM, Gianetti BA, Witchley JN. 2017. *Candida albicans* cell-type switching and functional plasticity
in the mammalian host. Nat Rev Microbiol. 15(2):96–108. doi:10.1038/nrmicro.2016.157.

Pathirana RU, Boone C, Nickerson KW. 2020. Longer Ubiquinone Side Chains Contribute to Enhanced
Farnesol Resistance in Yeasts. Microorganisms. 8(11):1641. doi:10.3390/microorganisms8111641.

Pfaller MA, Diekema DJ. 2007. Epidemiology of Invasive Candidiasis: a Persistent Public Health Problem.
Clin Microbiol Rev. 20(1):133–163. doi:10.1128/CMR.00029-06.

Rauceo JM, Blankenship JR, Fanning S, Hamaker JJ, Deneault J-S, Smith FJ, Nantel A, Mitchell AP. 2008.
Regulation of the *Candida albicans* cell wall damage response by transcription factor Sko1 and PAS
kinase Psk1. Mol Biol Cell. 19(7):2741–2751. doi:10.1091/mbc.e08-02-0191.

Romano A. 1966. Dimorphism. The Fungi, vol. 2. 2:181–209. Academic Press, New York, N.Y.

Saville SP, Lazzell AL, Monteagudo C, Lopez-Ribot JL. 2003. Engineered Control of Cell Morphology In
Vivo Reveals Distinct Roles for Yeast and Filamentous Forms of *Candida albicans* during Infection.
Eukaryot Cell. 2(5):1053–1060. doi:10.1128/EC.2.5.1053-1060.2003.

Sellam A, Chaillot J, Mallick J, Tebbji F, Richard Albert J, Cook MA, Tyers M. 2019. The p38/HOG stress-
activated protein kinase network couples growth to division in *Candida albicans.* PLoS Genet.
15(3):e1008052. doi:10.1371/journal.pgen.1008052.

Semighini CP, Hornby JM, Dumitru R, Nickerson KW, Harris SD. 2006. Farnesol-induced apoptosis in
Aspergillus nidulans reveals a possible mechanism for antagonistic interactions between fungi. Mol
Microbiol. 59(3):753–764. doi:10.1111/j.1365-2958.2005.04976.x.

Shchepin R, Hornby JM, Burger E, Niessen T, Dussault P, Nickerson KW. 2003. Quorum sensing in
Candida albicans: probing farnesol's mode of action with 40 natural and synthetic farnesol analogs.
Chem Biol. 10(8):743–750. doi:10.1016/s1074-5521(03)00158-3.

Wang H, Song W, Huang G, Zhou Z, Ding Y, Chen J. 2011. *Candida albicans* Zcf37, a zinc finger protein, is
required for stabilization of the white state. FEBS Lett. 585(5):797–802.
doi:10.1016/j.febslet.2011.02.005.

Wang X, Wang Y, Zhou Y, Wei X. 2014. Farnesol induces apoptosis-like cell death in the pathogenic
fungus *Aspergillus flavus*. Mycologia. 106(5):881–888. doi:10.3852/13-292.

Xie JL, Qin L, Miao Z, Grys BT, Diaz JDLC, Ting K, Krieger JR, Tong J, Tan K, Leach MD, et al. 2017. The
Candida albicans transcription factor Cas5 couples stress responses, drug resistance and cell cycle
regulation. Nat Commun. 8(1):499. doi:10.1038/s41467-017-00547-y.

Zordan RE, Miller MG, Galgoczy DJ, Tuch BB, Johnson AD. 2007. Interlocking transcriptional feedback
loops control white-opaque switching in *Candida albicans*. PLoS Biol. 5(10):e256.
doi:10.1371/journal.pbio.0050256.

Tables

Table 1. Summary of transcription regulator mutants with statistically significant differences in farnesol accumulation in YPD media at $30°C$.[a/]

Mutant	*S. cerevisiae* ortholog or best hit from CGD	Mean whole culture E,E-farnesol $\log_2$ fold change vs WT control ± SD (n=6)	Mean supernatant fraction E,E-farnesol $\log_2$ fold change vs WT control ± SD (n=6)	Predicted or known biological process(es) in *Candida albicans* adapted from CGD (Skrzpek et al. 2017)
\multicolumn{5}{c}{E,E-farnesol over accumulating mutants}
ΔΔswi4	SWI4	2.57*** ± 0.39	3.11*** ± 0.50	Component of SBF complex, G1/S progression
ΔΔahr1	LYS14	2.27*** ± 0.42	2.56 ***± 0.32	Regulation of adhesion genes, white/opaque switch, repressor of START
ΔΔnrg1	NRG2	1.73*** ± 0.45	1.93*** ± 0.49	Regulator of hyphal gene induction, virulence, chlamydospore formation, stress response
ΔΔrap1	RAP1	1.15*** ± 0.62	2.42*** ± 0.48	Telomere maintenance
ΔΔpho4	PHO4	1.07*** ± 0.53	1.68*** ± 0.23	Phosphate acquisition
ΔΔtye7	TYE7	0.95** ± 0.21	1.83*** ± 0.07	Regulator of glycolysis and carbohydrate metabolism
ΔΔash1	ASH1	0.82* ± 0.43	0.92** ± 0.28	GATA-like transcription factor required for WT virulence and filamentous growth
ΔΔaaf1	none	0.76* ± 0.31	1.21*** ± 0.19	Expression in *S. cerevisiae* increases adherence
ΔΔrtg1	RTG1	0.63 ± 0.49	0.96** ± 0.20	Regulation of sphingolipid homeostasis, galactose catabolism

ΔΔssn6	CYC8	0.20 ± 0.90	0.99*** ± 0.73	Forms a corepressor complex with Tup1, regulates a distinct set of genes different than Tup1 in *C. albicans*
ΔΔwor2	UME6	0.43 ± 0.82 (p=0.999)	0.59 ± 0.83 (p=0.217)	Regulator of white/opaque switching, maintenance of opaque state
ΔΔtac1[#]	HAL9	0.23 ± 0.56 (p=0.999)	0.36 ± 0.26 (p=0.919)	Transcriptional activator of drug-responsive genes *CDR1* and *CDR2*
ΔΔcsr1[#]	ZAP1	0.32 ± 0.31 (p=0.984)	0.60 ± 0.13 (p=0.195)	Transcription factor, role in zinc homeostasis
ΔΔupc2[#]	UPC2	0.11 ± 0.43 (p=0.999)	0.31 ± 0.46 (p=0.980)	Regulator of ergosterol biosynthetic genes and sterol uptake
ΔΔtup1[#]	TUP1	0.0 ± 0.15 (p=0.999)	-0.02 ± 0.47 (p=0.999)	Transcriptional corepressor with *Ssn6p*, regulates morphological switching, farnesol response, germ tube formation
E,E-farnesol under accumulating mutants				
ΔΔ orf19.6874 (C2_05640 W_A)	BAS1	-2.10*** ± 0.73	-2.65*** ± 1.03	Transcription factor, null mutant exhibits increase in invasive growth
ΔΔmsn4	MSN2	-1.89*** ± 0.37	-2.40*** ± 0.65	Zinc finger transcription factor, not a significant stress response regulator in *C. ablicans*
ΔΔcph2	HMS1	-1.93*** ± 0.63	-2.04*** ± 0.91	Transcriptional activator of hyphal growth
ΔΔfcr1	CAT8	-1.97*** ± 0.88	-3.68*** ± 3.83	Negative regulator of fluconazole/ketoconazole/brefeldin A resistance
ΔΔsko1	SKO1	-1.93*** ± 0.39	-1.41*** ± 0.46	Cell wall damage response, morphological switching, oxidative stress response
ΔΔgzf3	GZF3	-1.32*** ± 0.32	-1.63*** ± 0.54	GATA type transcription factor, ortholog regulates nitrogen catabolic gene expression

ΔΔcas5	*MOT3*	-1.16** ± 0.13	-0.51 ± 0.32	Cell wall homeostasis, adherence, stress response, cell cycle
ΔΔcup9	*TOS8*	-1.08** ± 0.63	-0.85 ± 0.86	Represses SOK1 expression in response to farnesol inhibition
ΔΔtec1	*TEC1*	-0.75* ± 0.66	-0.58 ± 0.76	Hyphal gene regulator, white cell pheromone response

a/ Log_2 fold change production values ± SD represent the mean farnesol accumulation value of 3 X and 3
Y (n=6) independently derived mutant compared to the mean farnesol accumulation value of the paired
SN152 parent (n=22. Both over and under accumulating mutants are ordered by how much they exceed
the SN152 parent. The four mutants added because of literature precedent are indicated by #.
Differences between mutants (n=6) and the SN152[+] parent (n=22 batches) were accessed by One-way
ANOVA with Dunnett's multiple comparisons test using GraphPad Prism. Differences were considered
significant at p<0.05 (*p<0.05, **p<0.01, ***p<0.001).

Figure Captions

Fig. 1. Initial screen E,E-farnesol supernatant to pellet ratio vs E,E- farnesol whole culture accumulation

at 24 hours post inoculation. Data represents the accumulation values for each of 164 transcription

regulator mutants (black) (n=1) and each SN152[+] replicate (red) (n=34). The red X indicates the mean

value for the SN152[+] replicates (n=34). Dotted lines represent a log2 fold change increase or decrease of

1 for farnesol from the mean for SN152[+].

Fig. 2. Transcription regulator mutants differing in farnesol accumulation and localization. Accumulation

values were measured 24 hours post inoculation at 30°C in YPD for each transcription regulator mutant

in both the X (3) and Y (3) independent mutants (total n=6) and the SN152[+] parent (n=22 batches). Data

represents the mean accumulation value ± SD (A,B) or mean farnesol supernatant to pellet ratio ± SD

(C). The mutants added because of literature precedent are indicated by #. Differences between groups

that were normally distributed and that had equal variances (C) were accessed by One-way ANOVA with

Dunnett's multiple comparisons test. Differences between groups that were not normally distributed or

heteroscedastic (A,B) were accessed by Kruskal-Wallis test with Dunn's multiple comparisons tests.

Differences were considered significant at p<0.05 (*p<0.05, **p<0.01, ***p<0.001).

Fig. 3. Temporal dynamics of farnesol accumulation in *ΔΔswi4, ΔΔahr1, ΔΔrap1, and ΔΔnrg1.* Farnesol

accumulation was accessed at 12, 18, 24, 36, 48, 80 hours post inoculation at 30°C in YPD for each

transcription regulator mutant in two independent growth curves (n=2) and 3 independent growth

curves for SN152[+] (n=3). Dry weights were accessed as each time point and are presented as the mean

dry weight ± SEM (A). Farnesol accumulation data represents the mean dry weight normalized farnesol

accumulation value ± SEM in either the whole culture (B) or the supernatant fraction (D). To access

overall farnesol production, area under the curve analyses (AUC) were performed and data are the

mean AUC ± SE for the whole culture (C) or supernatant fraction (E). Differences in the AUC between

groups were accessed by One-way ANOVA with Dunnett's multiple comparisons test. Differences were

considered significant at p<0.05 (*p<0.05, **p<0.01, ***p<0.001).

Fig. 4. Temporal dynamics of farnesol accumulation in *ΔΔorf19.6874, and ΔΔmsn4.* Farnesol

accumulation was accessed at 12, 18, 24, 36, 48, 80 hours post inoculation at 30°C in YPD for each

transcription regulator mutant in two independent growth curves (n=2) and 3 independent growth

curves for SN152[+] (n=3). Dry weights were accessed as each time point and are presented as the mean

dry weight ± SEM (A). Farnesol accumulation data represents the mean dry weight normalized farnesol

accumulation value ± SEM in either the whole culture (B) or the supernatant fraction (D). To access

overall farnesol production, area under the curve analyses (AUC) were performed and data are the

mean AUC ± SE for the whole culture (C) or supernatant fraction (E). Differences in the AUC between

groups were accessed by One-way ANOVA with Dunnett's multiple comparisons test. Differences were

considered significant at p<0.05 (*p<0.05, **p<0.01, ***p<0.001).

Fig. 5. Characterization of *ΔΔtup1* morphology in conditions used for the screen. (A) Representative

micrographs of the colony morphology of SN152$^+$ (white) and *ΔΔtup1* (pink) on YPD agar supplemented

with phloxine B grown at 30°C . (B) Representative micrographs of cellular morphology of SN152$^+$ (yeast)

and *ΔΔtup1* (hyphae and elongated cells).

A

SN152⁺

ΔΔtup1

B

SN152⁺

ΔΔtup1

50 µm

50 µm