Research Advances:

Five years later, with double the demographic data, naked mole-rat
mortality rates continue to defy Gompertzian laws by not increasing
with age

J. Graham Ruby[1], Megan Smith[1], Rochelle Buffenstein[1,2,3]

1: Calico Life Sciences LLC, South San Francisco, United States
2: current affiliation: Department of Biological Sciences, University of Illinois, Chicago
3: for correspondence: rbuffen@uic.edu

<u>Abstract</u>

The naked mole-rat (*Heterocephalus glaber*) is a mouse-sized rodent species, notable
for its eusociality and long lifespan. Previously, we reported that demographic aging,
i.e., the exponential increase of mortality hazard that accompanies advancing age in
mammals and other organisms, does not occur in naked mole-rats (Ruby et al, 2018).
The demographic data supporting that conclusion had taken over three decades to
accumulate, starting with the original rearing of *H.glaber* in captivity. In the five years
following that study, we ~doubled our quantity of demographic data. Here, we re-
evaluated our prior conclusions in light of these new data and found them to be
supported and indeed strengthened. We additionally provided insight into the social
dynamics of captive *H.glaber* with data and analyses of body weight and colony size
versus mortality. Finally, we provide a phylogenetically-proximal comparator in the form
of lifespan data from our Damaraland mole-rat (*Fukomys damarensis*) colony and
demographic meta-analysis of those data along with published data from Ansell's mole-
rat (*Fukomys anselli*). We found *Fukomys* mortality hazard to increase gradually with
age, an observation with implications on the evolution of exceptional lifespan among
mole-rats and the ecological factors that may have accompanied that evolution.

<u>Introduction</u>

The species *Heterocephalus glaber*, commonly known as the naked mole-rat, is a
eusocial mammal endemic to the arid and semi-arid regions of northeast Africa. In the
wild, naked mole-rats live an almost completely subterranean lifestyle, in colonies of up

to 295 animals (average size is 60 animals/colony; Brett, 1991) that cohabitate a
network of tunnels that the mole-rats dig themselves with their large, ever-growing
incisors (Buffenstein et al, 2021). Each colony is usually led by a single, socially-
dominant queen (breeding female), living with a small number of breeding males and a
large number of non-breeding progeny. New breeding females arise either through
conflict following the death of a queen or isolation of part of the colony, e.g. through
tunnel collapse (Brett, 1991; Jarvis, 1981; Buffenstein et al., 2022).
44
Naked mole-rats are notable for their extreme lifespans (Buffenstein 2005; Lewis and
Buffenstein 2016), living longer than any other documented rodent (Buffenstein and
Jarvis, 2002), with the longest previously-reported lifespan of 37 years (Lee et al.,2020)
and many animals living beyond 30 years (Lewis et al., 2018; Can et al., 2022). These
values are notable in the context of this species' small body size due to the strong
correlation across species between that value and mammalian lifespan: maximum
lifespan potential (MLSP) increases by 16% for each doubling of average species body
mass (Hulbert et al., 2007). Notably, naked mole-rats live approximately five times
longer than their size-corrected, allometric expectation using the equations of (de
Magalhães et al., 2007). For most mammalian species, lifespan is limited by an
exponential increase in the per-day risk of death (i.e. mortality hazard) with age
(Gavrilova & Gavrilov, 2015; Cohen, 2018) in accord with a statistical distribution first
defined by Gompertz based on human mortality (Gompertz, 1825). The increase in
mortality hazard with age is referred to as "demographic aging". We have previously
shown *H.glaber* to achieve its exceptional longevity through defiance of this trend,
exhibiting ~constant mortality hazard across the full spectrum of observed lifespans,
with no hazard increase evident even many-fold beyond their expected MLSP (Ruby et
al, 2018 & 2019).
63
Naked mole-rats are members of the the African mole-rat superfamily, Bathyergidae,
that includes more than 30 species of subterranean rodents and appeared to split from
the other bathyergids ~26 million years ago (MYA) (Fang et al., 2014; Visser et al.,
2019; Patterson and Upham, 2014; Faulkes and Bennett 2021). Bathyergid species
span the spectrum of social behavior, from solitary (e.g., *Georychus, Heliophobus* and
Bathyergus) to the only other known eusocial mammals, members of the genus
Fukomys (Jarvis and Bennett, 1993). Along with eusociality, *H.glaber* also share
above-expected longevity with members of the *Fukomys* genus, namely *Fukomys*
damarensis and *Fukomys anselli,* living ~twice as long as their size-corrected, allometric
expectations (see Methods for details). The co-occurrence of these unusual properties
in divergent mole-rat clades motivates a question about the similarity of age-related
hazard patterns.
76

For naked mole-rats, the lack of demographic aging is accompanied by seemingly-
indefinite maintenance of many physiological characteristics that typically change with
age. Naked mole-rats are resistant to age-related diseases such as cancer,
neurodegeneration, and cardiovascular disease (Edrey et al., 2011; Hadi et al., 2021;
Can et al., 2022) and show signs of tissue regeneration and remodeling preventing the
deterioration of age-associated physiological function (Buffenstein et al., 2020, Can et
al., 2022; O'Connor et al., 2002; Buffenstein and Craft, 2021). However, controversy
regarding the perpetual neoteny of *H.glaber* (Orr et al., 2016; Grimes et al., 2017;
Skulachev et al., 2017; Buffenstein et al., 2020 remains, supported by evidence for
chronological aging of the epigenetic (DNA methylation) clock, albeit showing that
breeders maintain a more youthful methyl clock (Kerepesi et al., 2022). This is further
stoked by a commentary on our original demographic analysis (Ruby et al, 2018) by
(Dammann et al, 2019), asserting that loss of records from our early years of our
vivarium rearing of *H.glaber* could have confounded our analysis of aging
demographics. The concern of that critique has been addressed using left-censorship
(Ruby et al, 2019). However, the critique also highlights the unique nature of our
collection as a resource for powered demographic analysis of naked mole-rats,
extending out to advanced ages. The lack of other collections of a similar scale,
worldwide, presents a practical challenge to the reproducibility of our results. To that
end, we were motivated to re-evaluate our original conclusions, using demographic data
for *H.glaber* collected over a five-year period since the data freeze from (Ruby et al,
2018).

Here, we re-visited the demographic analysis of our naked mole-rat collection, with
husbandry data now extended by five years across an expanded set of animals. We
found our original conclusions of naked mole-rat mortality hazard being age-
independent to be reproduced, using either the total sum of all historical data or only
those data collected after our previous study - the latter qualifying as a replication study.
We further conducted an analysis of naked mole-rat mortality hazard versus colony size
which suggested that social competition was greater for non-breeders in smaller
colonies. Finally, we provide demographic data for Damaraland mole-rats: combining
those data with other publicly available data for *Fukomys* species, we analyzed age-
specific mortality hazard and showed it to increase, in contrast to the naked mole-rats.

Results

The expanded data set revealed that the hazard of naked mole-rat mortality did
not increase as a function of age

Our expanded data set drew from 7,536 historically catalogued naked mole-rats from
our own collection; of these, 6,957 had sufficiently high (single-day) resolution records

for date-of-birth and date-of-death/censorship to be included (see Methods for details).
Our analyses of survival and mortality hazard began at T_{sex} for naked mole-rats: 6
months (183 days; Edrey et al., 2011); 6893 animals had aged up to T_{sex} and therefore
were included in our downstream analyses. Of those, 755 had recorded deaths and
6138 were right-censored (see Methods for censorship criteria). Kaplan-Meier survival,
along with death and censorship distributions, are plotted in Figure 1A. The last
recorded death was at 11,281 days-of-life (30.9 years), at which point 10 animals
remained alive; the last of those was censored at 12,873 days-of-life (35.2 years). At
that event, Kaplan-Meier survival remained at 55.5%.
124
Using the survival data, we calculated age-specific mortality hazard across the same
age-range windows as in (Ruby et al, 2018) (Figure 1B). As in the prior analysis,
mortality hazard estimates remained consistently low across the ~35 years of age
encompassed by the observations, never exceeding a per-day probability of 1/10,000.

Analysis of only data collected after our prior survey independently confirmed
constant-hazard demography.

The large size of our naked mole-rat collection across the five years of additional
observation time since our previous report (Ruby et al, 2018) provided substantial
additional power in the form of number of animals observed at most ages (Fig. 2A). For
many age groups, the amount of lifespan observation time since our previous snapshot
matched or exceeded that prior to it. This allowed us to use left-censorship at a date
after our original dates of record aggregation to perform an independent analysis of
naked mole-rat lifespan demographics and critically re-evaluate our original
conclusions.
139
For our previous report, lifespan records were aggregated in three batches: on April 14,
2016; April 20, 2016; and May 17, 2016 (Ruby et al, 2018). Animal counts as a function
of age, Kaplan-Meier survival, and age bin-specific hazard are shown for that data set in
magenta in Fig. 2A, B, and C, respectively. The possibility that incomplete record
survival during the earliest years of our keeping naked mole-rats in captivity could have
skewed mortality hazard trends was raised by Dammann et al (2019), and we
previously addressed that concern by excluding data from those early years of naked
mole-rat captivity, via left-censorship on January 1, 2008 (Ruby et al, 2019). No
additional data were provided at that time. The scale, Kaplan-Meier survival, and age-
specific hazard from that left-censored version of our original data set are shown in
orange in Fig. 2A, B, and C, respectively.
151
The scale, Kaplan-Meier survival, and age-specific hazard from our current, full data set
– also depicted in Figure 1 – are shown in green in Figures 2A, 2B, and 2C,

respectively. These data were aggregated on March 15, 2021 and encapsulated all
historical data. In order to perform an analysis of naked mole-rat lifespan demographics
that would be independent of our original analysis, we left-censored these data on May
17, 2016. The scale, Kaplan-Meier survival, and age-specific hazard from this left-
censored data set are shown in blue in Fig. 2A, B, and C, respectively. For each age
bin, the new, left-censored hazards were either lower or statistically indistinguishable
from the original data set, either full or left-censored (Fig. 2C). We therefore maintain
our original conclusion: that naked mole-rat mortality hazard does not increase with age,
with that assertion being statistically robust to the beginning of our final estimate bin
(6,529 days, or ~18 years).

Non-increasing mortality hazard was exhibited by breeders and non-breeders of
both sexes

The analyses described above were performed on our full population of animals. Sub-
populations defined by sex and breeding-status were analyzed individually for Kaplan-
Meier survival (Figure 3A). As previously observed (Ruby et al, 2018), there was a
notable survival difference between breeding versus non-breeding status, with breeders
enjoying longer survival than non-breeders. As with the aggregate population, mortality
hazard remained consistent with age, for both non-breeders (Figure 3B) and breeders
(Figure 3C), with the mortality hazard of breeders only ~20% that of non-breeders. As
previously observed, male non-breeders exhibited enhanced survival (Figure 3A) and
reduced mortality hazard (Figure 3B) at ages beyond 2000 days. This concurred with
the expectations of our model that multiple males can enjoy breeder status in a colony
despite not being labeled as such (only the founding breeders are so labeled; Ruby et
al, 2018).

Non-breeder mortality hazard was influenced by colony size

Some of the mortality hazard difference between breeders and non-breeders may
derive from social struggle between these hierarchies within the colony. To explore the
potential effects of intra-colony social dynamics on mortality hazard, we performed
analyses of the relationships between body weight or colony size on the mortality of
non-breeders.
184
For body weights, we analyzed a data set that had been aggregated to accompany our
originally published demographic data (Ruby et al, 2018; Supplementary Table 2).
These data included 19,118 individual body weight estimates from 1,940 animals,
captured longitudinally for animals of a variety of ages, with data becoming sparse after
~3500 days of age (Figure 4A). These data are provided in Supplemental Data File 2.
A great deal of variation existed across both the breeder and non-breeder populations,

in both males and females, though the age-specific averages were similar and stable
after ~500 days of age (Figure 4B).
193
For the analysis of mortality hazard versus body weight, only non-breeding animals
were included. The last available body weight measurement was used, requiring that
measurement was taken after 500 days of age, to avoid confounding with the social
status of animals that were not yet fully grown. There were 6,785 qualifying
measurements, derived from 942 animals. For each animal, mortality hazard was
analyzed across one year following that final measurement. Non-breeding animals of
each sex were grouped into quartiles based on body weight, and the aggregate
mortality hazard of each quartile is plotted, along with the aggregate body weight
values, in Figure 4C. There was a general trend of lighter/smaller animals having
greater mortality hazard, especially in males, though the data set was insufficiently
powered to statistically support this conclusion.
205
For colony sizes, we used our records' annotations of "colony ID" to determine the
number of non-breeding animals of each sex in each colony, at each analytical time
point (see Methods). As for body weight, animals were separated into quartiles based
on the number of non-breeders of the same sex in the colony, and each quartile for
each sex was cumulatively evaluated for mortality hazard across one year following the
analytical time point. This method allowed independent analysis of non-overlapping
time periods: we performed five such analyses using our new data set and four using
our prior data set, from Ruby et al, 2018 (examples in Figure 4D; all results shown in
Supplemental Figure S2). Analyses of individual time blocks were statistically
inconclusive, but there was a general trend of greater mortality hazard for non-breeders
in smaller colonies. Meta-analysis of the results from across all nine time-block
analyses supported this trend to a statistically significant degree, for both males (p-
value: $9.9 * 10^{-4}$) and, to a lesser degree, females (p-value: $1.8 * 10^{-2}$).
219

220 **The mortality hazard of mole-rats of the Fukomys genus increased substantially**
221 **with age**
222
We have previously placed the mortality demographics of naked mole-rats into the
context of demographic data from other mammals, which highlights the unique absence
of an exponential increase in hazard with age in naked mole-rats (Ruby et al, 2018). To
provide a more phylogenetically-proximal comparator, we performed a demographic
analysis of mole-rats from the *Fukomys* genus. For those species, survival analyses all
began at an estimated T_{sex} of 270 days of age, based upon our unpublished data and

personal communication from Nigel C. Bennett (MRI, University of Pretoria) and

Christopher G. Faulkes (Queen Mary University of London).

In their response to our previous publication, Dammann et al (2019) provide lifespan

data for 339 *Fukomys anselli* individuals, of which 332 aged past T_{sex} and therefore

were included in our downstream analyses (Supplemental Figure S2A: yellow). In their

analyses of lifespan, Dammann et al (2019) censor accidental deaths and deaths from

fighting: in that case, 170 animals' lifespans are censored and 162 died. Here and

previously (Ruby et al 2018,2019), we treat such instances as non-censored deaths: in

that case, 145 animals' lifespans are censored and 187 had died. Comparing Kaplan-

Meier survival analyses, the death versus censorship of these 25 individuals had a

small effect (Supplemental Figure S2B: yellow versus orange), only reducing the

median lifespan from 3,141 to 2,830 days; nor were any statistical differences observed

for age-specific mortality hazard estimates effect (Supplemental Figure S2C: yellow

versus orange). For downstream analyses, we considered those 25 individuals to be

dead rather than censored.

We provide additional demographic data for another mole-rat species within the

Fukomys genus (*Fukomys damarensis*). These data were compiled on March 15, 2021

and included lifespans for 702 animals, of which 634 aged past T_{sex} and therefore were

included in downstream analyses (Supplemental Figure S2A: cyan). These included

502 animals' lifespans that were censored and 132 deaths. Versus the *F.anselli* data

set, these data for *F.damarensis* were richer for young and sparser for old ages

(Supplemental Figure S2A). Nonetheless, these two species exhibited similar

demographic properties, both in terms of Kaplan-Meier (Supplemental Figure S2B) and

age-specific mortality hazard (Supplemental Figure S2C). For additional downstream

analyses, we combined the data from *F.anselli* and *F.damarensis* into a single data set,

representing the *Fukomys* genus (Supplemental Figure S2A: green). The combined

Fukomys data set included lifespan data from 1041 animals, 966 of which survived past

T_{sex} (270 days) so were included in downstream analyses; these included 319 deaths

and 647 censorships.

Kaplan-Meier survival analysis revealed a *Fukomys* median lifespan of 3,622 days

(Figure 5A). Unlike the survival curve for *H.glaber*, the *Fukomys* survival curve

approached zero before being depleted of animals, allowing for calculation of a maximal

lifespan, as previously defined (95th percentile; Ruby et al, 2018) of 7,369 days (Figure

5A), or 20.2 years. Age-specific mortality hazard was calculated for *Fukomys*, and

again unlike *H.glaber*, substantial increase was observed with age (Figure 5B). Though

less statistically robust, this trend was consistent with estimates taken from either the

F.damarensis data alone, or from the *F.anselli* data alone, with or without
accidental/fighting death censored (Supplemental Figure S2C).

Mole-rat mortality hazards showed lesser increase with age versus other
mammals, with *H.glaber* still uniquely constant.

Age of reproductive maturity (T_{sex}) for a species is also generally predictive of species
MLSP (Hamilton, 1966; Rose et al., 2007; Prothero, 1993). As described in Ruby et al
(2018) and shown again here (Figure 6A), the failure of naked mole-rat mortality hazard
to increase with age, even at many-fold the species' T_{sex}, is unique among the
mammals analyzed. These included laboratory mice (*Mus musculus*) using the raw
data for the control mice cohort from the published paper of Miller (Miller et al., 2014),
with T_{sex} at 42 days post-natal (Tacutu et al., 2018) and a median lifespan of 847 days;
humans (*Homo sapiens*), as reported for a 1900 birth cohort (Bell and Miller, 2005) and
with T_{sex} defined as 16 years (Apter, 1980; Anderson et al., 2003); and horses (*Equus*
ferus caballus), using insurance tables (Valgren, 1945) and T_{sex} at three years (Tacutu
et al., 2018). All of these species are anciently diverged from *H.glaber*, with its closest
relative being *M.musculus*, that diverged from *H.glaber* ~73 MYA (Kim et al., 2011). In
contrast, *H.glaber* diverged from humans ~88 MYA and from horses ~95 MYA (Springer
et al, 2003). These evolutionary relationships are depicted in Figure 6B.

Here, we added *Fukomys* mole-rat mortality hazard, age normalized to a T_{sex} of 270
days, to our comparative mammalian hazard plot (Figure 6A). *Fukomys* diverged from
H.glaber 26 MYA (Figure 6B; Fang et al, 2014) and displayed an intermediate rate of
hazard increase between the non-increasing hazard of *H.glaber* and the exponentially-
increasing hazard of other mammals. *Fukomys* mortality hazard appeared to increase
linearly, rather than with a Gompertzian exponential. The sparsity of late-life data for
that genus made conclusions tentative. Additionally the uncertainty of age at T_{sex} made
these conclusions tentative: we based our value of 270 days on our own experience,
but alternative T_{sex} values are reported for *F.anselli* and *F.damarensis* on the AnAge
website (Tacutu et al, 2018). However, with only 2.0% of the population surviving until
the collection of the final datum (Figure 5A), it was unlikely that trend derived from
observation through an insufficient portion of the genus' lifespan.

Discussion

The elusion of demographic aging by *H.glaber* was replicated in an additional 5
years of new data

305
In this study, we re-visited our analyses and conclusions from (Ruby et al, 2018),
bringing five years' worth of additional data collection to bear. That period represented
a period of unprecedented scale and stability for our collection: while animals continued
to be killed for research (KFR) purposes, there were fewer constraints on husbandry
resources to require culling of the population. Animals remained at the same
institutional vivarium across that period (Calico Life Sciences, LLC), thus avoiding the
potentially confounding husbandry challenges of institutional and geographic relocation.
Especially given the mature status of the collection across this period (i.e., many
animals of advanced age already present), these conditions facilitated an approximate
doubling of the available observation numbers for demographic analysis across most
age categories (Figure 2A).
317
These additional data provided two opportunities: first, to repeat our original analyses
on the full, updated historical data, which added resolution and statistical confidence to
our original claims (Figure 1). Second: the scale of data collected since our last study
facilitated an independent analysis of *H.glaber* lifespan demography, performed entirely
on data collected after the completion of data collection for our prior study (navy in
Figure 2). Importantly, that independent analysis validated our original claims about
naked mole-rat demography.
325
326 **Elevated hazard for non-breeders in small colonies**
327
We previously reported that breeders of both sexes of naked mole-rats have
substantially lower mortality hazards than non-breeders at all ages (Ruby et al, 2018),
and that result was reproduced here (Figure 3). This result was not limited to *H.glaber*:
our analysis of *F.damarensis* survival also revealed an advantage for breeders over
non-breeders. This trait is shared with many cooperative breeding mammals (Sharp
and Clutton-Brock, 2010; Dammann et al., 2011, Cram et al., 2018). These
observations conflict with the disposable soma theory of aging (Kirkwood, 1977) but do
not negate it: there are also multiple biological and sociological explanations for the
survival advantage of socially-dominant individuals (e.g. Clarke and Faulkes, 1997;
Medger et al., 2019; Sahm et al., 2021; Coen et al., 2021; Berger et al., 2018) that may
confound one another and variably contribute to the fates of breeders versus non-
breeders.
340
341
In order to gain deeper insight into the health consequences of naked mole-rat social
hierarchy amongst non-breeders, we analyzed historical records of body weight and
colony membership in terms of their impacts on mortality hazard (Figure 4). Dominance

in naked mole-rat colonies is largely determined by aggressive pushing, tail-tugging,
and shoving behaviors, so smaller animals are generally lower in the social hierarchy
than larger (Clarke and Faulkes 1997; Toor et al., 2020; Holmes and Goldman, 2021).
It was therefore unsurprising to observe that smaller adult animals had higher mortality
hazard than bigger animals (Figure 4C), though that trend did not attain statistical
significance in our analysis.
351
More surprising was the observation that non-breeders who were members of large
colonies had lower mortality hazard than non-breeders who were members of small
colonies, an observation that was statistically significant for both sexes (Figure 4E).
One might have expected fewer colony members to result in less aggressive
competition for, and therefore better access to, limited resources, e.g., food and heating
pads (Clarke and Faulkes, 2001). However, such concerns may have been less
relevant to our captive colonies than in the wild: animal husbandry rules necessitate the
scaling up of all resources in parallel with colony growth (Smith & Buffenstein, 2021).
Further, even in the wild, intra-group competition is often trumped by cooperative
advantages to produce overall survival advantages for members of larger social groups
across a variety of species (Jarvis et al., 1998; Silk et al., 2010; Broom et al, 2009;
Clutton-Brock, 2021). Larger colonies can also provide greater stability, with a well-
established successfully breeding female and a cohort of older siblings having already
resolved their dominance status and keeping the younger animals in check (Smith &
Buffenstein, 2021).
367
368 **The mortality hazard of *Fukomys* mole-rats increased gradually with age**
369
We sought to provide new evolutionary context for the exceptional longevity of naked
mole-rats through demographic analysis of mole-rats from the *Fukomys* genus. Though
substantially diverged from *H.glaber* 26 MYA (Fang et al., 2014), *Fukomys* mole-rats
offered a phylogenitically closer comparator versus other mammals for which data are
available (Figure 6B). Further, *Fukomys* mole-rats share the property of eusociality with
naked mole-rats, which is thought to be connected to exceptional lifespan by
evolutionary pressures (Kim et al., 2011; Fang et al., 2014). Demographic data were
published for Ansell's mole-rat, *F.anselli* (Dammann et al., 2019); we supplemented
those data with lifespan data from the closely-related Damaraland mole-rat,
F.damarensis. Those two species are estimated to have diverged ~1-1.5 MYA (Van
Daele et al, 2007), and they shared a similar ecological niche in the wild and statistically
consistent life history patterns in captivity (Supp Figure S2). We combined the data
from these species for a *Fukomys* genus-wide demography analysis and observed
hazard to increase with age (Figure 5B).
384

While the increase of mortality hazard with age among *Fukomys* mole-rats contrasted
with the age-independent hazard of *H.glaber*, it was gradual versus the Gompertzian
(i.e. exponential) increase typically exhibited by mammals (Figure 6A). Various
evolutionary theories highlight multiple aspects of mole-rat biology and lifestyle that
could either drive or derive from the evolution of such extended longevity.
Subterranean rodents, regardless of clade or whether they are solitary or social, live
longer than expected for their size, with environmental protection from extrinsic mortality
(Buffenstein 2005; Healy et al., 2014). Similar protection from extrinsic mortality is also
provided by herd protection/social lifestyle (Buffenstein 2005;Healy et al., 2015). Low
extrinsic hazards give evolutionary tinkering with aging rates the opportunity to manifest
selective advantage (Jacob, 1977), a premise backed up by several cross-species
studies (Gaillard & Lemaître 2017). In the context of eusociality, kin selection and the
maintenance of stable social hierarchies can provide further selective advantages to the
extension of lifespan for breeders (Clarke and Faulkes, 1997; Medger et al., 2019; Sahm et
al., 2021; Coen et al 2021). The converse may also be true, with enhanced longevity
facilitating a eusocial lifestyle (Ross et al., 2015): increased longevity may facilitate the
cohabiting of parents and multiple litters of offspring to form a eusocial colony,
enhancing mutual beneficial opportunities for cooperative living and overall fitness of the
social group (Carey, 2001; Lucas and Keller, 2020).
404
Regardless of which evolutionary theory best applies to the natural history of *H.glaber*
and *Fukomys* mole-rats, their shared (though unequal) defiance of Gompertzian aging
demonstrates that exponentially-increasing hazard with advancing age is not a fixed
property of mammalian biology. While it seems to be the default, evolutionary
mechanisms have managed to dampen or eliminate it when provided with the right
selective incentives. It is not clear the extent to which both *H.glaber* and *Fukomys*
benefit from longevity genes that arose in their common ancestor, versus evolved
independent longevity-promoting mechanisms in the 26 million years since their
divergence. However, their unequal successes suggest that multiple mechanisms are
needed to counteract Gompertzian aging, and that not all mechanisms are at play in
both lineages. The perspective that some age-slowing mechanisms are likely to be
conserved between *H.glaber* and *Fukomys*, and that others are likely to be specific to
H.glaber, should help the comparative biology community make maximal use of these
two valuable natural resources for the science of aging.

Materials and Methods

Animal husbandry

Husbandry practices for *H.glaber* were identical to those described in (Ruby et al,
2018). A similar husbandry protocol was used for the Damaraland mole-rats.

The original progenitors of our Damaraland mole-rat colony were trapped near the town
of Hotazel, South Africa (27°58′S, 17°41′E), and near the town of Dordabis in Namibia
(22°58′S, 17°41′E). Like the naked mole-rat, all animals were microchipped at 90 days
with a unique RFID number. Similarly, Damaraland mole-rats were group-housed in
interconnected transparent multi-cage systems, joined together by plexiglass clear
tubing in a climatically controlled room maintained at 26-28^{o}C, 21% oxygen and with RH
ranging between 20-50% and with a 12L:12D light cycle. Animals were fed daily a
variety of fresh fruit and vegetables, including apples, yams, carrots, pumpkin, and other
seasonal vegetables, replacing the natural geophytes upon which they feed. No water
was provided as the animals meet all their water requirements through their diet or
metabolic water production.

H.glaber demographic data

Demographic analysis of _H.glaber_ was performed on a newly-compiled aggregation of
lifespan data, from across the collection of naked mole-rats described in (Ruby et al,
2018), which had been maintained and expanded under the same husbandry conditions
described there. For this study, a snapshot of records was compiled on March 15,
2021, at which point they contained information for 7,536 animals from across the full
history of the collection. Of those, complete birth and death/right-censorship information
(single-day resolution), were available for 6,949 animals. For those animals, and versus
our prior snapshot of records from this collection – provided as "Supplementary file 1" in
(Ruby et al, 2018) – 3,329 could be matched to the prior records. Of those, 2,806 had
perfect matches to RFID tags, birth dates, sex, breeder status, colony name, and
logically consistent death/censorship data (for animals alive at the prior snapshot, new
death/censorship dates from after those dates; otherwise, perfect matches). For an
additional 94 animals, all data listed above were identical except for either colony name
or breeding status: both of these parameters are expected to change for animals moved
out of a colony to establish a new colony as a founding breeder. Combined, 3,222
animals matched for all vital data (RFID, birth date, sex, and consistent
death/censorship). For an additional 94 animals, one of the four vital data had been
updated: 11 for RFID (consistent with occasional replacement of RFID chips, or
placement of RFID chips in previously un-chipped animals); 32 for sex (consistent with
inaccurate initial sex determination for young animals); 40 for date of birth (consistent
with review of birth data at the time of RFID implantation); and 11 with updated and
inconsistent death/censorship dates. For all 94 of those animals, the updated values
were used for analysis. For an additional five animals, two pieces of vital data differed:
one had inconsistent death/censorship and sex; one had inconsistent death/censorship
and date of birth; and three had different dates of birth and sexes. For those animals as

well, the updated values were used for analysis. For an additional eight animals, exact
dates of death/censorship were missing from current records but found in the prior
snapshots, with all other vital data matching. For those animals, the death/censorship
values and dates from the (Ruby et al, 2018) snapshot were used. An additional 26
animals had RFID matches to animals that had missing or low-resolution birth and/or
death/censorship data in the (Ruby et al, 2018) snapshot; those animals were assigned
their original table ID's and current data values. In total, 3,355 animals were paired to
individuals cataloged in "Supplementary file 1" of (Ruby et al, 2018): all of those are
presented in Supplemental Data File 1 using their original ID's (e.g. "Hg-02241").
473
An additional 3,602 animals with day-resolution birth and death/censorship data were
newly added to the roster presented in Supplemental Data File 1, versus the catalog
from (Ruby et al, 2018). Those animals were all assigned ID numbers ≥5000 (e.g. "Hg-
05072"). The vast majority of those animals were born after (3,244) or during (33) the
prior data consolidation period (April 14 – May 17, 2016). Of the remaining 325
animals, the large majority (285) were born within a year prior to April 14, 2016,
reflecting a normal lag for husbandry records to be entered into our database.
481
In summary: 6,949 naked mole-rats were cataloged with day-resolution birth and
death/censorship data, 3,355 of which could be matched to animals from (Ruby et al,
2018). An additional eight animals with missing death/censorship data in the current
inventory were included using those data from (Ruby et al, 2018), bringing the total
number of animals analyzed to 6,957. For still-surviving animals, right censorship was
applied at the date of record aggregation: March 15, 2021.
488

489   **Analysis of lifespan and age-specific mortality hazard**
490

Except as otherwise specified in the text, right censorship status was applied to animal
lifespans in three cases: if the animal was (1) transferred to another collection, with the
date of transfer considered as the censorship event; (2) euthanized for research
purposes when still healthy, with the date of sacrifice as the censorship event; or (3) still
alive when the process of compiling records was completed, with the date of completion
as the censorship event. Left censorship was applied to entire data sets as specified in
the text, generally at specific historical dates in order to exclude data collected prior to
that date.
499

Kaplan-Meier survival was calculated using their method (Kaplan and Meier, 1958).
Mortality hazards were calculated for each specified age bin/group of animals as the
number of observed death events (i.e., number of days on which a naked mole-rat was
observed to have died) divided by the total number of days that naked mole-rats were

under observation, limited to that age bin/population. Confidence intervals were
calculated using the Wilson score interval with continuity correction (Newcombe, 1998).
506
507 **_H.glaber_ body weight analysis**
508
Historical records of body weights, collected in the course of animal health checks, were
aggregated concurrently with the lifespan data from (Ruby et al, 2018). These records
were less complete than the demographic data, but nonetheless included 19,118
measurements taken from 1,940 animals with complete lifespan data. For most
analyses, only weights taken after 500 years of age were used: there were 6,785 such
measurements, taken from 942 animals. The demographic data used for these
analyses was as compiled for (Ruby et al, 2018), and it is provided along with colony
membership and body weight data in Supplemental Data File 2. Animals from (Ruby et
al, 2018) with full demographic data but no body weight data are also included in that
file. Animal ID's in that file are matched to (Ruby et al, 2018) and Supplemental Data
File 1.
520
For mortality versus body weight, analysis was performed for survival across a single
year. For each animal with body weight(s) measured after 500 days of age, survival
data were analyzed starting from the date of the last recorded body weight (i.e. left-
censored on that date). If the animal survived uncensored for over a year, then right-
censorship was applied one year after the weight measurement. Animals with
qualifying body weight values were sorted into quartiles based on those final
measurements, and mortality hazard was estimated across all of the animals in each
quartile. Confidence intervals were calculated using the Wilson score interval with
continuity correction (Newcombe, 1998).
530
531 **_H.glaber_ colony size analysis**
532
For mortality versus colony size, analyses were performed of survival across a single
year. The results for each year are shown in Supplemental Figure S1: for years 2012-
2015, demographic records from (Ruby et al, 2018; also provided in Supplemental Data
File 2) were used, and April 14 was used as the cut-off date; for years 2016-2020, the
updated demographic records from this manuscript were used (Supplemental Data File
1), and March 15 was used as the cut-off date. Cut-off dates were selected to match
the data compilation date for each data set, ensuring a full year's worth of data
availability for each year's analysis. The shift to an older record compilation for earlier-
year analyses was to minimize the effect of movement out of colonies as a confounder:
colony affiliations for each compiled data set reflected colony membership at the time of
compilation and did not include prior membership history. Importantly, this caveat

should have had minimal impact: due to violence against intruders, animals are never
introduced into existing colonies, only removed from colonies in order to establish new
colonies as breeders (Smith and Buffenstein, 2021), and breeders were excluded from
these analyses.
548
For analysis of colony sizes, only animals at least 500 days old by each cut-off date
were considered. This decision was justified by animals younger than that age being
not fully grown and therefore potentially presenting less competitive rivalry; and by the
inconsistent integration of animals much younger than one year (pre-chipped) old into
our demography database. But it may have added variance in the form of colony-size
error in the cases of colonies with highly skewed age distributions. Analyses were also
conducted in a sex-specific manner, i.e., considering only the counts of same-sex, non-
breeding animals for colony size. This decision was justified under the assumption that
competition and hierarchy may be different between the two sexes (Reeve & Sherman,
1991), but may have added variance in cases where the sex ratio was skewed in a
colony, to the extent that social hierarchy among non-breeders is sex-independent.
560
For each cut-off date, for each sex, all qualifying animals were used to generate a
census (animal count) for each represented colony name. Individual animals were
sorted into quartiles based on the sizes of the colonies of which they were members;
tied animals were assigned to adjacent quartiles arbitrarily. For members of each
quartile, the mean and standard deviation of colony sizes for each animal were
calculated, and the mortality hazard for those animals was calculated across one year,
beginning at the cut-off date (i.e. left-censored on the cut-off date, and right censored
one year after the cut-off date for animals that were not deceased or otherwise
censored during the year). Confidence intervals were calculated using the Wilson score
interval with continuity correction (Newcombe, 1998).
571
For regression of mean colony size versus mortality hazard, for each sex: for all
quartiles across all years shown in Supplemental Figure S1, mean colony sizes were
treated as x-axis values and mortality hazard estimates as y-axis values, and linear
regression was performed using the "scipy.stats.linregress" function from Scipy
(Virtanen et al., 2020), which in addition to slope and intercept returns Pearson R and
regression p-value.
578
579   **_F.damarensis_ demographic data**
580
581   As for _H.glaber_, lifespan data from our longstanding collection of Damaraland mole-rats,
582   sp. _F.damarensis_, were compiled on March 15, 2021, with that date used as the right-
583   censorship point for still-living animals. Of 754 animals with records, 702 had day-

584 resolution birth and death/censorship data and were therefore included in these
585 analyses. Per-animal demographic data are provided in Supplemental Data File 3.
586
587 **Calculation of maximum versus observed expected lifespan**
588
589 Allometric expectations were determined using the equation from (de Magalhães et al.,
590 2007):
591
592 $T_{max} = 4.88 * m^{0.153}$
593
where body mass m is in grams and $T_{max,}$ predicted maximum lifespan, is in years. The
longevity quotient (LQ) was defined as the ratio of observed MLSP to T_{max}. For
H.glaber: using a body mass = 35g, T_{max} = 8.4 years; and with observed MLSP = 37
years, LQ = 4.4. For *F.anselli*: using a body mass = 90g, T_{max} = 9.7 years; and with
observed MLSP = 22 years, LQ = 2.3. For *F.damarensis*: using a body mass = 160g,
T_{max} = 10.6 years; and with observed MLSP = 20 years, LQ = 1.9.
600

601 <u>**Funding**</u>
602 This study was funded by Calico Life Sciences LLC.
603
604 <u>**Acknowledgements**</u>
605 We thank the numerous animal attendants who have diligently cared for the naked
606 mole-rats and Damaraland mole-rats over the past three-and-a-half decades.

607
<u>**Ethics**</u>
Animal experimentation: All of the animals were handled according to approved
institutional animal care and use committee (IACUC) protocols, most recently of the
Buck Institute (A10138). IACUC approval for housing these colonies of naked mole-rats
and Damaraland mole-rats were also obtained from all institutions at which these
collections were historically housed.
614
615 **Competing Interests**
616 The research was funded by Calico Life Sciences LLC, where all authors were
617 employees at the time the study was conducted. The authors declare no other
618 competing financial interests.
619

620 <u>**Supplemental Files**</u>
621 **Supplemental Figures:** two Supplemental Figures (S1 & S2), with legends, as one
622 PDF.

623
Supplemental Data File 1:
Full *H.glaber* lifespan data, compiled in 2021 (tab-separated text file).
Supplemental Data File 2:
Full *H.glaber* lifespan data, as compiled in 2016, with body weight measurements (tab-
separated text file).
Supplemental Data File 3:
Full *F.damarensis* lifespan data, compiled in 2021 (tab-separated text file).
Supplemental Data File 4:
Wherever practical: numerical values that are plotted across all Figures and
Supplemental Figures, organized by Figure (Excel notebook file).

<u>Figure Legends</u>

Fig 1: The mortality hazard of naked mole-rats failed to increase with age.
(A) Kaplan-Meier survival curve for naked mole-rats (green) after reaching
reproductive maturity (T_{sex}; 6 months from birth; 183 days; red). Plotted as in Fig.
1A of (Ruby et al., 2018), with a histogram of death and censorship (pink and
purple, respectively) in the inset, up until 5,306 days of age, at which point 174
animals remained in the population; and with censorship events after that
indicated by vertical ticks (purple), the size of which is proportional to the number
of animals censored at each day-of-life.
(B) Mortality-hazard estimates (solid green) with 95% confidence intervals (dotted
green) across the observed naked mole-rat lifespan. Estimates were calculated
across time intervals of increasing size (demarked by grey lines) to compensate
for decreasing accuracy-per-unit-time as the population size decreased. The
borders of the age bins (black), along with the number of deaths (pink) and total
years of animal life observed (green) per bin, are indicated at the bottom.
649
Fig 2: Exclusion of neither pre-2008 nor pre-2016 lifespan data through left-censorship
modified the observed lifespan demographics of *H.glaber*.
(A) The number of naked mole-rats observed (y-axis) at each age (x-axis) for four
datasets: i) the original data set presented in (Ruby et al., 2018) (pink); ii) that
data set left-censored on January 1, 2008, as presented in (Ruby et al., 2019)
(yellow); iii) the full data set compiled in 2021 for this manuscript (green); iv)
that data set left-censored on May 17, 2016 (navy), i.e. data collected after
the compilation of data for (Ruby et al., 2018).
(B) Kaplan–Meier survival curves starting at T_{sex} (183 days; red) for the four
demographic data sets, as colored and described in panel (A).
(C) Mortality hazard estimates (y-axis) across each of the lifespan bins from
Figure 1B (indicated on the x-axis), for each of the four demographic data

sets, as colored and described in panel (A). Horizontal bars indicate hazard
estimates, and vertical bars indicate 95% confidence intervals.

Fig. 3 Breeding and non-breeding male and female naked mole-rats exhibited non-
increasing mortal hazard as a function of age.

     667 (A) Kaplan-Meier survival starting at T_{sex} (183 days; red) for each of four
     668   reproductive categories of naked mole-rats: male breeders (navy), female
     669   breeders, (red), male non-breeders (cyan), and female non-breeders (pink).
     670 (B) Mortality hazard estimates (y-axis) for the indicated lifespan bins (x-axis),
     671   calculated for non-breeding males (cyan) and females (pink). The numbers of
     672   observed death events per bin are indicated at the top of each bin.
     673 (C) Mortality hazard estimates (y-axis) for the indicated lifespan bins (x-axis),
     674   calculated for breeding males (navy) and females (red). The numbers of
     675   observed death events per bin are indicated at the top of each bin.

677   **Fig. 4** Non-breeding naked mole-rat mortality hazard decreased with increasing body
678   weight and colony size.

     679 (A) Individual body weight values (y-axis) plotted longitudinally versus age (x-axis)
     680   for breeding males (navy), non-breeding males (cyan), breeding females (red),
     681   and non-breeding females (pink).
     682 (B) Mean body-weights for each of the categories from panel (A), for all weight
     683   measurements taken across each of the following age bins: 50-120 days, 120-
     684   200 days, 200-400 days, 400-700 days, 700-1000 days, 1000-1500 days, and
     685   1500-2000 days; inclusive of the 1st day and non-inclusive of the last day for
     686   each bin. Solid lines indicate mean body weights; dotted lines indicate standard
     687   deviations.
     688 (C) Top panel: body weight quartiles for male (left) and female (right) non-breeding
     689   animals, organized based on the last recorded weight measurement for each
     690   animal and only included if that measurement was taken at an older age than
     691   500 days. Horizontal bars indicate the mean body weight for each quartile;
     692   vertical bars indicate standard deviations. Bottom panel: mortality hazard for
     693   each body weight quartile in the top panel, calculated for the year following
     694   weight measurement. See Methods for details. Horizontal bars indicate mortality
     695   hazard estimates; vertical bars indicate 95% confidence intervals.
     696 (D) Examples of colony-size versus mortality-hazard plots. For each indicated year
     697   (2019 on the left; 2015 on the right), and for each indicated non-breeding sex
     698   (males on the left; females on the right): animals were organized into quartiles
     699   based on the number same-sex, non-breeding animals in their colony. Top
     700   panels: the mean colony sizes for animals in each quartile; vertical bars indicate
     701   standard deviations. Bottom panels: mortality hazard estimates for each colony-

size quartile from the top panel; vertical bars indicate 95% confidence intervals.
Similar analyses were performed for each year, 2012-2020; all plots are provided
in Supplemental Figure S1. See Methods for details.
(E) For all colony-size quartiles, from all years analyzed in Supplemental Figure S1:
a meta-analysis of quartile mean colony sizes (x-axis) versus one-year mortality
hazards (y-axis). Non-breeding males (cyan) and females (pink) were meta-
analyzed separately. Regression lines for each sex are shown (p-values: 9.8 *
10^{-4} for males; 0.018 for females).

Fig 5: The mortality hazard of *Fukomys* mole-rats increased with age.
(A) Kaplan-Meier survival curve for combined data from the *F.anselli and*
F.damarensis species of mole-rats (green) after reaching reproductive maturity
(T_{sex}; 9 months from birth; 270 days; red). Plotted as in Fig. 1A, with a histogram
of death and censorship (pink and purple, respectively) in the inset, up until 3,400
days of age, at which point 120 animals remained in the population; and with
censorship events after that indicated by vertical ticks (purple), the size of which
is proportional to the number of animals censored at each day-of-life. Equivalent
per-species analyses are provided in Supplemental Figure S2.
(B) Mortality-hazard estimates (solid green) with 95% confidence intervals (dotted
green) across the observed *Fukomys* mole-rat lifespan. Estimates were
calculated across time intervals of increasing size (demarked by grey lines) to
compensate for decreasing accuracy-per-unit-time as the population size
decreased. The borders of the age bins are indicated in gray, and the number of
deaths (pink) and total years of animal life observed (green) per bin are indicated
at the bottom of the panel. Equivalent per-species analyses are provided in
Supplemental Figure S2.
(D) For *F.damarensis* only: Kaplan-Meier survival starting at T_{sex} (270 days; red) for
each of four reproductive categories of Damaraland mole-rats: male breeders
(navy), female breeders, (red), male non-breeders (cyan), and female non-
breeders (pink).

Fig. 6: In contrast to the mortality hazards of other mammals, which increased with
chronological age, the mortality hazard of naked mole-rats remained constant.
(A) Age-specific mortality hazard for humans (navy), mice (cyan), horses (orange),
Fukomys mole-rats (purple), and naked mole-rats (green); with age (x-axis)
normalized to T_{sex} for each species. Reproduced from (Ruby et al, 2018), with
Fukomys data added and *H.glaber* data updated using results from this study.
See Results for references to original data sources.
(B) A phylogenic tree for the species from panel (A). See Results for references to
data sources.

References

744
745 Anderson, S. E., Dallal, G. E., & Must, A. (2003). Relative weight and race influence average age at
746 menarche: results from two nationally representative surveys of US girls studied 25 years
747 apart. *Pediatrics*, *111*(4), 844-850.

748
749 Apter, D. (1980). Serum steroids and pituitary hormones in female puberty: a partly longitudinal
750 study. *Clinical endocrinology*, *12*(2), 107-120.

751
752 Bell, F. C., & Miller, M. L. (2005). *Life tables for the United States social security area, 1900-2100* (No.
753 120). Social Security Administration, Office of the Chief Actuary.

754
755 Berger, V., Lemaître, J. F., Allainé, D., Gaillard, J. M., & Cohas, A. (2018). Early and adult social
756 environments shape sex-specific actuarial senescence patterns in a cooperative breeder. *The American*
757 *Naturalist*, *192*(4), 525-536.

758
759 Brett, R. in *The Biology of the Naked Mole-Rat* (eds Sherman, P. W., Jarvis, J. U. M. & Alexander, R. D.)
760 97–136 (Princeton Univ. Press, NJ, 1991).

761
762 Broom, M., Koenig, A., & Borries, C. (2009). Variation in dominance hierarchies among group-living
763 animals: modeling stability and the likelihood of coalitions. *Behavioral Ecology*, *20*(4), 844-855.

764
765 Buffenstein, R., & Jarvis, J. U. (2002). The naked mole rat--a new record for the oldest living
766 rodent. *Science of Aging Knowledge Environment*, *2002*(21), pe7-pe7.

767
768 Buffenstein, R. (2005). The naked mole-rat: a new long-living model for human aging research. *The*
769 *Journals of Gerontology Series A: Biological Sciences and Medical Sciences*, *60*(11), 1369-1377.

770
771 Buffenstein, R., Lewis, K. N., Gibney, P. A., Narayan, V., Grimes, K. M., Smith, M., ... & Brown-Borg, H.
772 M. (2020). Probing pedomorphy and prolonged lifespan in naked mole-rats and dwarf
773 mice. *Physiology*, *35*(2), 96-111.

774
775 Buffenstein, R., & Craft, W. (2021). The idiosyncratic physiological traits of the naked mole-rat; a resilient
776 animal model of aging, longevity, and healthspan. *The extraordinary biology of the naked mole-rat*, 221-
777 254.

778
779 Buffenstein, R., Park, T. J., & Holmes, M. M. (Eds.). (2021). *The Extraordinary Biology of the Naked Mole-*
780 *Rat*. Cham: Springer.

781
Buffenstein, R., Amoroso, V., Andziak, B., Avdieiev, S., Azpurua, J., Barker, A. J., ... & Smith, E. S. J.
(2022). The naked truth: a comprehensive clarification and classification of current 'myths' in naked mole-
rat biology. *Biological Reviews*, *97*(1), 115-140.

Can, E., Smith, M., Boukens, B. J., Coronel, R., Buffenstein, R., & Riegler, J. (2022). Naked mole-rats
maintain cardiac function and body composition well into their fourth decade of life. *GeroScience*, *44*(2),
731-746.
789
790   Carey, J. R. (2001). Demographic mechanisms for the evolution of long life in social insects. *Experimental*
791   *gerontology*, *36*(4-6), 713-722.
792
793   Clarke, F. M., & Faulkes, C. G. (1997). Dominance and queen succession in captive colonies of the
794   eusocial naked mole–rat, Heterocephalus glaber. *Proceedings of the Royal Society of London. Series B:*
795   *biological sciences*, *264*(1384), 993-1000.
796
797   Clarke, F. M., & Faulkes, C. G. (2001). Intracolony aggression in the eusocial naked mole-rat,
798   Heterocephalus glaber. *Animal Behaviour*, *61*(2), 311-324.
799
800   Clutton-Brock, T. (2021). Social evolution in mammals. *Science*, *373*(6561), eabc9699.
801
802   Coen, C. W., Bennett, N. C., Holmes, M. M., & Faulkes, C. G. (2021). Neuropeptidergic and
803   neuroendocrine systems underlying eusociality and the concomitant social regulation of reproduction in
804   naked mole-rats: a comparative approach. *The extraordinary biology of the naked mole-rat*, 59-103.
805
806   Cohen, A. A. (2018). Aging across the tree of life: The importance of a comparative perspective for the
807   use of animal models in aging. *Biochimica et Biophysica Acta (BBA)-Molecular Basis of Disease*, *1864*(9),
808   2680-2689.
809
810   Cram, D. L., Monaghan, P., Gillespie, R., Dantzer, B., Duncan, C., Spence-Jones, H., & Clutton-Brock, T.
811   (2018). Rank-related contrasts in longevity arise from extra-group excursions not delayed senescence in
812   a cooperative mammal. *Current Biology*, *28*(18), 2934-2939.
813
814   Dammann, P., Šumbera, R., Maßmann, C., Scherag, A., & Burda, H. (2011). Extended longevity of
815   reproductives appears to be common in Fukomys mole-rats (Rodentia, Bathyergidae). *PLoS One*, *6*(4),
816   e18757.
817
818   Dammann, P., Scherag, A., Zak, N., Szafranski, K., Holtze, S., Begall, S., ... & Platzer, M. (2019).
819   Comment on'Naked mole-rat mortality rates defy Gompertzian laws by not increasing with age'. *Elife*, *8*,
820   e45415.
821
822   Edrey, Y. H., Hanes, M., Pinto, M., Mele, J., & Buffenstein, R. (2011). Successful aging and sustained
823   good health in the naked mole rat: a long-lived mammalian model for biogerontology and biomedical
824   research. *ILAR journal*, *52*(1), 41-53.
825
826   Fang, X., Seim, I., Huang, Z., Gerashchenko, M. V., Xiong, Z., Turanov, A. A., ... & Gladyshev, V. N.
827   (2014). Adaptations to a subterranean environment and longevity revealed by the analysis of mole rat
828   genomes. *Cell reports*, *8*(5), 1354-1364.
829
830   Faulkes, C. G., & Bennett, N. C. (2021). Social evolution in African mole-rats–a comparative
831   overview. *The Extraordinary Biology of the Naked Mole-Rat*, 1-33.
832
833   Gaillard, J. M., & Lemaître, J. F. (2017). The Williams' legacy: A critical reappraisal of his nine predictions
834   about the evolution of senescence. *Evolution*, *71*(12), 2768-2785.

836 Gavrilova, N. S., & Gavrilov, L. A. (2015). Biodemography of old-age mortality in humans and
837 rodents. *Journals of Gerontology Series A: Biomedical Sciences and Medical Sciences*, *70*(1), 1-9.
838

839 Gompertz B (1825) On the nature of the function expressive of the law of human mortality, and on a new
840 mode of determining the value of life contingencies *Philosophical Transactions of the Royal Society of*
841 *London* 115:513–583.
842

843 Grimes, K. M., Barefield, D. Y., Kumar, M., McNamara, J. W., Weintraub, S. T., de Tombe, P. P., ... &
844 Buffenstein, R. (2017). The naked mole-rat exhibits an unusual cardiac myofilament protein profile
845 providing new insights into heart function of this naturally subterranean rodent. *Pflügers Archiv-European*
846 *Journal of Physiology*, *469*, 1603-1613.
847

848 Hadi, F., Smith, E. S. J., & Khaled, W. T. (2021). Naked mole-rats: resistant to developing cancer or good
849 at avoiding it?. *The extraordinary biology of the naked mole-rat*, 341-352.
850

851 Hamilton, W. D. (1966). The moulding of senescence by natural selection. *Journal of theoretical*
852 *biology*, *12*(1), 12-45.
853

854 Healy, K., Guillerme, T., Finlay, S., Kane, A., Kelly, S. B., McClean, D., ... & Cooper, N. (2014). Ecology
855 and mode-of-life explain lifespan variation in birds and mammals. *Proceedings of the Royal Society B:*
856 *Biological Sciences*, *281*(1784), 20140298.
857

858 Holmes, M. M., & Goldman, B. D. (2021). Social behavior in naked mole-rats: individual differences in
859 phenotype and proximate mechanisms of mammalian eusociality. *The extraordinary biology of the naked*
860 *mole-rat*, 35-58.
861

862 Jacob, F. (1977). Evolution and tinkering. *Science*, *196*(4295), 1161-1166.
863

864 Jarvis, J. U. M. (1981). Eusociality in a mammal: cooperative breeding in naked mole-rat
865 colonies. *Science*, *212*(4494), 571-573.
866

867 Jarvis, J. U. M., & Bennett, N. C. (1993). Eusociality has evolved independently in two genera of
868 bathyergid mole-rats—but occurs in no other subterranean mammal. *Behavioral Ecology and*
869 *Sociobiology*, *33*, 253-260.
870

871 Jarvis, J. U., Bennett, N. C., & Spinks, A. C. (1998). Food availability and foraging by wild colonies of
872 Damaraland mole-rats (Cryptomys damarensis): implications for sociality. *Oecologia*, 290-298.
873

874 Kaplan, E. L., & Meier, P. (1958). Nonparametric estimation from incomplete observations. *Journal of the*
875 *American Statistical Association*, 53, 457–481.
876

877 Kerepesi, C., Meer, M. V., Ablaeva, J., Amoroso, V. G., Lee, S. G., Zhang, B., ... & Gladyshev, V. N.
878 (2022). Epigenetic aging of the demographically non-aging naked mole-rat. *Nature*
879 *communications*, *13*(1), 355.
880

Kim, E. B., Fang, X., Fushan, A. A., Huang, Z., Lobanov, A. V., Han, L., ... & Gladyshev, V. N. (2011).
Genome sequencing reveals insights into physiology and longevity of the naked mole
rat. *Nature*, *479*(7372), 223-227.

Kirkwood, T. B. (1977). Evolution of ageing. *Nature, 270*(5635), 301-304.

Lee, B. P., Smith, M., Buffenstein, R., & Harries, L. W. (2020). Negligible senescence in naked mole rats
may be a consequence of well-maintained splicing regulation. *GeroScience, 42*(2), 633-651.

Lewis, K. N., & Buffenstein, R. (2016). The naked mole-rat: a resilient rodent model of aging, longevity,
and healthspan. In *Handbook of the Biology of Aging* (pp. 179-204). Academic press.

Lewis, K. N., Rubinstein, N. D., & Buffenstein, R. (2018). A window into extreme longevity; the circulating
metabolomic signature of the naked mole-rat, a mammal that shows negligible
senescence. *Geroscience, 40*(2), 105-121.

Lucas, E. R., & Keller, L. (2020). The co-evolution of longevity and social life. *Functional Ecology, 34*(1),
76-87.

de Magalhães, J. P. D., Costa, J., & Church, G. M. (2007). An analysis of the relationship between
metabolism, developmental schedules, and longevity using phylogenetic independent contrasts. *The
Journals of Gerontology Series A: Biological Sciences and Medical Sciences, 62*(2), 149-160.

Medger, K., Bennett, N. C., Ganswindt, S. B., Ganswindt, A., & Hart, D. W. (2019). Changes in prolactin,
cortisol and testosterone concentrations during queen succession in a colony of naked mole-rats
(Heterocephalus glaber): a case study. *The Science of Nature, 106*, 1-7.

Miller, R. A., Harrison, D. E., Astle, C. M., Fernandez, E., Flurkey, K., Han, M., ... & Strong, R. (2014).
Rapamycin-mediated lifespan increase in mice is dose and sex dependent and metabolically distinct from
dietary restriction. *Aging cell, 13*(3), 468-477.

Newcombe, R. G. (1998). Two-sided confidence intervals for the single proportion: comparison of seven
methods. *Statistics in medicine, 17*(8), 857-872.

O'Connor, T. P., Lee, A., Jarvis, J. U., & Buffenstein, R. (2002). Prolonged longevity in naked mole-rats:
age-related changes in metabolism, body composition and gastrointestinal function. *Comparative
Biochemistry and Physiology Part A: Molecular & Integrative Physiology, 133*(3), 835-842.

Orr, M. E., Garbarino, V. R., Salinas, A., & Buffenstein, R. (2016). Extended postnatal brain development
in the longest-lived rodent: prolonged maintenance of neotenous traits in the naked mole-rat
brain. *Frontiers in neuroscience, 10*, 504.

Patterson, B. D., & Upham, N. S. (2014). A newly recognized family from the Horn of Africa, the
Heterocephalidae (Rodentia: Ctenohystrica). *Zoological Journal of the Linnean Society, 172*(4), 942-963.

Prothero, J. (1993). Adult life span as a function of age at maturity. *Experimental gerontology, 28*(6), 529-
536.

Reeve, H. K., & Sherman, P. W. (1991). Intracolonial aggression and nepotism by the breeding female
naked mole-rat. *The biology of the naked mole-rat*, 337-357.

Rose, M. R., Rauser, C. L., Benford, G., Matos, M., & Mueller, L. D. (2007). Hamilton's forces of natural
selection after forty years. *Evolution*, *61*(6), 1265-1276.
934
935   Ross, C., Rychtář, J., & Rueppell, O. (2015). A structured population model suggests that long life and
936   post-reproductive lifespan promote the evolution of cooperation. *Journal of Theoretical Biology*, *369*, 85-
937   94.
938
939   Ruby, J. G., Smith, M., & Buffenstein, R. (2018). Naked mole-rat mortality rates defy Gompertzian laws by
940   not increasing with age. *elife*, *7*, e31157.
941
942   Ruby, J. G., Smith, M., & Buffenstein, R. (2019). Response to comment on'Naked mole-rat mortality rates
943   defy Gompertzian laws by not increasing with age'. *Elife*, *8*, e47047.
944
945   Sahm, A., Platzer, M., Koch, P., Henning, Y., Bens, M., Groth, M., ... & Dammann, P. (2021). Increased
946   longevity due to sexual activity in mole-rats is associated with transcriptional changes in the HPA stress
947   axis. *Elife*, *10*, e57843.
948
949   Sharp, S. P., & Clutton-Brock, T. H. (2010). Reproductive senescence in a cooperatively breeding
950   mammal. *Journal of Animal Ecology*, *79*(1), 176-183.
951
952   Silk, J. B., Beehner, J. C., Bergman, T. J., Crockford, C., Engh, A. L., Moscovice, L. R., ... & Cheney, D.
953   L. (2010). Strong and consistent social bonds enhance the longevity of female baboons. *Current
954   biology*, *20*(15), 1359-1361.
955
956   Skulachev, V. P., Holtze, S., Vyssokikh, M. Y., Bakeeva, L. E., Skulachev, M. V., Markov, A. V., ... &
957   Sadovnichii, V. A. (2017). Neoteny, prolongation of youth: from naked mole rats to "naked
958   apes"(humans). *Physiological reviews*.
959
960   Smith, M., & Buffenstein, R. (2021). Managed care of naked mole-rats. *The extraordinary biology of the
961   naked mole-rat*, 381-407.
962
963   Springer, M. S., Murphy, W. J., Eizirik, E., & O'Brien, S. J. (2003). Placental mammal diversification and
964   the Cretaceous–Tertiary boundary. *Proceedings of the National Academy of Sciences*, *100*(3), 1056-
965   1061.
966
967   Tacutu, R., Thornton, D., Johnson, E., Budovsky, A., Barardo, D., Craig, T., ... & De Magalhães, J. P.
968   (2018). Human ageing genomic resources: new and updated databases. *Nucleic acids research*, *46*(D1),
969   D1083-D1090.
970
971   Toor, I., Edwards, P. D., Kaka, N., Whitney, R., Ziolkowski, J., Monks, D. A., & Holmes, M. M. (2020).
972   Aggression and motivation to disperse in eusocial naked mole-rats, Heterocephalus glaber. *Animal
973   Behaviour*, *168*, 45-58.
974
975   Valgren, V. N. (1945). Livestock insurance and mortality data. *Journal of American Insurance*, 16-22.
976
Van Daele, P. A., Verheyen, E., Brunain, M., & Adriaens, D. (2007). Cytochrome b sequence analysis
reveals differential molecular evolution in African mole-rats of the chromosomally hyperdiverse genus
Fukomys (Bathyergidae, Rodentia) from the Zambezian region. *Molecular phylogenetics and
evolution*, *45*(1), 142-157.

Virtanen, P., Gommers, R., Oliphant, T. E., Haberland, M., Reddy, T., Cournapeau, D., ... & Van
Mulbregt, P. (2020). SciPy 1.0: fundamental algorithms for scientific computing in Python. *Nature*
methods, *17*(3), 261-272.

Visser, J. H., Bennett, N. C., & van Vuuren, B. J. (2019). Phylogeny and biogeography of the African
Bathyergidae: a review of patterns and processes. *PeerJ*, *7*, e7730.