1

Comparative Analysis of Tylosema esculentum Mitochondrial DNA Revealed Two Distinct Genome Structures

Comparative Analysis of Marama Mitogenome Structures

Jin Li, Christopher Cullis

Department of Biology, Case Western Reserve University, Cleveland, OH, United States

Corresponding Author:

Christopher Cullis

cac5@case.edu

Orchid id: 0000-0002-9768-7763

Contact for Jin Li: jxl1269@case.edu

Number of figures: 15

Number of tables: 7

Word count: 5036

Supplementary materials: 3 tables and 32 figures

Comparative Analysis of *Tylosema esculentum* Mitochondrial DNA Revealed Two Distinct Genome Structures

Highlight

The analysis of 84 marama mitogenomes revealed two germplasms and the structural variation affects certain gene copy numbers. Soil moisture levels may have played important roles in the mitogenome divergence.

Abstract

Tylosema esculentum (marama bean), an underutilized legume with edible and nutritious seeds, has the potential to improve local food security in southern Africa. This study investigated the diversity of marama mitogenomes by mapping sequencing data from 84 individuals to the previously published reference mitogenome. Two distinct germplasms were found, and a new mitogenome structure containing three circular molecules and one long linear chromosome was identified, with a unique 2,108 bp fragment and primers were designed on that for marama mitogenome typing. This structural variation increases copy number of certain genes, including *nad9*, *rrns* and *rrn5*. The two mitogenomes also differed at 230 loci, with only one nonsynonymous substitution in *matR*. The evolutionary analysis suggested that the divergence of marama mitogenomes may be related to soil moisture level. Heteroplasmy in the marama mitogenome was concentrated at specific loci, including 127,684 bp to 127,686 bp on chromosome LS1 (OK638188), and may be crucial in the evolution. Additionally, the mitogenomes of marama contained a cpDNA insertion of over 9 kb with numerous polymorphisms, resulting in the loss of function of the genes on that segment. This comprehensive analysis of marama mitogenome diversity may provide valuable insight for future improvement of the bean.

Keywords: Legume mitogenome, mitogenome assembly, PacBio HiFi sequencing, mitogenome diversity, genome structural variation, heteroplasmy, horizontal gene transfer, phylogeny

Introduction

Tylosema esculentum, also known as marama bean and gemsbok bean, is a non-nodulated legume from the Fabaceae family, Cercidoideae subfamily (LPWG, 2017; Sinou et al., 2020). Marama is endemic to the Kalahari Desert and the surrounding arid and semi-arid regions of Botswana, Namibia, and South Africa (Bower et al., 1988). Marama has developed a special drought avoidance strategy by growing giant tubers that can weigh more than 500 pounds to store water, helping them survive the harsh conditions of long-term drought and little rainfall (Cullis et al., 2018). The seeds of marama are edible and nutritious, with a protein content of approximately 30-39% dry matter (dm) and a lipid content of 35-48% dm, comparable to those found in commercial crops soybean and peanuts, respectively (Amarteifio, 1998; Belitz et al., 2004; Holse et al., 2010). Marama also contains many micronutrients and phytochemicals that are beneficial to human health (Jackson et al., 2010; Khan, 2018; Omotayo and Aremu, 2021). However, marama is still mainly collected from wild plants, and the domestication of marama

has long been thought to have the potential to improve local food security. Marama usually does
not flower and produce seeds until two to four years after planting, making traditional breeding
very inefficient (Cullis et al., 2019). Therefore, selecting improved marama individuals and
exploring the underlying genetic diversity is considered of great significance for the
improvement of the bean (Cullis et al., 2022).

Mitochondria in eukaryotic cells are generally considered to have originated from the
endosymbiosis of alpha-proteobacteria, although a number of changes have occurred since then,
including the loss of many genes and their transfer to the nuclear genome (Andersson et al.,
2002). The mitochondrial genomes of animals and plants have been found to vary greatly.
Animal mitogenomes are usually small, only about 16 kb in size, and contain 37 genes (Boore,
1999). Plant mitogenomes are commonly larger, with 50 to 60 genes and expanded intergenic
non-coding regions that result from DNA transfer from other cellular compartments or even from
different organisms (Unseld et al., 1997; Kazuyoshi and Kubo, 2010; Gualberto et al., 2014). In
land plants, they range in size from 66 kb in *Viscum scurruloideum* to 11.3 Mb in *Silene conica*
(Sloan et al., 2012; Skippington et al., 2015).

Plant mitochondrial genes are very conserved, are considered to play important roles in ATP
synthesis, and are also related to plant fertility and environmental adaptation (Hanson, 1990;
Budar and Roux, 2011; Heng et al., 2014). The base substitution rate of mitochondrial genes is
lower than that of chloroplast genes, and it is far lower, only about one-tenth, of the rate
observed in nuclear genes (Wolfe et al., 1987; Drouin et al., 2008). Plant mitogenomes evolve
even up to 100 times slower than animal mitogenomes (Palmer and Herbon; 1988). However, the
structure of plant mitochondrial genomes is very dynamic, with a large number of sequence
rearrangements, and repeat-mediated homologous recombination plays an important role in its
structural variations (Gualberto and Newton, 2017; Cole et al., 2018).

In the mitochondrial genomes of many angiosperms, repetitive sequences account for 5-10% of
the total genome size, and in a few plants such as *S. conica* and *Nymphaea colorata*, the
proportion can exceed 40% with repetitive fragments up to 80 kb in length (Sloan et al., 2012;
Dong et al., 2018). Recombination mediated by short or intermediate length repeats is considered
to be less frequent, but recombination on long repeats (>1 kb) is thought to occur more
frequently and usually generates equimolar recombined molecules in the plant mitogenomes
(Arrieta-Montiel, 2009; Guo et al., 2016; Li and Cullis, 2021).

The third-generation sequencing technology, such as PacBio, provides long reads with an
average length of 10-25 kb, spanning the long repeats in the plant mitogenome. This makes the
study of structural variations caused by the long repeat mediated recombination possible (Kozik
et al., 2019; Hon et al., 2020). High sequencing coverage is also important, not only to make the
genome assembly more reliable, but also to determine the proportion of different chromosome
structures. It is also indispensable for the accurate assessment of nucleotide polymorphisms
(Telenti et al., 2016). The latest PacBio HiFi sequencing with an extremely high accuracy of
99.9%, which needs little correction by the data generated from other sequencing platforms,
further promotes the genome assembly (Miga et al., 2020; Naish et al., 2021).

Although plant mitochondrial genomes are often reported as one master circular chromosome, in
reality, they often exist as multipartite structures. This includes a combination of linear
molecules, branched molecules, and subgenomic circular molecules (Oldenburg and Bendich,

1996; Manchekar et al., 2006). For example, the mitogenome assembly of *Solanum tuberosum*
was found to contain at least three autonomous chromosomes, including two small circular
molecules and a long linear chromosome of 312,491 bp in length (Varré et al., 2019).

In the previously published study on the mitogenome of *T. esculentum*, two different equimolar
structures were found to coexist in the same individual. There are two autonomous rings with a
total length of 399,572 bp and five smaller circular molecules (Li and Cullis, 2021). These two
structures are believed to be interchangeable by recombination on three pairs of long direct
repeats (3-5 kb in length). As described in the study of *Brassica campestris* mitogenome,
recombination on a pair of 2 kb repeats was postulated to split the 218 kb master chromosome
into two subgenomic circular molecules of 135 kb and 83 kb in length (Palmer and Shields,
1984).

The previous comparative analysis of 84 *T. esculentum* chloroplast genomes has found two
distinct germplasms. The two types of chloroplast genomes are different from each other at 122
loci and at a 230 bp inversion (Li and Cullis, 2023). Among many of these loci, heteroplasmy,
the existence of two or more different alleles, could be seen, albeit one generally at a frequency
below 2%. The occasional paternal leakage is considered to cause this phenomenon (Kvist et al.,
2003; Luo et al., 2018). The reason for its stable inheritance is thought to be related to the
developmental genetic bottleneck, but the specific mechanism is still unclear (Floros et al., 2018).

Research on heteroplasmy should avoid interference from homologous sequences of
mitochondrial DNA in other organelles. In fact, the horizontal gene transfer of DNA from
chloroplast genome to mitogenome and nuclear genome, or between mitogenome and nuclear
genome are very common (Woloszynska et al., 2004; Keeling and Palmer, 2008). The transfer of
DNA from mitogenome to chloroplast genome is very rare, but it has been reported in a few
studies (Goremykin et al., 2008; Straub et al., 2013). Many chloroplast genes have been found to
become pseudogenes and lost function after being inserted into the mitogenome, but the reason
behind it is still unclear (Li et al., 2022).

Although the mechanisms underlying the effects of cytoplasmic activities on agronomic traits are
not well understood, previous studies on potato cytoplasmic diversity have found that
cytoplasmic types are directly related to traits such as tuber yield, tuber starch content, disease
resistance, and cytoplasmic male sterility (Sanetomo and Gebhardt, 2015; Anisimova and
Gavrilenko, 2017; Smyda-Dajmund et al., 2020). Furthermore, certain mtDNA types and certain
chloroplast DNA types were found to be linked (Lössl et al., 1999). A comparative genomics
analysis based on the chloroplast genomes of 3,018 modern domesticated rice cultivars found
that their genotypes fall into two distinct clades, suggesting that the domestication of these
cultivars may have followed two distinct evolutionary paths (Moner et al., 2020). These studies
have the potential to reveal important selections occurring in organelle genomes, improve
understanding of plant adaptation to different environments, and provide a basis for crop
breeding to increase yield in corresponding environments.

The chloroplast genome is widely used in research on plant evolution, but the comparative
analysis based on plant mitochondrial genome is not so extensive, and there is even less research
on the mitochondrial diversity within the same species (Nikiforova et al., 2013; Carbonell-
Caballero et al., 2015). In this study, the diversity of marama mitochondrial genome was
analyzed by mapping the WGS reads of 84 *T. esculentum* individuals to the previously
assembled reference mitogenome aiming to: (1) discover possible mitogenome structural

diversity and the impact of structural variations on gene sequence and copy numbers; (2) compare the differential loci in mitogenomes of 43 independent marama individuals collected from different geographical locations in Namibia and South Africa to explore the divergences that have occurred and possible decisive environmental factors behind them; (3) look at heteroplasmy, the co-existence of multiple types of mitogenomes within the same individuals, and compare allele frequencies in related individuals to better understand the underlying cytoplasmic inheritance; (4) track polymorphisms accumulated in the chloroplast DNA insertion and interpret the fate of the inserted gene residues. In addition, the conserved protein-coding genes from the mitogenomes of *T. esculentum* and other Fabaceae species were compared to explore the evolutionary relationship between them.

Materials and Methods

Plant materials and DNA extraction

Samples 4 and 32 were two individuals in the greenhouse of Case Western Reserve University grown from seeds collected in Namibia at undocumented locations. They were identified by PCR amplification to have type 2 and type 1 germplasms, respectively. 1 g fresh young leaves were collected from the two plants and ground thoroughly with a pestle in a mortar containing liquid nitrogen. DNA was then extracted using a Quick-DNA HMW MagBead kit (Zymo Research) following the protocol. The double-stranded DNA was quantified by the Invitrogen^{TM} Qubit^{TM} 3.0 Fluorometer after mixing 5 μl DNA with 195 μl working solution, and 200 ng DNA was electrophoresed on a 1.5% agarose TBE gel at 40 V for 24 hours. The plant materials included another 84 marama individuals, 44 of which were plants grown in different geographical locations in Namibia and South Africa, and the remaining 40 were progeny plants grown from seeds collected there, as described in the previous study (Table S1) (Li and Cullis, 2023). The WGS Illumina reads of these 84 individuals are available and stored in the NCBI SRA database (PRJNA779273).

High-throughput sequencing

DNA extracted from Samples 4 and 32 (10 micrograms or more per sample) was sent to the Genomics Core Facility at the Icahn School of Medicine at Mount Sinai for sequencing. The HiFi sequencing libraries were prepared using the SMRTbell® express template prep kit 2.0 (Pacific Biosciences, Menlo Park, CA, USA). SMRT sequencing was performed on four 8M SMRT® Cells (two per sample) on the Sequel® II system. 2,184,632 PacBio HiFi reads with a total length of 21.6 G bases were generated for Sample 4 and 498 Mb reads for Sample 32. In addition, whole-genome sequencing of DNA from fresh young leaves or embryonic axis of germinating seeds from the 84 samples collected in Namibia and South Africa was performed using the Illumina platform, as described in the previous study (Li and Cullis, 2023).

Mitogenome assembly and annotation

The PacBio HiFi long reads were assembled using the HiCanu assembler (Nurk et al., 2020; https://canu.readthedocs.io/en/latest/quick-start.html#assembling-pacbio-hifi-with-hicanu). The input genome size was set to 2 Mb to obtain more complete organelle genome contigs. The PacBio long reads spanning the ends of the contigs were used to further scaffold the assembly. The assembled mitogenome was annotated using MITOFY (Alverson et al., 2010; https://dogma.ccbb.utexas.edu/mitofy/), BLAST (Johnson et al., 2008;

https://blast.ncbi.nlm.nih.gov/Blast.cgi), and tRNAscan-se 2.0 (Chan et al., 2021;
http://lowelab.ucsc.edu/tRNAscan-SE/). The assembly was verified by mapping Illumina reads
and PacBio HiFi reads from different individuals to the generated mitogenome sequences using
Bowtie 2 v2.4.4 (Langmead and Salzberg, 2012; https://github.com/BenLangmead/bowtie2) and
pbmm2 v1.10.0 (https://github.com/PacificBiosciences/pbmm2), respectively. The alignment
was visualized and checked in IGV (Robinson et al., 2011;
https://software.broadinstitute.org/software/igv/), showing no ambiguity. The mitochondrial gene
arrangement maps of *T. esculentum* were drawn by OGDRAW (Greiner et al., 2019;
https://chlorobox.mpimp-golm.mpg.de/OGDraw.html).

Mitogenome polymorphism

The program NUCmer in MUMmer4 (Marçais et al., 2018;
https://mummer4.github.io/index.html) was used to locate highly conserved regions in the two
types of mitochondrial genomes of *T. esculentum*. The alignment was visualized via a synteny
plot drawn by the RIdeogram (Hao et al., 2020;
https://github.com/TickingClock1992/RIdeogram) package in R.

The 2,108 bp type 2 unique fragment was blasted in the NCBI database for potential origin. Two
pairs of primers were designed for mitogenome typing, using NCBI Primer-BLAST (Ye et al.,
2012; https://www.ncbi.nlm.nih.gov/tools/primer-blast/), to amplify across the two ends of this
fragment by PCR. The primers to amplify across the left end included the left forward primer
(GAGACCGAGCGCAAGAACTA) and the left reverse primer
(TCAGATGGCTAAACAGGCGG), and the product size was 990 bp. The primers to amplify
across the right end included the right forward primer (CGCTCGTGACTCATTGAGGA) and
the right reverse primer (TTGGTAAGCGGATGCTCTGG), and the product size was 289 bp.

20 uL of mixtures were prepared by separately mixing DNA from six randomly selected *T.*
esculentum samples and Promega GoTaq Green Master Mix. The amplifications started with
denaturation at 95 ° C for 5 min, followed by 30 cycles of 95 ° C for 45 s, 54 ° C for 45 s, and
72 ° C for 1 min, and a final 72 ° C for 5 min.

The Illumina reads from the 84 individuals were mapped to the type 1 reference mitogenome of
T. esculentum (OK638188 and OK638189) using Bowtie 2 v2.4.4. The alignments were searched
for SNPs and indels using SAMtools 1.7 mpileup (Li, 2011; http://www.htslib.org/) and
BCFtools 1.8 call (Li, 2011; http://www.htslib.org/doc/1.8/bcftools.html), and visualized in IGV
to find heteroplasmy manually. To minimize the interference of sequencing errors and strand
bias, only alleles with a frequency of at least 2%, a Phred score above 20, and presence in strands
in both orientations were recorded. Alleles in the mtDNA homologous reads in the nuclear
genome were excluded. The alleles from the mitochondrial or chloroplast genomes were
distinguished by their frequency at the differential loci on the 9,798 bp chloroplast DNA
insertion.

Mitogenome divergence

A pairwise comparison was performed on the divergent mitogenomes of *T. esculentum*
(OK638188 and OK638189) and six other Fabaceae species, including *Cercis canadensis*
(MN017226.1), *Lotus japonicus* (NC_016743.2), *Medicago sativa* (ON782580.1), *Millettia*
pinnata (NC_016742.1), *Glycine max* (NC_020455.1), and *Vigna radiata* (NC_015121.1) using

the program PROmer (Delcher et al., 2002) in MUMmer4 to detect the syntenic regions. The
alignments were then visualized by the RIdeogram package in R. A synteny block diagram
between these seven mitogenomes was drawn by Mauve 2.4.0 (Darling et al., 2004;
https://darlinglab.org/mauve/mauve.html) and the genes contained in each block were marked on
the plot.

SSR analyses

244 Microsatellites were analyzed by MISA (Beier et al., 2017; https://webblast.ipk-
245 gatersleben.de/misa) on the type 1 reference mitogenome of *T. esculentum*, looking for repeats of
246 motifs one to six base pairs long. The minimum number of repetitions were set to 10, 6, 5, 5, 5, 5
247 for mono-, di-, tri-, tetra-, penta-, and hexanucleotide repeats, respectively.

Phylogenetic tree construction

Two phylogenetic trees were constructed separately, one based on the mitogenome conserved
gene sequences to explore the evolutionary relationship between *T. esculentum* and several other
selected legumes, and the other tree was built on all differential loci found in the mitogenomes of
T. esculentum to explore the inter-population and intra-population relationship among the 43
independent samples.

The 24 conserved mitochondrial genes, *atp1*, *atp4*, *atp6*, *atp8*, *atp9*, *nad3*, *nad4*, *nad4L*, *nad6*,
nad7, *nad9*, *mttB*, *matR*, *cox1*, *cox3*, *cob*, *ccmFn*, *ccmFc*, *ccmC*, *ccmB*, *rps3*, *rps4*, *rps12*, and
rpl16 from the mitogenomes of *T. esculentum* (OK638188 and OK638189), *Arabidopsis*
thaliana (NC_037304.1), *C. canadensis* (MN017226.1), *L. japonicus* (NC_016743.2), *M. sativa*
(ON782580.1), *M. pinnata* (NC_016742.1), *G. max* (NC_020455.1), and *V. radiata*
(NC_015121.1) were concatenated to make artificial chromosomes. A Maximum Likelihood
(ML) phylogenetic tree using the Jukes-Cantor model was built in Mega 11 (Tamura et al., 2021;
https://www.megasoftware.net/) after the chromosomes were aligned by Muscle v5 (Edgar, 2022;
https://www.drive5.com/muscle/). The topology was validated by a Bayesian inference
phylogenetic tree drawn by BEAST v1.10.4 (Suchard et al., 2017; https://beast.community/) and
FigTree v1.4.4 (http://tree.bio.ed.ac.uk/software/figtree/).

Artificial chromosomes concatenated by 40 bp segments at each of the 254 differential loci
found in the mitogenomes of *T. esculentum* were prepared for the 43 independent individuals and
aligned by Muscle v5. A Maximum Likelihood (ML) phylogenetic tree using the Jukes-Cantor
model was drawn on the 43 chromosomes. Frequencies from 1000 bootstrap replicates were
labeled on the branches with 40% as cutoff. The topology was verified by a neighbor-joining tree
in Mega 11.

Genetic information exchange between organelles

The reference chloroplast genome sequence of *T. esculentum* (KX792933.1) was blasted to its
type 1 reference mitogenome (OK638188 and OK638189) and visualized by the Advanced
Circos function in TBtools (Krzywinski et al., 2009; Chen et al., 2020; https://github.com/CJ-
Chen/TBtools). Primers were designed to verify the presence of the 9,798bp long homologous
fragment in both the mitochondrial and chloroplast genomes. Four pairs of primers were
designed, using NCBI Primer-BLAST, to amplify the products spanning the two ends of the
9,798 bp cpDNA insertion in the mitochondrial and chloroplast genomes, respectively. The two
pairs of primers designed based on the mitogenome sequence included: MitoLL Forward

(ACGCAGAAAAGAGGCCGAA) and MitoLR Reverse
(CCTTCGTTTAAGAGAATGTTTTTGG), and the product size was 117 bp. MitoRL Forward
(TCTTTGCTACAGCTGATAAAAATCG) and MitoRR Reverse
(CCTATGTTCGTTTTCGCCCTG), and the product size was 120 bp. The two pairs of primers
designed based on the plastome sequence included: ChlLL Forward
(CGTAGTCGGTCTGGCCC) and MitoLR Reverse (CCTTCGTTTAAGAGAATGTTTTTGG),
and the product size was 117 bp. MitoRL Forward (TCTTTGCTACAGCTGATAAAAATCG)
and ChlRR Reverse (GCTTTTAATAATATGGCCGTGATCT), and the product size was 120 bp.

20 uL of mixtures were prepared by separately mixing DNA from two randomly selected *T.*
esculentum samples and Promega GoTaq Green Master Mix. The amplifications started with
denaturation at 95 ° C for 5 min, followed by 32 cycles of 95 ° C for 30 s, 55 ° C for 30 s, and
72 ° C for 30 s, and a final 72 ° C for 5 min.

Results

Genome structure and rearrangement

When the WGS Illumina reads from the 84 individuals were mapped to the two chromosomes of
the reference mitogenome of *T. esculentum*, LS1 (OK638188) and LS2 (OK638189), two
distinct mitogenomes were found. The mitogenomes of 45 individuals were similar to the
reference mitogenome, with only a few substitutions and indels seen, termed type 1. However,
the mitogenomes of the other 39 individuals were very different from the reference, with a large
number of sequence and structural differences, termed type 2 (Table S1). This is consistent with
the previously published study of *T. esculentum* chloroplast genomes, where these 84 individuals
were found to contain two distinct germplasms (Li and Cullis, 2023). These two cytotypes
actually differ not only in the chloroplast genome but also in the mitochondrial genome. The
PacBio HiFi reads from the type 2 individual Sample 4 were assembled by Canu to generate
three circular molecules M1 (OP795449), M2 (OP795450), and M4 (OP795447), and one linear
chromosome M3 (OP795448), with a total length of 436,568 bp and a GC content of 44.8%
(Table 1) (Figures S1-5). The four chromosomes consisted of 21 contigs assembled directly from
Illumina reads of type 2 individuals, containing four double-copy regions and one triple-copy
region (Figure 1 and Table 2). Among these multi-copy regions, homologous sequences of
contigs H and I were also doubled in coverage in the type 1 mitogenome of *T. esculentum*, but
the rest were present as single-copy sequences in the type 1 mitogenome (Li and Cullis, 2021).

Both ends of the chromosome M3 were found to be long repeats that were homologous to parts
of other chromosomes (Figure 2). In addition, in a very long range of 15 to 20 kb, the sequencing
depth gradually decreased towards both ends, and many reads were found to stop in this range.
This furthers confirms that this is a linear chromosome that exists in different lengths in cells
because of the lack of telomere protection. Linear chromosomes have been found to stably exist
in eukaryotic cells even in the absence of telomeres, through strand-invasion between terminal
sequences and their homologous internal sequences to form t-loops to protect the chromosomes
from degradation (de Lange, 2015). Because the repeats at both ends of M3 are very long, the
PacBio reads we obtained cannot span them to verify whether this linear molecule recombines
with other chromosomes. Long range PCR amplification that can amplify sequences above 20 kb
can be considered here to answer this question.

Numerous differences were found between the type 1 and type 2 mitogenome structures (Figure 3). One is a rare recombination on a pair of *trnfM* genes on the large circular molecule M1, which inverts the 12,846 bp sequence in between. The sequence before inversion also appeared in type 2 plants, with a frequency less than 2%. Furthermore, the three small rings of the type 1 mitogenome exist in different forms in the type 2 mitogenome, and four gaps have been found on them. New type 2 exclusive fragments were discovered, including a 2,108 bp segment, which connected originally distantly located contigs C1 and B. In addition, the recombination on a pair of 35 bp direct repeats joined contigs C1 and D2 to form a new circular molecule M2. C2 was found to connect with K1 and then further extended to N and L2 to form a linear chromosome, but the mechanism behind this is unclear. It can be seen that the deletion and insertion of the entire DNA segment, alongside repeat mediated recombination, can lead to dramatic changes in the mitogenome structure.

The two types of *T. esculentum* mitogenomes were compared by NUCmer alignment and then visualized by a synteny diagram, showing a high degree of similarity (Figure 4). The basic blocks making up the two mitogenomes are highly similar, except for the 2,108 bp type 2 unique fragment and some other short insertions and deletions, but the order of these blocks has been changed by recombination. When the WGS Illumina reads from the 84 individuals were mapped to the region where the 2,108 bp type 2 unique fragment resides, 39 samples were found to contain this fragment while 45 plants did not (Figures S19-28). Furthermore, in type 1 mitogenome, LS1 and LS2 are two autonomous circular chromosomes that do not recombine into one master circle, but in type 2 mitogenome, a linear chromosome M3 was found to contain homologous sequences from both LS1 and LS2, suggesting that these two molecules may have been related in evolution.

Blast results showed that this 2,108 bp type 2 mitogenome exclusive fragment was highly similar to the mitochondrial sequences of Fabaceae species *Lupinus albus* and *Indigofera tinctoria*, suggesting that this fragment was possibly derived in the evolution from *Lupinus* or *Indigofera* (Figure 5A). Two pairs of primers were designed and found to effectively identify the 2,108 bp fragment. As shown in Figure 5B and 5C, sample B and C, out of the six randomly selected samples, contained both the 990 bp left end band and the 289 bp right end band after amplification, indicating that only these two of the six had type 2 mitogenomes.

Gene annotation

Both type 1 and type 2 mitogenomes of *T. esculentum* were found to contain 35 unique protein-coding genes, 3 unique rRNA genes, and 16 different tRNA genes (Figure 6; Table 3) (Li and Cullis, 2021). The type 2 mitogenomes have two copies of *nad9* and *atp8*. The gene *nad9* is located on contig K1, a long repeat possessed by both chromosomes M3 and M4 in type 2 mitogenomes. Whereas, there is only one copy of K1 in type 1 mitogenomes. The gene *atp8* is located on a pair of long repeats J, so its copy number is doubled as is the case in both types of mitogenomes. The copy number of exon 3 and 4 of gene *nad5* is also doubled in type 2 mitogenomes but not in type 1, and it is not known whether this affects its expression level. In addition, there are two copies of *rrn5* and *rrnS* in type 2 mitogenomes but only one copy in type 1. A total of 26 tRNA genes were found in type 2 mitogenomes, including four copies of *trnfM-CAT*, three copies of *trnM-CAT*, three copies of *trnC-GCA*, two copies of *trnP* and *trnQ*, and 12 single-copy tRNA genes.

The atypical start codon ACG was used by three genes *nad1*, *nad4L*, and *rps10*, and ATT was used by gene *mttB*. This is consistent with the research on the mitogenome of common beans from which these four genes were all reported to use an alternative initiation codon ACG (Bi et al., 2020). C-to-U editing was found to be widely used in mitochondrial and chloroplast genes in land plants (Takenaka et al., 2013). ATT is also usually used as an alternative start codon in the mitogenome. For example, *mttB* in *Salix purpurea* was reported to use an ATT start codon as well (Wei et al., 2016).

Mitogenome divergence

The mitogenome of *T. esculentum* was highly divergent from those of the six selected legume species, which covered 23% to 48% of the marama mitogenome, ranging in length from 91.9-191.8 kb (Figure 7). Of these species, *C. canadensis* was most closely related to marama, while *M. sativa* was the least similar to marama. The mitogenomes of *G. max*, *L. japonicus*, and *V. radiata* all contain homologous sequences covering 25% of the marama mitogenome, equal to 99.89 kb in length. *Bauhinia variegata* is closer to marama than *C. canadensis* in the phylogenetic tree, but its mitogenome sequence is not available in NCBI GenBank (Wunderlin, 2010).

The loss of mitochondrial protein-coding genes and the functional transfer of these genes from the organelle genome to the nuclear genome are common in the evolution of angiosperms, but some plants tend to retain a more complete set of mitochondrial genes (Palmer et al., 2000). The mitogenomes of Cercidoideae species *T. esculentum* and *C. canadensis* contain functional protein-coding genes *sdh3*, *sdh4*, and *rpl10*, which have been lost in many other legumes (Table 4). Rare gene losses were also seen in the mitogenomes of legumes, such as *rpl5* in *G. max*, *cox2* in *V. radiata* and *rps1* in *L. japonicus*, but these genes all remain intact and functional in *T. esculentum* and *C. canadensis* (Alverson et al., 2011; Kazakoff et al., 2012; Chang et al., 2013).

In a pairwise comparison of the mitogenomes of seven legume species, the synteny plot revealed numerous rearrangements and a high degree of divergence among the mitogenomes of even closely related species (Figure 8). The mitogenomes of *C. canadensis* and *T. esculentum* contain many distinct regions but also some long homologous segments.

The phylogenetic tree shown in Figure 9 was built on the 24 conserved mitochondrial protein-coding genes *atp1*, *atp4*, *atp6*, *atp8*, *atp9*, *nad3*, *nad4*, *nad4L*, *nad6*, *nad7*, *nad9*, *mttB*, *matR*, *cox1*, *cox3*, *cob*, *ccmFn*, *ccmFc*, *ccmC*, *ccmB*, *rps3*, *rps4*, *rps12*, and *rpl16*, which are present in all these eight species. This tree is consistent with previously published phylogenetic trees constructed on chloroplast protein-coding genes (Kim and Cullis, 2017; Wang et al., 2018). As another species of Cercidoideae, *C. canadensis* was expected to be the closest relative of these plants to *T. esculentum*. Among Faboideae species, *M. sativa* and *L. japonicus* are closely related, and *V. radiata*, *G. max*, and *M. pinnata* belong another clade.

Nucleotide polymorphism

17 haplotypes were found in these 47 plants, which could be clearly divided into two groups: namely the type 2 plants from the Namibian farm and the UP farm and the remaining type 1 plants (Figure 10). The mitogenomes of type 2 plants are relative conserved, which may be caused by sampling errors, and a larger sample size is needed to verify. The only differences between type 2 plants were four deletions at four closely located loci on the chromosome LS2.

However, the type 1 plants can be divided into many groups according to the variations. Some
geographical patterns can be seen in the distribution of variation. For example, in the four plants
from Osire, there were three substitutions on the chromosome LS1, A>C at 146,140 bp, C>A at
173,053 bp, and C>A at 217,508 bp, and a deletion at 128,192 bp on the chromosome LS2.
These are variations unique to Osire plants. Similar geographic-specific variation can be seen in
plants elsewhere. However, our data may still have sampling issues. For example, only a single
sample was collected in some regions including Otjiwarongo and Okamatapati. In addition,
although distant wild individuals in each geographic region were intentionally selected, there is
no guarantee that they are not related. The findings here still need to be validated by sequencing
more samples and studied alongside the phenotypic performance of the plants to determine
whether any of these variations are the result of plants evolving to better adapt to different
environments.

Type 2 refers to 7 Index plants excluding Index8 from the Pretoria Farm, 29 M descendent plants
originally from the Namibia Farm excluding M40, and two individuals R1R2 and nar16 of
unknown origin. Type 1 represents all remaining plants, including A plants, S plants, Index8, and
M40. Numbers in parentheses indicate counts of exclusive variants of this type. LS1 and LS2 are
type 1 marama reference mitochondrial chromosomes in GenBank with accession numbers
OK638188 and OK638189.

A total of 254 differential loci were found in the mitogenomes of the 84 *T. esculentum*
individuals, including 143 SNPs, 52 insertions, and 59 deletions (Table 5). Type 1 and type 2
mitogenomes differed at 230 loci, including 129 substitutions, 50 deletions, and 51 insertions.
The mitogenomes of type 2 plants differed at only 4 loci, whereas that of type 1 plants differed at
24 loci.

The mitochondrial gene sequence of *T. esculentum* is very conserved. A total of 11 variations
were found in the mitochondrial gene sequence, and only 1 of them was in the coding sequence,
which was a 2368A>G substitution and resulted in a N303D change in the gene *matR* (Table 6).
Furthermore, 10 of the 11 variations were found on one subgenomic ring LS2 of the reference
mitogenome of *T. esculentum*. Whether the chromosome LS1 is more conserved than the
chromosome LS2 is unknown. Although the intergenic spacer of LS1 contained more variations
than LS2, the gene sequence on LS1 appeared to be more conserved, and cpDNA insertions were
also found rarely in LS1, but abundantly in chromosome LS2.

In the phylogenetic tree constructed on the differential loci of the mitogenomes of *T. esculentum*,
the two germplasms fell into two clusters as expected (Figure 11). Furthermore, the type 1 plants
were then divided into groups. Tsumkwe and Tsjaka were two distant sampling sites, but the
plants from these two locations were clustered in one clade, which was also seen in the
phylogenic tree built on the complete chloroplast genome. One wondered whether there were
factors other than geographical distance that determined the grouping of these plants. On the
other hand, plants from Epukiro, Osire, Ombujondjou, and Otjiwarongo belonged to one clade,
and these sites were geographically close together and were all arid areas with soil moisture
anomalies (SMA) typically below -.04 m^3/m^3 (NASA, n.d.). In contrast, plants from another
clade seemed to grow in less arid regions. Notable differences between the mitogenomes of
plants from the two clades included three consecutive substitutions from GCC to TTA at
positions 127,684 to 127,686 on chromosome LS1 (OK638188), although the function is

unknown. Of course, the sample size of this study is still relatively small, and more plants need
to be collected in areas with different soil moisture for verification.

SSRs and heteroplasmy analyses

A total of 48 SSR motifs were found by MISA in the reference mitogenome of *T. esculentum*,
LS1 (OK638188) and LS2 (OK638189), of which 38 were simple mononucleotide
microsatellites, accounting for 79.2% of all discovered SSR motifs (Figure 12). Among them, 37
are A/T mononucleotide repeats, and only one is a G/C repeat. There are 8 dinucleotide repeats
and 2 trinucleotide repeats. No simple sequence repeats with core motifs of four nucleotides or
longer were found. There are three microsatellites in the coding sequence of the gene, including
two 10 bp A/T repeats, one located at the boundary of the coding sequence of the gene *sdh3*, the
other located in the exon 4 of the gene *nad1*, and a 12 bp AT repeat siting in the coding sequence
of *mttB*.

An individual collected from Aminuis, A11 had approximately 96% type 1 alleles and 4% type 2
alleles at all differential loci between the two types of marama mitogenomes (Figure 13). It has
been confirmed that these minor alleles are not from homologous sequences in the nuclear
genome or the chloroplast genome, suggesting the presence of two different mitogenomes in the
same individual. It was previously reported that heteroplasmy also exists in the chloroplast
genome of A11, with a minor genome frequency reaching 11% (Jin and Cullis, 2023). This
means that in the chloroplasts and mitochondria of A11, both major and minor genomes exist,
and the frequency of the minor genome is higher than that of the other studied plants. This is
most likely to be caused by an accidentally occurred paternal leakage. The proportions of
mitochondrial and chloroplast minor genomes differed in A11, indicating that this may be true
heterogeneity rather than due to accidental mixing of samples. However, being the only sample
with high overall organelle genome heterogeneity is not convincing, and more plants from
Aminuis need to be sequenced and studied.

In the mitogenomes of other individuals, heteroplasmy was found to be less common than in
their chloroplast genomes, and even low proportions of base substitutions below 2% were
relatively rare, and even if present, many were found from mitochondrial homologous segments
in the nuclear genome. However, at a few differential loci, including the three consecutive
substitutions at positions 127,684 to 127,686 and another substitution at 140,922 on chromosome
LS1 (OK638188), obvious heteroplasmy could be seen in multiple individuals, with the
proportion of minor alleles even up to 35%. These loci may have played important roles in the
evolution of these individuals under environmental selection.

Sequence transfer between chloroplast and mitochondrial genomes

It was interesting to find that the chloroplast DNA insertions were concentrated in a subgenomic
ring of the mitogenome (Figure 14). A low collinearity between the two genomes was seen, and
the inserted cpDNA fragments seemed to be completely rearranged. These contained a 9,798 bp
fragment, in which a large number of variations were observed. Primers were designed to
amplify across the two ends of this fragment in both plastome and mitogenome to verify its
presence (Figure 15).

20 variations were found in this long homologous DNA fragment, of which 13 mutations
occurred in the mitogenome and 7 occurred in the chloroplast genome (Table 7). These do not

include the differences between the two organelle genomes at another 72 loci in this segment
(this included 22 deletions, 6 insertions, and 44 SNPs.), which are the same for both germplasms,
making it difficult to tell where the mutation occurred. 10 variations were found in the gene
sequence on this segment, of which only 2 synonymous substitutions occurred in the chloroplast
genome, and the remaining 8 were in the mitogenome, which had a great impact on transcription,
including the introduction of early stop codons (Table S3).

The mitogenome of *T. esculentum* was 399,572 bp (type 1) in length, and a total of 254
variations were found in the mitogenomes of the 84 individuals. The length of the chloroplast
genome was 161,537 bp (type 1), and a total of 147 variations were found in the plastomes of
these plants. The value of variation per nucleotide of the chloroplast genome is even higher than
that of the mitogenome, so it is speculated that these chloroplast genes *psbC*, *rps14*, *psaB*, and
psaA are protected by some mechanism in the chloroplast genome, but after being transferred to
the mitogenome, the protection mechanism disappears and these genes begin to accumulate
mutations that render them nonfunctional and pseudogenes.

Conclusion

The comparative analysis of the organelle genomes of 84 *T. esculentum* individuals revealed two
germplasms with distinct mitochondrial and chloroplast genomes. The type 1 mitogenome
contains two autonomous rings, or five smaller subgenomic circular molecules. These two
equimolar structures are thought to be interchangeable through recombination on three pairs of
long direct repeats (Li and Cullis, 2021). The type 2 mitogenome contains three circular
molecules and one linear chromosome. It also has a unique fragment of 2,108 bp in length, likely
derived from the mitogenome of *Lupinus*. Primers were designed to amplify on this fragment for
germplasm typing. Both ends of the linear chromosome are repetitive sequences, also present in
other molecules, on which recombination may occur to protect the linear chromosome from
degradation.

The structural variation resulted in increased copy number of the genes *atp8*, *nad5* (exon3 and
exon4), *nad9*, *rrnS*, *rrn5*, *trnC*, and *trnfM* in the type 2 mitogenome, but it is unknown whether
this is reflected in the gene expression level. The genes *nad1*, *nad4L*, *rps10* and *mttB* were found
to use alternative start codons ACG and ATT, similar to the mitogenome of common bean (Bi et
al., 2020). A total of 254 differential loci were found in the mitogenomes of the 84 *T. esculentum*
individuals. Type 1 and type 2 mitogenomes differ from each other at 230 of these loci. Only one
of these 254 variations was found in the mitochondrial gene coding sequence, which altered the
amino acid sequence synthesized by *matR*.

The evolutionary study of the differential loci in the mitogenomes of *T. esculentum* found that
the two types of plants fell into two clusters, as expected. The type 1 plants can be further
divided into two clades, and the divergence is thought to be possibly related to soil moisture
content, although this still needs to be verified by studies with larger sample sizes. Three
consecutive substitutions at positions 127,684 to 127,686 on chromosome LS1 (OK638188) may
have played an important role here. The phylogenetic tree constructed based on the
mitochondrial genes of *T. esculentum* and other related Fabaceae species was consistent with the
previously published one built on chloroplast genes. *C. canadensis* and *T. esculentum* were found
to be closely related, and they have more complete sets of mitochondrial genes than the other
legumes.

Heteroplasmy in the mitochondrial genome of *T. esculentum* is not as prevalent as in its
chloroplast genome, but higher levels do exist at certain loci, such as loci 127,684 to 127,686 on
chromosome LS1. Among all samples, only one individual A11 from Aminuis had a generally
high degree of heteroplasmy at most differential loci, consistent with what was seen in its
chloroplast genome. Whether it is because of some of its own characteristics that A11 escaped
the genetic bottleneck at the developmental stage is still unclear.

Chloroplast insertions are thought to be concentrated in one of the subgenomic rings of the
mitogenome of *T. esculentum*. The mitogenomes of all marama individuals were found to
contain a long chloroplast DNA insertion over 9 kb in length. The study of the polymorphisms
on this segment infers that the sequence is protected by some mechanism in the chloroplast
genome, so only a small amount of synonymous substitutions is retained, but after being inserted
into the mitogenome, a large number of mutations have accumulated on it, making the genes on
it lose their function.

Supplementary data

549   Supplementary materials have been uploaded to JXB including 3 tables and 32 figures.

Acknowledgments

551   The authors would like to thank K. Logue for help with the initial genome assembly, to P.
552   Chimwamurombe, M. Takundwa, J. Vorster, and K. Kunert for providing marama samples from
553   Namibia and from the University of Pretoria Farm, to students in BIOL 301/401 in 2015 for
554   assistance with DNA isolation and in BIOL 309 in 2018 for sample collection.

Author Contributions

556   J.L. carried out the bioinformatics assembly and data analysis and drafted the manuscript. C.C.
557   conceived initial ideas for the project, provided extracted DNA, and assisted in writing and
558   editing the manuscript. All authors contributed to the submitted article.

Conflict of interest

560   The authors affirm that the research was carried out independent of any commercial or financial
561   associations that could be perceived as a potential conflict of interest.

Funding

563   This work was supported by teaching resources from the Department of Biology, Case Western
564   Reserve University.

Data availability

566   The sequencing data used in this study are available and have been deposited under BioProject
567   PRJNA897263 (https://www.ncbi.nlm.nih.gov/bioproject/897263).

References

Alverson, A. J., Wei, X., Rice, D. W., Stern, D. B., Barry, K., and Palmer, J. D. (2010). Insights
into the evolution of mitochondrial genome size from complete sequences of *Citrullus lanatus*
and *Cucurbita pepo* (Cucurbitaceae). Molecular Biology and Evolution, 27(6), 1436-1448.
https://doi.org/10.1093/molbev/msq029

Alverson, A. J., Zhuo, S., Rice, D. W., Sloan, D. B., and Palmer, J. D. (2011). The mitochondrial genome of the legume *Vigna radiata* and the analysis of recombination across short mitochondrial repeats. PLoS ONE, 6(1), e16404. https://doi.org/10.1371/journal.pone.0016404.

Amarteifio, J. O., and Moholo, D. (1998). The chemical composition of four legumes consumed in Botswana. Journal of Food Composition and Analysis, 11(4), 329-332. https://doi.org/10.1006/jfca.1998.0595

Andersson, S. G. E., Karlberg, O., Canbäck, B., and Kurland, C. G. (2003). On the origin of mitochondria: a genomics perspective. Philosophical Transactions of the Royal Society B, 358(1429), 165-179. https://doi.org/10.1098/rstb.2002.1193

Anisimova, I. N., and Gavrilenko, T. (2017). Cytoplasmic male sterility and prospects for its utilization in potato breeding, genetic studies and hybrid seed production. Russian Journal of Genetics: Applied Research, 7(7), 721-735. https://doi.org/10.1134/s2079059717070024

Arrieta-Montiel, M. P., Shedge, V., Davila, J. I., Christensen, A. J., and Mackenzie, S. A. (2009). Diversity of the Arabidopsis mitochondrial genome occurs via nuclear-controlled recombination activity. Genetics, 183(4), 1261-1268. https://doi.org/10.1534/genetics.109.108514

Beier, S., Thiel, T., Münch, T., Scholz, U., and Mascher, M. (2017). Misa-web: a web server for microsatellite prediction. Bioinformatics, 33, 2583-2585. https://doi:10.1093/bioinformatics/btx198.

Bi, C., Lu, N., Xu, Y., He, C., and Lu, Z. (2020). Characterization and analysis of the mitochondrial genome of common bean (*Phaseolus vulgaris*) by Comparative Genomic Approaches. International Journal of Molecular Sciences, 21(11), 3778. https://doi.org/10.3390/ijms21113778

Belitz, H. D., Grosch, W., and Schieberle, P. (2004). Food Chemistry. Berlin: Springer. http://doi: 10.1007/978-3-662-07279-0

Boore, J. L. (1999). Animal mitochondrial genomes. Nucleic Acids Research, 27(8), 1767-1780. https://doi.org/10.1093/nar/27.8.1767

Bower, N., Hertel, K., Oh, J., and Storey, R. (1988). Nutritional evaluation of marama bean (*Tylosema esculentum*, Fabaceae): analysis of the seed. Economic Botany, 42(4), 533-540. https://doi.org/10.1007/bf02862798

Budar, F., and Roux, F. (2011). The role of organelle genomes in plant adaptation. Plant Signaling & Behavior, 6(5), 635-639. https://doi.org/10.4161/psb.6.5.14524

Carbonell-Caballero, J., Alonso, R., Ibanez, V., Terol, J., Talon, M., and Dopazo, J. (2015). A phylogenetic analysis of 34 chloroplast genomes elucidates the relationships between wild and domestic species within the genus *Citrus*. Molecular Biology and Evolution, 32(8), 2015-2035. https://doi.org/10.1093/molbev/msv082

Chan, P., Lin, B., Mak, A., and Lowe, T. (2021). tRNAscan-SE 2.0: improved detection and functional classification of transfer RNA genes. Nucleic Acids Research, 49(16), 9077-9096. https://doi.org/10.1093/nar/gkab688

Chang, S., Wang, Y., Lu, J., Gai, J., Li, J., Chu, P., Guan, R., and Zhao, T. (2013). The mitochondrial genome of soybean reveals complex genome structures and gene evolution at intercellular and phylogenetic levels. PLoS ONE, 8(2), e56502. https://doi.org/10.1371/journal.pone.0056502

Chen, C., Chen, H., Zhang, Y., Thomas, H. R., Frank, M. H., He, Y., and Xia, R. (2020). TBtools: An Integrative Toolkit Developed for Interactive Analyses of Big Biological Data. Molecular Plant, 13(8), 1194–1202. https://doi.org/10.1016/j.molp.2020.06.009

Cole, L. W., Guo, W., Mower, J. P., and Palmer, J. D. (2018). High and variable rates of repeat-mediated mitochondrial genome rearrangement in a genus of plants. Molecular Biology and Evolution. https://doi.org/10.1093/molbev/msy176

Cullis, C. A., Chimwamurombe, P. M., Barker, N. P., Kunert, K. J., and Vorster, J. (2018). Orphan legumes growing in dry environments: marama bean as a case study. Frontiers in Plant Science, 9. https://doi.org/10.3389/fpls.2018.01199

Cullis, C. A., Chimwamurombe, P. M., Kunert, K. J., and Vorster, J. (2022). Perspective on the present state and future usefulness of marama bean (*Tylosema esculentum*). Food and Energy Security. https://doi.org/10.1002/fes3.422

Cullis, C. A., Lawlor, D. W., Chimwamurombe, P. M., Bbebe, N., Kunert, K. J., and Vorster, J. (2019). Development of marama bean, an orphan legume, as a crop. Food and Energy Security, 8(3). https://doi.org/10.1002/fes3.164

Darling, A. E., Mau, B., Blattner, F. R., and Perna, N. T. (2004b). Mauve: Multiple alignment of conserved genomic sequence with rearrangements. Genome Research, 14(7), 1394-1403. https://doi.org/10.1101/gr.2289704

de Lange, T. (2015). A loopy view of telomere evolution. *Frontiers in Genetics*, 6. https://doi.org/10.3389/fgene.2015.00321

Delcher, A. L., Phillippy, A., Carlton, J., and Salzberg, S.L. (2002). Fast algorithms for large-scale genome alignment and comparison. Nucleic Acids Research, 30(11), 2478-2483. https://doi.org/10.1093/nar/30.11.2478

Dong, S., Zhao, C., Chen, F., Liu, Q. H., Zhang, S., Wu, H., Zhang, L., and Liu, Y. (2018). The complete mitochondrial genome of the early flowering plant *Nymphaea colorata* is highly repetitive with low recombination. BMC Genomics, 19(1). https://doi.org/10.1186/s12864-018-4991-4

Drouin, G., Daoud, H., and Xia, J. (2008). Relative rates of synonymous substitutions in the mitochondrial, chloroplast and nuclear genomes of seed plants. Molecular Phylogenetics and Evolution, 49(3), 827-831. https://doi.org/10.1016/j.ympev.2008.09.009

Edgar, R. C. (2022). Muscle5: High-accuracy alignment ensembles enable unbiased assessments of sequence homology and phylogeny. Nature Communications, 13(1). https://doi.org/10.1038/s41467-022-34630-w

Floros, V. I., Pyle, A., Dietmann, S., Wei, W., Tang, W. W. C., Irie, N., Payne, B. a. I., Capalbo, A., Noli, L., Coxhead, J., Hudson, G., Crosier, M., Strahl, H., Khalaf, Y., Saitou, M., Ilic, D., Surani, M. A., and Chinnery, P. F. (2018). Segregation of mitochondrial DNA heteroplasmy through a developmental genetic bottleneck in human embryos. Nature Cell Biology, 20(2), 144-151. https://doi.org/10.1038/s41556-017-0017-8

Goremykin, V. V., Salamini, F., Velasco, R., and Viola, R. (2008). Mitochondrial DNA of *Vitis vinifera* and the issue of rampant horizontal gene transfer. Molecular Biology and Evolution, 26(1), 99-110. https://doi.org/10.1093/molbev/msn226

Greiner, S., Lehwark, P., and Bock, R. (2019). OrganellarGenomeDRAW (OGDRAW) version 1.3.1: expanded toolkit for the graphical visualization of organellar genomes. Nucleic Acids Research, 47(W1), W59-W64. https://doi.org/10.1093/nar/gkz238

Gualberto, J. M., Mileshina, D., Wallet, C., Niazi, A., Weber-Lotfi, F., and Dietrich, A. (2014). The plant mitochondrial genome: dynamics and maintenance. Biochimie, 100, 107-120. https://doi.org/10.1016/j.biochi.2013.09.016

Gualberto, J. M., and Newton, K. J. (2017). Plant mitochondrial genomes: Dynamics and mechanisms of mutation. Annual Review of Plant Biology, 68(1), 225-252. https://doi.org/10.1146/annurev-arplant-043015-112232

Guo, W., Grewe, F., Fan, W., Young, G. J., Knoop, V., Palmer, J. D., and Mower, J. P. (2016). *Ginkgo* and *Welwitschia* mitogenomes reveal extreme contrasts in Gymnosperm mitochondrial evolution. Molecular Biology and Evolution, 33(6), 1448-1460. https://doi.org/10.1093/molbev/msw024

Hanson, M. R. (1990). Plant mitochondrial mutations and male sterility. Annual Review of Genetics, 25(1), 461-486. https://doi.org/10.1146/annurev.ge.25.120191.002333

Hao, Z., Lv, D., Ge, Y., Shi, J., Weijers, D., Yu, G., and Chen, J. (2020). RIdeogram: drawing SVG graphics to visualize and map genome-wide data on the idiograms. PeerJ Computer Science, 6, e251. https://doi.org/10.7717/peerj-cs.251

Holse, M., Husted, S., and Hansen, Å. (2010). Chemical composition of marama bean (*Tylosema esculentum*) - a wild African bean with unexploited potential. Journal of Food Composition and Analysis, 23(6), 648-657. https://doi.org/10.1016/j.jfca.2010.03.006

Hon, T., Mars, K., Young, G., Tsai, Y., Karalius, J. W., Landolin, J. M., Maurer, N., Kudrna, D., Hardigan, M. A., Steiner, C. C., Knapp, S. J., Ware, D., Shapiro, B., Peluso, P. R., and Kingan, S. B. (2020). Highly accurate long-read HiFi sequencing data for five complex genomes. Scientific Data, 7(1). https://doi.org/10.1038/s41597-020-00743-4

Kozik, A., Rowan, B. A., Lavelle, D., Berke, L., Schranz, M. E., Michelmore, R. W., and Christensen, A. J. (2019). The alternative reality of plant mitochondrial DNA: One ring does not rule them all. PLOS Genetics, 15(8), e1008373. https://doi.org/10.1371/journal.pgen.1008373

Kvist, L., Martens, J., Nazarenko, A. Y., and Orell, M. (2003). Paternal leakage of mitochondrial DNA in the great tit (*Parus major*). Molecular Biology and Evolution, 20(2), 243-247. https://doi.org/10.1093/molbev/msg025

Langmead, B., and Salzberg, S. L. (2012). Fast gapped-read alignment with Bowtie 2. Nature Methods, 9(4), 357-359. https://doi.org/10.1038/nmeth.1923

Li, H. (2011). A statistical framework for SNP calling, mutation discovery, association mapping and population genetical parameter estimation from sequencing data. Bioinformatics, 27(21), 2987-2993. https://doi.org/10.1093/bioinformatics/btr509

Li, J., and Cullis, C. (2021). The multipartite mitochondrial genome of marama (*Tylosema esculentum*). Frontiers in Plant Science, 12. https://doi.org/10.3389/fpls.2021.787443

Li, J., Li, J., Ma, Y., Kou, L., Wei, J., and Wang, W. (2022). The complete mitochondrial genome of okra (*Abelmoschus esculentus*): using nanopore long reads to investigate gene transfer from chloroplast genomes and rearrangements of mitochondrial DNA molecules. BMC Genomics, 23(1). https://doi.org/10.1186/s12864-022-08706-2

Li, J., and Cullis, C. (2023). Comparative analysis of 84 chloroplast genomes of *Tylosema esculentum* reveals two distinct cytotypes. Frontiers in Plant Science, 13:1025408. https://doi.org/10.3389/fpls.2022.1025408

Lössl, A. G., Adler, N., Horn, R., Frei, U., and Wenzel, G. (1999). Chondriome-type characterization of potato: Mt α, β, γ, δ, □ and novel plastid-mitochondrial configurations in somatic hybrids. Theoretical and Applied Genetics, 99(1-2), 1-10. https://doi.org/10.1007/s001220051202

Luo, S., Valencia, C. A., Zhang, J., Lee, N., Slone, J., Gui, B., Wang, X., Li, Z., Dell, S., Brown, J., Chen, S. X., Chien, Y., Hwu, W., Fan, P., Wong, L. C., Atwal, P. S., and Huang, T. (2018). Biparental inheritance of mitochondrial DNA in humans. Proceedings of the National Academy of Sciences of the United States of America, 115(51), 13039-13044. https://doi.org/10.1073/pnas.1810946115

Jackson, J. C., Duodu, K. G., Holse, M., De Faria, M. D. L., Jordaan, D., Chingwaru, W., Hansen, A. L., et al. (2010). The morama Bean (*Tylosema esculentum*): A potential crop for southern Africa. Advances in Food and Nutrition Research, 187-246. https://doi.org/10.1016/b978-0-12-374468-5.00005-2

Johnson, M., Zaretskaya, I., Raytselis, Y., Merezhuk, Y., McGinnis, S., and Madden, T. L. (2008). NCBI BLAST: a better web interface. Nucleic Acids Research, 36(Web Server), W5–W9. https://doi.org/10.1093/nar/gkn201

Heng, S., Wei, C., Jing, B., Wan, Z., Wen, J., Yi, B., Ma, C., Tu, J., and Fu, T. (2014). Comparative analysis of mitochondrial genomes between the *hau* cytoplasmic male sterility (CMS) line and its iso-nuclear maintainer line in *Brassica juncea* to reveal the origin of the CMS-associated gene *orf288*. BMC Genomics, 15(1). https://doi.org/10.1186/1471-2164-15-322

Kazakoff, S. H., Imelfort, M., Edwards, D., Koehorst, J., Biswas, B., Batley, J., Scott, P. T., and Gresshoff, P. M. (2012). Capturing the biofuel wellhead and powerhouse: The chloroplast and

mitochondrial genomes of the leguminous feedstock tree *Pongamia pinnata*. PLoS ONE, 7(12), e51687. https://doi.org/10.1371/journal.pone.0051687

Keeling, P. J., and Palmer, J. D. (2008). Horizontal gene transfer in eukaryotic evolution. Nature Reviews Genetics, 9(8), 605-618. https://doi.org/10.1038/nrg2386

Khan, A. (2018). Phytochemicals and their role in curing fatal diseases: A review. Pure and Applied Biology, 7(4). https://doi.org/10.19045/bspab.2018.700193

Kim, Y., and Cullis, C. (2017). A novel inversion in the chloroplast genome of marama (*Tylosema esculentum*). Journal of Experimental Botany, 68(8), 2065-2072. https://doi.org/10.1093/jxb/erw500

Kitazaki, K., and Kubo, T. (2010). Cost of Having the Largest Mitochondrial Genome: Evolutionary Mechanism of Plant Mitochondrial Genome. Journal of Botany, 2010, 1-12. https://doi.org/10.1155/2010/620137

Krzywinski, M., Schein, J., Birol, N., Connors, J., Gascoyne, R., Horsman, D., Jones, S. J., and Marra, M. A. (2009). Circos: An information aesthetic for comparative genomics. *Genome Research*, *19*(9), 1639–1645. https://doi.org/10.1101/gr.092759.109

Marçais, G., Delcher, A. L., Phillippy, A. M., Coston, R., Salzberg, S. L., and Zimin, A. (2018). MUMmer4: A fast and versatile genome alignment system. PLOS Computational Biology, 14(1), e1005944. https://doi.org/10.1371/journal.pcbi.1005944

Manchekar, M., Scissum-Gunn, K., Song, D., Khazi, F. R., McLean, S. L., and Nielsen, B. L. (2006). DNA recombination activity in soybean mitochondria. Journal of Molecular Biology, 356(2), 288-299. https://doi.org/10.1016/j.jmb.2005.11.070

Miga, K. H., Koren, S., Rhie, A., Vollger, M. R., Gershman, A., Bzikadze, A., Brooks, S., et al. (2020). Telomere-to-telomere assembly of a complete human X chromosome. Nature, 585(7823), 79-84. https://doi.org/10.1038/s41586-020-2547-7

Moner, A. M., Furtado, A., and Henry, R. J. (2020). Two divergent chloroplast genome sequence clades captured in the domesticated rice gene pool may have significance for rice production. BMC Plant Biology, 20(1). https://doi.org/10.1186/s12870-020-02689-6

Naish, M., Alonge, M., Wlodzimierz, P., Tock, A. J., Abramson, B. W., Schmücker, A., Mandáková, et al. (2021). The genetic and epigenetic landscape of the *Arabidopsis* centromeres. Science, 374(6569). https://doi.org/10.1126/science.abi7489

NASA. (n.d.). *Drought harms corn crops in southern Africa*. NASA. Retrieved February 11, 2023, from https://www.earthobservatory.nasa.gov/images/144704/drought-harms-corn-crops-in-southern-africa

Nikiforova, S. V., Cavalieri, D., Velasco, R., and Goremykin, V. V. (2013). Phylogenetic analysis of 47 chloroplast genomes clarifies the contribution of wild species to the domesticated apple maternal line. Molecular Biology and Evolution, 30(8), 1751-1760. https://doi.org/10.1093/molbev/mst092

Nurk, S., Walenz, B. P., Rhie, A., Vollger, M. R., Logsdon, G. A., Grothe, R., Miga, K. H., Eichler, E. E., Phillippy, A. M., and Koren, S. (2020). HiCanu: accurate assembly of segmental

duplications, satellites, and allelic variants from high-fidelity long reads. Genome Research, 30(9), 1291-1305. https://doi.org/10.1101/gr.263566.120

Oldenburg, D. J., and Bendich, A. J. (1996). Size and structure of replicating mitochondrial DNA in cultured tobacco cells. The Plant Cell, 447-461. https://doi.org/10.1105/tpc.8.3.447

Omotayo, A. O., and Aremu, A. O. (2021). Marama bean [*Tylosema esculentum* (Burch.) A. Schreib.]: an indigenous plant with potential for food, nutrition, and economic sustainability. Food & Function, 12(6), 2389-2403. https://doi.org/10.1039/d0fo01937b

Palmer, J. D., Adams, K. L., Cho, Y., Parkinson, C. L., Qiu, Y. L., and Song, K. (2000). Dynamic evolution of plant mitochondrial genomes: Mobile genes and introns and highly variable mutation rates. Proceedings of the National Academy of Sciences, 97(13), 6960-6966. https://doi.org/10.1073/pnas.97.13.6960

Palmer, J. D., and Herbon, L. A. (1988). Plant mitochondrial DNA evolved rapidly in structure, but slowly in sequence. Journal of Molecular Evolution, 28(1-2), 87-97. https://doi.org/10.1007/bf02143500

Palmer, J. D., and Shields, C. R. (1984). Tripartite structure of the *Brassica campestris* mitochondrial genome. Nature, 307(5950), 437-440. https://doi.org/10.1038/307437a0

Robinson, J. A., Thorvaldsdottir, H., Winckler, W., Guttman, M., Lander, E. S., Getz, G., and Mesirov, J. P. (2011). Integrative genomics viewer. Nature Biotechnology, 29(1), 24-26. https://doi.org/10.1038/nbt.1754

Sanetomo, R., and Gebhardt, C. (2015). Cytoplasmic genome types of European potatoes and their effects on complex agronomic traits. BMC Plant Biology, 15(1). https://doi.org/10.1186/s12870-015-0545-y

Skippington, E., Barkman, T. J., Rice, D. W., and Palmer, J. D. (2015). Miniaturized mitogenome of the parasitic plant *Viscum scurruloideum* is extremely divergent and dynamic and has lost all nad genes. Proceedings of the National Academy of Sciences of the United States of America, 112(27). https://doi.org/10.1073/pnas.1504491112

Sinou, C., Cardinal-McTeague, W. M., and Bruneau, A. (2020). Testing generic limits in Cercidoideae (Leguminosae): Insights from plastid and duplicated nuclear gene sequences. Taxon, 69(1), 67-86. https://doi.org/10.1002/tax.12207

Sloan, D. B., Alverson, A. J., Chuckalovcak, J. P., Wu, M., McCauley, D. E., Palmer, J. D., and Taylor, D. (2012). Rapid evolution of enormous, multichromosomal genomes in flowering plant mitochondria with exceptionally high mutation rates. PLOS Biology, 10(1), e1001241. https://doi.org/10.1371/journal.pbio.1001241

Smyda-Dajmund, P., Śliwka, J., Janiszewska, M., and Zimnoch-Guzowska, E. (2020). Cytoplasmic diversity of potato relatives preserved at Plant Breeding and Acclimatization Institute in Poland. Molecular Biology Reports, 47(5), 3929-3935. https://doi.org/10.1007/s11033-020-05486-4

Straub, S. C. K., Cronn, R., Edwards, C., Fishbein, M., and Liston, A. (2013). Horizontal transfer of DNA from the mitochondrial to the plastid genome and its subsequent evolution in milkweeds

(Apocynaceae). Genome Biology and Evolution, 5(10), 1872-1885. https://doi.org/10.1093/gbe/evt140

Suchard, M. A., Lemey, P., Baele, G., Ayres, D. L., Drummond, A. J., and Rambaut, A. (2017). Bayesian phylogenetic and phylodynamic data integration using BEAST 1.10. Virus Evolution, 4(1). https://doi.org/10.1093/ve/vey016

Takenaka, M., Zehrmann, A., Verbitskiy, D., Härtel, B., and Brennicke, A. (2013). RNA editing in plants and its evolution. Annual Review of Genetics, 47(1), 335-352. https://doi.org/10.1146/annurev-genet-111212-133519

Telenti, A., Pierce, L. C. T., Biggs, W. H., Di Iulio, J., Wong, E. B., Fabani, M. M., Kirkness, E. F., Moustafa, A. A., Shah, N., Xie, C., Brewerton, S. C., Bulsara, N., Garner, C., Metzker, G., Sandoval, E., Perkins, B. A., Och, F. J., Turpaz, Y., and Venter, J. C. (2016). Deep sequencing of 10,000 human genomes. Proceedings of the National Academy of Sciences of the United States of America, 113(42), 11901-11906. https://doi.org/10.1073/pnas.1613365113

The Legume Phylogeny Working Group (LPWG). (2017). A new subfamily classification of the Leguminosae based on a taxonomically comprehensive phylogeny. Taxon, 66(1), 44-77. https://doi.org/10.12705/661.3

Unseld, M., Marienfeld, J., Brandt, P., and Brennicke, A. (1997). The mitochondrial genome of *Arabidopsis thaliana* contains 57 genes in 366,924 nucleotides. *Nature Genetics*, *15*(1), 57-61. https://doi.org/10.1038/ng0197-57

Varré, J., D'Agostino, N., Touzet, P., Gallina, S., Tamburino, R., Cantarella, C., Ubrig, E., Cardi, T., Drouard, L., Gualberto, J. M., and Scotti, N. (2019). Complete sequence, multichromosomal architecture and transcriptome analysis of the *Solanum tuberosum* mitochondrial genome. International Journal of Molecular Sciences, 20(19), 4788. https://doi.org/10.3390/ijms20194788

Wang, Y. H., Wicke, S., Wang, H., Jin, J. J., Chen, S. Y., Zhang, S. D., Li, D. Z., and Yi, T. S. (2018). Plastid genome evolution in the early-diverging legume subfamily Cercidoideae (Fabaceae). Frontiers in Plant Science, 9. https://doi.org/10.3389/fpls.2018.00138

Ward, B. J., Anderson, R. H., and Bendich, A. J. (1981). The mitochondrial genome is large and variable in a family of plants (Cucurbitaceae). Cell, 25(3), 793-803. https://doi.org/10.1016/0092-8674(81)90187-2

Wei, S., Wang, X., Bi, C., Xu, Y., Wu, D., and Ye, N. (2016). Assembly and analysis of the complete *Salix purpurea* L. (Salicaceae) mitochondrial genome sequence. SpringerPlus, 5(1), 1894. https://doi.org/10.1186/s40064-016-3521-6

Wolfe, K. H., Li, W., and Sharp, P. M. (1987). Rates of nucleotide substitution vary greatly among plant mitochondrial, chloroplast, and nuclear DNAs. Proceedings of the National Academy of Sciences of the United States of America, 84(24), 9054-9058. https://doi.org/10.1073/pnas.84.24.9054

Woloszynska, M., Bocer, T., Mackiewicz, P., and Janska, H. (2004). A fragment of chloroplast DNA was transferred horizontally, probably from non-eudicots, to mitochondrial genome of *Phaseolus*. Plant Molecular Biology, 56(5), 811-820. https://doi.org/10.1007/s11103-004-5183-y

Wunderlin, R. P. (2010). Reorganization of the Cercideae (Fabaceae: Caesalpinioideae). Phytoneuron, 48, 1-5.

Ye, J., Coulouris, G., Zaretskaya, I., Cutcutache, I., Rozen, S., and Madden, T. L. (2012). Primer-BLAST: A tool to design target-specific primers for polymerase chain reaction. BMC Bioinformatics, 13(1). https://doi.org/10.1186/1471-2105-13-134

Table

Table 1. Chromosome base composition of the type 2 *T. esculentum* mitogenome.

Molecule	A (%)	C (%)	G (%)	T (%)	G~C (%)	Length (bp)
M1	27.90	22.18	22.22	27.70	44.40	169,406
M2	27.60	21.63	23.26	27.52	44.90	56,355
M3	27.46	22.25	22.73	27.55	45.00	97,120
M4	27.70	22.58	22.44	27.29	45.00	113,687

Table 2. Length of the primary scaffolds constituting the type 2 *T. esculentum* mitogenome.

Unit number	Length (bp)	Unit number	Length (bp)
A	82,874	K1	4,052
B	39,273	K2	52,730
C1	30,017	L1	20,069
C2	5,272	L2	22,301
D1	5,866	N	27,581
D2	2,722	O	4,881
D3	26,671	M	9,654
H	5,212	Q	2,108
E1	1,768		
E2	25,384		
F	8,590		
G	5,798		
I	2,265		

Table 3. Gene annotation of the type 2 mitogenome of *T. esculentum*.

Category	Gene Name			
	M1	M2	M3	M4
Complex I (NADH dehydrogenase)	*nad1#, nad4L, nad6*	*nad2#, nad3, nad5#*	*nad2#, nad5#, nad7#, nad9*	*nad1#, nad4#, nad5#, nad9*
Complex II (succinate dehydrogenase)	*sdh4*	*sdh3*		
Complex III (ubiquinol cytochrome-c	*cob*			

reductase)

Complex IV (cytochrome-c oxidase)	cox1, cox3	cox2#		
Complex V (ATP synthase)	atp1, atp4, atp6, atp8*(2)	atp9		
Cytochrome c biogenesis	ccmFn		ccmC	ccmB, ccmFc#
Large subunit ribosomal proteins	rpl5, rpl10		rpl16	
Small subunit ribosomal proteins	rps1, rps4, rps10#, rps14	rps12	rps3#	
Maturases				matR
Transport membrane protein				mttB
Ribosomal RNAs	rrnL		rrn5, rrnS	rrn5, rrnS
Transfer RNAs	trnD-GTC, trnC-GCA, trnQ-TTG, trnH-GTG, trnfM-CAT*(2), trnF-GAA, trnP-TGG, trnS-GCT	trnC-GCA, trnfM-CAT	trnN-GTT, trnC-GCA, trnfM-CAT, trnM-CAT, trnY-GTA	trnQ-TTG, trnE-TTC, trnG-GCC, trnK-TTT, trnM-CAT*(2), trnP-TGG, trnS-TGA, trnT-TGT#, trnW-CCA

Genes with introns are marked with #. Genes with multiple copy numbers on the same chromosome are labeled with *, and the numbers in parentheses indicate the corresponding copy numbers.

Table 4. List of mitochondrial protein-coding genes lost during the evolution of some Fabaceae species.

Gene	T. esculentum	C. canadensis	L. japonicus	M. sativa	M. pinnata	V. radiata	G. max
sdh3	+	+	-	-	+	-	-
sdh4	+	+	-	-	-	-	-
cox2	+	+	+	+	+	-	+
rpl2	-	-	-	-	-	-	-
rpl5	+	+	+	+	+	+	-
rpl10	+	+	-	-	-	-	-
rps1	+	+	-	+	+	+	+
rps2	-	-	-	-	-	-	-
rps7	-	-	-	-	-	-	-
rps8	-	-	-	-	-	-	-
rps11	-	-	-	-	-	-	-
rps13	-	-	-	-	-	-	-
rps19	-	-	-	-	-	-	-

Table 5. Total number of variations found when mapping the WGS reads of all 84 individuals to the type 1 *T. esculentum* reference mitochondrial genomes OK638188 and OK638189.

Variation Type	Type 2 vs. Ref.	Type 1 vs. Ref.	Total
LS1			
Deletion	29 (29)	3 (3)	32
Insertion	34 (34)	0	34
SNP	79 (75)	13 (9)	88
LS2			
Deletion	25 (21)	7 (3)	27
Insertion	18 (17)	1 (0)	18
SNP	54 (54)	1 (1)	55
	239 (230)	24 (16)	254

Table 6. Variations found in *T. esculentum* mitochondrial gene sequences of the 84 individuals.

Position	Variant	Gene		Product
LS1 54113	Indel	*nad7*	Intron	NADH dehydrogenase subunit 7
LS2 2368	SNP	*matR*	Exon	maturase
LS2 35483	Indel	*nad4*	Intron	NADH dehydrogenase subunit 4
LS2 37645	SNP	*nad4*	Intron	NADH dehydrogenase subunit 4
LS2 39597	Indel	*nad4*	Intron	NADH dehydrogenase subunit 4
LS2 40927	SNP	*nad4*	Intron	NADH dehydrogenase subunit 4
LS2 41027	Indel	*nad4*	Intron	NADH dehydrogenase subunit 4
LS2 56879	SNP	*rps3*	Intron	ribosomal protein S3
LS2 69748	SNP	*nad2*	Intron	NADH dehydrogenase subunit 2
LS2 71388	Indel	*nad2*	Intron	NADH dehydrogenase subunit 2
LS2 71633	Indel	*nad2*	Intron	NADH dehydrogenase subunit 2

LS1 = OK638188, LS2 = OK638189

Table 7. Variations found in the 9,798 bp homologous segment within the organelle genomes of the 84 *T. esculentum* individuals.

Position	Variation	Type 1 Chloro	Type 1 Mito	Type 2 Chloro	Type 2 Mito	Localization
35570*	SNP	Ref	**Alt**	Ref	Ref	Mito
36347*	DEL	Ref	**Alt**	Ref	Ref	Mito
36942	SNP	Ref	Ref	**Alt**	Ref	Chloro
37257	SNP	**Ref**	Alt	Alt	Alt	Chloro
37813	SNP	**Alt** (some plants)	Ref	Ref	Ref	Chloro
38753*	DEL	Ref	Ref	Ref	**Alt**	Mito
38975	SNP	Ref	Ref	**Alt**	Ref	Chloro

39429*	SNP	Ref	**Alt**	Ref	Ref	Mito
40061*	SNP	Ref	**Alt**	Ref	Ref	Mito
40544*	SNP	Ref	**Alt**	Ref	Ref	Mito
41243*	SNP	**Ref**	Alt	Alt	Alt	Chloro
41716*	SNP	Ref	Ref	Ref	**Alt**	Mito
41718*	SNP	Ref	Ref	Ref	**Alt**	Mito
42587*	SNP	Ref	Ref	**Alt**	Ref	Chloro
43559	SNP	Ref	Ref	Ref	**Alt**	Mito
44059	INS	**Ref**	Alt	Alt	Alt	Chloro
44235	DEL	Ref	**Alt**	Ref	Ref	Mito
44302	SNP	Ref	**Alt**	Ref	Ref	Mito
44552	SNP	Ref	Ref	Ref	**Alt**	Mito
44762	DEL	Ref	Ref	Ref	**Alt**	Mito

Refs refer to alleles identical to the type 1 chloroplast reference genome sequence. Alt refers to an allele that differs from the type 1 chloroplast reference genome sequence. Only two alleles, Ref or Alt, were seen at each locus. Comparing alleles in the same row, the locations where the mutations occurred can be estimated and written in the last column. This table does not include differences between plastome and mitogenome at another 72 loci on this fragment (no inter-type differences were seen at these loci). Positions in the gene sequence are marked with an asterisk*.

Figure

Figure 1. The assembly graph of the type 2 mitogenome of *T. esculentum*. The type 2 mitogenome consists of three circular molecules M1, M2, and M4, and one linear molecule M3, and they were built on 21 long scaffolds (Table 1). Blue blocks show double-copy regions that are identical between two chromosomes, and orange blocks indicate triple-copy regions owned by three chromosomes. Close sequencing coverage was found for single-copy regions of the four chromosomes. Recombination on a pair of long inverted repeats I can change the junction of the upper and lower halves of M1 to that indicated by the yellow dashed arrows. The two structures before and after recombination have been confirmed by PacBio long reads, and their frequencies were close in the same individual (Figure S6-9). A long chloroplast insertion was found at the position of the gray segment M, about 9,798 bp, and its length varied slightly among different individuals. This chloroplast insertion and long repeats H, I, and O are also present in the type 1 mitogenome of *T. esculentum*. The type 1 mitogenome also has two rings, very similar to the M1 and M4 here, but other molecules have undergone dramatic changes.

Figure 2. Changes in sequencing coverage on the type 2 mitogenome chromosome M3 of *T. esculentum*. PacBio HiFi reads from the type 2 individual Sample 4 were aligned to the multicopy regions of the chromosome M3 using pbmm2. Chromosome M3 contains some long repeats that are identical to parts of other chromosomes, thus increasing the sequencing coverage of these regions in the alignment (Figures S10-12). At the positions of scaffolds C1, K1-O, and L2, with blue shading, the read depth was found to be doubled. At the location of scaffold H shaded in orange, the sequence depth was increased to 3-fold. However, a progressive decrease

in coverage indicated by red triangles was also seen at both ends of M3, as some linear M3 chromosomes had degenerated at both ends without telomere protection.

Figure 3. Step-by-step analysis of the structural differences between the two types of *T. esculentum* mitogenomes. i. The two autonomous circular chromosomes of type 1 *T. esculentum* mitogenome, LS1 (OK638188) (left) and LS2 (OK638189) (right). Colored arrows indicate the four pairs of long repeats (>1 kb). ii. Recombination on the direct repeats split the two large rings into five small circular molecules. Both conformations before and after recombination have been confirmed by PacBio reads to exist in type 1 individuals. iii. A rare recombination on a pair of *trnfM* genes at the junctions of D1 and D2, and E1 and E2, inverted the sequence D1-H-E1 in between. iv. Gaps with or without sequence deletions resulted in the three small circular chromosomes of the type 1 mitogenome present as different forms in type 2 individuals (Figures S13-18). v. New DNA fragments, including a 2,108 bp contig Q unique to type 2 individuals, joined originally remotely located sequences to form new structures. vi. The final type 2 mitogenome of *T. esculentum* with three circular and one linear chromosomes.

Figure 4. Synteny visualization of the two types of mitogenomes of *T. esculentum* by the R package RIdeogram after NUCmer alignment. The red triangle indicates the 2,108 bp type 2 mitogenome exclusive fragment of *T. esculentum*.

Figure 5. Homology analysis of the 2,108 bp fragment unique to type 2 *T. esculentum* mitogenome and design of primers for its PCR identification. (A) The 2,108 bp type 2 mitogenome exclusive sequence was blasted as a query in the NCBI database. Red horizontal bars indicate where database sequences are aligned, and separately aligned regions from the same database subject are connected by thin gray lines. (B) Two pairs of primers were designed to amplify products across both ends of the 2,108 bp fragment. The estimated size of the left end product is 990 bp, and the right end product is 289 bp. (C) Gel image of PCR amplification of DNA from six randomly selected samples with the two pairs of primers designed separately. The PCR products were electrophoresed on a 1.5% agarose gel at 80V for 1 hour.

Figure 6. The map of type 2 *T. esculentum* mitogenome gene arrangement drawn by OGDRAW. The annotation was performed by MITOFY and BLAST on the mtDNA of individual Index1 from UP Farm and deposited in GenBank under accession numbers OP795447-OP795450. All type 2 plants were found to have a similar gene arrangement. The dark gray pattern in the inner circle indicates GC content. Genes are colored according to their function. The decimal part after the gene name indicates the order of the exons. Genes inside the circle are transcribed clockwise, while those outside the circle are transcribed counterclockwise. Blue blocks represent two-copy regions that are identical between two chromosomes, and orange blocks show three-copy regions that are the same across three chromosomes.

Figure 7. Synteny block diagram of the Mauve alignment between the mitogenomes of *T. esculentum* and six other Fabaceae species, *Cercis canadensis* (MN017226.1), *Lotus japonicus* (NC_016743.2), *Medicago sativa* (ON782580.1), *Millettia pinnata* (NC_016742.1), *Glycine max* (NC_020455.1), and *Vigna radiata* (NC_015121.1). Similarities (percentage of marama mitogenome covered) are labeled above the bars. Genes contained in the synteny blocks are marked by gene symbols.

Figure 8. Synteny maps of the mitochondrial genomes of seven legume species. The colored bars represent the mitochondrial chromosomes of *C. canadensis* (MN017226.1), *L. japonicus* (NC_016743.2), *M. sativa* (ON782580.1), *M. pinnata* (NC_016742.1), *V. radiata* (NC_015121.1), *G. max* (NC_020455.1), and *T. esculentum*, which contains two chromosomes, LS1 (OK638188) and LS2 (OK638189). The gray ribbons indicate homologous sequences between the two neighboring species. The species were ordered according to the phylogenetic tree in Figure 9. Promer was used to detect syntenic regions between highly divergent genomes, which were then visualized by RIdeogram package in R.

Figure 9. Maximum Likelihood (ML) phylogenetic tree with the Jukes-Cantor model based on artificial chromosomes concatenated by 24 conserved mitochondrial genes, *atp1*, *atp4*, *atp6*, *atp8*, *atp9*, *nad3*, *nad4*, *nad4L*, *nad6*, *nad7*, *nad9*, *mttB*, *matR*, *cox1*, *cox3*, *cob*, *ccmFn*, *ccmFc*, *ccmC*, *ccmB*, *rps3*, *rps4*, *rps12*, and *rpl16* from *Arabidopsis thaliana* (NC_037304.1), *Cercis canadensis* (MN017226.1), *Lotus japonicus* (NC_016743.2), *Medicago sativa* (ON782580.1), *Millettia pinnata* (NC_016742.1), *Glycine max* (NC_020455.1), and *Vigna radiata* (NC_015121.1) in NCBI. The tree was drawn in Mega 11 after sequence alignment with Muscle v5. Percentage probabilities based on 1000 bootstrap replications are labeled on the branches. The topology was validated by the Bayesian inference phylogenetic tree drawn by BEAST (Figure S29).

Figure 10. Nucleotide matrices showing the distribution of mitochondrial genome variations in the 43 independent individuals and 4 additional samples of unknown origin. The first row indicates the alleles of the type 1 reference mitogenome of *T. esculentum* (LS1:OK638188 and LS2: OK638189). From the second row onwards, only bases different from the reference are shown, and bases identical to the reference are replaced by dashes. The two types of mitogenomes also differ from each other at another 170 loci, not shown here to save space (no within-type differences were found at those 170 loci). The full variation distribution is shown in Figure S30 and S31. All insertions are represented by the first two bases and deletions by the letter "D". The color bar to the left of the plant ID shows the source of the sample and is left blank for unknown sources.

Figure 11. Maximum Likelihood (ML) phylogenetic tree with the Jukes-Cantor model built on artificial chromosomes concatenated by 40 bp fragments at each of the 254 differential loci in the mitogenomes of *T. esculentum* according to the mitogenome sequences of the 43 independent individuals. Frequencies from 1000 bootstrap replicates were labeled on the branches with 40%

as cutoff. The topology was verified by the neighbor-joining method in Mega 11. Individuals with the same background color came from the same geographic location, and the sampling points were marked on the map of Namibia.

Figure 12. Distribution of SSR motifs of different repeat types in the type 1 reference mitogenome of *T. esculentum* analyzed by MISA. (A) Chromosome LS1 (OK638188). (B). Chromosome LS2 (OK638189).

Figure 13. Allele frequency plot of all differential loci between the two types of mitogenomes of *T. esculentum* in Aminuis individual A11. The x-axis of the left panel indicates the position on the chromosome concatenated by the reference marama mitogenomes LS1 (OK638188) and LS2 (OK638189).

Figure 14. Map of chloroplast DNA insertions in the mitochondrial genome of *T. esculentum* drawn by TBtools Advanced Circos. The reference mitogenome chromosomes LS1 (OK638188) and LS2 (OK638189) were aligned with the chloroplast genome (KX792933) of *T. esculentum* using BLAST (Figure S32; Table S2). The curves in the middle connect the homologous sequences of the plastome and mitogenome. The cpDNA insertions in the two mitochondrial chromosomes are colored orange and purple, respectively. cpDNA insertions were concentrated on the mitochondrial chromosome LS2, with only four short fragments on LS1. All the three chromosomes are circular but are shown here as linear, with numbers next to them indicating their length and orientation. The red triangle marks the position of the 9,798 bp long homologous fragment.

Figure 15. Amplification across the two ends of the 9,798 bp homologous fragment of the mitochondrial and chloroplast genomes in two random samples A and B. Lane1, HyperLadder II; Lanes 2-5, DNA from sample A; Lanes 6-9, DNA from sample B. The products in lanes 2 and 6 were amplified with primers MitoLL and MitoLR, in lane 3 and 7 were amplified with primers ChlLL and MitoLR, in lanes 4 and 8 were amplified with primers MitoRL and MitoRR, and in lanes 5 and 9 were amplified with primers MitoRL and ChlRR. The gel was run on a 1.5% agarose gel for 40 min at 100V. The annealing temperature of 55 ° C was a bit high for the primer ChlRR, resulting in low yields and faint bands in lanes 5 and 9. Decreasing the annealing temperature by 1 °C and repeating the PCR resulted in normal amplification.

(A)

(B)

(C)

Left   Right

C   2108 bp type 2 unique   B1

990 bp -

289 bp -

Sample A | Sample B | Sample C | Sample D | Sample E | Sample F | Sample A | Sample B | Sample C | Sample D | Sample E | Sample F

2k Left   2k Right

LS1   LS2