Exploring the Spatiotemporal Influence of Climate on American Avian Migration with Random Forests

I. Avery Bick[1,2], Vegar Bakkestuen[1], Marius Pedersen[2], Kiran Raja[2], Sarab Sethi[3]

[1]Norwegian Institute for Nature Research, Oslo, Norway
[2]Department of Computer Science, Norwegian University of Science and Technology, Gjøvik
[3]Department of Life Sciences, Imperial College London, London, UK

Abstract

North and South American Birds have adapted to climatic and ecological patterns to inform their

Spring and Fall migration timings. Temperature and precipitation patterns are shifting under

anthropogenic climate change, causing downstream effects on plant flowering cycles, insect

populations, and habitat availability. Understanding how these cues trigger migration could

improve the effectiveness and timing of bird surveys, as well as informing habitat protection and

creation efforts to lessen biodiversity loss due to climate change. Here, we employ a modeling

approach to explore how climate spatiotemporally affects accuracy of predicting bird

occurrence. Specifically, we train an ensemble of random forests on subsets of North and South

American climate data to predict distributions of historical bird occurrence probability for

passerine bird species in a North American forested region on eBird citizen science surveys

from 2008-2018. We further investigate the relevance of each feature, region, and temporal lag

for predicting the observed bird occurrence in a forested region in Northeast America, finding

that both temperature and precipitation facilitate accurate prediction. For predicting species in

October, when many passerines have begun their southward winter migration, we achieve more

accurate predictions of bird occurrence using lagged, rather than current, climate features alone

to predict communities in October. We also note significant higher random forest feature

26 importance for some lagged North American climate features than South American features.

27 These results suggest that machine learning models may be useful for identifying

28 spatiotemporal climatic cues that affect migratory behavior. Lastly, we explore the application

29 and limitations of random forests for prediction of future bird occurrence using 2021-2040

30 climate projections.

Introduction

Across the globe, birds undertake Spring migrations to breeding grounds and Fall migrations to

winter in warmer climates. In the Western Hemisphere specifically, many bird species migrate

northward from their winter homes in the Southeastern USA, the Caribbean, Mexico, and

Central America to breeding grounds in the Northern forests of Canada and the United States

each Spring, then again southward in the Fall (1,2). Bird species rely on climatic and ecological

conditions such as temperature, pressure, precipitation, and food availability to trigger the

timings of their migrations (3–6) in addition to their endogenous rhythms (7–9). However, these

climatic and ecological triggers are shifting with climate change. In North America, these are

projected to be characterized by warming of northern breeding regions, increased rainfall during

Spring migration, and less rainfall in southern regions during non-breeding season (10).

The broad effects of climate change on migrations have already been demonstrated: shifts to

earlier Spring first arrival dates (2,6,7,11,12), smaller and more dispersed migratory cohorts (8),

and a northward shifts in breeding grounds (13). Bird species will have to adapt behavioral

patterns more quickly to these habitat and climatic changes than has been done before

anthropogenic climate change began (2). However, the sensitivities to environmental and

ecological cues have been genetically shaped by natural selection to correctly time behaviors

such as egg laying to the availability of food, and species often over- or under-correct their

timings (5,14). Understanding the fine details of these shifts in behavior is key to conserving

biodiversity through the era of climate change. For example, the optimal times and places to

conduct bird surveys may shift. Additionally, understanding the climatic and ecological factors

underlying changing bird migration behavior could support habitat protection and creation.

Many studies have explored relationships between migration timings and climatic features

through linear-based models. For example, linking warmer Spring temperatures in the

Northeastern USA to earlier arrivals (7,8), or demonstrating effects of local weather on autumn

migration timing and stopovers (15,16). The literature demonstrates clearly that changes in bird

migratory behavior are correlated with, and to some degree explained by, the changing climate.

Still, many papers (7,8,16) show trends from post-hoc analyses and do not directly use climatic

features to predict how migratory patterns may change. Machine learning methods such as

random forests can leverage non-linear and complex trends in ecological data (17–19) to model

historical trends and make predictions of future behavior. Random forest models are high-

performing, efficient to train compared to neural networks under limited data, have built-in

feature importance metrics, and have been applied in ecological settings to relate climate and

ecological features to biodiversity (18,20) and animal movement and migration (19).

Here, we seek to build upon the existing literature with a predictive modeling approach for

species occurrence and migration patterns. Specifically, we use a random forest multi-

regression to predict eBird (21) checklist occurrences of common migrating passerine species in

a large North American forested region, using monthly averages of climate features (maximum

temperature, minimum temperature, and precipitation) across North, Central, and South

America as feature sets. By repeating this analysis across lags in the features, we explore

which regions and times are critical for predicting species Spring arrival and Fall departure

dates in North American forests. Through this approach, we make use of model error and

feature importance metrics to explore which features, at which times and places, are most

useful for prediction of migration. We demonstrate that random forest regressors can accurately

predict passerine species occurrence, even when presented with limited training data. We also

show that lagged temperature and precipitation features are as predictive, and often more

predictive, of species occurrence probability than non-lagged features. Finally, we train a

random forest regression model on historical data, then project future checklist occurrences

using several climate models projections averaged from 2021-2040, comparing our results to

the general literature consensus of earlier species Spring arrival times and later Fall departure

times due to climate change. The data and Python notebooks required to run these analyses

are available online (https://doi.org/10.5281/zenodo.8412210).

Methods

eBird Data

We used the eBird Basic Dataset from 2008-2018 (21) to derive the monthly occurrence rate of

bird species across surveys in our study area. The dataset consists of checklists of observed

species from citizen scientists, representing an "encounter rate" metric for each target species,

which can be thought of as the product of that species' detectability and occurrence probability

(22). We implement several data pre-processing steps to limit spatiotemporal variability in

species detectability to create an "apparent distribution" (22) of encounter rate that resembles

actual occurrence probability as much as possible: we restrict surveys to forested areas, only

consider the most common migrating passerine species, and drop high-effort surveys. To

account for increases in eBird usage throughout the study period, we also subsample surveys

(23) annually.

Checklists were first filtered to our study area, the Level 2 Ecoregion "Atlantic Highlands" within

the Level 1 "Northern Forests" Ecoregion (Fig 1) (24). These Environmental Protection Agency

(EPA) Ecoregions seek to aggregate areas of similar plant and animal species, as well as

abiotic characteristics such as water availability (25). The "Northern Forests" ecoregion is

characterized by extensive boreal forests and lakes (26), as well as extensive coverage by

eBird surveys. This spatial aggregation allowed us to limit the number of climate predictor

variables in our random forest models and thus more easily study the temporal effects of

regional climate on migration.

Checklists were further processed according to methods recommended by Johnston shown to

improve the performance of species occupancy and detection models when using eBird data,

especially in models with large survey sample sizes (22). Specifically, we included only

'complete' checklists, removing any surveys where an unknown species was observed, and

thus retaining information on species absence. Surveys with very high effort, specifically those

lasting over three hours and traveling more than 5 km, as well as area counts, were excluded to

retain spatial information of observations. Group surveys, which contain redundant checklists,

were condensed to a single checklist.

We next limited the species of interest to the 50 most common passerine species, as such birds

in general have a lower body mass and higher cold sensitivity, which can induce migratory

behaviors (27). Of these species, we filtered down to 41 species (mean observations: 3,571,

std: 2,044) (Table S1) which are either short-distance or long-distance migrants rather than

resident species, based on distribution maps from Birds of the World (28) and expert opinion. To

further refine our analysis to only forested regions, we filtered out all checklists with less than

50% surrounding tree cover based on 100m resolution tree cover density data (29). From 2008

until 2018, the number of annual surveys in eBird has greatly increased. To avoid bias in

inferred species richness in later years which have a greater number of surveys, we randomly

subsampled 1601 surveys at the annual scale, the number of surveys in 2008 after filtering for

forest cover and survey type (Fig S1). We then calculated the percentage of surveys on which

each species appears on a monthly scale to create our target features.

129   Climate Data

To study historical patterns in migration and climate, we utilized WorldClim monthly historical

raster data (30) downscaled with WorldClim 2.1 (31) for monthly minimum temperature,

maximum temperature, and precipitation at 2.5 minute spatial resolution across North and South

America. The raster data were averaged into EPA Level 1 Ecoregions for North and South

America (Fig 1A) that fall within typical migration pathways for birds that spend their breeding

season in the eBird study region. Features were then individually normalized by setting a mean

of 0 and scaling features to unit variance. Each of the three feature sets (maximum temperature,

minimum temperature, and precipitation) had a total of 15 features per month, from eight North

American Ecoregions and seven South American Ecoregions. One feature set containing all

three climate features was also created, with 45 features in total.

Figure 1: Map of targeted EPA Ecoregions and eBird survey locations. a) North and South American Level 1 Ecoregions for averaging of climate variables. b) The eBird study region Ecoregion 5.3, a subset of Level 1 Ecoregion 5 (NA5), outlined in light green with individual eBird survey coordinates in black. c) Data flow through random forest model ensembles. Monthly climate features are aggregated to the Level 1 Ecoregion spatial scale and monthly eBird species occurrence fractions across checklists are calculated for the Ecoregion 5.3 study region. An ensemble of ten models is trained and tested for each month and climate feature combination to predict checklist occurrence fractions, and the results of the ensemble are then averaged.

Similarly, to generate projected climate features, we utilized WorldClim monthly projected data

from the Coupled Model Intercomparison Project Phase 6 (CMIP6) (31), aggregated and

averaged to the same EPA Level 1 Ecoregions. The CMIP6 framework forms the standard for

international climate assessments, such as those from the Intergovernmental Panel on Climate

Change (32) and includes an ensemble of climate models from various research groups that

rely on common historical training data. For each CMIP6 model, we also tested several radiative

forcing pathways, dubbed the Shared Socioeconomic Pathways (SSP) (33), which compared

the effects of various greenhouse gas emissions scenarios. Due to our use of a standard

random forest regressor, which averages across previously seen training samples to make

predictions and cannot extrapolate, our projections of species occurrence fall within the bounds

of observed species occurrences on eBird checklists from 2008-2018. Still, in order to avoid

projecting species occurrence rates in radically different climatic conditions, we limited our study

to the 2021-2040 averaged WorldClim datasets. This consists of the mean projected monthly

minimum temperature, maximum temperature, and precipitation from 2021 to 2040. We used

four CMIP6 models available from WorldClim which performed best on tests comparing

probability distributions of projections to historical observations and the regularity of climate

fluctuations (34).

Analysis of Historical Avian Migration

We trained Random Forest regressors on the historical climate features to predict the eBird

checklist target features (Fig 1C). Twelve ensembles of models were trained in total, one for

prediction of eBird checklist occurrences in each month of the year. This separate training of

models for each month of the year and lag on the climate feature dataset enabled us to study

seasonal patterns in the error rates and feature importance. Each ensemble of models consists

of ten 1000-tree Random Forest models trained with varying 30:70 test-train splits across the

ten years of data being studied (2009-2018). For example, the first model was trained on 2009,

2010, and 2011 data, and tested on 2012-2018 data, the second model was trained on 2010,

2011, and 2012 data, and tested on 2013-2018 and 2009 data, and so forth. Each model is

trained on 7 years x 15 aggregated climate features, or 7 x 45 when using all temperature and

precipitation features. This process was then repeated while lagging the climate features from

one to six months. For example, the climate features from March were used to train models to

predict the checklist occurrences in April, and so forth.

From each group of ten models (an ensemble), we recorded the mean and variance of the test

set predictions and mean absolute errors for all species. We also extracted random forest

feature importance for each climate feature and Ecoregion. The features ranking highest in

importance were those which best split the distribution of target features across all nodes in the

decision trees. For a regression problem such as predicting checklist occurrences, this

decrease in impurity is related to the reduction of variance that each feature provides when

splitting training set samples. These forest importance metrics are relative, and sum to one (35).

As these ensembles of models generate distributions of error and feature importance, we were

able to compare their performances in Kolmogorov-Smirnov tests. These statistical tests can

evaluate whether one distribution is significantly greater or less than another, allowing us to test

whether certain features facilitate better model performance based on their errors, or if certain

groups of features have higher importance.

Projections of Avian Migration Shifts

We trained scikit-learn random forest multi-regression models (each with 1,000 trees) on the

combined data set of 45 climate features aggregated to the Ecoregion scale (minimum

temperature, maximum temperature, and precipitation) to predict the target eBird data from

2009-2015. Validation was performed on 2016-2018 data (Fig S4). We then used this model to

project on the 2021-2040 monthly average climate projections from the four CMIP6 models (EC-

Earth3-Veg, INM-CM4-8, MPI-ESM1-2-HR, and GISS-E2-1-H.) and four SSP projections (SSP

1, 2, 3, and 5). As with the historical ensemble of models, a separate model was trained to

predict each month of the year. Projected species-wise shifts in monthly occurrence probability

were calculated by subtracting the mean monthly eBird checklist occurrences (2009-2018) from

the mean projected occurrences for each of the four CMIP6 models and four projections.

Results

We found accurate per-species predictions of eBird checklists for the 41 passerines studied (Fig

2A) were made with monthly mean climate data (Fig 2D) as well as with lagged climate data up

to 6 months (Fig 2F). Average monthly errors in predictions were generally low across all

months and across climate features, ranging from 1.7 to 3.8 percent error in eBird checklist

occurrence fraction averaged across all species (Fig 3, Fig S2), with higher errors in the Spring

and Summer months when species richness was also higher. We noted a trend in the mean

annual error across species where the model tended to under-predict species occurrence

fraction in earlier years and over-predict species occurrence fraction in later years (Fig 3A-D).

This pattern in error followed a trend in the underlying eBird data; on average, each species in

the study occurred on 2.6% fewer eBird checklists from 2009 to 2018 (Fig 2C), a trend apparent

for most species (Fig 2 E, G).

Figure 2: Aggregated climate data supports accurate predictions of eBird species checklist occurrences - a) List of 41 common migrating passerine bird species observed in forested areas of the study region, representing the y-axis of the following heatmaps. b) Actual eBird checklist occurrence fractions for all species. c) Mean annual eBird checklist occurrence across all species (blue) and the mean annual error in occurrence fraction predictions (red) across all species. d) The Random Forest-predicted eBird checklist occurrence fractions for all species utilizing all available climate variables (minimum temperature, maximum temperature, and precipitation), e) eBird checklists minus the predicted checklists using current climate feature sets, showing over-predictions in blue and under-predictions in red, f) predicted eBird checklist occurrences utilizing all climate variables lagged by six months, and g) eBird checklists minus the predicted checklists using six month lagged climate variables.

Figure 3 illustrates the accuracy of the random forest ensembles for prediction of species

occurrence fractions, but also how the temporal trends in the underlying eBird features impacted

model performance. The model increasingly tended to overpredict individual species checklist

occurrence towards later years in the analysis (Fig 3 C, D). Following this, the derived species

richness was also increasingly over-predicted (Fig 3 A, B). Figure 3E and 3F focus on April and

October specifically, as these months are transitory periods between breeding and winter

seasons (Fig 2B) when migratory species are arriving and departing our study area. We noted

higher volatility and no apparent trend in model error for April (Fig 3E). In October, however, the

lowest error occurred when climate feature sets were lagged by one to three months (Fig 3F).

We performed Kolmogorov-Smirnov tests on the distributions of errors across the model

ensembles to determine whether the lower error of these lagged datasets was significant. After

correcting for multiple hypotheses, we found that lagging climate features for two and three

months produced significant reductions in model MAE for predicting eBird checklist occurrences

in October (p=0.00981) (Table S2). No such significant changes in error were found for lagged

datasets in April. We applied Kolmogorov-Smirnov tests between distributions of random forest

feature importance of the North and South American ecoregions (Fig S5) to determine if the

model was identifying relevant regional features. For predicting checklist occurrences in October

in our northeastern study region, we found that North American climate features were

significantly more important only at a lag of 3 ($p < 0.001$) (Table S2). No significant effects were

seen for importance in predicting April occurrences.

Figure 3: Model predictions across years reveal underlying shifts in eBird species occurrence and strength of lagged climate data for predictions. a) 2009 and b) 2018 eBird species richness versus mean and standard deviation of predicted species richness across the model ensemble. Species richness shown includes all species appearing on more than two percent of actual and predicted eBird checklists. c) 2009 and d) 2018 eBird ground truth versus predicted checklist occurrence fraction, showing mean and standard deviation of predictions across the model ensemble. e) Average random forest mean absolute test set errors by month of feature lag for April and f) October across all species and climate features.

By training on the 2008-2018 eBird checklists with all climate features and testing on CMIP6 climate projections (making broad assumptions of no change in habitat or species abundance), we projected higher positive shifts in checklist occurrence probability for April through August (ranging from 0.36% to 0.75%), with less change projected for Fall months (ranging from 0.011 to 0.19%) (Fig 4A). We found an increase in species occurrence probability across models projected for 25 of 41 species in April (mean: 0.80%), 15 species having a projected decrease (mean: -0.36%), and one species with no change (Fig 4B). Results were more mixed for October, with projected occurrence probability increases for 20 of 41 species (mean: 0.67%), decreases for 19 (mean: -0.46%), and no change for two species (Fig 4C). In contrast to the variation within eBird training data, the variance of occurrence probability for all species was

effectively zero across all four CMIP6 models and four SSP projections tested, indicating that

the random forest model consistently reached the bounds of known distributions of checklist

occurrences across the available climate training data.

Figure 4: Projections of eBird checklist occurrences from 2021-2040 show earlier Spring
occurrence probability for passerine species - a) Violin plot showing mean projected shifts of
occurrence probability for 41 passerine species in 2021-2040 compared to historical eBird
monthly means from 2008-2018 in Ecoregion 5.3 b) April and c) October per-species difference
between mean historical eBird occurrence probability and mean 2021-2040 projected occurrence
probability.

Discussion

In this section, we begin by highlighting the effectiveness of aggregated climate features from

both North and South America in accurately predicting eBird checklist occurrence probability for

41 passerine species in our North American forest study region, as well as species richness

(depicted in Fig 2B and Fig 3 A, B). Importantly, the accuracy of our predictions remained

consistent regardless of the specific climate variable chosen as a training feature (as illustrated

in Fig 3E, 3F, and Fig S2). Additionally, combining multiple climate features resulted in reduced

variability in prediction error (as indicated in Fig 3E, F). Furthermore, we observed that the trend

in mean annual model error closely mirrored the trend in mean annual checklist occurrence over

time across all species (as shown in Fig 2C). This alignment in error trends might be attributed

to underlying patterns of decreasing biodiversity and total bird counts observed during our study

period, a phenomenon that has also been documented in previous studies (36,37). It's worth

noting that potential errors could also arise from systematic shifts in the accuracy of the

underlying eBird data. However, we attempted to mitigate this risk by using randomly selected

complete checklists, as recommended by Johnston et al. (2021).

During October, a period when species are actively migrating southward from our study region

(as depicted in Fig 3A), we observed a notable reduction in prediction error when climate

features were lagged by two or three months, in comparison to using current climate features

(as illustrated in Fig 3B, Table S2). The results suggest that our model ensembles may be

effectively capturing delayed temporal patterns in climate that influence migratory departure.

Furthermore, in October, the random forest analysis highlighted that North American climate

features were significantly more influential than South American features at a lag of three

months, indicating that the model may be discerning spatial climate cues contributing to

migratory behavior. Conversely, in April, when birds are arriving in our Northern study area from

their Southern wintering grounds (as shown in Fig 2B), the model did not identify any statistically

significant temporal or spatial differences in prediction error or feature importance.

These findings align with existing research that highlights the influence of time lagged effects of

climate on migration triggers (14,38). Our projections of bird species from 2021-2040 were not

designed to model future changes in biodiversity and population abundance. Instead, they serve

the purpose of exploring how bird species might respond if sudden shifts in climatic conditions

were to occur in the near future, such as within the next year. Essentially, we aimed to uncover

insights from historical climate and bird migration patterns that could shed light on potential

future trends. Furthermore, it's important to note that our approach utilizes a random forest

model architecture. As a result, our projections of future species occurrence probability remain

constrained within the bounds of the historical training data, rendering them inherently

conservative. These projections do not account for factors like changes in land-use and

development (10), alterations in the timing of resource availability related to insects and

flowering plants (14,38–41), or shifts in inter-species competition (42), all of which affect bird

populations and distributions. These aspects are known to exert significant influences on bird

populations and distributions.

Despite these limitations, we were still able to discern potential future trends in the timing of bird

arrivals and departures, consistent with existing literature and empirical research. Notably, our

findings indicate an increased occurrence of species arriving in North American forests during

the spring (as depicted in Fig 4B). However, the trends for autumn months were more varied (as

shown in Fig 4C), aligning with previous studies (12,16). Several aspects of our results suggest

a higher sensitivity to climate and greater volatility in the Spring migration than the Fall

migration. This includes the higher test set error observed in April compared to October (Fig 3 E,

F) and the more pronounced and extensive shifts in projected future species occurrences in

April and May than in Fall months (as displayed in Fig 4 A-C). It is important to note that test set

error for the climate projection model exceeded the training set error (covering the period from

2008 to 2015) and displayed an upward trend from 2016 to 2018 (as presented in Figs S6, 7).

This trend culminated in a monthly maximum mean absolute error of 5.3% for August 2018.

The monthly scale of data resolution and the resulting small dataset size in this analysis

constrained our ability to employ more deep learning techniques, like recurrent neural networks,

which potentially could capture nuanced temporal influences of climate on migration.

Nonetheless, the utilization of random forests decision trees offered several advantages,

including built-in feature importance methods and the ability to confine future predictions within

the bounds of the training dataset. This contrasts with simpler models such as multiple linear

regression, which might produce projections outside the historical range.

For future research, there are several avenues to explore for enhancing our understanding of

bird migration in relation to climate. One promising direction involves improving the temporal

and spatial resolution of the climate datasets, which could enable machine learning models to

provide a more comprehensive view of climate influence on migration patterns. Additionally,

focusing on factors related to population abundance might yield valuable insights, particularly

considering observed decreases in bird populations. Furthermore, incorporating other climate

features that are known to affect migrations such as wind speed and direction (as highlighted by

(16)) could contribute to a more comprehensive model. Finally, the inclusion of indices derived

from satellite data such as normalized difference vegetation and water indices, which are

associated with food availability and drought conditions (as mentioned by (18,43)), could

provide a more holistic understanding of the ecological factors impacting bird migrations.

355   Conclusions

In this study, we present a modeling approach to identify spatiotemporal climate features that

influence bird migratory behavior. Our results demonstrate that random forest models trained on

these features accurately predicted eBird checklist occurrence probability throughout the year in

a northern forested region. Our findings indicate that lagged feature sets were more effective in

predicting passerine species occurrence probability in October compared to current climate

conditions. The models also assigned greater spatiotemporal importance to North American

features over South American features in October, coinciding with the southward migration of

species to their wintering sites.

Furthermore, we explored potential future trends in migratory behavior using mean climate

projections from CMIP6 for the period 2021-2040. Our results suggest an increased occurrence

probability in Spring and Summer months, but a reduced effect in the Fall and Winter. These

findings align with existing studies, supporting the hypothesis of a shifting of migratory timings

with climate change, characterized by earlier spring arrivals and later autumn departures. Our

pilot tests demonstrate that thorough investigation into model performance and feature

importance within machine learning models can contribute to our understanding of complex

ecological processes.

373   Acknowledgements

We acknowledge the World Climate Research Programme, which, through its Working Group

on Coupled Modelling, coordinated and promoted CMIP6. We thank the climate modeling

groups for producing and making available their model output, the Earth System Grid

Federation (ESGF) for archiving the data and providing access, and the multiple funding

agencies who support CMIP6 and ESGF.

Data preprocessing computations were performed on resources provided by Sigma2 - the

National Infrastructure for High Performance Computing and Data Storage in Norway.

SS was partially funded by a Herchel Smith Postdoctoral Research Fellowship.

IAB was funded by doctoral research grant 323294 from the Research Council of Norway.

Author Contributions

385 IAB led manuscript writing and methodology design and ran the analyses. VB, MP, KR, and SS

386 advised on methodology design and research questions and made significant contributions to

387 the manuscript. All authors reviewed and approved the manuscript.

Conflicts of Interest

389 The authors have no conflicts of interest to declare.
390

References

1. Jahn AE, Levey DJ, Smith KG. Reflections Across Hemispheres: A System-wide Approach To New
World Bird Migration. The Auk. 2004 Oct 1;121(4):1005–13.

2. Faaborg J, Holmes RT, Anders AD, Bildstein KL, Dugger KM, Gauthreaux SA, et al. Recent advances in
understanding migration systems of New World land birds. Ecol Monogr. 2010 Feb;80(1):3–48.

3. Sutherland WJ. Evidence for Flexibility and Constraint in Migration Systems. J Avian Biol. 1998
Dec;29(4):441.

4. Lemoine N, Schaefer HC, Böhning-Gaese K. Species richness of migratory birds is influenced by global
climate change. Glob Ecol Biogeogr. 2007 Jan;16(1):55–64.

5.   Visser ME, Caro SP, van Oers K, Schaper SV, Helm B. Phenology, seasonal timing and circannual
rhythms: towards a unified framework. Philos Trans R Soc B Biol Sci. 2010 Oct 12;365(1555):3113–27.

6.   Koleček J, Adamík P, Reif J. Shifts in migration phenology under climate change: temperature vs.
abundance effects in birds. Clim Change. 2020 Mar;159(2):177–94.

7.   Marra PP, Francis CM, Mulvihill RS, Moore FR. The influence of climate on the timing and rate of spring
bird migration. Oecologia. 2005 Jan;142(2):307–15.

8.   Miller-Rushing AJ, Lloyd-Evans TL, Primack RB, Satzinger P. Bird migration times, climate change, and
changing population sizes: BIRD MIGRATIONS, CLIMATE, AND POPULATIONS. Glob Change Biol.
2008 Sep;14(9):1959–72.

9.   Åkesson S, Helm B. Endogenous Programs and Flexibility in Bird Migration. Front Ecol Evol. 2020 Mar
26;8:78.

10. La Sorte FA, Fink D, Blancher PJ, Rodewald AD, Ruiz-Gutierrez V, Rosenberg KV, et al. Global change
and the distributional dynamics of migratory bird populations wintering in Central America. Glob Change
Biol. 2017 Dec;23(12):5284–96.

11. Butler CJ. The disproportionate effect of global warming on the arrival dates of short-distance migratory
birds in North America. Ibis. 2003 Jul;145(3):484–95.

12. Mills AM. Changes in the timing of spring and autumn migration in North American migrant passerines
during a period of global warming: Passerine migration and global warming. Ibis. 2005 Mar
1;147(2):259–69.

13. La Sorte FA, Thompson FR. Poleward Shifts In Winter Ranges Of North American Birds. Ecology. 2007
Jul;88(7):1803–12.

14. Visser ME, Holleman LJM, Gienapp P. Shifts in caterpillar biomass phenology due to climate change and
its impact on the breeding biology of an insectivorous bird. Oecologia. 2006 Feb;147(1):164–72.

15. Shipley JR, Kelly JF, Frick WF. Toward integrating citizen science and radar data for migrant bird
conservation. Williams R, Wegmann M, editors. Remote Sens Ecol Conserv. 2018 Jun;4(2):127–36.

16. Haest B, Hüppop O, Pol M, Bairlein F. Autumn bird migration phenology: A potpourri of wind,
precipitation and temperature effects. Glob Change Biol. 2019 Dec;25(12):4064–80.

17. Cutler DR, Edwards TC, Beard KH, Cutler A, Hess KT, Gibson J, et al. RANDOM FORESTS FOR
CLASSIFICATION IN ECOLOGY. Ecology. 2007 Nov;88(11):2783–92.

18. Hunt ML, Blackburn GA, Siriwardena GM, Carrasco L, Rowland CS. Using satellite data to assess spatial
drivers of bird diversity. Sankey T, Caravaggi A, editors. Remote Sens Ecol Conserv. 2022 Dec
24;rse2.322.

19. Wijeyakulasuriya DA, Eisenhauer EW, Shaby BA, Hanks EM. Machine learning for modeling animal
movement. Zhang J, editor. PLOS ONE. 2020 Jul 27;15(7):e0235750.

20. Martín B, Onrubia A, González-Arias J, Vicente-Vírseda JA. Citizen science for predicting spatio-
temporal patterns in seabird abundance during migration. Paiva VHR, editor. PLOS ONE. 2020 Aug
14;15(8):e0236631.

21. eBird Basic Dataset. Ithaca, New York: Cornell Lab of Ornithology; 2022.

22. Johnston A, Hochachka WM, Strimas-Mackey ME, Ruiz Gutierrez V, Robinson OJ, Miller ET, et al.
Analytical guidelines to increase the value of community science data: An example using eBird data to
estimate species distributions. Fourcade Y, editor. Divers Distrib. 2021 Jul;27(7):1265–77.

23. Fink D, Auer T, Johnston A, Ruiz-Gutierrez V, Hochachka WM, Kelling S. Modeling avian full annual
cycle distribution and population trends with citizen science data. Ecol Appl [Internet]. 2020 Apr [cited
2021 Nov 16];30(3). Available from: https://onlinelibrary.wiley.com/doi/10.1002/eap.2056

24. Level I and II Ecoregions of North America [Internet]. U.S. Environmental Protection Agency.; 2010.
Available from: ftp://ftp.epa.gov/wed/ecoregions/cec_na/NA_CEC_Eco_Level2.zip

25. Omernik JM, Griffith GE. Ecoregions of the Conterminous United States: Evolution of a Hierarchical
Spatial Framework. Environ Manage. 2014 Dec;54(6):1249–66.

26. Ecological regions of North America: toward a common perspective [Internet]. Montréal, Québec:
Commission for Environmental Cooperation; 1997. Available from:
http://ecologicalregions.info/data/cec_na/CEC_NAeco.pdf

27. Dawson WR, Marsh RL, Yacoe ME. Metabolic adjustments of small passerine birds for migration and
cold. Am J Physiol-Regul Integr Comp Physiol. 1983 Dec 1;245(6):R755–67.

28. Billerman SM, Keeney BK, Rodewald PG, Schulenberg TS. Birds of the World. Ithaca, NY, USA:
Cornell Laboratory of Ornithology; 2022.

29. Copernicus 2018 100m Forest Cover Fraction [Internet]. Copernicus Global Land Service; 2020.
Available from: doi:10.5281/zenodo.4723921

30. Harris I, Jones PD, Osborn TJ, Lister DH. Updated high-resolution grids of monthly climatic observations
- the CRU TS3.10 Dataset: UPDATED HIGH-RESOLUTION GRIDS OF MONTHLY CLIMATIC
OBSERVATIONS. Int J Climatol. 2014 Mar 15;34(3):623–42.

31. Fick SE, Hijmans RJ. WorldClim 2: new 1-km spatial resolution climate surfaces for global land areas. Int
J Climatol. 2017 Oct;37(12):4302–15.

32. Eyring V, Bony S, Meehl GA, Senior CA, Stevens B, Stouffer RJ, et al. Overview of the Coupled Model
Intercomparison Project Phase 6 (CMIP6) experimental design and organization. Geosci Model Dev. 2016
May 26;9(5):1937–58.

33. O'Neill BC, Tebaldi C, van Vuuren DP, Eyring V, Friedlingstein P, Hurtt G, et al. The Scenario Model
Intercomparison Project (ScenarioMIP) for CMIP6. Geosci Model Dev. 2016 Sep 28;9(9):3461–82.

34. Papalexiou SM, Rajulapati CR, Clark MP, Lehner F. Robustness of CMIP6 Historical Global Mean
Temperature Simulations: Trends, Long-Term Persistence, Autocorrelation, and Distributional Shape.
Earths Future [Internet]. 2020 Oct [cited 2022 Sep 14];8(10). Available from:
https://onlinelibrary.wiley.com/doi/10.1029/2020EF001667

35. Feature Importances with a Forest of Trees [Internet]. scikit-learn; 2023. Available from: https://scikit-
learn.org/stable/auto_examples/ensemble/plot_forest_importances.html

36. Anders AD, Post E. Distribution-wide effects of climate on population densities of a declining migratory
landbird. J Anim Ecol. 2006 Jan;75(1):221–7.

37. Sauer JR, Link WA. Analysis of the North American Breeding Bird Survey Using Hierarchical Models.
The Auk. 2011 Jan;128(1):87–98.

38. Evans TR, Salvatore D, van de Pol M, Musters CJM. Adult firefly abundance is linked to weather during
the larval stage in the previous year: Firefly abundance and weather. Ecol Entomol. 2019 Apr;44(2):265–
73.

39. Sparks TH, Jeffree EP, Jeffree CE. An examination of the relationship between flowering times and
temperature at the national scale using long-term phenological records from the UK. Int J Biometeorol.
2000 Aug 16;44(2):82–7.

40. Keatley MR, Fletcher TD, Hudson IL, Ades PK. Phenological studies in Australia: potential application in
historical and future climate analysis. Int J Climatol. 2002 Nov 30;22(14):1769–80.

41. Rawal DS, Kasel S, Keatley MR, Nitschke CR. Herbarium records identify sensitivity of flowering
phenology of eucalypts to climate: Implications for species response to climate change: Herbarium
Records Identify Phenological Shift. Austral Ecol. 2015 Apr;40(2):117–25.

42. Freeman BG, Strimas-Mackey M, Miller ET. Interspecific competition limits bird species' ranges in
tropical mountains. Science. 2022 Jul 22;377(6604):416–20.

43. Alessandrini C, Scridel D, Boitani L, Pedrini P, Brambilla M. Remotely sensed variables explain
microhabitat selection and reveal buffering behaviours against warming in a climate-sensitive bird species.
Pettorelli N, Kuemmerle T, editors. Remote Sens Ecol Conserv. 2022 Oct;8(5):615–28.

493
494