Proteostasis and lysosomal quality control deficits in Alzheimer's disease neurons

Ching-Chieh Chou[1],*, Ryan Vest[4,5,6], Miguel A. Prado[7,8], Joshua Wilson-Grady[7], Joao A. Paulo[7],

Yohei Shibuya[9,10], Patricia Moran-Losada[3,5], Ting-Ting Lee[1], Jian Luo[11], Steven P. Gygi[7], Jeffery

W. Kelly[12,13], Daniel Finley[7], Marius Wernig[9,10], Tony Wyss-Coray[3,5], Judith Frydman[1,2,3,14]*

[1]Department of Biology, [2]Department of Genetics, Stanford University, Stanford, California, USA

[3]Wu Tsai Neurosciences Institute, Stanford University School of Medicine, Stanford, California,
USA

[4]Department of Chemical Engineering, Stanford University, Stanford, California, USA

[5]Department of Neurology and Neurological Sciences, Stanford University, Stanford, California,
USA

[6]Qinotto, Inc. San Carlos, California, USA

[7]Department of Cell Biology, Harvard Medical School, Boston, Massachusetts, USA

[8]Instituto de Investigación Sanitaria del Principado de Asturias (ISPA), Oviedo, Spain

[9]Departments of Pathology, Stanford University School of Medicine, Stanford, California, USA

[10]Institute for Stem Cell Biology and Regenerative Medicine, Stanford University School of
Medicine, Stanford, California, USA

[11]Palo Alto Veterans Institute for Research, Inc. (PAVIR), Palo Alto, California, USA

[12]Department of Chemistry, The Scripps Research Institute, La Jolla, California, USA.

[13]The Skaggs Institute for Chemical Biology, The Scripps Research Institute, La Jolla,
California, USA.

[14]Lead contact

*Corresponding authors: cchou6@stanford.edu (C.-C.C.), jfrydman@stanford.edu (J.F.)

Summary:

The role of proteostasis and organelle homeostasis dysfunction in human aging and Alzheimer's

disease (AD) remains unclear. Analyzing proteome-wide changes in human donor fibroblasts

and their corresponding transdifferentiated neurons (tNeurons), we find aging and AD

synergistically impair multiple proteostasis pathways, most notably lysosomal quality control

(LQC). In particular, we show that ESCRT-mediated lysosomal repair defects are associated

with both sporadic and PSEN1 familial AD. Aging- and AD-linked defects are detected in

fibroblasts but highly exacerbated in tNeurons, leading to enhanced neuronal vulnerability,

unrepaired lysosomal damage, inflammatory factor secretion and cytotoxicity. Surprisingly,

tNeurons from aged and AD donors spontaneously develop amyloid-β inclusions co-localizing

with LQC markers, LAMP1/2-positive lysosomes and proteostasis factors; we observe similar

inclusions in brain tissue from AD patients and APP-transgenic mice. Importantly, compounds

enhancing lysosomal function broadly ameliorate these AD-associated pathologies. Our findings

establish cell-autonomous LQC dysfunction in neurons as a central vulnerability in aging and

AD pathogenesis.

Keywords:

Lysosome, Proteostasis, Transdifferentiated neuron, Aging, Alzheimer's disease, Proteomics

INTRODUCTION

Aging is the most prominent risk factor for AD(Alzheimer's Association, 2020; Harvey et al.,

2003; Qiu et al., 2009). Most cases are sporadic AD (sAD) and manifest late-onset symptoms.

In contrast, less than 5% of cases arise from familial AD (fAD) caused by pathogenic mutations

in causative genes with age of onset before 65 years.

The molecular bases of AD dysfunction are poorly understood. Given that accumulation of

amyloid plaques and neurofibrillary tangles in the brain are hallmarks of AD(DeTure and

Dickson, 2019), it has been proposed that loss of proteostasis is a central contributing factor.

AD has also been linked to collapse of mitochondrial and endoplasmic reticulum

homeostasis(Chakravorty et al., 2019; Nijholt et al., 2011), and autophagic pathology, including

the presence of dystrophic neurites with accumulations of lysosomes, in the brain of AD

patients(Gowrishankar et al., 2015; Nixon and Yang, 2011; Piras et al., 2016). Another less-

studied AD hallmark, granulovacuolar degeneration (GVD), is characterized by the presence of

membrane-bound vacuoles immuno-positive for endocytic vesicle and lysosome markers, such

as CHMP2B and LAMP1/2. The formation of GVD bodies is selective to pyramidal neurons and

associated with the early stages of amyloid-β (Aβ) and tau pathology in AD(Riku et al., 2019; Su

et al., 2002; Wiersma et al., 2019; Yamoah et al., 2020). Taken together, these lines of evidence

implicate defective proteostasis and/or organelle homeostasis as early pathogenic events

preceding overt neuronal loss in AD. However, whether and how proteostasis and organelle

homeostasis defects are directly linked to human aging and AD, and if so, whether they occur in

aged or AD neurons in a cell autonomous manner, remains unclear.

The maintenance of proteostasis requires many chaperones and protein quality control systems

that uphold the delicate balance between protein synthesis, folding and degradation(Hipp et al.,

2014). Additionally, the proteostasis network is critically required for the biogenesis, repair and

quality control of cellular organelles(Douglas and Dillin, 2010; Galluzzi et al., 2017; Kaushik and

Cuervo, 2015). The inter-relationship between protein and organelle quality control is

highlighted by the role of soluble proteostasis factors such as Hsp70 and the ubiquitin-

proteasome pathway in organelle quality control(Craig, 2018; Pohl and Dikic, 2019) as well as

the role of organelle functions, such as lysosomal degradation or mitochondrial energy

production, in proteostasis maintenance(Anding and Baehrecke, 2017; Gottschling and

Nyström, 2017). The inter-relatedness of these pathways likely renders cellular health

vulnerable to extensive dysfunction of a given proteostasis pathway or organelle, which would

predispose cells to damage accumulation over time.

While model systems have offered insights into the link between aging, amyloid disease and

disruption of proteostasis or organelle function (Cuervo and Wong, 2014; Douglas and Dillin,

2010; Hughes and Gottschling, 2012; Labbadia and Morimoto, 2015), studying the cell biology

of aging and AD in human systems has been extremely challenging. Since induced pluripotent

stem (iPS) cell reprogramming rejuvenates the epigenome, transcriptome and

proteome(Hansson et al., 2012; Mertens et al., 2015; Miller et al., 2013; Studer et al., 2015),

neurodegeneration models using embryonic stem cells or iPS cells to generate induced

neuronal systems (iNs) forgo the key contribution of aging to AD phenotypes. In contrast, direct

transdifferentiation of human adult somatic cells into neurons (herein tNeurons since iNs is also

used to refer to iPS cell-derived neurons) enables the preservation of aging signatures in

cellular models(Huh et al., 2016; Kim et al., 2018b; Mertens et al., 2015; Pang et al., 2011;

Vierbuchen et al., 2010). In recent years, specific tNeuron subtypes have been generated by

independent laboratories for modeling neurodegenerative diseases(Caiazzo et al., 2011;

Mertens et al., 2021; Tang et al., 2017; Victor et al., 2018). tNeurons with characteristics of

cortical neurons derived from AD patients were shown to recapitulate sAD-related alterations in

transcriptome and metabolome(Mertens et al., 2021; Traxler et al., 2022). Hence, we

hypothesized that tNeurons are amenable to define how aging and AD change the proteome.

We here undertook the proteomic and cell biological characterization of proteostasis and

organelle homeostasis changes in AD. We obtained human adult fibroblasts from young, aged

and AD donors and used them to generate tNeurons. We then assessed how aging and AD

change the cellular proteome of fibroblasts and their derived tNeurons (Figure 1A). Cellular and

molecular characterization of neuronal phenotypes were extended and supported by histological

validation in post-mortem mouse and human brain tissue. Strikingly, we found that aging and

AD lead to substantial alterations in the proteomes and cell biological functions of both patient-

derived fibroblasts and tNeurons. Quantitative proteomics analysis indicated deficits in

proteostasis and organelle homeostasis, with dramatic changes in the endosome-lysosome

pathway. Our findings indicate aging and AD act synergistically to impair proteostasis and

lysosomal homeostasis. Notably, tNeurons were significantly more vulnerable than fibroblasts to

deficits in lysosomal repair and recovery from damage. Thus, AD tNeurons contained

constitutively damaged lysosomes, as well as Aβ and hyper-phosphorylated tau (pTau)

deposits; they also secreted pro-inflammatory factors and were susceptible to neuronal death.

Highlighting the centrality of lysosomal function to AD-linked phenotypes, pharmacological

rescue of damaged lysosomes in both fibroblasts and tNeurons alleviated cell-based aging and

AD pathologies. We conclude that human tNeurons provide a model for assessing cell-

autonomous aging and AD phenotypes and a powerful platform to explore the cell biology and

therapeutic targets of AD.

RESULTS

Proteostasis deficits in human fibroblasts derived from aged and AD donors

To develop cellular models for aging and AD, we collected fibroblasts from healthy young (age

27.2 ± 4.0) and aged donors (age 71.1 ± 5.5), as well as aged patients with sAD (herein

aged/sAD, age 67.7 ± 6.9) and fAD patients with PSEN1 mutations (herein fAD-PSEN1, age

41.3 ± 9.0) (Table S1). We observed a correlation of donor's age and disease status with age-

and disease-associated markers, including those previously described in AD brain

tissue(Mansuroglu et al., 2016; Nativio et al., 2018; Salminen et al., 2012). Compared with

fibroblasts of young donors, fibroblasts from aged and aged/sAD donors showed an increase in

DNA damage measured by γ-H2AX puncta (Figure 1B) and a global loss of epigenetic marker,

histone 3 lysine 9 trimethylation (H3K9me3) (Figure 1C). Histone H4 lysine 16 acetylation

(H4K16ac) became enriched during aging but reduced with AD (Figure 1C). This recapitulated

findings in a genome-wide assessment of histone modifications of the human brain, which

observed H4K16ac enriched in normal aging but reduced in AD(Nativio et al., 2018).

To assess if human fibroblasts exhibit aging- and AD-linked proteostasis defects, we examined

the accumulation of ubiquitin-positive puncta as well as autophagy receptor p62/SQSTM1. No

inclusions were observed under basal conditions, but the treatment with proteasome inhibitor

Bortezomib (BTZ) or autophagy inhibitor Chloroquine (CQ) led to a moderate increase in the

accumulation of ubiquitin and p62/SQSTM1 puncta in aged fibroblasts (Figures 1D and 1E).

Strikingly, the protein accumulations were significantly exacerbated in aged/sAD fibroblasts,

suggesting that proteostasis dysfunction and susceptibility to proteotoxic stress is more acute in

aged/AD cells than in normal aged cells.

Human fibroblasts reveal proteome-wide changes related to aging and AD

To understand how aging and sAD affect the proteome of primary human fibroblasts, we carried

out a global quantification of proteins in our donor set via multiplex Tandem Mass Tag (TMT)

labeling. A total of 7,314 proteins were quantified from each sample. The proteins with fewer

than two unique peptides were excluded, resulting in 5,747 proteins used for further analysis.

Hierarchical and differential expression analysis identified changes in relative protein

abundance across samples (Figures S1A and S1B) and top proteomic hits (Figure S1C). These

analyses revealed aging and sAD impact proteostasis. Top-ranked pathways impacted by aging

and sAD revealed by gene ontology (GO) (Figure S1D) and integrated network analysis

(Figures 1F and S1E), corresponded to endosome-lysosome function and repair, protein folding,

inflammation, and cytoskeleton remodeling. In proteome of aging fibroblasts, we observed the

up-regulation of endosome-lysosomal proteins targeted to endosomal-sorting complex (e.g.

CHMP6 and RAB43) and lysosomal degradation (e.g. CTSB and CTSL), and down-regulation of

proteins for forming the ferritin complex (e.g. FTL and FTH1) (Figure 1F and 1G). In the

proteome of aging/sAD fibroblasts, endosome-lysosomal proteins were down-regulated,

particularly those involved in the molecular machinery for regulating lysosomal pH (e.g. v-

ATPases: ATP6V0D1 and ATP6AP1). In addition, the aging/sAD proteome exhibited significant

changes in mitochondrial proteins (Fig. 1F). To better understand the divergent trajectories in

proteome changes in aging versus aging/sAD, we performed a cluster analysis based on the

similarity of protein expression across young, aged and aged/sAD fibroblasts (Figure 1H). The

similarity matrix identified three main cluster trajectories in proteostasis pathways changing

during healthy aging versus aging/sAD, which impacted most lysosomal proteolysis pathways

and subsets of protein folding and chaperone factors, as well as translation machinery

components. These data indicate that aging changes the cellular proteome to alter proteostasis

and organelle homeostasis pathways. Notably, aging with sAD exacerbates these proteome

changes. These findings are consistent with the progressive increase in vulnerability observed

from aged to aged/sAD fibroblasts to proteotoxic stress.

Enhanced proteostasis deficits in human tNeurons are linked to aging and AD protein

pathologies

We next transdifferentiated the primary fibroblasts from young, aged, aged/sAD and fAD-

PSEN1 donors characterized above into neurons. Introducing transcription factors Brn2, Ascl1,

Myt1l, and Ngn2 (herein BAMN factors) yielded a majority of tNeurons with cortical neuron

characteristics (Figures 2A and S2B and & Table S1), as previous studies reported(Pang et al.,

2011; Tanabe et al., 2018). At post-induction day (PID) 35 to 42, tNeurons expressed neuronal

markers Tuj1, MAP2, NeuN, Tau, GAP43 and Synapsin-1 (Syn-1) (Figures 2A and S2C). The

aged and aged/sAD donors showed a slight reduction in fibroblast-to-neuron reprogramming

efficiency and synaptic density identified by Syn-1 puncta (Figures 2A and S2C). Direct

conversion to tNeurons retained aging hallmarks, since aged/sAD tNeurons also exhibited the

age-related increase in DNA damage (Figure 2B), global loss of H3K9me3 and H4K16ac

redistribution (Figure 2C) observed in the donor's fibroblasts. We next examined whether

tNeurons exhibit aging- and AD-linked proteostasis deficits. Notably, even under basal

conditions, the tNeurons derived from aged and AD donors contained p62/SQSTM1 and

ubiquitin puncta (Figures 2D and S2D). This contrasted to their corresponding fibroblasts, where

these proteostasis deficits were only uncovered upon BTZ or CQ treatment (Figures 1D and

1E). Further, we observed significantly more p62/SQSTM1 puncta in both aged/sAD and fAD-

PSEN1 tNeurons than in aged tNeurons.

The above analyses suggest neuronal proteostasis is more vulnerable to aging and AD than

that of the parental fibroblasts. This led us to examine formation of AD-associated protein

pathologies in tNeurons (Figures 2D and S2D). Remarkably, aging tNeurons contained a higher

burden of intracellular total Aβ, the toxic isoform Aβ42, pTau, hyper-phosphorylated TDP-43

(pTDP-43) in tNeurons (Figures 2D and S2D). Strikingly, these pathological protein deposits

were significantly increased in tNeurons from aged/sAD donors (Figures 2D and S2D). The

clear progressive enrichment in these AD phenotypes between aged and aged/AD tNeurons,

suggests a cooperative contribution of aging to sAD dysfunction. The strongest enrichment in

protein pathology was observed in fAD-PSEN1 tNeurons, suggesting that genetic mutations can

have a great impact on neuronal proteostasis. Of note, TDP-43 pathology is the hallmark of

amyotrophic lateral sclerosis (ALS) and frontotemporal dementia (FTD)(Chou et al., 2018). The

increase in pTDP-43 and loss of nuclear TDP-43 in aged, aged/sAD and fAD-PSEN1 tNeurons

(Figures 2D and S2D) resonates with findings that TDP-43 pathology concurrence in 23-50% of

AD as a potential co-factor (Amador-Ortiz et al., 2007; Josephs et al., 2014) and suggests that a

cell-intrinsic deficit in proteostasis can lead to multiple protein pathologies. These experiments

uncover the increased fragility of neuronal proteostasis to aging and AD as compared to

fibroblasts and establish that aging and AD cooperatively promote cell-intrinsic accumulations of

multiple AD-linked protein pathologies, including Aβ, pTau and pTDP-43.

Quantitative proteomics uncovers how aging and AD alter the neuronal proteome

While transcriptomic changes in aged and AD derived neurons have been extensively

characterized(Grubman et al., 2019; Mathys et al., 2019; Raj et al., 2018), how these conditions

affect the proteome of patient-derived neurons has not been examined. We carried out

quantitative analyses of proteome changes in young, aged and AD-derived tNeurons at PID 40.

TMT-proteomics revealed a total of 6,015 proteins with more than two unique peptides, allowing

us to identify the top proteomic hits related to aging, aging/sAD and fAD-PSEN1 (Figures S3A

and S3B). Combining GO and integrated network analysis revealed the common, top-ranked

pathways altered by aging and AD, which include endosome-lysosome function, synapses,

chaperones, mitochondrial function, lipid signaling and inflammation (Figures 2E and S3C).

Comparing the proteomes of aged and young tNeurons identified age-related up-regulation of

proteins for modulating synaptic function (e.g. RAB27B and CADPS) and mitochondria (e.g.

HK2). In contrast, proteins for endosome-lysosomes (e.g. CLU, CD63, CTSC and TMEM175)

and chaperones (e.g. SACS) were mostly down-regulated with aging (Figure 2E). To evaluate

sAD-related changes, we compared the proteomes of aged/sAD and aged tNeurons and found

that the expression of proteins for endosome-lysosomes (e.g. CLU and TMEM175),

mitochondria (e.g. CHDH), inflammation (e.g. PYCARD and CRLF1) as well as synapses (e.g.

SORCS2) becomes enriched in aged/sAD tNeurons. We also observed down-regulation of

LGALS3BP and RAB27B, suggesting a deficit in membrane and vesicular trafficking in sAD

(Figure 2E). Since familiar PSEN1 mutations elicit early development of AD, we also examined

how fAD PSEN1 mutations affect the neuronal proteome versus that of aged/sAD derived

neurons. Comparative analyses showed that the components of the endosome-lysosome (e.g.

ABCA8, PLD3 and GALC), synapse (e.g. STON2) and membrane raft (e.g. AHNAK) pathways

are up-regulated in fAD-PSEN1. fAD-PSEN1 tNeurons also downregulated many mitochondrial

proteins, including mitoribosomes (Figure 2E). These proteomic analyses reveal that aging and

AD synergistically affect the architecture of the neuronal proteome by impacting organelle

homeostasis and proteostasis, thus suggesting a molecular rationale for the progressive

severity of amyloid and proteostasis phenotypes observed in aged and aged/sAD tNeurons.

Notably, proteins with changing expression in aged/sAD tNeurons were strongly associated with

the risk genes for AD (Figure 2F). This suggests that proteome changes in aged neurons may

synergize with risk gene mutations to bring about age-related neurodegenerative diseases.

Cluster analysis of our proteomic data further clarified how proteostasis pathways change across young, aged, aged/sAD and fAD-PSEN1 tNeurons (Figure 2G). Major Clusters implicated proteins collectively involved in proteolytic pathways, additional pathways corresponded to autophagy, amyloid processing and translation. Comparing young, aged, aged/sAD and fAD-PSEN1 tNeurons, we identified six proteolysis-related clusters of protein trajectories, which ranged in size from 5 to 30 proteins. Cluster A reflected the relative enrichment for proteins associated with fAD-PSEN1. Cluster B and C indicated the relative enrichment for proteins in aged/sAD tNeurons, such as PYCARD and CTSC, whereas Cluster D and F showed the relative enrichment for proteins in aged tNeurons, such as CTSK and ERAP1. Notably, proteins in the autophagy pathway showed distinct trajectories of change for aged/sAD. Cluster analysis of all protein whose abundance was altered across tNeurons categories identified twelve clusters that exhibited distinct, yet concerted changes in biological processes during normal aging and AD processes (Figure S3D).

Previous transcriptomic studies sowed that aging-related transcriptomic signatures can be shared by multiple cell types or specific to a single cell type(Zhang et al., 2021). We thus asked whether specific aging and AD proteome signatures are shared between the donor fibroblasts and their matching tNeurons. These analyses revealed an overlap of pathways and protein hits shared between human fibroblast and tNeurons but also cell-type-specific processes impacted by aging and AD. We found 44 shared protein hits in the aging proteome and 33 shared proteins in the aged/sAD proteome of both cell types (Figure 2H). For healthy aging, the lysosome, ER and cytoskeleton pathways were similarly affected in both cell types, whereas the axon and late endosome pathways were uniquely affected in neurons. For sAD, the lysosome and ER pathways were commonly altered in both cell types, whereas proteins involved in

postsynaptic density and mitochondria matrix were specifically affected in neurons. These

results reveal that aging and AD can broadly impact the proteome of different cell types, but

also induce specific neuronal deficits.

Defective responses to lysosomal damage emerge in aged and AD fibroblasts

Proteomic analysis of human fibroblasts indicated that aging and aged/sAD affect the

endosome-lysosome pathway (Figures 1F, 1G and S1E). Cluster analysis provided a deeper

understanding of molecular signatures in the endosome-lysosome pathway across young, aged

and aged/sAD fibroblasts. For instance, ferritin (Cluster A) decreased in aging, whereas proteins

involved in lysosomal acidification (Cluster B) and lysosomal proteolysis and LQC (Cluster C)

(e.g. CHMP6 and LGALS8) were enriched in aging but decreased in sAD (Figures 3A and 3B).

These analyses were consistent with our finding that aged/sAD cells are highly sensitive to

lysosomal stress whereby CQ treatment of fibroblasts lead to strong accumulation of

autophagosome-interacting protein LC3B in aged/sAD cells (Figure S4A).

Based on these analyses, we next employed a battery structural and cell biological assays to

characterize the impact of aging and AD on lysosomal function in patient fibroblasts.

Transmission electron microscopy (TEM) showed that even under basal conditions, individual

lysosomes in aged/sAD fibroblasts are significantly larger than those in aged fibroblasts (by

50.7%, Figure 3C). To investigate the cellular response to lysosomal damage stress, fibroblasts

were treated with L-leucyl-L-leucine O-methyl ester (LLOME), a lysosomotropic compound that

is commonly used to induce lysosomal membrane permeabilization(Aits et al., 2015; Maejima et

al., 2013). LLOME induced lysosomal enlargement in all groups of fibroblasts (Figure 3C).

However, immunogold detection in TEM of lysosomal damage marker protein Galectin-3

showed it was frequently detected upon LLOME treatment of young and aged fibroblasts, but

barely in aged/sAD fibroblasts (Figure 3D), suggesting the potential dysregulation of lysosomal

damage response in AD.

Protein aggregates and pathogens are known factors to trigger damage to the lysosomal

membrane (Figure 3E). Leakage of intraluminal contents from ruptured lysosomes activates cell

death pathways, which are strongly associated with neurodegenerative diseases(Nixon et al.,

2008). Several pathways of LQC were recently identified as cytoprotective mechanisms to

resolve damaged lysosomes to avert cytotoxicity(Radulovic et al., 2018; Skowyra et al., 2018).

Mildly damaged lysosomes are repaired by successive recruitment of ESCRT proteins

(ESCRTs), whereas if the damage is severe, glycan-binding proteins, Galectins, bind to

glycosylated proteins residing in the lysosomal lumen and promote lysosomal clearance by

autophagy (i.e. lysophagy). In young healthy fibroblasts, LLOME-induced lysosomal damage

led to the expected time-dependent increase in ESCRT-III protein CHMP2B and Galectin-3

accumulation on lysosomes, leading to characteristic punctate structures (Figure 3E).

Scatterplots correlating the immunostaining intensity for CHMP2B and Galectin-3 showed these

puncta increased during LLOME treatment but were subsequently dissipated and reduced after

washout of the LLOME, as expected for lysosomal repair from damage (Figure 3F). We then

examined if the lysosomal damage response was different between healthy and diseased

fibroblasts. In young fibroblasts, CHMP2B-stained puncta became enriched during LLOME

treatment and rapidly dissipated after the LLOME washout. In contrast, aged and fAD-PSEN1

fibroblasts formed significantly higher number of CHMP2B puncta during LLOME treatment and

retained them for a longer time after washout (Figures 3G and S4B). Importantly, LLOME

treatment of aged/sAD fibroblasts led to lower densities of CHMP2B puncta on lysosomes,

suggestive of an impaired lysosome damage response. Indeed, a defective lysosomal damage

response in aged/sAD fibroblasts was also detected for additional ESCRT subunits, including

ESCRT-I TSG101, ESCRT-IIII CHMP1A and CHMP6 and ESCRT-III accessory factor ALIX

(Figure S4C). Further indicating deficits in lysosomal repair, the dynamics of Galectin-3

recruitment to lysosomes was also affected in aged and AD fibroblasts. LLOME treatment led to

higher levels of Galectin-3 puncta in aged and fAD-PSEN1 fibroblasts compared to young

healthy fibroblasts, as well as much slower dissipation rates during LLOME washout (Figures

3G and S4B). Of note, the Galectin-3 response to LLOME damage in aged/sAD fibroblasts was

more anomalous since high-density accumulation of Galectin-3 puncta was observed only 1 hr

after the LLOME washout, accompanied by slow dissipation of Galectin-3-puncta. Taken

together, these results indicate that aged and AD fibroblasts have a greater vulnerability to

lysosomal damage and likely due to deficits in LQC and recovery pathways.

We next asked if the dysregulated LQC in aged and AD fibroblasts is associated with enhanced

cytotoxicity. To this end, we evaluated the induction of cell death following the treatment using

two lysosome-damaging agents, LLOME and silica nanoparticles, with different mechanisms of

action (Figure 3H). LLOME-induced lysosomal membrane permeabilization is mediated by

lysosomal protease Cathepsin-C, while silica nanoparticles can directly damage the lysosomal

membrane and activate NLRP3 inflammasomes independent of Cathepsin-C(Mulay and

Anders, 2017; Skowyra et al., 2018). As compared with young fibroblasts, a 24-hr mild LLOME

treatment induced a moderate increase in cell death in aged and fAD-PSEN1 fibroblasts but a

dramatically rise in cell death in aged/sAD fibroblasts. Similarly, uptake of silica nanoparticles

led to a moderate increase in cell death in aged fibroblasts but a substantial increase in

aged/sAD and fAD-PSEN1 fibroblasts. These results indicate that aged/AD fibroblasts, which

have pronounced defects in LQC, are more prone to lysosomal damage-mediated cell death. In

addition to ESCRTs and Galectins, other proteostasis pathways participate in lysosome repair;

for instance, calcium homeostasis and the chaperone Hsp70 display essential roles in ESCRT

assembly and lysosomal stabilization(Kirkegaard et al., 2010; Scheffer et al., 2014). We find that

inhibition of calcium signaling by calcium chelator BAPTA-AM or Hsp70 function by VER-

155008 significantly perturbed LQC, impacting both CHMP2B and Galectin-3 recruitment during

LLOME treatment and dissipation after LLOME washout (Figure S4D). Consequently, both

calcium chelation and Hsp70 inhibition enhanced cell death associated with lysosomal damage

(Figure S4E). These results highlight the importance of chaperone function for LQC and

resonate with previous links between lysosomal calcium signaling and Hsp70 function with

AD(Lackie et al., 2017; Lee et al., 2015; Tong et al., 2018).

We next investigated how LQC deficits in aged and AD fibroblasts affect lysosomal function. To

assess the extent of lysosomal leakage following damage in these cells we preloaded

lysosomes with LysoTracker Red DND-99. LysoTracker Red fluorescence was significantly

increased in aged/sAD fibroblasts, indicative of lysosomal enlargement. LLOME treatment led to

loss of fluorescence, indicative of dye leakage from lysosomes, in all groups as expected from

lysosomal damage (Figure 3I). Because leakage of lysosomal protons will reduce lysosomal

acidification and intraluminal hydrolase activity, we next evaluated how lysosomes maintain or

repair these functions during and after damage. We measured lysosomal acidity changes in

fibroblasts using pH-sensitive, fluorescein (FITC)-conjugated Dextran (40 kDa). FITC-Dextran is

self-quenched in an acidic environment and increases fluorescence intensity upon

deacidification due to lysosomal damage (Figure S4F). At basal conditions, no obvious

endocytic deficiency was observed in aged and AD fibroblasts, assessed by an uptake of

tetramethylrhodamine (TMR)-conjugated dextran (70 kDa) (Figure S4F). Lysosomes of aged

and AD fibroblasts exhibited mild deacidification at the basal conditions (Figure 3J). Upon

damage, we found that young fibroblasts better resist pH perturbation and rapidly restore acidity

after lysosomal damage; in contrast aged fibroblasts showed intermediate acidity loss while AD

fibroblasts showed substantial lysosomal acidity loss and poor acidity recovery after damage.

Since lysosomal hydrolases, such as Cathepsins, are optimally active at pH 4-5, we used

Cathepsin-B substrates to measure changes in lysosomal proteolytic activity during lysosomal

damage. As expected, Cathepsin-B activity was dramatically lost in the presence of LLOME

(Figure 3K). One hr after LLOME washout, young fibroblasts were able to recover ~88% of

Cathepsin-B activity, whereas aged and AD fibroblasts only recovered ~68% and 45% of

activity, respectively. While the Cathepsin-B activity of young and aged fibroblasts was almost

fully recovered 8 hr after LLOME washout, AD fibroblasts restored less than 80% of Cathepsin-

B activity. We also examined the localization of transcription factor EB (TFEB), which

translocates to the nucleus upon lysosomal damage and activates the regulatory gene network

for lysosomal biogenesis(Settembre et al., 2011). While this was the case in young fibroblasts,

we observed a deficit in TFEB translocation into the nucleus following lysosomal damage in

aged/sAD fibroblasts (Figure S4G), perhaps due to aberrant nucleocytoplasmic transport in

AD(Eftekharzadeh et al., 2018). Interestingly, the delayed recovery from lysosomal damage in

AD fibroblasts may lead to detrimental effects beyond lysosomal dysfunction, as suggested by

recent proteomics analyses of damaged lysosomes(Eapen et al., 2021; Kravić et al., 2022).

Indeed, we find LLOME-damaged lysosomes do not only recruit ESCRTs and Galectins, but

also proteins involved in proteostasis (e.g. Hsp70), RNA processing (e.g. TDP43) and nuclear

pore complex (e.g. Nup98) (Figure 3L). Of note, calcium chelation by BAPTA-TA suppressed

the recruitment of ESCRTs and TDP-43 to the damaged lysosomes, but unexpectedly

enhanced Hsp70 association with lysosomes (Figure S4H). These results provide unprecedent

insights into how aging and AD increase lysosomal fragility and impair lysosomal repair, which

could jointly promote failures in proteostasis, lysosomal function and cell survival in AD patient

fibroblasts.

Aging- and AD-linked deficits in lysosomal structure and repair are exacerbated in

tNeurons

We next examined how fibroblast-to-tNeuron lineage reprogramming affects endosome-

lysosomal function. TEM comparison of lysosomal ultrastructure for young versus aged versus

aged/sAD tNeurons revealed a progressive increase in the size of individual lysosomes within

the cell body and proximal neurites from young to aged to aged/AD (Figure 4A). In addition, we

observed small electron-dense granules proximal to the lysosomal membrane in aged tNeurons

and large electron-dense granules in the swollen lysosomes of aged/sAD tNeurons. Strikingly,

the enlarged lysosomes within the cell body of aged/sAD tNeurons were closely surrounded by

mitochondria, suggesting increased mitochondria-lysosome contacts in AD neurons.

We next examined if lysosomal recovery from damage is also impaired in AD tNeurons as seen

in fibroblasts. Notably, even under basal conditions, we observed significantly higher number of

CHMP2B and Galectin-3 puncta in the lysosomes of aged/sAD tNeurons (Figures 4B, 4C, S5A

and S5B), consistent with spontaneously damaged lysosomes. Damage markers were less

frequently observed in young, aged and fAD-PSEN1 tNeurons. This suggests aging and AD

combine to create a synergistic deficit in lysosomal homeostasis in tNeurons. Interestingly, a

portion of aged/sAD tNeurons also showed aberrant accumulations of CHMP2B adjacent to the

plasma membrane and neurite branch points (Figure S5A), consistent with reports that

CHMP2B can participate in the repair of both plasma and lysosomal membrane

damage(Jimenez et al., 2014; Radulovic et al., 2018; Skowyra et al., 2018). There was no

significant difference in neuronal apoptosis at the baseline between groups (Figure S5C).

To understand the dynamics of lysosomal repair, we monitored the change in ESCRT-0 HGS

during LLOME treatment and washout. We chose ESCRT component HGS because its

distribution was comparable across all groups at baseline (Figures 4D and 4E). During LLOME

treatment, the number of HGS puncta moderately increased in aged tNeurons compared with

young tNeurons but substantially increased in aged/sAD and fAD-PSEN1 tNeurons. Monitoring

repair after LLOME washout showed that HGS puncta rapidly dissipate in young tNeurons

returning to baseline levels after 2 hr. In contrast, aged tNeurons exhibited a moderate delay in

the dissipation of HGS puncta while AD tNeurons exhibited a severe delay returning to baseline.

Relative to aged tNeurons, the half-life estimate of HGS granule dissipation increased by ~3-

fold in aged/sAD and ~1.5-fold in fAD-PSEN1 tNeurons (Figure 4E). Of note, we observed

recruitment of TDP-43 and Hsp70 to damaged lysosomes in aged tNeurons, which was further

exacerbated in aged/sAD tNeurons (Figure S5D), illustrating that deficits in lysosomal repair can

disturb additional proteostasis pathways.

We next examined lysosomal function in aged and AD neurons. Under basal conditions,

aged/sAD tNeurons exhibited a slight loss of lysosomal acidification which was exacerbated in

the presence of LLOME (Figure 4F). The enhanced lysosomal damage and reduced repair

observed in aged/sAD tNeurons was accompanied by reduced Cathepsin-B activity, indicative

of compromised lysosomal proteolytic function (Figure 4G). Since we observed spontaneous

formation of aberrant Aβ42 deposits in aged/sAD tNeurons (Figures 2D and S2D) and previous

finding showed Aβ preferentially accumulates in poorly acidified lysosomes at early AD

stages(Lee et al., 2022), we examined the relation of Aβ42 deposits with lysosomes in

tNeurons. Interestingly, we found Aβ42 colocalized with LAMP1 and LC3B in aged/sAD

tNeurons (Figure 4H), consistent with a link between lysosomal damage and amyloid

accumulation. Our data suggests that accumulation of damaged lysosomes in aged and AD

neurons leads to an overload of ESCRT and proteostasis systems, thus diminishing cellular

resilience to stress, and leading to increased accumulations of amyloid deposits and

susceptibility to neuronal cell death in AD.

Lysosomal damage markers associate with amyloid pathology in brain tissue of AD

patients and mouse models

To extend our observations from aged and AD derived tNeurons to brain tissues, we carried out

histopathological analysis of post-mortem cerebral cortex of AD patients and mouse models. In

sections of AD patients, we observed a global and intra-neuronal elevation of LAMP2, CHMP2B

and Galectin-3 immunoreactivity as compared with age-matched healthy control (HC)

individuals (Figures 5A and S5E). We observed two distinct patterns of CHMP2B staining in the

AD brain. While CHMP2B primarily colocalized with LAMP2-positive lysosomes in the

perinuclear region of neurons, we also observed thin thread-like CHMP2B-positive structures

with variable length and devoid of LAMP2 positivity in patient cerebral cortex. We also observed

increased colocalization of Galectin-3 with LAMP2 in the neurons of AD patients (Figures 5A

and S5E). Of note, Galectin-3 is highly up-regulated in microglia of AD patients brains and

5xFAD mice and strongly associated with amyloid plaques(Boza-Serrano et al., 2019; Grubman

et al., 2021). Also consistent with our finding that Aβ42 colocalizes with lysosomes in AD

tNeurons (Figure 4H), we observed Aβ deposits associated with LAMP2-positive lysosomes in

the extra-cellular and intra-neuronal plaques using 6E10 antibody in AD patient's cortex.

To explore how aging contributes to aberrant lysosomal signatures in the brain, we conducted

histological analysis of Galectin-3 and LAMP1 in the neocortex of young (3-6 month) and aged

(20-24 month) mice (Figure S6A). In aged mice, we observed a moderate increase in Galectin-3

immunoreactivity which colocalized with enlarged LAMP1-positive lysosomes in neurons (Figure

S6B). Together with our tNeuron analyses, these data implicate aging-linked lysosomal deficits

as contributing early factors to AD prior to neurodegeneration. Therefore, we examined

lysosomal and proteostasis pathologies in a mouse model for AD(Faizi et al., 2012). Transgenic

mice expressing human APP carrying Swedish (K670N/M671L) and London (V717I) mutations

(APPLon/Swe) exhibited a widespread accumulation of LAMP1-positive lysosomes in the

neocortex as compared with the non-transgenic (NTg) mice (Figures 5B, S6A, S6C and S6D).

Aberrant lysosomal accumulation forming various sizes of protein clumps and colocalizing with

lysosomal damage markers, such as CHMP2B and Galectin-3 (Figures 5B and S6C) and

chaperone Hsp70 (Figure S6D), were observed in the perinuclear region of individual pyramidal

neurons as well asthe extra-cellular space (Figure S6E). Similar to our findings in AD tNeurons,

the buildup of Aβ42-containing amyloid plaques was observed inside these lysosome clumps

(Figure 5B). Extra-cellular Aβ depositions were associated with an increase in the size of protein

clumps immuno-positive for CHMP2B and Galectin-3, as well as the severity of brain damage

regions devoid of MAP2 staining (Figures 5B, S6F and S6G). Of note, the pyramidal neurons of

APPLon/Swe mice proximal to Aβ plaques also contained intra-neuronal Aβ42 deposits within

lysosomes (Figure S6H). Taken together, the global increase in immunoreactivity of lysosomal

damage markers accompanying with co-aggregation with amyloid plaques and neuritic

degeneration suggest a critical role of lysosomal damage in the development of AD

neuropathology.

Lysosomal damage potentiates inflammatory activation in AD tNeurons

Our tNeuron proteomic datasets revealed that inflammasome adaptor protein PYCARD, also

referred to as ASC is up-regulated in aged/sAD tNeurons (Figure 6A). Inflammasomes are

multimolecular protein complexes involved in the innate immune response and composed of

NLRP3 (receptor), ASC (adaptor) and protease caspase-1 (effector). While NLRP3

inflammasomes are primarily activated in microglia as a pathogenic mechanism of AD(Heneka

et al., 2013), our data implicated this inflammatory response may also be upregulated in AD

neurons. We thus examined inflammasome activation via immunostaining of NLRP3 and ASC.

Under basal conditions, aged and fAD-PSEN1 tNeurons contained 2.7% inflammasome-

positive neurons per image field while aged/sAD tNeurons contained 4.6% inflammasome-

positive neurons per image field (Figure 6B).However, a 3-hr LLOME treatment led to

widespread activation of NLRP3 inflammasomes in tNeurons. Strikingly, the occurrence of

inflammasome-positive neurons was increased by 35-40% in aged/sAD tNeurons compared

with those of aged and fAD-PSEN1 donors. These results indicate that AD neurons possess a

relatively low-level inflammasome activation, but their high vulnerability to lysosomal damage

can lead to a potent activation of inflammasomes (Figure 6B).

We next examined whether tNeurons secrete inflammatory factors under basal and lysosomal

damage conditions. We thus compared cytokine secretion in the conditioned medium obtained

from tNeurons of young, aged, aged/sAD and fAD-PSEN1 donors (Figures 6C and 6D). To

assess if cytokine secretion is enhanced by lysosomal damage, we also compared the condition

medium of tNeurons of young donors with or without a low-dose LLOME treatment for 7 days

(Figures 6C and 6D) Cytokine profiling showed an age-dependent increase in levels of pro-

inflammatory cytokines (e.g. IL-6 and IFN-γ) and sAD-dependent increase in pro- and anti-

inflammatory cytokines (e.g. IL-1β, IL-4 and IL-15) and chemokines (e.g. CCL2 and CXCL12)

(Figures 6C and S7A). Importantly, extended treatment of young tNeurons with a low-dose of

LLOME increased the secretion of pro-inflammatory cytokines (e.g. IL-1β, IL-6 and IFN-γ) and

chemokines (e.g. CCL2) (Figures 6D and S7A). Pearson's correlation analysis identified a

moderate and positive correlation between total cytokine levels in the conditioned medium of

aged and aged/sAD tNeurons, as well as between young tNeurons treated with LLOME and AD

tNeurons (Figure 6E).

To put these observations in a disease relevant context, we examined cytokine expression in

existing human datasets and cerebrospinal fluid (CSF) samples. First, we examined cytokine

expression in single-nucleus RNA-seq analysis of neurons, astrocytes, microglia and

oligodendrocytes obtained from post-mortem cortex from HC donors. Cytokine and chemokine

transcripts were commonly found in neurons and glial cells. As expected, the transcript levels of

cytokines, chemokines and other inflammatory molecules are much higher in microglia than in

neurons (Figure 6F). Because CSF cytokines have been increasingly recognized as potential

biomarkers for diagnosing AD(Bettcher et al., 2018), we conducted an unbiased analysis of CSF

proteins in 50 HC and 29 AD donors in a search for biomarkers. Among them, we found higher

CSF levels of IL-15 and IL-20, and lower CSF levels of IL-1A, IL-1RA, VEGFA and TNFRSF6B

in AD than in HC (Figure S7B, Table S3). The increased concentration of IL-15 was previously

observed in CSF of AD patients and positively correlated with age of onset(Rentzos et al.,

2006). These data suggest that lysosomal dysfunction in aging and AD may lead to increased

secretion of a subset of cytokines that may serve as potential biomarkers of disease.

Rescuing defective lysosomes ameliorates AD pathologies in fibroblasts and tNeurons

An important question raised by our findings is whether aging and AD-linked lysosomal deficits

and vulnerabilities discovered in patient fibroblasts and tNeurons can be ameliorated by

pharmacological agents. Several small molecules have been shown to target specific aspects of

lysosomal homeostasis (Figure 7A). We previously identified a brain-penetrant lysosomal v-

ATPase complex activator, C381, that promotes lysosomal acidification. C381 can reduce

neuroinflammation, protein aggregates and neurodegeneration in mouse models of PD and

FTD(Vest et al., 2022). Another compound, thioperamide, increases a late endosomal and

lysosomal phospholipid, bis(monoacylglycero)phosphate (BMP), and rescues defects in

lysosomal lipid metabolism and membrane permeabilization in lysosomal storage

disorders(Moreau et al., 2019).

We tested whether C381 can ameliorate lysosomal dysfunction in human fibroblasts. Pre-

incubation of C381 at concentrations up to 12.5 µM for 24 hr reduced damaged lysosome-

mediated cell death in young, aged and AD fibroblasts (Figure 7B). We next examined if C381

reduces the loss of lysosomal acidification and proteolytic capacity upon LLOME treatment of

AD fibroblasts and subsequent recovery after LLOME washout (Figures 3J and 3K). Pre-treating

AD fibroblasts with 6.2 µM C381 improved their capacity to resist the loss of lysosomal

acidification and Cathepsin-B activity during LLOME treatment and accelerated the recovery

after LLOME washout (Figures 7C and 7D). Consistent with these functional recovery data,

C381 led to a dramatic reduction in number of CHMP2B and Galectin-3 puncta caused by

LLOME treatment (Figure 7E).

Next, we asked whether C381 ameliorates lysosomal deficits in tNeurons. First, we examined

the effect of pre-treatment of aged/sAD and fAD-PSEN1 tNeurons with 3.1 µM C381 for 7 days

on LLOME-mediated apoptotic cell death (Figure 7F). C381 was significantly neuroprotective

from LLOME-mediated apoptosis, and Caspase-3/7 activation by ~51% in AD tNeurons. As

observed in fibroblasts, C381 pretreatment reduced several aspects of lysosomal dysfunction in

neurons and improved their resilience to lysosomal damage. For instance, we find that aged

and aged/sAD tNeurons exhibited lower Cathepsin-B activity compared you young tNeurons,

even under basal condition (Figure 7G); the loss of Cathepsin-B activity is strongly exacerbated

by LLOME treatment, particularly for aged/sAD tNeurons. However, pre-treating tNeurons with

C381 restored Cathepsin-B activity to near-normal levels and ameliorated lysosomal proteolytic

deficits. Consistent with the improved recovery of lysosomal pH and proteolytic function, pre-

treatment of tNeurons with 3.1 µM C381 simultaneously reduced the number of detectable

CHMP2B and Galectin-3 puncta on lysosomes in aged and AD tNeurons by 15-39% and 34-

60%, respectively (Figure 7H).

The above analyses show that C381 preserves lysosomal functionality and facilitates the repair

of damaged lysosomes in aged and AD fibroblasts and tNeurons. We next investigated whether

improving lysosomal health impacts the AD-related pathologies observed in tNeurons, including

appearance of Aβ42 deposits (Figures 2D and S2D) and increased secretion of pro-

inflammatory cytokines (Figures 6C and S7A). Consistent wit lysosomal dysfunction leading to

accumulation of intra-neuronal Aβ42 levels, damaging lysosomal integrity by LLOME treatment

or blocking lysosomal repair with BAPTA-AM led to significant increases in intra-neuronal Aβ42

levels in aged/sAD and fAD-PSEN1 tNeurons (Figure 7I). Of note, LLOME had a marginal effect

on intra-neuronal Aβ42 levels in aged tNeurons. These experiments suggest that impairing

lysosomal function and repair beyond a specific threshold leads to Aβ42 accumulation. We next

tested if ameliorating lysosomal dysfunction by treating tNeurons with either C381 or

thioperamide ameliorates amyloid accumulation. Strikingly, both compounds significantly

reduced intra-neuronal Aβ42 levels, in aged tNeurons by ~32-36%, and AD tNeurons by ~34-

46%. Similar experiments causally linked lysosomal damage to inflammatory activation and

secretion of pro-inflammatory cytokines in AD tNeurons (Figures 6C, 6D and S7A). Thus,

amelioration of lysosomal damage by C381 treatment in aged/sAD and fAD-PSEN1 tNeurons

significantly reduced the secretion of pro-inflammatory cytokines (e.g. IL-6 and IL-15) and

chemokines (e.g. CCL2 and CXCL12) (Figures 7J and S7C). Taken together, these

experiments link the aging- and AD-related increase in lysosomal vulnerability and LQC

dysfunction to amyloid Aβ42 deposition and inflammatory cytokine secretion, and demonstrate

that improving lysosomal integrity and function has protective effects on aged/sAD and fAD-

PSEN1 neurons.

DISCUSSION

Here we address how aging and AD affect the human proteome using patient derived primary

fibroblast and their matching directly transdifferentiated tNeurons. These models have been

shown to preserve epigenetic and transcriptomic aging signatures(Huh et al., 2016; Mertens et

al., 2015; Miller et al., 2013). Our unbiased proteomic analyses were complemented and

extended by a battery of molecular and functional analyses. We find that aging and AD lead to

dramatic changes in several components of the proteostasis and organelle homeostasis

machinery. In particular, we identify the decline in endosome-lysosomal function and repair as a

cell-autonomous feature of sAD and fAD-linked mutations in PSEN1 gene. Deficits in lysosomal

repair lead to enhanced sensitivity to lysosomal damage stress in aged and AD. In neurons,

LQC defects are exacerbated, leading to damaged lysosomes even at basal conditions, which

predispose neurons to increased protein pathologies, such as intra-neuronal Aβ deposits,

inflammasome activation and cytotoxicity (Figure 7K). Consistent with our findings in tNeuron

models for AD, we identify massive accumulations of lysosomal damage markers across the

cerebral cortex of AD mice and patients, frequently colocalized with Aβ pathology. Finally, we

show that small molecules rescuing lysosomal pH regulation and integrity promote cellular

resilience to lysosomal damage and reduce both amyloid loads and pro-inflammatory cytokine

secretion in AD neurons. Importantly, our tNeuron models of aging and AD recapitulate AD-

related protein pathologies, including the spontaneous deposits of Aβ42 and p-Tau (Figures 2D

and S2D). These observations, together with our quantitative proteomic analyses support the

intrinsic dysregulation in proteostasis in AD neurons, and show it can lead to amyloid deposits

without seeding or overexpression of human disease proteins. Overall, this study demonstrates

the utility of transdifferentiation of fibroblasts from AD patients to neurons as a valuable

approach to understand molecular mechanisms of aging and AD and to advance

neurodegenerative disease therapeutic approaches.

Proteomic and proteostasis signatures in aging and AD

Our analyses reveal cellular proteome changes in human cell models of aging and AD. Of note,

such cytosolic and organelle-specific proteostasis changes are often not apparent from genomic

and transcriptomic data(Laskowska et al., 2019). Proteomic analyses of donor fibroblasts

indicate an age-dependent up-regulation of LQC proteins and and AD-dependent down-

regulation of v-ATPases (Figures 1F, 1G and 3A). In tNeurons, the levels of lysophagy-related

proteins were altered in aging (e.g. CNN2, HspB1) and AD (e.g. DPP7, PLBD2, PLD3, TAGLN)

(Figure 2E)(Eapen et al., 2021; Kravić et al., 2022). Our proteomic and functional analyses

inAD tNeurons highlight the functional interconnection between endosome-lysosomal,

proteostasis and inflammation subcellular systems. While the interconnectivity of different

quality control pathways establishes robustness and adaptability under normal conditions, the

impairment in one of these subsystems, for instance due to aging or disease-mutation, can drive

a cascade of decline to inter-connected subsystems (Gottschling and Nyström, 2017; Petkovic

et al., 2021). We propose that such a vicious cycle of dysfunction may contribute to the

synergistic defects observed for aging and AD in our cell models. Our data comparing two cell

types also suggest that neurons are particularly vulnerable to this cycle of homeostatic decline,

thus underlies their sensitivity to aging and mutation and their higher incidence of protein

pathologies accompanying cellular death.

Intriguingly, we find that AD patient fibroblasts exhibit an increased tendency to lysosomal

damage in absence of protein pathology. However, the more severe level of lysosome damage

in aged/sAD tNeurons is accompanied by protein deposits. Of note, the intracellular Aβ deposits

were significantly higher in fAD-PSEN1 than aged/sAD tNeurons. These lines of evidence

suggest that aging, genetic mutations and stochastic disease factors synergistically and

adversely impact lysosomal repair and recovery after damage; and beyond a threshold, this

may lead to a cell-autonomous mechanism for neurodegeneration in AD. While, defective

autophagy in microglia has been extensively studied and linked to AD pathogenesis (Wang et

al., 2022); the role of neuronal autophagy in this pathological event is less understood. Neurons

are particularly sensitive to lysosomal dysfunction compared with other types of brain cells (Tian

et al., 2021). Of note, lysosomes are interconnected hubs linked to both proteostasis and

homeostasis of other organelles (such as mitochondria and ER), it is clear to envision how such

deficits impair cell survival. Since internalized aggregated proteins are thought to be an

important mechanism of endosome-lysosomal membrane rupture(Ditaranto et al., 2001; Jiang

et al., 2017), the presence of protein pathologies likely enhances further lysosome dysfunction.

Previous studies found that the depletion of lysosomal repair factors, such as ESCRT-III

subunits or PI4K2A, in immortalized cell lines impacts lysosomal stability and causes an escape

of tau fibrils from lysosomes to trigger cytoplasmic aggregation (Chen et al., 2019; Tan and

Finkel, 2022). The knockdown of ESCRTs or expression of mutant VPS4A enhanced

intracellular Aβ accumulation and endosomal membrane lesion, mimicking AD pathogenesis

(Fruhmann et al., 2018; Willén et al., 2017). Hence, our work provides a direct link between

defective LQC and AD neuropathology (Figure 7K). The failure to resolve lysosomal damage in

conjunction with other well-studied autophagy-lysosomal defects can collectively contribute to

AD pathogenesis(Cortes and La Spada, 2019; Whyte et al., 2017; Wolfe et al., 2013).

Genetic risk factors link to endosomal and autophagy-lysosomal dysfunction in neurons

Our proteomic analyses resonate with GWAS studies identifying AD casual genes and risk

genes for late-onset AD to the endosomal and autophagy-lysosomal pathway, such as BIN1,

CLU, GRN, PICALM, PLD3 and SORL1(Zhang et al., 2022). We find PSEN1 mutations

promote defects in lysosomal acidification and increase susceptibility to lysosome-mediated

death. Our data is consistenty with, and explains, a number of previous studies. PSEN1

pathogenic mutations or knockout impair lysosomal v-ATPase assembly and autophagosome-

lysosome fusion and elevate Aβ42/40 ratio (Hung and Livesey, 2018; Kwart et al., 2019; Lee et

al., 2015; Lee et al., 2010). A recent study characterized different transgenic mouse models for

AD and identified accumulation of APP-βCTF/Aβ in poorly acidified lysosomes of cortical

neurons as a very early event in AD (Lee et al., 2022). Before amyloid deposits were observed,

neurons had developed low v-ATPase activities, lysosomal acidification deficits and higher

membrane permeabilization in mouse and Braak-stage-II human brains. The build-up of APP-

βCTF/Aβ in the perinuclear deacidified lysosomes progressed with age, consistent with our

findings. Our findings further implicate intraneuronal accumulation of amyloid peptides resulting

from neuronal lysosomal defects as the origin of extracellular amyloid plaques in the brain

(Gouras et al., 2000; Masters et al., 1985; Mochizuki et al., 2000).

Lysosomal pathology links to GVD and neuroinflammation in AD

The presence of cytoplasmic granule-containing vacuoles is a neuropathological feature of GVD

in AD(Köhler, 2016). Similar to our findings in cells, GVD bodies structurally resemble endocytic

and autophagic vesicles and contain various proteins, including the typical hallmarks CHMP2B,

LAMP1 and CK1δ, proteins involved in endocytosis (CHMP1A) and proteostasis (Hsp70), and

AD-related amyloidogenic proteins, such as Aβ, pTau and TDP-43(Hondius et al., 2021). Thus,

our analyses of AD tNeurons as well as human and mouse brain sections suggested lysosomal

damage underlying GVD etiology in AD.

Our study indicates that loss of lysosomal integrity in neurons plays critical roles in

inflammasome activation in pathological conditions. Consistent with this observation, a recent

study indicated the senescence-like neurons may be a novel source of neuroinflammation in AD

(Herdy et al., 2022). Previous findings also indicate lysosomal membrane permeabilization in

microglia activates NLRP3 inflammasomes and promotes the secretion of pro-inflammatory

cytokines, thus aggravating neuronal damage(Wang et al., 2017). Our cytokine profiling of

conditioned medium from AD tNeurons revealed that the increased secretion of pro-

inflammatory cytokines is associated with chronic lysosomal damage. Human CSF proteomics

further supported IL-15 as the potential biomarker for AD (Figure S7B).

Comparing transdifferentiated neurons with iPS cell-derived neurons

One key finding in our study is that aging-linked proteostasis and organelle homeostasis

impairment sensitizes cells, particularly neurons, to further damage due to stochastic AD events

or to AD-causing mutations. These findings are enabled because directly transdifferentiated

neurons retain aging signatures while stem cell reprogramming has demonstrated a remarkable

ability to rejuvenate cells by reversing age-related global changes (Mahmoudi and Brunet,

2012). Many lines of evidence indicate that cell-based models of age-dependent

neurodegenerative diseases benefit from retaining these aging marks. In Huntington's disease,

mutant huntingtin aggregates are often detected in human tNeurons and post-mortem tissue,

but rarely in iPS cell-derived neurons from the fibroblasts of same patient cohorts(Victor et al.,

2018). Similarly, tNeurons from donors harboring the FTD-linked MAPT IVS10+16 mutation

produced more aggregation-competent tau than iPS cell-derived neurons from the same

donors(Capano et al., 2022). These lines of evidence highlight the importance of aging in the

manifestation of proteostasis dysfunction. Transcriptomic and proteomic analysis of iPS cell-

derived neurons derived from sAD patients reported increased expression of genes and

proteins involved in synaptic vesicle regulation and microtubule organization and exhibited

pathological signatures of increased synaptic loss, Aβ42 levels and reactive oxygen species

production(Birnbaum et al., 2018; Israel et al., 2012; Lagomarsino et al., 2021; Zhao et al.,

2020). However, the iPS cell-derived sAD neurons did not reveal the strong disease- and aging-

related changes in proteostasis and lysosomal homeostasis observed with tNeurons. Since sAD

tNeurons and iPS cell-derived neurons both show an elevated expression of synaptic proteins,

perhaps synaptic dysregulation is an early, hyper-vulnerable event that is faithfully reflected by

both neuronal models, while proteostasis and organelle homeostasis deficits require the

contribution of aging.

722 Pharmacological rescue of lysosomal deficits ameliorates AD neuropathology

Our data suggests that drugs targeting lysosomal function may provide potential therapeutic

benefit to aging and AD. Lysosomal dysfunction is commonly observed across

neurodegenerative diseases and several strategies for improving autophagy-lysosomal function

have been tested in different model systems(Cao et al., 2021b). Thus, loss of lysosomal

acidification is central to aging and to several forms of neurodegenerative diseases and

neuronal ceroid lipofuscinosis. Based on these findings, targeting lysosomal acidification has

become a desired strategy. Re-acidifying lysosomes by v-ATPase-targeting compounds (for

example, EN6 and C381) or endocytosed nanoparticles (for example, PLGA-aNP and NP-1)

showed promising effects on promoting the clearance of protein aggregates, rescue of

autophagic dysfunction and attenuation of neuroinflammation and neurodegeneration in

different pathological contexts(Bourdenx et al., 2016; Chung et al., 2019; Lee et al., 2015; Vest

et al., 2022). The v-ATPase-targeting compound, C381 can rescue deficits in lysosomal

acidification and disease-relevant pathologies in both neurons and glial cells in disease models

for AD, PD and FTD. Further, C381 is orally bioavailable and brain penetrant, reflecting good

translational potential (Vest et al., 2022). Administration of mTOR inhibitor, rapamycin, was

shown to reduce neuropathology in mouse models for AD (Talboom et al., 2015), although its

associated immunosuppression effects reduces translation potential for AD. A large-scale drug

screening for mTOR-independent autophagy activators identified additional molecules that

promotes tau and Aβ clearance in neuronal models for AD and FTD(Yoon et al., 2022).

Moreover, ameliorating lysosomal membrane permeabilization prevents inflammatory

responses and cell death. Overexpression or exogenous administration of Hsp70 stabilized

lysosomes against membrane rupture in patient fibroblasts of Niemann-Pick disease(Kirkegaard

et al., 2010). Peripheral delivery of progranulin exhibited a promising effect on the correction of

BMP deficiency and reduction of lysosomal damage and dysfunction, lipofuscinosis and

inflammatory responses in the mouse model for FTD(Logan et al., 2021). Despite these

promising avenues, compounds that directly enhance lysosomal repair have not yet been

reported. One of the reasons is that these repair processes involve the concerted recruitment of

more than 30 proteins, including ESCRT and proteostasis factors to determine the timing and

hierarchy of events leading to membrane curvature, scission and fusion. The depletion of single

ESCRT subunit affects lysosomal repair, whereas the overexpression of single subunit has no

obvious improvement(Skowyra et al., 2018). Future studies leading to discovery of small

molecules that accelerate lysosomal repair may exert beneficial effects for treating

neurodegenerative diseases.

Study limitation: Our study uses monocultures of tNeurons without glial cell support and short-

term culture conditions (i.e. 5-6 weeks) which are quite distinct from the complex tissue

organization and prolonged aging processes in human brain. Thus, our cell-based models may

not entirely reflect the physiological signatures of aging and pathology of AD. The neurons were

derived from human adult fibroblasts of several donors to strive to account for patient

heterogeneity. The challenging conversion and cell culture system limits the experimental

scalability for larger donor cohort sizes and large-scale biochemical analyses. In addition, the

healthy young and aged fibroblasts were collected from donors who are clinically normal at that

moment, but we cannot rule out their possibility for developing dementia due to AD after

decades. Finally, it will be important to determine the mechanism of action of lysosome-

targeting compounds and investigate how these compounds change cell proteomes and

promote lysosomal activation.

Acknowledgements:

We thank Stanford Alzheimer's Disease Research Center (ADRC), D. Channappa and M.

Greicius with kind support for collecting and identifying human fibroblast lines, post-mortem

brain tissue and CSF samples. Y. Rosenberg-Hasson and J. Liang from Stanford Human

Immune Monitoring Center for help with cytokine profiling; J. Perrino from Stanford Cell

Sciences Imaging Core Facility for help with TEM experiments; P. Chu from Stanford Human

Pathology/Histology Service Center for help with tissue processing; M. Weglarz from Stanford

Shared FACS Facility for FACS support. We thank T.K. Rainbolt, V.B. Masto and S. Taguwa of

the Frydman laboratory for helpful discussions on experimental design; J. Kelly, R.I. Morimoto,

S. Finkbeiner and D. Gate for helpful inputs to the manuscript. This work was supported by NIA

P01AG054407 (J.F., D.F. & J.W.K.) and the Michael J. Fox Foundation for Parkinson's

Research and the Aligning Science Across Parkinson's (ASAP) initiative (J.F.). The Michael J

Fox Foundation administers the grant ASAP-000282 on behalf of ASAP and itself; the Glenn

Foundation for Medical Research Postdoctoral Fellowship in Aging Research (C.-C.C.), the Life

Sciences Research Foundation Postdoctoral Fellowship (C.-C.C.) and the Alzheimer's Disease

Research program in BrightFocus Foundation Grant (C.-C.C.); the NIA-funded Stanford ADRC

P50AG047366 and P30AG066515 (T.W.-C. & P.M.L.); the Goldberg Fund Fellowship and the

Miguel Servet Program (CP21/00017) (M.A.P.).

Author contributions:

C.-C.C. and J.F. conceptualized the study, designed experiments, interpreted data and wrote

the manuscript with inputs from all authors. C.-C.C. conducted cell culture, immunofluorescence

staining, microscopy, flow cytometry, conditioned medium experiments with the help of T.-T.L..

C.-C.C. performed transdifferentiation with the help of Y.S. under the supervision of M.W.. C.-

C.C., R.V. and J.L. carried out mice and drug experiments under the supervision of T.W.-C..

M.A.P. and J.W.-G. set up the proteomic data-processing pipelines and performed bioinformatic

data analysis with the help of C.-C.C. and J.A.P. under the supervision of S.P.G. and D.F..

P.M.L. processed and analyzed human CSF data under the supervision of T.W.-C.. J.W.K.

provided key reagents and helped design lysosome function assays. All authors read and

approved the manuscript.

Declaration of interests:

R.V., J.L. and T.W.-C. are co-founders of Qinotto Inc.

FIGURES AND FIGURE LEGENDS

Figure 1 Molecular and quantitative proteomic analysis identify signatures of aging and Alzheimer's disease (AD) in human adult fibroblasts.

(A) Experimental layouts for this study. Human fibroblasts are collected from donors of healthy young and aged, aged with sporadic AD (aged/sAD) and familial AD with PSEN1 mutations (fAD-PSEN1). Fibroblasts and transdifferentiated neurons (tNeurons) are used for molecular and proteomic characterization of aging and disease signatures and tested for cellular responses to small molecules. Validation of cellular findings is performed in post-mortem brain tissue from AD mouse models and patients.

(B) Levels of DNA damage measured by increased numbers of the nuclear foci of γ-H2AX
immunocytochemistry shown in the representative images of human fibroblasts.

(C) Age- and AD-related epigenetic alterations measured by histone modifications H3K9me3
and H4K16ac immunocytochemistry in human fibroblasts.

(D) Representative images of endogenous ubiquitin and p62/SQSTM1 in young fibroblasts
treated with DMSO control (Ctrl), Bortezomib (BTZ, 25 nM) or Chloroquine (CQ, 25 μM) for 24
hr. Insert: higher magnification view of ubiquitin and p62/SQSTM1. Scale bar: 40 μm.

(E) Quantification of proteostasis markers shown in panel (D) for detecting changes in ubiquitin
and p62/SQSTM1 under proteasomal and autophagic stress caused by the treatment of BTZ
and CQ, respectively, in young, aged and aged/sAD fibroblasts.

(F) Network analysis of fibroblast proteomes including proteins in the top-ranked pathways
associated with aging and sAD by GeneMANIA and Gene ontology (GO). Comparison between
the fibroblasts from aged and young donors (Aging proteome) and between the fibroblasts from
aged/sAD and aged donors (sAD proteome). Colored circles represent the enrichment of
identified proteins revealing by log2-fold change: increase in red and decrease in blue.

(G) Volcano plots showing differentially expressed proteins in the endosome-lysosomal
pathway. The plots display the log2-fold change (x-axis) and -log10 statistical P value (y-axis).
Proteins with significantly changed levels are shown in dark grey. Among them, the endosome-
lysosomal proteins are labeled in blue-green color.

(H) Clustering representation based on the Pearson correlation of the enrichment of fibroblast
proteins in the proteostasis pathways. GO analysis of the differentially expressed proteins,
defined by the relative protein abundance (z-score), reveals the top proteostasis terms
associated with aging and sAD, including proteolysis, protein folding and translation. Black dots
within the violin graph represent the expression levels of individual proteins in each group.

Data show box-and-whisker plots of 2 to 4 independent experiments and cells from independent
healthy control (HC) and AD donors. Box-and-whisker plots depict 5th, 25th, 50, 75th and 95th
percentiles. Differences between groups are compared with One-Way ANOVA followed by
Bonferroni post-hoc analysis. **P < 0.01 and ***P < 0.001.

See also Figure S1 and Table S1.

Figure 2 Human tNeurons carry molecular and proteomic signatures of aging, sAD and fAD harboring PSEN1 mutations.

(A) Representative images of human fibroblasts immunostained for S100A4 and Vimentin, and tNeurons immunostained for GAP43, Tuj1, MAP2 and NeuN with DAPI counterstaining. Quantification of Tuj1-, MAP2- and NeuN-positive cells at post-induction day (PID) 35. Scale bar: 100 µm.

(B) Measurement of DNA damage in tNeurons quantified by numbers of the nuclear foci of γ-H2AX immunocytochemistry.

(C) Measurement of immunocytochemical changes in histone modifications H3K9me3 and H4K16ac in tNeurons.

(D) Representative images and quantification of proteostasis- and disease-associated protein markers in tNeurons, including amyloid-β 42 (Aβ42), hyper-phosphorylated tau (pTau) and TDP-43 (pTDP-43), autophagy adaptor p62/SQSTM1 and ubiquitin. Scale bar: 20 µm.

(E) Differential expression of proteins detected in tNeurons from healthy young (n = 3) and aged (n = 3) individuals, and patients with sAD (n = 6) and fAD-PSEN1 (n = 2) at PID 40. The top pathways for aging, sAD and fAD-PSEN1 proteomes are analyzed using GO databases and annotated in color. The aging proteome is reflected by comparing proteomic data between aged and young, the sAD proteome by comparing between aged/sAD and aged, and the fAD-PSEN1 proteome by comparing between fAD-PSEN1 and aged/sAD. The size of the filled circles reflects log2-fold change. A circle with a thick black outline indicates that the protein is enriched in aged (Aging proteome), aged/sAD (sAD proteome) and fAD-PSEN1(fAD-PSEN1 proteome) tNeurons. Protein name with underline represents the previously known protein associated with AD.

(F) List of differentially expressed proteins associated with the risk genes for age-related neurodegenerative diseases. AD: Alzheimer's disease. PD: Parkinson's disease. ALS/FTD: Amyotrophic lateral sclerosis and frontotemporal dementia.

(G) Clustering heatmap of Pearson correlation coefficients of tNeuron protein expression in the proteostasis pathways. Top GO terms associated with aging and AD, including proteolysis, autophagy, amyloid processing and translation. In the proteolysis term, Cluster A to F show distinct protein expression patterns and the differentially expressed protein lists, defined by the relative protein abundance (z-score) across young, aged, aged/sAD and fAD-PSEN1 group. Bar graph represents the average expression levels of proteins in each group.

(H) Diagram identifying aging- and AD-related protein hits and pathways that are unique to or

shared by human fibroblasts and tNeurons.

Data show box-and-whisker plots of 2 to 4 independent experiments and cells from independent

HC and AD donors. Box-and-whisker plots depict 5th, 25th, 50, 75th and 95th percentiles.

Differences between groups are compared with One-Way ANOVA followed by Bonferroni post-

hoc analysis. *P < 0.05, **P < 0.01 and ***P < 0.001.

See also Figure S2, Figure S3 and Table S2.

Figure 3 Defects in lysosomal quality control (LQC) exacerbate lysosomal dysfunction

and cell death in AD fibroblasts.

(A) Clustering heatmap of Pearson correlation coefficients of the fibroblast protein expression in

the endosome-lysosomal pathway. Cluster B and C show the reduction in the relative

abundance of proteins involved in lysosomal pH regulation (Cluster B), and lysosomal

proteolysis and LQC (Cluster C) in aged/sAD fibroblasts.

(B) Network of fibroblast proteins and the relative abundance in the endosome-lysosomal

pathway. Protein abundance increases in aged/sAD shown in red and decreases in blue. The

circle size is represented by -log10 statistical P value.

(C) Transmission electron microscopy (TEM) for analyzing the ultrastructural morphology of

lysosomes in fibroblasts. Experiments were performed in 2 cell lines from each group and

repeated twice independently. Scale bar: 1 µm.

(D) Immunogold staining of lysosomal damage marker Galectin-3 for TEM in fibroblasts treated

with 1 mM LLOME for 10 min reveals Galectin-3 translocation to the damaged lysosomes.

Experiments were performed in 2 cell lines from each group and repeated twice independently.

Insert: higher magnification view of gold particles. Size of gold particles: 10 nm. Scale bar: 500

nm.

(E) Schematic for LQC machinery. While undergoing lysosomal damage stress, the lysosomal

membrane is subjected to rupture. Lysosomes recruit ESCRT proteins (ESCRTs) to repair the

mildly damaged membrane and Galectins to target the massively damaged lysosomes for

lysophagy. Immunostaining of ESCRT-III protein CHMP2B (green), Galectin-3 (magenta) and

lysosome-associated membrane protein LAMP2 (grey), during different time courses of

lysosomal damage caused by 1 mM LLOME treatment. Insert: higher magnification view of

CHMP2B, Galectin-3 and LAMP2. Hollow arrowhead: colocalization of CHMP2B and LAMP2.

Solid arrowhead: colocalization of CHMP2B, Galectin-3 and LAMP2. Scale bar: 20 µm.

(F) 2D Histogram shows the colocalization patterns of CHMP2B (x-axis) and Galectin-3 (y-axis)

immunofluorescence in young fibroblasts during 1 mM LLOME treatment and washout. Each

dot means the pixel intensity in both CHMP2B and Galectin-3 channels of the image.

(G) Plot profiles represent the dynamics of CHMP2B and Galectin-3, indicated by the changes

in puncta number as shown in panel (E), in fibroblasts of young (Y), aged (A), aged/sAD (sAD)

and fAD-PSEN1 (PSEN1) donors during 1 mM LLOME treatment and washout. The puncta

number represents the tendency of lysosomes toward damage and the efficiency of lysosomal

recovery from damage.

(H) Lysosome-mediated death in fibroblasts caused by the treatment of LLOME (upper) or Silica

nanoparticles (bottom) for 24 hr, detected by the SYTOX Green assay. The SYTOX Green

intensities are normalized to Hoechst and divided by the data with no compound treatment as

fold change.

(I) The detection of leakage of intraluminal contents from fibroblast lysosomes during 1 mM

LLOME treatment.

(J) Changes in lysosomal acidification, indicated by the pH-sensitive Fluorescein isothiocyanate

(FITC)-conjugated Dextran, in fibroblasts during 1 mM LLOME treatment for 30 min and after

washout for up to 8 hr, revealing the functional recovery of lysosomes after damage.

(K) Measurement of lysosomal proteolytic capacity by monitoring the lysosomal protease

Cathepsin-B activity of fibroblasts during 1 mM LLOME treatment for 30 min and washout.

(L) Immunostained fibroblasts for endogenous Hsp70, TDP-43 and Nup98 in the presence of

1mM LLOME. Insert: higher magnification view of Hsp70, TDP-43 and Nup98 and LAMP2.

Yellow arrowhead: protein colocalization with LAMP2. Scale bar: 20 μm.

Data show mean and SD or box-and-whisker plots of 3 to 8 independent experiments and 3 cell

lines from independent HC and AD donors. Box-and-whisker plots indicate 5th, 25th, 50, 75th

and 95th percentiles. Differences between groups are compared with One-Way or Two-Way

ANOVA followed by Bonferroni post-hoc analysis. $*P < 0.05$, $**P < 0.01$ and $***P < 0.001$. $\#P <$

0.05 and $\#\#\#P < 0.001$ as comparing to 0 min of each group in panel (I).

See also Figure S4.

Figure 4 Deficits in lysosomal membrane repair drive lysosomal dysfunction and Aβ

pathology in AD tNeurons.

(A) TEM for analyzing ultrastructural morphology of lysosomes and mitochondria in human

tNeurons. Changes in lysosomal size, electron-dense material abundance and mitochondria-

lysosome contacts defined by the contact length are detected in tNeurons. Red arrowhead:

electron-dense materials in lysosomes. Scale bar (i): 20 µm. Scale bar (ii-iv): 1 µm.

(B) Representative images of tNeurons immunostained for LAMP2 (green), CHMP2B and

Galectin-3 (magenta) and Tuj1 (blue) at PID 35. Insert: higher magnification view of CHMP2B

and Galectin-3 colocalization with LAMP2. Scale bar: 10 µm.

(C) Plots indicate the quantification of numbers of CHMP2B and Galectin-3 puncta in the cell

body of tNeurons at the basal condition as shown in panel (B).

(D) Representative images of fAD-PSEN1 tNeurons immunostained for LAMP2 (green),

ESCRT-0 HGS protein (magenta) and Tuj1 (blue) during 0.25 mM LLOME treatment for 30 min

and washout at PID 36. Insert: higher magnification view of HGS and LAMP2 in the cell body

and neurites. Yellow arrowhead: HGS colocalization with LAMP2. Scale bar: 10 µm.

(E) Plot reflects the lysosomal recovery from damage by measuring the numbers of HGS puncta

in the cell body during LLOME treatment and washout as shown in panel (D). The half-life (t1/2)

represents the time required for lysosomal recovery in each group.

(F) Changes in lysosomal acidification by detecting the intensity of preloaded pH-sensitive

FITC-conjugated Dextran in tNeurons at the basal condition and during 0.25 mM LLOME

treatment for 30 min.

(G) Lysosomal proteolysis reflected by the fold change of Cathepsin-B activity in tNeurons

calculated by the value in 0.25 mM LLOME treatment for 30 min relative to the basal condition.

(H) Immunofluorescent analysis of colocalization of Aβ42 with LAMP1 and LC3B in aged/sAD

tNeurons. Insert: higher magnification view of Aβ42, LAMP1 and LC3B. Scale bar: 10 µm.

Data show box-and-whisker plots of 3 independent experiments and 3 cell lines from

independent HC and AD donors. Differences between groups are compared with One-Way

ANOVA followed by Bonferroni post-hoc analysis. *P < 0.05 and ***P < 0.001. #P < 0.05 and

###P < 0.001.

See also Figure S5.

Figure 5 Lysosomal damage is exacerbated in AD and linked to amyloid accumulation in post-mortem brain tissue.

(A) Double immunofluorescent analysis of CHMP2B, Galectin-3 or Aβ (magenta) and LAMP2 (cyan) in the post-mortem cerebral cortex of HC and AD donors. Insert: higher magnification view of CHMP2B, Galectin-3 and Aβ colocalization with LAMP2. Yellow arrowhead: CHMP2B-positive fibril structures. Scale bar: 20 μm.

(B) Immunofluorescent analysis of CHMP2B, Galectin-3, LAMP1 and Aβ42 in the neocortex of non-transgenic wild-type mice (NTg) and transgenic mice expressing mutant human APP with the Swedish (K670N/M671L) and London (V717I) mutations (APPLon/Swe) for modeling AD. The brain tissue is co-stained with MAP2 and Hoechst. Quantitative analysis shows changes in the immunoreactivity of LAMP1, CHMP2B and Galectin-3 in APPLon/Swe mice relative to NTg mice. Insert: higher magnification view of colocalization between CHMP2B, Galectin-3, LAMP1 and Aβ42. Scale bar: 50 μm.

988   Data show mean and SD plots of 3 independent experiments in mice (NTg: n = 8 to 10; APP: n

989   = 10 to 13). Differences between groups are compared with unpaired t test. ***$P < 0.001$.

990   See also Figures S5 and S6.

Figure 6 Lysosomal damage mediates inflammatory responses in AD tNeurons.

(A) Proteomic network of tNeuron proteins and the relative abundance in the inflammatory response pathway. The relative abundance indicated by log2-fold change is shown by color: increase in red and decrease in blue.

(B) Representative images of tNeurons immunostained for inflammasome markers NLRP3 (cyan) and ASC (magenta) with or without 0.25 mM LLOME treatment for 3 hr at PID 40. Quantification of the percentage of tNeurons with inflammasomes per image. Schematic for the association between lysosomal vulnerability/damage and inflammasome activation in aged and aged/sAD tNeurons. Yellow arrowhead: colocalization of NLRP3 and ASC. Scale bar: 10 μm.

(C) Experimental schematic for inflammatory profiling of conditioned medium from tNeurons at

the basal condition at PID 40. Heatmap represents the changes in cytokine and chemokine

levels. Exact log2-fold change (Log2FC) and P values can be found in Figure S7A.

(D) Schematic and heatmap for inflammatory profile of conditioned medium from young

tNeurons treated with or without lysosomal damage stress (0.1 mM LLOME starting at PID 33

for 7 days). Exact log2-fold change (Log2FC) and P values can be found in Figure S7A.

(E) Heatmap and Pearson correlation analysis for identified cytokines and chemokines of

tNeurons at the basal and chronic lysosomal damage conditions from panel (C) and (D).

Number in each cell indicates the correlation coefficient.

(F) Single-cell transcriptomic analysis of selected cytokines and chemokines from panel (C) and

(D) in major cell types of human brain, including neuron, astrocyte, microglia and

oligodendrocyte. The transcript expression scores (H: high; M: medium; L: low) is determined by

the expression level of each transcript in public datasets retrieved from the Human Cell Atlas

and Allen Brain Map.

Violin plots depict the median and distributions of data from 3 independent experiments and 3

cell lines from independent healthy controls and AD patients. Inflammatory profiling is

conducted in technical replicates and 2 to 6 cell lines. Log2-fold change in mean fluorescence

intensity (MFI) is compared to the median of young tNeurons. Differences between groups are

compared with One-Way ANOVA followed by Bonferroni post-hoc analysis. $*P < 0.05$ and $***P$

< 0.001.

See also Figure S7.

1024

1025

1026

Figure 7 Pharmacological rescue of lysosomal resilience to damage ameliorates AD phenotypes in fibroblasts and tNeurons.

(A) Schematic for small molecules that modulate lysosomal function. C381: regulates lysosomal pH by targeting v-ATPases. Thioperamide: elevates the levels of bis(monoacylglycero)phosphate (BMP). LLOME: triggers lysosomal membrane permeabilization. BAPTA-AM: cell-permeable selective Ca2+ chelator that blocks Ca2+-dependent recruitment of ESCRT-III/Vps4 complex to the damaged lysosome.

(B) Plot profile of a dose-dependent reduction in LLOME-mediated cell death in fibroblasts by C381 pre-treatment for 24 hr. Cells are treated with 1 mM LLOME for 24 hr. The SYTOX intensity is normalized to Hoechst and divided by the value at 0 µM C381 to reflect the fold change by C381.

(C) FITC-dextran intensity is indicative of lysosomal acidity. The pre-treatment of 6.25 µM C381

for 24 hr in fibroblasts is followed by 1 mM LLOME treatment for 30 min and washout (refer to

left y-axis). Dysregulation of lysosomal acidification in aged/sAD fibroblasts (dash line, refer to

right y-axis) shown as a reference control acquired from Figure 4J.

(D) Plot indicates the restoration of lysosomal proteolytic dysfunction in fibroblasts by the pre-

treatment of 6.25 µM C381 for 24 hr. Lysosomal protease Cathepsin-B activity is measured

during LLOME treatment and washout (refer to left y-axis). Defective recovery of Cathepsin-B

activity in aged/sAD fibroblasts (dash line, refer to right y-axis) serving as a reference control

acquired from Figure 4K.

(E) The effect of pre-treatment of 6.25 µM C381 for 24 hr on lysosomal damage in fibroblasts,

indicated by the numbers of CHMP2B and Galectin-3 puncta after 1 mM LLOME treatment for

30 min.

(F) Caspase-3/7 activation, a canonical upstream marker of apoptosis, is found in aged/sAD

and fAD-PSEN1 tNeurons after 0.5 mM LLOME treatment for 1 hr at PID 42. The pre-treatment

of 3.1 µM C381 is continued for 7 days.

(G) Lysosomal proteolysis in tNeurons is reflected by Cathepsin-B activity. The changes in

Cathepsin-B activity can be observed after 0.25 mM LLOME treatment for 30 min at PID 35 or

the pre-treatment of 3.1 µM C381 for 7 days.

(H) 2D Histogram represents the molecular signatures of lysosomal damage by measuring

CHMP2B (x-axis) and Galectin-3 (y-axis) puncta number in aged, aged/sAD and fAD-PSEN1

tNeurons after 0.25 mM LLOME treatment at PID 35. Each dot represents a single neuron and

the number of detectable CHMP2B and Galectin-3 puncta in the cell body. Pre-treatment with

3.1 µM C381 for 7 days alters the CHMP2B and Galectin-3 puncta number.

(I) Lysosomal modulation by small molecules influences the intra-neuronal Aβ42 levels

measured in aged, aged/sAD and fAD-PSEN1 tNeurons. Representative images of aged/sAD

tNeurons immunostained for Aβ42 (magenta) and Tuj1 (green) at PID 35. Quantification of fold-

changes in Aβ42 levels during treatment relative to DMSO control. Scale bar: 50 µm.

(J) Rescue of lysosomal damage reduces the secretion of inflammatory factors from AD

tNeurons. Inflammatory profiling of conditioned medium from aged/sAD and fAD-PSEN1

tNeurons at PID 35 after the treatment of 3.1 µM C381 for 7 days. Exact Log2FC and P values

can be found in Figure S7C.

(K) Summary of aging and AD signatures in fibroblasts and tNeurons, including the aging- and

AD-dependent proteome-wide remodeling and neuron-specific protein pathologies. Proposed

model for the link of lysosomal vulnerability and damage to neurotoxicity in AD and the strategy

for rescuing lysosomal defects in neurons.

Data show box-and-whisker plots of 3 to 6 independent experiments and 3 cell lines from

independent healthy controls and AD patients. Box-and-whisker plots depict 5th, 25th, 50, 75th

and 95th percentiles. Data show mean and SD plots of 3 to 5 independent experiments.

Inflammatory profiling is conducted in technical replicates and 2 cell lines. Differences between

groups are compared with One-Way or Two-Way ANOVA followed by Bonferroni post-hoc

analysis. *$P < 0.05$, **$P < 0.01$ and ***$P < 0.001$.

See also Figure S7.

1080

1081 **STAR Methods**

1082 **Lead Contact for Reagent and Resource Sharing**

1083 Further requests for resources and reagents should be directed to the Lead Contact, Judith
1084 Frydman (jfrydman@stanford.edu).

1085

Data and Code availability

All proteomic source data of human fibroblasts and tNeurons have been deposited and are
publicly available at ProteomeXchange (accession number: PXD040834). The mass
spectrometry parameters, sample information, raw data and the comparison between our
datasets and public genomic, transcriptomic and proteomic repositories are provided in Tables
S2 and S4.

1092

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Human subjects

De-identified human fibroblasts, post-mortem prefrontal cortex and CSF samples were acquired
from individuals of various ages and disease conditions from Stanford Alzheimer's Disease
Research Center (ADRC). Cells and tissue samples were obtained under written consents from
all subjects approved by Institutional Review Board of Stanford University. The cell and tissue
samples collected by ADRC were not specifically for this study. For histological experiments, of
these subjects, 8 were assessed as healthy control (HC) and 10 were patients with cognitive
impairment (dementia due to AD). For CSF proteomics experiments, 50 were assessed as HC
and 29 were patients with cognitive impairment (MCI or dementia due to AD). Age and sex
demographics are detailed in Tables S1 and S3. In the ADRC, all HC and individuals with
cognitive impairment received neurological examinations and cognitive tests to determine
cognitive status and consensus diagnosis by a team of neuropathologists. The pathological
diagnosis of post-mortem tissues was made by microscopic examination of multiple brain
regions using Amyloid score, Braak neurofibrillary degeneration score and CERAD neuritic
plaque score.

1109

Cell culture

HEK293T cells

The HEK293T cells are the cell line derived human embryonic kidney and were acquired directly

from ATCC (CRL-1573). Cells were grown in culture medium (DMEM supplemented with

GlutaMAX, 10% FBS, 1% Penicillin-Streptomycin, 1% HEPES and 1% Sodium pyruvate)

(ThermoFisher Scientific) sterilized by a 0.22 μm vacuum filter (ThermoFisher Scientific) in a

37°C incubator with 5% CO_2 in the air. The HEK293T cells were used for lentiviral production

by transfecting lentiviral vector of interest mixed with packaging and envelope plasmid.

Passaging the cells was performed with Trypsin-EDTA (ThermoFisher Scientific) every three

days.

1120

Human fibroblasts

The primary human dermal fibroblasts derived from clinically healthy adults and individuals

diagnosed with AD were collected from the shared resources in the Stanford ADRC and Coriell

Institute for Medical Research, which operates the NIGMS, NIA, NINDS cell repository. Cells

were grown in culture medium (DMEM supplemented with GlutaMAX, 10% FBS, 1% Penicillin-

Streptomycin, 1% MEM NEAA, 1% Sodium pyruvate and 0.1% β-mercaptoethanol)

(ThermoFisher Scientific) sterilized by a 0.22 μm vacuum filter (ThermoFisher Scientific) and

maintained in a 37°C, 5% CO_2 incubator. The subculture of proliferating fibroblasts used for

regular experiments and neuronal transdifferentiation was typically within 3 to 7 passages.

Some fibroblast lines obtained with slightly higher passage numbers were used for neuronal

transdifferentiation no more than 12 passages. Passaging the cells was performed with Trypsin-

EDTA every 6-7 days.

1133

Human tNeurons

The tNeurons were transdifferentiated from human fibroblasts with low passage numbers using

a combination of transcription factors and small molecules. The method was derived from the

protocols as previously described(Pang et al., 2011; Tanabe et al., 2018). Briefly, the lentiviral

expression of Brn2, Ascl1, Myt1l and Ngn2 in human fibroblasts initiated neuronal

reprogramming. The transduced cells underwent puromycin and PSA-NCAM selection and

cultured in reprogramming medium (DMEM/F12:Neurobasal (1:1) Medium, 2% B-27, 1% N-2,

0.25% GlutaMAX, 1% Penicillin-Streptomycin) (ThermoFisher Scientific) for 15 days and then

switched to maturation medium (BrainPhys Neuronal Medium (STEMCELL Technologies), 2%

B-27, 1% N-2, 0.25% GlutaMAX, 1% Penicillin-Streptomycin) for additional 15 to 22 days. The

culture media were supplemented with BDNF and NT-3 (neurotrophic factors; Peprotech),

Doxycycline (effector for the Tet-On system; Cayman), Forskolin (cAMP activator; Sigma-

Aldrich), SB 431542 (TGF-β/Activin/Nodal inhibitor; TOCRIS), Dorsomorphin (BMP inhibitor;

TOCRIS) and XAV939 (removal from maturation medium) (Wnt inhibitor; Stemgent). The

tNeurons resembled cortical glutamatergic neurons and were used for the evaluation of

reprogramming efficiency, aging and disease signatures, and small molecule treatments after 5

weeks in culture in a 37°C, 5% CO_2 incubator. The medium was half-changed every 2-4 days

throughout the lifetime of the culture. Detailed protocol was described in Method Details > Direct

generation of neurons from human fibroblasts.

1153

Mice

All mice used in this study were C57BL/6 genetic background. Mice of old age (20-24-month

old) were obtained from the National Institute on Aging rodent colony and young age (3-month

old) were obtained from Jackson Laboratories or Charles River Laboratories. All experiments

used male mice. A transgenic mouse model with the expression of high levels of human

APP751 carrying V717I, K670M/N671L mutations (also referring to APPLon/Swe) in neurons

under control of a Thy1.2 promoter has been studied in different laboratories. The APPLon/Swe

mice developed amyloid plagues associated with an overproduction of Aβ42 in the neocortex

and working memory deficits at 3 months old and the plaque formation spread to the

hippocampus and thalamus region at 5-7 months old(Faizi et al., 2012; Rockenstein et al.,

2001). This study used 3-6-month-old APPLon/Swe and the age-matched non-transgenic mice.

All mice were kept in a 12-hr light/dark cycle in a temperature- and humidity-controlled

environment and provided ad libitum access to food and water. All animal care and procedures

complied with the Animal Welfare Act and were in accordance with institutional guidelines and

approved by the V.A. Palo Alto Committee on Animal Research and the institutional

administrative panel of laboratory animal care at Stanford University.

1170

METHOD DETAILS

Lentivirus preparation

The FUW lentiviral vector expressing single transcription factor, including Brn2, Ascl1, Myt1l,

and Ngn2-puro (also referring as BAMN factors), or EGFP is under the control of TetO promoter

and M2rtTA under the control of ubiquitin promoter. The HEK293T cells (ATCC) were plated at

a density of 6 x 106 in a poly-L-ornithine-coated 10-cm dish and the next day co-transfected

with 5 µg lentiviral transfer vector, 4 µg packaging plasmid (psPAX2) and 2.5 µg envelope

plasmid (pMD2.G) using Lipofectamine 2000 in OptiMEM (ThermoFisher Scientific). After 6-hr

incubation of Lipofectamine/DNA mixture in OptiMEM, the transfection medium was replaced

with fresh DMEM supplemented with GlutaMAX, 2% FBS, 1% Penicillin-Streptomycin, 1%

HEPES and 0.1% BME (ThermoFisher Scientific). The cell supernatants containing lentiviral

particles were harvested after 24 hr and stored at 4°C. Cells were replenished with fresh DMEM

medium with 2% FBS and cultured for additional 24 hr. The supernatants were then harvested

and pooled with the first collection. To remove cell debris, the supernatants were centrifugated

at 400 x G for 5 min and passed through 0.45 µm syringe filters. The clear virus-containing

media can be stored at 4°C for about 1 week. For long-term storage, alternatively, the virus-

containing media were spun by ultracentrifugation at 25,000 rpm for 90 min at 4°C to pellet the

viruses. The viruses were resuspended in DMEM medium with 2% FBS and snap-frozen in

small aliquots to store at -80°C.

1190

Direct generation of neurons from human fibroblasts

Human adult fibroblasts were plated at a density of 200 × 103 per well of a 6-well plate coated

with poly-L-ornithine. The next day, fibroblasts were infected with lentiviruses expressing BAMN

factors and M2rtTA with or without EGFP by incubating with the diluted virus-containing medium

in DMEM supplemented with GlutaMAX, 2% FBS, 1% Penicillin-Streptomycin, 1% HEPES and

0.1% BME plus 4 µg/mL Polybrene for 24 hr. The virus-containing medium was then discarded

and replaced by fresh fibroblast culture medium plus 1 µg/mL Doxycycline, followed by

puromycin (0.5 µg/mL) selection for 48 hr. The transduced cells were subjected to PSA-NCAM+

selection following the manufacturer's instructions. Briefly, 0.05% Trypsin-EDTA was added to

the cells for 5 min at 37°C to dissociate the cells from surface and neutralized by fibroblasts

culture medium, followed by the centrifugation at 300 x G for 5 min at room temperature to pellet

cells. The cells were resuspended in autoMACS buffer and labeled with anti-PSA-NCAM-APC

(Miltenyi Biotec) for 10 min at 4°C in the dark. After a wash by autoMACS buffer and

centrifugation at 300 x G for 10 min, the cells were incubated with anti-APC MicroBeads

(Miltenyi Biotec) for 15 min at 4°C in the dark. Unbound beads were then washed off and the

cells were resuspended in autoMACS buffer for subsequent flow cytometry analysis and

separation of magnetically PSA-NCAM-labeled and unlabeled cells. The PSA-NCAM+ cells

were re-plated at a density of 50 × 103 cells per cm2 to the plate coated with vitronectin (VTN-

N, 5 µg/mL; ThermoFisher Scientific) and laminin (rhLaminin-521, 1 µg/mL; Corning). Cells were

cultured in the fibroblast culture medium plus 1 µg/mL Doxycycline and the next day switched to

the reprogramming medium (DMEM/F12:Neurobasal (1:1) Medium, 2% B-27, 1% N-2, 0.25%

GlutaMAX, 1% Penicillin-Streptomycin) (ThermoFisher Scientific) supplemented with small

molecules: 1 µg/mL Doxycycline (Cayman), 5 µM Forskolin (Sigma-Aldrich), 10 µM SB 431542

(TOCRIS), 2 µM Dorsomorphin (TOCRIS) and 2 µM XAV939 (Stemgent). After one week, 10

ng/mL BDNF and NT-3 (Peprotech) were added to the reprogramming medium. Half of the

medium was changed every 2 to 3 days. After 8 days, cells were switched to maturation

medium (BrainPhys Neuronal Medium (STEMCELL Technologies), 2% B-27, 1% N-2, 0.25%

GlutaMAX, 1% Penicillin-Streptomycin) supplemented with 1 µg/mL Doxycycline, 5 µM

Forskolin, 10 µM SB 431542, 2 µM Dorsomorphin and 10 ng/mL BDNF and NT-3 and cultured

for additional 15 to 22 days. Half of the medium was changed every 3 to 4 days. The efficiency

of transdifferentiation of human fibroblasts into tNeurons was measured by the percentage of

remaining transduced cells that express Tuj1, NeuN and MAP2 and the percentage of EGFP-

positive cells showing neuron-like morphology.

1224

Quantitative proteomics (TMT, Tandem Mass Tag)

Flash frozen cell pellets were lysed in 8M urea buffer (8M urea, 150 mM NaCl, 50 mM HEPES

pH 7.5, 1x EDTA-free protease inhibitor cocktail (Roche), 1x PhosSTOP phosphatase inhibitor

cocktail (Roche)). Lysates were clarified by centrifugation at 17,000 x g for 15 min at 4°C.

Protein concentration of the supernatant was quantified by bicinchroninic acid assay (BCA,

Pierce). To reduce and alkylate cysteines, 100 µg of protein was sequentially incubated with 5

mM TCEP for 30 min, 14 mM iodoacetamide for 30 min, and 10 mM DTT for 15 min. All

reactions were performed at room temperature. Next, proteins were chloroform-methanol

precipitated and the pellet resuspended in 200 mM EPPS pH 8.5. Then, LysC (Wako) was

added at 1:100 (LysC:protein) ratio and incubated overnight at room temperature in an orbital

shaker at 1,500 rpm. The day after, samples were further digested for 5 hr at 37ºC with trypsin

at 1:75 (trypsin:protein) ratio in an orbital shaker at 1,500 rpm. After digestion, samples were

clarified by centrifugation at 17,000 x g for 10 min. Peptide concentration of the supernatant was

quantified using a quantitative colorimetric peptide assay (ThermoFisher). For TMT labelling we

used 2 different kits to label each of the TMT experiments included in this manuscript. In TMT-

01, peptides from fibroblast's samples were labelled with TMT-6plex tags. In TMT-02, peptides

from tNeuron's samples were labelled with TMTpro-16plex tags. Same method was followed to

label both TMT sets(Cao et al., 2021a; Liu et al., 2022; Nguyen et al., 2017). Briefly, 25 g of

peptides was brought to 1 µg/µl with 200 mM EPPS (pH 8.5), acetonitrile (ACN) was added to a

final concentration of 30% followed by the addition of 50 µg of each TMT reagents. After 1 hr of

incubation at room temperature, the reaction was stopped by the addition of 0.3%

hydroxylamine (Sigma) for 15 min at room temperature. Extra information regarding both TMT

sample labels is included in Table S4. After labelling, samples of each TMT were combined,

desalted with tC18 SepPak solid-phase extraction cartridges (Waters), and dried in the

SpeedVac. Next, desalted peptides were resuspended in 5% ACN, 10 mM NH4HCO3 pH 8.

Both TMT were fractionated in a basic pH reversed phase chromatography using a HPLC

equipped with a 3.5 µm Zorbax 300 Extended-C18 column (Agilent). Fractions were collected in

a 96-well plate, then combined into 24 samples. Twelve of them were desalted following the

C18 Stop and Go Extraction Tip (STAGE-Tip)(Rappsilber et al., 2003) and dried down in a

SpeedVac. Finally, peptides were resuspended in 1% formic acid, 3% ACN, and analyzed by

LC-MS3 in an Orbitrap Fusion Lumos (ThermoFisher). For TMT-01 the MS was running in SPS-

MS3 mode(McAlister et al., 2014). For TMT-02, instrument was equipped with FAIMS and

running in RTS-MS3 mode(Schweppe et al., 2020a; Schweppe et al., 2019; Schweppe et al.,

2020b). More information regarding all MS parameters used for both TMT are included in Table

S4. A suite of in-house pipeline (GFY-Core Version 3.8, Harvard University) was used to obtain

final protein quantifications from all RAW files collected. RAW data were converted to mzXML

format using a modified version of RawFileReader (5.0.7) and searched using the search

engine Sequest or Comet (for TMT-01 and TM-02 respectively)(Eng et al., 2008; Eng et al.,

2013; Eng et al., 1994) against a mouse target-decoy protein database (downloaded from

UniProt in June 2019) that included the most common contaminants. Precursor ion tolerance

was set at 20 ppm and product ion tolerance at 1 Da. Cysteine carbamidomethylation (+57.0215

Da) and TMT tag (+229.1629 Da or +304.2071 Da for TMT-6plex or TMTpro-16pex,

respectively) on lysine residues and peptide N-termini were set as static modifications. Up to 2

variable methionine oxidations (+15.9949 Da) and 2 miss cleavages were allowed in the

searches. Peptide-spectrum matches (PSMs) were adjusted to a 1% FDR with a linear

discriminant analysis(Huttlin et al., 2010) and proteins were further collapsed to a final protein-

level FDR of 1%. TMT quantitative values we obtained from MS3 scans. Only those with a

signal-to-noise ratio > 100 and an isolation specificity > 0.7 were used for quantification. Each

TMT was normalized to the total signal in each column. Quantifications included in Table S4 are

represented as relative abundances. RAW files will be made available upon request. The data

have been deposited in the ProteomeXchange Consortium via the PRIDE(Vizcaíno et al., 2014)

partner repository with the data set identifier PXD040834. Biological pathway and gene ontology

enrichment analysis were performed using the ClueGo (Cytoscape plug-in), Enrichr or STRING.

1278

SYTOX Green assay for measuring cell death

The nucleic acid dye SYTOX Green (ThermoFisher Scientific) was diluted to 100 µM in DMSO

and stored at -20°C. Human fibroblasts were seeded at 5 x 103 per well of a 96-well plate

treated with DMSO or LLOME at a series of concentrations to evaluate the lysosome-mediated

cell death or pre-treated with C381 followed by 2 mM LLOME to evaluate the rescue of

lysosome-mediated cell death. The uptake of membrane-impermeant SYTOX Green dye occurs

while cell membrane is compromised. The fluorescent nucleic acid-intercalating dyes then emit

a significant increase in fluorescence intensity as dead-cell markers. SYTOX Green staining

solution was prepared at a final concentration of 100 nM in HBSS and incubated with cells for

30 min at 37°C. After rinse once with PBS, cells were monitored for SYTOX Green intensity

changes in CLARIOstar plate reader. Then, the cells were rinsed once with PBS, counterstained

with 5 µg/mL Hoechst (ThermoFisher Scientific) for 10 min at room temperature and rinsed with

PBS. The measurement of Hoechst fluorescence is performed in plate reader and allowed

SYTOX Green levels to be normalized to the cell density.

1293

Caspase-3/7 activation

To detect neuronal apoptosis, we added the Caspase-3/7 substrate FAM-DEVD-FMK

(ImmunoChemistry), one of the fluorochrome-labeled inhibitors of caspases that covalently and

irreversibly binds to the active caspases. The green fluorescence intensity is a direct

measurement of Caspase-3/7 activity. Human tNeurons were treated with DMSO or 0.5 mM

LLOME for 1 hr to measure the lysosome-mediated apoptosis or pretreated with 3.1 µM C381

followed by LLOME treatment to assess the rescuing effect on lysosome-mediated apoptosis.

The FAM-DEVD-FMK reagent was reconstituted in DMSO and stored at -20°C. When cells

were available, the FAM-DEVD-FMK reagent was diluted with PBS 1:5 ratio and added to cell

culture medium at a dilution of 1:30 to form 1X staining solution. Cells were incubated with the

FAM-DEVD-FMK solution for 30 min at 37°C. After a rinse by Apoptosis Buffer, the cells were

fixed by 4% PFA for 15 min and counterstained with 5 µg/mL Hoechst for confocal microscope

imaging.

1307

Lysotracker

LysoTracker Red DND-99 was reconstituted in DMSO and stored at -20°C. Human adult

fibroblasts were seeded at 5 x 103 per well of a 96-well plate and pre-incubated with 0.1 µM

LysoTracker Red for 30 min at 37°C. After washing off excess dye by PBS, cells were treated

with 1 mM LLOME for up to 60 min to evaluate the lysosomal activity. After a wash by PBS,

cells were prepared for imaging by CLARIOstar plate reader.

1314

Fluorescence-conjugated Dextran assay for measuring endocytosis and lysosomal

acidification

Fluorescein isothiocyanate (FITC)-conjugated Dextran at 40 kDa and tetramethylrhodamine

(TMR)-conjugated Dextran at 70 kDa (ThermoFisher Scientific) were reconstituted in H2O and

stored at -20°C. Human adult fibroblasts were seeded at 4 x 104 per well of a 24-well plate or 5

x 103 per well of a 96-well plate. Cells were incubated with FITC- and TMR-Dextran at 0.5

mg/ml for 4 hr at 37°C and rinsed by PBS, followed by 20-hr chase in fresh culture medium to

accumulate Dextran in the late endosomes and lysosomes. Then, cells were treated with DMSO

or 1 mM LLOME for 30 min and washed out to assess the cellular endocytosis and lysosomal

recovery function, or pre-treated with 6.25 µM C381 for 24 hr, followed by LLOME treatment and

washout to evaluate the rescuing effect of C381 in fibroblasts. The cells were fixed by 4%

paraformaldehyde (PFA) for 15 min and prepared for imaging by a confocal microscope and

CLARIOstar plate reader.

1328

Magic Red Cathepsin-B assay for measuring lysosomal proteolysis

Magic Red Cathepsin-B substrate, MR-(RR)2, containing arginine-arginine (RR) sequence was

reconstituted in DMSO and stored at -20°C. When cells were available for experiment, MR-

(RR)2 was diluted with H2O 1:10 ratio and added to cell culture medium at a dilution of 1:25 to

form 1X staining solution. Active Cathepsin-B cleaves MR-(RR)2 and emits fluorescence with

optimal excitation of 592 nm and emission of 628 nm. To test lysosomal proteolytic capacity,

human adult fibroblasts were incubated MR-(RR)2 staining solution with for 30 min at 37°C,

followed by DMSO or 1 mM LLOME treatment for 30 min and washout for lysosomal recovery

experiment. To evaluate the pharmacological rescuing effect, the C381 treatment was 24 hr

before the cells were loaded with MR-(RR)2. Cells were treated with 1 mM LLOME for 30 min

and washed out, and then fixed by 4% PFA for 15 min for imaging by a confocal microscope

and CLARIOstar plate reader.

1341

Mice brain perfusion and tissue processing

Mice were anaesthetized with 2.5% (v/v) Avertin (Sigma-Aldrich). Transcardial perfusion with

50 ml cold PBS was performed using a peristaltic pump with the perfusate flow rate not

exceeding 10 ml/min. Brain tissue processing was performed as described previously(Vest et

al., 2022; Villeda et al., 2014). Hemibrains were isolated and fixed in 4% PFA overnight at 4°C

before transferring to 30% sucrose in PBS at 4 °C for preservation. Hemibrains were

cryosectioned coronally at a thickness of 40 μm on a freezing–sliding microtome, and the free-

floating sections were stored in cryoprotectant (40% PBS, 30% glycerol, 30% ethylene glycol)

and kept at -20 °C until staining.

1351

Immunofluorescence and image acquisition and analysis

Human adult fibroblasts were fixed with 4% PFA for 15 minutes at room temperature,

permeabilized with 0.3% Triton X-100 in PBS for 5 minutes or cold Methanol on ice for 20

minutes, blocked with pre-warmed 5% bovine serum albumin (BSA) in PBS and shook for 45

min at room temperature. Fibroblasts were incubated with primary antibody solution in 5% BSA

overnight at 4°C in a moisture chamber: anti-ALIX (1:200; Santa Cruz, sc-53540), anti-Amyloid-

β (1:100; Cell Signaling, 8243), anti-Amyloid-β (1-42) (1:200; Enzo Life Sciences, ADI-905-804-

100), anti-CHMP1A (1:200; Santa Cruz, sc-271617), anti-CHMP2B (1:400; Proteintech, 12527-

1-AP), anti-CHMP6 (1:200; Santa Cruz, sc-398963), anti-Galectin-3 (1:1000; Biolegend,

125401), anti-γ-H2AX (1:200; Millipore Sigma, 05-636), anti-H3K9me3 (1:500; Abcam, ab8898),

anti-H4K16ac (1:300; Thermo Fisher Scientific, MA5-27794), anti-Hsp70 (1:1000; Abcam,

ab45133), anti-LAMP1 (1:2000; Cell Signaling, 9091), anti-LAMP2 (1:200; DSHB, H4B4), anti-

LC3B (1:200; Cell Signaling, 2775)., anti-Nup98 (1:500; Cell Signaling, 2598), anti-

p62/SQSTM1 (1:400; Abcam, ab56416), anti- S100A4 (1:200; Abcam, ab124805), anti-TDP-43

(1:1000; Proteintech, 12892-1-AP), anti-TFEB (1:100; Proteintech, 13372-1-AP), anti-TSG101

(1:200; Santa Cruz, sc-7964), anti-Ubiquitin (1:200; LifeSensors, AB120), and anti-Vimentin

(1:400; Cell Signaling, 5741). Fibroblasts were washed with PBS three times and incubated with

fluorophore-conjugated secondary antibody solution in the dark for 1 hr at room temperature:

anti-mouse, anti-rabbit and anti-rat (1:500, Thermo Fisher Scientific). Finally, fibroblasts were

washed with PBS three times, mounted with DAPI containing ProLong Glass Antifade Mountant

(Thermo Fisher Scientific) and air-dried for overnight prior to imaging.

1373

Human tNeurons were undergone two-step fixation: 1) remove half the culture medium and add

equivalent to half the volume of 4% PFA for 5 min at room temperature; 2) remove the old

solution and add 4% PFA for 15 min at room temperature. The tNeurons were permeabilized

with 0.3% Triton X-100 in PBS for 5 minutes at room temperature or cold Methanol on ice for 20

minutes, blocked with pre-warmed 5% bovine serum albumin (BSA) in PBS and shook for 45

min at room temperature. Fibroblasts were incubated with primary antibody solution in 5% BSA

overnight at 4°C in a moisture chamber: anti-ASC/PYCARD (1:100; Santa Cruz, sc-514414),

anti-GAP-43 (1:300; Novus Biologicals, NB300-143), anti-HGS (1:200; Genetex, GTX101718),

anti-MAP2 (1:1000; Biolegend, 822501), anti-NeuN (1:500; Abcam, ab177487), anti-NLRP3

(1:200; R&D systems, MAB7578), anti-pTau (AT8) (1:100; Thermo Fisher Scientific, MN1020),

anti-pTDP-43 Ser409/410 (1:300; Cosmo Bio, CAC-TIP-PTD-M01), anti-Synapsin-1 (Syn-1)

(1:200; Abcam, ab64581), anti-Tau (1:100; Aves Labs), and anti-Beta tubulin III (Tuj1) (1:500;

Neuromics, CH23005 & 1:1000; Biolegend, 801201). The tNeurons were washed with PBS

three times and incubated with fluorophore-conjugated secondary antibody solution in the dark

for 1 hr at room temperature: anti-mouse, anti-rabbit and anti-rat, and anti-chicken (1:500,

Thermo Fisher Scientific), washed with PBS three times, mounted with DAPI containing

ProLong Glass Antifade Mountant (Thermo Fisher Scientific) and air-dried for overnight prior to

imaging.

1392

Free-floating mouse tissue sections at 30-50 μm were collected and proceed for

immunostaining in multi-well plates. Mouse tissue sections were first rinsed with PBS three

times and permeabilized with 0.3% Triton X-100 in PBS for 20 min at room temperature,

followed by incubation with blocking buffer (10% Normal Donkey Serum and 0.03% Triton-X-

100 in PBS) and shook for 1 hr at room temperature. After PBS rinsing, the tissue sections were

incubated with primary antibody solution in 10% Normal Donkey Serum in PBS overnight at 4°C

in a moisture chamber: anti-Amyloid-β (1-42) (1:100; Enzo Life Sciences, ADI-905-804-100),

anti-CHMP2B (1:100; Proteintech, 12527-1-AP), anti-Galectin-3 (1:50; R&D systems, AF1197),

anti-Hsp70 (1:300; Abcam, ab45133), anti-LAMP1 (1:50; Santa Cruz, sc-19992), and anti-MAP2

(1:500; Biolegend, 822501). Fluorophore-conjugated secondary antibody solution was

incubated for 1 hr at room temperature: anti-mouse, anti-rabbit, anti-goat and anti-chicken

(1:500, Thermo Fisher Scientific). Tissue sections were washed with PBS two times,

counterstained with Hoechst (1:2000) and washed with PBS two times again before mounting

with ProLong Glass Antifade Mountant (Thermo Fisher Scientific). Samples were air-dried for

overnight prior to imaging.

1408

Paraffin-embedded human brain tissues of the cerebral cortex were sectioned at 10 μm

thickness. Deparaffinization was achieved by washing slides through xylenes twice, each for 5

min and rehydrating via gradient ethanol (100%　95%　70%　50%) into water, each for 10

min. Following heat-mediated antigen retrieval using Citrate buffer, pH 6.0 (Sigma-Aldrich) for

30 min at 95° C, tissue sections were rinsed with PBS once and incubated with blocking buffer

(10% Normal Donkey Serum and 0.03% Triton-X-100 in PBS) for 2 hr at room temperature.

Slides were incubated with antibody cocktails in blocking buffer overnight at 4°C in a moisture

chamber: Amyloid-β (6E10) (1:100; Biolegend, 803001), anti-CHMP2B (1:100; GeneTex,

GTX118181), anti-Galectin-3 (1:50; R&D systems, AF1197 & 1:100; Biolegend, 125401), anti-

Hsp70 (1:300; Abcam, ab45133), anti-LAMP2 (1:200; Abcam, ab213294), and anti-MAP2

(1:500; Biolegend, 822501). Tissue sections were incubated with fluorophore-conjugated

secondary antibody solution for 1 hr at room temperature: anti-mouse, anti-rabbit, anti-rat, anti-

goat and anti-chicken (1:500, Thermo Fisher Scientific). After a PBS rinse, slides were

counterstained with Hoechst (1:2000) and washed with PBS two times. The complete air-dried

tissue sections were mounted with ProLong Glass Antifade Mountant (Thermo Fisher Scientific)

and subjected to dry in the dark overnight prior to imaging.

1425

Imaging was acquired at Z-series (10-30 sections, 0.2-1 μm steps) according to experimental

paradigm using Zeiss LSM 700 confocal fluorescence microscope with 20x, 63x and 100x

objectives. In each experiment, all groups were subjected to image using the same acquisition

settings. The z-stack images were performed Maximum Intensity Projection to analyze the

mean pixel intensity and determine a threshold to quantify puncta number in the cells by Fiji. For

quantitative histology, three to five separate sections were sampled using a 20x objective and

fluorescence signals were measured from entire image field to the mean fluorescence change.

Tissue sections were imaged and analyzed by blinded observers.

1434

Transmission electron microscopy

Cells were grown on Ibidi dishes: μ-Dish 35 mm, high Grid-50 Glass Bottom is a 35 mm then

fixed in Karnovsky's fixative: 2% Glutaraldehyde (EMS Cat# 16000) and 4% pFormaldehyde

(EMS Cat# 15700) in 0.1M Sodium Cacodylate (EMS Cat# 12300) pH 7.4 for 1 hr, chilled and

sent to Stanford's CSIF on ice. They were then post-fixed in cold 1% Osmium tetroxide (EMS

Cat# 19100) in water and allowed to warm for 2 hr in a hood, washed 3X with ultra-filtered

water, then en bloc stained 2 hr in 1% Uranyl Acetate at room temperature. Samples were then

dehydrated in a series of ethanol washes for 10 min each at room temperature beginning at

30%, 50%, 70%, 95%, changed to 100% 2X, then Propylene Oxide (PO) for 10 min. Samples

were infiltrated with EMbed-812 resin (EMS Cat#14120) mixed 1:1, and 2:1 with PO for 2 hr

each. The samples were then placed into EMbed-812 for 2 hours opened then placed into flat

molds w/labels and fresh resin and placed into 65o C oven overnight. Cells of interest were

located using the grid pattern and cut out with a gem saw and remounted on pre-labeled resin

blocks with fresh resin and polymerized overnight again. Once full polymerized the glass

coverslip was etched away using hydrofluoric acid for 20 min. Using the finder grid pattern left

behind the block faces were trimmed down allowing for serial sectioning of the cells of interest.

Sections were taken around 90 nm, picked up on formvar/Carbon coated slot Cu grids, stained

for 40seconds in 3.5% Uranyl Acetate in 50% Acetone followed by staining in 0.2% Lead Citrate

for 6 min. Observed in the JEOL JEM-1400 120kV and photos were taken using a Gatan Orius

2k X 2k digital camera.

1455

Cytokine profiling analysis on neuronal conditioned medium using Luminex multiplex
analysis

Secretion of inflammatory factors was analyzed using cytokine profiling of the conditioned

medium from tNeurons of healthy donors and AD patients. Conditioned medium was collected

48 hr after the last medium change in a 12-well plate with 1 mL of neuronal maturation medium

at PID 38, and centrifuged at 10,000 x G for 10 min at room temperature to pellet out

particulates. For Human 80 plex panel (EMD-Millipore), a minimum of 200 μL of supernatants

was stored at -80°C. Cell free medium was also collected to monitor the background

fluorescence. Cell numbers were determined by an automated cell counter for normalization of

cytokine levels. The setup of cytokine profiling assay was performed according to the

manufacturer's instructions. Briefly, samples were mixed with antibody-linked magnetic beads

on a 96-well plate and incubated overnight at 4°C with shaking. Cold and room temperature

incubation steps were performed on an orbital shaker at 500-600 rpm. Plates were washed

twice with wash buffer in a Biotek ELx405 washer. Following one hr incubation at room

temperature with biotinylated detection antibody, streptavidin-PE was added for 30 minutes with

shaking. Plates were washed as above and PBS added to wells for reading in the Luminex

FlexMap3D Instrument with a lower bound of 50 beads per sample per cytokine. Each sample

was measured in duplicate. Custom Assay Chex control beads were purchased from Radix

Biosolutions, Georgetown, Texas, and are added to all wells. The analyses of all conditioned

medium samples were performed using raw data (mean fluorescence intensity (MFI)) rather

than concentration (pg/mL) to avoid calculating bias per as per recommendation of the Stanford

Human Immune Monitoring Center.

1478

CSF samples and protein discovery

We used the SOMAScan assay platform(Gold et al., 2010; Kim et al., 2018a) (SomaLogic Inc.)

to measure the relative levels of 76 human proteins in CSF. This platform is based on modified

single-stranded DNA aptamers (SOMAmer) capable of binding to specific protein targets with

high sensitivity and specificity. We collected 79 CSF samples (50 HC and 29 AD samples) from

a multi-ethnic cohort of older American adults (age range 60-87 years) between 2015 and 2020.

Samples were stored at -80 °C and 150 µl aliquots of CSF were sent on dry ice to SomaLogic.

CSF samples were analyzed via SOMAScan assay in five batches. To account for variation

within and across batches, control, calibrator and buffer samples are added in each 96-well

plate. Data normalization was conducted by the manufacturer following three stages. First, in

Hybridization Control Normalization, hybridization control probes are used to remove individual

sample variance. Second, Intraplate Median Signal Normalization, median normalization

removed inter-sample differences within the plate. Last, Plate Scaling and Calibration, this final

step removed variance across assay runs.

1493

QUANTIFICATION AND STATISTICAL ANALYSIS

Quantification of fluorescence images was performed by CLARIOstar plate reader software and

open-source Fiji software. For each technical replicate, the fluorescence intensity of the

background from cell-free solution or cell-free area in the image field was subtracted from

intensity measurements. For most experiments, a power analysis was performed to pre-

determine the sample sizes based on results from initial experiments. For quantifications of

cytokine levels and human and mouse brain samples, the experiments were blinded. Statistical

analysis was performed with unpaired t-test, one- way ANOVA or two-way ANOVA based on

the experimental design using GraphPad Prism Software. All values were expressed as the

box-and-whisker plots or mean ± SD. Differences between two groups were analyzed using

two-sided unpaired t-test with Welch's correction. Differences between multiple groups were

analyzed using one- way or two-way ANOVA followed by Bonferroni post-hoc analysis.

Differences were considered statistically significant for P values < 0.05.

1507

REFERENCES:

Aits, S., Kricker, J., Liu, B., Ellegaard, A.M., Hamalisto, S., Tvingsholm, S., Corcelle-Termeau, E., Hogh, S., Farkas, T., Holm Jonassen, A., et al. (2015). Sensitive detection of lysosomal membrane permeabilization by lysosomal galectin puncta assay. Autophagy 11, 1408-1424.

Alzheimer's Association (2020). 2020 Alzheimer's disease facts and figures. Alzheimers Dement.

Amador-Ortiz, C., Lin, W.L., Ahmed, Z., Personett, D., Davies, P., Duara, R., Graff-Radford, N.R., Hutton, M.L., and Dickson, D.W. (2007). TDP-43 immunoreactivity in hippocampal sclerosis and Alzheimer's disease. Annals of neurology 61, 435-445.

Anding, A.L., and Baehrecke, E.H. (2017). Cleaning House: Selective Autophagy of Organelles. Developmental cell 41, 10-22.

Bettcher, B.M., Johnson, S.C., Fitch, R., Casaletto, K.B., Heffernan, K.S., Asthana, S., Zetterberg, H., Blennow, K., Carlsson, C.M., Neuhaus, J., et al. (2018). Cerebrospinal Fluid and Plasma Levels of Inflammation Differentially Relate to CNS Markers of Alzheimer's Disease Pathology and Neuronal Damage. Journal of Alzheimer's disease : JAD 62, 385-397.

Birnbaum, J.H., Wanner, D., Gietl, A.F., Saake, A., Kundig, T.M., Hock, C., Nitsch, R.M., and Tackenberg, C. (2018). Oxidative stress and altered mitochondrial protein expression in the absence of amyloid-beta and tau pathology in iPSC-derived neurons from sporadic Alzheimer's disease patients. Stem cell research 27, 121-130.

Bourdenx, M., Daniel, J., Genin, E., Soria, F.N., Blanchard-Desce, M., Bezard, E., and Dehay, B. (2016). Nanoparticles restore lysosomal acidification defects: Implications for Parkinson and other lysosomal-related diseases. Autophagy 12, 472-483.

Boza-Serrano, A., Ruiz, R., Sanchez-Varo, R., Garcia-Revilla, J., Yang, Y., Jimenez-Ferrer, I., Paulus, A., Wennstrom, M., Vilalta, A., Allendorf, D., et al. (2019). Galectin-3, a novel endogenous TREM2 ligand, detrimentally regulates inflammatory response in Alzheimer's disease. Acta neuropathologica.

Caiazzo, M., Dell'Anno, M.T., Dvoretskova, E., Lazarevic, D., Taverna, S., Leo, D., Sotnikova, T.D., Menegon, A., Roncaglia, P., Colciago, G., et al. (2011). Direct generation of functional dopaminergic neurons from mouse and human fibroblasts. Nature 476, 224-227.

Cao, C., Prado, M.A., Sun, L., Rockowitz, S., Sliz, P., Paulo, J.A., Finley, D., and Fleming, M.D. (2021a). Maternal Iron Deficiency Modulates Placental Transcriptome and Proteome in Mid-Gestation of Mouse Pregnancy. The Journal of nutrition 151, 1073-1083.

Cao, M., Luo, X., Wu, K., and He, X. (2021b). Targeting lysosomes in human disease: from basic research to clinical applications. Signal Transduct Target Ther 6, 379.

Capano, L.S., Sato, C., Ficulle, E., Yu, A., Horie, K., Kwon, J.S., Burbach, K.F., Barthelemy, N.R., Fox, S.G., Karch, C.M., et al. (2022). Recapitulation of endogenous 4R tau expression and formation of insoluble tau in directly reprogrammed human neurons. Cell Stem Cell 29, 918-932 e918.

Chakravorty, A., Jetto, C.T., and Manjithaya, R. (2019). Dysfunctional Mitochondria and Mitophagy as Drivers of Alzheimer's Disease Pathogenesis. Front Aging Neurosci 11, 311.

Chen, J.J., Nathaniel, D.L., Raghavan, P., Nelson, M., Tian, R., Tse, E., Hong, J.Y., See, S.K., Mok, S.A., Hein, M.Y., et al. (2019). Compromised function of the ESCRT pathway promotes

endolysosomal escape of tau seeds and propagation of tau aggregation. The Journal of biological chemistry 294, 18952-18966.

Chou, C.C., Zhang, Y., Umoh, M.E., Vaughan, S.W., Lorenzini, I., Liu, F., Sayegh, M., Donlin-Asp, P.G., Chen, Y.H., Duong, D.M., et al. (2018). TDP-43 pathology disrupts nuclear pore complexes and nucleocytoplasmic transport in ALS/FTD. Nature neuroscience 21, 228-239.

Chung, C.Y., Shin, H.R., Berdan, C.A., Ford, B., Ward, C.C., Olzmann, J.A., Zoncu, R., and Nomura, D.K. (2019). Covalent targeting of the vacuolar H(+)-ATPase activates autophagy via mTORC1 inhibition. Nature chemical biology 15, 776-785.

Cortes, C.J., and La Spada, A.R. (2019). TFEB dysregulation as a driver of autophagy dysfunction in neurodegenerative disease: Molecular mechanisms, cellular processes, and emerging therapeutic opportunities. Neurobiology of disease 122, 83-93.

Craig, E.A. (2018). Hsp70 at the membrane: driving protein translocation. BMC Biol 16, 11.

Cuervo, A.M., and Wong, E. (2014). Chaperone-mediated autophagy: roles in disease and aging. Cell Res 24, 92-104.

DeTure, M.A., and Dickson, D.W. (2019). The neuropathological diagnosis of Alzheimer's disease. Molecular neurodegeneration 14, 32.

Ditaranto, K., Tekirian, T.L., and Yang, A.J. (2001). Lysosomal membrane damage in soluble Abeta-mediated cell death in Alzheimer's disease. Neurobiology of disease 8, 19-31.

Douglas, P.M., and Dillin, A. (2010). Protein homeostasis and aging in neurodegeneration. The Journal of cell biology 190, 719-729.

Eapen, V.V., Swarup, S., Hoyer, M.J., Paulo, J.A., and Harper, J.W. (2021). Quantitative proteomics reveals the selectivity of ubiquitin-binding autophagy receptors in the turnover of damaged lysosomes by lysophagy. Elife 10.

Eftekharzadeh, B., Daigle, J.G., Kapinos, L.E., Coyne, A., Schiantarelli, J., Carlomagno, Y., Cook, C., Miller, S.J., Dujardin, S., Amaral, A.S., et al. (2018). Tau Protein Disrupts Nucleocytoplasmic Transport in Alzheimer's Disease. Neuron 99, 925-940.e927.

Eng, J.K., Fischer, B., Grossmann, J., and Maccoss, M.J. (2008). A fast SEQUEST cross correlation algorithm. Journal of proteome research 7, 4598-4602.

Eng, J.K., Jahan, T.A., and Hoopmann, M.R. (2013). Comet: an open-source MS/MS sequence database search tool. Proteomics 13, 22-24.

Eng, J.K., McCormack, A.L., and Yates, J.R. (1994). An approach to correlate tandem mass spectral data of peptides with amino acid sequences in a protein database. J Am Soc Mass Spectrom 5, 976-989.

Faizi, M., Bader, P.L., Saw, N., Nguyen, T.V., Beraki, S., Wyss-Coray, T., Longo, F.M., and Shamloo, M. (2012). Thy1-hAPP(Lond/Swe+) mouse model of Alzheimer's disease displays broad behavioral deficits in sensorimotor, cognitive and social function. Brain Behav 2, 142-154.

Fruhmann, G., Marchal, C., Vignaud, H., Verduyckt, M., Talarek, N., De Virgilio, C., Winderickx, J., and Cullin, C. (2018). The Impact of ESCRT on Abeta1-42 Induced Membrane Lesions in a Yeast Model for Alzheimer's Disease. Frontiers in molecular neuroscience 11, 406.

Galluzzi, L., Baehrecke, E.H., Ballabio, A., Boya, P., Bravo-San Pedro, J.M., Cecconi, F., Choi, A.M., Chu, C.T., Codogno, P., Colombo, M.I., et al. (2017). Molecular definitions of autophagy and related processes. The EMBO journal 36, 1811-1836.

Gold, L., Ayers, D., Bertino, J., Bock, C., Bock, A., Brody, E.N., Carter, J., Dalby, A.B., Eaton,
B.E., Fitzwater, T., et al. (2010). Aptamer-based multiplexed proteomic technology for biomarker
discovery. PloS one 5, e15004.
Gottschling, D.E., and Nyström, T. (2017). The Upsides and Downsides of Organelle
Interconnectivity. Cell 169, 24-34.
Gouras, G.K., Tsai, J., Naslund, J., Vincent, B., Edgar, M., Checler, F., Greenfield, J.P.,
Haroutunian, V., Buxbaum, J.D., Xu, H., et al. (2000). Intraneuronal Abeta42 accumulation in
human brain. The American journal of pathology 156, 15-20.
Gowrishankar, S., Yuan, P., Wu, Y., Schrag, M., Paradise, S., Grutzendler, J., De Camilli, P.,
and Ferguson, S.M. (2015). Massive accumulation of luminal protease-deficient axonal
lysosomes at Alzheimer's disease amyloid plaques. Proceedings of the National Academy of
Sciences of the United States of America 112, E3699-3708.
Grubman, A., Chew, G., Ouyang, J.F., Sun, G., Choo, X.Y., McLean, C., Simmons, R.K.,
Buckberry, S., Vargas-Landin, D.B., Poppe, D., et al. (2019). A single-cell atlas of entorhinal
cortex from individuals with Alzheimer's disease reveals cell-type-specific gene expression
regulation. Nature neuroscience 22, 2087-2097.
Grubman, A., Choo, X.Y., Chew, G., Ouyang, J.F., Sun, G., Croft, N.P., Rossello, F.J.,
Simmons, R., Buckberry, S., Landin, D.V., et al. (2021). Transcriptional signature in microglia
associated with Abeta plaque phagocytosis. Nature communications 12, 3015.
Hansson, J., Rafiee, M.R., Reiland, S., Polo, J.M., Gehring, J., Okawa, S., Huber, W.,
Hochedlinger, K., and Krijgsveld, J. (2012). Highly coordinated proteome dynamics during
reprogramming of somatic cells to pluripotency. Cell reports 2, 1579-1592.
Harvey, R.J., Skelton-Robinson, M., and Rossor, M.N. (2003). The prevalence and causes of
dementia in people under the age of 65 years. Journal of neurology, neurosurgery, and
psychiatry 74, 1206-1209.
Heneka, M.T., Kummer, M.P., Stutz, A., Delekate, A., Schwartz, S., Vieira-Saecker, A., Griep,
A., Axt, D., Remus, A., Tzeng, T.C., et al. (2013). NLRP3 is activated in Alzheimer's disease
and contributes to pathology in APP/PS1 mice. Nature 493, 674-678.
Herdy, J.R., Traxler, L., Agarwal, R.K., Karbacher, L., Schlachetzki, J.C.M., Boehnke, L.,
Zangwill, D., Galasko, D., Glass, C.K., Mertens, J., et al. (2022). Increased post-mitotic
senescence in aged human neurons is a pathological feature of Alzheimer's disease. Cell Stem
Cell 29, 1637-1652.e1636.
Hipp, M.S., Park, S.H., and Hartl, F.U. (2014). Proteostasis impairment in protein-misfolding and
-aggregation diseases. Trends Cell Biol 24, 506-514.
Hondius, D.C., Koopmans, F., Leistner, C., Pita-Illobre, D., Peferoen-Baert, R.M., Marbus, F.,
Paliukhovich, I., Li, K.W., Rozemuller, A.J.M., Hoozemans, J.J.M., et al. (2021). The proteome
of granulovacuolar degeneration and neurofibrillary tangles in Alzheimer's disease. Acta
neuropathologica 141, 341-358.
Hughes, A.L., and Gottschling, D.E. (2012). An early age increase in vacuolar pH limits
mitochondrial function and lifespan in yeast. Nature 492, 261-265.
Huh, C.J., Zhang, B., Victor, M.B., Dahiya, S., Batista, L.F., Horvath, S., and Yoo, A.S. (2016).
Maintenance of age in human neurons generated by microRNA-based neuronal conversion of
fibroblasts. Elife 5.

Hung, C.O.Y., and Livesey, F.J. (2018). Altered gamma-Secretase Processing of APP Disrupts
Lysosome and Autophagosome Function in Monogenic Alzheimer's Disease. Cell reports 25,
3647-3660 e3642.
Huttlin, E.L., Jedrychowski, M.P., Elias, J.E., Goswami, T., Rad, R., Beausoleil, S.A., Villén, J.,
Haas, W., Sowa, M.E., and Gygi, S.P. (2010). A tissue-specific atlas of mouse protein
phosphorylation and expression. Cell 143, 1174-1189.
Israel, M.A., Yuan, S.H., Bardy, C., Reyna, S.M., Mu, Y., Herrera, C., Hefferan, M.P., Van Gorp,
S., Nazor, K.L., Boscolo, F.S., et al. (2012). Probing sporadic and familial Alzheimer's disease
using induced pluripotent stem cells. Nature 482, 216-220.
Jiang, P., Gan, M., Yen, S.H., McLean, P.J., and Dickson, D.W. (2017). Impaired endo-
lysosomal membrane integrity accelerates the seeding progression of alpha-synuclein
aggregates. Scientific reports 7, 7690.
Jimenez, A.J., Maiuri, P., Lafaurie-Janvore, J., Divoux, S., Piel, M., and Perez, F. (2014).
ESCRT machinery is required for plasma membrane repair. Science (New York, NY) 343,
1247136.
Josephs, K.A., Murray, M.E., Whitwell, J.L., Parisi, J.E., Petrucelli, L., Jack, C.R., Petersen,
R.C., and Dickson, D.W. (2014). Staging TDP-43 pathology in Alzheimer's disease. Acta
neuropathologica 127, 441-450.
Kaushik, S., and Cuervo, A.M. (2015). Proteostasis and aging. Nature medicine 21, 1406-1415.
Kim, C.H., Tworoger, S.S., Stampfer, M.J., Dillon, S.T., Gu, X., Sawyer, S.J., Chan, A.T.,
Libermann, T.A., and Eliassen, A.H. (2018a). Stability and reproducibility of proteomic profiles
measured with an aptamer-based platform. Scientific reports 8, 8382.
Kim, Y., Zheng, X., Ansari, Z., Bunnell, M.C., Herdy, J.R., Traxler, L., Lee, H., Paquola, A.C.M.,
Blithikioti, C., Ku, M., et al. (2018b). Mitochondrial Aging Defects Emerge in Directly
Reprogrammed Human Neurons due to Their Metabolic Profile. Cell reports 23, 2550-2558.
Kirkegaard, T., Roth, A.G., Petersen, N.H., Mahalka, A.K., Olsen, O.D., Moilanen, I., Zylicz, A.,
Knudsen, J., Sandhoff, K., Arenz, C., et al. (2010). Hsp70 stabilizes lysosomes and reverts
Niemann-Pick disease-associated lysosomal pathology. Nature 463, 549-553.
Köhler, C. (2016). Granulovacuolar degeneration: a neurodegenerative change that
accompanies tau pathology. Acta neuropathologica 132, 339-359.
Kravić, B., Bionda, T., Siebert, A., Gahlot, P., Levantovsky, S., Behrends, C., and Meyer, H.
(2022). Ubiquitin profiling of lysophagy identifies actin stabilizer CNN2 as a target of VCP/p97
and uncovers a link to HSPB1. Molecular cell 82, 2633-2649.e2637.
Kwart, D., Gregg, A., Scheckel, C., Murphy, E.A., Paquet, D., Duffield, M., Fak, J., Olsen, O.,
Darnell, R.B., and Tessier-Lavigne, M. (2019). A Large Panel of Isogenic APP and PSEN1
Mutant Human iPSC Neurons Reveals Shared Endosomal Abnormalities Mediated by APP β-
CTFs, Not Aβ. Neuron 104, 256-270.e255.
Labbadia, J., and Morimoto, R.I. (2015). The biology of proteostasis in aging and disease.
Annual review of biochemistry 84, 435-464.
Lackie, R.E., Maciejewski, A., Ostapchenko, V.G., Marques-Lopes, J., Choy, W.Y., Duennwald,
M.L., Prado, V.F., and Prado, M.A.M. (2017). The Hsp70/Hsp90 Chaperone Machinery in
Neurodegenerative Diseases. Frontiers in neuroscience 11, 254.
Lagomarsino, V.N., Pearse, R.V., 2nd, Liu, L., Hsieh, Y.C., Fernandez, M.A., Vinton, E.A., Paull,
D., Felsky, D., Tasaki, S., Gaiteri, C., et al. (2021). Stem cell-derived neurons reflect features of

protein networks, neuropathology, and cognitive outcome of their aged human donors. Neuron
109, 3402-3420.e3409.

Laskowska, E., Kuczynska-Wisnik, D., and Lipinska, B. (2019). Proteomic analysis of protein
homeostasis and aggregation. J Proteomics 198, 98-112.

Lee, J.H., McBrayer, M.K., Wolfe, D.M., Haslett, L.J., Kumar, A., Sato, Y., Lie, P.P., Mohan, P.,
Coffey, E.E., Kompella, U., et al. (2015). Presenilin 1 Maintains Lysosomal Ca(2+) Homeostasis
via TRPML1 by Regulating vATPase-Mediated Lysosome Acidification. Cell reports 12, 1430-
1444.

Lee, J.H., Yang, D.S., Goulbourne, C.N., Im, E., Stavrides, P., Pensalfini, A., Chan, H.,
Bouchet-Marquis, C., Bleiwas, C., Berg, M.J., et al. (2022). Faulty autolysosome acidification in
Alzheimer's disease mouse models induces autophagic build-up of Aβ in neurons, yielding
senile plaques. Nature neuroscience 25, 688-701.

Lee, J.H., Yu, W.H., Kumar, A., Lee, S., Mohan, P.S., Peterhoff, C.M., Wolfe, D.M., Martinez-
Vicente, M., Massey, A.C., Sovak, G., et al. (2010). Lysosomal proteolysis and autophagy
require presenilin 1 and are disrupted by Alzheimer-related PS1 mutations. Cell 141, 1146-
1158.

Liu, X., Li, J., Gygi, S.P., and Paulo, J.A. (2022). Profiling Yeast Deletion Strains Using Sample
Multiplexing and Network-Based Analyses. Journal of proteome research 21, 1525-1536.

Logan, T., Simon, M.J., Rana, A., Cherf, G.M., Srivastava, A., Davis, S.S., Low, R.L.Y., Chiu,
C.L., Fang, M., Huang, F., et al. (2021). Rescue of a lysosomal storage disorder caused by Grn
loss of function with a brain penetrant progranulin biologic. Cell 184, 4651-4668 e4625.

Maejima, I., Takahashi, A., Omori, H., Kimura, T., Takabatake, Y., Saitoh, T., Yamamoto, A.,
Hamasaki, M., Noda, T., Isaka, Y., et al. (2013). Autophagy sequesters damaged lysosomes to
control lysosomal biogenesis and kidney injury. The EMBO journal 32, 2336-2347.

Mahmoudi, S., and Brunet, A. (2012). Aging and reprogramming: a two-way street. Current
opinion in cell biology 24, 744-756.

Mansuroglu, Z., Benhelli-Mokrani, H., Marcato, V., Sultan, A., Violet, M., Chauderlier, A.,
Delattre, L., Loyens, A., Talahari, S., Begard, S., et al. (2016). Loss of Tau protein affects the
structure, transcription and repair of neuronal pericentromeric heterochromatin. Scientific
reports 6, 33047.

Masters, C.L., Multhaup, G., Simms, G., Pottgiesser, J., Martins, R.N., and Beyreuther, K.
(1985). Neuronal origin of a cerebral amyloid: neurofibrillary tangles of Alzheimer's disease
contain the same protein as the amyloid of plaque cores and blood vessels. The EMBO journal
4, 2757-2763.

Mathys, H., Davila-Velderrain, J., Peng, Z., Gao, F., Mohammadi, S., Young, J.Z., Menon, M.,
He, L., Abdurrob, F., Jiang, X., et al. (2019). Single-cell transcriptomic analysis of Alzheimer's
disease. Nature 570, 332-337.

McAlister, G.C., Nusinow, D.P., Jedrychowski, M.P., Wühr, M., Huttlin, E.L., Erickson, B.K.,
Rad, R., Haas, W., and Gygi, S.P. (2014). MultiNotch MS3 enables accurate, sensitive, and
multiplexed detection of differential expression across cancer cell line proteomes. Analytical
chemistry 86, 7150-7158.

Mertens, J., Herdy, J.R., Traxler, L., Schafer, S.T., Schlachetzki, J.C.M., Böhnke, L., Reid, D.A.,
Lee, H., Zangwill, D., Fernandes, D.P., et al. (2021). Age-dependent instability of mature

neuronal fate in induced neurons from Alzheimer's patients. Cell Stem Cell 28, 1533-
1548.e1536.
Mertens, J., Paquola, A.C., Ku, M., Hatch, E., Bohnke, L., Ladjevardi, S., McGrath, S.,
Campbell, B., Lee, H., Herdy, J.R., et al. (2015). Directly Reprogrammed Human Neurons
Retain Aging-Associated Transcriptomic Signatures and Reveal Age-Related
Nucleocytoplasmic Defects. Cell Stem Cell 17, 705-718.
Miller, J.D., Ganat, Y.M., Kishinevsky, S., Bowman, R.L., Liu, B., Tu, E.Y., Mandal, P.K., Vera,
E., Shim, J.W., Kriks, S., et al. (2013). Human iPSC-based modeling of late-onset disease via
progerin-induced aging. Cell Stem Cell 13, 691-705.
Mochizuki, A., Tamaoka, A., Shimohata, A., Komatsuzaki, Y., and Shoji, S. (2000). Abeta42-
positive non-pyramidal neurons around amyloid plaques in Alzheimer's disease. Lancet 355, 42-
43.
Moreau, D., Vacca, F., Vossio, S., Scott, C., Colaco, A., Paz Montoya, J., Ferguson, C.,
Damme, M., Moniatte, M., Parton, R.G., et al. (2019). Drug-induced increase in
lysobisphosphatidic acid reduces the cholesterol overload in Niemann-Pick type C cells and
mice. EMBO reports 20, e47055.
Mulay, S.R., and Anders, H.J. (2017). Crystal nephropathies: mechanisms of crystal-induced
kidney injury. Nat Rev Nephrol 13, 226-240.
Nativio, R., Donahue, G., Berson, A., Lan, Y., Amlie-Wolf, A., Tuzer, F., Toledo, J.B., Gosai,
S.J., Gregory, B.D., Torres, C., et al. (2018). Dysregulation of the epigenetic landscape of
normal aging in Alzheimer's disease. Nature neuroscience 21, 497-505.
Nguyen, A.T., Prado, M.A., Schmidt, P.J., Sendamarai, A.K., Wilson-Grady, J.T., Min, M.,
Campagna, D.R., Tian, G., Shi, Y., Dederer, V., et al. (2017). UBE2O remodels the proteome
during terminal erythroid differentiation. Science (New York, NY) 357.
Nijholt, D.A., de Graaf, T.R., van Haastert, E.S., Oliveira, A.O., Berkers, C.R., Zwart, R., Ovaa,
H., Baas, F., Hoozemans, J.J., and Scheper, W. (2011). Endoplasmic reticulum stress activates
autophagy but not the proteasome in neuronal cells: implications for Alzheimer's disease. Cell
death and differentiation 18, 1071-1081.
Nixon, R.A., and Yang, D.S. (2011). Autophagy failure in Alzheimer's disease--locating the
primary defect. Neurobiology of disease 43, 38-45.
Nixon, R.A., Yang, D.S., and Lee, J.H. (2008). Neurodegenerative lysosomal disorders: a
continuum from development to late age. Autophagy 4, 590-599.
Pang, Z.P., Yang, N., Vierbuchen, T., Ostermeier, A., Fuentes, D.R., Yang, T.Q., Citri, A.,
Sebastiano, V., Marro, S., Sudhof, T.C., et al. (2011). Induction of human neuronal cells by
defined transcription factors. Nature 476, 220-223.
Petkovic, M., O'Brien, C.E., and Jan, Y.N. (2021). Interorganelle communication, aging, and
neurodegeneration. Genes Dev 35, 449-469.
Piras, A., Collin, L., Grüninger, F., Graff, C., and Rönnbäck, A. (2016). Autophagic and
lysosomal defects in human tauopathies: analysis of post-mortem brain from patients with
familial Alzheimer disease, corticobasal degeneration and progressive supranuclear palsy. Acta
Neuropathol Commun 4, 22.
Pohl, C., and Dikic, I. (2019). Cellular quality control by the ubiquitin-proteasome system and
autophagy. Science (New York, NY) 366, 818-822.

Qiu, C., Kivipelto, M., and von Strauss, E. (2009). Epidemiology of Alzheimer's disease:
occurrence, determinants, and strategies toward intervention. Dialogues in clinical neuroscience
11, 111-128.
Radulovic, M., Schink, K.O., Wenzel, E.M., Nahse, V., Bongiovanni, A., Lafont, F., and
Stenmark, H. (2018). ESCRT-mediated lysosome repair precedes lysophagy and promotes cell
survival. The EMBO journal.
Raj, T., Li, Y.I., Wong, G., Humphrey, J., Wang, M., Ramdhani, S., Wang, Y.C., Ng, B., Gupta,
I., Haroutunian, V., et al. (2018). Integrative transcriptome analyses of the aging brain implicate
altered splicing in Alzheimer's disease susceptibility. Nature genetics 50, 1584-1592.
Rappsilber, J., Ishihama, Y., and Mann, M. (2003). Stop and go extraction tips for matrix-
assisted laser desorption/ionization, nanoelectrospray, and LC/MS sample pretreatment in
proteomics. Analytical chemistry 75, 663-670.
Rentzos, M., Zoga, M., Paraskevas, G.P., Kapaki, E., Rombos, A., Nikolaou, C., Tsoutsou, A.,
and Vassilopoulos, D. (2006). IL-15 is elevated in cerebrospinal fluid of patients with
Alzheimer's disease and frontotemporal dementia. Journal of geriatric psychiatry and neurology
19, 114-117.
Riku, Y., Duyckaerts, C., Boluda, S., Plu, I., Le Ber, I., Millecamps, S., Salachas, F., Brainbank
Neuro, C.E.B.N.N., Yoshida, M., Ando, T., et al. (2019). Increased prevalence of
granulovacuolar degeneration in C9orf72 mutation. Acta neuropathologica 138, 783-793.
Rockenstein, E., Mallory, M., Mante, M., Sisk, A., and Masliaha, E. (2001). Early formation of
mature amyloid-beta protein deposits in a mutant APP transgenic model depends on levels of
Abeta(1-42). Journal of neuroscience research 66, 573-582.
Salminen, A., Kaarniranta, K., Haapasalo, A., Hiltunen, M., Soininen, H., and Alafuzoff, I.
(2012). Emerging role of p62/sequestosome-1 in the pathogenesis of Alzheimer's disease.
Progress in neurobiology 96, 87-95.
Scheffer, L.L., Sreetama, S.C., Sharma, N., Medikayala, S., Brown, K.J., Defour, A., and
Jaiswal, J.K. (2014). Mechanism of Ca(2)(+)-triggered ESCRT assembly and regulation of cell
membrane repair. Nature communications 5, 5646.
Schweppe, D.K., Eng, J.K., Yu, Q., Bailey, D., Rad, R., Navarrete-Perea, J., Huttlin, E.L.,
Erickson, B.K., Paulo, J.A., and Gygi, S.P. (2020a). Full-Featured, Real-Time Database
Searching Platform Enables Fast and Accurate Multiplexed Quantitative Proteomics. Journal of
proteome research 19, 2026-2034.
Schweppe, D.K., Prasad, S., Belford, M.W., Navarrete-Perea, J., Bailey, D.J., Huguet, R.,
Jedrychowski, M.P., Rad, R., McAlister, G., Abbatiello, S.E., et al. (2019). Characterization and
Optimization of Multiplexed Quantitative Analyses Using High-Field Asymmetric-Waveform Ion
Mobility Mass Spectrometry. Analytical chemistry 91, 4010-4016.
Schweppe, D.K., Rusin, S.F., Gygi, S.P., and Paulo, J.A. (2020b). Optimized Workflow for
Multiplexed Phosphorylation Analysis of TMT-Labeled Peptides Using High-Field Asymmetric
Waveform Ion Mobility Spectrometry. Journal of proteome research 19, 554-560.
Settembre, C., Di Malta, C., Polito, V.A., Garcia Arencibia, M., Vetrini, F., Erdin, S., Erdin, S.U.,
Huynh, T., Medina, D., Colella, P., et al. (2011). TFEB links autophagy to lysosomal biogenesis.
Science (New York, NY) 332, 1429-1433.
Skowyra, M.L., Schlesinger, P.H., Naismith, T.V., and Hanson, P.I. (2018). Triggered
recruitment of ESCRT machinery promotes endolysosomal repair. Science (New York, NY) 360.

Studer, L., Vera, E., and Cornacchia, D. (2015). Programming and Reprogramming Cellular Age
in the Era of Induced Pluripotency. Cell Stem Cell 16, 591-600.
Su, J.H., Kesslak, JP., Head, E., and Cotman, C.W.(2002) Caspase-cleaved amyloid precursor
protein and activated caspase-3 are co-localized in the granules of granulovacuolar
degeneration in Alzheimer's disease and Down's syndrome brain. Acta neuropathologica104,1-
6.
Talboom, J.S., Velazquez, R., and Oddo, S. (2015). The mammalian target of rapamycin at the
crossroad between cognitive aging and Alzheimer's disease. NPJ Aging Mech Dis 1, 15008.
Tan, J.X., and Finkel, T. (2022). A phosphoinositide signalling pathway mediates rapid
lysosomal repair. Nature 609, 815-821.
Tanabe, K., Ang, C.E., Chanda, S., Olmos, V.H., Haag, D., Levinson, D.F., Sudhof, T.C., and
Wernig, M. (2018). Transdifferentiation of human adult peripheral blood T cells into neurons.
Proceedings of the National Academy of Sciences of the United States of America.
Tang, Y., Liu, M.L., Zang, T., and Zhang, C.L. (2017). Direct Reprogramming Rather than iPSC-
Based Reprogramming Maintains Aging Hallmarks in Human Motor Neurons. Frontiers in
molecular neuroscience 10, 359.
Tian, R., Abarientos, A., Hong, J., Hashemi, S.H., Yan, R., Dräger, N., Leng, K., Nalls, M.A.,
Singleton, A.B., Xu, K., et al. (2021). Genome-wide CRISPRi/a screens in human neurons link
lysosomal failure to ferroptosis. Nature neuroscience 24, 1020-1034.
Tong, B.C., Wu, A.J., Li, M., and Cheung, K.H. (2018). Calcium signaling in Alzheimer's disease
& therapies. Biochim Biophys Acta Mol Cell Res 1865, 1745-1760.
Traxler, L., Herdy, J.R., Stefanoni, D., Eichhorner, S., Pelucchi, S., Szücs, A., Santagostino, A.,
Kim, Y., Agarwal, R.K., Schlachetzki, J.C.M., et al. (2022). Warburg-like metabolic
transformation underlies neuronal degeneration in sporadic Alzheimer's disease. Cell
metabolism 34, 1248-1263.e1246.
Vest, R.T., Chou, C.C., Zhang, H., Haney, M.S., Li, L., Laqtom, N.N., Chang, B., Shuken, S.,
Nguyen, A., Yerra, L., et al. (2022). Small molecule C381 targets the lysosome to reduce
inflammation and ameliorate disease in models of neurodegeneration. Proceedings of the
National Academy of Sciences of the United States of America 119, e2121609119.
Victor, M.B., Richner, M., Olsen, H.E., Lee, S.W., Monteys, A.M., Ma, C., Huh, C.J., Zhang, B.,
Davidson, B.L., Yang, X.W., et al. (2018). Striatal neurons directly converted from Huntington's
disease patient fibroblasts recapitulate age-associated disease phenotypes. Nature
neuroscience 21, 341-352.
Vierbuchen, T., Ostermeier, A., Pang, Z.P., Kokubu, Y., Sudhof, T.C., and Wernig, M. (2010).
Direct conversion of fibroblasts to functional neurons by defined factors. Nature 463, 1035-1041.
Villeda, S.A., Plambeck, K.E., Middeldorp, J., Castellano, J.M., Mosher, K.I., Luo, J., Smith,
L.K., Bieri, G., Lin, K., Berdnik, D., et al. (2014). Young blood reverses age-related impairments
in cognitive function and synaptic plasticity in mice. Nature medicine 20, 659-663.
Vizcaíno, J.A., Deutsch, E.W., Wang, R., Csordas, A., Reisinger, F., Ríos, D., Dianes, J.A., Sun,
Z., Farrah, T., Bandeira, N., et al. (2014). ProteomeXchange provides globally coordinated
proteomics data submission and dissemination. Nature biotechnology 32, 223-226.
Wang, D., Zhang, J., Jiang, W., Cao, Z., Zhao, F., Cai, T., Aschner, M., and Luo, W. (2017). The
role of NLRP3-CASP1 in inflammasome-mediated neuroinflammation and autophagy

dysfunction in manganese-induced, hippocampal-dependent impairment of learning and
memory ability. Autophagy 13, 914-927.
Wang, Z., Wang, Q., Li, S., Li, X.J., Yang, W., and He, D. (2022). Microglial autophagy in
Alzheimer's disease and Parkinson's disease. Front Aging Neurosci 14, 1065183.
Whyte, L.S., Lau, A.A., Hemsley, K.M., Hopwood, J.J., and Sargeant, T.J. (2017). Endo-
lysosomal and autophagic dysfunction: a driving factor in Alzheimer's disease? Journal of
neurochemistry 140, 703-717.
Wiersma, V.I., van Ziel, A.M., Vazquez-Sanchez, S., Nolle, A., Berenjeno-Correa, E., Bonaterra-
Pastra, A., Clavaguera, F., Tolnay, M., Musters, R.J.P., van Weering, J.R.T., et al. (2019).
Granulovacuolar degeneration bodies are neuron-selective lysosomal structures induced by
intracellular tau pathology. Acta neuropathologica 138, 943-970.
Willén, K., Edgar, J.R., Hasegawa, T., Tanaka, N., Futter, C.E., and Gouras, G.K. (2017). Aβ
accumulation causes MVB enlargement and is modelled by dominant negative VPS4A.
Molecular neurodegeneration 12, 61.
Wolfe, D.M., Lee, J.H., Kumar, A., Lee, S., Orenstein, S.J., and Nixon, R.A. (2013). Autophagy
failure in Alzheimer's disease and the role of defective lysosomal acidification. Eur J Neurosci
37, 1949-1961.
Yamoah, A., Tripathi, P., Sechi, A., Kohler, C., Guo, H., Chandrasekar, A., Nolte, K.W., Wruck,
C.J., Katona, I., Anink, J., et al. (2020). Aggregates of RNA Binding Proteins and ER
Chaperones Linked to Exosomes in Granulovacuolar Degeneration of the Alzheimer's Disease
Brain. Journal of Alzheimer's disease : JAD 75, 139-156.
Yoon, L., Botham, R.C., Verhelle, A., Sanz-Martinez, P., Xu, J., Cole, C.M., Tan, E.P., Chou, C.-
C., Cuoco, C.A., Massey, L.A., et al. (2022). mTOR inhibitor-independent Autophagy Activator
Ameliorates Tauopathy and Prionopathy Neurodegeneration Phenotypes. bioRxiv,
2022.2009.2029.509997.
Zhang, M.J., Pisco, A.O., Darmanis, S., and Zou, J. (2021). Mouse aging cell atlas analysis
reveals global and cell type-specific aging signatures. Elife 10.
Zhang, W., Xu, C., Sun, J., Shen, H.M., Wang, J., and Yang, C. (2022). Impairment of the
autophagy-lysosomal pathway in Alzheimer's diseases: Pathogenic mechanisms and
therapeutic potential. Acta Pharm Sin B 12, 1019-1040.
Zhao, J., Fu, Y., Yamazaki, Y., Ren, Y., Davis, M.D., Liu, C.C., Lu, W., Wang, X., Chen, K.,
Cherukuri, Y., et al. (2020). APOE4 exacerbates synapse loss and neurodegeneration in
Alzheimer's disease patient iPSC-derived cerebral organoids. Nature communications 11, 5540.
1886
1887