Complex Formation of Immunoglobulin Superfamily Molecules Side-IV and Beat-

IIb Regulates Synaptic Specificity in the *Drosophila* Visual System

Jiro Osaka[1], Arisa Ishii[1], Xu Wang[1], Satoko Hakeda-Suzuki[1,2], Riku Iwanaga[1], Hinata

Kawamura[1], Takashi Suzuki[1*]

1. School of Life Science and Technology, Tokyo Institute of Technology, Yokohama,

226-8501, Japan

2. Research Initiatives and Promotion Organization, Yokohama National University,

Yokohama, 240-8501, Japan

*: Corresponding author.

suzukit@bio.titech.ac.jp (T.S.)

SUMMARY

Neurons express many cell surface proteins as mutually binding "key-lock molecules"

that can create synapses. However, the molecular mechanisms of how neurons make

synapses only with preferred targets are not completely understood. Here we identified

Side-IV and Beat-IIb, belonging to the *Drosophila* immunoglobulin superfamily, as a

new key-lock combination capable of inducing synapse formation. Side-IV interaction

with Beat-IIb transduces bifurcated signaling to Side-IV's co-receptor, Kirre, and a

synaptic scaffold protein, Dsyd-1. Localization and genetic interaction analyses revealed

that Side-IV localizes subcellularly at synapse formations defined by Beat-IIa/b and

anchors Dsyd-1. Our data demonstrate that a complex made up of Side-IV, Beat-IIb,

Kirre, and Dsyd-1 not only narrows neuronal binding specificity but also recruits synapse

formation factors Kirre and Dsyd-1 to restrict synapse formation loci and inhibit

miswiring. We propose a mechanism by which key-lock molecules set a hierarchy of

preference among neuronal pairs in a complex circuit in vivo.

Keywords: synaptic specificity, synapse formation, key-lock molecules, transmembrane

clustering, subcellular localization, Beat and Side proteins, irre cell recognition module,

Dsyd-1-Liprin-α

INTRODUCTION

All neurons can potentially form synapses with many other neurons, but still demonstrate

specificity in synapse partner selection.[1-4] During development, neurons stochastically

extend filopodia, searching for synaptic partners.[5-7] Synaptogenesis begins only when

filopodia find a synaptic partner through specific molecular interactions.[2,5] By the study

of synapse formation in cultured cells, several cell surface proteins were found to have

"key-lock molecules" that can induce synapse formation.[8] In vivo studies have shown

that neurons have the capability to form a constant number of synapses indiscriminately;

however, they rank target synapses according to a preferred order.[3,4] The molecular

mechanisms that drive synaptic preferences and specificity have not been fully explored.

54   The *Drosophila* visual system consists of a compound eye and four optic

ganglia, compactly composed of about 200 diverse neurons with well-documented

connections.[9,10] In addition, single-cell analyses in the last few years have revealed cell-

type and time-specific gene expression profiles of the *Drosophila* visual system, which

have been compiled into a database such as Gene Expression Omnibus.[11-15] This makes

the *Drosophila* visual system an ideal model for connectome and transcriptome-based

analyses. Several transmembrane protein pairs with binding interactions have been

reported to be expressed between synaptic partners.[13,16-18] The most studied is the DIP-

Dpr families of the immunoglobulin superfamily (IgSF), which forms heterophilic or

homophilic interaction networks.[16,19] In recent years, it has been shown that DIP-Dpr

protein interactions are involved in neural connections, axon guidance, and cell death

signaling.[3,17,18,20-24] Another member of IgSF, the Beat-Side families, have similar

properties to DIP-Dpr, although its function is less well understood. Therefore, we

considered the Beat-Side families as a candidate for the molecules involved in neural

circuit formation.

69   Both *beaten path Ia* (*beat-Ia*) and *sidestep* (*side*) were identified by the genetic

screening of mutants showing abnormal axon guidance in neuromuscular junctions

(NMJ).[25-27] The *beat-Ia* and *side* mutants had similar phenotypes and were shown to

interact genetically and biochemically.[16,28,29] Genomic and interactome analyses revealed

that *beat-Ia* and *side* have 14 and 8 paralogs, respectively, with a heterophilic interaction

network.[16,29-32] Furthermore, single-cell RNAseq data showed that Beat-Side families are

expressed in a cell-specific manner throughout the nervous system.[11,13] Although several

works have reported the function of Beat-Ia/Side interaction in NMJ, motor function, and

leg nerve development,[28,33-35] the functions of other Beat-Side families are still elusive,

especially in the central nervous system.

In the present study, we report a new key-lock combination in synaptic

specificity, Side-IV and Beat-IIb. Overexpression of *side-IV* can induce synapses in the

R7 photoreceptor in *Drosophila*. Triggered by *trans*-interaction with Side-IV, Beat-IIb

made a cluster complex that can transduce bifurcated signaling to induce synapses with

its co-receptor, Kirre, and a synaptic scaffold protein, Dsyd-1. Surprisingly, in *side-IV*

loss of function analysis, against our intuitive guess, we saw more ectopic synapse

formation in the distal part of lamina monopolar neurons. Combining loss-of-function

analyses and subcellular protein localization analyses, we demonstrated that the newly

characterized Side-IV/Beat-IIb/Kirre/Dsyd-1 complex has two functions. One is to

narrow neuronal binding specificity via Side-IV/Beat-IIb extracellular interactions. The

second is to recruit synapse formation factors Kirre and Dsyd-1 to restrict synapse

formation loci and inhibit miswiring. These two functions can explain how redundant

key-lock molecules can rank preferred interactions among neuronal pairs and accomplish

synaptic specificity in a complex circuit.

RESULTS

Side-IV and Beat-IIb are a New Key-Lock Combination for Synaptic Induction

To identify a new key-lock molecule, we focused on the function of the Beat-Side

families. Photoreceptor R7 projects its axons to the medulla and mainly synapses on the

dendrite of Dm8 in the M4–M6 layer.[10,36] The presynaptic structure of R7 is limited to

the M4–M6 layer.[6,37] By utilizing synaptic tagging with recombination (STaR),[37] which

can label the active zone protein Bruchpilot (Brp) in a cell-specific manner, we

specifically visualized R7's presynaptic structures with R7-specific FLPase 20C11-FLP

(Figure 1A).

Beat-Side family members were either knocked down (Figures S1A and B) or

overexpressed (Figures S1C and D) in R7. We found that ectopic synapses were induced

in the M1–M3 layer of R7 by overexpression of *side-IV* (Figures 1B and B', yellow allows,

and S1C). In addition, some axons overshot toward the M10 layer (Figures 1B and B'',

yellow arrowheads). These phenotypes suggested that R7 was miswired around M1–M3

and M10 layers other than M4–M6 due to expression of *side-IV*. Endogenous tags of

Side-IV revealed R7 doesn't endogenously expresses *side-IV* (Figures S1E–I'). Since

Side-IV biochemically interacts with Beat-IIa and Beat-IIb,[16,29] we tested if Side-IV-

Beat-IIa/b interactions were responsible for the miswiring phenotypes. Both synaptic

induction and overshoot phenotypes generated by *side-IV* overexpression were

significantly suppressed in a *beat-IIa* or *beat-IIb* mutant background (Figures 1C–G). An

endogenous Beat-IIb marker[38] showed layer-specific expression in the M1–M2 and M9–

M10 layers where miswiring phenotypes occurred (Figures 1H–I). Furthermore, we

found that the Beat-IIb signals became punctate when *side-IV* was expressed (Figures 1J–

K''), and presynaptic Brp puncta were induced adjacently (Figures 1K' and K'', yellow

arrows). The punctate signal suggested clustering transmembrane proteins, a common

phenomenon seen with other synapse-organizing molecules.[39] The concurrent

visualization of Side-IV, Beat-IIb, and Brp also revealed clustered signals of Side-IV,

about 70% of which co-localized with Beat-IIb clusters (Figures 1L–L"). Furthermore,

about 70% of Side-IV/Beat-IIb clustered signals co-localized with ectopic synapses

(Figures 1L−L", yellow arrows). These data indicated that Side-IV induced Beat-IIb

clustering, and that this cluster complex induced synapse (Figure 1M). We therefore have

identified Side-IV/Beat-IIb as a new key-lock combination capable of inducing synapses

by establishing cluster complexes.

Side-IV Transduces Signals Both by its Co-Receptors and the Cytoplasmic Domain

to Induce Synapses

To uncover the mechanism of synaptic induction by Side-IV, we investigated the impact

of each domain. Side-IV has five extracellular Ig domains (Ig1–5), a fibronectin type III

domain (FnIII), and a cytoplasmic domain (Cyto) with a PDZ-binding motif (PDZb)

(Figure 2A). We constructed domain-deficient variants (Figure S2A) and chimeras of

Side-IV with a non-functional membrane protein, mCD8 (Figures 2A and B). With these

constructs, we analyzed the Side-IV/Beat-IIb clustering and ectopic synapse/overshoot

induction phenotypes in R7 (Figures 2 and S2B–K). We found that Side-IV could not

induce synapse without an extracellular Ig domain (Figures S2B–E, and I). In *Drosophila*

S2 cells, cell aggregations were formed via *trans*-interactions when *side-IV* expressing

cells were mixed with *beat-IIb* expressing cells (Figures S2L–L", yellow dotted lines).

The cell aggregations were not formed by Side-IV $^{\Delta Ig1}$ or Side-IV $^{\Delta Ig1-5}$ (Figures S2M–

O"), indicating that the Ig domain is necessary for *trans*-interaction. The chimeric protein

with the extracellular domain of Side-IV fused with the transmembrane and cytoplasmic

domains of mCD8 (Side-IV^{Ecto}::mCD8$^{TM+Cyto}$) showed a similar degree of synapse

induction as Side-IV $^{\Delta Cyto}$ (Figures 2E–F''', and H–K), suggesting that the transmembrane

domain of Side-IV is not important for synaptic induction. On the other hand, the chimeric

protein with the extracellular domain of Side-IV fused with mCD8 and the FnIII domain

deleted (Side-IV $^{Ecto\Delta FnIII}$::mCD8) showed no cluster signal in the M1–M3 layer of R7 and

failed to induce synapses (Figures 2G–K). This indicates that the FnIII domain is essential

for Side-IV clustering. *Side-IV$^{Ecto\Delta FnIII}$::mCD8* expressing cells demonstrate strong cell

aggregation with *beat-IIb* expressing cells, to a similar degree as *side-IV ^{FL}* (Figures S2P–

Q). This indicates that the chimera has a strong affinity for Beat-IIb but has lost clustering

ability. These data suggest that the Ig domains are important for *trans*-interaction with

Beat-IIb, and the FnIII domains are important for Side-IV clustering; both are required to

establish the Side-IV/Beat-IIb clustering complex, which is crucial for synaptic induction.

Introducing Nrxs-Nlgs interaction induced synapse formation in M1–M3, but

the homophilic interactions of epithelial cadherins or human-derived sidekick cell

adhesion molecule 2 (SDK2) did not (Figure S3). This indicates that increased affinity

between R7 and postsynaptic neurons[40] is not sufficient for synaptic induction. Therefore,

we hypothesized that the Side-IV/Beat-IIb cluster complex could transduce signals. Side-

IV could induce synapses without the cytoplasmic domain (Figures 2E–E''', and J). Side-

IV may transduce signal via its co-receptors. The synaptic induction phenotype of Side-

IV$^{\Delta Cyto}$ or Side-IV$^{\Delta PDZb}$ was much weaker than that of full-length Side-IV (Figures 2D–

E''', and J; Figures S2C, G–I), indicating that signaling through the PDZ-binding motif

also contributed to synaptic induction. Side-IV may have two distinct signaling pathways

transduced by its co-receptors and the cytoplasmic domain. Taken together, these data

show that the extracellular Ig and FnIII domains of Side-IV are essential for establishing

the Side-IV/Beat-IIb cluster complex, and the cytoplasmic domain of Side-IV and/or its

co-receptors may transduce signals for synapse organization.

Kirre Can Form a Complex with Side-IV with the Extracellular Domain as a Co-

Receptor

To identify the co-receptor of Side-IV, we searched for transmembrane proteins that can

enhance Side-IV signaling. The immunoglobulin superfamily protein Kirre[41] can enhance

Side-IV's phenotypes (Figures 3A–F). Kirre expression doubled the Side-IV overshoot

phenotype, while Kirre itself could not induce an overshoot at all, indicating that the

enhancement is not additive, but synergistic (Figure 3F). By analyzing the overshoot

phenotype, we tested the impact of the cytoplasmic domains on the synergistic effects

(Figures 3G–P). Side-IV and Kirre had little synergy when each lacked a cytoplasmic

domain (Figures 3O and P). On the other hand, they were strongly synergistic when either

cytoplasmic domain remained (Figures 3L, N, and P). This suggests that Kirre can

enhance Side-IV signaling, through the cytoplasmic domains, as a co-receptor.

If Kirre works as a co-receptor of Side-IV, the two proteins would likely form

a Side-IV/Kirre complex. As predicted, Side-IV co-immunoprecipitated with Kirre, and

vice versa (Figure S4A). Since Side-IV and Kirre did not form cell aggregation via *trans*-

interactions (Figures S4B–D), they may form a complex by their *cis*-interaction. In

Drosophila S2 cells, Flamingo polarity protein accumulates at cell contact surfaces and

can recruit transmembrane proteins Vang, Frizzled, and Gogo by *cis*-interactions.[42,43]

Using S2 cells, we assessed the *cis*-interaction between Side-IV and Kirre. Homophilic

self-interaction of Kirre causes Kirre-transfected S2 cells to form aggregates, often

accumulating at sites of cell-cell contact (Figures 3Q and Q'). We noted Side-IV co-

accumulation with Kirre (Figures 3Q" and R). Since Side-IV co-accumulation did not

occur when *cadE*-expressing cells aggregated via their homophilic interactions (Figures

3S–T), the effect was Kirre-specific. These data suggest that Side-IV and Kirre form a

complex at the cell membrane through their *cis*-interaction. Furthermore, the

accumulation requires Side-IV's extracellular domain but not its cytoplasmic domain

(Figures 3U and V). This result was consistent with the domain requirement for synergy

between Side-IV and Kirre in the overshoot phenotype (Figure S4 E–M). These data

suggest that Kirre may work as a co-receptor of Side-IV, and that they form a Side-

IV/Kirre complex established by *cis* interaction through their extracellular domains.

The Cytoplasmic Domain of Side-IV May Establish a Dsyd-1-Liprin-α Complex that

Functions Downstream

The synaptic induction of Side-IV$^{\Delta Cyto}$ suggests that signaling through the cytoplasmic

domain is also functionally required for signaling. To identify the downstream signaling

targets of Side-IV, we searched for synapse related molecules necessary for Side-IV

synaptic induction. By the RNAi approach, we found that knocking down *dsyd-1* or

liprin-α could significantly suppress synaptic induction by *side-IV* overexpression in R7

(Figures 4A–F), suggesting that Dsyd-1 and Liprin-α work downstream of Side-IV.

Using S2 cells, we analyzed the co-localization of Side-IV with Dsyd-1 and

Liprin-α (Figures 4G–L). Side-IV showed substantial co-localization with Dsyd-1

(Figures 4G and I). Since Dsyd-1 has a PDZ domain that interacts with synapse-

organizing molecules such as Nrx-1,[44] we hypothesized that Side-IV interacts with Dsyd-

1 through Side-IV's PDZ-binding motif. The co-localization index between Side-IV and

Dsyd-1 decreased by almost half in the absence of Side-IV's PDZ-binding motif (Figures

4H–I), indicating its necessity for the Side-IV/Dsyd-1 interaction. Although Side-IV and

Liprin-α did not show strong co-localization, their co-localization index almost doubled

when *dsyd-1* was also expressed in the S2 cells (Figures 4J–L). This indicates that Side-

IV can form a complex with Liprin-α through Dsyd-1. These data suggest that the PDZ-

binding motif of Side-IV interacts with Dsyd-1, and establish Dsyd-1/Liprin-α as a

complex important for active zone formation.[44,45]

To validate the idea that Dsyd-1 and Liprin-α work downstream of the

cytoplasmic domain of Side-IV, we utilized Side-IV$^{\Delta Cyto}$, which cannot transduce

cytoplasmic signaling. Knockdown of *dsyd-1* suppressed synaptic induction by Side-IV^{FL},

but not Side-IV$^{\Delta Cyto}$ (Figure 4M), indicating that Dsyd-1 acts exclusively on signaling

through the Side-IV's cytoplasmic domain. On the other hand, knockdown of *liprin-α*

suppressed synaptic induction by both Side-IV^{FL} and Side-IV$^{\Delta Cyto}$ (Figure 4M). This

indicates that Liprin-α affects signaling both through the Side-IV's cytoplasmic domain

and the co-receptor(s). The knockdown of *kirre* suppressed synaptic induction by Side-

IV$^{\Delta Cyto}$, but not Side-IV^{FL} (Figure 4M), indicating that Kirre affects signaling independent

of the cytoplasmic domain. These data indicate that Side-IV forms a signaling complex

with Kirre as a co-receptor, and additionally with Dsyd-1/Liprin-α downstream of the

cytoplasmic domain (Figure 4N).

***side-IV* and *side-VII* are Required for Synapse Formation of L2 Neurons Restricted**

in the Proximal Region

By toggled expression experiments of *side-IV* in R7, we found that Side-IV/Beat-IIb is a

new key-lock combination for synaptic induction at the presynapse of R7. To analyze

their endogenous function, we searched for synaptic partners where Side-IV/Beat-IIb

interaction functions by utilizing single-cell RNAseq data in the *Drosophila* visual

system.[11,17] The Side-IV/Beat-IIa/b and Side-VII/Beat-IV interactions were predicted

between L2 and L4 neurons (Figures 5A and B).[11,17] Furthermore, endogenous tags of

Side-IV showed signals in the M2 layer (Figures S1F and F'), indicating that L2 expressed

side-IV. Although there are five subtypes of lamina neurons, only L2 and L4 make

reciprocal synapses restricted to the proximal region of the lamina (Figures 5A, C, and

C').[46] We generated *side-IV* or *side-VII* mutants (see Methods) and analyzed the function

of Beat-Side interactions for the synapse formation (Figures 5C–H).

247   Despite Side-IV's strong synapse-organizing ability in R7, the *side-IV* mutant

did not show any decrease in synapse number in the proximal region (Figures 5C–D', and

G). Potentially, the loss of synapse-organizing molecules did not result in synaptic loss

because of molecular redundancy[47,48]; disrupting a few genes may not be sufficient for

decreasing synapse numbers. At least ten transmembrane interactions were predicted

between L2 and L4,[13] indicating that these synaptic connections may have high molecular

redundancy. However, the *side-IV* mutant showed ectopic synapse formation in the distal

region (Figures 5C–D', and H), similar to those reported in the *dip-β* mutant.[3] We

conducted rescue experiments of *side-IV* and found the Side-IV [FL] could rescue the

phenotype completely while Side-IV [ΔCyto] could partially (Figures. S5A–D), indicating

that regaining the physical strength of cell-cell contact is not sufficient, but cytoplasmic

signaling is also important. The *side-VII* mutants also showed ectopic synapse formation

in the distal region, which was significantly enhanced in the *side-IV* / *side-VII* double

mutants (Figures 5E–H). Knockdown of *side-IV* or *side-VII,* specifically in L2, resulted

in ectopic synapse formation in the distal region, indicating that *side-IV* and *side-VII* are

required in L2 (Figures S5E–H). Furthermore, downregulation of *beat-IIa/b* or *beat-IV*

phenocopied the ectopic synapse formation in the distal region (Figures S5I–P). To assess

which lamina neuron actually induced the ectopic synapse formation by diminishing

Beat-Side interactions, we analyzed the synapse of each subtype of lamina neuron in *side-*

IV / side-VII double mutant flies. L2 showed ectopic synapse formation in the distal

region (Figures 5I–L), whereas other subtypes did not (Figure S6). These data indicate

that Side-IV/Beat-IIa/b and Side-VII/Beat-IV interactions are necessary for proper

synapse formation, restricted to the proximal region, of L2. Side-IV and Side-VII may

regulate the synaptic specificity of L2 by indicating the location of synapse formation.

Side-IV and Side-VII Demonstrate Beats-Dependent Subcellular Localization to the

Proximal Region

If Side-IV and Side-VII determine the location of synapse formation of L2, both should

localize to the proximal region where presynaptic structures are formed. We analyzed the

endogenous localization of Side-IV and Side-VII with FLPase-based cell-specific tags

(Figure S1E). The co-localization of Side-IV, Side-VII, and Brp at 72 hours after

puparium formation (APF) revealed that Side-IV and Side-VII localization was restricted

to the proximal region (Figures 6A–A'''). Both *side-IV* and *side-VII* were expressed in L2

neurons, from developmental to adult stages (Figure S7). At the 72 hour APF and adult

stages, endogenous Side-IV and Side-VII showed subcellular localization restricted to the

proximal region (Figures S7C–D' and G–H'). This characteristic subcellular localization

was also observed when the GFP-tagged *side-IV* or *side-VII* was expressed in L2 neurons

(Figures 6B–C'). Since the dendritic arbors of the L2 neuron are uniform from the distal

to proximal regions,[49,50] we speculated that the subcellular localization of Side-IV and

Side-VII is regulated in a non-cell-autonomous manner. We hypothesized that Beats

expressed in L4 might define the subcellular localization of Sides.

We analyzed the localization dependency of Side-IV and Side-VII on Beat-

IIa/b and Beat-IV, respectively (Figures 6D–M). In the wild type, Side-IV localized from

the proximal to the distal region at 48 hours APF (Figures 6D and D'). Subcellular

localization to the proximal region was observed at 72 hours APF (Figures 6F and F').

With downregulated *beat-IIa*/*b*, Side-IV localized similarly to the wild type condition at

48 hours APF (Figures 6E and E'). However, it failed to localize to the proximal region

at 72 hours APF (Figures 6G–H). The localization of Side-VII in the wild type was similar

to that of Side-IV, appearing throughout the lamina at 48 hours APF (Figures 6I and I')

and only the proximal region at 72 hours APF (Figures 6L and L'). Similar to the impact

of *beat-IIa/b* downregulation on Side-IV localization, the downregulation of *beat-IV* did

not affect Side-VII localization at 48 hours APF (Figures 6J and J') but disrupted its

localization at 72 hours APF (Figures 6L–M). These data suggest that the subcellular

localization of Side-IV and Side-VII are Beat-IIa/b and Beat-IV-dependent, respectively.

If Beat-IIa/b and Beat-IV defined the localization of Side-IV and Side-VII, they

should also localize to the proximal area. We labeled the L4 morphology and analyzed

the localization of Beat-IIb and Beat-IV with MiMIC-GFP transgenics.[38] We mainly

focused on Beat-IV here, since Beat-IIb-MiMIC-GFP showed no localization on axons

(Figures S8A–A", yellow arrowheads), but only in cell bodies (Figures S8A–A", yellow

arrows), perhaps due to some defects in axonal transport. Beat-IV-MiMIC-GFP signal

was seen only in the proximal area of L4 already at 72 hours APF (Figures S8B–B"). We

then analyzed the co-localization of Side-VII and Beat-IV, and found Beat-IV localized

to the proximal region before Side-VII (Figures S8C–D"). Futhermore, the localization

of Beat-IV is not dependent on Side-VII (Figures S8E–H'). These data suggest that Beat-

IV localizes to the proximal region first, and then dictates Side-VII localization. In

summary, Side-IV and Side-VII showed subcellular localization to the proximal region

in a Beat-dependent manner.

Complex Formation of Side-IV with Kirre and Dysd-1 Restricts Loci of Synapse

Formation and Inhibits Miswiring

The functional and localization analyses of Side-IV and Side-VII suggest that they

instruct L2 to form synapses in the proximal region. The remaining question was why

side-IV or *side-VII* mutants showed ectopic synapse formation in the distal region. A

previous study reported that DIP-β causes L4 to prefer L2 as its synaptic partner over

other neurons. The *dip-β* mutant resulted in miswiring to alternative neurons with ectopic

synapse formation in the distal region.[3] Similarly, Side-IV and Side-VII may help L2

choose L4 as its synaptic partner and inhibit its miswiring to alternative neurons. In this

case, there may be molecules that instruct miswiring of L2 to alternative neurons when

side-IV or *side-VII* are absent.

Our analysis of *side-IV* in R7 suggested that Kirre can work as a co-receptor of

Side-IV (Figure 3). Previous research reported that the *kirre* mutant resulted in total loss

of postsynaptic structures of L4,[51] indicating that Kirre is a major presynaptic regulator

in L2. Furthermore, single-cell-RNAseq data predicted unbiased interactions, homophilic

self-interactions of Kirre and heterophilic Kirre-Rst interactions, between L2 and all other

subtypes of lamina neurons.[13,16] We hypothesized that Kirre cooperates with Side-IV to

establish L2-L4 connectivity but induces synaptic connection indiscriminately without

Side-IV. In L2, Kirre showed the strongest localization in the proximal region, similar to

Side-IV (Figures 7A–B). However, Kirre localized more broadly than Side-IV, and some punctate signals were observed in the distal region (Figures 7A''', yellow arrowheads, B, and C). This indicates that Kirre localizes and interacts with Kirre/Rst not only in the proximal but also in the distal region. To assess whether Kirre is responsible for the ectopic synapse formation in the *side-IV* mutants, we tested genetic interactions between *kirre* mutant[52] and *side-IV* or *side-IV / side-VII* double mutants (Figures 7D–K). Deleting one copy of *kirre* decreased the number of synapses in the proximal region (Figures 7D–G and J–K), as well as suppressed ectopic synapse formation in the distal region (Figures 7E–I and K). These data indicate that Kirre promotes L2 miswiring with alternative neurons in the distal region in the *side-IV* or *side-IV / side-VII* double mutant.

The mechanism by which Side-IV can inhibit ectopic synapse formation in the distal region of the wild type is still elusive. One possibility is that some competitive mechanisms inhibit ectopic synapse formation. Since Side-IV and Side-VII localize specifically to the proximal region (Figure 6), they may compete with the molecules working in the distal region. A competitive synapse formation mechanism has been reported in R7.[5] While neurons extend several filopodia to search for synaptic partners, only one or two filopodia that recruit synapse seeding factors are stabilized at each time point.[5] Synaptic loci compete for the factors, since the amounts are limited and Dsyd-1 and Liprin-α are potential candidates for synapse seeding factors.[5] To test whether Dsyd-1 plays a role as a competing factor here, we knocked in a FLPase-based cell-specific GFP-tag to the Dsyd-1 locus and analyzed its endogenous localization (Figures 7L–T; see Methods). Dsyd-1 localized to the proximal region at 48 hours APF (Figures 7L–L"), and signals became punctate at 72 hours APF (Figures 7P and P'). *side-IV* or *side-IV / side-VII* double mutant flies showed leaky Dsyd-1 signals in the distal region at 48 hours APF

(Figures 7M–N''', yellow arrows, and O). Overall, the number of Dsyd-1 puncta decreased

in the proximal region and increased in the distal region (Figures 7Q–T). These data

indicate that Side-IV can anchor Dsyd-1 localization to the proximal region, not allowing

the distally distributed molecules a chance to form synapses. Taken together, our results

indicate that Side-IV may establish L2-L4 synapse specificity by cooperating with Kirre

and Dsyd-1 in the proximal region, which can restrict synapse formation loci and inhibit

miswiring.

DISCUSSION

This study identified Side-IV/Beat-IIb as a new key-lock combination that can induce

synapse formation by establishing a cluster complex and signaling through its co-receptor

Kirre or cytoplasmic signaling via Dsyd-1/Liprin-α in the *Drosophila* R7 photoreceptor.

Side-IV regulates the loci of L2 synapse formation by anchoring Dsyd-1 to inhibit

miswiring, with subcellular localization specified by Beat-IIa/b. We therefore propose a

new key-lock complex formation model between the Beat and Side families for

establishing synaptic specificity (Figure. 8).

Side-IV Establishes a Signaling Complex with Kirre and Dsyd-1

An essential observation is that Side-IV has the potential to induce synapse formation in

the absence of its cytoplasmic domain (Figures 2E–E''', and J). Based on this observation,

we hypothesized that Side-IV could induce signaling through co-receptors. We

demonstrated that Side-IV and its co-receptor Kirre synergistically mediate signal

transduction through the extracellular domain. Kirre is a well-known cell adhesion

molecule and a member of the irre cell recognition module (IRM) family.[41] Its functions

range from nervous system-related processes such as axon guidance and synapse formation to myoblast aggregation and fusion, cell sorting, and other diverse biological phenomena.[41,51] *Cis*-regulation of Side-IV by such well-known cell adhesion molecules was surprising. Since co-expressing *side-IV* with other IRM members, *hbs* and *sns,* did not enhance the phenotype (data not shown), Kirre may function as a unique co-receptor for Side-IV. It is noted that there were some redundancies in the signaling of Side-IV/Kirre. The co-overexpression experiment in R7 showed that if either Side-IV or Kirre signaling pathway is active, meaning that the cytoplasmic domain of Side-IV or Kirre is present, the complex can induce a stronger phenotype (Figures 3G–P). This study indicates that Beat-Side family proteins may have unique co-receptors with cytoplasmic redundancy, and that their interplay could generate diverse connections.

Our study suggests that Dsyd-1 and Liprin-α work downstream of Side-IV. Side-IV strongly co-localized with Dsyd-1, but this co-localization significantly decreased when the PDZ-binding motif of Side-IV was deleted (Figures 4G–I). Furthermore, knockdown of *dsyd-1* specifically suppressed synaptic induction by cytoplasmic signaling of Side-IV (Figure 4M). Liprin-α co-localized with Side-IV through Dsyd-1, and knockdown of *liprin-a* strongly suppressed synaptic induction by Side-IV (Figures 4J–M). These data suggest that Dsyd-1 interacts with Side-IV's PDZ-binding motif and works with Liprin-α. In *Drosophila,* other synapse organization molecules Nrx-1 and Lar interact with Dysd-1 and Liprin-α, respectively.[44,53] Side-IV may compete with these molecules to recruit Dsyd-1 or Liprin-α.

Subcellular Localization of Side-IV and Side-VII Establishes L2-L4 Connectivity

Although Side-IV exhibits strong synapse induction abilities in R7, the *side-IV* mutant

did not demonstrate any decrease in the number of synapses in lamina neurons (Figures

5C–D', and G). Such observation is quite common with other synapse organizers, such as

Neurexins and Neuroligins; even knockdown or mutation of their three paralogues only

slightly impacts synapse number despite their strong synapse-organizing abilities.[47,48]

Conversely, notable ectopic synapse formation of *side-IV* in the distal region was

observed (Figures 5C–D', and H).

Based on the expression data, three Beat-Side interactions are predicted

between L2 and L4 (Figures 5A and B),[11,17] and the number of induced ectopic synapses

in the distal region increased with disruption of these interactions. We confirmed that

ectopic synapses were formed in L2 but not in other subtypes of lamina neurons (Figures

5I–L and S6). These findings indicate that several members of the Beat and Side families

function redundantly to determine the loci of synapse formation in L2, specifically in the

proximal region. This is supported by the localization analysis in which both Side-IV and

Side-VII showed distinct subcellular localization to the proximal region in L2 which is

defined by Beat-IIa/b and Beat-IV, respectively (Figures 6B–M). Despite uniform

dendritic arborization of L2 from the proximal to distal areas,[49,50] the interplay between

Beat-Side interactions and the non-uniform cellular morphology of L4 (dendrite

distribution to the proximal region)[49] appears to establish non-uniform synapse formation

restricted to the proximal region of L2. The DIP-β protein, which works at loci of L4

synapse formation, also showed substantial accumulation in the proximal region at 72

hours APF,[3] resembling the localization of Side-IV and Side-VII. The collaboration

between DIP-Dpr and Beat-Side families is crucial in establishing L2-L4 connectivity.

Complex Formation of Side-IV Links Synaptic Specificity and Synaptogenesis

Synaptogenesis has a two-faced nature that is robust in establishing neural circuits, but can also be indiscriminate.[3,4,6,7] It is plausible that molecular coding, such as recognizing key-lock molecules, determines synaptic partners.[2,8] However, the loss of key-lock molecules sometimes results in miswiring with alternative partners.[3,4] Neurons can synapse with various partners, and key-lock molecules can bias synaptic partner preference. We initially sought to understand how preference for a synaptic partner is determined. Increasing the affinity between R7 and postsynaptic partners by homophilic interactions of CadE or hSDK2 alone did not induce synapse formation (Figure S3). Moreover, Side-IV $^{\Delta Cyto}$ couldn't completely rescue the ectopic synapse formation in *side-IV* (Figures S5A–D). These data indicate that some intracellular signaling mechanisms are required to increase preference and generate synaptic connection.[54]

Previous studies have demonstrated that DIP-β facilitates the preferential formation of synapses between L4 and L2 neurons specifically.[3] However, the exact intracellular signaling mechanisms remain elusive due to the absence of apparent intracellular signaling motifs in DIP-β.[16,55] The similarity in phenotypes between *dip-β* and *side-IV* mutants suggests that Side-IV also plays a role in facilitating L2 neurons' preference to form synapses with L4 neurons. Our study demonstrated that Dsyd-1 and Liprin-α, important signaling factors for active zone formation,[44,45] work downstream of Side-IV. Furthermore, Side-IV can anchor localization of Dsyd-1 to the proximal region, indicating that Side-IV causes accumulation of Dsyd-1 in L2-L4 contact sites and increases the L2 preference for L4. Simultaneously, competition for the synapse seeding factor Dsyd-1 is a limiting resource and can ensure relative preference for a synapse,[5,56] resulting in the inhibition of miswiring of L2. Similarly, Kirre, another major driver of

L2-L4 synaptogenesis,[51] can increase miswiring if not properly regulated (Figures 7D–

K). These results suggest that establishing a signaling complex between key-lock

molecules and synapse formation factors, with the appropriate ratio and subcellular

localization, is crucial to achieving synaptic specificity. Our study proposes that one

example of this is Side-IV complex formation with Kirre and Dsyd-1, which initiates

synaptogenesis, coupled with the *trans*-synaptic interaction of Side-IV and Beat-IIb,

which determines synaptic specificity.

ACKNOWLEDGMENT

We gratefully acknowledge Dr. Filipe Pinto-Teixeira at Molecular, Cellular and

Developmental biology unit (MCD) in Center for Integrative Biology for providing the

side-IV[MI10049]-T2AGal4 flies. We thank the Bloomington Drosophila Stock Center,

Vienna Drosophila Resource Center (VDRC), Kyoto Drosophila Stock Center (Kyoto-

Fly), and FlyORF for providing fly stocks. We thank Enago (www.enago.jp) for the

English language review. This work was supported by JSPS KAKENHI #21J12660 (J.

O.), a Grant-in-Aid for Scientific Research (B) and a Grant-in-Aid for Scientific Research

on Innovative Areas from MEXT (#16H06457, #21H05682 and #21H02483 (T. S.)), a

Takeda Visionary Research Grant from the Takeda Science Foundation (T. S.).

AUTHOR CONTRIBUTIONS

J.O., A.I., S.H.S., H.K., and T.S. designed experiments. J.O., A.I., S.H.S., and H.K.,

performed experiments and analyzed data. J.O., A.I., X.W., and R.I. generated transgenic

reagents. J.O. and T.S. wrote the manuscript with reviewing from all co-authors.

DECLARATION OF INTERESTS

The authors declare that we have no competing interests.

REFERENCES

1. Sperry, R.W. (1963). CHEMOAFFINITY IN THE ORDERLY GROWTH OF NERVE FIBER PATTERNS AND CONNECTIONS. Proceedings of the National Academy of Sciences of the United States of America *50*, 703-710. 10.1073/pnas.50.4.703.

2. Sanes, J.R., and Zipursky, S.L. (2020). Synaptic Specificity, Recognition Molecules, and Assembly of Neural Circuits. Cell *181*, 536-556. 10.1016/j.cell.2020.04.008.

3. Xu, C., Theisen, E., Maloney, R., Peng, J., Santiago, I., Yapp, C., Werkhoven, Z., Rumbaut, E., Shum, B., Tarnogorska, D., et al. (2019). Control of Synaptic Specificity by Establishing a Relative Preference for Synaptic Partners. Neuron *103*, 865-877.e867. 10.1016/j.neuron.2019.06.006.

4. Kiral, F.R., Dutta, S.B., Linneweber, G.A., Hilgert, S., Poppa, C., Duch, C., von Kleist, M., Hassan, B.A., and Hiesinger, P.R. (2021). Brain connectivity inversely scales with developmental temperature in Drosophila. Cell Reports *37*, 110145. https://doi.org/10.1016/j.celrep.2021.110145.

5. Özel, M.N., Kulkarni, A., Hasan, A., Brummer, J., Moldenhauer, M., Daumann, I.M., Wolfenberg, H., Dercksen, V.J., Kiral, F.R., Weiser, M., et al. (2019). Serial Synapse Formation through Filopodial Competition for Synaptic Seeding Factors. Developmental cell *50*, 447-461.e448. 10.1016/j.devcel.2019.06.014.

6. Kiral, F.R., Linneweber, G.A., Mathejczyk, T., Georgiev, S.V., Wernet, M.F., Hassan, B.A., von Kleist, M., and Hiesinger, P.R. (2020). Autophagy-dependent filopodial kinetics restrict synaptic partner choice during Drosophila brain wiring. Nature communications *11*, 1325. 10.1038/s41467-020-14781-4.

7. Hiesinger, P.R. (2021). Brain wiring with composite instructions. BioEssays : news and reviews in molecular, cellular and developmental biology *43*, e2000166. 10.1002/bies.202000166.

8. Südhof, T.C. (2021). The cell biology of synapse formation. The Journal of cell biology *220*. 10.1083/jcb.202103052.

9. Takemura, S.Y., Bharioke, A., Lu, Z., Nern, A., Vitaladevuni, S., Rivlin, P.K., Katz, W.T., Olbris, D.J., Plaza, S.M., Winston, P., et al. (2013). A visual motion detection circuit suggested by Drosophila connectomics. Nature *500*, 175-181. 10.1038/nature12450.

10. Takemura, S.-y., Xu, C.S., Lu, Z., Rivlin, P.K., Parag, T., Olbris, D.J., Plaza, S., Zhao, T., Katz, W.T., Umayam, L., et al. (2015). Synaptic circuits and their variations within different columns in the visual system of Drosophila. Proceedings of the National Academy of Sciences of the United States of America *112*, 13711-13716. 10.1073/pnas.1509820112.

11. Davis, F.P., Nern, A., Picard, S., Reiser, M.B., Rubin, G.M., Eddy, S.R., and
Henry, G.L. (2020). A genetic, genomic, and computational resource for
exploring neural circuit function. eLife *9*, e50901. 10.7554/eLife.50901.
12. Cosmanescu, F., Katsamba, P.S., Sergeeva, A.P., Ahlsen, G., Patel, S.D.,
Brewer, J.J., Tan, L., Xu, S., Xiao, Q., Nagarkar-Jaiswal, S., et al. (2018).
Neuron-Subtype-Specific Expression, Interaction Affinities, and Specificity
Determinants of DIP/Dpr Cell Recognition Proteins. Neuron *100*, 1385-
1400.e1386. 10.1016/j.neuron.2018.10.046.
13. Kurmangaliyev, Y.Z., Yoo, J., Valdes-Aleman, J., Sanfilippo, P., and Zipursky,
S.L. (2020). Transcriptional Programs of Circuit Assembly in the Drosophila
Visual System. Neuron *108*, 1045-1057.e1046. 10.1016/j.neuron.2020.10.006.
14. Özel, M.N., Simon, F., Jafari, S., Holguera, I., Chen, Y.C., Benhra, N., El-
Danaf, R.N., Kapuralin, K., Malin, J.A., Konstantinides, N., and Desplan, C.
(2021). Neuronal diversity and convergence in a visual system developmental
atlas. Nature *589*, 88-95. 10.1038/s41586-020-2879-3.
15. Jain, S., Lin, Y., Kurmangaliyev, Y.Z., Valdes-Aleman, J., LoCascio, S.A.,
Mirshahidi, P., Parrington, B., and Zipursky, S.L. (2022). A global timing
mechanism regulates cell-type-specific wiring programmes. Nature *603*, 112-
118. 10.1038/s41586-022-04418-5.
16. Özkan, E., Carrillo, R.A., Eastman, C.L., Weiszmann, R., Waghray, D.,
Johnson, K.G., Zinn, K., Celniker, S.E., and Garcia, K.C. (2013). An
extracellular interactome of immunoglobulin and LRR proteins reveals receptor-
ligand networks. Cell *154*, 228-239. 10.1016/j.cell.2013.06.006.
17. Tan, L., Zhang, K.X., Pecot, M.Y., Nagarkar-Jaiswal, S., Lee, P.-T., Takemura,
S.-Y., McEwen, J.M., Nern, A., Xu, S., Tadros, W., et al. (2015). Ig Superfamily
Ligand and Receptor Pairs Expressed in Synaptic Partners in Drosophila. Cell
163, 1756-1769. 10.1016/j.cell.2015.11.021.
18. Carrillo, R.A., Özkan, E., Menon, K.P., Nagarkar-Jaiswal, S., Lee, P.T., Jeon,
M., Birnbaum, M.E., Bellen, H.J., Garcia, K.C., and Zinn, K. (2015). Control of
Synaptic Connectivity by a Network of Drosophila IgSF Cell Surface Proteins.
Cell *163*, 1770-1782. 10.1016/j.cell.2015.11.022.
19. Sergeeva, A.P., Katsamba, P.S., Cosmanescu, F., Brewer, J.J., Ahlsen, G.,
Mannepalli, S., Shapiro, L., and Honig, B. (2020). DIP/Dpr interactions and the
evolutionary design of specificity in protein families. Nature communications
11, 2125. 10.1038/s41467-020-15981-8.
20. Barish, S., Nuss, S., Strunilin, I., Bao, S., Mukherjee, S., Jones, C.D., and
Volkan, P.C. (2018). Combinations of DIPs and Dprs control organization of
olfactory receptor neuron terminals in Drosophila. PLoS genetics *14*, e1007560.
10.1371/journal.pgen.1007560.
21. suppXu, S., Xiao, Q., Cosmanescu, F., Sergeeva, A.P., Yoo, J., Lin, Y.,
Katsamba, P.S., Ahlsen, G., Kaufman, J., Linaval, N.T., et al. (2018).
Interactions between the Ig-Superfamily Proteins DIP-α and Dpr6/10 Regulate
Assembly of Neural Circuits. Neuron *100*, 1369-1384.e1366.
10.1016/j.neuron.2018.11.001.
22. Courgeon, M., and Desplan, C. (2019). Coordination between stochastic and
deterministic specification in the Drosophila visual system. Science (New York,
N.Y.) *366*. 10.1126/science.aay6727.

23.   Venkatasubramanian, L., Guo, Z., Xu, S., Tan, L., Xiao, Q., Nagarkar-Jaiswal,
S., and Mann, R.S. (2019). Stereotyped terminal axon branching of leg motor
neurons mediated by IgSF proteins DIP-α and Dpr10. eLife *8*, e42692.
10.7554/eLife.42692.
24.   Bornstein, B., Meltzer, H., Adler, R., Alyagor, I., Berkun, V., Cummings, G.,
Reh, F., Keren-Shaul, H., David, E., Riemensperger, T., and Schuldiner, O.
(2021). Transneuronal Dpr12/DIP-δ interactions facilitate compartmentalized
dopaminergic innervation of Drosophila mushroom body axons. The EMBO
journal *40*, e105763. 10.15252/embj.2020105763.
25.   Vactor, D.V., Sink, H., Fambrough, D., Tsoo, R., and Goodman, C.S. (1993).
Genes that control neuromuscular specificity in Drosophila. Cell *73*, 1137-1153.
10.1016/0092-8674(93)90643-5.
26.   Fambrough, D., and Goodman, C.S. (1996). The Drosophila beaten path gene
encodes a novel secreted protein that regulates defasciculation at motor axon
choice points. Cell *87*, 1049-1058. 10.1016/s0092-8674(00)81799-7.
27.   Sink, H., Rehm, E.J., Richstone, L., Bulls, Y.M., and Goodman, C.S. (2001).
sidestep encodes a target-derived attractant essential for motor axon guidance in
Drosophila. Cell *105*, 57-67. 10.1016/s0092-8674(01)00296-3.
28.   Siebert, M., Banovic, D., Goellner, B., and Aberle, H. (2009). Drosophila motor
axons recognize and follow a Sidestep-labeled substrate pathway to reach their
target fields. Genes Dev *23*, 1052-1062. 10.1101/gad.520509.
29.   Li, H., Watson, A., Olechwier, A., Anaya, M., Sorooshyari, S.K., Harnett, D.P.,
Lee, H.-K.P., Vielmetter, J., Fares, M.A., Garcia, K.C., et al. (2017).
Deconstruction of the beaten Path-Sidestep interaction network provides insights
into neuromuscular system development. eLife *6*, e28111. 10.7554/eLife.28111.
30.   Pipes, G.C., Lin, Q., Riley, S.E., and Goodman, C.S. (2001). The Beat
generation: a multigene family encoding IgSF proteins related to the Beat axon
guidance molecule in Drosophila. Development (Cambridge, England) *128*,
4545-4552. 10.1242/dev.128.22.4545.
31.   Aberle, H. (2009). Searching for guidance cues: follow the Sidestep trail. Fly *3*,
270-273. 10.4161/fly.9790.
32.   Zinn, K. (2009). Choosing the road less traveled by: a ligand-receptor system
that controls target recognition by Drosophila motor axons. Genes &
development *23*, 1042-1045. 10.1101/gad.1803009.
33.   Kinold, J.C., Pfarr, C., and Aberle, H. (2018). Sidestep-induced neuromuscular
miswiring causes severe locomotion defects in Drosophila larvae. Development
(Cambridge, England) *145*. 10.1242/dev.163279.
34.   Kind, E., Longden, K.D., Nern, A., Zhao, A., Sancer, G., Flynn, M.A.,
Laughland, C.W., Gezahegn, B., Ludwig, H.D.F., Thomson, A.G., et al. (2021).
Synaptic targets of photoreceptors specialized to detect color and skylight
polarization in Drosophila. eLife *10*, e71858. 10.7554/eLife.71858.
35.   Heymann, C., Paul, C., Huang, N., Kinold, J.C., Dietrich, A.C., and Aberle, H.
(2022). Molecular insights into the axon guidance molecules Sidestep and
Beaten path. Frontiers in physiology *13*, 1057413. 10.3389/fphys.2022.1057413.
36.   Karuppudurai, T., Lin, T.-Y., Ting, C.-Y., Pursley, R., Melnattur, Krishna V.,
Diao, F., White, Benjamin H., Macpherson, Lindsey J., Gallio, M., Pohida, T.,
and Lee, C.-H. (2014). A Hard-Wired Glutamatergic Circuit Pools and Relays

UV Signals to Mediate Spectral Preference in Drosophila. Neuron *81*, 603-615.
https://doi.org/10.1016/j.neuron.2013.12.010.
37. Chen, Y., Akin, O., Nern, A., Tsui, C.Y., Pecot, M.Y., and Zipursky, S.L.
(2014). Cell-type-specific labeling of synapses in vivo through synaptic tagging
with recombination. Neuron *81*, 280-293. 10.1016/j.neuron.2013.12.021.
38. Nagarkar-Jaiswal, S., Lee, P.-T., Campbell, M.E., Chen, K., Anguiano-Zarate,
S., Gutierrez, M.C., Busby, T., Lin, W.-W., He, Y., Schulze, K.L., et al. (2015).
A library of MiMICs allows tagging of genes and reversible, spatial and
temporal knockdown of proteins in Drosophila. eLife *4*, e05338.
10.7554/eLife.05338.
39. Tanaka, H., Miyazaki, N., Matoba, K., Nogi, T., Iwasaki, K., and Takagi, J.
(2012). Higher-order architecture of cell adhesion mediated by polymorphic
synaptic adhesion molecules neurexin and neuroligin. Cell reports *2*, 101-110.
10.1016/j.celrep.2012.06.009.
40. Gao, S., Takemura, S.Y., Ting, C.Y., Huang, S., Lu, Z., Luan, H., Rister, J.,
Thum, A.S., Yang, M., Hong, S.T., et al. (2008). The neural substrate of spectral
preference in Drosophila. Neuron *60*, 328-342. 10.1016/j.neuron.2008.08.010.
41. Fischbach, K.F., Linneweber, G.A., Andlauer, T.F., Hertenstein, A., Bonengel,
B., and Chaudhary, K. (2009). The irre cell recognition module (IRM) proteins.
Journal of neurogenetics *23*, 48-67. 10.1080/01677060802471668.
42. Strutt, H., and Strutt, D. (2008). Differential stability of flamingo protein
complexes underlies the establishment of planar polarity. Current biology : CB
18, 1555-1564. 10.1016/j.cub.2008.08.063.
43. Hakeda-Suzuki, S., Berger-Müller, S., Tomasi, T., Usui, T., Horiuchi, S.Y.,
Uemura, T., and Suzuki, T. (2011). Golden Goal collaborates with Flamingo in
conferring synaptic-layer specificity in the visual system. Nature neuroscience
14, 314-323. 10.1038/nn.2756.
44. Owald, D., Khorramshahi, O., Gupta, V.K., Banovic, D., Depner, H., Fouquet,
W., Wichmann, C., Mertel, S., Eimer, S., Reynolds, E., et al. (2012).
Cooperation of Syd-1 with Neurexin synchronizes pre- with postsynaptic
assembly. Nature neuroscience *15*, 1219-1226. 10.1038/nn.3183.
45. Fouquet, W., Owald, D., Wichmann, C., Mertel, S., Depner, H., Dyba, M.,
Hallermann, S., Kittel, R.J., Eimer, S., and Sigrist, S.J. (2009). Maturation of
active zone assembly by Drosophila Bruchpilot. The Journal of cell biology *186*,
129-145. 10.1083/jcb.200812150.
46. Rivera-Alba, M., Vitaladevuni, S.N., Mishchenko, Y., Lu, Z., Takemura, S.Y.,
Scheffer, L., Meinertzhagen, I.A., Chklovskii, D.B., and de Polavieja, G.G.
(2011). Wiring economy and volume exclusion determine neuronal placement in
the Drosophila brain. Current biology : CB *21*, 2000-2005.
10.1016/j.cub.2011.10.022.
47. Varoqueaux, F., Aramuni, G., Rawson, R.L., Mohrmann, R., Missler, M.,
Gottmann, K., Zhang, W., Südhof, T.C., and Brose, N. (2006). Neuroligins
determine synapse maturation and function. Neuron *51*, 741-754.
10.1016/j.neuron.2006.09.003.
48. Gokce, O., and Südhof, T.C. (2013). Membrane-tethered monomeric neurexin
LNS-domain triggers synapse formation. The Journal of neuroscience : the

official journal of the Society for Neuroscience *33*, 14617-14628.
10.1523/jneurosci.1232-13.2013.

49.   Fischbach, K.F., and Dittrich, A.P.M. (1989). The optic lobe of Drosophila
melanogaster. I. A Golgi analysis of wild-type structure. Cell and Tissue
Research *258*, 441-475. 10.1007/BF00218858.

50.   Kolodziejczyk, A., Sun, X., Meinertzhagen, I.A., and Nässel, D.R. (2008).
Glutamate, GABA and acetylcholine signaling components in the lamina of the
Drosophila visual system. PLoS One *3*, e2110-e2110.
10.1371/journal.pone.0002110.

51.   Lüthy, K., Ahrens, B., Rawal, S., Lu, Z., Tarnogorska, D., Meinertzhagen, I.A.,
and Fischbach, K.F. (2014). The irre cell recognition module (IRM) protein
Kirre is required to form the reciprocal synaptic network of L4 neurons in the
Drosophila lamina. Journal of neurogenetics *28*, 291-301.
10.3109/01677063.2014.883390.

52.   Weavers, H., Prieto-Sánchez, S., Grawe, F., Garcia-López, A., Artero, R.,
Wilsch-Bräuninger, M., Ruiz-Gómez, M., Skaer, H., and Denholm, B. (2009).
The insect nephrocyte is a podocyte-like cell with a filtration slit diaphragm.
Nature *457*, 322-326. 10.1038/nature07526.

53.   Kaufmann, N., DeProto, J., Ranjan, R., Wan, H., and Van Vactor, D. (2002).
Drosophila liprin-alpha and the receptor phosphatase Dlar control synapse
morphogenesis. Neuron *34*, 27-38. 10.1016/s0896-6273(02)00643-8.

54.   Jiang, X., Sando, R., and Südhof, T.C. (2021). Multiple signaling pathways are
essential for synapse formation induced by synaptic adhesion molecules.
Proceedings of the National Academy of Sciences of the United States of
America *118*. 10.1073/pnas.2000173118.

55.   Lobb-Rabe, M., Nawrocka, W.I., Carrillo, R.A., and Özkan, E. (2023). Neuronal
wiring receptors Dprs and DIPs are GPI anchored and this modification
contributes to their cell surface organization. bioRxiv, 2023.2003.2002.530872.
10.1101/2023.03.02.530872.

56.   Simon, F., and Konstantinides, N. (2021). Single-cell transcriptomics in the
Drosophila visual system: Advances and perspectives on cell identity regulation,
connectivity, and neuronal diversity evolution. Developmental biology *479*,
107-122. 10.1016/j.ydbio.2021.08.001.

57.   Kondo, S., and Ueda, R. (2013). Highly improved gene targeting by germline-
specific Cas9 expression in Drosophila. Genetics *195*, 715-721.
10.1534/genetics.113.156737.

58.   Trush, O., Liu, C., Han, X., Nakai, Y., Takayama, R., Murakawa, H., Carrillo,
J.A., Takechi, H., Hakeda-Suzuki, S., Suzuki, T., and Sato, M. (2019). N-
Cadherin Orchestrates Self-Organization of Neurons within a Columnar Unit in
the Drosophila Medulla. The Journal of Neuroscience *39*, 5861-
5880. 10.1523/jneurosci.3107-18.2019.

59.   Thorpe, H.M., and Smith, M.C. (1998). In vitro site-specific integration of
bacteriophage DNA catalyzed by a recombinase of the resolvase/invertase
family. Proceedings of the National Academy of Sciences of the United States of
America *95*, 5505-5510. 10.1073/pnas.95.10.5505.

60.   Hakeda-Suzuki, S., Takechi, H., Kawamura, H., and Suzuki, T. (2017). Two
receptor tyrosine phosphatases dictate the depth of axonal stabilizing layer in the
visual system. eLife *6*, e31812. 10.7554/eLife.31812.
61.   Owald, D., Fouquet, W., Schmidt, M., Wichmann, C., Mertel, S., Depner, H.,
Christiansen, F., Zube, C., Quentin, C., Körner, J., et al. (2010). A Syd-1
homologue regulates pre- and postsynaptic maturation in Drosophila. Journal of
Cell Biology *188*, 565-579. 10.1083/jcb.200908055.
62.   Sugie, A., Hakeda-Suzuki, S., Suzuki, E., Silies, M., Shimozono, M., Möhl, C.,
Suzuki, T., and Tavosanis, G. (2015). Molecular Remodeling of the Presynaptic
Active Zone of Drosophila Photoreceptors via Activity-Dependent Feedback.
Neuron *86*, 711-725. 10.1016/j.neuron.2015.03.046.
63.   Bolte, S., and Cordelières, F.P. (2006). A guided tour into subcellular
colocalization analysis in light microscopy. Journal of microscopy *224*, 213-232.
10.1111/j.1365-2818.2006.01706.x.
64.   Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch,
T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an
open-source platform for biological-image analysis. Nature methods *9*, 676-682.
10.1038/nmeth.2019.

LEGENDS TO FIGURES

Figure 1. The Interaction of Side-IV and Beat-IIb is Instructive for Synaptic Induction

(A−B") The synapse and axon of R7. Control (A−A"), *side-IV* OE (B−B"). Dotted lines indicate M1−M3 layer. Arrows and arrowheads indicate ectopic synapses and overshoot axons, respectively.

(C−E') Brp-GFP puncta of R7. *side-IV* OE in Wild Type (C and C'), *beat-IIa* mutant (D and D'), and *beat-IIb* mutant (E and E') background. Dotted lines indicate M1−M3 layer.

(F) Quantification of the number of the ectopic synapse (n > 60 axons).

(G) Quantification of the rate of overshoot axon (n >10 brains).

(H−H''') The localization of Beat-IIb and synapses of R7. Dotted lines indicate M1−M3

layer.

(I) The phenotypic scheme of *side-IV* OE.

(J−K'') The punctate signal of Beat-IIb by *side-IV* OE. Control (J−J''), *side-IV* OE (K−

K''). Arrows indicate punctate Beat-IIb signal.

(L) Mutual localization of Side-IV, Beat-IIb, and Brp of R7. Dotted lines indicate M1−

M3 layer. Arrows and arrowhead (L−L''') indicate co-localized Side-IV and Beat-IIb

puncta with and without Brp-mCherry puncta, respectively.

(M) The summary scheme of Side-IV clusters, Beat-IIb clusters, and synaptic induction.

Statistical treatment of the number synapse and the rate of overshoot axon were analyzed

with the Mann-Whitney test. In this and all following statistical significance. n.s. $P > 0.05$,

$* P \leq 0.05$, $** P \leq 0.01$, $*** P \leq 0.001$, $**** P \leq 0.0001$.

Figure 2. The Chimeric Analysis of Side-IV and mCD8

(A) The domain structure of Side-IV and mCD8.

(B) The variants of Side-IV domain deletion or chimera.

(C−G''') Mutual localization of Side-IV, Beat-IIb, and Brp of R7. Dotted lines indicate

M1−M3 layer.

(H) Quantification of the rate of Side-IV-V5 variants clustered axon (n > 28 axons).

(I) Quantification of the rate of Beat-IIb clustered axon (n > 47 axons).

(J) Quantification of the number of the ectopic synapse (n > 32 axons).

(K) Quantification of the rate of overshot axon (n > 6 brains).

Statistical treatment of the rate of Side-IV-V5 variants or Beat-IIb clustered axon were

analyzed with the Chi-square test. Statistical treatment of the number of the ectopic

synapse and the rate of overshoot axon were analyzed with the Mann-Whitney test.

Figure 3. Kirre Interacts with Side-IV through the Extracellular Domain and

Works as a Co-receptor

(A–E) The synapse and axon of R7. Control (A and A'), *kirre* OE (B and B'), Control

UAS (*luciferase* RNAi) + *side-IV* OE (C and C'), *kirre* OE + *side-IV* OE (D and D').

Dotted lines indicate M1−M3 layer.

(E) Quantification of the number of the ectopic synapse (n > 44 axons).

(F) Quantification of the rate of overshoot axon (n > 4 brains).

(G−O) Photoreceptors in medulla with co-expression of *side-IV* and *kirre* variants.

Control (G), *kirre* OE (H), *kirre* $^{\Delta Cyto}$ OE (I), *side-IV* OE (J), *side-IV* + *kirre* OE (K),

side-IV+kirre $^{\Delta Cyto}$ OE (L), *side-IV* $^{\Delta Cyto}$ OE (M), *side-IV* $^{\Delta Cyto}$ + *kirre* OE (N), *side-IV*

$^{\Delta Cyto}$ + *kirre* $^{\Delta Cyto}$ OE (O).

(P) Quantification of the rate of overshoot axon (n > 4 brains).

(Q−V) Protein accumulation at cell boundaries in transiently transfected S2 cells

subjected to aggregation assay. The S2 cells were co-transfected with Act5c-gal4. UAS-

side-IV-mCherry and UAS-*kirre*-GFP (Q–Q"). (R) Distribution of accumulation ratio of

Side-IV and Kirre. UAS-*side-IV*-mCherry and UAS-*cadE*-GFP (S–S"). (T) Distribution

of accumulation ratio of Side-IV and CadE. (U) Distribution of accumulation ratio of

Side-IV co-transfected with Kirre, Kirre $^{\Delta Cyto}$, or CadE. (V) Distribution of

accumulation ratio of Side-IV variants co-transfected with Kirre. The sample size is

indicated (n).

Statistical treatment of the number of synapse and the rate of overshoot axon were

analyzed with the Mann-Whitney test. Distribution of the cell-cell contact accumulation

ration was analyzed with the unpaired t-test.

Figure 4. Side-IV Transduces a Signal with Dsyd-1 and Liprin-α through the

Cytoplasmic Tail

(A−D) The synapse and axon of R7.Control (A and A', same images as 3A and A'),

Control RNAi (*luciferase* RNAi) + *side-IV* OE (B and B', same images as 3C and C'),

dsyd-1 RNAi + *side-IV* OE (C and C'), *liprin-α* RNAi + *side-IV* OE (D and D'). Dotted

lines indicate M1−M3 layer.

(E) Quantification of the number of the ectopic synapse (n > 69 axons).

(F) Quantification of the rate of overshoot axon (n > 6 brains).

(G−H'') The localization of Side-IV and Dsyd-1. The S2 cells were co-transfected with

Act5c-gal4, UAS-*dsyd-1*-mCherry, and UAS-*side-IV* ^{FL}-GFP or UAS-*side-IV* $^{\Delta PDZb}$.

Side-IV ^{FL}-GFP and Dsyd-1-mCherry (G−G''), Side-IV $^{\Delta PDZb}$-GFP and Dsyd-1-mCherry

(H−H'').

(I) The quantification of the co-localization (n > 35 images).

(J−K'') The localization of Side-IV and Liprin-α. The S2 cells were co-transfected with

UAS-*side-IV*-GFP, UAS-*liprin-a*-mCherry with and without UAS-*dsyd-1*-V5. Side-IV-

GFP and Liprin-α-mCherry (J−J"), Side-IV-GFP, Liprin-α-mCherry, and Dsyd-1-V5

(K−K").

(L) The quantification of the co-localization (n > 35 images).

(M) Quantification of the number of ectopic synapses (n > 43 axons) represented as

mean ± SD.

(N) The working model of synaptic induction by Side-IV.

Statistical treatment of the number of synapse and the rate of overshoot axons were

analyzed with the Mann-Whitney test. Statistical treatment of the Co-localization index

was analyzed with the unpaired t-test.

Figure 5. *side-IV* **and** *side-VII* **are Required for Proper Synapse Formation**

Patterns of the L2 Neurons

(A) Scheme of synaptic connection patterns of lamina neurons in lamina.

(B) Scheme of predicted Beat-Side interactions between L2 and L4 neurons.

(C−F') The presynaptic structures and cell morphology of Lamina neuron (LN). Control

(C and C'), *side-IV*$^{\Delta 4-1}$ mutant (D and D'), *side-VII*$^{\Delta E3-4-1}$ mutant (E and E'), *side-IV* /

side-VII double mutant (F and F'). Dotted lines indicate the proximal and distal

boundary.

(G) Quantification of the number of synapses in the proximal region (n > 82 axons).

(H) Quantification of the number of synapses in the distal region (n > 107 axons).

(I−J') The synapse and axon of L2. Control (I and I'), *side-IV*$^{\Delta 4-1}$ / *side-VII*$^{\Delta E3-4-1}$ double

mutant (J and J'). Dotted lines indicate the proximal and distal boundary.

(K) Quantification of synapses in the proximal region of the L2 neuron (n > 88 axons).

(L) Quantification of synapses in the distal region of the L2 neuron (n > 84 axons).

Statistical treatment of the number of synapses in the proximal region and distal region

were analyzed with the unpaired t-test and Mann-Whitney test, respectively.

Figure 6. Side-IV and Side-VII Showed Beats-dependent Subcellular Localization

in the Proximal Region

(A−A''') Mutual localization of endogenous Brp, Side-IV, and Side-VII at 72h after

pupal formation (APF). Merge (A), Brp (A'), Side-IV (A''), and Side-VII (A'''). Dotted

lines indicate the proximal and distal boundary.

(B−C') Localization of Side-IV (B and B') and Side-VII (C and C') in the L2 neuron.

Dotted lines indicate the boundary between the proximal and distal region.

(D−G') Endogenous localization of Side-IV in the lamina neuron. Control 48h APF (D

and D'), *beat-IIa^{+2}* mutant + *beat-IIb* RNAi 48h APF (E and E'). Control 72h APF (F

and F'), *beat-IIa^{+2}* mutant + *beat-IIb* RNAi 72h APF (G and G'). Dotted lines indicate

the proximal and distal boundary.

(H) Quantification of the rate of the axons with Side-IV localization in the proximal

region in the 72h APF (n > 149 axons).

(I−M') Endogenous localization of Side-VII in the lamina neuron. Control 48h APF (I

and I'), *beat-IV* RNAi 48h APF (J and J'). Control 72h APF (L and L'), *beat-IV* RNAi

72h APF (M and M'). Dotted lines indicate the proximal and distal boundary.

(H) Quantification of the rate of the axons with Side-VII localization in the proximal

region in the 72h APF (n > 147 axons).

Statistical treatment of the rate of Side-IV or Side-VII positive axon was analyzed with

the Chi-square test.

Figure. 7 Side-IV Limit the Dysd-1 Localization in the Proximal and Inhibit the

Synapse Formation by Kirre in the Distal Region.

(A) The mutual localization Kirre and Side-IV on L2 neuron at 72h APF. Merge (A),

magnified Merge (A'), Side-IV-GFP (A"), Kirre-HA (A'''). Compared with Side-IV

localization, Kirre puncta signals were observed in the distal region (yellow arrowheads

in A'''). Dotted lines indicate the proximal and distal boundary.

(B) The fluorescent intensity of Side-IV-GFP (green) and Kirre-HA (magenta)

measured from proximal to distal across L2 axons as shown in A" and A''' (dotted lines).

The average intensity of 29 axons was calculated.

(C) Schematic of Side-IV (green) and Kirre (magenta) localization in the L2 neuron.

(D−I) The presynaptic structures of lamina neurons. Control (D, same image as 5C'),

side-IV mutant (E), *side-IV / side-VII* double mutant (F, same image as 5F'), *kirre*

heterozygous mutant (G), *kirre* heterozygous and *side-IV* mutant (H), *kirre*

heterozygous and *side-IV / side-VII* double mutant. Dotted lines indicate the proximal

and distal boundary.

(H) Quantification of synapses in the proximal region (n > 69 axons).

(H) Quantification of synapses in the distal region (n > 79 axons).

(L−N") The localization of endogenous Dsyd-1 in lamina neurons at 48h APF. Control

(L−L"), *side-IV* mutant (M−M"), *side-IV / side-VII* double mutant (N−N"). Dotted lines

indicate the proximal and distal boundary.

(O) The rate of Dsyd-1 diffused axon in the distal region (n > 92 axons).

(P−R') The localization of Dsyd-1 in lamina neuron at 72h APF Control (P and P'), *side-*

IV mutant (Q and Q'), *side-IV* / *side-VII* double mutant (R and R'). Dotted lines indicate

the proximal and distal boundary.

(S) Quantification of the number of Dsyd-1 puncta in the proximal region (n > 120

axons).

(T) Quantification of the number of Dsyd-1 puncta in the distal region (n > 96 axons).

Statistical treatment of the number of Brp-GFP or Dsyd-1 puncta in the proximal region

and distal region were analyzed with the unpaired t-test and Mann-Whitney test,

respectively. The rate of Dsyd-1 diffused axon was analyzed with by the Chi-square

test.

Figure. 8 Side-IV-Beat-IIb's Complex Formation Model.

The new key-lock combination Side-IV-Beat-IIb made cluster complex and transduced

bifurcated signaling through co-receptor, Kirre and scaffold protein complex, Dsyd-1-

Liprin-α. Side-IV can specific the loci of synapse formation and anchor the Dsyd-1 to

inhibit miswiring.

LEGENDS TO SUPPLEMENTAL FIGURES

Figure S1. Mis-expression of *side-IV* in R7 induced ectopic synapse formation in the

M1–M3 layer. (A and B) The Beat/Side double knockdown experiments. Quantification

of the number of synapses in the M1−M3 layer (A) and M4−6 layer (B) per axon. Control

(n=119 axons/4 brains), *beat-IIb* & *beat-IIIc* RNAi (n=70/4), *side* & *side-III* RNAi

(n=164/6), *side* & *side-V* RNAi (n=184/7), *side* & *side-VII* RNAi (n=108/4), *side-III* &

side-V RNAi (n=118/4), *side-III* & *side-VII* RNAi (n=96/4), *side-V* & *side-VII* (n=100/4).

(C and D) The Beat/Side overexpression (OE) experiments. Quantification of the number

of synapses in the M1−M3 layer (C) and M4−6 layer (D) per axon. Control (n=119

axons/4 brains), *beat-IIa* OE (n=145/6), *beat-IIb* OE (n=145/6), *beat-VI* OE (n=71/4),

side OE (n=132/4), *side-IV* OE (n=40/2), *side-VII* OE (n=34/2), *side-VIII* OE (n=104/4).

(E) Scheme of tissue-specific visualization of endogenous Side-IV. The FRT-stop-FRT-

GFP cassette was knocked in just before the stop codon of *side-IV*. By expressing FLPase,

endogenous localization of Side-IV can be observed in a tissue-specific manner. (F and

F') Localization of Side-IV in the optic lobe at the adult stage. heatshock-FLP was used

to visualize Side-IV in all cells. (G−I') Expression of endogenous *side-IV* of R7 in a

developmental stage. The endogenous Side-IV on R7 was visualized with *side-IV*-FsF-

GFP and 20C11-FLP. Axons of photoreceptor R7 were visualized with U-myr-RFP

(magenta, GMR-FsF-Gal4 with 20C11-FLP). 24 h APF (G−G'), 48 h APF (H−H'), 72 h

APF (I−I'). No Side-IV-GFP signals were observed at any stages, indicating that R7

doesn't endogenously express *side-IV*. Statistical treatment of the number of Brp-GFP

puncta in the M1−M3 layer and M4−M6 layer were analyzed with the Mann-Whitney test

and unpaired t-test, respectively. In this and all following statistical treatment; n.s. $P >$

0.05, * $P ≤ 0.05$, ** $P ≤ 0.01$, *** $P ≤ 0.001$, **** $P ≤ 0.0001$.

Figure S2. The Domain deletion analysis of *side-IV* in R7 and S2 cultural cells. (A)

Domain structure of Side-IV. (B−H') The Brp-GFP signal of R7 with overexpression (OE)

of *side-IV* domain deletion variants. The Brp-GFP puncta (green) in R7 is visualized by

combining the STaR method with 20C11-FLP. Photoreceptors were visualized by 24B10

(magenta). Control (B and B'), *side-IV ^{FL}* OE (C and C'), *side-IV $^{\Delta Ig1-5}$* OE (D and D'),

side-IV $^{\Delta Ig1}$ OE (E and E'), *side-IV $^{\Delta FnIII}$* OE (F and F'), *side-IV $^{\Delta Cyto}$* OE (G and G'), *side-*

IV $^{\Delta PDZb}$ OE (H and H'). Orange dotted lines indicate region of M1−M3 layer. (I)

Quantification of the number of synapses in the M1−M3 layer per axon. Control (n= 84

axons/5 brains), *side-IV ^{FL}* OE (n= 60/5), *side-IV $^{\Delta Ig1-5}$* OE (n= 57/5), *side-IV $^{\Delta Ig1}$* OE (n=

70/5), *side-IV $^{\Delta FnIII}$* OE (n= 86/7), *side-IV $^{\Delta Cyto}$* OE (n= 33/3), *side-IV $^{\Delta PDZ}$* OE (n= 45/4).

(J) Quantification of the rate of overshoot axon per optic lobe. Control (n=12 brains),

side-IV ^{FL} OE (n= 10), *side-IV $^{\Delta Ig1-5}$* OE (n= 12), *side-IV $^{\Delta Ig1}$* OE (n= 10), *side-IV $^{\Delta FnIII}$* OE

(n= 10), *side-IV $^{\Delta Cyto}$* OE (n=13), *side-IV $^{\Delta PDZ}$* OE (n= 12). (K) Quantification of the rate

of Beat-IIb clustered per axon. Control (n= 81 axons/4 brains), *side-IV ^{FL}* OE (n= 161/7),

side-IV $^{\Delta Ig1-5}$ OE (n= 91/4), *side-IV $^{\Delta Ig1}$* OE (n= 133/6), *side-IV $^{\Delta FnIII}$* OE (n= 82/4), *side-IV*

$^{\Delta Cyto}$ OE (n= 68/4), *side-IV $^{\Delta PDZ}$* OE (n= 80/4). (L−P") Cell aggregation assay with

Drosophila S2 cell. S2 cells were individually transfected with UAS-*side-IV* variants or

mcd8-mCherry or UAS-*beat-IIb*-GFP. All cells were co-transfected with Actin5C-gal4.

After the transfection, cells were mixed together. Side-IV ^{FL}-mCherry vs Beat-IIb-GFP

(L−L"), Side-IV $^{\Delta Ig1}$-mCherry vs Beat-IIb-GFP (M−M"), Side-IV $^{\Delta Ig1−5}$-mCherry vs Beat-

IIb-GFP (N−N"), mCd8-mCherry vs Beat-IIb-GFP (O−O"), and Side-IV $^{Ecto\Delta FnIII}$::mCD8

vs Beat-IIb-GFP (P−P''). Yellow dotted blankets represent cell aggregations. (Q) The

quantification of the number of cells per cell aggregation. Side-IV ^{FL}-mCherry vs Beat-

IIb-GFP (n=70 cell aggregations), Side-IV $^{\Delta Ig1}$-mCherry vs Beat-IIb-GFP (n=51), Side-

IV $^{\Delta Ig1-5}$-mCherry vs Beat-IIb-GFP (n=40), mCd8-mCherry vs Beat-IIb-GFP (n=48),

Side-IV $^{Ecto\Delta FnIII}$::mCD8 vs Beat-IIb-GFP (n=86). Statistical treatment of the number of

synapses in the M1−M3 layer, the rate of overshoot axons, and the number of cell per cell

aggregation were analyzed with the Mann-Whitney test. Statistical treatment of the rate

of Beat-IIb clustered axon was performed with the Chi-square test.

Figure S3. Synapse was induced in the M1−M3 layer in R7 by introducing Nrxs-Nlgs

interactions but not CadE or hSDK2 homophilic interactions. (A−E') The Brp-GFP

signal of R7 with overexpression (OE) of *nrx-1* or *nlg1−3* with FlySAM in both R7 and

postsynaptic neuron by combining GMR-gal4 and Ort-gal4. The Brp-GFP puncta (green)

in R7 is visualized by combining the STaR method with 20C11-FLP. Photoreceptors were

visualized by 24B10 (magenta) Control (A and A'), *nrx-1* OE (B and B'), *nlg-1* OE (C and

C'), *nlg-2* OE (D and D'), *nlg-3* OE (E and E'). (F) Quantification of the number of

synapses in the M1-M3 layer per axon. Control (n= 114 axons/5 brains), *nrx-1* OE (n=

157 /5), *nlg-1* OE (n= 169/5), *nlg-2* OE (n= 58/3), *nlg-3* OE (n= 114/4). (G−I) The Brp-

GFP signal of R7 by overexpressing the molecules which have homophilic interaction in

R7 and postsynaptic neuron by combining GMR-gal4 and Ort-gal4. Control (G and G'),

cadE OE (H and H'), *hsdk2* OE (I and I'). Orange dotted lines indicate region of M1−M3

layer. (I) Quantification of the number of synapses in the M1−M3 layer per axon. Control

(n= 109 axons/7 brains), *cadE* OE (n= 67/5), *hSDK2* OE (n=73/5). Statistical treatment

of the number of synapses in the M1−M3 layer was analyzed with the Mann-Whitney test.

Figure S4. The biochemical and genetic interaction between Side-IV and Kirre.

(A) The result of Co-IP for Side-IV and Kirre. S2 cell were transfected with Actin5C-gal4

and UAS-*side-IV*-V5/GFP and/or UAS-*kirre*-GFP/V5. Immunoprecipitation (IP) and

Immunoblotting (blot) were performed with α-GFP and α-V5, respectively. The bands

between 100 kDa and 150 kDA were detected only when cells expressed both *side-IV* and

kirre, showing they make complex with interaction. Predicted molecular weights of Side-

IV and Kirre are about 110 kDa and 101kDa, respectively. (B) Aggregation assay of S2

cell transfected Actin5C-gal4 and UAS-*mcd8*-mCherry against Actin5C-gal4 and UAS-

kirre-GFP. Homophilic interaction of Kirre causes Kirre-transfected S2 cells to form

aggregates (Kirre-GFP cluster). (C) Aggregation assay of S2 cell transfected Actin5C-

gal4 and UAS-*side-IV*-mCherry against Actin5C-gal4 and UAS-*kirre*-GFP. (D)

Quantification of the rate of Kirre-GFP cluster that contains mCherry positive cell. If

Side-IV interacts with Kirre in *trans*, many of Kirre-GFP clusters should contain *side-IV*-

mCherry expressing cells. There was no significance of the rate of Kirre-GFP cluster that

has mCherry positive cell between aggregation assay of mCd8-mCherry vs Kirre-GFP

and Side-IV-mCherry vs Kirre-GFP, indicating Side-IV didn't interact with Kirre in *trans*.

The sample size is indicated (n). Statistical treatment was performed with the Chi-square

test. (E−L) Photoreceptors in medulla with co-expression of *side-IV* variant *kirre* with

GMR-gal4. *side-IV* $^{\Delta Ig1-5}$ OE (E), *side-IV* $^{\Delta Ig1-5}$ + *kirre* OE (F), *side-IV* $^{\Delta Ig1}$ OE (G), *side-*

IV $^{\Delta Ig1}$ + *kirre* OE (H), *side-IV* $^{\Delta FnIII}$ OE (I), *side-IV* $^{\Delta FnIII}$ + *kirre* OE (J), *side-IV* $^{\Delta PDZb}$ OE

(K), *side-IV* $^{\Delta PDZb}$ + *kirre* OE (L). (M) Quantification of the rate of overshoot axon with

side-IV variants and *kirre*. Control (n=12 brains), *kirre* OE (n=4), *side-IV* ^{FL} OE (n=10),

side-IV ^{FL} + *kirre* OE (n=6), *side-IV* $^{\Delta Ig1-5}$ OE (n=12), *side-IV* $^{\Delta Ig1-5}$ + *kirre* OE (n=6), *side-*

IV $^{\Delta Ig1}$ OE (n=10), *side-IV* $^{\Delta Ig1}$ + *kirre* OE (n=6), *side-IV* $^{\Delta FnIII}$ OE (n=10), *side-IV* $^{\Delta FnIII}$ OE

+ *kirre* OE (n=7), *side-IV* $^{\Delta Cyto}$ OE (n=13), *side-IV* $^{\Delta Cyto}$ + *kirre* OE (n=6), *side-IV* $^{\Delta PDZb}$

OE (n=12), *side-IV* $^{\Delta PDZb}$ + *kirre* OE (n=5). Statistical treatment of the rate of the Kirre-

GFP cell cluster with mCherry positive cell and the rate of overshoot axon was performed

with the Chi-square test and the Mann-Whitney test, respectively.

Figure S5. Disruption of Side-IV-Beat-IIa/b or Side-VII-Beat-IV interactions results

in ectopic synapse formation in the distal region. (A−C') Brp-GFP of lamina neuron in

rescue experiment. *side-IV* $^{\Delta 4-1/T2A-gal4}$ (A and A'), *side-IV* $^{\Delta 4-1/T2A-gal4}$ + Side-IV ^{FL} (B and

B'), *side-IV* $^{\Delta 4-1/T2A-gal4}$ + Side-IV $^{\Delta Cyto}$ (C and C'). Orange dotted lines indicate the

boundary between the proximal and distal region. (D) Quantification of Brp-GFP puncta

in the distal region in the rescue experiment. *side-IV* $^{\Delta 4-1/+}$ (n= 88 axons/ 3 brains), *side-IV*

$^{T2A-gal4/+}$ (n=90/3), *side-IV* $^{\Delta 4-1/T2A-gal4}$ (n= 134/4), *side-IV* $^{\Delta 4-1/T2A-gal4}$ + Side-IV ^{FL} (n= 90/3),

side-IV $^{\Delta 4-1/T2A-gal4}$ + Side-IV $^{\Delta Cyto}$ (n= 86/3). (E−G') Brp-GFP of lamina neuron in L2

specific *side-IV* or *side-VII* mutant. L2 neuron (magenta, L2) were visualized by

combining UAS-myr-RFP with L2-specific gal4 driver (GMR16H03-gal4). L2-specific

side-IV or *side-VII* mutant means *side-IV*$^{Δ4\text{-}1}$ or *side-VII*$^{ΔE3\text{-}4\text{-}1}$ heterozygote with L2-

specific RNAi [GMR16H03-Gal4, UAS-RNAi]). Control (E and E'), L2 specific *side-IV*

mutant (F and F'), L2 specific *side-VII* mutant (G and G'). Orange dotted lines indicate

the boundary between the proximal and distal region. (H) Quantification of Brp-GFP

puncta in L2 specific *side-IV* or *side-VII* mutant. Control (n=96 axons/3 brains), L2

specific *side-IV* mutant (n=160/4), L2 specific *side-VII* mutant (n=129/3). (I−L') Brp-GFP

of lamina neuron in *beat* mutants. Control (I and I'), *beat-IIa*$^{+2}$ mutant (J and J'), *beat-*

IIb^{Δ14} mutant (K and K'), *beat-IV*^{Δ65} mutant (L and L'). Brp-GFP puncta (green) of all

lamina neurons are visualized by combining the STaR method and R27G05-FLP. And

Lamina neuron (LN) is visualized with LexAop-myrtdTOM (magenta). Orange dotted

lines indicate the boundary between the proximal and distal region. (M−N') Brp-GFP of

lamina neurons when *beat-IIb* were knocked down on *beat-IIa* mutant background.

Control (M and M'), *beat-IIa*$^{+2}$ mutant + *beat-IIb* RNAi (J and J'). *beat-IIb* was knocked

down specifically in lamina neurons (Ay-gal4 & R27G05-FLP) and lamina neurons were

visualized with UAS-myr-RFP (magenta). Orange dotted lines indicate the boundary

between the proximal and distal region. (O) Quantification of Brp-GFP puncta in the

distal region in *beat* single mutants. Control (n=161 axons/3 brains), *beat-IIa*$^{+2}$ mutant

(n=191/3), *beat-IIb*^{Δ14} mutant (n=102/3), *beat-IV*^{Δ65} mutant (n=133/3). (P) Quantification

of Brp-GFP puncta in the distal region of lamina neurons upon simultaneous loss of *beat-*

IIa and *beat-IIb*. Control (n=144 axons/3 brains), *beat-IIa*$^{+2}$ mutant + *beat-IIb* RNAi

(n=129/3). Statistical processing of the number of Brp-GFP puncta in the distal region of

the lamina neuron was analyzed with the Mann-Whitney test.

1017

Figure S6. *side-IV* and *side-VII* double mutants didn't form ectopic synapses in the

Distal region of L1, L3, L4, and L5 neurons. (A−H') Brp-GFP (green) and axons

(magenta) of lamina neurons with different sub-type. (A−B') L1 neuron. Control (A and

A'), *side-IV^{Δ2}* / *side-VII$^{ΔE3-4-1}$* double mutant (B and B') expressing FLPase specifically in

the L1 neuron (R48A08-AD (II), R66A01-DBD (III) L1 split-GAL4 + UAS-FLP). (C−

D') L3 neuron. Control (C and C'), *side-IV^{Δ2}* / *side-VII$^{ΔE3-4-1}$* double mutant (D and D')

expressing FLPase specifically in L3 neuron (R64B03-AD (II), R14B07-DBD (III) L3

split-GAL4 + UAS-FLP). (E−F') L4 neuron. Control (E and E'), *side-IV^{Δ2}* / *side-VII$^{ΔE3-4-}$*

^{1} double mutant (F and F') expressing FLPase specifically in L4 neuron (R31C06-AD (II),

R34G07-DBD (III) L4 split-GAL4 + UAS-FLP). Orange dotted lines indicate the

boundary between the proximal and distal region. (G−H') L5 neuron. Control (G and G'),

side-IV^{Δ2} / *side-VII$^{ΔE3-4-1}$* double mutant (H and H') expressing FLPase specifically in L5

neuron (R64D07-AD (II), R37E10-DBD (III) L5 split-GAL4 + UAS-FLP). (I)

Quantification of Brp-GFP puncta in the proximal region of the L4 neuron. Control

(n=124 axons/3 brains), *side-IV^{Δ2}* / *side-VII$^{ΔE3-4-1}$* double mutant (n=123/3). (J)

Quantification of Brp-GFP puncta in the distal region of the L4 neuron. Control (n=191

axons/3 brains), *side-IV^{Δ2}* / *side-VII$^{ΔE3-4-1}$* double mutant (n=191/3). Statistical treatment

of the number of Brp-GFP puncta in the proximal region and distal region of the L4

neuron were analyzed with the unpaired t-test and Mann-Whitney test, respectively.

1037

1038 **Figure S7. L2 neurons express *side-IV* and *side-VII* from developmental to adult.** (A−

H') Endogenous expression of *side-IV* or *side-VII* in L2 neuron. Side-IV and Side-VII were visualized by combining UAS-FLP with L2-gal4 (R16H03-gal4) or L2-split-gal4 (R53G02-AD (II), R29G11-DBD (III)) against *side-IV*-FsF-GFP or *side-VII*-FsF-GFP. *side-IV* expression in L2 neuron at 24h APF (A and A'), 48h APF (B and B'), 72h APF (C and C'), 0d after eclosion (D and D'). Orange dotted lines indicate the boundary between the proximal and distal region. *side-VII* expression in L2 neuron at 24h APF (E and E'), 48h APF (F and F'), 72h APF (G and G'), 0d after eclosion (H and H'). Orange dotted lines indicate the boundary between the proximal and distal region.

Figure S8. Localization Beat-IIb and Beat-IV in the lamina neuron. (A−A″) Localization of Beat-IIb in the lamina neuron at the 72 hours APF. Localization of Beat-IIb was visualized by *beat-IIb*-MiMIC-GFP (green). The morphology of the L4 neuron was visualized with the L4 neuron-specific gal4 driver (GMR31C06-gal4) against UAS-myr-RFP (magenta). No Beat-IIb-GFP signal was observed in the axon of the L4 neuron (yellow arrowheads), although the Beat-IIb-GFP signal was confirmed in the cell body of the L4 neuron (yellow arrows). It suggests that *beat-IIb*-MiMIC-GFP is expressed in the L4 neuron, however, it may not be transported correctly to the axon. (B−B″) Localization of Beat-IV in the lamina neurons at the 72 hours APF. Localization of Beat-IV was visualized by *beat-IV*-MiMIC-GFP (green). A strong Beat-IV-GFP signal was observed on the axon in the proximal region of the L4 neuron. (C−D″) Analysis of the mutual localization of Side-VII and Beat-IV. Side-VII was visualized by combining *side-VII*-FsF-mCherry (magenta) and R27G05FLP. Beat-IV was visualized by *beat-IV*-MiMIC-

GFP (green). Orange dotted lines indicate the boundary between the proximal and distal

region. At the 60h APF (C−C"). Side-VII localized from proximal to distal (C") while

Beat-IV showed a subcellular localization in the proximal region (C"). At the 66h APF

(D−D"). Side-VII is strongly localized in the proximal region (D") with strong

colocalization with Beat-IV (D"). (E−H') Analysis of the Beat-IV localization dependency

to Side-VII in the lamina neurons. Control 48h APF (E and E'), *side-VII* RNAi 48h APF

(F and F'), Control 72h APF (G and G'), *side-VII* RNAi 72h APF (H and H'). *side-VII* was

knocked down in lamina neurons (Ay-gal4 & R27G05-FLP). Beat-IV similarly localized

in the proximal region, indicating the localization of Beat-IV is not Side-VII dependent.

1070

1071   **SUPPLEMENTALY TABLE**

1072

1073   **Table S1, The list of genotypes of flies used in this study.**

1074   **Table S2, The list of oligonucleotides used in this study.**

1075   **Table S3, The list of mutant alleles generated in this study.**

1076

1077   **RESOURCE AVAILABILITY**

1078

1079   **Lead Contact**

1080   Further information and requests for resources and reagents should be directed to and will

1081   be fulfilled by the Lead Contact, Takashi Suzuki (suzukit@bio.titech.ac.jp).

1082

1083 **Material Availability**

1084 All unique reagents generated in this study are available from the Lead Contact without

1085 restriction.

1086

1087 <u>**METHODS**</u>

1088

1089 **Key Resource Table**

1090

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
mouse anti 24B10	DSHB	Cat#24B10, RRID: AB_528161
rat anti cadN	DSHB	Cat#DN-Ex#8, RRID:AB_528121
mouse V5 Tag Monoclonal Antibody	ThermoFisher	Cat#R960-25, RRID:AB_2556564
Anti-HA High Affinity; Rat monoclonal antibody (clone 3F10)	Roche	Cat#11867423001, RRID:AB_390918
Anti-Myc Tag Antibody, clone 4A6	Millipore	Cat# 05-724MG, RRID:AB_568800
rabbit GFP Polyclonal Antibody	ThermoFisher	Cat#A-11122, RRID:AB_221569
Alexa Fluor 488 Goat anti-Mouse	ThermoFisher	Cat#A11029, RRID:AB_2534088
Alexa Fluor 568 Goat anti-Mouse	ThermoFisher	Cat#11031, RRID:AB_144696
Alexa Fluor 633 Goat anti-Mouse	ThermoFisher	Cat#A-21052, RRID:AB_2535719
Alexa Fluor 488 Goat anti-Rat	ThermoFisher	Cat#A-11006, RRID:AB_141373
Alexa Fluor 568 Goat anti-Rat	ThermoFisher	Cat#A-11077, RRID:AB_141874
Goat anti-Mouse IgG (H+L) Secondary Antibody, HRP	ThermoFisher	Cat#32430, RRID:AB_1185566
Biological Samples		
RE17794 (cDNA for *beat-IIa*)	Drosophila Genomics Resource Center	RRID:DGRC_8501
RE17755 (cDNA for *beat-IIb*)	Drosophila Genomics Resource Center	RRID:DGRC_1090247
GH14967 (cDNA for *side-VII*)	Drosophila Genomics Resource Center	RRID:DGRC_4843
Critical Commercial Assays		
KOD One® PCR Master Mix	TOYOBO	Cat#KMM-101
NEBuilder® HiFi DNA Assembly Master Mix	BioLabs	Cat#E2621X
FastGene Plasmid Mini Kit	NIPPON Genetics	Cat#FG-90502
FastGene Gel/PCR Extraction Kit	NIPPON Genetics	Cat#FG-91302
Pierce™ Protein A Magnetic Beads	ThermoFisher	Cat#88845
7.5% Mini-PROTEAN® TGX™ Precast Protein Gels	BIO-RAD	Cat#4561025
Trans-Blot Turbo Mini 0.2 µm PVDF Transfer Packs	BIO-RAD	Cat#1704156

Experimental Models: Organisms/Strains		
GMR-gal4	Bloomington Drosophila Stock Center	BDSC_8605, RRID:BDSC_8605
GMR-FsF-gal4	Chen et al., 2014	N/A
ort-gal4 (X)	Gao et al., 2008	N/A
ort-gal4 (III)	Bloomington Drosophila Stock Center	BDSC_56519, RRID:BDSC_56519
Ay-gal4	Bloomington Drosophila Stock Center	BDSC_4413, RRID:BDSC_4413
R48A08-p65.AD (L1-AD)	Bloomington Drosophila Stock Center	BDSC_70707, RRID:BDSC_70707
R66A01-GAL4.DBD (L1-DBD)	Bloomington Drosophila Stock Center	BDSC_68999, RRID:BDSC_68999
R48A08-p65.AD (L2-AD)	Bloomington Drosophila Stock Center	BDSC_70707, RRID:BDSC_70707
R66A01-GAL4.DBD (L2-DBD)	Bloomington Drosophila Stock Center	BDSC_68999, RRID:BDSC_68999
GMR16H03-Gal4 (L2-gal4)	Bloomington Drosophila Stock Center	BDSC_48744, RRID:BDSC_48744
R64B03-p65.AD (L3-AD)	Bloomington Drosophila Stock Center	BDSC_75895, RRID:BDSC_75895
R14B07-GAL4.DBD (L3-DBD)	Bloomington Drosophila Stock Center	BDSC_68940, RRID:BDSC_68940
R31C06-p65.AD (L4-AD)	Bloomington Drosophila Stock Center	BDSC_70646, RRID:BDSC_70646
R34G07-GAL4.DBD (L4-DBD)	Bloomington Drosophila Stock Center	BDSC_69029, RRID:BDSC_69029
GMR31C06-gal4 (L4-gal4)	Bloomington Drosophila Stock Center	BDSC_49883, RRID:BDSC_49883
R64B07-p65.AD (L5-AD)	Bloomington Drosophila Stock Center	BDSC_71106, RRID:BDSC_71106
R37E10-GAL4.DBD (L5-DBD)	Bloomington Drosophila Stock Center	BDSC_69000, RRID:BDSC_69000
side-IV[MI10049]-T2AGal4	Gift From Filipe Pinto-Teixeira	N/A
kirre^{sps}	Weavers et al., 2009	N/A
P{SAM.dCas9.GS05146}attP40 (Nrx-1)	Bloomington Drosophila Stock Center	BDSC_82741, RRID:BDSC_82741
P{SAM.dCas9.GS04053}attP40 (Nlg-1)	Bloomington Drosophila Stock Center	BDSC_81419, RRID:BDSC_81419
P{SAM.dCas9.GS04054}attP40 (Nlg-2)	Bloomington Drosophila Stock Center	BDSC_81420, RRID:BDSC_81420
P{SAM.dCas9.GS05201}attP40 (Nlg-3)	Bloomington Drosophila Stock Center	BDSC_82936, RRID:BDSC_82936
UAS-FLP (X)	Bloomington Drosophila Stock Center	BDSC_8208, RRID:BDSC_8208
UAS-FLP (II)	Bloomington Drosophila Stock Center	BDSC_4539, RRID:BDSC_4539
UAS-myr-RFP (III)	Bloomington Drosophila Stock Center	BDSC_7119, RRID:BDSC_7119
UAS-myr-RFP (II)	Bloomington Drosophila Stock Center	BDSC_7118, RRID:BDSC_7118
UAS-*luciferase*-RNAi (Control RNAi)	Bloomington Drosophila Stock Center	BDSC_35788, RRID:BDSC_35788

UAS-*beat-IIb*-RNAi	Vienna Drosophila Resource Center	VDRC_104935, RRID:FlyBase_FBst0479711
UAS-*beat-IIIc*-RNAi	Vienna Drosophila Resource Center	VDRC_27137, RRID:FlyBase_FBst0456776
UAS-*beat-IV*-RNAi	Vienna Drosophila Resource Center	VDRC_110257, RRID:FlyBase_FBst0481835
UAS-*side*-RNAi	Vienna Drosophila Resource Center	VDRC_1283, RRID:FlyBase_FBst0450699
UAS-*side-III*-RNAi (II)	Vienna Drosophila Resource Center	VDRC_101967, RRID:FlyBase_FBst0482037
UAS-*side-III*-RNAi (III)	Vienna Drosophila Resource Center	VDRC_22742, RRID:FlyBase_FBst0454672
UAS-*side-IV*-RNAi	Vienna Drosophila Resource Center	VDRC_16636, RRID:FlyBase_FBst0452368
UAS-*side-V*-RNAi (II)	Vienna Drosophila Resource Center	VDRC_44997, RRID:FlyBase_FBst0465867
UAS-*side-V*-RNAi (III)	Vienna Drosophila Resource Center	VDRC_44997, RRID:FlyBase_FBst0465867
UAS-*side-VII*-RNAi (II)	Vienna Drosophila Resource Center	VDRC_106353, RRID:FlyBase_FBst0472832
UAS-*side-VII*-RNAi (III)	Vienna Drosophila Resource Center	VDRC_106353, RRID:FlyBase_FBst0472832
UAS-*kirre*-RNAi	Vienna Drosophila Resource Center	VDRC_27227, RRID:FlyBase_FBst0456824
UAS-*dsyd-1*-RNAi	Bloomington Drosophila Stock Center	BDSC_32946, RRID:BDSC_32946
UAS-*liprin-α*-RNAi	Bloomington Drosophila Stock Center	BDSC_53868, RRID:BDSC_53868
P{EP}*side-VIII*	Bloomington Drosophila Stock Center	BDSC_17248, RRID:BDSC_17248
UAS-*beat-VI-HA*	FlyORF	RRID:FlyBase_FBst0502390
UAS-*shg (cadE)*	Bloomington Drosophila Stock Center	BDSC_58494, RRID:BDSC_58494
UAS-*hSDK2-HA*	Bloomington Drosophila Stock Center	BDSC_78487, RRID:BDSC_78487
UAS-*kirre-HA*	Bloomington Drosophila Stock Center	BDSC_92196, RRID:BDSC_92196
UAS-myc-*liprin-α*	Bloomington Drosophila Stock Center	BDSC_63809, RRID:BDSC_63809
UAS-*dsyd-1*	Owald et al., 2010	N/A
brp-FsF-GFP (III)	Bloomington Drosophila Stock Center	BDSC_55753, RRID:BDSC_55753
brp-FsF-V5-LexA, LexAop-tdTomato	Bloomington Drosophila Stock Center	BDSC_56142, RRID:BDSC_56142
20C11-FLPG5	Bloomington Drosophila Stock Center	BDSC_55766, RRID:BDSC_55766
R27G05-FLPG5	Bloomington Drosophila Stock Center	BDSC_55765, RRID:BDSC_55765
hs-FLP	Bloomington Drosophila Stock Center	BDSC_5258, RRID:BDSC_5258
Mi{PT-GFSTF.0}beat-IIb[MI03102-GFSTF.0]	Bloomington Drosophila Stock Center	BDSC_59406, RRID:BDSC_59406
Mi{PT-GFSTF.1}beat-IV[MI05715-GFSTF.1]	Bloomington Drosophila Stock Center	BDSC_66506, RRID:BDSC_66506

nos-Cas9 (attp40)	Bloomington Drosophila Stock Center	BDSC_78781, RRID:BDSC_78781
nos-Cas9 (attp2)	Bloomington Drosophila Stock Center	BDSC_78782, RRID:BDSC_78782
vas-int.Dm(ZH-2A); PBac{y[+]-attP-3B}(VK00037)	Bloomington Drosophila Stock Center	BDSC_24872, RRID:BDSC_24872
vas-int.Dm(ZH-2A); PBac{y[+]-attP-3B}(VK00033)	Bloomington Drosophila Stock Center	BDSC_24871, RRID:BDSC_24871
P{y[+t7.7] v[+t1.8]=TKO.GS04527}attP40 (TRiP-KO for *side-IV*)	Bloomington Drosophila Stock Center	BDSC_82064, RRID:BDSC_82064
P{y[+t7.7] v[+t1.8]=WKO.5-C1}attP40 (TRiP-KO for *beat-IV*)	Bloomington Drosophila Stock Center	BDSC_84159, RRID:BDSC_84159
beat-IIa^{+2}	This paper	N/A
beat-IIb$^{\Delta14}$	This paper	N/A
beat-IV$^{\Delta65}$	This paper	N/A
side-IV$^{\Delta4-1}$	This paper	N/A
side-IV$^{\Delta4-2}$	This paper	N/A
side-VII$^{\Delta E3-4-1}$	This paper	N/A
side-VII$^{\Delta E3-4-2}$	This paper	N/A
side-VII$^{\Delta E3-4-1}$, side-IV$^{\Delta2}$	This paper	N/A
UAS-*beat-IIa*	This paper	N/A
UAS-*beat-IIb*	This paper	N/A
UAS-*side*	This paper	N/A
UAS-*side-IV*	This paper	N/A
UAS-*side-IV*-V5	This paper	N/A
UAS-*side-IV*-GFP	This paper	N/A
UAS-*side-IV $^{\Delta Ig1-5}$*	This paper	N/A
UAS-*side-IV $^{\Delta Ig1}$*	This paper	N/A
UAS-*side-IV $^{\Delta FnIII}$*	This paper	N/A
UAS-*side-IV $^{\Delta Cyto}$* -V5	This paper	N/A
UAS-*side-IV $^{\Delta PDZb}$*	This paper	N/A
UAS-*side-IV$^{Ecto\Delta FnIII}::mCD8$-V5*	This paper	N/A
UAS-*side-IV$^{Ecto}::mCD8^{TM+Cyto}$-V5*	This paper	N/A
UAS-*side-VII*	This paper	N/A
UAS-*side-VII*-GFP	This paper	N/A
UAS-*kirre $^{\Delta Cyto}$*-HA	This paper	N/A
brp-FsF-GFP (II)	This paper	N/A
brp-FsF-mCherry (II)	This paper	N/A
side-IV-FsF-GFP	This paper	N/A
side-VII-FsF-GFP	This paper	N/A
side-VII-FsF-mCherry	This paper	N/A
dsyd-1-FsF-GFP	This paper	N/A
Oligonucleotides		
Please refer to Table S1.	This paper	N/A
Software and Algorithms		
Imaris 8.2.0	Bitplane	http://www.bitplane.com/imaris
NIS-elements VERSION 4.11	Nikon	https://www.nis-elements.cz/en
Prism	Graphpad	https://www.graphpad.com/ scientific-software/prism/

Fiji	Schindelin et al., 2012	https://fiji.sc/
Other		
All of the genotypes used in this paper are listed in Table S2.	This paper	N/A

1091

1092

Generation of transgenic flies

side-IV-FsF-GFP, *side-VII*-FsF-GFP (mCherry), *brp*-FsF-GFP (mCherry), and *dsyd-1*-

FsF-GFP knock-in allele were generated by CRISPR/Cas9 technology.[57] A knock-in

vector containing the homology arms, the flip-out cassette with GFP (FRT-stop-FRT-

GFP (mCherry), were generated as described previously.[58] The oligo DNAs used for

amplification of *side-IV*, *side-VII*, *brp*, and *dsyd-1* fragments and creating gDNA are

listed in Table S1. A gRNA vectors were created and cloned into pBFv-U6.2 vector and

were injected to eggs of nos-Cas9 (attp40) or nos-Cas9 (attp2) flies at WellGenetics or

Bestgene together with the knock-in vector. The precise integration of the knock-in vector

was verified by PCR and sequencing.

Mutant alleles of *beat-IIa*, *beat-IIb*, *beat-IV*, *side-IV*, and *side-VII* were generated

by CRISPR/Cas9 technology.[57] The gRNA vectors were created and cloned into pBFv-

U6.2 vector for *beat-IIa*, *beat-IIb*, and *side-VII*. TRiP-KO lines (BDSC 84159, BDSC

82064) were used to generate mutant alleles of *beat-IV* and *side-IV*. For generating *side-*

IV / *side-VII* double mutant, we introduced frameshift mutation *side-IV$^{\Delta 2}$* on *side-VII$^{\Delta E3-}$*

$^{4-1}$ allele by using TRiP-KO line (BDSC 82064). The oligo DNAs used for generating

mutant alleles and all the mutations of mutant alleles are shown in Table S1 and Table

S3, respectively.

UAS lines were generated φC31 integrase-mediated recombination.[59] The cDNAs

were cloned into pUAST-attB vector and injected to egg of vas-int.Dm(ZH-2A);

PBac{y[+]-attP-3B}(VK00037) or vas-int.Dm(ZH-2A); PBac{y[+]-attP-3B}(VK00033)

at Wellgenetics. The full length of cDNAs were amplified from cDNA clones (RE17794

for *beat-IIa*, RE17755 for *beat-IIb*, ECD01053 for *side-IV*, GH14967 for *side-VII*). The

available cDNA clones of *beat-IIb* (RE17755) and *side-IV* (ECD01053) didn't cover all

of the coding sequences. We amplified the lacking sequences from the fly genome and

conjected the full length of cDNAs. We conjected 2 fragments (F1 & F2) by overlap pcr

for generating insertions of Side-IV domain deletion variants, Side-IV-mCD8 chimeras,

and GFP tagged Side-IV or Side-VII. To generate the V5 or GFP tagged Side-IV and

Side-VII constructs, GS-linker were inserted before the V5 or GFP. Furthermore, the

sequence of PDZ binding motif of Side-IV or Side-VII were inserted in after the V5 or

GFP not to disrupt the function of PDZ binding motif. The primers used to generate the

UAS lines are shown in Table S1.

1125

Immunohistochemistry and imaging

The flies were maintained under standard conditions in Drosophila media at a temperature

of 25˚C. As described previously, the experimental procedures for brain dissection,

fixation, and immunostaining were performed as described previously.[60] Fly brains were

dissected in 0.1% PBT (PBS containing Triton X-100) and then fixed with 4%

paraformaldehyde at room temperature for 60 min. A primary antibody was added and

incubated overnight at 4 degrees on a shaker. The next day, the samples were washed

with 0.1 % PBT 3 times and incubated at room temperature for 60 min. The secondary

antibody was added and incubated at room temperature for 60 min. Samples were washed

with 0.1% PBT 3 times and incubated at room temperature for 2 hr. Finally, samples were

washed with PBS once and replaced by VECTASHIELD®. The following antibodies

were used for immunohistochemistry: mAb24B10 (1:50, DSHB), rat antibody to CadN

(Ex#8, 1:50, DSHB), mouse antibody to V5 (R960, 1:400, Invitrogen), rat antibody to

HA (3F10, 1:50, Roche). To enhance the signal intensity of Beat-IIb-MiMIC-GFP and

Beat-IV-MiMIC-GFP, rabbit antibody to GFP conjugated with Alexa488 (1:400, Life

technologies) was used. The secondary antibodies were Alexa488-, Alexa568- or

Alexa633-conjugated (1:400, Life Technologies). For image acquisition S2 cells, they

were fixed 4% paraformaldehyde at room temperature for 30 min and washed with PBS

for 30min.

1145

Image acquisition

All confocal images were taken by Nikon C2+ or A1 confocal microscope with NIS

elements (version 4.11). Each brain image was acquired between the depths of 30 and 50

μm from the surface of brain with a Z step size of 0.5 or 1μm. Images were acquired twice

and averaged. Images were processed with NIS elements (Nikon), Adobe Photoshop, and

Adobe illustrator.

1152

Cell culture transfection and Co-immunoprecipitation

We used a mixture of culture medium; 500mL of Schneider's Drosophila Medium (×1,

gibco), 5mL of Penicillin-Streptomycin Solution (×100, wako), and 50mL of fetal bovine

serum (Sigma). S2 cells were grown in 24 well plates (TrueLine) with 500μL of the

culture medium. For transfection, 1.5mg of UAS vectors and pActin5C-Gal4 were co-

transfected and incubated with 10mL of HilyMax (DOJINDO) and 30mL of Schneider's

Drosophila Medium. For co-immunoprecipitation, 3 wells of S2 cells were lysed in Lysis

Buffer (50 mM Tris at pH 6.8, 150 mM NaCl, 1% NP40, protease inhibitor cocktail

[Roche]), immunoprecipitated with 1 μg of GFP antibody (Life Technologies) and

Protein A beads (Pierce) according to the manufacture's procedure. SDS-PAGE was

performed on 7.5% SDS-PAGE gels (Bio-Rad). Immunoblotting was performed by V5

antibody (R960, 1:2500, Invitrogen) and HRP secondary antibody (REF32430, 1:1000,

Invitrogen). The images were taken by ImageQuant LAS 4000mini (cytiva) with

Chemiluminescent HRP Substrate (MILLIPORE).

pUAS plasmids were generated by cloning cDNAs into pUAST-GFP or mCherry

vectors. To amplify cDNA or domain deletion variants of *side-IV*, we used the same

primers as generate transgenic flies. Genome extractions were performed to amplify the

cDNA of *kirre* (UAS-*kirre.C-HA,* BDSC 92196), *cadE* (UASp-*shg.R*, BDSC58494),

dsyd-1 (UAS-*dsyd-1*),[61] and *liprin-a* (UAS-MYC-*liprin-a.H*, BDSC63809). All of

primers used to generate the pUAS plasmids are shown in Table S1.

1173

QUANTIFICATION AND STATISTICAL ANALYSIS

All of the names of samples were randomized before the quantification. The number of

Brp puncta of lamina neurons and photoreceptor R7 was counted manually with IMARIS

(version 8.2.0) as described previously.[62] The Brp puncta on a 3D image were identified

with the spot function of IMARIS (diameter was set to 0.35μm). For the separation of the

region of proximal region and distal region of lamina neurons, we cropped each stack of

the proximal and distal ROI with NIS elements and a pen tablet (Cintiq 13HD, Wacom).

We first drew the lines on the top and bottom of the lamina cartridge. Next, we drew a

line on the midpoint of the lamina cartridge referring to the top and bottom lines. The

accept region function of NIS-elements was then used to select the regions above and

below the midpoint lines, which were then cropped as the proximal and distal regions,

respectively. The number of Dsyd-1 puncta in the proximal region and distal region at

72h APF were counted as the same method. The rate of overshoot axon was calculated

manually with NIS elements, by counting the number of photoreceptors that overshoot

beyond M6 layer and dividing the total number of axons.

In Figure 1M, the Side-IV cluster with Beat-IIb cluster was calculated manually

with IMARIS, by counting the number of Side-IV puncta that co-localized with Beat-IIb

puncta and dividing the total number of Side-IV puncta in the M1–M3 layer. The Side-

IV & Beat-IIb cluster with Brp was calculated, by counting the number of Side-IV puncta

co-localized with Beat-IIb and Brp puncta and dividing the total number of Side-IV

puncta co-localized with Beat-IIb puncta.

In Figures 2H, I, and S2K the rate of Side-IV or Beat-IIb clustered axon were

calculated manually with IMARIS, by the counting the number of axons with Side-IV or

Beat-IIb clustered signal in the M1–M3 layer and dividing the total number of axons.

In Figures 3R, T, and U–V, the cell-cell contact accumulation ratio was calculated

manually with NIS-elements as described previously.[43] Calculate the average intensity of

GFP or mCherry signal on cell-cell border by drawing the line ROI and dividing the

average intensity on the membrane, which did not contact other cells.

In Figures 4I and L, Pearson's Coefficient was calculated with the JACoP plugin

on Fiji.[63,64]

In Figures 6H and M, the rate of Side-IV or Side-VII positive axon was calculated

manually with NIS elements, by counting the number of the axon that has Side-IV or

Side-VII signal to the proximal region and dividing the total number of axons.

In Figure 7B, the average intensity along the Proximal to Distal regions of Side-

IV and Kirre in the L2 neuron was calculated by drawing a linear ROI using NIS-elements

and calculating the average intensity.

In Figure 7O, the rate of Dsyd-1 positive axon (Distal) was calculated manually

with NIS-elements, by counting the number of the axon that has Dsyd-1 signal on the

distal region and dividing the total number of the axon at 48h APF.

In Figure S2Q, the number of cells per cell aggregation was calculated manually

with NIS-elements, by counting the total number of cells on each cell aggregation. The

cell cluster with contains both GFP and mCherry expressing cells was identified as cell

aggregation.

In Figure S4D, the rate of Kirre-GFP cell cluster with and without mCherry

positive cell was calculated manually with NIS-elelments, by counting the rate of Kirre-

GFP cell cluster which had more than one mCherry positive cell. The cell cluster with

more than two GFP expressing cells was identified as Kirre-GFP cell cluster.

1221

1222   **Statistical analysis**

1223   All of the statistical analysis are described in the figure legends. Statistical significance.

1224   n.s. $P > 0.05$, * $P \leq 0.05$, ** $P \leq 0.01$, *** $P \leq 0.001$, **** $P \leq 0.0001$.

1225

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

Figure S1

Figure S2

Figure S3

Figure S4

Figure S5

Figure S6

side-IV expresstion in L2 neuron

side-VII expresstion in L2 neuron

Figure S7

Figure S8