A Pvr–AP-1–Mmp1 signaling pathway is activated in astrocytes upon

traumatic brain injury

Tingting Li[1], Wenwen Shi[1], Margaret S. Ho[2,*], Yong Q. Zhang[1,*]

[1]Key Laboratory of Molecular and Developmental Biology, Institute of Genetics and

Developmental Biology, University of Chinese Academy of Sciences, Chinese

Academy of Sciences, Beijing 100101, China.

[2] School of Life Science and Technology, ShanghaiTech University, Shanghai,

201210, China

*Correspondence: margareth@shanghaitech.edu.cn, yqzhang@genetics.ac.cn (lead

contact)

Main Points:

• The study provided RNA-seq data of astrocytes following traumatic brain injury

(TBI)

• Genes involved in endocytic trafficking are upregulated in astrocytes after TBI

• A new Pvr–AP-1–Mmp1 pathway is activated in astrocytes following TBI.

• Inhibition of endocytosis in astrocytes upregulates the Pvr–AP-1–Mmp1 signaling.

Abstract

Traumatic brain injury (TBI) caused by external mechanical forces is a major health burden worldwide, but the underlying mechanism in glia remains largely unclear. We report herein that *Drosophila* adults exhibit a defective blood-brain-barrier (BBB), elevated innate immune responses, and hypertrophy of astrocytes upon consecutive strikes with a high-impact trauma device. RNA sequencing (RNA-seq) analysis of these astrocytes revealed upregulated expression of genes encoding PDGF and VEGF receptor-related (Pvr, a receptor tyrosine kinase (RTK)), adaptor protein complex 1 (AP-1, a transcription factor complex of the c-Jun N-terminal Kinase (JNK) pathway) composed of Jun-related antigen (Jra) and kayak (kay), and matrix metalloproteinase 1 (Mmp1) following TBI. Interestingly, Pvr is both required and sufficient for AP-1 and Mmp1 upregulation, while knockdown of AP-1 expression in the background of Pvr overexpression in astrocytes rescued Mmp1 upregulation upon TBI, indicating that Pvr acts as the upstream receptor for the downstream AP-1–Mmp1 transduction. Moreover, dynamin-associated endocytosis was found to be an important regulatory step in downregulating Pvr signaling. Our results identify a new Pvr–AP-1–Mmp1 signaling pathway in astrocytes in response to TBI, providing potential targets for developing new therapeutic strategies of TBI.

Keywords: TBI, astrocytes, Pvr, RNA-seq, *Drosophila*

Introduction

Traumatic brain injury (TBI) is one of the leading causes for adult disability due to limited neurological recovery (Joy et al., 2019). About 43% of people discharged after hospitalization for acute TBI develop long-term locomotor disability (Ma et al., 2014). TBI triggers primary neural injuries via rapid physical forces that damage axons, glia and blood vessels (Chodobski et al., 2011; Daneshvar et al., 2015), followed by diverse secondary damage such as Wallerian axon degeneration (Büki and Povlishock, 2006) and blood-brain barrier (BBB) dysfunction (Chodobski et al., 2011) in the central nervous system (CNS). The outcome of TBI is affected by various factors including age and diet (Byrns et al., 2021; Katzenberger et al., 2016). In addition, glial cells are critical mediators in response to TBI in the mammalian brain (Mira et al., 2021). During TBI, astrocytes become active, a process characterized by swelling, scar tissue formation, and engulfment of neuronal debris (Liddelow et al., 2017; Sofroniew and Vinters, 2010), while microglia rapidly transform to their active state to phagocytose cell debris (Davalos et al., 2005).

Drosophila melanogaster is an ideal model organism for discovering new gene networks and investigating the molecular and cellular responses to CNS injury (Ayaz et al., 2008; Bier, 2005; Hoopfer et al., 2006; MacDonald et al., 2006). *Drosophila* astrocyte-like glia (henceforth astrocytes), the conserved counterparts for mammalian astrocytes, and ensheathing glia (ENG) constitute the major glial subclasses in the CNS (Doherty et al., 2009; Stork et al., 2014). A previous study revealed that *Drosophila* ENG perform the phagocytic function by the engulfment receptor Draper which interacts with downstream signaling components, including ced-6, Shark, Src42a, c-Jun N-terminal Kinase (JNK), and Mmp1 in the glial engulfment process in response to axotomy (Awasaki et al., 2006; Doherty et al., 2014; Lu et al., 2014; Lu et al., 2017; Purice et al., 2017). Although ENG function upon axotomy has been well characterized, the response of other types of glia such as astrocytes to neural injury remains unclear.

In mammals, signaling by platelet-derived growth factor (PDGF) directs cell migration, alters cell shape, and plays a role in wound healing (Hoch and Soriano, 2003; Sil et al., 2018). Meanwhile, vascular endothelial growth factor (VEGF) signaling is critical for cell proliferation and survival, and vascular endothelial cell mitogenesis and permeability (Apte et al., 2019; Sakurai et al., 2005). Pvr, short for PDGF- and VEGF receptor-related, is the sole *Drosophila* ortholog of the PDGF and VEGF receptor families (Cho et al., 2002). Pvr interacts with different signaling molecules in a context-dependent manner. For example, Pvr regulates thorax closure during metamorphosis via the c-Jun N-terminal Kinase (JNK) signaling pathway (Ishimaru et al., 2004), but the Pvr-mediated wound-induced epidermal cell migration does not depend on JNK (Wu et al., 2009). The JNK Activator Protein-1 (AP-1) signaling pathway is involved in different cellular processes, including cell migration (Pastor-Pareja et al., 2004), organ size control (Willsey et al., 2016), and cell death (Moreno et al., 2002). The JNK effector AP-1 complex, composed of heterodimeric Jun/Jun-related antigen (Jra) and Fos/kayak (kay), promotes transcription of genes including matrix metalloproteinase 1 (Mmp1) that functions in tissue remodeling, regulation of inflammatory processes, and metastasis of cancer cells (Egeblad and Werb, 2002; Page-McCaw et al., 2007; Parks et al., 2004).

Previous studies have implicated a role for ensheathing glia, Draper pathway, and the innate immune pathway in response to TBI (Doherty et al., 2009; Lu et al., 2017; Purice et al., 2017; Katzenberger et al., 2013). However, whether astrocytes also play a role in TBI response remains elusive. Herein, using a *Drosophila* TBI model induced with a high-impact trauma (HIT) device (Katzenberger et al., 2013), we found that TBI caused multiple defects including upregulated expression of innate immune genes and astrocyte hypertrophy. To uncover the molecular pathways activated by TBI, we performed RNA sequencing (RNA-seq) analysis of astrocytes sorted by fluorescence-activated cell sorting (FACS) to profile TBI-induced genes. Interestingly, we showed that the expression of Pvr, AP-1, and Mmp1 was upregulated

following TBI. Knockdown of Pvr or AP-1 expression in astrocytes rescued TBI-induced Mmp1 upregulation, while Pvr overexpression promoted Mmp1 expression via AP-1 upon TBI. Our findings identify a new Pvr–AP-1–Mmp1 signaling pathway that acts predominantly in astrocytes in response to TBI, and shed light on the mechanisms of astrocyte responses following TBI.

Materials and Methods:

Drosophila stocks and husbandry

Flies were maintained on standard cornmeal medium at 25°C otherwise indicated. The following *Drosophila* strains were used: w^{1118}, *UAS-mCD8::GFP*, *R86E01-Gal4* (#45914; Bloomington), *R54F03-Gal4* (#39157; Bloomington), *UAS-Redstinger* (#8547; Bloomington), *UAS-Jra* (#7216; Bloomington), *UAS-kay* (#7213; Bloomington), *UAS-Pvr* (#58429; Bloomington), *UAS-nLacZ* (#3955, Bloomington), *PCNA-GFP* (#25749; Bloomington) (Thacker et al., 2003), and *TRE-dsRed* (Chatterjee and Bohmann, 2012). *UAS-draper-RNAi*, *UAS-Pvr-RNAi*, *UAS-Jra-RNAi*, *UAS-kay-RNAi*, *UAS-Arpc2-RNAi*, *UAS-cpa-RNAi*, *UAS-cpb-RNAi*, *UAS-Stam –RNAi*, *UAS-Vps24-RNAi*, and *UAS-Chmp1-RNAi* flies were obtained from Tsinghua Fly Center (https://thfc.zzbd.org/en). *y w UAS-CD8::GFP hs-FLP; tubP-Gal80 FRT-40A/CyO; alrm-Gal4 UAS-CD8::GFP UAS-Dcr2/TM6 Tb Hu* (alrm-Gal4 is a specific driver in astrocytes (Doherty et al., 2009), a gift from Oren Schuldiner) and *FRT-40A* (a gift from Liqun Luo) were used for MARCM analysis of astrocytes. *UAS-shibire^{ts1}* (Pfeiffer et al., 2012) was provided by Y. Zhong at Tsinghua University.

Traumatic brain injury (TBI) Assay

Construction and use of the HIT device were described previously (Katzenberger et al., 2013). Deflection of the spring to 90° was the standard angle for strike. Mortality index at 24 h (MI_{24}) were determined by analyzing at least 120 flies (20 flies for each test).

Dye penetration assay

Dye penetration experiments were performed as previously described (Bainton et al., 2005). 10 kD Texas red-conjugated dextran (Molecular Probes) at 2.5 mM was injected into the soft tissue between two abdominal segments of the exoskeleton of flies with and without TBI. Flies were allowed to recover in fresh food tubes and photographed 2 h later on a Zeiss microscope to detect the integrity of brain-eye barrier (BEB). To detect the integrity of BBB, fly brains were dissected at 2 h after injection and examined under an Olympus FluoView FV1000 microscope.

MARCM analysis

Mosaic analysis with a repressible cell marker (MARCM) analysis in astrocytes was carried out largely following previous publications (Jia et al., 2019; Lee and Luo, 1999). Specifically, newly laid eggs were collected for a 4–5 h interval and cultured at 18°C. Clones were induced by heat shock at 24 h after larval hatching in a 37°C water bath for 15 min and flies were raised at 25°C after heat shock until immunohistochemistry analysis of adult brain.

Immunohistochemistry analysis and confocal microscopy

Brain tissues were dissected out from adult after specific treatments. Fixation and staining of adult brains were performed as previously described (Jia et al., 2019). Brains were mounted on slides in Vectashield mounting media (Vector Labs). Primary antibodies were used at the following dilutions: chicken anti-GFP (ab13970, Abcam) at 1:1000, mouse anti-Repo (8D12, DSHB) at 1:100, mouse anti-Elav (9F8A9, DSHB) at 1:100, mouse anti-Mmp1 (3A6B4, 3B8D12, and 5H7B11, DSHB) each at 1:20 for all three antibodies, rabbit anti-Pvr (McDonald et al., 2003) at 1:1000, rabbit anti-Dcp-1 (Asp216, Cell Signaling Technology (CST)) at 1:100, rabbit anti-pH3 (06-570, Millipore) at 1:1000, and mouse anti-β-gal (40-1a, DSHB) at 1:100. All secondary antibodies (Alexa Fluor 488, 568 or 633 labeled; Molecular Probes) were used at a dilution of 1:1000. The nuclei dye To-Pro-3 Iodide (T3605, ThermoFisher) was also used at a dilution of 1:1000. The cell-permeant Mitotracker (M22426,

ThermoFisher) was used at a concentration of 100 nM. For soma diameter analysis,

we measured the average of long and short axes of astrocytes. All images were

collected using an Olympus FV1000 laser scanning confocal microscope and

processed with ImageJ. Serial images of 25–30 µm (one image per 1 µm slice) were

projected for presentation and statistical analysis.

Astrocyte preparation for RNA-seq analysis

Flies were collected at 3 days after eclosion and treated with TBI. Brains from

adult flies (*R86E01-Gal4/UAS-Redstinger*) with no injury, 4 h after injury and 24 h

after injury were dissected out in freshly prepared adult hemolymph saline (AHS)

buffer, as previously described (Yang et al., 2016). Briefly, a total of ∼50 brains for

each group was immediately transferred to an Eppendorf tube (1.5 mL) containing

200 µL AHS with Pronase (1 mg/mL; P5147, SigmaAldrich) and Dispase (1 mg/mL;

LS02104, Worthington Biochemical Corporation). After tissue digestion for 30 min at

25°C, the enzyme medium was removed, followed by two brief, gentle washes with

fresh AHS containing 2% PBS. Brain samples were then triturated by gently pipetting

in 1 mL AHS through four fire-polished Pasteur pipettes with descending pore sizes

(400 µm, 300 µm, 200 µm, and 100 µm) for a total of ∼20 min. Next, the dissociated

cells were centrifuged for 10 min at $100 \times g$ and the AHS was replaced with the same

buffer to minimize any RNA content released from damaged cells during the

trituration process. Finally, dissociated tissues were filtered through 45 µm filter and

then put on ice. FACS sorting was performed according to the following gating

procedure with 70 µm nozzle by BD FACS Aria: (1) an initial SSC-A/FSC-A gate to

minimize debris; (2) a SSC-A/pe-Texa Ted-A gate to choose only red fluorescence

positive cells. At this step, 1600 cells were sorted directly into ice-cold cell lysis

buffer (oligo-dT$_{30}$VN primer, ribonuclease inhibitor, dNTP mix) and stored at −80°C

until all samples were ready for RNA extraction and cDNA amplification.

cDNA library preparation

A total of 5 biological replicates were collected for each condition (no injury, 4 h

after injury, and 24 h after injury). Cell samples were sent to the Anoroad Company

for RNA extraction. An Oligo-dT primer was introduced to the reverse transcription

reaction for first-strand cDNA synthesis, followed by PCR amplification to enrich the

cDNA and magbeads purification step was used to purify the products. Briefly,

Smart-Seq2 method was used to amplify cDNA. Concentration and integrity of cDNA

was assessed using Qubit 3.0 Flurometer (Life Technologies) and Agilent 2100

Bioanalyzer (Agilent Technologies) to ensure the cDNA length was around 1-2 kbp.

After quality control, 40 ng cDNA was used to be fragmented at 350 bp by

Bioruptor® Sonication System (Diagenode Inc.). To construct Illumina library, end

repair, 3' ends A-tailing, adaptor ligation, PCR amplification and library validation

were performed. PerkinElmer LabChip® GX Touch and Step OnePlus™ Real-Time

PCR System were used for library quality inspection. Qualified libraries were loaded

on Illumina Hiseq platform for PE150 sequencing.

RNA-seq and data analysis.

A total of 539 million reads (average 35.9 million reads per sample) was obtained.

The fastq data were first processed with Trimmomatic (Bolger et al., 2014) to remove

Illumina adaptor. The remaining data was aligned against the UCSC *D. melanogaster*

genome (dm6) using Hisat2 (Kim et al., 2015; Kim et al., 2019; Pertea et al., 2016)

with the annotation file of the same version. On average, 92% of the initial reads were

mapped to dm6. HTSeq was used to calculate the raw read counts of genes for each

sample. DESeq2 (Love et al., 2014) was used to normalize the data. Transcripts were

considered as significantly differentially expressed if their adjusted *p* values were less

than 0.01 ($p < 0.01$) by the Benjamini-Hochberg method. Following normalization of

the data, DESeq2 was used to perform pairwise comparisons between the control and

experimental groups using parametric tests. Genes with absolute value of logarithmic

fold change ($\log_2$FC) > 1.0 were qualified for the final set of DEGs. Volcano plots and

heatmap were constructed for visualization of the resulting expression intensity data

following the identification of DEGs and RNA-Seq data normalization by the

218   adjusted *p* value calculations.

Quantitative reverse transcription-PCR analysis.

220   RT-qPCR was performed largely based on previous protocols (Zhao et al., 2020).

Flies were treated with the same conditions as the RNA-seq experiment. RNA for

antimicrobial peptides (AMPs) analysis was extracted from whole fly brains by TriZol.

RNA for confirming RNA-seq findings was isolated from more than 10000 astrocytes

sorted by FACS using the PicoPure RNA isolation kit (KIT0204, Life Technologies).

cDNA was generated with SuperScript III First-Strand Synthesis SuperMix

(11752250, Invitrogen). For each gene, at least three pairs of qPCR primers were

screened by standard curve analysis, and the best primer pairs were chosen based on

their efficiency and specificity. Real-time qPCR was carried out by KAPA Library

Quantification Kit (KR0389, KAPA Biosystems). Primer sequences are provided in

Table S1.

Thermogenetic inhibition of endocytosis

232   For inhibition of endocytosis in astrocytes using *UAS-Shi^{ts}*, crosses were reared

and treated by HIT device at 23°C to avoid unintended inactivation. For

thermogenetic inactivation, flies after TBI were undergone intermittent temperature

treatment at 32°C (32°C for 30 min, and then back to 23°C for 30 min, for four

rounds, four hours in total) until dissection.

Statistical analysis

238   All statistical comparisons were performed using GraphPad Prism software, with

significance indicated by ∗∗∗ ($p \le 0.001$), ∗∗ ($p \le 0.01$), ∗ ($p \le 0.05$), and ns ($p > 0.05$).

One-way ANOVA was used to compare multiple group means. Student's t-tests were

used for statistical comparisons between two groups.

Results

TBI is induced in *Drosophila* adults by a high-impact trauma (HIT) device

245   To analyze how brains respond to TBI, we applied consecutive strikes to adult

flies using a HIT device to induce brain injury as described previously (Katzenberger et al., 2015). To quantify the degree of injury, we analyzed the mortality index at 24 h after TBI (MI_{24}), defined as the percentage of dead over the total flies tested. Consistent with a previous report (Katzenberger et al., 2013), MI_{24} of 3-day-old adult flies was $20 \pm 3.18\%$ after 4 HIT strikes (Figure 1A). MI_{24} after 6 strikes was $71.88 \pm 7.44\%$, suggesting that increasing the number of strikes results in a higher MI_{24} rate (Figure 1A). Given that TBI has been shown to disrupt the formation of the blood-brain barrier (BBB) in mammals and flies and blood-eye barrier (BEB) in flies (Rodríguez-Baeza et al., 2003; Sangiorgi et al., 2013; Katzenberger et al., 2015; Saikumar et al., 2020), we examined BBB/BEB integrity 2 h after injecting tetramethylrhodamine-conjugated dextran (molecular weight = 10,000 Da) into adult fly abdominal segments (Bainton et al., 2005). The dye crossed the barrier and penetrated the retina and brain, indicating that both BBB and BEB were disrupted after HIT strikes (Figure 1B–G). The percentage of flies exhibiting dextran-permeable BBB increased as the number of strikes increased from 0 to 6, with six strikes resulting in severe BBB disruption (Figure 1G) but maintaining a moderate viability providing enough samples for phenotypic analysis. Based on these results, six HIT strikes administered at 5 min intervals were applied to induce TBI throughout the present study.

In addition to disruption of BBB/BEB integrity, we also analyzed TBI-induced secondary damage such as local and systemic immune responses. Given that Toll and immune deficiency (Imd) are the two crucial innate immune response pathways in *Drosophila* (Valanne et al., 2011), and their common downstream genes encode antimicrobial peptides (AMPs) that protect flies from TBI injury (Alphen et al., 2022), we analyzed the expression of AMPs in adult fly brains at different time points after TBI. As shown by the real-time reverse transcriptase-polymerase chain reaction (RT-qPCR) results, expression of the AMP genes *AttC*, *DiptB* and *Mtk* was significantly upregulated at 1 h and reached a peak at 4 h after injury (Figure 1H),

consistent with previous reports (Barekat et al., 2016; Katzenberger et al., 2013).

Taken together, these results suggest that HIT strikes induce TBI reflected by

BBB/BEB disruption and elevated expression of innate immune response genes.

TBI induces neuronal but not glial apoptosis in adult fly brains

Given that neurons and glia have been shown to undergo cell death during both

primary and secondary injury responses upon TBI in humans (Ng and Lee, 2019), we

performed immunostaining with an antibody against Death caspase-1 (Dcp-1), the

homolog of human caspase-3, to detect apoptotic cells in our TBI model. Interestingly,

the number of Dcp-1$^+$ puncta was significantly increased in adult fly brains at 24 h

after TBI (Figures 1I and J), with a 2.47-fold increase compared to the number in

uninjured controls (Figure 1K). Furthermore, we found that almost all Dcp-1$^+$ puncta

were co-localized with Elav rather than Repo, suggesting that neurons but not glia are

apoptotic (Figures 1L−M"). These results suggest that, as in mammals, TBI also

induces neuronal apoptosis in fruit flies.

In addition, we analyzed cell proliferation following TBI using an antibody

against phospho-histone H3 (pH3), a marker for the M stage of the cell cycle, and fly

lines expressing the proliferating cell nuclear antigen (PCNA) fused with GFP

(PCNA-GFP), a reporter for the G1-S transition of mitosis (Thacker et al., 2003). As a

control, 3rd instar larval brains exhibited numerous GFP$^+$ and pH3$^+$ cells, suggesting

that larval brain cells proliferate vigorously (Figures S1A−A"). Interestingly,

essentially no GFP$^+$ or pH3$^+$ cells were detected in either uninjured or TBI-treated

adult fly brains at various time points (Figures S1B−I). These results indicate that

adult fly brain cells do not divide in response to TBI, whereas neurons and glia in

adult vertebrate brains proliferate after nervous system injury, such as stab injury and

TBI (Yu and He, 2019).

TBI leads to astrocyte activation

In addition to the TBI-induced neuronal apoptosis mentioned above, glial cells also become reactive after nervous system injury (Burda and Sofroniew, 2014). It has been shown that ENG phagocytose neuronal debris upon axotomy (Doherty et al., 2009), but whether astrocytes respond to TBI remains unknown. To explore the potential role of astrocytes in the TBI response, *R86E01-GAL4* (Kremer et al., 2017), an astrocyte-specific driver, was used to express *UAS-mCD8::GFP* to label astrocyte plasma membrane. In uninjured fly brains, GFP intensities in the cell bodies and processes of astrocytes were widely distributed throughout the brain (Figures 2A and 2D). While no significant change in GFP intensity was detected in brains at 4 h after TBI (Figures 2B and 2D), the GFP intensity was significantly increased and the cell bodies of astrocytes were significantly enlarged at 24 h after TBI in the central brain region, especially in the antennal lobes (ALs) (Figures 2C and 2D). Of note, these results were not affected by the presence of one or two UAS transgenes as the efficiency of *R86E01-GAL4* in driving the expression of either was comparable as shown by the similar GFP levels (Figures 2A–C, I, and J).

In mammals, the hallmarks of astrocyte activation are hypertrophy of astrocytes and upregulated level of the glial fibrillary acidic protein (GFAP) (Pekny et al., 2004). Given that there's no GFAP homolog in *Drosophila (*Doherty et al., 2009)*, we examined morphological changes in astrocytes at single-cell resolution following TBI by mosaic analysis with a repressible cell marker (MARCM) on adult fly brains at 3– 4 days after eclosion. The cell bodies of these astrocytes were located at the interface between neuropil and cortex regions, or in neuropils, and their cellular processes infiltrated into neuropils (Figure 2E). Interestingly, astrocytes in injured brains exhibited bright and distinct GFP$^+$ accumulations associated with their cellular processes at 24 h after TBI (Figures 2F and 2G). Quantitatively, the number of these accumulations per square nanometer in the cellular processes was significantly increased at 24 h after TBI; the increase in the number of accumulations might be

responsible for the increase in GFP intensity triggered by TBI (Figure 2G). Importantly, the diameter of astrocytes cell bodies was also increased at 24 h after injury (Figure 2H). Taken together, these results show that astrocytes become hypertrophic and reactive in adult brains at 24 h after TBI.

Axotomy induces expression of injury-related proteins such as extracellular proteases the matrix metalloprotease (MMP) (Purice et al., 2017). In addition to the morphological changes in astrocytes, we observed upregulated expression of Mmp1 in TBI-treated brains (Figures 2I–J'', 2M–N'', and 2Q). These Mmp1$^+$ signals were colocalized with membrane-bound *mCD8::GFP* driven by the astrocytes-specific driver *R86E01-GAL4* or ENG-specific driver *R56F03-GAL4* (Figures 2K, 2L, 2O, and 2P), suggesting that Mmp1 might be produced from both types of glia after TBI.

Transcriptional profiling of astrocytes in adult flies reveals elevated expression of injury-related genes upon TBI

To further investigate how astrocytes respond to TBI, we analyzed transcriptional changes in astrocytes by RNA-seq (Figure 3A). Due to the tiny soma and morphological complexity, the analysis of a purified population of glial cells has been difficult and technically challenging. We have optimized a protocol for purifying astrocytes from adult Drosophila of *R86E01-Gal4>UAS-Redstinger* by fluorescence-activated cell sorting (FACS) (see details in Materials and Methods). As FACS removes most cell masses and RFP-negative cells, the remaining RFP$^+$ cells are mostly viable, as shown by MitoTracker staining, and purity was 92.96% (RFP$^+$ and MitoTracker$^+$ cells/total cells) in uninjured brains (Figure S2 A-C'). Further analysis of the RNA-seq data showed that the astrocyte markers *Eaat1* (110359 reads) and *alrm* (186413 reads) were highly expressed, while *nSyb*, a neuronal marker, was hardly detected (125 reads) at 4 h after TBI, verifying the purity of FACS-sorted astrocytes. A principal component analysis (PCA) revealed that the variance between different groups (no injury vs. 4 h and 24 h after injury) was greater than the variance

within groups (Figure S2D). From analysis on five biological replicates for each time point (no injury, 4 h, and 24 h after injury), we obtained 539 million reads (an average of 36 million reads per sample) and aligned the RNA-seq reads to the *Drosophila* genome. We found that $93.67 \pm 0.89\%$ of reads mapped to the *Drosophila* genome (dm6 version). Using a cutoff of $\geq$ 2-fold differential expression (adj. *p* value <0.01, Benjamini-Hochberg procedure), we identified 522 and 357 genes with altered expression at 4 h (416 upregulated and 106 downregulated) and 24 h (229 upregulated and 128 downregulated) after injury, respectively (Figures 3B–D; Table S2).

Gene Ontology (GO) analysis of these upregulated genes revealed that a significant proportion were associated with endosomal transport, actin cytoskeleton, and wound healing (Figure S3). Furthermore, we detected upregulated expression of components functioning in the innate immune response after TBI. For instance, expression of the components Relish and Dif of the Toll and Imd pathways was upregulated in astrocytes after TBI (Figure 3D), in agreement with a previous finding that Relish is necessary for TBI-induced secondary injuries (Swanson et al., 2020). In addition, expression of injury-related genes including *chic*, *Jra*, *kay, rl*, *Cad96Ca*, *Idgf4*, *Stam*, *draper*, *Pvr*, *spict*, *Mmp1* and *NijA*, and receptor genes including *Pvr*, *draper*, *AdoR*, *PGRP-SA*, and *NimA* were upregulated in astrocytes at 4 h or 24 h after injury (Table S3). A set of endocytic trafficking genes was also identified, including *Arpc1*, *Arpc2*, *cpa*, *cpb*, *Stam*, and *Chmp1*. Subsequent RT-qPCR experiments confirmed the upregulation of all six genes tested at 4 h and/or 24 h after TBI (Figure 3E), supporting the robustness of the RNA-seq analysis. As a control, the astrocytes-specific *alrm* gene showed no upregulation upon TBI by RT-qPCR and RNA-seq. Taken together, these findings validate the approach of RNA-seq transcription profiling in detecting genes functionally related to injury responses in astrocytes following TBI.

Pvr and AP-1 are required for the upregulation of Mmp1 after TBI

Among the candidates exhibiting upregulated expression in astrocytes upon TBI, three receptors (Pvr, AdoR and Draper), all acting upstream of and activating the JNK pathway (Ishimaru et al., 2004; Macdonald et al., 2013; Poernbacher and Vincent, 2018), emerged as new factors with uncharacterized functions in astrocytes (Figure 3D). The *Drosophila* RNA interference (RNAi) Screening Center Integrative Ortholog Prediction Tool (DIOPT) identified orthologs of these three receptors, subunits of transcriptional factor AP-1 (composed of Jra and kay) of the JNK pathway, Mmp1, and endocytic trafficking genes in mouse and human (Table 1) (Hu et al., 2011). These proteins mediate essential injury responses in vertebrates (Borea et al., 2016; Motz and Coukos, 2013; Shi et al., 2021). To examine a potential role of the receptors and their downstream signaling components in astrocytes after TBI, we quantified the immunostaining intensity of these proteins from projected images of the whole brain (including 20-30 slices with 1 µm per slice). As our RNA-seq analysis revealed elevated expression of JNK-related injury genes such as *AP-1* and *Mmp1* (Figure 3D and Table 1), we analyzed the expression of Mmp1, a transcriptional target of AP-1 (Purice et al., 2017), as a readout for the activity of JNK signaling in response to TBI. We found that upregulated Mmp1 expression induced by TBI was potently suppressed by knocking down the expression of Pvr, Jra or kay (Figures 4A–D, 4F), indicating that Mmp1 upregulation is mediated by Pvr and AP-1, and predominantly occurrs in astrocytes. We note that the location and intensity of Mmp1 signals are variable among different individuals (Figure 4), probably because the brain damage caused by HIT is not stereotypical.

Even though flies with Draper knockdown in astrocytes exhibited high mortality, analysis of escapers indicated that Mmp1 upregulation upon TBI was not completely inhibited by Draper knockdown (Figure 4E and 4F). Knockdown of the other receptor AdoR in astrocytes caused a weak inhibition of AP-1 and Mmp1 upregulation (Figure S4A–A", S4G–H", S4I and S4J); *TRE-dsRed* that expresses the red fluorescent protein

dsRed under the control of 4 × AP-1 motifs (Chatterjee and Bohmann, 2012) was used

as a reporter for AP-1 activity. As Pvr showed the strongest regulation on Mmp1

expression among the three receptors (Figure 4 and S4), we focused on the potential

role of Pvr in astrocytes following TBI in the present study. We observed substantial

Pvr expression predominantly in astrocytes as RNAi knockdown of Pvr in astrocytes

showed greatly reduced level of Pvr in the whole brain at 24 h after TBI (Figure S5).

Conversely, TBI further increased the Pvr level in brains when Pvr was overexpressed

under the control of *R86E01-Gal4* (Figure S5). It is noteworthy to mention that

although Draper expression was upregulated upon axotomy (MacDonald et al., 2006)

and TBI (Figure 3D), inhibiting Draper expression in astrocytes did not suppress the

Mmp1 upregulation after TBI (Figure 4E and 4F). Taken together, these results

suggest that Pvr, but not Draper, and AP-1 are required for upregulated Mmp1 in

astrocytes after TBI.

Pvr positively regulates AP-1 signaling and Mmp1 expression in astrocytes after

TBI

We next investigated the genetic pathway by which Pvr functions in astrocytes in

response to TBI. We examined the activity of AP-1 using the *in vivo* reporter

TRE-dsRed (Chatterjee and Bohmann, 2012). In uninjured brains, we detected low

TRE-dsRed levels, suggesting a basal AP-1 activity (Figure 5A–A"). Similar to Mmp1,

TRE-dsRed levels were robustly upregulated at 24 h after TBI, suggesting activation

of JNK/AP-1 signaling (Figure 5B–B"). Importantly, TBI-induced expression of

TRE-dsRed and Mmp1 was suppressed or further upregulated when inhibiting or

overexpressing Pvr expression in astrocytes, respectively (Figure 5C–F").

Overexpression of Jra or kay alone did not affect expression of TRE-dsRed or Mmp1

(Figure S4A–F", 5I and 5J), suggesting that overexpressing a single subunit of the

AP-1 complex is not sufficient to upregulate AP-1 activity. While Pvr overexpression

caused an increase in TRE-dsRed and Mmp1 expression levels, *Jra* or *kay* RNAi

knockdown in the background of Pvr overexpression suppressed the increase (Figure 5G–L), indicating that Jra and kay act downstream of Pvr to mediate the injury response upon TBI. These results together show that Pvr is a receptor acting upstream of AP-1 signaling that promotes Mmp1 expression during the injury response in astrocytes (Figure 5M).

Disruption of endocytosis and endocytic trafficking enhances Pvr, AP-1, and Mmp1 upregulation upon TBI

How was the Pvr signalling induced by TBI regulated? It has been shown that VEGFR2 endocytosis proceeds in a clathrin-dependent manner (Simons et al., 2016). During clathrin-dependent endocytosis, scission needs Cdc42-interacting protein 4 (Cip4) to recruit dynamin and Wiskott-Aldrich syndrome protein (WASp), the activator of the Arp2/3 complex that mediates actin polymerization and promotes the movement of newly-formed clathrin-coated vesicles (Brüser and Bogdan, 2017). VEGFRs are transmembrane RTKs that are activated when they locate at the cell membrane (Simons et al., 2016). We first investigated whether endocytosis is involved in Pvr signaling upon TBI. *Shibire* (*shi*) encodes a GTPase called dynamin which is responsible for fission of the vesicle from the membrane (van der Bliek and Meyerowitz, 1991). We analysed the effect of endocytosis on Pvr signaling by blocking the activity of dynamin with a temperature sensitive mutant, shi^{ts}, in astrocytes. Interestingly, both Pvr (Figures 6A, 6B, and 6E) and TRE-dsRed (Figures 6C, 6D, and 6F) intensities were increased at 4 h after TBI when dynamin activity was blocked at the non-permissive 32°C. As a control, expression of nLacZ under the same conditions as mutated dynamin did not exhibit a significant difference in the intensities of Pvr and TRE-dsRed at the permissive 23°C and the non-permissive 32°C after TBI (Figures 6A, 6B, 6E, and 6F). These results suggest that Pvr signaling is facilitated when endocytosis is blocked.

In addition to Pvr, results from our transcription profiling analysis also revealed

upregulation of the expression of actin dynamics and endocytic trafficking genes including *Arpc2*, *cpa*, and *cpb* following TBI (Tables 1 and S2). To explore whether endocytic trafficking genes are involved in the injury response in astrocytes, we analyzed Mmp1 and TRE-dsRed expression when inhibiting the expression of Arpc2, cpa, or cpb. Interestingly, TBI-induced Mmp1 levels were further increased after knockdown of each of the three (Figures 7A–D, 7O); consistently, TRE-dsRed expression was also increased after knockdown of *cpa* or *cpb* (Figures 7H–K, 7P). These results indicate that genes involved in endocytosis and endocytic trafficking normally inhibit induction of AP-1 and Mmp1 by TBI in astrocytes.

In addition, we also observed upregulated expression of a set of endosomal sorting genes, including *Stam*, *Chmp1*, and *Vps24* that encode components of the endosomal sorting complex required for transport (ESCRT)in astrocytes in response to TBI (Tables 1 and S2). These genes are implicated in the formation of intraluminal vesicles (ILVs) from multivesicular bodies (MVBs). Inhibiting expression of the ESCRT-0 subunit Stam and the ESCRT-III subunits Chmp1 or Vps24 in astrocytes however did not result in a significant difference in Mmp1 or TRE-dsRed expression levels compared with controls following TBI (Figure 7E–G, 7L-N, 7O and 7P), suggesting that endosome sorting in MVB may not affect Pvr signaling activity.

Discussion

Using a previously established a *Drosophila* TBI model, we found that TBI causes neuronal apoptosis, BBB/BEB disruption, and astrocyte hypertrophy. Astrocytes respond to TBI by elevated Pvr expression, which in turn enhances the expression of the downstream effectors AP-1 and Mmp1. Our findings identify a new signaling pathway upregulated in astrocytes following TBI.

Unlike vertebrates, in which astrocytes become reactive (reactive astrogliosis) and actively clear neuronal and synaptic debris via phagocytosis upon axon injury (Liddelow and Barres, 2017), axon transection in *Drosophila* shows no immediate

astrocytic response to injury (Doherty et al., 2009; Purice et al., 2017). Instead, *Drosophila* ENG express key components of the phagocytic machinery and engulf neuronal debris in response to axotomy (Doherty et al., 2009). Here, we examined for the first time the molecular and cellular changes of astrocytes after TBI. Although no astrocytes-mediated phagocytosis was detected upon axotomy (Doherty et al., 2009) and TBI (Figure S6), astrocytes exhibited apparent damage response including enlarged soma (hypertrophy) and altered cellular processes containing more membrane-bound GFP$^+$ accumulations (Figure 2A–J). These findings suggest that astrocytes are activated upon TBI, demonstrating a critical role for astrocytes in the injury response.

The innate immune response in *Drosophila* relies on the activation of multiple signaling pathways including NF-κB/Relish-mediated Imd and Doral/Dif-mediated Toll signaling pathways. Interestingly, our RNA-seq data revealed elevated expression of genes *Relish* and *Dif* in both pathways upon TBI (Figure 3D). However, the expressions of *AMPs* were minimum in astrocytes before TBI and no AMPs genes were identified in differentially expressed genes. Thus, AMPs genes were not activated in astrocytes, though upregulated *AMP* expressions were detected by bulk RNA-seq after TBI. In addition, expression of AP-1 (composed of Jra and kay) and Mmp1 was also upregulated, indicating that the AP-1signaling pathway is activated in astrocytes following TBI. Even though upregulated Mmp expression in reactive glial responses is an evolutionarily conserved feature of *Drosophila* and mammalian glia (Cunningham et al., 2005; Purice et al., 2017), transmembrane receptors on astrocytes that regulate Mmp1 expression have not been identified. In mammals, knockdown of VEGF signaling in keratinocytes reduced angiogenesis and delayed wound healing (Rossiter et al., 2004). In *Drosophila,* studies of wound healing showed that Pvr likely acts upstream of actin regulatory proteins that promote actin polymerization to lead cells migration (Brock et al., 2012; Wu et al., 2009).

Through a combination of RNA-seq and genetic analyses, we identified Pvr as a

new receptor in astrocytes; Pvr promotes Mmp1 expression via the transcription factor complex AP-1. At present, we do not know how Pvr promotes expression of AP-1 and Mmp1. But previous findings show that Pvr phosphorylates JNK to trigger the downstream signaling containing AP-1 during thorax closure in metamorphosis (Ishimaru et al., 2004). During development, the midline glia survival and migration are regulated by the ventral unpaired median (VUM) and the midline precursor 1 (MP1) midline neurons, the sources of Pvr ligands, i.e., PDGF- and VEGF-related factor 1-3 (Pvf1-3) (Learte et al., 2008). Additionally, epidermal cell-derived Pvf1 may be sequestered in the blood and exposed upon injury to bind Pvr receptors on wound-edge epidermal cells and initiate the extension of cell processes into the wound gap (Wu et al., 2009). Specifically, Pvr ligands bind to Pvr receptor to induce autophosphorylation of tyrosine residues, leading to JNK phosphorylation. Phosphorylated JNK (Bsk) then phosphorylates Jun/Jra transcription factors, triggering Jun/Jra association with Fos/kay, to form the AP-1 complex (Bogoyevitch and Kobe, 2006). In addition, glial AP-1 is essential for recovery, ensuring brain integrity and animal survival in the early post-TBI period (Byrns et al., 2021). Given that Pvr and AP-1 genetically interact in astrocytes upon TBI, it is likely that Pvr acts upstream of AP-1 and regulates its activity via JNK phosphorylation following brain injury. We speculate that Pvr ligands that may secret from injured cells to activate Pvr-positive astrocytes after TBI; activated Pvr then increases Mmp1 expression via AP-1 and secretion to the intercellular space to participate in injury response.

In addition to Pvr, our RNA-seq data also identified other receptors potentially functioning in astrocytes in response to TBI. Specifically, AdoR expression was upregulated after TBI (Table S2), and knockdown of AdoR expression in astrocytes led to inhibition of AP-1 after TBI (Figure S4G–J), though at a weaker extent compared with Pvr knockdown. In mammals, $A_{2B}R$, a subtype of AdoR, was shown to stimulate VEGF production in various cell types (Borea et al., 2016). Pvr promotes the expression of adenosine deaminase-related growth factor A (Adgf-A) which

inactivates extracellular adenosine, the ligand of AdoR (Zurovec et al., 2002). Furthermore, Pvr and AdoR collaborate in differentiating cells to maintain the expression of Adgf-A (Mondal et al., 2011). Thus, it is possible that Pvr and AdoR act independently or in a coordinated manner to regulate the injury response to TBI in astrocytes in *Drosophila*.

Our RNA-seq profiling also revealed upregulation of Draper. Previous studies revealed a role for Draper-mediated phagocytosis of axonal debris in ENG after axotomy (Lu et al., 2017; MacDonald et al., 2006); axotomy induces Wallerian-like degeneration (Büki and Povlishock, 2006; Tang-Schomer et al., 2010). Specifically, damage-associated molecular pattern (DAMP) proteins released from injured neurons bind to Draper and activate the JNK and JAK/STAT pathways in ENG (Doherty et al., 2014; Freeman, 2015; MacDonald et al. 2013). Here, our findings showed that Draper does not affect Mmp1 expression in astrocytes upon TBI. Although an increase in Draper expression was detected in astrocytes by RNA-seq, knockdown of its expression did not significantly suppress Mmp1 upregulation after TBI. Thus, Draper activates JNK/AP-1 and JAK/STAT signaling in ENG in response to axotomy (Doherty et al., 2014; MacDonald et al. 2013), whereas astrocytes respond to TBI via Pvr-mediated AP-1 signaling. Despite different pathways are activated in different types of glia upon brain injury, they may cooperate in response to TBI.

Endocytosis and trafficking of VEGFR, the mammalian Pvr homolog, regulate the specificity as well as the duration and strength of the signaling output (Simons et al., 2016). Once activated on the plasma membrane and translocated into the cytoplasm, VEGFRs are either shuttled to lysosomes for degradation or recycled back to the plasm membrane to replenish the receptor pool (Simons et al., 2016). Our findings showed that disruption of endocytosis by disrupting dynamin activity and inhibiting the expression of Arpc2, cpa, or cpb that regulate endocytic transport into early endosomes enhanced the AP-1 and Mmp1 upregulation in astrocyte following TBI (Figures 7A–D, 7H–K, 7O, 7P). As receptor endocytosis attenuates the strength

or duration of many signaling processes by physically reducing the concentration of cell surface receptors accessible to the ligand (Jékely et al., 2005), blocking dynamin-mediated endocytosis and endocytic trafficking potentially results in accumulated Pvr receptors on the plasma membrane to activate the downstream effector AP-1, thereby enhancing the astrocytes response to TBI. On the other hand, ubiquitinated RTKs are recognized by ESCRT-0, followed by sequential recruitment of ESCRT-I, ESCRT-II, and ESCRT-III onto the membranes of MVBs to form intraluminal vesicles (ILVs) (Goh and Sorkin, 2013). Knockdown of the ESCRT-0 subunit Stam and the ESCRT-III subunits Chmp1 and Vps24 did not affect the expression of Pvr downstream signaling components (Figure 7E–G, 7L–N, 7O and 7P), suggesting that Pvr signal transduction was not regulated by endosomal sorting.

Considering the importance of VEGF signaling in a variety of cellular processes, including wound repair (Folkman and Klagsbrun, 1987), elucidating the downstream signaling mechanisms, as well as defining the full array of Pvr targets in healthy and diseased brains, will be critical for dissecting the protective or detrimental roles of VEGF in the context of brain injury. Given that the Pvr receptor, AP-1, and the MMP family of secreted proteases are all highly conserved across species, our work provides new insights into how VEGF signaling in glia mechanistically contributes to the TBI response in mammals.

Acknowledgements

This work was supported in part by the Ministry of Science and Technology of China (2019YFA0707100 and 2021ZD0203900 to Y.Q. Zhang), the National Natural Science Foundation of China (31921002, 31830036, and 31861143031 to Y.Q. Zhang, and 31871039 and 32170962 to M. Ho), and the Chinese Academy of Sciences Strategic Priority Research Program B grants (XDBS1020100 to Y.Q. Zhang). We thank Bloomington and Tsinghua stock centers for providing fly stocks. We are also grateful

References

Apte, R. S., Chen, D. S., Ferrara, N. (2019). VEGF in Signaling and Disease: Beyond

Discovery and Development. *Cell*, *176*(6), 1248-1264.

Awasaki, T., Tatsumi, R., Takahashi, K., Arai, K., Nakanishi, Y., Ueda, R., Ito, K.

(2006). Essential role of the apoptotic cell engulfment genes draper and ced-6 in

programmed axon pruning during Drosophila metamorphosis. *Neuron*, *50*(6),

855-867.

Ayaz, D., Leyssen, M., Koch, M., Yan, J., Srahna, M., Sheeba, V., Hassan, B. A.

(2008). Axonal injury and regeneration in the adult brain of Drosophila. *Journal of*

Neuroscience, *28*(23), 6010-6021.

Bainton, R. J., Tsai, L. T., Schwabe, T., DeSalvo, M., Gaul, U., Heberlein, U. (2005).

moody encodes two GPCRs that regulate cocaine behaviors and blood-brain barrier

permeability in Drosophila. *Cell*, *123*(1), 145-156.

Barekat, A., Gonzalez, A., Mauntz, R. E., Kotzebue, R. W., Molina, B.,

El-Mecharrafie, N., Ratliff, E. P. (2016). Using Drosophila as an integrated model to

study mild repetitive traumatic brain injury. *Sci Rep*, *6*, 25252.

Bier, E. (2005). Drosophila, the golden bug, emerges as a tool for human genetics.

Nature Reviews. Genetics, *6*(1), 9-23.

Bogoyevitch, M. A., Kobe, B. (2006). Uses for JNK: the many and varied substrates

of the c-Jun N-terminal kinases. *Microbiology and Molecular Biology Reviews*, *70*(4),

1061-1095.

Bolger, A. M., Lohse, M., Usadel, B. (2014). Trimmomatic: a flexible trimmer for

Illumina sequence data. *Bioinformatics*, *30*(15), 2114-2120.

Borea, P. A., Gessi, S., Merighi, S., Varani, K. (2016). Adenosine as a Multi-Signaling

Guardian Angel in Human Diseases: When, Where and How Does it Exert its

Protective Effects? *Trends in Pharmacological Sciences*, *37*(6), 419-434.

Brock, A. R., Wang, Y., Berger, S., Renkawitz-Pohl, R., Han, V. C., Wu, Y., Galko, M. J. (2012). Transcriptional regulation of Profilin during wound closure in Drosophila larvae. *Journal of Cell Science*, *125*(Pt 23), 5667-5676.

Brüser, L., Bogdan, S. (2017). Adherens Junctions on the Move-Membrane Trafficking of E-Cadherin. *Cold Spring Harbor Perspectives in Biology*, *9*(3).

Büki, A., Povlishock, J. T. (2006). All roads lead to disconnection?--Traumatic axonal injury revisited. *Acta Neurochirurgica*, *148*(2), 181-193; discussion 193-184.

Burda, J. E., Sofroniew, M. V. (2014). Reactive gliosis and the multicellular response to CNS damage and disease. *Neuron*, *81*(2), 229-248.

Byrns C. N., Saikumar J., Bonini N. M. (2021). Glial AP1 is activated with aging and accelerated by traumatic brain injury. *Nat. Aging*, *1*(7), 585-597.

Chatterjee, N., Bohmann, D. (2012). A versatile ΦC31 based reporter system for measuring AP-1 and Nrf2 signaling in Drosophila and in tissue culture. *PLoS ONE*, *7*(4), e34063.

Cho, N. K., Keyes, L., Johnson, E., Heller, J., Ryner, L., Karim, F., Krasnow, M. A. (2002). Developmental control of blood cell migration by the Drosophila VEGF pathway. *Cell*, *108*(6), 865-876.

Chodobski, A., Zink, B. J., Szmydynger-Chodobska, J. (2011). Blood-brain barrier pathophysiology in traumatic brain injury. *Transl Stroke Res*, *2*(4), 492-516.

Cunningham, L. A., Wetzel, M., Rosenberg, G. A. (2005). Multiple roles for MMPs and TIMPs in cerebral ischemia. *Glia*, *50*(4), 329-339.

Daneshvar, D. H., Goldstein, L. E., Kiernan, P. T., Stein, T. D., McKee, A. C. (2015). Post-traumatic neurodegeneration and chronic traumatic encephalopathy. *Molecular and Cellular Neurosciences*, *66*(Pt B), 81-90.

Davalos, D., Grutzendler, J., Yang, G., Kim, J. V., Zuo, Y., Jung, S., Gan, W. B. (2005). ATP mediates rapid microglial response to local brain injury in vivo. *Nature Neuroscience*, *8*(6), 752-758.

Doherty, J., Logan, M. A., Tasdemir, O. E., Freeman, M. R. (2009). Ensheathing glia

function as phagocytes in the adult Drosophila brain. *Journal of Neuroscience*, *29*(15),

4768-4781.

Doherty, J., Sheehan, A. E., Bradshaw, R., Fox, A. N., Lu, T. Y., Freeman, M. R.

(2014). PI3K signaling and Stat92E converge to modulate glial responsiveness to

axonal injury. *PLoS Biology*, *12*(11), e1001985.

Egeblad, M., Werb, Z. (2002). New functions for the matrix metalloproteinases in

cancer progression. *Nature Reviews. Cancer*, *2*(3), 161-174.

Folkman, J., Klagsbrun, M. (1987). Angiogenic factors. *Science*, *235*(4787), 442-447.

Freeman, M. R. (2015). Drosophila Central Nervous System Glia. *Cold Spring*

Harbor Perspectives in Biology, *7*(11).

Goh, L. K., Sorkin, A. (2013). Endocytosis of receptor tyrosine kinases. *Cold Spring*

Harbor Perspectives in Biology, *5*(5), a017459.

Hoch, R. V., Soriano, P. (2003). Roles of PDGF in animal development. *Development*,

130(20), 4769-4784.

Hoopfer, E. D., McLaughlin, T., Watts, R. J., Schuldiner, O., O'Leary, D. D., Luo, L.

(2006). Wlds protection distinguishes axon degeneration following injury from

naturally occurring developmental pruning. *Neuron*, *50*(6), 883-895.

Hu, Y., Flockhart, I., Vinayagam, A., Bergwitz, C., Berger, B., Perrimon, N., Mohr, S.

E. (2011). An integrative approach to ortholog prediction for disease-focused and

other functional studies. *BMC Bioinformatics*, *12*, 357.

Ishimaru, S., Ueda, R., Hinohara, Y., Ohtani, M., Hanafusa, H. (2004). PVR plays a

critical role via JNK activation in thorax closure during Drosophila metamorphosis.

EMBO Journal, *23*(20), 3984-3994.

Jékely, G., Sung, H. H., Luque, C. M., Rørth, P. (2005). Regulators of endocytosis

maintain localized receptor tyrosine kinase signaling in guided migration.

Developmental Cell, *9*(2), 197-207.

Jia, M., Meng, D., Chen, M., Li, T., Zhang, Y. Q., Yao, A. (2019). Drosophila

homolog of the intellectual disability-related long-chain acyl-CoA synthetase 4 is required for neuroblast proliferation. *J Genet Genomics*, *46*(1), 5-17.

Joy, M. T., Ben Assayag, E., Shabashov-Stone, D., Liraz-Zaltsman, S., Mazzitelli, J., Arenas, M., Carmichael, S. T. (2019). CCR5 Is a Therapeutic Target for Recovery after Stroke and Traumatic Brain Injury. *Cell*, *176*(5), 1143-1157.e1113.

Katzenberger, R. J., Chtarbanova, S., Rimkus, S. A., Fischer, J. A., Kaur, G., Seppala, J. M., Wassarman, D. A. (2015). Death following traumatic brain injury in Drosophila is associated with intestinal barrier dysfunction. *Elife*, *4*.

Katzenberger, R. J., Ganetzky, B., Wassarman, D. A. (2016). Age and diet affect genetically separable secondary injuries that cause acute mortality following traumatic brain injury in Drosophila. *G3 (Bethesda)*, *6*(12), 4151-66.

Katzenberger, R. J., Loewen, C. A., Wassarman, D. R., Petersen, A. J., Ganetzky, B., Wassarman, D. A. (2013). A Drosophila model of closed head traumatic brain injury. *Proceedings of the National Academy of Sciences of the United States of America*, *110*(44), E4152-4159. Kim, D., Langmead, B., Salzberg, S. L. (2015). HISAT: a fast spliced aligner with low memory requirements. *Nature Methods*, *12*(4), 357-360.

Kim, D., Paggi, J. M., Park, C., Bennett, C., Salzberg, S. L. (2019). Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype. *Nature Biotechnology*, *37*(8), 907-915.

Kremer, M. C., Jung, C., Batelli, S., Rubin, G. M., Gaul, U. (2017). The glia of the adult Drosophila nervous system. *Glia*, *65*(4), 606-638.

Learte, A. R., Forero, M. G., Hidalgo, A. (2008). Gliatrophic and gliatropic roles of PVF/PVR signaling during axon guidance. *Glia*, *56*(2), 164-176.

Lee, T., Luo, L. (1999). Mosaic analysis with a repressible cell marker for studies of gene function in neuronal morphogenesis. *Neuron*, *22*(3), 451-461.

Liddelow, S. A., Barres, B. A. (2017). Reactive Astrocytes: Production, Function, and Therapeutic Potential. *Immunity*, *46*(6), 957-967.

Liddelow, S. A., Guttenplan, K. A., Clarke, L. E., Bennett, F. C., Bohlen, C. J.,

Schirmer, L., Barres, B. A. (2017). Neurotoxic reactive astrocytes are induced by

activated microglia. *Nature*, *541*(7638), 481-487.

Love, M. I., Huber, W., Anders, S. (2014). Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. *Genome Biology*, *15*(12), 550.

Lu, T. Y., Doherty, J., Freeman, M. R. (2014). DRK/DOS/SOS converge with

Crk/Mbc/dCed-12 to activate Rac1 during glial engulfment of axonal debris. *Proc.*

Natl. Acad. Sci. USA, *111*(34), 12544-12549.

Lu, T. Y., MacDonald, J. M., Neukomm, L. J., Sheehan, A. E., Bradshaw, R., Logan,

M. A., Freeman, M. R. (2017). Axon degeneration induces glial responses through

Draper-TRAF4-JNK signaling. *Nature Communications*, *8*, 14355.

Ma, V. Y., Chan, L., Carruthers, K. J. (2014). Incidence, prevalence, costs, and impact

on disability of common conditions requiring rehabilitation in the United States:

stroke, spinal cord injury, traumatic brain injury, multiple sclerosis, osteoarthritis,

rheumatoid arthritis, limb loss, and back pain. *Archives of Physical Medicine and*

Rehabilitation, *95*(5), 986-995.e981.

MacDonald, J. M., Beach, M. G., Porpiglia, E., Sheehan, A. E., Watts, R. J., Freeman,

M. R. (2006). The Drosophila cell corpse engulfment receptor Draper mediates glial

clearance of severed axons. *Neuron*, *50*(6), 869-881.

Macdonald, J. M., Doherty, J., Hackett, R., Freeman, M. R. (2013). The c-Jun kinase

signaling cascade promotes glial engulfment activity through activation of draper and

phagocytic function. *Cell Death and Differentiation*, *20*(9), 1140-1148.

McDonald, J. A., Pinheiro, E. M., Montell, D. J. (2003). PVF1, a PDGF/VEGF

homolog, is sufficient to guide border cells and interacts genetically with Taiman.

Development, *130*(15), 3469-3478.

Mira, R. G., Lira, M., Cerpa, W. (2021). Traumatic Brain Injury: Mechanisms of Glial

Response. *Front Physiol*, *12*, 740939.

Mondal, B. C., Mukherjee, T., Mandal, L., Evans, C. J., Sinenko, S. A.,

Martinez-Agosto, J. A., Banerjee, U. (2011). Interaction between differentiating cell-

and niche-derived signals in hematopoietic progenitor maintenance. *Cell*, *147*(7),

1589-1600.

Moreno, E., Yan, M., Basler, K. (2002). Evolution of TNF signaling mechanisms:

JNK-dependent apoptosis triggered by Eiger, the Drosophila homolog of the TNF

superfamily. *Current Biology*, *12*(14), 1263-1268.

Motz, G. T., Coukos, G. (2013). Deciphering and reversing tumor immune

suppression. *Immunity*, *39*(1), 61-73. https://doi.org/10.1016/j.immuni.2013.07.005

Ng, S. Y., & Lee, A. Y. W. (2019). Traumatic Brain Injuries: Pathophysiology and

Potential Therapeutic Targets. *Front Cell Neurosci*, *13*, 528.

Page-McCaw, A., Ewald, A. J., Werb, Z. (2007). Matrix metalloproteinases and the

regulation of tissue remodelling. *Nature Reviews. Molecular Cell Biology*, *8*(3),

221-233.

Parks, W. C., Wilson, C. L., López-Boado, Y. S. (2004). Matrix metalloproteinases as

modulators of inflammation and innate immunity. *Nature Reviews. Immunology*, *4*(8),

617-629.

Pastor-Pareja, J. C., Grawe, F., Martín-Blanco, E., García-Bellido, A. (2004). Invasive

cell behavior during Drosophila imaginal disc eversion is mediated by the JNK

signaling cascade. *Developmental Cell*, *7*(3), 387-399.

Pekny M, Pekna M. (2004). Astrocyte intermediate filaments in CNS pathologies and

regeneration. *The Journal of Pathology, 204*(4):428-37.

Pertea, M., Kim, D., Pertea, G. M., Leek, J. T., Salzberg, S. L. (2016). Transcript-level

expression analysis of RNA-seq experiments with HISAT, StringTie and Ballgown.

Nature Protocols, *11*(9), 1650-1667.

Pfeiffer, B. D., Truman, J. W., Rubin, G. M. (2012). Using translational enhancers to

increase transgene expression in Drosophila. *Proceedings of the National Academy of

Sciences of the United States of America, 109*(17), 6626-6631.

Poernbacher, I., Vincent, J. P. (2018). Epithelial cells release adenosine to promote

local TNF production in response to polarity disruption. *Nature Communications*, *9*(1),

4675.

Purice, M. D., Ray, A., Munzel, E. J., Pope, B. J., Park, D. J., Speese, S. D., Logan, M.

A. (2017). A novel Drosophila injury model reveals severed axons are cleared through

a Draper/MMP-1 signaling cascade. *Elife*, *6*.

Rodríguez-Baeza, A., Reina-de la Torre, F., Poca, A., Martí, M., Garnacho, A. (2003).

Morphological features in human cortical brain microvessels after head injury: a

three-dimensional and immunocytochemical study. *Anatomical Record. Part A,*

Discoveries in Molecular, Cellular, and Evolutionary Biology, *273*(1), 583-593.

Rossiter, H., Barresi, C., Pammer, J., Rendl, M., Haigh, J., Wagner, E. F., Tschachler,

E. (2004). Loss of vascular endothelial growth factor a activity in murine epidermal

keratinocytes delays wound healing and inhibits tumor formation. *Cancer Research*,

64(10), 3508-3516.

Saikumar J, Byrns C. N., Hemphill M., Meaney D. F., Bonini N. M. (2020). Dynamic

neural and glial responses of a head-specific model for traumatic brain injury in

Drosophila. *Proceedings of the National Academy of Sciences of the United States of*

America, 117(29):17269-77.

Sakurai, Y., Ohgimoto, K., Kataoka, Y., Yoshida, N., Shibuya, M. (2005). Essential

role of Flk-1 (VEGF receptor 2) tyrosine residue 1173 in vasculogenesis in mice.

Proceedings of the National Academy of Sciences of the United States of America,

102(4), 1076-1081.

Sangiorgi, S., De Benedictis, A., Protasoni, M., Manelli, A., Reguzzoni, M., Cividini,

A., Balbi, S. (2013). Early-stage microvascular alterations of a new model of

controlled cortical traumatic brain injury: 3D morphological analysis using scanning

electron microscopy and corrosion casting. *Journal of Neurosurgery*, *118*(4), 763-774.

Shi, X., Luo, L., Wang, J., Shen, H., Li, Y., Mamtilahun, M., Yang, G. Y. (2021).

Stroke subtype-dependent synapse elimination by reactive gliosis in mice. *Nature*

Communications, *12*(1), 6943.

Sil, S., Periyasamy, P., Thangaraj, A., Chivero, E. T., Buch, S. (2018). PDGF/PDGFR

axis in the neural systems. *Molecular Aspects of Medicine*, *62*, 63-74.

Simons, M., Gordon, E., Claesson-Welsh, L. (2016). Mechanisms and regulation of

endothelial VEGF receptor signaling. *Nature Reviews. Molecular Cell Biology*, *17*(10),

611-625.

Sofroniew, M. V., Vinters, H. V. (2010). Astrocytes: biology and pathology. *Acta*

Neuropathologica, *119*(1), 7-35.

Stork, T., Sheehan, A., Tasdemir-Yilmaz, O. E., Freeman, M. R. (2014). Neuron-glia

interactions through the Heartless FGF receptor signaling pathway mediate

morphogenesis of Drosophila astrocytes. *Neuron*, *83*(2), 388-403.

Swanson, L. C., Trujillo, E. A., Thiede, G. H., Katzenberger, R. J., Shishkova, E.,

Coon, J. J., Ganetzky, B., Wassarman, D. A. (2020). Survival following traumatic

brain injury in Drosophila is increased by heterozygosity for a mutation of the

NF-kappaB innate immune response transcription factor Relish. *Genetics*, *216*(4),

1117-1136.

Tang-Schomer, M. D., Patel, A. R., Baas, P. W., Smith, D. H. (2010). Mechanical

breaking of microtubules in axons during dynamic stretch injury underlies delayed

elasticity, microtubule disassembly, and axon degeneration. *FASEB Journal*, *24*(5),

1401-1410.

Thacker, S. A., Bonnette, P. C., Duronio, R. J. (2003). The contribution of

E2F-regulated transcription to Drosophila PCNA gene function. *Current Biology*,

13(1), 53-58.

Valanne, S., Wang, J. H., Rämet, M. (2011). The Drosophila Toll signaling pathway.

Journal of Immunology, *186*(2), 649-656.

Van Alphen B., Stewart S., Iwanaszko M., Xu F, Li K., Rozenfeld S., Ramakrishnan

A., Itoh T. Q., Sisobhan S., Qin Z. et al., (2022). Glial immune-related pathways

mediate effects of closed head traumatic brain injury on behavior and lethality in

Drosophila. *PLoS Biololy*, *20*(1), e3001456.

Willsey, H. R., Zheng, X., Carlos Pastor-Pareja, J., Willsey, A. J., Beachy, P. A., Xu, T.

(2016). Localized JNK signaling regulates organ size during development. *Elife*, *5*.

A. M. van der Bliek, E. M. Meyerowitz (1991). Dynamin-like protein encoded by the

Drosophila shibire gene associated with vesicular traffic. *Nature, 351*(6325), 411-414.

Wu, Y., Brock, A. R., Wang, Y., Fujitani, K., Ueda, R., Galko, M. J. (2009). A

blood-borne PDGF/VEGF-like ligand initiates wound-induced epidermal cell

migration in Drosophila larvae. *Current Biology*, *19*(17), 1473-1477.

Yang, C. P., Fu, C. C., Sugino, K., Liu, Z., Ren, Q., Liu, L. Y., Lee, T. (2016).

Transcriptomes of lineage-specific Drosophila neuroblasts profiled by genetic

targeting and robotic sorting. *Development*, *143*(3), 411-421.

Yu, S., He, J. (2019). Stochastic cell-cycle entry and cell-state-dependent fate outputs

of injury-reactivated tectal radial glia in zebrafish. *Elife*, *8*.

Zhao, K., Hong, H., Zhao, L., Huang, S., Gao, Y., Metwally, E., Zhang, Y. Q. (2020).

Postsynaptic cAMP signaling regulates the antagonistic balance of Drosophila

glutamate receptor subtypes. *Development*, *147*(24).

Zurovec, M., Dolezal, T., Gazi, M., Pavlova, E., Bryant, P. J. (2002). Adenosine

deaminase-related growth factors stimulate cell proliferation in Drosophila by

depleting extracellular adenosine. *Proceedings of the National Academy of Sciences*

of the United States of America, *99*(7), 4403-4408.

    Figure legends

Figure 1. TBI is induced in adult flies by a high impact trauma (HIT) device.

(A) MI_{24} is the mortality index calculated from the percentage of dead over total flies.

Increasing the number of strikes from 4 to 6 increases the MI_{24} value. Error bars

denote standard error of the mean (SEM; ***p <0.001, Student's t-test). (B–E) The

intactness of blood-eye-barrier (BEB) and blood-brain-barrier (BBB) before or after 6

strikes is indicated by the accumulated fluorescence intensity at the border of the eye

(B, D) and brain (C, E), respectively. The examination was carried out at 2 h after

fluorescent dye injection. Arrows indicate the hemolymph exclusion line. Scale bar =

100 μm. (F) The percentage of permeable BEB with or without TBI (***p <0.001).

(G) The percentage of permeable BBB following different numbers of strikes. Error

bars denote SEM (**p <0.01 and ***p <0.001, Student's t-test). (H) Histogram

showing mRNA expression levels of innate immune response genes of the AMP

family (*Attc*, *DiptB* and *Mtk*) in the adult brain. Expression was increased in all cases

following treatment. The *RPL28* gene served as an internal reference. Results are

means ± SEM (***p <0.001, Student's t-test). (I, J) The number of Dcp-1^{+} puncta

(green) increased in injured adult brains 24 h after TBI. Scale bar = 50 μm. (K)

Quantification of Dcp-1^{+} puncta. Error bars denote SEM (**p <0.01, Student's t-tests).

Results are means ± SEM. (L–M") Apoptotic debris labelled with anti-Dcp-1

antibodies. Glial and neuronal nuclei are labelled with anti-Repo (L) and anti-Elav (M)

antibodies, respectively. Three representative images (L–L" and M–M") for each pair

of antibody staining are shown. Scale bar = 20 μm.

Figure 2. TBI alters glial morphology and upregulates Mmp1 expression.

(A–C) Representative images of astrocyte membranes labelled with *mCD8::GFP*

driven by the astrocyte driver *R86E01-Gal4*. Co-expression of *UAS-Redstinger* labels

astrocyte nuclei. GFP intensities are increased at 4 h (B) and 24 h (C) after injury. The

rectangle defined by white dash lines denotes antennal lobes. Scale bars = 100 μm. (D)

Quantification of GFP intensity at different timepoints after TBI. Results are

means $\pm$ SEM (**p <0.01, Student's t-test). (E, F) MARCM analysis reveals single

astrocytes with evenly distributed GFP$^+$ cellular processes without injury (E) and an

increased number of GFP$^+$ puncta at 24 h after injury (F). Arrows denote GFP$^+$

accumulations in astrocyte processes. White dots demarcate neuropils. Anti-Repo

antibodies labelled glial nuclei. TO-PRO labelled all nuclei. Insets show the

morphology of a glial soma before and after injury. Scale bar = 10 μm. (G)

Quantification of GFP$^+$ puncta in astrocyte processes before and after injury. (H)

Quantification of the astrocyte soma diameter surrounding ALs after injury. Results

are means $\pm$ SEM (***p <0.001, Student's t-test). (I–J", M–N") Representative images

of ALs co-labelled with GFP and Mmp1 (magenta) in uninjured and injured adult

brains. (J–J") GFP-labelled astrocyte membranes (green) overlapped with Mmp1

(magenta) after injury. (N–N") After injury, Mmp1 co-localizes with GFP-labelled

ENG (*R56F03-Gal4*) membranes (green). Arrows show Mmp1-GFP colocalization in

L" and P". Scale bar = 20 μm. (K, L) Graph showing fluorescence intensity per pixel

in each channel for the inset in I" and J" of the astrocyte membrane and Mmp1 along a

line drawn through a cell body. (O, P) Graph showing fluorescence intensity per pixel

in each channel for the inset in M" and N" of the ENG membrane and Mmp1 along a

line drawn through the process of ENG. (Q) Quantification of Mmp1 intensity in I–J"

and M–N". Ast is the short of astrocytes. Results are means $\pm$ SEM (**p <0.01,

Student's t-test).

Figure 3. Transcriptome analysis of astrocytes reveals upregulation of

injury-related and innate immunity genes.

(A) After dissection, *Drosophila* brains were disrupted and the cell suspension was

subjected to FACS. Astrocyte cDNAs were amplified by the Smart-seq2 method and a

cDNA library was constructed. A subset of upregulated genes from RNA-seq

screening was validated by quantitative real-time PCR. (B, C) Volcano plots of the

gene expression analysis; 522 differentially expressed genes (416 upregulated, 106

downregulated) in astrocytes were detected at 4 h after injury (B), and 357

differentially expressed genes (229 upregulated, 129 downregulated) in astrocytes

were detected at 24 h after injury (C). The threshold for differential expression was set

at $\log_2$FC = 1 with an adj. p <0.01. FC = fold change; adj. = adjusted (for multiple

comparisons by the Benjamin-Hochberg procedure). (D) Heatmap of the expression

of a small set of representative genes, including *draper*, *Jra* and *kay*, previously

identified as essential factors during glial clearance of severed axons. (E) Quantitative

real-time PCR validation of a subset of upregulated genes from RNA-seq screening.

Error bars represent SEM. Results are means ± SEM (*p <0.05, **p <0.01 and

***p <0.001 by Student's t tests).

Table 1. Upregulated genes related to receptors and endocytosis.

Figure 4. Pvr, Jra and kay are required for Mmp1 upregulation after TBI.

(A–F) Knockdown of Pvr, Jra or kay by *R86E01-Gal4* driven RNAi in astrocytes

suppresses Mmp1 upregulation after TBI. Scale bar = 100 μm. All images are

projections of 25–30 μm immunostaining slices (one image per 1 μm slice). Arrows

show Mmp1 signals. (F) Quantification of Mmp1 intensity in different genotypes.

Results are means ± SEM (*p <0.05 by Student's t-test).

Figure 5. Pvr is required for AP-1 upregulation after TBI.

(A–B") Adult brains showing increased *TRE-dsRed* (an AP-1 reporter) expression at

24 h after TBI. (C–D") Overexpression of Pvr in astrocytes leads to upregulated

dsREd expression compared with control flies. Scale bar = 100 μm. (E–F")

Knockdown of Pvr in astrocytes causes downregulation of dsRed expression

compared with control flies after TBI. (G–H") Overexpression of Pvr and knockdown

of Jra at the same time in astrocytes causes downregulation of dsRed and Mmp1 after

TBI compared with controls. (I–J'') Overexpression of Pvr and knockdown of kay in astrocytes at the same time causes downregulation of dsRed and Mmp1 after TBI compared with controls. (K, L) Quantification of Mmp1 (K) and dsRed (L) intensity. Results are means ± SEM **p <0.01, ***p <0.001, Student's t-tests.

Figure 6. Inhibition of dynamin activity increases the expression levels of Pvr and dsRed after TBI.

(A, B) Representative Pvr immunostaining (green), dsRed (magenta) and β-gal intensity (gray) in control animals (*TRE-dsRed;R86E01-Gal4>UAS-nLacZ*) at 4 h after TBI. All images are projections of 25–30 μm immunostaining slices (one image per 1 μm slice). (C, D) Representative Pvr immunostaining (green), dsRed (magenta) and Mmp1 intensity (gray) in *shi^{ts}*-expressing animals (*TRE-dsRed;R86E01-Gal4>UAS-shi^{ts}*) at 4 h after TBI. For thermogenetic inactivation of dynamin, flies after TBI were incubated at intermittent temperature at 32°C (32°C for 30 min, and then back to 23°C for 30 min, for four rounds, four hours in total) until dissection. (E, F) Quantification of Pvr and dsRed intensity in different genotypes. Results are means ± SEM (*p <0.05).

Figure 7. Knockdown of endocytic trafficking-related genes up-regulates expression of AP-1 and Mmp1upon TBI

(A, H) Representative Mmp1 immunostaining (magenta) and dsRed intensity (magenta) in control animals (*R86E01-Gal4*) at 24 h after TBI. (B–D, I–K) Knockdown of *Arpc2*, *cpa*, and *cpb* in astrocytes results in further upregulation of Mmp1 and dsRed upon TBI. All images are projections of 25–30 μm immunostaining slices (one image per 1 μm slice). (E–G, L–N) Knockdown of *Stam*, *Chmp1* or *Vps24*, components of ESCRT, in astrocytes had no influence on Mmp1 and dsRed upregulation compared with control after TBI (A, H). All images are projections of 25–30 μm immunostaining slices (one image per 1 μm slice). (O, P) Quantification of

Mmp1 and dsRed intensity in different genotypes. Results are means $\pm$ SEM

(*p <0.05, **p <0.01, ***p <0.001, Student's t-tests). (Q) Schematic diagram of Pvr–

AP-1–MMP1 signaling activated by TBI in astrocytes. AP-1 transcription factor

complex is composed of Jra and kay.

Figure S1. No cell proliferation is detected in adult brains.

(A–C") Drosophila brains expressing the S-phase activity reporter PCNA::GFP at

larval and adult stages were dissected and co-stained with the mitotic marker

phosphohistone H3 (pH3). As controls, uninjured larval brains exhibit both GFP$^+$ and

pH3$^+$ signals, indicating that cells are proliferative at this stage (A–A"). In both

uninjured and injured adult brains, no apparent GFP$^+$ and pH3$^+$ signals were detected

(B–C"). Scale bar = 50 μm. (D–F) No pH3$^+$ cells were observed in adult brains at

different time points of 48 h (D), 72 h (E) and 5 days (F) after TBI. Scale bar = 50

μm.

Figure S2. Sorting of adult astrocytes by FACS.

(A) To discriminate between healthy and damaged or dying cells, side scatter (SSC-A,

log) and forward scatter (FSC-A, log) were plotted. Higher SSC-A signals indicate

increased granularity and therefore likely dying cells. Thus, one should define the

sorting gate as the population with the lower SSC-A signal (P1). (P2) Exclude cell

clusters by sorting only events with a low FSC width (FSC-W) versus similar area

(FSC-A) signal to get exact single cell. (P3) To gate for RFP$^+$ cells, plotted the signal

from the excitation of cells with the 568-nm laser (PE-Texas Red-A, log) against the

signal from the SSC-A. The population with the higher PE-Texas Red-A signal was

the candidate cells cluster. The strategy for FACS is shown at the lower right corner.

(B–C') A population of cells with strong RFP signal was purified from adult flies of

R86E01-Gal4>UAS-Redstinger. (B and B') Confocal images of representative

astrocytes before and after sorting by FACS. Scale bar = 50 μm. (C and C') As

cell-permeant MitoTracker labels mitochondria of live cells, cells after FACS both

labelled by RFP and MitoTracker. Scale bar = 10 μm. (D) A Principal Component

Analysis of all biological replicates (uninjured versus injured) revealing genes

expressed at absolute value of logarithmic fold change (log$_2$FC) > 1.0.

Figure S3. GO analysis of differentially expressed genes at 4 h and 24 h after

injury.

Scatterplot showing GO functional enrichment results for differentially expressed

genes of astrocytes in the brain at 4 h (A) and 24 h (B) after injury, $p < 0.01$.

Figure S4. Overexpression of Jra or kay does not enhance the expression of

TRE-RFP and Mmp1 in adult brain after TBI.

(A, B") Fly brains showing TRE-RFP expression before and 24 h after TBI. (C–F")

overexpression of Jra or kay in astrocytes shows no apparent changes of RFP and

Mmp1 staining signals compared with control flies with or without brain injury

(*TRE-dsRed/UAS-Jra* or *kay*; *R86E01-Gal4/+*). Scale bar = 100 μm. (G, H)

Knockdown of AdoR in astrocytes inhibits the expression of Mmp1 after TBI.

However, Knockdown of AdoR in astrocytes does not inhibit the expression of dsRed.

(I, J) Quantification of Mmp1 and dsRed intensity before and after injury in different

genotypes. Data are present as means ± SEM; *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$ by

Student's *t* tests.

1010

Figure S5. Pvr is upregulated in astrocytes in the adult brains after TBI.

Immunostainings of Pvr in the control brains showed Pvr was expressed at low levels

in uninjured brains (A) and its expression was upregulated at 24 h after TBI (B). (C, D)

Knockdown of Pvr in astrocytes reduced Pvr expression upon TBI. (E, F) Pvr

overexpression in astrocytes increased Pvr expression upon TBI. Scale bar = 100 μm.

(G) Quantification of Pvr intensity in A–E. Data are present as means ± SEM;

*$p < 0.05$, ***$p < 0.001$ by Student's *t* tests.

1018

Figure S6. Astrocytes do not engulf debris in adult brain after TBI.

(A–C ") Confocal images are single z-confocal slices. Astrocytes membranes were

labeled with GFP (*UAS-mCD8::GFP* expression driven by *R86E01-Gal4*; green);

nuclei were labeled with TO-PRO3 (blue), while axons were labeled with Ncad (a cell

adhesion molecule; magenta). Inset, high-magnification view of the boxed region.

Scale bars = 10 μm. (D–F) The graph shows fluorescence intensity in each channel for

the inset in A–C, respectively, of GFP, Ncad and nuclear staining along a line drawn

through an astrocyte nucleus.

Figure 1

Figure 2

R86E01-Gal4>UAS-mCD8::GFP; UAS-Redstinger

No injury
A GFP · Redstinger · Merge

4 h after injury
B

24 h after injury
C 100 μm

D
GFP intensity (%)
□ No injury
▨ 4 h after TBI
▦ 24 h after TBI

E No injury
GFP
repo
T3605

F 24 h after TBI
10 μm

G
Puncta/μm²
***

H
Diameter of Ast. (um)
□ No injury
▦ 24 h after TBI
***

R86E01-GAL4>mCD8::GFP
No injury
I GFP **I'** Mmp-1 **I''** Merge

24 h after injury
J **J'** **J''**

K No injury

L 24 h after injury
Distance (μm)

R56F03-GAL4>mCD8::GFP
No injury
M **M'** **M''** 20 μm

24 h after injury
N GFP **N'** Mmp-1 **N''** Merge

O No injury

P 24 h after injury
GFP
Mmp-1
Distance (μm)

Q
Mmp1 intensity
ENG Ast.
□ No injury
▦ 24 h after TBI
**

Figure 3

Table 1

Genes in *Drosophila*	Conserved genes in mouse	Conserved genes in human	4 h after injury (FC)	24 h after injury (FC)
Pvr	pdgfr/vegfr	VEGFR	13.83	2.79
AdoR	Adora2a	ADORA2B	37.53	7.73
Mmp1	Mmp14	MMP14	26.35	4.79
Jra	Jun	JUN	3.58	1.18
kay	FOSL2	FOS	6.23	1.77
cpa	Capza2	CAPZA1	3.43	1.72
cpb	Capzb	CAPZB	2.01	1.25
Stam	Stam2	STAM	2.14	1.46
Chmp1	Chmp1b	CHMP1B	2.16	0.92
Vps24	Chmp3	CHMP3	2.11	0.87

Figure 4

Figure 5

Figure 6

TRE-dsRed;R86E01-Gal4>UAS-nLacZ

A No injury (4 h) — Pvr | TRE-dsRed | β-gal | Merge

B 4 h after TBI

TRE-dsRed;R86E01-Gal4>UAS-shi^{ts}

C No injury (4 h) — Pvr | TRE-dsRed | Mmp1 | Merge

D 4 h after TBI

100 μm

Reared and treated by HIT device at 23°C | 32°C | 23°C | 32°C | 23°C | 32°C | 23°C | 32°C | 23°C
30 min for each treatment

E Pvr intensity

F dsRed intensity

TRE-dsRed;R86E01-Gal4 >UAS-nLacZ

TRE-dsRed;R86E01-Gal4 >UAS-shi^{ts}

□ No injury
■ 4 h after TBI

Figure 7

Figure S1

Larva / No injury: A (GFP), A' (pH3), A'' (Merge)

Adult / No injury: B, B', B''

Adult / 24 h after TBI: C, C', C'' — 50 μm

D 48 h after injury (pH3)

E 72 h after injury

F 5 d after injury — 50 μm

Figure S2

Figure S3

Figure S4

Figure S5

Figure S6

R86E01-GAL4>mCD8::GFP