1

**Title: Susceptibility to Zika virus in a Collaborative Cross mouse strain is**

**induced by *Irf3* deficiency *in vitro* but requires other variants *in vivo*, not**

**linked to the type I interferon response**

5
Short title: *Irf3* deficiency induces susceptibility to ZIKV *in vitro*, not *in vivo*

Marie Bourdon[§1], Caroline Manet[§1], Laurine Conquet[1], Corentin Ramaugé-Parra[1], Etienne

Kornobis[2], Eliette Bonnefoy[3], Xavier Montagutelli[#1]

[1]Institut Pasteur, Université de Paris, Mouse Genetics Laboratory, F-75015 Paris, France

[2]Institut Pasteur, Université de Paris, Biomics, F-75015 Paris, France

[3]Université Paris Cité, Institut Cochin, Inserm, CNRS, F-75014 Paris, France

[§] M.B. and C. M. equally contributed to this work and are co-first authors

[#] Corresponding author: xavier.montagutelli@pasteur.fr

Keywords: Zika virus, mouse, Irf3, interferon response, Collaborative Cross

**Abstract**

Zika virus (ZIKV) is a Flavivirus responsible for recent epidemics in Pacific Islands and in the

Americas. In humans, the consequences of ZIKV infection range from asymptomatic infection

to severe neurological disease such as Guillain-Barré syndrome or fetal neurodevelopmental

defects, suggesting, among other factors, the influence of host genetic variants. We previously

reported similar diverse outcomes of ZIKV infection in mice of the Collaborative Cross (CC), a

collection of inbred strains with large genetic diversity. CC071/TauUnc (CC071) was the most

susceptible CC strain with severe symptoms and lethality. Notably, CC071 has been recently

reported to be also susceptible to other flaviviruses including dengue virus, Powassan virus,

West Nile virus, and to Rift Valley fever virus. To identify the genetic origin of this broad

susceptibility, we investigated ZIKV replication in mouse embryonic fibroblasts (MEFs) from

CC071 and two resistant strains. CC071 showed uncontrolled ZIKV replication associated with

delayed induction of type-I interferons (IFN-I). Genetic analysis identified a mutation in the

*Irf3* gene specific to the CC071 strain which prevents the protein phosphorylation required to

activate interferon beta transcription. We demonstrated that this mutation induces the same

defective IFN-I response and uncontrolled viral replication in MEFs as an *Irf3* knock-out allele.

By contrast, we also showed that *Irf3* deficiency did not induce the high plasma viral load and

clinical severity observed in CC071 mice and that susceptibility alleles at other genes, not

associated with the IFN-I response, are required. Our results provide new insight into the *in*

*vitro* and *in vivo* roles of *Irf3*, and into the genetic complexity of host responses to flaviviruses.

**Author summary**

Recent ZIKV outbreaks led to millions of infected people, with rare but severe complications

such as Guillain-Barré syndrome and encephalitis in adults suggesting that host genes

influence the susceptibility to severe forms of infection. We previously reported the

importance of host genes in ZIKV pathogenesis using a panel of genetically diverse mouse

strains and identified CC071 as the most susceptible strain. Importantly, this mouse strain has

been shown by others to be also susceptible to several other RNA viruses. Through a

combination of functional and genetic approaches in a cellular model, we identified a

mutation in the *Irf3* gene which plays a key role in activating the expression of interferon beta

to induce the type I interferon response, the first line of host defense against the virus. This

mutation fully explains the high viral replication observed in CC071 cells. However, it was not

able to induce the elevated viremia and the symptoms displayed by CC071 ZIKV-infected mice,

unraveling the implication of other host genes which are not associated with the type I

interferon response. Because of the broad susceptibility of CC071 to multiple viruses, our

results have implications beyond ZIKV infection and contribute to shedding light on the

plurality of host mechanisms fighting infectious diseases.

## Introduction

Zika virus (ZIKV) is a mosquito-borne virus of the *Flaviviridae* family identified in 1947 in Uganda. The first noticeable human outbreaks occurred in Micronesia in 2007 and in French Polynesia and New Caledonia in 2013-2014. In 2015-2016, ZIKV caused an epidemic in Brazil which rapidly spread across the Americas and the Caribbean. To date, 89 countries have reported evidence of mosquito-transmitted Zika virus infection (https://www.who.int/health-topics/zika-virus-disease#tab=tab_1).

While most people infected with ZIKV remain asymptomatic, some develop non-specific symptoms including rash, fever, conjunctivitis, muscle and joint pain, malaise and headache. Neurological complications have been described in adults such as Guillain-Barré syndrome (1) and encephalitis (2). Infection of pregnant women was associated with congenital Zika syndrome in the fetus, which can lead to neurodevelopmental deficiencies, brain malformation (3) or in some cases to fetal loss (4).

Many factors may contribute to this variable severity, including the viral strain, the infection route and dose, and the host genetic background (5,6). Indeed, mouse and human studies have shown that host genes influence flaviviral infections' outcomes (7). While human genetic studies are hampered by the variability of these multiple factors, they can be controlled in mouse models which have proven very valuable to identify susceptibility variants (8,9). Relevant ZIKV infection models have been developed in mice either using *Ifnar1* knock-out (KO) mice in which the IFNAR receptor to IFN-I has been inactivated (10,11), or by blocking this receptor using a monoclonal antibody targeting the IFNAR1 receptor subunit (MAR1-5A3 (12)).

We have previously explored the role of mouse natural genetic variants on ZIKV susceptibility in the Collaborative Cross (CC), a panel of recombinant inbred mice encompassing a genetic

diversity similar to that of the human population and capturing approximately 90% of the

mouse natural genetic variants (13,14). We reported that the CC genetic diversity enabled

large variations in the clinical severity of ZIKV disease, plasma viral load and intensity of brain

pathology, comparable to those observed in human cases (15). We specifically identified

CC071/TauUnc (CC071) mice as very susceptible, with high mortality and high peak plasma

viral load. Notably, the CC071 strain has been recently reported as also susceptible to other

flaviviruses including dengue virus (15), Powassan virus (16), West Nile virus (WNV) (15), and

to Rift Valley fever virus (17), emphasizing its value to decipher genetic factors controlling host

responses to RNA viruses.

We previously demonstrated that genetic background influenced ZIKV replication in CC mouse

embryonic fibroblasts (MEFs). Here, we investigated the mechanisms driving high viral

replication in CC071 MEFs. We found that, compared with the more resistant C57BL/6J (B6)

and CC001/Unc (CC001) strains, ZIKV-infected CC071 MEFs displayed a delayed expression of

the IFN-I response genes. Genetic and functional analyses identified a strain-specific variant

in the *Irf3* gene, the first transcription factor involved in interferon (IFN) β expression, which

mimics the effects of a null allele *in vitro* and fully explains the delayed IFN-I expression and

uncontrolled viral replication. By contrast, we showed, from the *in vivo* comparison of CC071,

*Irf3*-deficient and backcross mice, that the CC071 *Irf3* mutation is not sufficient to explain the

high susceptibility of CC071 mice to ZIKV infection and that other genes, not associated with

the IFN-I response, are involved. These findings provide new insights into the roles of *Irf3* in

viral diseases and exemplify how the study of CC strains allows deciphering the role of host

genes in viral pathogenesis.

## Results

*CC071 MEFs show defective control of viral replication and delayed IFN-I expression, but normal response to IFN-I stimulation*

We previously reported that, unlike CC001 MEFs, CC071 MEFs produced increasing quantities of viral particles during the first 72 hours post-infection (hpi) (15). Here, we confirmed and expanded this observation by infecting B6, CC001 and CC071 MEFs and by quantifying viral particles by FFA. After ZIKV infection, CC071 MEFs displayed high and increasing viral titers between 24 and 72 hpi, while CC001 and B6 MEFs showed stable and lower titers (Fig 1A).

To investigate the origin of the defective control of viral replication in CC071 MEFs, we measured the expression level of the *Ifnb1* gene coding for IFNβ in ZIKV-infected CC071, CC001 and B6 MEFs. *Ifnb1* expression is induced very rapidly after virus detection by sensors and triggers the innate antiviral response which is essential for limiting viral replication. In CC001 and B6 MEFs, *Ifnb1* expression was significantly induced at 24 hpi and remained stable and high until at least 72 hpi (Fig 1B). In contrast, its expression in CC071 MEFs was low at 24 hpi and reached the level of CC001 only at 72 hpi. Similar results were obtained for *Ifn4a* which encodes one of the IFNα proteins (Sup Fig 1). Notably, *Ifnb1* expression in CC071 MEFs at 72 hpi was significantly higher than in B6 MEFs, showing that CC071 MEFs were delayed but not intrinsically hampered in their ability to induce strong IFN-I expression.

To test whether this defective induction of *Ifnb1* expression was specific to ZIKV infection, MEFs were then transfected with the influenza A virus-derived 3-phosphate-hairpin-RNA (3p-hpRNA), an agonist of the RIG-I/MDA5-MAVS pathway, or treated with polyinosine-polycytidylic acid (poly (I:C)), that activates both Toll-like receptor (TLR) 3 (TLR3) and the RIG-I/MDA5-MAVS pathway (18). Here again, CC071 MEFs showed a delayed expression of *Ifnb1* after both stimulations by comparison with B6 and CC001 MEFs (Sup Fig 2), indicating that the

defect in IFN-I genes expression in CC071 MEFs was not specific to ZIKV infection. This result

suggested a defect in the molecular cascade between cellular sensors of pathogen-associated

molecular patterns (PAMP) and the *Ifnb1* gene transcription machinery.

To evaluate the capacity of CC071 MEFs to respond to IFN-I, they were treated with

recombinant IFNα. The expression of IFN-stimulated genes (ISGs) such as *Ifitm3* was induced

with the same kinetics and level as in B6 and CC001 MEFs (Fig 1C), showing that CC071 MEFs

are able to respond normally to IFN-I stimulation and that their defect is limited to the

induction of *Ifnb1* gene expression.

*CC071's delayed Ifnb1 expression is strain-specific*

To gain insight into the mechanisms responsible for defective *Ifnb1* induction, we investigated

the expression levels of all genes involved in *Ifnb1* expression on CC071, B6 and CC001 MEFs

at 16, 24 and 32 hpi. Mock-infected MEFs were analyzed at 24 hours as controls. Expression

levels were measured by RNA sequencing (RNAseq) which provided a comprehensive analysis

of transcriptomic changes. In CC001 MEFs, the expression of many genes rapidly increased

after infection (160 at 16h, 821 at 24h and 971 at 32h; log2 fold-change > 1, FDR = 0.05),

reflecting a robust innate antiviral response (Sup Fig 3A). A similar pattern was observed in B6

MEFs. By contrast, the expression of only 38 genes was increased in CC071 MEFs at 32hpi (34

of which were also activated in CC001), consistent with the delayed induction of *Ifnb1*

expression. Among the genes that are involved in the pathway between PAMP sensors and

*Ifnb1* transcription, ISGs such as *Tlr3*, *Ddx58* (coding for RIG-I sensor) or *Irf7* were not activated

upon infection in CC071, while constitutively expressed genes such as *Mavs*, *Ticam1* (coding

for the TRIF adaptor), *Traf3* or *Irf3*, showed comparable levels of expression in the three

strains (Sup Fig 3B). Therefore, this analysis did not provide new clues for identifying the gene

responsible for the defect observed in CC071.

We then leveraged the genetic architecture of the CC which genomes are patchworks of

haplotypes inherited from the eight founder strains (19). Although CC071 was the only strain

with severe ZIKV disease, we hypothesized that, if the delayed activation of *Ifnb1* resulting in

uncontrolled viral replication observed in CC071 MEFs was due to an allele at one of the genes

involved in the *Ifnb1* induction pathway inherited from a parental strain, ZIKV-infected MEFs

of CC strains carrying the same allele would present similarly high viral titers. We therefore

derived MEFs from each CC strain available to us carrying the same ancestral haplotype as

CC071 at one of the 13 genes of the pathway (Fig 2A). Upon ZIKV infection, none of these CC

MEFs showed viral titer kinetics resembling that observed in CC071 MEFs (Fig 2B). These

results suggested two alternative hypotheses. Either the delayed *Ifnb1* activation involved two

members of the pathway with a CC071-specific allelic combination leading to a non-functional

interaction, or CC071 was carrying a strain-specific allele at one of these genes, resulting from

a mutation proper to CC071 that probably arose on an ancestral haplotype during the CC071

inbreeding. However, the sequencing of one male of each CC strain reported in 2017 (20) did

not identify such "private" variants with high predicted impact in CC071 for any of these genes.

Whatever the molecular mechanism, our results indicated that it was specific to CC071.

*Genetic analysis identifies Irf3 as a candidate gene in a haplotype shared between CC071 and*

*CC001*

We then turned to a genetic mapping approach. We first established that (CC001xCC071)F1

MEFs responded to infection with as rapid induction of *Ifnb1* expression as CC001 MEFs (Sup

Fig 4), suggesting that this CC071 trait was recessively inherited. F1 mice were therefore

backcrossed with CC071. MEFs were produced from each of 51 backcross (BC) embryos,

infected with ZIKV, and analyzed for viral titer and *Ifnb1* expression as above. One CC001 and

one CC071 MEF lines were included in each infection experiment as controls. BC MEFs

displayed either rapid and high *Ifnb1* expression with low viral titer (like CC001), or high viral

titer and delayed *Ifnb1* expression (like CC071, Fig 3A-B), showing that these two traits

correlated across the BC diverse genetic backgrounds.

To confirm this apparently binary distribution, we conducted linear discriminant analysis (LDA)

on CC001 and CC071 MEFs using *Ifnb1* expression at the three time points as variables.

Applying the LDA coefficients to backcross MEFs data classified individuals either in a CC001-

like group (n=31; 61%) or in a CC071-like group (n=20; 39%), with a mean probability of

prediction of 0.975 and 0.991, respectively (Fig 3C). Quantitative trait locus (QTL) mapping

was performed using LDA classification as a binary trait. Genome scan identified a peak on

chromosome 7 with a LOD (logarithm of the odd) score of 9.138 ($p < 0.001$, Fig 4A) located in

a region centered on the *Irf3* gene which, given its main role in the regulation of *Ifnb1*

expression, appeared as an obvious candidate. However, both CC001 and CC071 inherited the

CAST/EiJ haplotype in this region (Fig 4B), strongly suggesting that CC071's susceptibility was

caused by a variant proper to this strain.

*Abnormal splicing of Irf3 mRNA in CC071 leads to a loss of IRF3 transcriptional function*

To identify the CC071-specific mutation, we re-analyzed the RNAseq data and investigated the

splicing events between *Irf3* exons. As shown in Fig 5A, no splicing was observed between

exons 6 and 7 in CC071 MEFs, while a short cryptic exon was added to exon 6 (red box). This

aberrant splicing resulted in an mRNA lacking the last two exons. Notably, exon 8 encodes the

serin-rich region of the protein with the phosphorylation sites necessary for IRF3 activation

and nuclear translocation leading to *Ifnb1* transcription (Fig 5B). Neither long-range PCRs nor

sequencing could identify the exact nature of *Irf3* genetic alteration in CC071 but suggested

the insertion of a repeated sequence between exons 6 and 7. Nevertheless, the functional

consequence of this mutation was confirmed by Western blot using a specific C-terminal IRF3

antibody which showed that full-length IRF3 protein was absent in CC071 MEFs (Fig 5C).

Moreover, immunofluorescence using an antibody directed against phosphorylated IRF3

detected a positive signal in the nucleus of many ZIKV-infected CC001 and B6 MEFs, but not

in CC071 MEFs (Fig 5D). Altogether, these results show that CC071 carries a mutation in *Irf3*

that prevents IRF3 phosphorylation which is required to induce *Ifnb1* expression. Whether the

altered mRNA sequence prevents the production of the protein or alters its activity, this

mutation results in the loss of IRF3 transcriptional function.

CC071 *Irf3 mutation is responsible for uncontrolled viral replication in MEFs*

To test if the delayed *Ifnb1* expression resulting in uncontrolled viral replication in CC071 MEFs

was caused exclusively by the *Irf3* mutation, we performed a quantitative complementation

test by producing compound heterozygous MEFs carrying a knockout *Irf3* allele ($Irf3^{KO}$) and a

CC071 allele ($Irf3^{71}$). These $Irf3^{KO/71}$ MEFs were compared with CC071 and B6-$Irf3^{KO/KO}$ MEFs,

and with heterozygous MEFs carrying a B6 wildtype allele and either an $Irf3^{KO}$ or an $Irf3^{71}$ allele

($Irf3^{+/KO}$ or $Irf3^{+/71}$, respectively, Fig 6A). While $Irf3^{+/KO}$ and $Irf3^{+/71}$ ZIKV-infected MEFs showed

the same pattern as CC001 or B6 MEFs (rapid induction of *Ifnb1* expression and controlled

viral replication, see Fig 1A for comparison), CC071 ($Irf3^{71/71}$), B6-$Irf3^{KO/KO}$ and $Irf3^{KO/71}$ MEFs

carrying two defective alleles at *Irf3* showed similar results (Fig 6B-C). These data demonstrate

that, since the $Irf3^{KO}$ did not complement the $Irf3^{71}$ allele, the *Irf3* mutation in CC071

contributes to the defects observed in ZIKV-infected MEFs. Moreover, since the data obtained

on CC071 and on *Irf3*$^{KO/71}$ MEFs were identical, we conclude that the CC071 *Irf3* mutation is

sufficient to induce the defects observed in CC071 MEFs.

*CC071 Irf3 mutation is not sufficient to explain susceptibility in vivo*

We investigated whether this *Irf3* mutation was also responsible for the high susceptibility to

ZIKV of CC071 mice. We first evaluated its effects in a context allowing IFN-I response. To this

aim, we compared B6, B6-*Irf3* KO and CC071 mice infected without prior MAR1-5A3

treatment, which results in moderate and short-lasting plasma viral loads without clinical

signs, as we previously reported (15). Similar viral loads were observed in B6 and B6-*Irf3* KO

mice at all days p.i. (dpi), while CC071 mice showed significantly higher viral loads at days 1

and 2 p.i. (Fig 7A). This result indicates that, unlike in MEFs, the *Irf3* null mutation does not

induce elevated viral replication *in vivo* (as measured by plasma viral load) and suggests that

CC071 mice carry susceptibility alleles at other loci.

We then blocked IFN-I response by pre-treating mice with MAR1-5A3 and compared peak

plasma viral load and clinical signs in ZIKV-infected B6, B6-*Irf3* KO and CC071 mice. While no

differences were observed between B6 and B6-*Irf3* KO mice (moderate plasma viral loads

without clinical signs), as expected if the IFN-I response is neutralized, CC071 mice developed

significantly higher viral loads at day 2 p.i. and signs of disease (ruffled fur, hunched posture

and body weight loss) around 7 days p.i. (Fig 7B). This contrast could result either from

functional differences between the *Irf3*$^{KO}$ and *Irf3*$^{71}$ alleles or from CC071 alleles at other genes

not linked with the IFN-I response. F1 mice between CC071 and B6-*Irf3* KO mice gave similar

results as B6-*Irf3* KO mice, suggesting that CC071 susceptibility alleles were recessive.

Therefore, we generated a small cohort of (CC071 x B6-*Irf3* KO) x CC071 backcross mice and

genotyped them for the *Irf3* gene by PCR using primers that amplify intron 6 of the *Irf3*$^{KO}$ allele,

but not of the *Irf3^{71}* allele. (Fig 5A and Fig S5). We found no differences in plasma viral load

and clinical scores between *Irf3^{KO/71}* and *Irf3^{71/71}* backcross mice, indicating that the two alleles

are functionally equivalent *in vivo*. Moreover, *Irf3^{KO/71}* backcross mice were more severely

affected than *Irf3^{KO/71}* F1 mice, indicating the strong effects of recessive alleles from CC071

which likely explain the variability of plasma viral load and clinical signs among backcross mice.

Altogether, our results demonstrate that the CC071 mutation is a loss-of-function (LOF)

variant which does not confer susceptibility to ZIKV disease in the absence of other recessive

alleles carried by CC071 at genes not associated with the IFN-I response.

**Discussion**

An increasing number of publications have reported how the genetic diversity provided by the CC translates into an extended range of resistant and susceptible phenotypes in various infectious models. Since our first description of the high susceptibility of CC071 mice to ZIKV infection, with disease severity and peak plasma viral load almost as high as those of B6-*Ifnar1* KO mice, and higher clinical signs and mortality than 129-*Ifnar1* KO mice (15), several groups have reported similar observations for other viruses. High viral load was found in CC071 mice after dengue infection (15). Lethality after infection with West Nile virus (15) or Powassan virus (16) were also described, although there is likely contribution from the defective *Oas1b* allele that CC071 has inherited from 129S1/SvImJ. CC071 was also one of the most susceptible CC strains to Rift Valley Fever virus infection (17) and to hepacivirus with long-term viral persistence (21). Because of its unique genetic background, this strain will be increasingly useful for infectious diseases studies, which underlines the importance of deciphering the mechanisms of its susceptibility to each pathogen.

In this study, we leveraged from our previous observation that CC071 MEFs failed to control ZIKV replication by comparison with resistant strains. Although they cannot recapitulate the complex interactions between multiple pathways, cell types and tissues of a whole organism, MEFs are a convenient cellular model which can be easily derived from any mouse genetic background. We identified a delayed activation of the *Ifnb1* gene in CC071 MEFs, resulting in delayed stimulation of ISGs. We then used a combination of genetic approaches to find the causative gene defect. The observation that MEFs were normally responsive to IFN-I stimulation was consistent with the higher susceptibility of mice treated by the MAR1-5A3 antibody compared with untreated mice which did not develop symptoms and showed much lower plasma viral load (15). These results pointed at the pathway from PAMP sensors to *Ifnb1*

transcription factors. Transcriptional analysis did not identify reduced expression of non-ISGs

of this pathway. CC strains' genomic structure allows searching for haplotypes inherited from

the same founder in CC strains showing similar phenotypes. However, we did not identify such

haplotypes for genes of the *Ifnb1* induction pathway, suggesting that the defects observed in

CC071 MEFs were strain specific. It is finally the analysis of MEFs derived from backcross

embryos that established a monogenic inheritance, and genetic linkage unambiguously

pointed at the causative *Irf3* gene. The MEF experimental model was particularly appropriate

since we could derive cell lines from every backcross embryo. Our RNAseq data showed an

RNA splicing defect in the *Irf3* gene in CC071, which functional consequences could be

validated *in vitro*. The formal proof that the CC071 *Irf3* mutation was necessary and sufficient

to cause *Ifnb1* delayed activation and uncontrolled viral replication came from a quantitative

complementation test in which the $Irf3^{KO}$ allele was combined either with a CC071 or a B6

(functional) allele. Altogether, these results show that the *Irf3* defect in CC071 abrogates its

transcriptional activity. Unfortunately, none of the antibodies recognizing the N-terminal

protein domain that we tested worked in our hands in western blot. Therefore, we could not

establish whether the *Irf3* mRNA detected by RNAseq is translated into an abnormal protein

or not. However, the *in vivo* observation that $Irf3^{KO/71}$ and $Irf3^{71/71}$ backcross mice displayed

the same phenotypes upon ZIKV infection shows that the two alleles are functionally

equivalent in this context.

*Irf3* maps to a chromosomal region that CC071 inherited from CAST/EiJ, like the resistant

CC001 strain (and CC046/Unc, not available to us). The CC071-specific mutation may have

arisen during the inbreeding generations leading to this strain. Alternatively, the mutation

might have been segregating in the CAST/EiJ founders and transmitted by chance only to the

CC071 breeding funnel. Since all CAST/EiJ founders used in CC crosses originated from The

Jackson Laboratory colony, genotyping past and present breeders of this colony could not only

unravel the origin of the mutation but also allow eliminating this mutation to avoid unwanted

effects.

Notably, this is not the first example of a CC-strain-specific mutation. We (22) and others (23)

previously reported the extreme susceptibility of CC042/GeniUnc (CC042) to *Salmonella*

Typhimurium and to *Mycobacterium tuberculosis*, respectively, as a consequence of a *de novo*

15-nucleotide deletion in the *Itgal1* gene. In the case of *Salmonella* Typhimurium, CC042 was

standing out, with bacterial loads up to 1000 times higher than other CC and the susceptible

B6 strains. In our previous study on ZIKV, CC071 was the most susceptible strain, but its peak

viral load was just the highest in a continuous distribution of values. Additionally, this is not

the first example of a *Irf3* spontaneous variant identified in mice modulating susceptibility to

bacterial (24) or viral infections (25).

*Irf3* is an important transcription factor involved in the innate immune response. It is

constitutively expressed and, at rest, inactive IRF3 is present in the cytoplasm. Upon viral entry

(or other stimuli that activate TLRs such as TLR3 and 4 or RIG-I-like receptors), signal

transduction leads to the phosphorylation of IRF3, resulting in its dimerization and

translocation to the nucleus where it binds to the *Ifnb1* promoter (26). This mechanism leads

to the very fast production of IFN$\beta$ which is secreted by the cell and triggers the immediate

response to viral infection through the activation of ISGs with diverse antiviral functions (27).

In addition to its transcriptional activity, IRF3 has been shown to counter viral infection by two

additional mechanisms. IRF3 induces apoptosis through the RLR-induced IRF3-mediated

pathway of apoptosis (RIPA) in infected cells (28) and inhibits NF-κB-mediated inflammation

through the repression of IRF3-mediated NF-κB activity (RIKA) (29). These newly described

roles of IRF3 depend on other protein domains than those involved in its transcriptional

activity. Whether these activities are also abrogated by the CC071 *Irf3* mutation remains to be

established.

In mice, studies that used *Irf3*-deficient cellular models have consistently reported decreased

IFN-I production and/or increased viral replication. For example, after infection with WNV,

viral replication was increased in B6-*Irf3* KO bone marrow macrophages (BMMs) and

moderately in primary neurons (30). Higher viral replication was observed in herpes simplex

virus 1 (HSV-1)-infected B6-*Irf3* KO bone marrow-derived dendritic cells (BMDCs) and BMMs,

and IFNβ was reduced in BMDCs supernatants (31). Similarly, *Ifnb1* expression was reduced in

HSV-1-infected (32) and CHIKV-infected (33) B6-*Irf3* KO MEFs. In line with these findings, our

study provides, to our knowledge, the first evidence for a role of *Irf3* in the infection of murine

cells by ZIKV. Moreover, we recently reported that primary cultured neurons (PCNs) show a

delayed activation of the *Ifnb1* expression upon ZIKV exposure compared with MEFs, and that

this delay is even longer in PCNs derived from CC071 compared with CC001 (34). The

identification of the *Irf3* mutation in CC071 provides an explanation for this observation,

although it would require confirmation in B6-*Irf3* KO PCNs.

In contrast, *in vivo* studies have reported inconsistent consequences of *Irf3* deficiency

between viral infections. Here, we did not observe any differences in clinical signs nor in

plasma viral loads between B6 and B6-*Irf3* KO mice (pre-treated or not with the MAR1-5A3

antibody), consistently with a previous study that reported neither mortality, nor body weight

loss after ZIKV infection (10). Likewise, B6-*Irf3* KO mice have been reported to show no

mortality and low virus load in the circulation following dengue virus (35) and CHIKV infection

(33). Contrastingly, WNV infection was lethal in all B6-*Irf3* KO mice while 65% of infected WT

mice survived (30). Differences were also associated with the route of infection. For example,

all WT and B6-*Irf3* KO mice survived after intravenous inoculation with HSV-1 (32), while

intranasal inoculation led to 90% mortality in B6-*Irf3* KO but only 30% in WT mice (36).In

humans, LOF mutations in *Irf3* have been associated with increased susceptibility to diseases

caused by WNV (37,38), HSV-1 (39,40) and more recently by SARS-CoV-2 (41). Discrepancies

between the consequences of *Irf3* deficiency in *in vitro* and *in vivo* experiments likely reflect

differences between a single cell type model with an intrinsic capacity to mount a more or less

IRF3-dependent IFN-I response, and a multicellular organism in which the paracrine effect of

IFN-I produced by the most reactive cells induces protection of other, less reactive, cell types.

In fact, we have recently reported such cooperation between neurons and microglia cells (34).

Our *in vivo* data, showing high plasma viral loads and clinical signs in all CC071 and half of

backcross mice after MAR1-5A3 treatment, demonstrated that the high susceptibility of

CC071 mice requires the contribution of other recessive alleles. Since their effect was

observed in mice which IFNAR receptor had been blocked, we conclude that their mode of

action is not dependent on an intact IFN-I response. Moreover, the comparison between the

results of the complementation test on MEFs, with CC071 and *Irf3*$^{KO/71}$ yielding very similar

data, and the *in vivo* observation that CC071 showed higher viremia than *Irf3*$^{KO/71}$ F1 mice,

suggests that these other susceptibility alleles may not act at the cell level, at least during the

first 72 hpi, but rather at a more integrated or systemic level. Whether or not these alleles

require *Irf3* deficiency to induce the susceptibility observed in the CC071 mice has yet to be

determined. While our limited understanding of the severe disease developing in CC071 ZIKV-

infected mice does not point at potential mechanisms, identifying these alleles will allow

disentangling the genetic factors controlling ZIKV and other viruses' pathogenesis.

375 Material and methods
376

*Mice and crosses*

C57BL/6J (B6) mice were purchased from Charles River Laboratories France. Collaborative

Cross strains (CC001/Unc, CC071/TauUnc, CC005/TauUnc, CC011/Unc, CC026/GeniUnc,

CC061/GeniUnc, CC021/Unc, CC006/TauUnc, CC025/GeniUnc, CC039/Unc, CC060/Unc) were

purchased from the Systems Genetics Core Facility, University of North Carolina and bred at

the Institut Pasteur. *Irf3 Irf7* double KO mice (C57BL/6J-*Bcl2l12/Irf3$^{tm1Ttg}$ Irf7$^{tm1Ttg}$*,(32,42))

were bred at the Institut Pasteur and backcrossed to B6 mice to generate *Irf3* single KO mice

(B6-*Irf3* KO ). Genetic mapping was performed on MEFs derived from (CC001 x CC071) x CC071

and CC071 x (CC001 x CC071) backcross embryos. For the quantitative complementation test,

MEFs were derived from CC071, B6-*Irf3* KO and (B6-*Irf3$^{+/KO}$* x CC071) embryos.. *In vivo*

experiments were performed on B6, B6-*Irf3* KO, CC071, (CC071 x *Irf3* KO) F1 and CC071 x

(CC071 x *Irf3* KO)backcross mice. All crosses are described as female x male. All mice were

maintained as described previously (15). Mouse experiments were approved by the Institut

Pasteur Ethics Committee and authorized by the French Ministry of Research (project #19469),

in compliance with French and European regulations.

*ZIKA virus*

The FG15 Asian Zika virus (ZIKV) strain, isolated from a patient during ZIKV outbreak in French

Guiana in December 2015, was obtained from the Virology Laboratory of the Institut Pasteur

of French Guiana. Viral stock (passage 5) was prepared from supernatant of infected C6/36

cells, clarified by centrifugation at 800g and titrated on Vero cells by focus-forming assay.

*Mouse infection*

All infection experiments were performed in a biosafety level 3 animal facility and mice were

kept in isolators. Six- to 10-week-old, male or female mice were injected intraperitoneally with

$10^7$ PFU of ZIKV FG15. In some *in vivo* experiments, mice received an IP injection of 2 mg of

MAR1-5A3 anti-IFNAR antibody (Euromedex, Cat#BX-BE0241) one day prior infection. Mouse

numbers are indicated in figure legends. Both males and females were used since no

differences between sexes were detected in our previous and present experiments. Clinical

signs and body weight loss were recorded for up to seven days post infection. Blood samples

were collected on EDTA from the retromandibular vein for plasma viral load assessment.

Quantification of ZIKV viral copies by qPCR was previously described (15).

*MEFs isolation*

Pregnant females were euthanized at day 13.5-15.5 of gestation. For B6, CC001, CC071 and

crosses used for genetic mapping and the complementation test, MEFs were isolated from

individual fetuses to obtain biological replicates. For other CC strains, MEFs were derived from

individual or pooled fetuses. Fetus bodies were chopped and digested with trypsin (Gibco

Cat#25300054), then cultured at 37°C and 5% $CO_2$ in complete medium (DMEM Gibco Cat#

31966047, 10% fetal bovine serum PAA Laboratories Cat#A15-101, 1% penicillin/streptomycin

Sigma Cat#P4333). MEFs were used until passage 2. For the backcross experiment, MEF lines

were isolated from 51 backcross fetuses. Heads were used to prepare DNA for whole-genome

genotyping.

*MEFs infection*

MEFs were seeded at $5.10^4$ cells per well in 24-well plates the day before infection. They were

exposed to ZIKV FG15 strain at a MOI of 5 for 2 hours after which the inoculum was replaced

with fresh complete medium and MEFs were incubated for up to 72 hours. For kinetics studies,

different wells were used for each time point. Backcross MEFs were infected in 6 infection

experiments, each of which included one CC001 and one CC071 MEF lines.

*MEFs IFNα stimulation*

MEFs were seeded at $5.10^4$ cells per well in 24-well plates one day before stimulation and

treated with 300 IU/mL IFNα (Miltenyi Biotec Cat#130-093-131) and incubated for up to 24

hours.

*Focus forming assay*

Quantification of ZIKV particles was performed by focus forming assay on Vero cells (ATCC

CRL-1586) as previously described (15).

*RNA extraction from cells*

MEFs were lysed in 350µL of RLT buffer (Qiagen) with 1% β-mercaptoethanol. RNA was

extracted using RNeasy Mini Kit (Qiagen Cat#74104) according to the manufacturer's

instructions, with addition of DNase I (Qiagen Cat# 79254) to prevent genomic DNA

contamination.

*Reverse transcription and qPCR*

Reverse-transcription was performed on 200ng of RNA using Superscript II polymerase

(Invitrogen Cat#18064022) and RNaseOUT ribonuclease inhibitor (Invitrogen Cat#10777019).

qPCR was performed on 20ng of cDNAs using Power SYBR Green PCR Master Mix (Applied

Biosystems Cat#4367659) and 6pmol of each primer, on a QuantStudio 12K Flex or a ViiA 7

(ThermoFisher Scientific). Primers (Eurofins) sequences are provided in Sup Table 1. Gene

expression was expressed on a Log10 scale of relative expression to the reference *Tbp* gene.

*Genotyping*

Genomic DNA was prepared from backcross fetuses' heads by proteinase K digestion, phenol-

chloroform extraction and ethanol precipitation according to standard protocols. Whole-

genome genotyping was performed at Neogen (Neogen/Geneseek, Inc, Lincoln, NE, USA)

using the MiniMUGA array containing 11,125 SNP markers. For the quantitative

complementation test, *Irf3* and *Irf7* genotyping was performed by Transnetyx (Cordova, TN)

by real-time PCR on fetuses' heads. (CC071 x *Irf3* KO) x CC071 backcross individuals were

genotyped by PCR using primers amplifying the intron 6 of $Irf3^+$ and $Irf3^{KO}$ but not $Irf3^{71}$ alleles.

Primers (Eurofins) sequences are provided in Sup Table 1.

*RNA splicing analysis from RNA sequencing data*

RNA sequencing data were produced as described in the Supplementary methods. Splicing

analysis of the *Irf3* gene was performed using Majiq 2.4 (43) with default parameters to

investigate alternative transcripts between genotypes.

*Immunofluorescence*

MEFs were plated on glass coverslips before infection, fixed with 4% paraformaldehyde for 20

min and permeabilized with 100% methanol for 10 min at -20°C. Cells were incubated with

blocking buffer (5% FBS 0.3% triton in PBS) for 1 hour, with primary antibodies diluted in

antibody incubation buffer (AIB: 1% BSA 0.3% triton in PBS) overnight at 4°C and with

secondary antibodies and Hoechst (dilution 1:1000) diluted in AIB for 1 hour. Coverslips were

mounted on slides and imaged with a widefield microscope (Zeiss Axio Observer.Z1 with a

Plan-Apochromat 20x/0.8 M27 objective and a Hamamatsu sCMOS ORCA-Flash 4.0 v3

camera). ZEN blue 2012 software (ZEISS) imaging software was used for image capture and

Image J software (National Institutes of Health) to adjust brightness and contrast. Primary and

secondary antibodies are indicated in Sup Table 2.

*Western blot*

MEFs were trypsinized for 5 min, washed and lysed in a protein extraction buffer (10mM

TrisHCl pH7.5, 5mM EDTA, 150mM NaCl, 30mM $Na_2HPO_4$, 50mM NaF, 10% glycerol, 1% NP40,

1X cOmplete (Roche #11873580001), 1X PhosSTOP (Roche #4906845001), 1/1000 benzonaze

(Sigma Cat#E1014)) for 30 min at 4°C. Proteins diluted in Laemmli were resolved on 4-12% Bis-

Tris gels (Invitrogen Cat#NP0323BOX) in MOPS buffer (Invitrogen Cat#NP0001) and

transferred to nitrocellulose membranes (Bio-Rad Cat#1620112) in a 25mM Tris 200mM

glycine 20% ethanol buffer. Blots were blocked in 5% milk in TBS-T (0.1% Tween20 in Tris Base

Sodium), incubated with primary antibodies diluted in 3% milk in TBST overnight at 4°C, and

incubated with secondary antibodies diluted in 3% milk in TBST for 90 min. Blots were revealed

with ECL substrate (Thermo Scientific Cat#32132) and imaged with X-ray films. Primary and

secondary antibodies are indicated in Sup Table 2.

*Statistical analysis*

Statistical analyses were performed using R version 4.1.0 (44). Viral titers, gene expression

and genome copies were log-transformed for graphs and statistical tests. One way ANOVA

followed by Tukey HSD were used for testing multiple comparisons. For *in vivo* studies without

MAR1-5A3 treatment, non-parametric Kruskal-Wallis followed by Wilcoxon tests with Holm

correction for multiple testing were used to handle values below the limit of detection. Linear

discriminant analysis (LDA) was conducted using the MASS package (45). The LDA was trained

on the phenotypes of the two parental CC001 and CC071 strains from each infection batch.

LDA coefficients were applied to backcross mice for assignment to "CC071-like" or "CC001-

like" groups.

*Genetic analysis*

Raw genotypes were curated using the stuart package (46). QTL mapping was performed using

R/qtl (47). LDA prediction was used as a binary trait. Statistical significance thresholds were

computed by data permutation (n=1000). 95% confidence interval was estimated using the

Bayesian method.

**Acknowledgements**

We thank Matthieu Prot, Maxime Chazal and Sandrine Vandormael-Pournin for technical

advice and for providing reagents. We are grateful to Tommy Penel and Rachid Chennouf of

the Institut Pasteur Central Animal Facility of the C2RA (Center for Animal Resources and

Research) for the careful breeding of CC strains and for the maintenance of mice in the BSL-3

animal facility, respectively. We thank Etienne Simon-Loriere and Nolwenn Jouvenet for

scientific advice and continuous support along this project.

**Author contributions**

517 X.M. conceived and supervised this study. M.B. and C.M. designed and performed the

518 experiments and analyzed data. L.C. and C.R.P. performed experiments. E.K. analyzed RNAseq

519 data. X.M. designed experiments and analyzed data. E.B. provided conceptual advice. M.B.,

520 C.M. and X.M. wrote the paper. All authors commented on and edited the manuscript.

**Funding**

523 This work was supported by grants from the Agence Nationale de la Recherche (ANR)

524 NeuroZika (ANR-20-CE16-0017) and from the French Government's Investissement d'Avenir

525 program, Laboratoire d'Excellence: IBEID (Integrative Biology of Emerging Infectious Diseases,

526 ANR-10-LABX-62-IBEID). C.M. and M.B. were supported by doctoral fellowships from grant

527 ANR-10-LABX-62-IBEID.

**Availability of data and materials**

530    RNAseq primary data are available from the European Nucleotide Archive (accession number

531    E-MTAB-12765). All other experimental data that support the finding of this study are

532    available from the corresponding authors upon reasonable request.

534    **Competing interests**

535    The authors declare that they have no competing interests.

**Figure legends**

**Figure 1. CC071 MEFs fail to control viral replication, with delayed *Ifnb1* expression but normal response to IFN-I.** (A, B) MEFs derived from B6 (black circles), CC001 (blue triangles) and CC071 (red squares) were infected with ZIKV at a MOI of 5 and analyzed 24, 48 and 72 hpi. (A) Viral titer in supernatants was quantified by FFA. (B) *Ifnb1* expression was normalized to *Tbp* reference gene. Data are mean +/- sem from 3 to 4 biological replicates per strain (MEFs derived from individual embryos). (C) MEFs were stimulated with recombinant IFNα. *Ifitm3* relative expression normalized to *Tbp* reference gene is shown as an example of ISG. Data are mean +/- sem from 2 to 3 biological replicates. Blue asterisks and black hashes show statistical significance of CC071 compared to CC001 and to B6, respectively (ANOVA followed by Tukey HSD, */# p < 0.05, **/## p < 0.01, ***/### p < 0.001).

**Figure 2. Haplotype analysis fails to identify a gene from the *Ifnb1* induction pathway associated with uncontrolled viral replication.** (A) Identification of CC strains which carry the same ancestral haplotype as CC071 at the genes involved in the pathway leading to *Ifnb1* expression. Colored boxes indicate matched haplotypes between CC071 and other CC strains. Letters and colors designate the eight CC founder strains. A: A/J (yellow); B: C57BL/6J; C: 129S1/SvImJ (pink); D: NOD/ShiLtJ; E: NZO/HILtJ (light blue); F: CAST/EiJ (green); G: PWK/PhJ (red); H: WSB/EiJ (purple). Doubled letters (eg AA) indicate homozygous genotypes. Heterozygous genotypes are indicated by the two corresponding letters (eg AD). (B) Kinetics of viral titer in MEFs from CC071 and the 8 CC strains shown in (A). Experimental conditions were as in Figure 1A. n: number of technical replicates for each strain. Data are mean +/- sem

from the technical replicates. Letters show statistical significance between CC071 and other

strains. (ANOVA followed by Tukey HSD, a: p < 0.05, b: p < 0.01, c: p < 0.001).

**Figure 3. Backcross MEFs display either a CC001-like or a CC071-like phenotype.** MEFs

derived from CC001 (blue triangles), CC071 (red squares) and backcross (gray circles) embryos

were infected with ZIKV at a MOI of 5. (A) Viral titer and (B) *Ifnb1* expression in ZIKV-infected

MEFs from 9 backcross embryos. Experimental conditions were as in Figure 1A-B. Red and

blue curves show the results for CC071 and CC001 MEFs, respectively, from the same infection

experiment. (C) Results of LDA on the backcross MEFs. LDA coefficients were calculated from

*Ifnb1* expression data in CC001 and CC071 infected MEFs, and applied to backcross MEFs. The

graph shows the probability of each BC MEF to belong to the "CC001-like" group, resulting in

two distinct populations shown in blue ("CC001-like") and in red ("CC071-like"). n: number of

BC MEFs assigned to each group.

**Figure 4. Genetic analysis of *Ifnb1* expression in backcross MEFs identifies a major**

**determinant mapping to the *Irf3* locus.** (A) Genome-wide linkage analysis of the LDA

classification of backcross MEFs performed with R/qtl. X-axis: genomic location. Y-axis: LOD

score of the phenotype-genotype association. Genome-wide significance thresholds (P = 0.05,

P = 0.1 and P = 0.63) were computed from 1000 data permutation. The chromosome 7 peak

has a LOD score of 9.138. (B) Schematic representation of CC001 and CC071 chromosome 7

haplotypes, from https://csbio.unc.edu/CCstatus/CCGenomes/. Colors represent the CC

ancestral haplotypes (same colors as in Figure 2A). Thick vertical black lines show the peak's

95% Bayesian confidence interval (25.9-31.3 cM, corresponding to 40.1-50.6 Mb). The red line

shows the position of *Irf3* gene.

**Figure 5. CC071 *Irf3* mRNAs show abnormal splicing, resulting in defective IRF3 protein.** (A) Schematic representation of the exons of the *Irf3* gene with the number of reads spanning successive exons in the CC001 and CC071 RNAseq data (one sample of each strain). The red box between CC071's exons 6 and 7 depicts a novel exon resulting from abnormal splicing. (B) Schematic representation of the IRF3 protein structural domains (exon 1 is untranslated). Exon 8 encodes the serine rich region containing the phosphorylation sites for IRF3 activation. (C) Western blot using an anti-C-terminal IRF3 antibody from mock-infected and ZIKV-infected B6, CC001 and CC071 MEFs at 2 hpi, showing the absence of full-length IRF3 in CC071 MEFs. Vinculin was used as a loading control. (D) Immunofluorescence using an anti-phosphorylated IRF3 (pIRF3, green) in ZIKV-infected B6, CC001 and CC071 MEFs at 24 hpi, showing the absence of pIRF3 in the nucleus of CC071 MEFs upon infection. Red-labeled 4G2 antibody labels ZIKV-infected cells. Cell nuclear DNA labeled by Hoechst (blue). Quantification of the number of infected and pIRF3 positive cells is presented in the table. Proportions were established on 420, 428 and 551 cells for CC001, CC071 and B6, respectively.

**Figure 6. Quantitative complementation test confirms that *Irf3* LOF in CC071 is responsible for uncontrolled viral replication and delayed *Ifnb1* expression.** (A) Mice heterozygous for an inactivated *Irf3* allele (B6-*Irf3*$^{+/KO}$) were mated with CC071 mice to produce four types of embryos from which MEFs were derived. The genetic background and *Irf3* genotype is shown below each type of embryos. (B) Viral titer and (C) *Ifnb1* expression in ZIKV-infected MEFs. Experimental conditions were as in Figure 1A-B. Data are mean +/- sem from 6 biological replicates for CC071 and 3 for the other groups. The asterisks represent the results of ANOVA test between all groups (* $p < 0.05$). Results of the Tukey HSD post-hoc are as follows. Viral

titer at 72h: CC071 vs *Irf3^{+/71}* and vs *Irf3^{+/KO}*: $p < 0.001$. *Irf3^{KO/71}* vs *Irf3^{+/71}* and vs *Irf3^{+/KO}*: $p <$

$0.01$. *Irf3^{KO/KO}* vs *Irf3^{+/71}* and vs *Irf3^{+/KO}*: $p < 0.05$. *Ifnb1* expression at 24hpi: CC071 vs *Irf3^{+/71}*

and vs *Irf3^{+/KO}*, *Irf3^{KO/71}* vs *Irf3^{+/71}* and vs *Irf3^{+/KO}*, *Irf3^{KO/KO}* vs *Irf3^{+/71}* and vs *Irf3^{+/KO}*: $p < 0.001$.

*Ifnb1* expression at 48hpi: CC071 vs *Irf3^{+/71}*and vs *Irf3^{+/KO}*, *Irf3^{KO/71}* vs *Irf3^{+/71}* and vs *Irf3^{+/KO}*,

*Irf3^{KO/KO}* vs *Irf3^{+/71}* and vs *Irf3^{+/KO}*: $p < 0.01$.

**Figure 7. *Irf3* LOF is not sufficient to explain CC071 susceptibility to ZIKV infection *in vivo*.**

(A) B6, B6-*Irf3* KO and CC071 mice were infected IP with $10^7$ FFUs of ZIKV and monitored for 7

days, without prior IP injection of MAR1-5A3 IFNAR-blocking monoclonal antibody. Plasma

viral load was quantified at days 1 to 3 p.i. by RT-qPCR. The two lines of the group legend (X-

axis) indicate the genetic background of the mice and their genotype at the *Irf3* locus,

respectively. Each dot represents one mouse. Groups were compared by Kruskal-Wallis

followed by Wilcoxon test with Holm correction for multiple testing (* $p < 0.05$). (B) B6, B6-

*Irf3* KO, CC071, (CC071 x *Irf3* KO) F1 and (CC071 x *Irf3* KO) x CC071 BC mice were infected IP

with $10^7$ FFUs of ZIKV after IP injection of 2 mg of MAR1-5A3 IFNAR-blocking monoclonal

antibody 24 h before infection, and monitored for 7 days. The graph shows plasma viral loads

quantified at day 2 p.i. by RT-qPCR. The two lines of the group legend (X-axis) indicate the

genetic background of the mice and their genotype at the *Irf3* locus, respectively. BC mice are

separated into two groups depending on the genotype at the *Irf3* gene: homozygous for the

CC071 mutant allele (*Irf3^{71/71}*), or heterozygous for the CC071 and the KO alleles (*Irf3^{KO/71}*). *Irf3*

genotyping results are presented in Figure S5*.* Each dot represents one mouse. Groups were

compared by ANOVA followed by Tukey HSD (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$). Below

the graph is shown the distribution of clinical scores at day 7 p.i. in the same groups of mice

as above (0, no symptoms; 1, slight hunched posture; 2, ruffled fur, hunched posture and/or

mild ataxia; 3: prostration, ataxia, partial paralysis).**Supplemental information**

**Legend to supplemental figures**

**Figure S1.** *Ifna4* **expression is delayed in ZIKV-infected CC071 MEFs.** MEFs derived from B6 (gray circles), CC001 (blue triangles) and CC071 (red squares) were infected with ZIKV at a MOI of 5. *Ifna4* expression was determined by RT-qPCR on MEFs total RNA by normalizing to *Tbp* housekeeping gene. Data are mean +/- sem from 3 biological replicates. For one CC001, one B6 and one CC071 replicates at 0 hpi and one CC071 replicate at 24 hpi, gene expression was below the limit of detection. Blue asterisks and black hashes show statistical significance of CC071 compared to CC001 and to B6, respectively (ANOVA followed by post-hoc Tukey HSD, */# $p < 0.05$, *** $p < 0.001$).

**Figure S2. Delayed IFN-I expression in CC071 MEFs results from a constitutive defect in the transcription activation cascade.** MEFs were transfected with either (A) poly(I:C), which activates both TLR and RIG-I pathways, or (B) 3p-hpRNA, a RIG-I agonist. *Ifnb1* expression was determined as in Figure 1B. Data are mean +/- sem from 3 biological replicates for CC001 and CC071 (2 at 4 hours) or 2 biological replicates for B6 (1 at 4 hours). Blue asterisks and black hashes show statistical significance of CC071 compared to CC001 and to B6, respectively (ANOVA followed by post-hoc Tukey HSD, */# $p < 0.05$).

**Figure S3**. **Expression of genes involved in the *Ifnb1* induction pathway in ZIKV-infected MEFs of B6, CC001 and CC071 strains.** MEFs derived from B6 (black circles), CC001 (blue triangles) and CC071 (red squares) were infected with ZIKV at a MOI of 5. mRNA expression levels were measured by RNAseq in non-infected (NI) and ZIKV-infected MEFs. (A) Number of upregulated

genes per strain at 16, 24 and 32 hpi (log2 fold-change > 1, FDR = 0.05). (B-G) Genes

constitutively expressed. (H-N) Genes which expression is induced by the IFN-I response

(ISGs). Expression levels are shown on a logarithmic scale. For *Ifnb1* expression, null counts

were transformed to 1. Data are mean +/- sem from 3 biological replicates. Blue asterisks and

black hashes show statistical significance of CC071 compared to CC001 and to B6, respectively

(ANOVA followed by post-hoc Tukey HSD, */# p < 0.05, **/## p < 0.01, ***/### p < 0.001).

**Figure S4. (CC001xCC071)F1 MEFs show normal induction of *Ifnb1*.** *Ifnb1* expression upon

ZIKV infection in CC001, CC071 and (CC001xCC071)F1 MEFs determined as in Figure 1B. Data

are mean +/- sem from 3 biological replicates. Blue asterisks and orange hashes show

statistical significance of CC071 compared to CC001 and to F1, respectively (ANOVA followed

by post-hoc Tukey HSD, **/## p < 0.01).

**Figure S5. Genotyping of (CC071 x Irf3 KO) x CC071 backcross individuals.** Backcross

individuals were genotyped by PCR with primers amplifying the intro 6 of Irf3 (sequences in

Sup Table 1). The *Irf3^{KO}* allele results in a band at 877pb while the *Irf3^{71}* allele results in no

band. Individuals 1, 5, 11, 14, 15, 16, 17, 19, 21, 22 and 23 are *Irf3^{KO/71}* and individuals 2, 6, 7,

8, 9, 10, 12, 13 and 24 are *Irf3^{71/71}*.

**References**

1. Cao-Lormeau VM, Blake A, Mons S, Lastère S, Roche C, Vanhomwegen J, et al.
Guillain-Barré Syndrome outbreak associated with Zika virus infection in French
Polynesia: a case-control study. Lancet. 9 avr 2016;387(10027):1531‑9.

2. Carod-Artal FJ. Neurological complications of Zika virus infection. Expert Review
of Anti-infective Therapy. 4 mai 2018;16(5):399‑410.

3. Oliveira Melo AS, Malinger G, Ximenes R, Szejnfeld PO, Alves Sampaio S, Bispo
de Filippis AM. Zika virus intrauterine infection causes fetal brain abnormality and
microcephaly: tip of the iceberg?: Physician Alert. Ultrasound Obstet Gynecol. janv
2016;47(1):6‑7.

4. Paixao ES, Cardim LL, Costa MCN, Brickley EB, de Carvalho-Sauer RCO, Carmo
EH, et al. Mortality from Congenital Zika Syndrome — Nationwide Cohort Study in
Brazil. N Engl J Med. 24 févr 2022;386(8):757‑67.

5. Rossi SL, Ebel GD, Shan C, Shi PY, Vasilakis N. Did Zika Virus Mutate to Cause
Severe Outbreaks? Trends in Microbiology. 1 oct 2018;26(10):877‑85.

6. Liu ZY, Shi WF, Qin CF. The evolution of Zika virus from Asia to the Americas. Nat
Rev Microbiol. mars 2019;17(3):131‑9.

7. Manet C, Roth C, Tawfik A, Cantaert T, Sakuntabhai A, Montagutelli X. Host
genetic control of mosquito-borne Flavivirus infections. Mamm Genome.
2018;29(7‑8):384‑407.

8. Rasmussen AL, Okumura A, Ferris MT, Green R, Feldmann F, Kelly SM, et al. Host
genetic diversity enables Ebola hemorrhagic fever pathogenesis and resistance.
Science. 21 nov 2014;346(6212):987‑91.

9. Gralinski LE, Menachery VD, Morgan AP, Totura AL, Beall A, Kocher J, et al. Allelic
Variation in the Toll-Like Receptor Adaptor Protein Ticam2 Contributes to SARS-
Coronavirus Pathogenesis in Mice. G3 (Bethesda). 7 juin 2017;7(6):1653‑63.

10. Lazear HM, Govero J, Smith AM, Platt DJ, Fernandez E, Miner JJ, et al. A Mouse
Model of Zika Virus Pathogenesis. Cell Host Microbe. 11 mai 2016;19(5):720‑30.

11. Grant A, Ponia SS, Tripathi S, Balasubramaniam V, Miorin L, Sourisseau M, et al.
Zika Virus Targets Human STAT2 to Inhibit Type I Interferon Signaling. Cell Host
& Microbe. juin 2016;19(6):882‑90.

12. Sheehan KCF, Lai KS, Dunn GP, Bruce AT, Diamond MS, Heutel JD, et al.
Blocking Monoclonal Antibodies Specific for Mouse IFN-$\alpha$/$\beta$ Receptor Subunit 1
(IFNAR-1) from Mice Immunized by *In Vivo* Hydrodynamic Transfection. Journal of
Interferon & Cytokine Research. nov 2006;26(11):804‑19.

13. Roberts A, Pardo-Manuel de Villena F, Wang W, McMillan L, Threadgill DW. The
polymorphism architecture of mouse genetic resources elucidated using genome-

wide resequencing data: implications for QTL discovery and systems genetics.
Mamm Genome. juill 2007;18(6-7):473-81.

14. Keane TM, Goodstadt L, Danecek P, White MA, Wong K, Yalcin B, et al. Mouse
genomic variation and its effect on phenotypes and gene regulation. Nature. sept
2011;477(7364):289-94.

15. Manet C, Simon-Lorière E, Jouvion G, Hardy D, Prot M, Conquet L, et al. Genetic
Diversity of Collaborative Cross Mice Controls Viral Replication, Clinical Severity,
and Brain Pathology Induced by Zika Virus Infection, Independently of Oas1b. J
Virol     [Internet].     17     janv     2020;94(3).     Disponible     sur:
https://journals.asm.org/doi/10.1128/JVI.01034-19

16. Jasperse BA, Mattocks MD, Noll KM, Ferris MT, Heise MT, Lazear HM.
Neuroinvasive flavivirus pathogenesis is restricted by host genetic factors in
Collaborative Cross mice, independently of Oas1b [Internet]. bioRxiv; 2022 [cité 26
oct                 2022].                 Disponible                 sur:
https://www.biorxiv.org/content/10.1101/2022.10.24.513634v1

17. Cartwright HN, Barbeau DJ, Doyle JD, Klein E, Heise MT, Ferris MT, et al. Genetic
diversity of collaborative cross mice enables identification of novel rift valley fever
virus encephalitis model. PLOS Pathogens. 14 juill 2022;18(7):e1010649.

18. McCartney S, Vermi W, Gilfillan S, Cella M, Murphy TL, Schreiber RD, et al. Distinct
and complementary functions of MDA5 and TLR3 in poly(I:C)-mediated activation
of    mouse    NK    cells.    Journal    of    Experimental    Medicine.    21    déc
2009;206(13):2967-76.

19. Collaborative Cross Consortium. The Genome Architecture of the Collaborative
Cross    Mouse    Genetic    Reference    Population.    Genetics.    1    févr
2012;190(2):389-401.

20. Srivastava A, Morgan AP, Najarian ML, Sarsani VK, Sigmon JS, Shorter JR, et al.
Genomes of the Mouse Collaborative Cross. Genetics. 1 juin 2017;206(2):537-56.

21. Brown AJ, Won JJ, Wolfisberg R, Fahnøe U, Catanzaro N, West A, et al. Host
genetic variation guides hepacivirus clearance, chronicity, and liver fibrosis in mice
[Internet]. Microbiology; 2023 mars [cité 24 mars 2023]. Disponible sur:
http://biorxiv.org/lookup/doi/10.1101/2023.03.18.533278

22. Zhang J, Teh M, Kim J, Eva MM, Cayrol R, Meade R, et al. A Loss-of-Function
Mutation in the Integrin Alpha L (Itgal) Gene Contributes to Susceptibility to
Salmonella enterica Serovar Typhimurium Infection in Collaborative Cross Strain
CC042. Infect Immun. 17 2019;88(1).

23. Smith CM, Proulx MK, Lai R, Kiritsy MC, Bell TA, Hock P, et al. Functionally
Overlapping Variants Control Tuberculosis Susceptibility in Collaborative Cross
Mice. Vance R, éditeur. mBio. 24 déc 2019;10(6):e02791-19.

24. Garifulin O, Qi Z, Shen H, Patnala S, Green MR, Boyartchuk V. Irf3 Polymorphism
Alters Induction of Interferon Beta in Response to Listeria monocytogenes
Infection. PLOS Genetics. 7 sept 2007;3(9):e152.

25. Moore TC, Al-Salleeh FM, Brown DM, Petro TM. IRF3 polymorphisms induce
different innate anti-Theiler's virus immune responses in RAW264.7 macrophages.
Virology. sept 2011;418(1):40-8.

26. Bourdon M, Manet C, Montagutelli X. Host genetic susceptibility to viral infections:
the role of type I interferon induction. Genes Immun. déc 2020;21(6-8):365-79.

27. Schoggins JW. Interferon-Stimulated Genes: What Do They All Do? Annu Rev
Virol. 29 sept 2019;6(1):567-84.

28. Chattopadhyay S, Kuzmanovic T, Zhang Y, Wetzel JL, Sen GC. Ubiquitination of
the Transcription Factor IRF-3 Activates RIPA, the Apoptotic Pathway that Protects
Mice from Viral Pathogenesis. Immunity. 17 mai 2016;44(5):1151-61.

29. Popli S, Chakravarty S, Fan S, Glanz A, Aras S, Nagy LE, et al. IRF3 inhibits
nuclear translocation of NF-κB to prevent viral inflammation. Proc Natl Acad Sci
USA. 13 sept 2022;119(37):e2121385119.

30. Daffis S, Samuel MA, Keller BC, Gale M Jr, Diamond MS. Cell-specific IRF-3
responses protect against West Nile virus infection by interferon-dependent and -
independent mechanisms. PLoS Pathog. 27 juill 2007;3(7):e106.

31. Menachery VD, Leib DA. Control of Herpes Simplex Virus Replication Is Mediated
through an Interferon Regulatory Factor 3-Dependent Pathway. J Virol. déc
2009;83(23):12399-406.

32. Honda K, Yanai H, Negishi H, Asagiri M, Sato M, Mizutani T, et al. IRF-7 is the
master regulator of type-I interferon-dependent immune responses. Nature. 7 avr
2005;434(7034):772-7.

33. Schilte C, Buckwalter MR, Laird ME, Diamond MS, Schwartz O, Albert ML. Cutting
Edge: Independent Roles for IRF-3 and IRF-7 in Hematopoietic and
Nonhematopoietic Cells during Host Response to Chikungunya Infection. JI. 1 avr
2012;188(7):2967-71.

34. Manet C, Mansuroglu Z, Conquet L, Bortolin V, Comptdaer T, Segrt H, et al. Zika
virus infection of mature neurons from immunocompetent mice generates a
disease-associated microglia and a tauopathy-like phenotype in link with a delayed
interferon beta response. J Neuroinflammation. 20 déc 2022;19(1):307.

35. Chen HW, King K, Tu J, Sanchez M, Luster AD, Shresta S. The roles of IRF-3 and
IRF-7 in innate antiviral immunity against dengue virus. J Immunol. 15 oct
2013;191(8):4194-201.

36. Canivet C, Rhéaume C, Lebel M, Piret J, Gosselin J, Boivin G. Both IRF3 and
especially IRF7 play a key role to orchestrate an effective cerebral inflammatory
response in a mouse model of herpes simplex virus encephalitis. J Neurovirol. déc
2018;24(6):761-8.

37. Bigham AW, Buckingham KJ, Husain S, Emond MJ, Bofferding KM, Gildersleeve
H, et al. Host genetic risk factors for West Nile virus infection and disease
progression. PLoS One. 2011;6(9):e24745.

38. Cahill ME, Conley S, DeWan AT, Montgomery RR. Identification of genetic variants
associated with dengue or West Nile virus disease: a systematic review and meta-
analysis. BMC Infect Dis. déc 2018;18(1):282.

39. Mork N, Kofod-Olsen E, Sorensen KB, Bach E, Orntoft TF, Ostergaard L, et al.
Mutations in the TLR3 signaling pathway and beyond in adult patients with herpes
simplex encephalitis. Genes Immun. déc 2015;16(8):552-66.

40. Andersen LL, Mork N, Reinert LS, Kofod-Olsen E, Narita R, Jorgensen SE, et al.
Functional IRF3 deficiency in a patient with herpes simplex encephalitis. J Exp
Med. 24 août 2015;212(9):1371-9.

41. Zhang Q, Bastard P, Liu Z, Le Pen J, Moncada-Velez M, Chen J, et al. Inborn errors
of type I IFN immunity in patients with life-threatening COVID-19. Science
[Internet].        24        sept        2020;        Disponible        sur:
https://www.ncbi.nlm.nih.gov/pubmed/32972995
https://science.sciencemag.org/content/sci/370/6515/eabd4570.full.pdf

42. Sato M, Suemori H, Hata N, Asagiri M, Ogasawara K, Nakao K, et al. Distinct and
essential roles of transcription factors IRF-3 and IRF-7 in response to viruses for
IFN-alpha/beta gene induction. Immunity. oct 2000;13(4):539-48.

43. Vaquero-Garcia J, Barrera A, Gazzara MR, González-Vallinas J, Lahens NF,
Hogenesch JB, et al. A new view of transcriptome complexity and regulation
through the lens of local splicing variations. eLife. 1 févr 2016;5:e11752.

44. R Core Team. R: A Language and Environment for Statistical Computing [Internet].
2021. Disponible sur: https://www.R-project.org/

45. Venables WN, Ripley BD. Modern Applied Statistics with S. Fourth Edition
[Internet].        Fourth.        Springer;        2002.        Disponible        sur:
https://www.stats.ox.ac.uk/pub/MASS4/

46. Bourdon M, Montagutelli X. stuart: an R package for the curation of SNP genotypes
from experimental crosses. G3 (Bethesda). 24 août 2022;jkac219.

47. Broman KW, Wu H, Sen S, Churchill GA. R/qtl: QTL mapping in experimental
crosses. Bioinformatics. 1 mai 2003;19(7):889-90.

**A**  Viral titer

**B**  *Ifnb1*

**C**  *Ifitm3* post IFNα stimulation

Figure 1

**A**

## CC strains haplotypes

| Gene | Protein | CC071 | CC001 | CC005 | CC026 | CC021 | CC006 | CC025 | CC039 | CC060 |
|------|---------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| *Ddx58* | RIG-I | HH | BB | EE | CC | HH | AA | HH | AD | EE |
| *Ifih1* | LGP2 | GG | AA | DD | HH | GG | BB | AA | BB | CC |
| *Dhx58* | MDA−5 | FF | EE | DD | HH | FF | DD | FF | FF | EE |
| *Tlr3* | TLR3 | AA | AA | BB | CC | EE | AA | BB | FF | CC |
| *Mavs* | MAVS | CC | AA | DD | HH | GG | CC | DD | HH | DD |
| *Ticam1* | TRIF | CC | AA | EE | BB | CC | DD | HH | AA | EE |
| *Azi2* | NAP1 | EE | DD | FF | BB | FF | EE | GG | CC | EE |
| *Traf3* | TRAF3 | HH | HH | GG | CC | FF | CC | CC | AA | BB |
| *Tbk1* | TBK1 | HH | DD | GG | HH | AA | FF | AA | GG | BB |
| *Ikbke* | IKKε | CC | EE | CC | AA | EE | CC | CC | GH | AA |
| *Irf3* | IRF3 | FF | FF | HH | BB | CC | DD | EE | HH | BB |
| *Irf7* | IRF7 | HH | BB | DD | DD | HH | EE | DD | HH | DD |
| *Ifnb1* | IFNβ | HH | EE | DD | CC | HH | BB | HH | CC | DE |

**B**

Figure 2

Figure 3

Figure 4

**A**

CC001:

| 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 |

66  122  81  76  58  69  68

CC071:

| 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 |

53  38  32  32  23  26  0  15

Number of splicing counts

**B**

IRF3

| 2 | 3 | 4 | 5 | 6 | 7 | 8 |

DNA binding domain (DBD)  Regulatory domain (RD)
Nuclear export signal (NES)  Serine rich region (SRR)

**C**

B6  CC001  CC071
mock  2hpi  mock  2hpi  mock  2hpi

130kDa — Vinculin

55kDa — IRF3 Cter

**D**

B6  CC001  CC071

Hoechst/pIRF3/ZIKV

30µm

| Count | B6 | CC001 | CC071 |
|---|---|---|---|
| Infected cells | 24 | 91 | 76 |
| pIRF3 positive cells | 5 | 28 | 0 |
| **Percentage** | 21% | 31% | 0% |

Figure 5

**A**

CC071 → CC071 *Irf3*<sup>71/71</sup>

(B6xCC071)F1 *Irf3*<sup>KO/71</sup>

(B6xCC071)F1 *Irf3*<sup>+/71</sup>

*Irf3*<sup>+/KO</sup> → B6 *Irf3*<sup>+/KO</sup>

*Irf3* KO → B6 *Irf3*<sup>KO/KO</sup>

**B** Viral titer

Log$_{10}$ FFU/mL

Hours post infection
24    48    72

**C** *Ifnb1*

Log$_{10}$ relative expression

Hours post infection
24    48    72

■ *Irf3*<sup>71/71</sup> (n=6)  ▲ *Irf3*<sup>+/71</sup> (n=3)  ◆ *Irf3*<sup>KO/KO</sup> (n=3)

◆ *Irf3*<sup>KO/71</sup> (n=3)  ● *Irf3*<sup>+/KO</sup> (n=3)

Figure 6

Figure 7

**Supplementary methods**

*MEFs stimulation with poly(I :C) and 3p-hpRNA*

For intracellular stimulation with Poly(I:C) or with 3p-hpRNA, MEFs were seeded at $1.10^5$ cells per well in 12-well plates the day before stimulation, transfected with 1 µg/mL Poly(I:C) (InvivoGen Cat#vac-pic) or 0.5 µg/mL 3p-hpRNA (InvivoGen Cat#tlrl-hprna) using 5 µL Lipofectamine LTX and 1 µL for poly(I:C) stimulation or 0.5 µL for 3p-hpRNA stimulation of Plus Reagent (ThermoFischer Scientific Cat#15338100), according to the manufacturer's instructions. After stimulation, MEFs were incubated for 8 to 24 hours.

*RNA sequencing*

MEF RNA was prepared as described in the main text. RNA integrity and quantification were assessed using the RNA Nano 6000 Assay Kit of the Bioanalyzer 2100 system (Agilent Technologies, CA, USA). One microgram of high-quality RNA samples (RIN > 9.2) representing biological triplicates were submitted to Novogene for RNA-sequencing (Novogene Beijing, China). Poly-A selected RNA was used for paired-end library preparation and transcriptome sequencing. Sequencing libraries were generated using NEBNext® UltraTM RNA Library Prep Kit for Illumina® (NEB, USA) following manufacturer's instructions. The library preparations were sequenced on an Illumina platform and paired-end reads were generated. The RNA-seq analysis was performed with Sequana 0.9.8 (1). In particular, we used the RNA-seq pipeline https://github.com/sequana/sequana_rnaseq) built on top of Snakemake 6.1.1 (2). Briefly, reads were trimmed from adapters using Cutadapt 2.7 then mapped to the *Mus musculus* genome assembly GCA_000001635.8 from NCBI using STAR 2.7.3a (3). FeatureCounts 1.6.4 (4) was used to produce the count matrix, assigning reads to features using corresponding annotation v92 from NCBI with strand-specificity information. Quality control statistics were summarized using MultiQC 1.6 (5). Clustering of transcriptomic profiles were controlled using Principal Component Analysis (PCA). Differential expression testing was conducted using DESeq2 library 1.24.0 (6) scripts indicating the significance (Benjamini-Hochberg adjusted p-values, false discovery rate FDR < 0.05) and the effect size (fold-change) for each comparison.

1. Cokelaer T, Desvillechabrol D, Legendre R, Cardon M. 2017. "Sequana": a Set of Snakemake NGS pipelines. JOSS 2:352.

2. Köster J, Rahmann S. 2012. Snakemake—a scalable bioinformatics workflow engine. Bioinformatics 28:2520–2522.

3. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR. 2013. STAR: ultrafast universal RNA-seq aligner. Bioinformatics 29:15–21.

4. Liao Y, Smyth GK, Shi W. 2014. featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. Bioinformatics 30:923–930.

5. Ewels P, Magnusson M, Lundin S, Käller M. 2016. MultiQC: summarize analysis results for multiple tools and samples in a single report. Bioinformatics 32:3047–3048.

6. Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biol 15:550.

Figure S1

Figure S2

Figure S3

Figure S4

Figure S5

| Gene | Forward | Reverse |
|------|---------|---------|
| qPCR | | |
| *Ifnb1* | CAGCTCCAAGAAAGGACGAAC | GGCAGTGTAACTCTTCTGCAT |
| *Ifna4* | TGATGAGCTACTACTGGTCAGC | GATCTCTTAGCACAAGGATGGC |
| ZIKA *Env* | CCGCTGCCCAACACAAG | CCACTAACGTTCTTTTGCAGACAT |
| *Ifitm3* | ACTGTGATCAACATGCCCAGAG | CTTCCGATCCCTAGACTTCACG |
| *Tbp* | AGAACAATCCAGACTAGCAGC | GGGAACTTCACATCACAGCTC |
| PCR genotyping | | |
| *Irf3* | CACTCCCACGCCTTCTGG | ACAAGCCTCTTGACCCATGG |

S1 table

**Immunofluorescence antibodies**

| Primary/secondary | Antibody | Species | Dilution | Reference |
|---|---|---|---|---|
| Primary | ZIKV envelop protein | Mouse | 1:200 | 4G2 antibody from ATCC hybridoma |
| Primary | Phosphorylated IRF3 | Rabbit | 1:250 | Cell Signaling Technology Cat#29047 |
| Secondary | Anti-rabbit AlexaFluor-488-conjugated IgG | Goat | 1:500 | Invitrogen Cat#A-11034 |
| Secondary | Anti-mouse AlexaFluor-594-conjugated IgG | Donkey | 1:500 | Invitrogen Cat#A-21203 |

**Western blot antibodies**

| Primary/secondary | Antibody | Species | Dilution | Reference |
|---|---|---|---|---|
| Primary | IRF3 Cter | Rabbit | 1:1000 | Cell Signaling Cat#4302 |
| Primary | Vinculin | Rabbit | 1:1000 | Proteintech Cat# 26520-1-AP |
| Secondary | Rabbit IgG HRP Linked Whole Ab | Donkey | 1:10,000 | Sigma Cat# GENA934-100UL |

S2 table