Title: Collaborative Cross strains reveal that *Irf3* deficiency results in

uncontrolled Zika virus replication *in vitro* but is insufficient to induce disease

susceptibility in mice.

Running title: *Irf3* deficiency promotes ZIKV replication

Marie Bourdon[§1], Caroline Manet[§1], Laurine Conquet[1], Corentin Ramaugé-Parra[1], Etienne

Kornobis[2], Eliette Bonnefoy[3], Xavier Montagutelli[#1]

[1]Institut Pasteur, Université de Paris, Mouse Genetics Laboratory, F-75015 Paris, France

[2]Institut Pasteur, Université de Paris, Biomics, F-75015 Paris, France

[3]Université Paris Cité, Institut Cochin, Inserm, CNRS, F-75014 Paris, France

[§] M.B. and C. M. equally contributed to this work and are co-first authors

[#] Corresponding author: xavier.montagutelli@pasteur.fr

Keywords: Zika virus, mouse, Irf3, interferon response, Collaborative Cross, susceptibility

Abstract

Zika virus (ZIKV) is a Flavivirus responsible for recent epidemics in Pacific Islands and in the

Americas. In humans, the consequences of ZIKV infection range from asymptomatic infection

to severe neurological disease such as Guillain-Barré syndrome or fetal neurodevelopmental

defects, suggesting the influence of host genetic factors. We previously reported similar

diverse outcomes of ZIKV infection in mice of the Collaborative Cross, a collection of inbred

strains with large genetic diversity. CC071 was the most susceptible strain with severe

symptoms and lethality. Here, we investigated viral replication in mouse embryonic fibroblasts

from CC071 and two resistant strains. CC071 showed enhanced viral replication associated

with delayed induction of type-I interferons (IFN-I). Using a combination of genetic

approaches, we identified a loss of function (LOF) mutation in the *Irf3* gene, specific to CC071,

and demonstrated that it fully explains the defective IFN-I response and uncontrolled viral

replication in CC071 MEFs. However, this mutation was not sufficient to induce the high

plasma viral load and clinical severity observed in CC071 mice, indicating the involvement of

other susceptibility genes *in vivo*. Considering the susceptibility of the CC071 strain to multiple

viruses, our results provide new insight into the role of *Irf3* in innate antiviral response.

Importance

Recent ZIKV outbreaks led to millions of infected people, with rare but severe complications

such as Guillain-Barré syndrome and encephalitis in adults suggesting host genetic control.

We previously reported the importance of host genes in ZIKV pathogenesis using a panel of

genetically diverse mouse strains and identified CC071 as the most susceptible strain. Through

a combination of functional and genetic approaches in a cellular model, we identified a

mutation in the *Irf3* gene which plays a key role in the expression of interferon beta. This

mutation fully explains the high viral replication observed in CC071 cells. However, it was not

sufficient to explain the symptoms displayed by CC071 ZIKV-infected mice, unraveling the

implication of other host genes. Since CC071 has been reported as a valuable model for several

viral infections, our results have implications beyond ZIKV infection and contribute to

shedding light on host mechanisms fighting infectious diseases.

Introduction

Zika virus (ZIKV) is a mosquito-borne virus of the *Flaviviridae* family identified in 1947 in Uganda. The first noticeable human outbreaks occurred in Micronesia in 2007 and in French Polynesia and New Caledonia in 2013-2014. In 2015-2016, ZIKV caused an epidemic in Brazil which rapidly spread across the Americas and the Caribbean. To date, 89 countries have reported evidence of mosquito-transmitted Zika virus infection (https://www.who.int/health-topics/zika-virus-disease#tab=tab_1).

While most people infected with ZIKV remain asymptomatic, some develop non-specific symptoms including rash, fever, conjunctivitis, muscle and joint pain, malaise and headache. Neurological complications have been described in adults such as Guillain-Barré syndrome (1) and encephalitis (2). Infection of pregnant women was associated with congenital Zika syndrome in the fetus, which can lead to neurodevelopmental deficiencies, brain malformation (3) or in some cases to fetal loss (4).

Many factors may contribute to this variable severity, including the viral strain, the infection route and dose, and the host genetic background (5, 6). Indeed, mouse and human studies have shown that host genes influence flaviviral infections' outcomes (7). While human genetic studies are hampered by the variability of these multiple factors, they can be controlled in mouse models which have proven very valuable to identify susceptibility variants (8, 9). Relevant ZIKV infection models have been developed in mice either using *Ifnar1* knock-out (KO) mice in which the IFNAR receptor to IFN-I has been inactivated (10, 11), or by blocking this receptor using a monoclonal antibody targeting the IFNAR1 receptor subunit (MAR1-5A3 (12)).

We have previously explored the role of mouse natural genetic variants on ZIKV susceptibility in the Collaborative Cross (CC), a panel of recombinant inbred mice encompassing a genetic diversity similar to that of the human population and capturing approximately 90% of the mouse natural genetic variants (13, 14). We reported that the CC genetic diversity enabled large variations in the clinical severity of ZIKV disease, plasma viral load and intensity of brain pathology (15). We specifically identified CC071 mice as very susceptible, with high mortality and high peak plasma viral load. We also demonstrated that genetic background influenced the control of ZIKV replication in CC mouse embryonic fibroblasts (MEFs).

Here, we investigated the mechanisms driving high viral replication in CC071 MEFs. We found that, compared with the more resistant C57BL/6J (B6) and CC001 strains, ZIKV-infected CC071

MEFs displayed a delayed expression of the IFN-I response genes. Genetic and functional

analyses identified a strain-specific, LOF variant in the *Irf3* gene, the main transcription factor

involved in IFN-I expression. *In vivo* comparison of CC071 and B6-*Irf3* KO mice showed that

Irf3 deficit is not sufficient to explain the high susceptibility of CC071 mice to ZIKV infection,

unraveling the plurality of genetic factors controlling susceptibility to viral infections. These

findings provide new insights on the role of *Irf3* in viral diseases and exemplify how the study

of CC strains allows deciphering the role of host genes in ZIKV and other viruses' pathogenesis.

Results

CC071 MEFs show defective control of viral replication and delayed IFN-I expression, but normal response to IFN-I stimulation

We previously reported that, unlike CC001 MEFs, CC071 MEFs produced increasing quantities of viral particles during the first 72 hours post-infection (hpi) (15). Here, we confirmed and expanded this observation by infecting B6, CC001 and CC071 MEFs and by quantifying viral particles by FFA. After ZIKV infection, CC071 MEFs displayed high and increasing viral titers between 24 and 72 hpi, while CC001 and B6 MEFs showed stable and lower titers (Fig 1A).

To investigate the origin of the defective control of viral replication in CC071 MEFs, we measured the expression level of the *Ifnb1* gene coding for IFNβ in ZIKV-infected CC071, CC001 and B6 MEFs. *Ifnb1* expression is induced very rapidly after virus detection by sensors and triggers the innate antiviral response which is essential for limiting viral replication. In CC001 and B6 MEFs, *Ifnb1* expression was significantly induced at 24 hpi and remained stable and high until at least 72 hpi (Fig 1B). In contrast, its expression in CC071 MEFs was low at 24 hpi and reached the level of CC001 only at 72 hpi. Similar results were obtained for *Ifn4a* which encodes one of the IFNα proteins (Sup Fig 1). Notably, *Ifnb1* expression in CC071 MEFs at 72 hpi was significantly higher than in B6 MEFs, showing that CC071 MEFs were delayed but not intrinsically hampered in their ability to induce a strong IFN-I expression.

To test whether this defective induction of *Ifnb1* expression was specific to ZIKV infection, MEFs were then transfected with the influenza A virus-derived 3-phosphate-hairpin-RNA (3p-hpRNA), an agonist of the RIG-I/MDA5-MAVS pathway, or treated with polyinosine-polycytidylic acid (poly (I:C)), that stimulates both Toll-like receptor 3 (TLR3) and the RIG-I/MDA5-MAVS pathway (16). Here again, CC071 MEFs showed a delayed expression of *Ifnb1* by comparison with B6 and CC001 MEFs (Sup Fig 2), indicating that the defect in IFN-I genes expression in CC071 MEFs was not specific to ZIKV infection. This result suggested a defect in the molecular cascade between cellular sensors of pathogen-associated molecular patterns (PAMP) and the *Ifnb1* gene transcription machinery.

To evaluate the capacity of CC071 MEFs to respond to IFN-I stimulation, they were treated with recombinant IFNα. The expression of IFN-stimulated genes (ISGs) such as *Ifitm3* was induced with the same kinetics and level as in B6 and CC001 MEFs (Fig 1C), showing that CC071

MEFs are able to respond normally to IFN-I stimulation and that their defect is limited to the induction of *Ifnb1* gene expression.

CC071's delayed Ifnb1 expression is strain-specific

To gain insight into the mechanisms responsible for defective *Ifnb1* induction, we investigated the expression levels of all genes involved in *Ifnb1* expression on CC071, B6 and CC001 MEFs at 16, 24 and 32 hpi. Mock-infected MEFs were analyzed at 24 hours as controls. Expression levels were measured by RNA sequencing (RNAseq) which provided a comprehensive analysis of transcriptomic changes. In CC001 MEFs, the expression of many genes rapidly increased after infection (160 at 16h, 821 at 24h and 971 at 32h; log2 fold-change > 1, FDR = 0.05), reflecting a robust innate antiviral response (Sup Fig 3A). A similar pattern was observed in B6 MEFs. By contrast, the expression of only 38 genes was increased in CC071 MEFs at 32hpi (34 of which were also activated in CC001), consistent with the delayed induction of *Ifnb1* expression. Among the genes that are involved in the pathway between PAMP sensors and *Ifnb1* transcription, ISGs such as *Tlr3*, *Ddx58* (coding for RIG-I sensor) or *Irf7* were not activated upon infection in CC071, while constitutively expressed genes such as *Mavs*, *Ticam1* (coding for the TRIF adaptor), *Traf3* or *Irf3*, showed comparable levels of expression in the three strains (Sup Fig 3B). Therefore, this analysis did not provide new clues for identifying the gene responsible for the defect observed in CC071.

We then leveraged the genetic architecture of the CC which genomes are patchworks of haplotypes inherited from the eight founder strains (17). Although CC071 was the only strain with severe ZIKV disease, we hypothesized that, if the delayed activation of *Ifnb1* resulting in uncontrolled viral replication observed in CC071 MEFs was due to an allele at one of the genes involved in the *Ifnb1* induction pathway inherited from a parental strain, ZIKV-infected MEFs of CC strains carrying the same allele would present similarly high viral titers. We therefore derived MEFs from each available CC strain carrying the same ancestral haplotype as CC071 at one of the 13 genes of the pathway (Fig 2A). Upon ZIKV infection, none of these CC MEFs showed viral titer kinetics resembling that observed in CC071 MEFs (Fig 2B). These results suggested two alternative hypotheses. Either the delayed *Ifnb1* activation involved two members of the pathway with a CC071-specific allelic combination resulting in non-functional interaction, or CC071 was carrying a strain-specific allele at one of these genes, resulting from a mutation that arose on an ancestral haplotype during the CC071 inbreeding. However, the

sequencing of one male of each CC strain reported in 2017 (18), did not identify such "private" variants with high predicted impact in CC071 for any of these genes. Whatever the molecular mechanism, our results indicated that it was specific to CC071.

Genetic analysis identifies Irf3 as a candidate gene in a haplotype shared between CC071 and CC001

We then turned to a genetic mapping approach. We first established that (CC001xCC071)F1 MEFs responded to infection with rapid induction of *Ifnb1* expression, as CC001 (Sup Fig 4), suggesting that the CC071 trait was recessively inherited. F1 mice were therefore backcrossed with CC071. MEFs lines were produced from 51 backcross (BC) embryos, infected with ZIKV, and analyzed for viral titer and *Ifnb1* expression as above. One CC001 and one CC071 MEF lines were included in each infection experiment as controls. BC MEFs displayed either high *Ifnb1* expression and low viral titer (like CC001), or high viral titer and delayed *Ifnb1* expression (like CC071, Fig 3A-B), showing that these two traits correlated across the BC diverse genetic backgrounds.

To confirm this apparently binary distribution, we conducted linear discriminant analysis (LDA) on CC001 and CC071 MEFs using *Ifnb1* expression at the three time points. Applying the LDA coefficients to backcross MEFs data classified individuals either in a CC001-like group (n=31; 61%) or in a CC071-like group (n=20; 39%), with a mean probability of 0.975 and 0.991, respectively (Fig 3C). Quantitative trait locus (QTL) mapping was performed using LDA classification as a binary trait. Genome scan identified a peak on chromosome 7 with a LOD (logarithm of the odd) score of 9.138 ($p < 0.001$, Fig 4A) located in a region centered on the *Irf3* gene which, given its main role in the regulation of *Ifnb1* expression, appeared as an obvious candidate. However, both CC001 and CC071 inherited the CAST/EiJ haplotype in this region (Fig 4B), strongly suggesting that CC071's defect was caused by a *de novo* mutation.

Abnormal splicing of Irf3 mRNA in CC071 leads to a loss of function

To identify the CC071-specific mutation, we re-analyzed the RNAseq data and investigated the splicing events between *Irf3* exons. As shown in Fig 5A, no splicing was observed between exons 6 and 7 in CC071 MEFs, while a short cryptic exon was added to exon 6 (red in Fig 5A). This aberrant splicing resulted in an mRNA lacking the last two exons. Notably, exon 8 encodes the serin-rich region of the protein with the phosphorylation sites necessary for IRF3

activation and nuclear translocation leading to *Ifnb1* transcription (Fig 5B). A combination of long-range PCRs and sequencing did not identify the exact nature of *Irf3* genetic alteration in CC071 but suggested the insertion of a repeated sequence between exons 6 and 7. Nevertheless, the functional consequence of this mutation was confirmed by Western blot using a specific C-terminal IRF3 antibody which showed that full-length IRF3 protein was absent in CC071 MEFs (Fig 5C). Moreover, immunofluorescence using an antibody directed against phosphorylated IRF3 detected a positive signal in the nucleus of many ZIKV-infected CC001 and B6 MEFs, but not in CC071 MEFs (Fig 5D). Altogether, these results show that CC071 carries a mutation in *Irf3* that prevents IRF3 dimerization and phosphorylation which are required to induce *Ifnb1* expression. Whether the altered mRNA sequence prevents the production of the protein or alters its activity, this mutation results in IRF3 loss of function.

Irf3 LOF is responsible for uncontrolled viral replication in CC071

To test if the delayed *Ifnb1* expression resulting in uncontrolled viral replication in CC071 MEFs was caused exclusively by *Irf3* LOF, we performed a quantitative complementation test by producing compound heterozygous MEFs carrying a knockout *Irf3* allele ($Irf3^{KO}$) and the allele from CC071 (denoted $Irf3^{KO}/Irf3^{71}$). These MEFs were compared with CC071 MEFs and with heterozygous MEFs carrying a B6 wildtype allele and either an $Irf3^{KO}$ or an $Irf3^{71}$ allele ($Irf3^{+}/Irf3^{KO}$ or $Irf3^{+}/Irf3^{71}$, respectively, Fig 6A). While $Irf3^{+}/Irf3^{KO}$ and $Irf3^{+}/Irf3^{71}$ ZIKV-infected MEFs showed the same pattern as CC001 or B6 MEFs (rapid induction of *Ifnb1* expression and controlled viral replication, see Fig 1A for comparison), CC071 and $Irf3^{KO}/Irf3^{71}$ MEFs carrying two defective alleles at *Irf3* showed similar results (Fig 6B-C). These data demonstrate that, since the $Irf3^{KO}$ did not complement the $Irf3^{71}$ allele, the *Irf3* mutation in CC071 contributes to the defects observed in ZIKV-infected MEFs. Moreover, since the data obtained on CC071 and on $Irf3^{KO}/Irf3^{71}$ MEFs were identical, we conclude that the *Irf3* mutation is sufficient to induce the defects observed in CC071 MEFs.

Irf3 LOF is not sufficient to explain CC071's susceptibility in vivo

Lastly, we investigated whether the *Irf3* LOF was also responsible for the high susceptibility to ZIKV of CC071 mice. We first compared clinical signs and plasma viral load in ZIKV-infected B6, B6-$Irf3^{KO/KO}$ and CC071 mice pre-treated with MAR1-5A3. While no differences were observed between B6 and B6-$Irf3^{KO/KO}$ (no clinical signs and moderate plasma viral loads), as expected

under conditions of neutralization of the IFN-I response, CC071 mice developed symptoms

(ruffled fur, hunched posture and body weight loss) around 7 days p.i. (Fig 7A), and

significantly higher viral loads at days 2 and 6 p.i. (Fig 7B) , indicating that genes not associated

to the IFN-I response play a major role in the susceptibility of CC071 mice to ZIKV infection.

We then measured plasma viral load in mice of the same genotypes infected without MAR1-

5A3 treatment. CC071 mice showed significantly higher viral loads at days 1 and 2 p.i., while

similar values were again observed in B6 and B6-*Irf3*^{KO/KO} mice (Fig 7C). These results show

that, unlike at the cell level, *Irf3* LOF does not result in elevated viral replication in infected

mice (as measured by plasma viral load), either with or without IFNAR blockade, and therefore

is not sufficient to explain the high susceptibility of CC071 mice, suggesting the involvement

of other host genes.

Discussion

Among the CC strains we tested, CC071 mice pre-treated with anti-IFNAR antibody were the most susceptible to ZIKV infection, with disease severity and peak plasma viral load almost as high as those of B6-*Ifnar1* KO mice, and higher clinical signs and mortality than 129-*Ifnar1* KO mice (15). In this susceptible model, IFN-I response is not permanently and ubiquitously abrogated as in *Ifnar1* deficient strains, but delayed, temporarily blocked and not in tissues unreached by the antibody (probably such as the brain). It is therefore particularly appropriate to identify host susceptibility genes. However, traits measured *in vivo* such as viremia, symptoms and death result from complex interactions between multiple pathways and cell types, from which distinct mechanisms are difficult to dissect. Cultured MEFs offer a lower level of complexity although they cannot provide the level of integration of a whole organism. To decipher the susceptibility of CC071 mice, we used MEFs as a convenient cellular model which can be easily derived from any mouse genetic background.

Starting from the confirmed observation of uncontrolled viral replication in CC071 MEFs, we identified a delayed activation of the *Ifnb1* gene resulting in delayed stimulation of ISGs and used a combination of genetic approaches to find the causative gene defect. The observation that MEFs were normally responsive to IFN-I stimulation was consistent with the higher susceptibility of mice treated by the MAR1-5A3 antibody compared with untreated mice (15). These results pointed at the pathway from PAMP sensors to *Ifnb1* transcription factors. Transcriptional analysis did not identify reduced expression of non-ISGs of this pathway. CC strains' genomic structure allows searching for haplotypes inherited from the same founder in CC strains showing similar phenotypes. However, we did not identify such haplotypes for genes of the *Ifnb1* induction pathway, suggesting that the defects observed in CC071 MEFs were strain-specific. It is finally the analysis of MEFs derived from backcross embryos that established a monogenic inheritance, and genetic linkage unambiguously pointed at the causative *Irf3* gene. The MEF experimental model was particularly appropriate since we could derive cell lines from every backcross embryo. Our RNAseq data informed on the RNA splicing defect in the *Irf3* gene, which functional consequences could be validated *in vitro*. The formal proof that the CC071 *Irf3* mutation was necessary and sufficient to cause *Ifnb1* delayed activation and uncontrolled viral replication came from a quantitative complementation test using an *Irf3* KO allele.

Notably, this is not the first example of a *de novo* mutation responsible for a CC-strain-specific trait. We (19) and others (20) previously reported the extreme susceptibility of CC042 to *Salmonella* Typhimurium and to *Mycobacterium tuberculosis*, respectively, as a consequence of a *de novo* 15-nucleotide deletion in the *Itgal1* gene. In the case of *Salmonella* Typhimurium, CC042 was standing out, with bacterial loads up to 1000 times higher than other CC and the susceptible B6 strains. In our previous study on ZIKV, CC071 was the most susceptible strain, but its peak viral load was just the highest in a continuous distribution of values. Additionally, this is not the first example of *Irf3* spontaneous polymorphism identified in mice modulating susceptibility to bacterial (21) or viral infections (22).

Irf3 is the main transcription factor involved in the innate immune response. It is constitutively expressed and, at rest, inactive IRF3 is present in the cytoplasm. Upon viral entry (or other stimuli that activate TLRs such as TLR3 and 4 or RIG-like receptors), signal transduction leads to the phosphorylation of IRF3, leading to its dimerization and translocation to the nucleus where it binds to the IFN-I genes promoters (23). This mechanism leads to the very fast production of IFNβ which is secreted by the cell and triggers the immediate response to viral infection through the activation of ISGs with diverse antiviral functions (24). In humans, LOF mutations in *Irf3* have been associated with increased susceptibility to West Nile virus (WNV) (25, 26), herpes simplex virus 1 (HSV-1) (27, 28) and more recently to SARS-CoV-2 (29).

In mice, many studies have used *Irf3*-deficient models. After infection with WNV, viral replication was increased in *Irf3^{KO/KO}* bone marrow macrophages (BMMs) and primary neurons (30). Higher viral replication was observed in HSV-1-infected *Irf3^{KO/KO}* bone marrow-derived dendritic cells (BMDCs) and BMMs, and IFNβ was reduced in BMDCs supernatants (31). Similarly, *Ifnb1* expression was reduced in HSV-1-infected (32) and CHIKV-infected (33) *Irf3^{KO/KO}* MEFs. In line with these findings, our study provides, to our knowledge, the first evidence for a role of *Irf3* in the infection of murine cells by ZIKV.

In vivo studies led to contrasting results depending on the virus and infection route. *Irf3^{KO/KO}* mice were reported to be resistant to ZIKV with no mortality nor body weight loss (10). Moreover, we found that plasma viral loads were similar in B6-*Irf3^{KO/KO}* and B6 mice, whether mice were treated with anti-IFNAR antibody, or not. Likewise, *Irf3^{KO/KO}* mice showed no mortality and low virus in the circulation following dengue virus (34) and CHIKV infection (33). Contrastingly, WNV infection was lethal in *Irf3^{KO/KO}* mice while 65% of infected WT mice survived, suggesting that *Irf3* LOF may impact clinical parameters (survival, body weight loss)

only in the case of highly pathogenic viruses. The route of infection may also be a critical parameter. After intravenous inoculation with HSV-1, all WT and $Irf3^{KO/KO}$ mice survived (32), while intranasal inoculation led to 30% and 90% mortality in WT and $Irf3^{KO/KO}$ mice, respectively (35).

Our *in vivo* data showing higher viral load in CC071 than in B6-$Irf3^{KO/KO}$ mice, with clinical signs, indicate that other genes control the susceptibility of CC071. The results of the complementation test on MEFs, with CC071 and $Irf3^{KO}$/$Irf3^{71}$ yielding very similar data, suggest that these genes may not act at the cell level, at least during the first 72 hpi. Our poor understanding of the severe disease developing in CC071 ZIKV-infected mice does not point at potential mechanisms. As in this study, appropriate crosses will be necessary to detect genomic regions from which candidate genes could be identified and validated. It will then be possible to assess the specific contribution of the *Irf3* LOF mutation to the susceptibility of CC071 mice.

CC071 has been reported to be susceptible to other flaviviruses. High viral load was found after dengue infection (15). Lethality after infection with WNV (15) or Powassan virus (36) were also described, although there is likely contribution from the defective *Oas1b* allele that CC071 has inherited from 129S1/SvImJ. CC071 was also one of the most susceptible CC strains to Rift Valley Fever virus infection (37) and to hepacivirus, with long-term viral persistence (38). Because of its unique genetic background, this strain will be increasingly useful for infectious disease studies, which justifies deciphering its mechanisms of susceptibility.

Material and methods

Mice and crosses

C57BL/6J mice were purchased from Charles River Laboratories France. Collaborative Cross strains (CC001/Unc, CC071/TauUnc, CC005/TauUnc, CC011/Unc, CC026/GeniUnc, CC061/GeniUnc, CC021/Unc, CC006/TauUnc, CC025/GeniUnc, CC039/Unc, CC060/Unc) were purchased from the Systems Genetics Core Facility, University of North Carolina and bred at the Institut Pasteur. *Irf3 Irf7* double KO mice (C57BL/6J-*Bcl2l12/Irf3^{tm1Ttg} Irf7^{tm1Ttg}*,(32, 39); B6-*Irf3^{KO}* and B6-*Irf7^{KO}* thereafter) were bred at the Institut Pasteur. All mice were maintained as described previously (15). (CC001 x CC071) x CC071 backcross embryos were obtained from reciprocal crosses (F1 female x CC071 male and vice-versa). For the quantitative complementation test, B6-*Irf3$^{+/KO}$* B6-*Irf7$^{+/KO}$* females were crossed with CC071 males and B6-*Irf3$^{+/KO}$* B6-*Irf7$^{+/KO}$* males were crossed with C57BL/6J females. Genotype at the *Irf7* loci was not considered as only B6-*Irf7$^{+/KO}$* of B6-*Irf7$^{+/+}$* mice were used. For *in vivo* infections, B6-*Irf3* simple KO mice were obtained from an F2 between B6-*Irf3 Irf7* double KO and B6 mice. Mouse experiments were approved by the Institut Pasteur Ethics Committee and authorized by the French Ministry of Research (project #19469), in compliance with French and European regulations.

ZIKA virus

The FG15 Asian Zika virus (ZIKV) strain, isolated from a patient during ZIKV outbreak in French Guiana in December 2015, was obtained from the Virology Laboratory of the Institut Pasteur of French Guiana. Viral stock (passage 5) was prepared from supernatant of infected C6/36 cells, clarified by centrifugation at 800g and titrated on Vero cells by focus-forming assay.

Mouse infection

All infection experiments were performed in a biosafety level 3 animal facility and mice were kept in isolators. Six- to 10-week-old, male or female mice were injected intraperitoneally with 10^7 PFU of ZIKV FG15. Mice received an IP injection of 2 mg of MAR1-5A3 anti-IFNAR antibody (Euromedex, Cat#BX-BE0241) one day prior infection. Mouse numbers are indicated in figure legends. Both males and females were used since no differences between sexes were detected in our previous and present experiments. Clinical signs and body weight loss were

recorded for up to seven days post infection. Blood samples were collected on EDTA at two- and six-days post infection from the retromandibular vein for plasma viral load assessment. Quantification of ZIKV viral copies by qPCR was previously described (15).

MEFs isolation

Pregnant females were euthanized at day 13.5-15.5 of gestation. For B6, CC001 and CC071, MEFs were isolated from individual fetuses to obtain biological replicates. For other CC strains, MEFs were derived from individual or pooled fetuses. Fetus bodies were chopped and digested with trypsin (Gibco Cat#25300054), then cultured at 37°C and 5% CO_2 in complete medium (DMEM Gibco Cat# 31966047, 10% fetal bovine serum PAA Laboratories Cat#A15-101, 1% penicillin/streptomycin Sigma Cat#P4333). MEFs were used until passage 2. For the backcross experiment, MEF lines were isolated from 52 backcross fetuses. Heads were used to prepare DNA for whole-genome genotyping.

MEFs infection

MEFs were seeded at 5.10^4 cells per well in 24-well plates the day before infection. They were exposed to ZIKV FG15 strain at a MOI of 5 for 2 hours after which the inoculum was replaced with fresh complete medium and MEFs were incubated for up to 72 hours. For kinetics studies, different wells were used for each time point. Backcross MEFs were infected in 6 infection experiments, each of which included a CC001 and a CC071 MEF lines.

MEFs IFNα stimulation

MEFs were seeded at 5.10^4 cells per well in 24-well plates the day before stimulation and treated with 300 IU/mL IFNα (Miltenyi Biotec Cat#130-093-131) and incubated for up to 24 hours.

Focus forming assay

Quantification of ZIKV particles was completed by focus forming assay on Vero cells (ATCC CRL-1586) as previously described (15).

RNA extraction from cells

MEFs were lysed in 350μL of RLT buffer (Qiagen) with 1% β-mercaptoethanol. RNA was extracted using RNeasy Mini Kit (Qiagen Cat#74104) according to the manufacturer's instructions, with DNase I (Qiagen Cat# 79254) to prevent genomic DNA contamination.

Reverse transcription and qPCR

Reverse-transcription was performed on 200ng of RNA using Superscript II polymerase (Invitrogen Cat#18064022) RNaseOUT ribonuclease inhibitor (Invitrogen Cat#10777019). qPCR was performed on 20ng of cDNAs using Power SYBR Green PCR Master Mix (Applied Biosystems Cat#4367659) and 6pmol of each primer, on a QuantStudio 12K Flex or a ViiA 7 (ThermoFisher Scientific). Primers (Eurofins) sequences are provided in Sup Table 1. Gene expression was expressed on a Log10 scale of relative expression to the reference *Tbp* gene.

Genotyping

Genomic DNA was prepared from backcross fetuses' heads by proteinase K digestion, phenol-chloroform extraction and ethanol precipitation according to standard protocols. DNAs were sent to Neogen (Neogen/Geneseek, Inc, Lincoln, NE, USA) for whole-genome genotyping using the MiniMUGA array containing 11,125 SNP markers. For the quantitative complementation test, *Irf3* and *Irf7* genotyping was performed by Transnetyx (Cordova, TN) by real-time PCR on fetuses' heads.

RNA splicing analysis from RNA sequencing data

RNA sequencing data were produced as described in the Supplementary methods. Splicing analysis was performed using Majiq 2.4 (40) with default parameters to investigate alternative transcripts between conditions, focusing on the *Irf3* gene.

Immunofluorescence

MEFs were plated on glass coverslips before infection, fixed with 4% paraformaldehyde for 20 min and permeabilized with pure methanol for 10min at -20°C. Cells were incubated with blocking buffer (5% FBS 0.3% triton in PBS) for 1 hour, with primary antibodies diluted in antibody incubation buffer (AIB: 1% BSA 0.3% triton in PBS) overnight at 4°C and with secondary antibodies and Hoechst (dilution 1:1000) diluted in AIB for 1 hour. Coverslips were mounted on slides and imaged with a widefield microscope (Zeiss Axio Observer.Z1 with a

Plan-Apochromat 20x/0.8 M27 objective and a Hamamatsu sCMOS ORCA-Flash 4.0 v3

camera). ZEN blue 2012 software (ZEISS) imaging software was used for image capture and

Image J software (National Institutes of Health) to adjust brightness and contrast. Primary and

secondary antibodies are indicated in Sup Table 2.

Western blot

MEFs were trypsinized for 5 min, washed and lysed in a protein extraction buffer (10mM

TrisHCl pH7.5, 5mM EDTA, 150mM NaCl, 30mM Na_2HPO_4, 50mM NaF, 10% glycerol, 1% NP40,

1X cOmplete (Roche #11873580001), 1X PhosSTOP (Roche #4906845001), 1/1000 benzonaze

(Sigma Cat#E1014)) for 30 min at 4°C. Proteins diluted in Laemmli were resolved on 4-12% Bis-

Tris gels (Invitrogen Cat#NP0323BOX) in MOPS buffer (Invitrogen Cat#NP0001) and

transferred to nitrocellulose membranes (Bio-Rad Cat#1620112) in a 25mM Tris 200mM

glycine 20% ethanol buffer. Blots were blocked in 5% milk in TBS-T (0.1% Tween20 in Tris Base

Sodium), incubated with primary antibodies diluted in 3% milk in TBST overnight at 4°C, and

incubated with secondary antibodies diluted in 3% milk in TBST for 90 min. Blots were revealed

with ECL substrate (Thermo Scientific Cat#32132) and imaged with X-ray films. Primary and

secondary antibodies are indicated in Sup Table 2.

Statistical analysis

Statistical analyses were performed using R (41) version 4.1.0. Viral titers, gene expression

and genome copies were Log10-transformed for graphs and statistical tests. One way ANOVA

followed by Tukey HSD were used for testing multiple comparisons. For *in vivo* studies, non-

parametric Kruskal-Wallis followed by Wilcoxon tests with Bonferroni correction for multiple

testing were used to handle values below the limit of detection. Linear discriminant analysis

(LDA) was conducted using the MASS package (42). The LDA was trained on the phenotypes

of the two parental CC001 and CC071 strains from each infection batch. LDA coefficients were

applied to backcross mice for assignment to "CC071-like" or "CC001-like" groups.

Genetic analysis

Raw genotypes were curated using the stuart package (43). QTL mapping was performed using

R/qtl (44). LDA prediction was used as a binary trait. Statistical significance thresholds were

computed by data permutation (n=1000). 95% confidence interval was estimated using the

Bayesian method.

Acknowledgements

We thank Matthieu Prot, Maxime Chazal and Sandrine Vandormael-Pournin for technical advice and for providing reagents. We are grateful to Tommy Penel and Rachid Chennouf of the Institut Pasteur Central Animal Facility of the C2RA (Center for Animal Resources and Research) for the careful breeding of CC strains and for the maintenance of mice in the BSL-3 animal facility, respectively. We thank Etienne Simon-Loriere and Nolwenn Jouvenet for scientific advice and continuous support along this project.

Author contributions

X.M. conceived and supervised this study. M.B. and C.M. designed and performed the experiments and analyzed data. L.C. and C.R.P. performed experiments. E.K. analyzed RNAseq data. X.M. designed experiments and analyzed data. E.B. provided conceptual advice. M.B., C.M. and X.M. wrote the paper. All authors commented on and edited the manuscript.

Funding

This work was supported by grants from the Agence Nationale de la Recherche (ANR) NeuroZika (ANR-20-CE16-0017) and from the French Government's Investissement d'Avenir program, Laboratoire d'Excellence: IBEID (Integrative Biology of Emerging Infectious Diseases, ANR-10-LABX-62-IBEID). C.M. and M.B. were supported by doctoral fellowships from grant ANR-10-LABX-62-IBEID.

Availability of data and materials

RNAseq primary data are available from the European Nucleotide Archive (accession number E-MTAB-12765). All other experimental data that support the finding of this study are available from the corresponding authors upon reasonable request.

Competing interests

The authors declare that they have no competing interests.

Figure legends

Figure 1. CC071 MEFs fail to control viral replication, with delayed *Ifnb1* expression but

normal response to IFN-I. (A,B) MEFs derived from B6 (black circles), CC001 (blue triangles)

and CC071 (red squares) were infected with ZIKV at a MOI of 5 and analyzed 24, 48 and 72 hpi

(A) Viral titer in supernatants was quantified by FFA. (B) *Ifnb1* expression was normalized to

Tbp reference gene. Data are mean +/- sem from 3 to 4 biological replicates per strain (MEFs

derived from individual embryos). (C) MEFs were stimulated with recombinant IFNα. *Ifitm3*

relative expression normalized to *Tbp* reference gene is shown as an example of ISG. Data are

mean +/- sem from 2 to 3 biological replicates. Blue asterisks and black hashes show statistical

significance of CC071 compared to CC001 and to B6, respectively (ANOVA followed by Tukey

HSD, */# $p < 0.05$, **/## $p < 0.01$, *** $p < 0.001$).

Figure 2. Haplotype analysis fails to identify a genes from the IFN-I induction pathway

associated with uncontrolled viral replication. (A) Identification of CC strains which carry the

same ancestral haplotype as CC071 at the genes involved in the pathway leading to IFN-I

expression. Colored boxes indicate matched haplotypes between CC071 and other CC strains.

Letters and colors designate the eight CC founder strains. A: A/J (yellow); B: C57BL/6J; C:

129S1/SvImJ (pink); D: NOD/ShiLtJ; E: NZO/HILtJ (light blue); F: CAST/EiJ (green); G: PWK/PhJ

(red); H: WSB/EiJ (purple). Doubled letters (eg AA) indicate homozygous genotypes.

Heterozygous genotypes are indicated by the two corresponding letters (eg AD). (B) Kinetics

of viral titer in MEFs from CC071 and the 8 CC strains shown in (A). Experimental conditions

were as in Figure 1A. n: number of technical replicates for each strain. Data are mean +/- sem

from the technical replicates. Letters show statistical significance between CC071 and other

strains. (ANOVA followed by Tukey HSD, a: $p < 0.05$, b: $p < 0.01$, c: $p < 0.001$).

Figure 3. Backcross MEFs display either a CC001-like or a CC071-like phenotype. MEFs

derived from CC001 (blue triangles), CC071 (red squares) and backcross (gray circles) embryos

were infected with ZIKV at a MOI of 5. (A) Viral titer and (B) *Ifnb1* expression in ZIKV-infected

MEFs from 9 backcross embryos. Experimental conditions were as in Figure 1A-B. Red and

blue curves show the results for CC071 and CC001 MEFs, respectively, from the same infection

experiment. (C) Results of LDA on the backcross MEFs. LDA coefficients were calculated from *Ifnb1* expression data in CC001 and CC071 infected MEFs and applied to backcross MEFs. The graph shows the probability of each BC MEF to belong to the "CC001-like" group, resulting in two distinct populations shown in blue ("CC001-like") and in red ("CC071-like"). n: number of BC MEFs assigned to each group.

Figure 4. Genetic analysis of *Ifnb1* expression in backcross MEFs identifies a major determinant mapping to the *Irf3* locus. (A) Genome-wide linkage analysis of the LDA classification of backcross MEFs performed with R/qtl. X-axis: genomic location. Y-axis: LOD score of the phenotype-genotype association. Genome-wide significance thresholds (P = 0.05, P = 0.1 and P = 0.63) were computed from 1000 data permutation. The chromosome 7 peak has a LOD score of 9.138. (B) Schematic representation of CC001 and CC071 chromosome 7 haplotypes, from https://csbio.unc.edu/CCstatus/CCGenomes/. Colors represent the CC ancestral haplotypes (same colors as in Figure 2A). Thick vertical black lines show the peak's 95% Bayesian confidence interval (25.9-31.3 cM, corresponding to 40.1-50.6 Mb). The red line shows the position of *Irf3* gene.

Figure 5. CC071 *Irf3* mRNAs show abnormal splicing, resulting in defective IRF3 protein. (A) Schematic representation of the exons of the *Irf3* gene with the number of reads spanning successive exons in the CC001 and CC071 RNAseq data (one sample of each strain). The red box between CC071's exons 6 and 7 depicts a novel exon resulting from abnormal splicing. (B) Schematic representation of the IRF3 protein structural domains (exon 1 is untranslated). Exon 8 encodes the serine rich region containing the phosphorylation sites for IRF3 activation. (C) Western blot using an anti-C-terminal IRF3 antibody from mock-infected and ZIKV-infected B6, CC001 and CC071 MEFs at 2 hpi, showing the absence of full-length IRF3 in CC071 MEFs. Vinculin was used as a loading control. (D) Immunofluorescence using an anti-phosphorylated IRF3 (pIRF3, green) in ZIKV-infected B6, CC001 and CC071 MEFs at 24 hpi, showing the absence of pIRF3 in the nucleus of CC071 MEFs upon infection. Red-labeled 4G2 antibody labels ZIKV-infected cells. Cell nuclear DNA labeled by Hoechst (blue). Quantification of the number of infected and pIRF3 positive cells is presented in the table. Proportions were established on 420, 428 and 551 cells for CC001, CC071 and B6, respectively.

Figure 6. Quantitative complementation test confirms that *Irf3* LOF in CC071 is responsible

for uncontrolled viral replication and delayed *Ifnb1* expression. (A) Mice heterozygous for an

inactivated *Irf3* allele (B6-*Irf3*$^+$/*Irf3*^{KO}) were mated with B6 and with CC071 mice to produce

four types of embryos from which MEFs were derived. The genetic background and *Irf3*

genotype is shown below each type of embryos. (B) Viral titer and (C) *Ifnb1* expression in ZIKV-

infected MEFs. Experimental conditions were as in Figure 1A-B. Data are mean +/- sem from

3 biological replicates for each group. At every time point, the comparison of either CC071 or

Irf3^{KO}/*Irf3*^{71} with either *Irf3*$^+$/*Irf3*^{71} or *Irf3*$^+$/*Irf3*^{KO} gave identical statistical significance (ANOVA

followed by post-hoc Tukey HSD, * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$).

Figure 7. *Irf3* LOF is not sufficient to explain CC071 susceptibility to ZIKV infection *in vivo*.

CC071, B6-*Irf3*^{KO} and B6 mice were infected IP with 10^7 FFUs of ZIKV and monitored for 7 days,

with (A,B) or without (C) IP injection of 2 mg of MAR1-5A3 IFNAR-blocking monoclonal

antibody 24 h before infection. (A) Average clinical score of anti-IFNAR-treated at day 6 p.i.

(B,C) Plasma viral load measured by RT-qPCR. Each dot represents one mouse. Groups were

compared by Kruskal-Wallis test followed by Wilcoxon test with Bonferroni correction for

multiple testing (* $p < 0.05$).

Supplemental information

Legend to supplemental figures

Figure S1. *Ifna4* **expression is delayed in ZIKV-infected CC071 MEFs.** MEFs derived from B6 (gray circles), CC001 (blue triangles) and CC071 (red squares) were infected with ZIKV at a MOI of 5. *Ifna4* expression was determined by RT-qPCR on MEFs total RNA by normalizing to *Tbp* housekeeping gene. Data are mean +/- sem from 3 biological replicates. One CC001, one B6 and one CC071 replicates at 0 hpi and one CC071 replicate at 24 hpi were below the limit of detection. Blue asterisks and black hashes show statistical significance of CC071 compared to CC001 and to B6, respectively (ANOVA followed by post-hoc Tukey HSD, */# $p < 0.05$, *** $p < 0.001$).

Figure S2. Delayed IFN-I expression in CC071 MEFs results from a constitutive defect in innate antiviral response. MEFs were transfected with either (A) poly(I:C), which activates both TLR and RIG-I pathways, or (B) 3p-hpRNA, a RIG-I agonist. *Ifnb1* expression was determined as in Figure 1B. Data are mean +/- sem from 3 biological replicates for CC001 and CC071 (2 at 4 hours) or 2 biological replicates for B6 (1 at 4 hours). Blue asterisks and black hashes show statistical significance of CC071 compared to CC001 and to B6, respectively (ANOVA followed by post-hoc Tukey HSD, */# $p < 0.05$).

Figure S3. **Expression of genes involved in the** *Ifnb1* **induction pathway in ZIKV-infected MEFs of B6, CC001 and CC071 strains.** MEFs derived from B6 (black circles), CC001 (blue triangles) and CC071 (red squares) were infected with ZIKV at a MOI of 5. mRNA expression levels were measured by RNAseq in non-infected (NI) and ZIKV-infected MEFs. (A) Number of upregulated genes per strain at 16, 24 and 32 hpi (B-G) Genes constitutively expressed. (H-N) Genes which expression is induced by the IFN-I response (ISGs). Expression levels are shown on a logarithmic scale. For *Ifnb1* expression, null counts were transformed to 1. Data are mean +/- sem from 3 biological replicates. Blue asterisks and black hashes show statistical significance of CC071 compared to CC001 to B6, respectively (ANOVA followed by post-hoc Tukey HSD, */# $p < 0.05$, **/## $p < 0.01$, ***/### $p < 0.001$).

585 **Figure S4. (CC001xCC071)F1 MEFs show normal induction of *Ifnb1*.** *Ifnb1* expression upon

586 ZIKV infection in CC001, CC071 and (CC001xCC071)F1 MEFs determined as in Figure 1B. Data

587 are mean +/- sem from 3 biological replicates. Blue asterisks and orange hashes show

588 statistical significance of CC071 compared to CC001 and to F1, respectively (ANOVA followed

589 by post-hoc Tukey HSD, **/## $p < 0.01$).

References

1. Cao-Lormeau V-M, Blake A, Mons S, Lastère S, Roche C, Vanhomwegen J, Dub T, Baudouin L, Teissier A, Larre P, Vial A-L, Decam C, Choumet V, Halstead SK, Willison HJ, Musset L, Manuguerra J-C, Despres P, Fournier E, Mallet H-P, Musso D, Fontanet A, Neil J, Ghawché F. 2016. Guillain-Barré Syndrome outbreak associated with Zika virus infection in French Polynesia: a case-control study. Lancet 387:1531–1539.

2. Carod-Artal FJ. 2018. Neurological complications of Zika virus infection. Expert Review of Anti-infective Therapy 16:399–410.

3. Oliveira Melo AS, Malinger G, Ximenes R, Szejnfeld PO, Alves Sampaio S, Bispo de Filippis AM. 2016. Zika virus intrauterine infection causes fetal brain abnormality and microcephaly: tip of the iceberg?: Physician Alert. Ultrasound Obstet Gynecol 47:6–7.

4. Paixao ES, Cardim LL, Costa MCN, Brickley EB, de Carvalho-Sauer RCO, Carmo EH, Andrade RFS, Rodrigues MS, Veiga RV, Costa LC, Moore CA, França GVA, Smeeth L, Rodrigues LC, Barreto ML, Teixeira MG. 2022. Mortality from Congenital Zika Syndrome — Nationwide Cohort Study in Brazil. N Engl J Med 386:757–767.

5. Rossi SL, Ebel GD, Shan C, Shi P-Y, Vasilakis N. 2018. Did Zika Virus Mutate to Cause Severe Outbreaks? Trends in Microbiology 26:877–885.

6. Liu Z-Y, Shi W-F, Qin C-F. 2019. The evolution of Zika virus from Asia to the Americas. Nat Rev Microbiol 17:131–139.

7. Manet C, Roth C, Tawfik A, Cantaert T, Sakuntabhai A, Montagutelli X. 2018. Host genetic control of mosquito-borne Flavivirus infections. Mamm Genome 29:384–407.

8.   Rasmussen AL, Okumura A, Ferris MT, Green R, Feldmann F, Kelly SM, Scott DP, Safronetz

D, Haddock E, LaCasse R, Thomas MJ, Sova P, Carter VS, Weiss JM, Miller DR, Shaw GD,

Korth MJ, Heise MT, Baric RS, de Villena FP-M, Feldmann H, Katze MG. 2014. Host genetic

diversity enables Ebola hemorrhagic fever pathogenesis and resistance. Science

346:987–991.

9.   Gralinski LE, Menachery VD, Morgan AP, Totura AL, Beall A, Kocher J, Plante J, Harrison-

Shostak DC, Schafer A, Pardo-Manuel de Villena F, Ferris MT, Baric RS. 2017. Allelic

Variation in the Toll-Like Receptor Adaptor Protein Ticam2 Contributes to SARS-

Coronavirus Pathogenesis in Mice. G3 (Bethesda) 7:1653–1663.

10.   Lazear HM, Govero J, Smith AM, Platt DJ, Fernandez E, Miner JJ, Diamond MS. 2016. A

Mouse Model of Zika Virus Pathogenesis. Cell Host Microbe 19:720–730.

11.   Grant A, Ponia SS, Tripathi S, Balasubramaniam V, Miorin L, Sourisseau M, Schwarz MC,

Sánchez-Seco MP, Evans MJ, Best SM, García-Sastre A. 2016. Zika Virus Targets Human

STAT2 to Inhibit Type I Interferon Signaling. Cell Host & Microbe 19:882–890.

12.   Sheehan KCF, Lai KS, Dunn GP, Bruce AT, Diamond MS, Heutel JD, Dungo-Arthur C,

Carrero JA, White JM, Hertzog PJ, Schreiber RD. 2006. Blocking Monoclonal Antibodies

Specific for Mouse IFN- α / β Receptor Subunit 1 (IFNAR-1) from Mice Immunized by *In*

Vivo Hydrodynamic Transfection. Journal of Interferon & Cytokine Research 26:804–819.

13.   Roberts A, Pardo-Manuel de Villena F, Wang W, McMillan L, Threadgill DW. 2007. The

polymorphism architecture of mouse genetic resources elucidated using genome-wide

resequencing data: implications for QTL discovery and systems genetics. Mamm Genome

18:473–481.

14.   Keane TM, Goodstadt L, Danecek P, White MA, Wong K, Yalcin B, Heger A, Agam A, Slater

635   G, Goodson M, Furlotte NA, Eskin E, Nellåker C, Whitley H, Cleak J, Janowitz D,

636   Hernandez-Pliego P, Edwards A, Belgard TG, Oliver PL, McIntyre RE, Bhomra A, Nicod J,

637   Gan X, Yuan W, van der Weyden L, Steward CA, Bala S, Stalker J, Mott R, Durbin R, Jackson

638   IJ, Czechanski A, Guerra-Assunção JA, Donahue LR, Reinholdt LG, Payseur BA, Ponting CP,

639   Birney E, Flint J, Adams DJ. 2011. Mouse genomic variation and its effect on phenotypes

640   and gene regulation. Nature 477:289–294.

641   15.   Manet C, Simon-Lorière E, Jouvion G, Hardy D, Prot M, Conquet L, Flamand M, Panthier

642   J-J, Sakuntabhai A, Montagutelli X. 2020. Genetic Diversity of Collaborative Cross Mice

643   Controls Viral Replication, Clinical Severity, and Brain Pathology Induced by Zika Virus

644   Infection, Independently of Oas1b. J Virol 94.

645   16.   McCartney S, Vermi W, Gilfillan S, Cella M, Murphy TL, Schreiber RD, Murphy KM,

646   Colonna M. 2009. Distinct and complementary functions of MDA5 and TLR3 in poly(I:C)-

647   mediated activation of mouse NK cells. Journal of Experimental Medicine 206:2967–

648   2976.

649   17.   Collaborative Cross Consortium. 2012. The Genome Architecture of the Collaborative

650   Cross Mouse Genetic Reference Population. Genetics 190:389–401.

651   18.   Srivastava A, Morgan AP, Najarian ML, Sarsani VK, Sigmon JS, Shorter JR, Kashfeen A,

652   McMullan RC, Williams LH, Giusti-Rodríguez P, Ferris MT, Sullivan P, Hock P, Miller DR,

653   Bell TA, McMillan L, Churchill GA, Villena FP-M de. 2017. Genomes of the Mouse

654   Collaborative Cross. Genetics 206:537–556.

655   19.   Zhang J, Teh M, Kim J, Eva MM, Cayrol R, Meade R, Nijnik A, Montagutelli X, Malo D,

656   Jaubert J. 2019. A Loss-of-Function Mutation in the Integrin Alpha L (Itgal) Gene

657   Contributes to Susceptibility to Salmonella enterica Serovar Typhimurium Infection in

658   Collaborative Cross Strain CC042. Infect Immun 88.

659   20.   Smith CM, Proulx MK, Lai R, Kiritsy MC, Bell TA, Hock P, Pardo-Manuel de Villena F, Ferris

660   MT, Baker RE, Behar SM, Sassetti CM. 2019. Functionally Overlapping Variants Control

661   Tuberculosis Susceptibility in Collaborative Cross Mice. mBio 10:e02791-19.

662   21.   Garifulin O, Qi Z, Shen H, Patnala S, Green MR, Boyartchuk V. 2007. Irf3 Polymorphism

663   Alters Induction of Interferon Beta in Response to Listeria monocytogenes Infection.

664   PLOS Genetics 3:e152.

665   22.   Moore TC, Al-Salleeh FM, Brown DM, Petro TM. 2011. IRF3 polymorphisms induce

666   different innate anti-Theiler's virus immune responses in RAW264.7 macrophages.

667   Virology 418:40–48.

668   23.   Bourdon M, Manet C, Montagutelli X. 2020. Host genetic susceptibility to viral infections:

669   the role of type I interferon induction. Genes Immun 21:365–379.

670   24.   Schoggins JW. 2019. Interferon-Stimulated Genes: What Do They All Do? Annu Rev Virol

671   6:567–584.

672   25.   Bigham AW, Buckingham KJ, Husain S, Emond MJ, Bofferding KM, Gildersleeve H,

673   Rutherford A, Astakhova NM, Perelygin AA, Busch MP, Murray KO, Sejvar JJ, Green S,

674   Kriesel J, Brinton MA, Bamshad M. 2011. Host genetic risk factors for West Nile virus

675   infection and disease progression. PLoS One 6:e24745.

676   26.   Cahill ME, Conley S, DeWan AT, Montgomery RR. 2018. Identification of genetic variants

677   associated with dengue or West Nile virus disease: a systematic review and meta-

678   analysis. BMC Infect Dis 18:282.

679   27.   Mork N, Kofod-Olsen E, Sorensen KB, Bach E, Orntoft TF, Ostergaard L, Paludan SR,

680   Christiansen M, Mogensen TH. 2015. Mutations in the TLR3 signaling pathway and

681   beyond in adult patients with herpes simplex encephalitis. Genes Immun 16:552–66.

682   28.   Andersen LL, Mork N, Reinert LS, Kofod-Olsen E, Narita R, Jorgensen SE, Skipper KA,

683   Honing K, Gad HH, Ostergaard L, Orntoft TF, Hornung V, Paludan SR, Mikkelsen JG, Fujita

684   T, Christiansen M, Hartmann R, Mogensen TH. 2015. Functional IRF3 deficiency in a

685   patient with herpes simplex encephalitis. J Exp Med 212:1371–9.

686   29.   Zhang Q, Bastard P, Liu Z, Le Pen J, Moncada-Velez M, Chen J, Ogishi M, Sabli IKD, Hodeib

687   S, Korol C, Rosain J, Bilguvar K, Ye J, Bolze A, Bigio B, Yang R, Arias AA, Zhou Q, Zhang Y,

688   Onodi F, Korniotis S, Karpf L, Philippot Q, Chbihi M, Bonnet-Madin L, Dorgham K, Smith

689   N, Schneider WM, Razooky BS, Hoffmann HH, Michailidis E, Moens L, Han JE, Lorenzo L,

690   Bizien L, Meade P, Neehus AL, Ugurbil AC, Corneau A, Kerner G, Zhang P, Rapaport F,

691   Seeleuthner Y, Manry J, Masson C, Schmitt Y, Schluter A, Le Voyer T, Khan T, Li J, Fellay J,

692   Roussel L, Shahrooei M, Alosaimi MF, Mansouri D, Al-Saud H, Al-Mulla F, Almourfi F, Al-

693   Muhsen SZ, Alsohime F, Al Turki S, Hasanato R, van de Beek D, Biondi A, Bettini LR,

694   D'Angio M, Bonfanti P, Imberti L, Sottini A, Paghera S, Quiros-Roldan E, Rossi C, Oler AJ,

695   Tompkins MF, Alba C, Vandernoot I, Goffard JC, Smits G, Migeotte I, Haerynck F, Soler-

696   Palacin P, Martin-Nalda A, Colobran R, Morange PE, Keles S, Colkesen F, Ozcelik T, Yasar

697   KK, Senoglu S, Karabela SN, Gallego CR, Novelli G, Hraiech S, Tandjaoui-Lambiotte Y,

698   Duval X, Laouenan C, Clinicians C-S, Clinicians C, Imagine CG, French CCSG, Co VCC,

Amsterdam UMCC, Biobank, Effort CHG, Niaid U, Group TCI, Snow AL, Dalgard CL, Milner

J, Vinh DC, Mogensen TH, Marr N, Spaan AN, Boisson B, Boisson-Dupuis S, Bustamante J,

Puel A, Ciancanelli M, Meyts I, Maniatis T, Soumelis V, Amara A, Nussenzweig M, Garcia-

Sastre A, Krammer F, Pujol A, Duffy D, Lifton R, Zhang SY, Gorochov G, Beziat V, Jouanguy

E, Sancho-Shimizu V, Rice CM, Abel L, Notarangelo LD, Cobat A, Su HC, Casanova JL. 2020.

Inborn errors of type I IFN immunity in patients with life-threatening COVID-19. Science

https://doi.org/10.1126/science.abd4570.

30. Daffis S, Samuel MA, Keller BC, Gale M Jr, Diamond MS. 2007. Cell-specific IRF-3

responses protect against West Nile virus infection by interferon-dependent and -

independent mechanisms. PLoS Pathog 3:e106.

31. Menachery VD, Leib DA. 2009. Control of Herpes Simplex Virus Replication Is Mediated

through an Interferon Regulatory Factor 3-Dependent Pathway. J Virol 83:12399–12406.

32. Honda K, Yanai H, Negishi H, Asagiri M, Sato M, Mizutani T, Shimada N, Ohba Y, Takaoka

A, Yoshida N, Taniguchi T. 2005. IRF-7 is the master regulator of type-I interferon-

dependent immune responses. Nature 434:772–7.

33. Schilte C, Buckwalter MR, Laird ME, Diamond MS, Schwartz O, Albert ML. 2012. Cutting

Edge: Independent Roles for IRF-3 and IRF-7 in Hematopoietic and Nonhematopoietic

Cells during Host Response to Chikungunya Infection. JI 188:2967–2971.

34. Chen HW, King K, Tu J, Sanchez M, Luster AD, Shresta S. 2013. The roles of IRF-3 and IRF-

7 in innate antiviral immunity against dengue virus. J Immunol 191:4194–201.

35.   Canivet C, Rhéaume C, Lebel M, Piret J, Gosselin J, Boivin G. 2018. Both IRF3 and especially IRF7 play a key role to orchestrate an effective cerebral inflammatory response in a mouse model of herpes simplex virus encephalitis. J Neurovirol 24:761–768.

36.   Jasperse BA, Mattocks MD, Noll KM, Ferris MT, Heise MT, Lazear HM. 2022. Neuroinvasive flavivirus pathogenesis is restricted by host genetic factors in Collaborative Cross mice, independently of Oas1b. bioRxiv https://doi.org/10.1101/2022.10.24.513634.

37.   Cartwright HN, Barbeau DJ, Doyle JD, Klein E, Heise MT, Ferris MT, McElroy AK. 2022. Genetic diversity of collaborative cross mice enables identification of novel rift valley fever virus encephalitis model. PLOS Pathogens 18:e1010649.

38.   Brown AJ, Won JJ, Wolfisberg R, Fahnøe U, Catanzaro N, West A, Moreira FR, Batista MN, Ferris MT, Linnertz CL, Leist SR, Nguyen C, De la Cruz G, Midkiff BR, Xia Y, Montgomery SA, Billerbeck E, Bukh J, Scheel TKH, Rice CM, Sheahan TP. 2023. Host genetic variation guides hepacivirus clearance, chronicity, and liver fibrosis in mice. preprint. Microbiology.

39.   Sato M, Suemori H, Hata N, Asagiri M, Ogasawara K, Nakao K, Nakaya T, Katsuki M, Noguchi S, Tanaka N, Taniguchi T. 2000. Distinct and essential roles of transcription factors IRF-3 and IRF-7 in response to viruses for IFN-alpha/beta gene induction. Immunity 13:539–48.

40.   Vaquero-Garcia J, Barrera A, Gazzara MR, González-Vallinas J, Lahens NF, Hogenesch JB, Lynch KW, Barash Y. 2016. A new view of transcriptome complexity and regulation through the lens of local splicing variations. eLife 5:e11752.

41. R Core Team. 2021. R: A Language and Environment for Statistical Computing.

https://www.R-project.org/.

42. Venables WN, Ripley BD. 2002. Modern Applied Statistics with S. Fourth EditionFourth.

Springer. https://www.stats.ox.ac.uk/pub/MASS4/.

43. Bourdon M, Montagutelli X. 2022. stuart: an R package for the curation of SNP genotypes

from experimental crosses. G3 (Bethesda) jkac219.

44. Broman KW, Wu H, Sen S, Churchill GA. 2003. R/qtl: QTL mapping in experimental

crosses. Bioinformatics 19:889–890.

749
750

Figure 1. CC071 MEFs fail to control viral replication, with delayed *Ifnb1* expression but normal response to IFN-I. (A,B) MEFs derived from B6 (black circles), CC001 (blue triangles) and CC071 (red squares) were infected with ZIKV at a MOI of 5 and analyzed 24, 48 and 72 h.p.i. (A) Viral titer in supernatants was quantified by FFA. (B) *Ifnb1* expression was normalized to *Tbp* reference gene. Data are mean +/- sem from 3 to 4 biological replicates per strain (MEFs derived from individual embryos). (C) MEFs were stimulated with recombinant IFNα. *Ifitm3* relative expression normalized to *Tbp* reference gene is shown as an example of ISG. Data are mean +/- sem from 2 to 3 biological replicates. Blue asterisks and black hashes show statistical significance of CC071 compared to CC001 and to B6, respectively (ANOVA followed by Tukey HSD, */# p < 0.05, **/## p < 0.01, *** p < 0.001).

A

Comparison of CC strains haplotypes of IFN−I pathway genes

Gene	Protein	CC071	CC001	CC005	CC026	CC021	CC006	CC025	CC039	CC060
Ddx58	RIG−I	HH	BB	EE	CC	HH	AA	HH	AD	EE
Ifih1	LGP2	GG	AA	DD	HH	GG	BB	AA	BB	CC
Dhx58	MDA−5	FF	EE	DD	HH	FF	DD	FF	FF	EE
Tlr3	TLR3	AA	AA	BB	CC	EE	AA	BB	FF	CC
Mavs	MAVS	CC	AA	DD	HH	GG	CC	DD	HH	DD
Ticam1	TRIF	CC	AA	EE	BB	CC	DD	HH	AA	EE
Azi2	NAP1	EE	DD	FF	BB	FF	EE	GG	CC	EE
Traf3	TRAF3	HH	HH	GG	CC	FF	CC	CC	AA	BB
Tbk1	TBK1	HH	DD	GG	HH	AA	FF	AA	GG	BB
Ikbke	IKKε	CC	EE	CC	AA	EE	CC	CC	GH	AA
Irf3	IRF3	FF	FF	HH	BB	CC	DD	EE	HH	BB
Irf7	IRF7	HH	BB	DD	DD	HH	EE	DD	HH	DD
Ifnb1	IFNβ	HH	EE	DD	CC	HH	BB	HH	CC	DE

B

Figure 2. Haplotype analysis fails to identify a genes from the IFN-I induction pathway associated with uncontrolled viral replication. (A) Identification of CC strains which carry the same ancestral haplotype as CC071 at the genes involved in the pathway leading to IFN-I expression. Colored boxes indicate matched haplotypes between CC071 and other CC strains. Letters and colors designate the eight CC founder strains. A: A/J (yellow); B: C57BL/6J; C: 129S1/SvImJ (pink); D: NOD/ShiLtJ; E: NZO/HILtJ (light blue); F: CAST/EiJ (green); G: PWK/PhJ (red); H: WSB/EiJ (purple). Doubled letters (eg AA) indicate homozygous genotypes. Heterozygous genotypes are indicated by the two corresponding letters (eg AD). (B) Kinetics of viral titer in MEFs from CC071 and the 8 CC strains shown in (A). Experimental conditions were as in Figure 1A-C. n: number of technical replicates for each strain. Data are mean +/- sem from the technicalreplicates. Letters show statistical significance between CC071 and other strains. (ANOVA followed by Tukey HSD, a: $p < 0.05$, b: $p < 0.01$, c: $p < 0.001$).

Figure 3. Backcross MEFs display either a CC001-like or a CC071-like phenotype. MEFs derived from CC001 (blue triangles), CC071 (red squares) and backcross (gray circles) embryos were infected with ZIKV at a MOI of 5. (A) Viral titer and (B) Ifnb1 expression in ZIKV-infected MEFs from 9 backcross embryos. Experimental conditions were as in Figure 1A-B. Red and blue curves show the results for CC071 and CC001 MEFs, respectively, from the same infection experiment. (C) Results of LDA on the backcross MEFs. LDA coefficients were calculated from Ifnb1 expression data in CC001 and CC071 infected MEFs and applied to backcross MEFs. The graph shows the probability of each BC MEF to belong to the "CC001-like" group, resulting in two distinct populations shown in blue ("CC001-like") and in red ("CC071-like"). n: number of BC MEFs assigned to each group.

Figure 4. Genetic analysis of *Ifnb1* expression in backcross MEFs identifies a major determinant mapping to the *Irf3* locus. (A) Genome-wide linkage analysis of the LDA classification of backcross MEFs performed with R/qtl. X-axis: genomic location. Y-axis: LOD score of the phenotype-genotype association. Genome-wide significance thresholds (P = 0.05, P = 0.1 and P = 0.63) were computed from 1000 data permutation. The chromosome 7 peak has a LOD score of 9.138. (B) Schematic representation of CC001 and CC071 chromosome 7 haplotypes, from https://csbio.unc.edu/CCstatus/CCGenomes/. Colors represent the CC ancestral haplotypes (same colors as in Figure 2A). Thick vertical black lines show the peak's 95% Bayesian confidence interval (25.9-31.3 cM, corresponding to 40.1-50.6 Mb). The red line shows the position of *Irf3* gene.

Count	B6	CC001	CC071
Infected cells	24	91	76
pIRF3 positive cells	5	28	0
Percentage	**21%**	**31%**	**0%**

Figure 5. CC071 *Irf3* mRNAs show abnormal splicing, resulting in defective IRF3 protein. Schematic representation of the exons of the *Irf3* gene with the number of reads spanning successive exons in the CC001 and CC071 RNAseq data (one sample of each strain). The red box between CC071's exons 6 and 7 depicts a novel exon resulting from abnormal splicing. (B) Schematic representation of the IRF3 protein structural domains (exon 1 is untranslated). Exon 8 encodes the serine rich region containing the phosphorylation sites for IRF3 activation. (C) Western blot using an anti-C-terminal IRF3 antibody from mock-infected and ZIKV-infected B6, CC001 and CC071 MEFs at 2 h.p.i., showing the absence of full-length IRF3 in CC071 MEFs. Vinculin was used as a loading control. (D) Immunofluorescence using an anti-phosphorylated IRF3 (pIRF3, green) in ZIKV-infected B6, CC001 and CC071 MEFs at 24 h.p.i., showing the absence of pIRF3 in the nucleus of CC071 MEFs upon infection. Red-labeled 4G2 antibody labels ZIKV-infected cells. Cell nuclear DNA labeled by Hoechst (blue). Quantification of the number of infected and pIRF3 positive cells is presented in the table. Proportions were established on 420, 428 and 551 cells for CC001, CC071 and B6, respectively.

Figure 6. Quantitative complementation test confirms that *Irf3* LOF in CC071 is responsible for uncontrolled viral replication and delayed *Ifnb1* expression. (A) Mice heterozygous for an inactivated *Irf3* allele (B6-$Irf3^+/Irf3^{KO}$) were mated with B6 and with CC071 mice to produce four types of embryos from which MEFs were derived. The genetic background and *Irf3* genotype is shown below each type of embryos. (B) Viral titer and (C) *Ifnb1* expression in ZIKV-infected MEFs. Experimental conditions were as in Figure 1A-B. Data are mean +/- sem from 3 biological replicates for each group. At every time point, the comparison of either CC071 or $Irf3^{KO}/Irf3^{71}$ with either $Irf3^+/Irf3^{71}$ or $Irf3^+/Irf3^{KO}$ gave identical statistical significance (ANOVA followed by post-hoc Tukey HSD, * p < 0.05, ** p < 0.01, *** p < 0.001).

A Clinical scores at 7 days p.i.

Figure 7. _Irf3_ LOF is not sufficient to explain CC071 susceptibility to ZIKV infection _in vivo._ CC071, B6-$Irf3^{KO}$ and B6 mice were infected IP with 10^7 FFUs of ZIKV and monitored for 7 days, with (A,B) or without (C) IP injection of 2 mg of MAR1-5A3 IFNAR-blocking monoclonal antibody 24 h before infection. (A) Average clinical score of anti-IFNAR-treated at day 6 p.i. (B,C) Plasma viral load measured by RT-qPCR. Each dot represents one mouse. Groups were compared by Kruskal-Wallis test followed by Wilcoxon test with Bonferroni correction for multiple testing (* $p < 0.05$).

Figure S1. *Ifna4* **expression is delayed in ZIKV-infected CC071 MEFs.** MEFs derived from B6 (gray circles), CC001 (blue triangles) and CC071 (red squares) were infected with ZIKV at a MOI of 5. *Ifna4* expression was determined by RT-qPCR on MEFs total RNA by normalizing to *Tbp* housekeeping gene. Data are mean +/- sem from 3 biological replicates. One CC001, one B6 and one CC071 replicates at 0 hpi and one CC071 replicate at 24 hpi were below the limit of detection. Blue asterisks and black hashes show statistical significance of CC071 compared to CC001 and to B6, respectively (ANOVA followed by post-hoc Tukey HSD, */# $p < 0.05$, *** $p < 0.001$).

Figure S2. Delayed IFN-I expression in CC071 MEFs results from a constitutive defect in innate antiviral response. MEFs were transfected with either (A) poly(I:C), which activates both TLR and RIG-I pathways, or (B) 3p-hpRNA, a RIG-I agonist. *Ifnb1* expression was determined as in Figure 1B. Data are mean +/- sem from 3 biological replicates for CC001 and CC071 (2 at 4 hours) or 2 biological replicates for B6 (1 at 4 hours). Blue asterisks and black hashes show statistical significance of CC071 compared to CC001 and to B6, respectively (ANOVA followed by post-hoc Tukey HSD, */# $p < 0.05$).

Figure S3. Expression of genes involved in the *Ifnb1* induction pathway in ZIKV-infected MEFs of B6, CC001 and CC071 strains. MEFs derived from B6 (black circles), CC001 (blue triangles) and CC071 (red squares) were infected with ZIKV at a MOI of 5. mRNA expression levels were measured by RNAseq in non-infected (NI) and ZIKV-infected MEFs. (A) Number of upregulated genes per strain at 16, 24 and 32 hpi (B-G) Genes constitutively expressed. (H-N) Genes which expression is induced by the IFN-I response (ISGs). Expression levels are shown on a logarithmic scale. For *Ifnb1* expression, null counts were transformed to 1. Data are mean +/- sem from 3 biological replicates. Blue asterisks and black hashes show statistical significance of CC071 compared to CC001 to B6, respectively (ANOVA followed by post-hoc Tukey HSD, */# p < 0.05, **/## p < 0.01, ***/### p < 0.001).

Figure S4. (CC001xCC071)F1 MEFs show normal induction of *Ifnb1*. *Ifnb1* expression upon ZIKV infection in CC001, CC071 and (CC001xCC071)F1 MEFs determined as in Figure 1B. Data are mean +/- sem from 3 biological replicates. Blue asterisks and orange hashes show statistical significance of CC071 compared to CC001 and to F1, respectively (ANOVA followed by post-hoc Tukey HSD, **/## $p < 0.01$).

Supplementary methods

MEFs stimulation with poly(I :C) and 3p-hpRNA

For intracellular stimulation with Poly(I:C) or with 3p-hpRNA, MEFs were seeded at 1.10^5 cells per well in 12-well plates the day before stimulation, transfected with 1 µg/mL Poly(I:C) (InvivoGen Cat#vac-pic) or 0.5 µg/mL 3p-hpRNA (InvivoGen Cat#tlrl-hprna) using 5 µL Lipofectamine LTX and 1 µL for poly(I:C) stimulation or 0.5 µL for 3p-hpRNA stimulation of Plus Reagent (ThermoFischer Scientific Cat#15338100), according to the manufacturer's instructions. After stimulation, MEFs were incubated for 8 to 24 hours.

RNA sequencing

MEF RNA was prepared as described in the main text. RNA integrity and quantification were assessed using the RNA Nano 6000 Assay Kit of the Bioanalyzer 2100 system (Agilent Technologies, CA, USA). One microgram of high-quality RNA samples (RIN > 9.2) representing biological triplicates were submitted to Novogene for RNA-sequencing (Novogene Beijing, China). Poly-A selected RNA was used for paired-end library preparation and transcriptome sequencing. Sequencing libraries were generated using NEBNext® UltraTM RNA Library Prep Kit for Illumina® (NEB, USA) following manufacturer's instructions. The library preparations were sequenced on an Illumina platform and paired-end reads were generated. The RNA-seq analysis was performed with Sequana 0.9.8 (1). In particular, we used the RNA-seq pipeline https://github.com/sequana/sequana_rnaseq) built on top of Snakemake 6.1.1 (2). Briefly, reads were trimmed from adapters using Cutadapt 2.7 then mapped to the Mus musculus genome assembly GCA_000001635.8 from NCBI using STAR 2.7.3a (3). FeatureCounts 1.6.4 (4) was used to produce the count matrix, assigning reads to features using corresponding annotation v92 from NCBI with strand-specificity information. Quality control statistics were summarized using MultiQC 1.6 (5). Clustering of transcriptomic profiles were controlled using a Principal Component Analysis (PCA). Differential expression testing was conducted using DESeq2 library 1.24.0 (6) scripts indicating the significance (Benjamini-Hochberg adjusted p-values, false discovery rate FDR < 0.05) and the effect size (fold-change) for each comparison.

1. Cokelaer T, Desvillechabrol D, Legendre R, Cardon M. 2017. "Sequana": a Set of Snakemake NGS pipelines. JOSS 2:352.

2. Köster J, Rahmann S. 2012. Snakemake—a scalable bioinformatics workflow engine. Bioinformatics 28:2520–2522.

3. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR. 2013. STAR: ultrafast universal RNA-seq aligner. Bioinformatics 29:15–21.

4. Liao Y, Smyth GK, Shi W. 2014. featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. Bioinformatics 30:923–930.

5. Ewels P, Magnusson M, Lundin S, Käller M. 2016. MultiQC: summarize analysis results for multiple tools and samples in a single report. Bioinformatics 32:3047–3048.

6. Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biol 15:550.

TabS1

Gene	Forward	Reverse
Ifnb1	CAGCTCCAAGAAAGGACGAAC	GGCAGTGTAACTCTTCTGCAT
Ifna4	TGATGAGCTACTACTGGTCAGC	GATCTCTTAGCACAAGGATGGC
ZIKA *Env*	CCGCTGCCCAACACAAG	CCACTAACGTTCTTTTGCAGACAT
Ifitm3	ACTGTGATCAACATGCCCAGAG	CTTCCGATCCCTAGACTTCACG
Tbp	AGAACAATCCAGACTAGCAGC	GGGAACTTCACATCACAGCTC

TabS2

Immunofluorescence antibodies

Primary/secondary	Antibody	Species	Dilution	Reference
Primary	ZIKV envelop protein	Mouse	1:200	4G2 antibody from ATCC hybridoma
Primary	Phosphorylated IRF3	Rabbit	1:250	Cell Signaling Technology Cat#29047
Secondary	Anti-rabbit AlexaFluor-488-conjugated IgG	Goat	1:500	Invitrogen Cat#A-11034
Secondary	Anti-mouse AlexaFluor-594-conjugated IgG	Donkey	1:500	Invitrogen Cat#A-21203

Western blot antibodies

Primary/secondary	Antibody	Species	Dilution	Reference
Primary	IRF3 Cter	Rabbit	1:1000	Cell Signaling Cat#4302
Primary	Vinculin	Rabbit	1:1000	Proteintech Cat# 26520-1-AP
Secondary	Rabbit IgG HRP Linked Whole Ab	Donkey	1:10,000	Sigma Cat# GENA934-100UL