Using photographs and deep neural networks to understand flowering phenology and diversity in mountain meadows

Aji John [1,3]*, Elli J. Theobald [1], Nicoleta Cristea [2,3], Amanda Tan [3] and Janneke Hille Ris Lambers [1,4]

[1]Department of Biology, University of Washington, Seattle, WA 98195, USA.
[2]Department of Civil and Environmental Engineering, University of Washington, Seattle, WA 98195, USA.
[3]eScience Institute, University of Washington, Seattle, WA 98195, USA.
[4]Plant Ecology, Institute of Integrative Biology, D-USYS, ETH Zürich, 8092 Zürich, Switzerland.

*Corresponding author – ajijohn@uw.edu

ABSTRACT

Mountain meadows are an essential part of the alpine-subalpine ecosystem; they provide

ecosystem services like pollination and are home to diverse plant communities. Changes

in climate affect meadow ecology on multiple levels, for example by altering growing

season dynamics. Tracking the effects of climate change on meadow diversity through the

impacts on individual species and overall growing season dynamics is critical to

conservation efforts. Here, we explore how to combine crowd sourced camera images with

machine learning to quantify flowering species richness across a range of elevations in

alpine meadows located in Mt Rainier National Park, Washington, USA. We employed

three machine learning techniques (Mask R-CNN, RetinaNet and YOLOv5) to detect

wildflower species in images taken during two flowering seasons. We demonstrate that

deep learning techniques can detect multiple species, providing information on flowering

richness in photographed meadows. The results indicate higher richness just above the

tree line for most of the species, which is comparable with patterns found using field

studies. We found that the two-stage detector Mask R-CNN was more accurate than

single-stage detectors like RetinaNet and YOLO, with the Mask R-CNN network

performing best overall with mean average precision (mAP) of 0.67 followed by RetinaNet

(0.5) and YOLO (0.4). We found that across the methods using anchor box variations in

multiples of 16 led to enhanced accuracy. We also show that detection is possible even

when pictures are interspersed with complex backgrounds and are not in focus. We found

differential detection rates depending on species abundance, with additional challenges

related to similarity in flower characteristics, labeling errors, and occlusion issues.

Despite these potential biases and limitations in capturing flowering abundance and

location-specific quantification, accuracy was notable considering the complexity of

flower types and picture angles in this data set. We therefore expect that this approach

can be used to address many ecological questions that benefit from automated flower

detection, including studies of flowering phenology and floral resources, and that this

approach can therefore complement a wide range of ecological approaches (e.g., field

observations, experiments, community science, etc.). In all, our study suggests that

ecological metrics like floral richness can be efficiently monitored by combining machine

learning with easily accessible publicly curated datasets (e.g., Flickr, iNaturalist).

Keywords: Alpine wildflowers; convolutional neural net, climate-change, phenology.

1. INTRODUCTION

Meadows are ecologically important, and are valuable ecosystems found in the montane regions of the world. They are in the transitional zone between forest and alpine areas, and at the elevations that support forbs, sedges, and wildflowers. Functionally, montane meadows offer essential services such as water regulation, carbon sequestration, and erosion prevention (Yongheng 2007, Hopping et al. 2018). They additionally provide habitat for wildlife and are thus crucial for restoration and conservation efforts (Hik et al. 2001, Marini et al. 2007, Rui et al. 2015). Finally, their captivating floral displays also draw visitors and hold cultural significance, particularly for indigenous communities. However, alpine environments housing montane meadows are also uniquely susceptible to climate change (Ganjurjav et al. 2016, Steinbauer et al. 2018).

Montane meadows are particularly sensitive to changes in temperature and precipitation patterns because biological activity is generally limited by a short growing window that is driven by late snowmelt or cold temperatures (Cardinale et al. 2007), making them particularly sensitive warming climate (Inouye 2008). Floral richness is one measure that provides an important yardstick on the fitness of Alpine wildflowers, as meadow species are generally perennial (Roslin et al. 2021). Furthermore, flowers are good indicators for measuring the effects of climate change (Dreyer et al. 2006, Miller-Rushing et al. 2008, Gallagher et al. 2009, Rawal et al. 2015). Studying montane meadows provides insights into ecosystem dynamics, plant and animal adaptations, and climate change effects (Stöcklin et al. 2009, Jiang et al. 2018, Duan et al. 2021). Their sensitivity to changing climate patterns makes them valuable indicators of climate change impacts, with shifts in plant communities and downstream effects offering early warnings.

Phenological shifts in flowering due to climate change could have dramatic

ecological impacts (Gottfried et al. 2012, Anthelme and Dangles 2012) as well as

sociological effects. An extremely warm year might hasten flowering and make flowers

susceptible to frost damage (Klein et al. 2018). On other hand, in a year with late

snowmelt (i.e., after an extremely snowy winter or an unusually chilly spring), flowering

might get delayed, resulting in less time for flowers to develop into fruits. Years with

earlier flowering phenological could also result in a mismatch between pollinators and

flower, if pollinators emerge after peak flowering or earlier before the onset of flowering

(Inouye 2008, Ogilvie et al. 2017, Vorkauf et al. 2021). Furthermore, the experience of

human visitors who visit to enjoy the wildflower displays could be at risk after these

mismatches (Breckheimer et al. 2020, Hille Ris Lambers et al. 2021). Hence, evaluating

patterns of floral richness spatiotemporally is important for understanding the factors

driving meadow species reproductive life cycle (Inouye et al. 2003).

Traditionally, wildflower richness and flowering phenology has been studied with

labor-intensive and time-consuming field observations. Detecting flowers from pictures

taken in natural settings (such as alpine meadows) could provide a solution but is a

complex task that recent developments in computer vision algorithms may aid in.

Isolating flowers in a picture of a typical mountain meadow is challenging because of the

diversity of meadow species, the wide range of floral structures, and because of the

backgrounds in which the flowers exist (like leaves, trees, rocks etc.). Flowers are

inherently complex because of the extraordinary diversity of floral morphologies resulting

from eco-evolutionary dynamics (Moyroud and Glover 2017). However, machine learning

(ML) vision-based algorithms show promise in detecting life stages or phenologies (e.g.,

identifying blooming), and in-depth evaluation of a single phenological stage for

particular species has been successful (e.g. fruit maturity in passion fruit (Tu et al. 2018),

strawberries (Chen et al. 2019) and tomatoes (Wan et al. 2018); ripeness in papayas

(Santos Pereira et al. 2018), and flowering in cotton plants (Jiang et al. 2020)). ML

approaches have also been used for classifying the progression of flowering in apples

(Dias et al. 2018, Wang et al. 2021). These studies showed that it is possible to evaluate

stages of a plant life event (for e.g., fruiting) from images with ML methods. However,

these approaches have not been attempted in more species rich and complex systems like

Alpine wildflower meadows, where issues related to occlusion, complex backgrounds and

a high floral richness may complicate these approaches (Osherov and Lindenbaum 2017,

Picek et al. 2022).

An evolving class of machine learning algorithms called deep neural networks

(dNNs) have been particularly useful for detecting and segmenting multiple objects in an

image. dNNs have found numerous applications in ecological conservation (Clark et al.

2023). For example, advanced dNN algorithms, such as convolutional neural networks

(CNNs) have been deployed for image and audio recognition while recurrent neural

networks (RNNs) have been used to analyze sequence data. Both contribute to

biodiversity monitoring by identifying species from images, audio recordings and genetic

sequences sometimes extracted from the environment (Aide et al. 2013, Khalighifar et al.

2021). These approaches also play a pivotal role in analyzing satellite imagery for land

cover classification and detecting deforestation, aiding in habitat mapping and

environmental monitoring (Chiang et al. 2020, Kattenborn et al. 2021). Thus, dNNs can

be harnessed to predict the impact of climate change on ecosystems, enabling more

accurate modeling of how changing conditions affect a wide variety of biodiversity and

ecosystem services (Anderson 2018). These applications highlight the potential of deep

learning to enhance conservation strategies by extracting insights from complex

ecological data.

Here, we explore the use of dNNs to facilitate conservation planning by assisting

in the species identification from multimodal data (images, in our case). Similar

applications exist, ranging from tree crown detection to identifying cancerous cells

(Abdolali et al. 2020, Hao et al. 2021). Like many of these applications, we employ dNN

methods to perform object detection (for e.g., a flower) and instance segmentation (e.g.,

detect all the instances of a flower). Our use of dNNs for mapping flowers is also motivated

by their ability to extract subtle features and learn from them efficiently (Dias et al. 2018).

This means flowers can be individually tagged and segmented in a meadow image with

dedicated dNNs. Thus, our work contributes to a growing integration of deep learning

techniques into ecological research which can provide opportunities to address crucial

conservation challenges.

Broadly, dNN approaches to identify objects (such as flowers) can be divided into

one-stage and two-stage techniques. One-stage techniques perform object detection in a

single pass by directly predicting object classes and bounding boxes, offering real-time

processing but potentially sacrificing precision for small or overlapping objects. In

contrast, two-stage techniques follow a multi-stage process involving region proposal and

refinement steps, enabling more accurate localization and handling of overlapping objects

at the cost of increased computational complexity and slower processing. A widely used

one-stage algorithm called YOLO (Redmon et al. 2016) is a well-known object instance

detection method, but has issues with objects with multiple sizes (pictures where objects

are taken at different distances), and background noise. RetinaNet is a comparable one-

stage algorithm that overcomes these limitations (Iglovikov and Shvets 2018). However,

these approaches do not fully utilize certain attributes for classification, especially if

objects are close together or small (Zhou et al. 2019). In such cases, a two-stage algorithm

like Mask R-CNN (He et al. 2017) becomes advantageous. By integrating feature

extraction, segmentation, and pixel-level classification, Mask R-CNN has been shown to

achieve higher accuracy in various studies (Jia et al. 2020, Machefer et al. 2020, Hao et

al. 2021).

In this paper, we therefore explore these three state-of-the-art dNN methods

(YOLO, RetinaNet, and Mask R-CNN) for the detection and localization of Alpine

wildflowers in complex settings – i.e., montane meadows. We define identification of a

species by its flowering phenophase only i.e., the stage where the species flower(s) are

visible. Specifically, we focus on two objectives; i) Evaluating the three dNN based

techniques in their ability to detect flowering species in alpine meadows to derive

diversity and abundance and ii) using the data to explore an ecological hypothesis on the

relationship between elevation and floral richness to demonstrate the utility of this

approach. Our expectation is that diversity declines with increasing elevation but in a

non-linear way, and that ML approaches applied to pictures can capture this relationship.

2. MATERIALS AND METHODS

2.1 Study Area

We conducted our study in the subalpine regions (upward of 1400 m) of the Mount Rainier National Park (MORA), which have an impressive wildflower display that is one of the main summertime visitor attractions. Meadows are snow covered most of the year, even late into July, with peak-flowering that lasts between 2 and 3 weeks between late July and early September – depending on snowmelt dates (Theobald et al. 2017). We collected data along an existing network of trails called the Reflections Lakes Trail, which has been a site for focused meadow phenology observations (Hille Ris Lambers et al. 2021, Manzanedo et al. 2022). The trail is on the south side of MORA and covers an elevational gradient of 1400 m to 1950 m (Fig. 1).

Figure 1: Study area at Mt Rainier National Park, with the red dots indicating the sites
along the Reflections Lakes trail at which the pictures were taken for the study (n=27).

172 The inset shows the location (in blue) of the study area in the context of Washington State,
173 USA.
174

2.2   Data

Volunteers participating in this study collected camera images using an iPhone camera

(version 11 Pro) during the flowering season of 2020 and 2021. Sixteen of the 42 species

that commonly occur at the study site were targeted in this study (see Table 1). The subset

was chosen based on species list in use by the community science program MeadoWatch

(Manzanedo et al. 2022). Photos were taken from the trail overlooking the meadows. The

curated dataset contained 1545 high resolution photos: 768 x 1024-pixel (landscape) and

1024 x 768-pixel (portrait). The portrait and landscape images were then tiled for easier

annotation after converting them to a common dimension of 1024 pixel x 1024 pixel.

Padding was done so that it was easier to tile the images in equal sizes. For e.g., tiling into

256 x 256 of a 1024 x 1024 format image results in 8 images. We partitioned the data into

training, test, and validation sets. Specifically, 70% of the data was allocated to the

training set, while the remaining 30% was evenly distributed between the test and

validation sets. In total, we used 1221 images for training and 324 images for validation

and testing.

For our training dataset, we annotated each individual image, identifying the

number and identity of flowers in each image. We used the LabelMe software to perform

the annotation (Russell et al. 2008), which was then outputted in a JSON (JavaScript

Object Notation) file format. These annotations included multiple labels (flowers) with

their annotation mask (see Fig. 2 for sample annotations). For the application of the Mask

R-CNN and the RetinaNet dNN algorithms, the annotation outputs were in JSON format,

which was converted to a COCO format by the training framework. Because the YOLO

algorithms require a specific format (the YOLO format), we converted the LabelMe

annotations to this format prior to analysis.

Table 1: The training dataset comprises a total of 1221 images having 4349 instances of flowers annotated. The validation dataset had a total of 166 images having 597 instances of flowers annotated, and the test dataset consisted of a total of 158 images having 407 instances of flowers annotated. All the images were 256 by 256 pixels and collectively represented 16 wildflower species.

Picture	Common Name	Scientific Name	Instances		
			Training	Validation	Test
	Western Anemone	*Anemone occidentalis*	508	75	24
	Bracted Lousewort	*Pedicularis bracteosa*	119	20	21
	Mountain Daisy	*Erigenon peregrinus*	265	39	27
	Subalpine Lupine	*Lupinus arcticus*	902	108	31
	Glacier Lily	*Erythronium grandiflorum*	103	23	21

Cascade Aster	*Aster ledophyllus*	177	55	20
American Bistort	*Polygonum bistortoides*	322	69	26
Gray's Lovage	*Ligusticum grayi*	123	23	28
North Microseris	*Microseris alpestris*	213	20	28
Scarlet Paintbrush	*Castilleja miniata*	364	21	20
Sharptooth Angelica	*Angelica arguta*	96	22	26
Avalanche Lily	*Erythronium montanum*	129	21	29

	Magenta Paintbrush	*Castilleja parviflora*	638	34	33
	Sitka Valerian	*Valeriana sitchensis*	95	19	27
	Tall Bluebell	*Mertensia paniculata*	145	26	26
	Broadleaf Arnica	*Arnica latifolia*	150	22	20

Figure 2: Example annotated images from the training dataset. (A) Magenta Paintbrush
(*Castilleja parviflora*) flowers (B) A flowering Cascade Aster (*Aster ledophyllus*) (C)
Avalanche lilies (*Erythronium montanum*); particularly found earlier in the season (D)
Cluster of Scarlet Paintbrush (*Castilleja miniata*) in full bloom. Collectively, these images
illustrate the complexity of the machine learning task - with different abundances, photo
angles, and backgrounds.

2.3 Machine Learning Methods

We explored how three dNN algorithms performed in detection and segmentation: YOLO, RetinaNet and Mask R-CNN. These three methods are inherently able to extract features and learn from them efficiently (Dias et al. 2018). In our study, we utilized YOLOv5 (here onwards referred to as YOLO), which solely supports bounding boxes. However, the latest version is YOLOv8 that also supports generation of pixel level mask (segmentation). YOLO and RetinaNet support bounding box detection only, whereas Mask R-CNN and YOLO give the mask of the detected object in addition to the bounding box. We briefly describe the three methods in Appendix A. We used the Detectron2 framework (an implementation of the PyTorch deep learning library) from Facebook Artificial Intelligence Research (FAIR) to train deep neural network models for flower detection for RetinaNet and Mask R-CNN. The Detectron2 has an out-of-the-box implementation of many of the latest CNN algorithms that perform detection and segmentation (Wu et al. 2019). Detectron2 allows writing reproducible code with a standardized way of dealing with input data and annotations. Detectron2 also provides boiler plate code for dataset handling, augmentations, training, visualization, and performance metrics. Augmentation methods enhance the versatility of the model. We applied two augmentation techniques during training: random image flipping and rotation (Xu et al. 2023). All three of these methods i.e., YOLO, RetinaNet, and Mask R-CNN apply bounding box detection (i.e., localize the object), but Mask R-CNN and the recent version of YOLO (v8), in addition, gives pixel level masks of the class. In addition, Mask R-CNN supports pluggable feature extraction network backbone like U-Net (Ronneberger et al. 2015) that could cover a wide range of domains. The development, training and inference

240 of the models were performed on Azure Cloud on a server having the configuration

241 Standard NC12_Promo (12 vCPUs, 112 GiB memory with 2 NVIDIA Tesla K80 GPUs).

2.4 Hyperparameter tuning

We performed hyperparameter tuning using the validation data composed of 597

instances (compared with 4349 instances used in training). Hyperparameter fine-tuning

of all three models was achieved with training for 1500 iterations on Google Colab's K80

GPU. Anchors are predefined with aspect ratio and scale to establish the location and size

of the bounding box for an object. We used different anchor sizes (Kalantar et al., 2020)

for the three algorithms. We kept the learning rate at the default value of 0.00025 with a

batch size of 128. We also kept the aspect ratio of 1:1 and scale to double with a total of 5

increments.

2.5 Evaluation Metrics

We used detection evaluation metrics as defined by the COCO dataset (Lin et al., 2014).

Intersection over Union (IoU) is a metric that evaluates how the ground truth and

predicted detections overlap. IoU of 1 means an ideal detection whereby there is an exact

overlap whereas IoU of zero means that there is no overlap. Average Precision (AP)

indicates how a detection method performed at a particular IoU. Here, we report AP at

0.5, 0.75 and 1 (as in Table 2). AP is calculated by taking average of AP of ten IoU

thresholds (t= [0.5,0.55,0.60,0.65,0.70,0.75,0.80,.0.85,90,.95] and when averaged

across all categories the metric provides us mean average precision (mAP). True positive

(TP) would mean IoU > = Threshold, and false positive (FP) is IoU < Threshold. We

further inspect the Precision (TP/TP+FP) and Recall (TP/TP+FN) to evaluate frequency

of false detections or missed ones; note that higher mAP indicates better detection

performance. Average Precision (AP) and Average Recall (AR) is further used to calculate

the F-1 score which helps with interpretability of the techniques. Mask R-CNN identifies

the anchor boxes based on the IoU, which is a critical step in the first part of the network

(RPN).

267
For each photograph, we computed two floral indices: species diversity, as

quantified by Shannon's diversity index, and species richness. The Shannon diversity

index was determined using the formula $H = -\sum_{k=0}^{n} p_i \ln p_i$, where p_i represents the

proportion of a specific flowering species in the photograph (Shannon 1948). Species

richness was determined as the count of distinct flowering species within a given

photograph.

3. RESULTS

3.1   Hyperparameter tuning results

We found that maintaining anchor box variations that are multiple of 16 i.e.,

[16,32,64,128,256] yielded better accuracy (Table 2). The results were consistent with

YOLO and RetinaNet (Appendix C).

Table 2: Results of anchor box variation with Mask R-CNN.

Anchor Box Size	AP	AP50	AP75	AP	AP50	AP75
	Bounding Box			Segmentation		
[4, 8, 16, 32, 64]	11.708	23.180	10.338	12.230	22.296	12.217
[8, 16, 32, 64, 128]	13.515	24.992	12.023	13.340	23.783	13.358
[16, 32, 64, 128, 256]	18.128	31.767	17.148	17.871	30.838	20.211

| [32, 64, 128, 256,512] | 17.485 | 31.555 | 16.395 | 17.031 | 30.935 | 16.596 |

3.2 Performance and training time of the flower detection methods

We tested the three dNNs algorithms using over 316 images with an average number of 40 labels per flower category. We used two augmentations in training: Resize Shortest Edge and Random Flip. Each of the augmentation methods were trained for 10000 epochs. The time needed for training differed between the three algorithms, ranging from 3 hours for YOLO to 12 hours in Mask R-CNN.

Table 3 shows the mAPs and different IoU thresholds for all the methods. In addition, Mask R-CNN provides a classification metric by identified object size as well. In our case, for small, medium, and large flowers, the mAP was 29.27, 42.76 and 48.80, respectively. Bounding box detection applies to all the methods, but segmentation (drawing of masks) only applies to Mask R-CNN, so segmentation scores are only reported for Mask R-CNN.

Table 3: Evaluation results for the BBox and Segmentation across the three algorithms. Mean Average precision (mAP) for the three methods. All three algorithms were trained for 10000 iterations (epochs).

	BBox			Seg				
Method	mAP	mAP50	mAP75	mAP	mAP50	mAP75	F-1 Score	Training time (hours)
Mask R-CNN	45.80	67.24	52.10	31.32	59.05	30.78	0.20	12
RetinaNet	23.59	51.28	18.63	N/A	N/A	N/A	0.12	8
YOLO	32.66	60.68	31.91	N/A	N/A	N/A	0.23	3

297   Table 4: Per category BBox evaluation (using mAP) results across all three methods.

	AP50		
Flower	Mask R-CNN	RetinaNet	YOLO
Western Anemone	58.45	35.22	27.90
American Bistort	42.79	23.55	56.80
Avalanche Lily	40.30	18.74	52.2
Bracted Lousewort	58.05	18.28	4.42
Gray's Lovage	36.02	20.07	43.90
Magenta Paintbrush	42.01	29.98	27.80
Mountain Daisy	51.42	23.31	67.70
North Microseris	68.22	39.10	77.80
Sitka Valerian	45.43	21.31	10.40
Subalpine Lupine	38.36	24.67	39.20
Scarlet Paintbrush	59.91	33.00	74.60
Tall Bluebell	46.04	14.45	83.00
Glacier Lily	29.93	11.56	14.70
Sharptooth Angelica	24.34	08.11	48.30
Broadleaf Arnica	61.09	21.80	41.50
Cascade Aster	26.83	34.34	07.62

All three methods were able to isolate the flowers within the visual complexity of a

meadow. Table 4 gives the breakdown of mAP by species for all the three methods. On

average, Mask R-CNN performs better for all species, but we note differences in

identifying individual species across the methods. For example, Scarlet Paintbrush, Tall

Bluebell and North Microseris are detected more accurately by YOLO than by Mask R-

CNN or RetinaNet. Regardless of the species, they are detected with higher accuracy if

they are towards the foreground of the image and if the image is in focus - see for example

Bracted Lousewort being isolated by Mask R-CNN while two species in the horizon are

missed (Fig. 3A, B). Similarly, occluded flowers are being missed for example some of the

Mountain Daisy flowers that are behind the stalks of other species (Fig. 3C, D). The

sample prediction results for YOLO and RetinaNet are in Appendix B.

Figure 3: Predictions from the Mask R-CNN method showing the fluency of the model; the percent next to the predicted class represents model confidence. (A and B) Meadow in peak flowering with predictions (B) corresponding original tile. (C and D) Another meadow in peak flowering with predictions (D) corresponding to the original tile.

3.3 Richness and abundance of flowering species

We found that floral species richness and abundance per photo varies along the elevational gradient. Specifically, we see lower floral richness at lower and higher elevations, and increased richness along mid elevations i.e., 1700-1800 m (Mask R-CNN shown in Fig. 4A). YOLO aligns with Mask R-CNN on elevation and richness relationship (Fig. 4A, C and Appendix D), but RetinaNet does not suggest any association (Fig 4E), We also see increased density of flowers at the mid-elevation compared to lower and higher elevations (Fig. 4B). RetinaNet showing elevated density of flowers compared to Mask R-CNN and YOLO (Fig. 4B, D, F). Floral abundance in pictures was also found to vary over time, even over very short periods with similar overestimation by RetinaNet (Fig. 5).

We found a significant and positive correlation between diversity, determined through annotations, and diversity as assessed by predictions from dNN methods using a separate labeled dataset (correlation coefficient R = 0.62 for Mask R-CNN, R = 0.52 for YOLO, and R = 0.1 for RetinaNet, all with a p-value of less than 0.001; refer to Fig. 6A). Among the dNN methods, this correlation in terms of species richness was evident for Mask R-CNN and YOLO (correlation coefficient R = 0.36 for Mask R-CNN, and R = 0.42 for YOLO, both with a p-value of less than 0.001; see Fig. 6B), but such a correlation was not observed for RetinaNet. Moreover, there existed a correlation between Shannon diversity and species richness of flowers when comparing Mask R-CNN and YOLO (correlation coefficient R = 0.51 for Shannon diversity and R = 0.8 for species richness, both with a p-value of less than 0.001; refer to Fig. 6C and Fig. 6D, respectively).

Figure 4: Detected species from all the collections using images collected in August 2020; predictions are from Mask R-CNN, YOLO and RetinaNet methods. Number of unique flower species detected per photo (A) by Mask R-CNN, (C) by YOLO, and by RetinaNet (E). Number of occurrences of each flower species per picture (B) by Mask R-CNN, (D) by YOLO, and (E) by RetinaNet.

346
347 Figure 5: Floral density of species by picture for all the collection dates in August 2020
348 from all three methods, (A) by Mask R-CNN, (B) YOLO (C) RetinaNet.

349
350
Figure 6: Comparison of diversity and richness using a labelled hold-out dataset across
all three methods, (A) Diversity, (B) Richness, (C) Diversity between YOLO and Mask R-
CNN, and (D) Richness between YOLO and Mask R-CNN. Inset showing Pearson's
correlation coefficient, and the corresponding p-value.

4. DISCUSSION

We demonstrate that deep learning techniques can detect, identify, and count Alpine

wildflowers in photographs, and thereby provide detailed information on flowering

occurrences in complex mountain meadow systems. In general, Mask R-CNN
demonstrates superior performance; however, YOLO and RetinaNet exhibit better results
for specific flower cases. Furthermore, we show that these approaches verify known
ecological patterns – that species flowering varies temporally and by elevation at our
study site (Kudo and Suzuki 1999, Prevéy 2020). Our results show that with timestamped,
geo-coded photographs, deep learning techniques can identify and count flowers,
potentially facilitating detailed analyses on flowering richness as well as floral abundance,
both key ecological metrics relevant to Alpine wildflower fitness and pollinator
performance.

The relative performance of the three dNN methods we tested depended on task-
specific requirements, performance-tradeoffs, and the necessity for detailed instance
segmentation. For object detection and instance segmentation, YOLO, and Mask R-CNN
stand out as equally implementable methodologies, with variation in performance.
YOLO's real-time efficiency derives from its single-pass approach, making it well-suited
for rapid detection tasks; however, challenges with this approach arise in accurately
identifying small and densely overlapping objects. RetinaNet addresses scale variability
by incorporating anchor boxes and feature pyramid networks, finding equilibrium
between speed and precision, though it might not excel in intricate instance
segmentation. In contrast, Mask R-CNN's offers unparalleled pixel-level accuracy for
object separation but demands greater computational resources and time. We discuss
these points in more detail below.

4.1 Differences in performance and methodologies for identifying flowers in camera images

In our application, the two-stage detector (Mask-R-CNN) was more accurate but slower in its predictions than the one stage detectors (YOLO, RetinaNet). This is consistent with other studies using this approach to count and identify individual flowers of a single species (Yu et al. 2019, Machefer et al. 2020, Hao et al. 2021). Single stage detectors perform classification and regression together on the predicted anchor boxes. The trade-off is that adding an additional stage incurs computation time (Suh et al. 2021). The YOLO algorithm which is a single stage detector is able to accurately draw boxes around the flowers as well as classify them correctly more speedily than Mask R-CNN (similar findings are presented in ((Prasetyo et al. 2020) and (Sumit et al. 2020)). However, it is crucial to emphasize the significance of annotations. Predictions made within images might inadvertently miss numerous instances of flowers, a situation expected due to the inherent complexity in annotating such occurrences during the training phase.

The images acquired for our study were in two dimensions (1024 x 768 and 768 x 1024). However, we found that breaking down these extensive format images into 256 x 256-pixel images resulted in the lowest classification error. The 256x256 images seem to offer sufficient dimensions for detecting objects like flowers of similar size within the dataset. We observed that dealing with larger format image samples demanded more computational resources. While analyzing the metrics, it is important to consider that larger images lead to higher occurrences of box and objectness errors. This is because sub-images are extracted from the same larger image, and if the sub-image is larger, it will contain more boxes within it (Safaldin et al. 2023). As these errors are aggregated for each

potential box, the cumulative box and objectness error tends to be greater for larger
images.

While our deep learning study was done with a minimal dataset, the initial results
are promising. Augmenting the dataset through more annotation is apparent, but the
resource-intensive nature of this step needs to be weighed against computation and
resource availability. Moreover, exploring algorithms designed for efficacy under weak
supervision (heuristically labeled data) could be a potential avenue for future
enhancement (Phan et al. 2023).

4.2   Flower characteristics, species abundance and implications for
detection

Several factors led to variation in our ability to classify and detect flowers. Most
wildflower meadows (and thus pictures) include large variations in the abundance of
species, like Subalpine Lupine, which tends to be more abundant than other flowers. This
results in Subalpine Lupine being detected more than other species like Mountain Daisy
which appears more sparsely, a finding that reflects real differences in floral abundance.
However, the algorithm also makes mistakes in species identity, and it is likely that the
similar characteristics of some flowers are responsible. For example, some meadow
wildflowers have an elongated floral structure (e.g., Bracted Lousewort and Subalpine
Lupine), while others are more circular in nature (e.g., Mountain Daisy and Cascade
Aster). Misclassification tended to occur when structures and color were similar (e.g.,
Sharptooth Angelica and Gray's Lovage). Unlike in other studies, in which data
enhancement helped prevent overfitting and improved accuracy (Tian et al. 2020), we did

not see improvement in accuracy with image augmentations. We also see the shortcoming

where occluded flowers were not being isolated (Fig. 3A, C).

Despite these challenges, we were encouraged by the accuracy we achieved in this

study as these pictures are inherently complex and therefore challenging for classifiers

(Fig. 6). Some challenges include inconsistent light levels (e.g., because of slope and

aspect) and heterogenous backgrounds (e.g., rocks, lots of plants, etc.). Additionally,

labeling errors, namely flowers that are missed while annotating could contribute to

decreased accuracy. For example, the YOLO method penalizes incorrectly drawn boxes -

if a flower is not labeled in the training set, YOLO would penalize the model for drawing

a box around the flower despite it being correct. These challenges are all ones that could

be addressed in studies that standardize photo acquisition and automation and increase

annotation effort, both factors feasible for many ecological studies.

There also is likely to be sampling bias in this study. In the iconic wildflower

meadows of Mt. Rainier National Park, the chances of picking a location and

photographing it is almost certainly positively correlated with the probability of a flower

existing in that place. That means that our pictures are disproportionately likely to

contain flowers, potentially predisposing our algorithm to bias toward flowers. Another

limitation of our study is that it did not capture the full flowering season and does not

quantify the flowers by location (e.g., elevation). Fortunately, these issues are easily

addressed with a planned study, for example with images are captured from intentionally

placed PhenoCams along an elevational gradient.

4.3   Applicability to ecological research

We believe that combining image analysis with automated floral detection can

provide important information on metrics of ecological interest, which can either

complement or in some cases even replace more traditional ecological sampling. For

example, our analyses show that floral richness and abundance are linked to elevation in

a non-linear way (Fig. 4A, B), similar to patterns shown by other studies (Fierer et al.

2011, Guo et al. 2013, Antonelli et al. 2018, Sponsler et al. 2022). Floral richness is driven

both by plant diversity (which is influenced by many biotic and abiotic factors), as well as

by the probability that species are flowering (Fabbro and Körner 2004, Sargent and

Ackerly 2008, Theobald et al. 2017) – an important indicator of plant fitness resources

for pollinators. Thus, this approach could be used to provide information on diversity

patterns across gradients, phenological patterns of flowering relative to climate, and floral

resources for pollinators.

For abundant species with recognizable flowers, these approaches may also allow

for the detection of distributional shifts, if conducted annually. For example, climate

change might alter the distribution of specific wildflowers found in montane

environments; possibly as the result of elevational shifts (Gottfried et al. 2012, Steinbauer

et al. 2018). Similarly, a rapid invasion of a non-native flowering species can be detected

using this approach. The combination of the floral detection approaches introduced here

with drone imagery could allow for assessment of such shifts over large spatial areas, or

in remote areas.

However, there are some caveats to using flowers in pictures to quantify richness

and / or abundance. For one, floral richness is not total species richness, as many

perennial plants do not flower every year or in every location. Moreover, different species

produce differing number of flowers per individual, meaning that a higher number of

flowers need not mean a greater density of individuals. A second important caveat in our

measures of floral abundance is that fields of view vary from picture to picture, which

means the total area available to flowers (and thus flowering abundance) will vary even if

underlying patterns detected by more traditional ecological approaches (e.g., m²

quadrats) do not.

In total, however, we believe the promise of these approaches for generating large

volumes of data with less manpower (as described in (Wilson et al. 2017)) outweigh these

issues. Moreover, these challenges can be overcome with large volumes of pictures / data

or by employing fixed cameras (e.g., webcams, PhenoCams) with common fields of view.

For example, if pictures are taken automatically via PhenoCams or TrailCams, these fixed

cameras allow for pictures to be taken in the same place repeatedly, which saves time and

gives an advantage over taking them oneself as one gets to control when and where the

pictures are to be taken. Then, differences in floral richness would not be due to

differences in field of view (the way the picture was taken). Regardless, we feel that floral

richness derived from photos is still of interest as it gives a quick proxy of site-specific

flowering patterns.

We also believe our approach can also be used to address questions about

phenology that otherwise would not be possible with traditional ecological sampling. As

photos are engrained with date/time and location information, we could also have linked

these measures to time to provide information on phenology (as in (Breckheimer et al.

2020)). In this case, crowd sourced pictures (e.g., Flickr, iNaturalist, Instagram)

increased the data volume (# of observations) and allowed for quantification of phenology

in previous years (i.e., going back in time). This and similar approaches, however, require

specific statistical approaches. For example, crowd sourced pictures (e.g., from Flickr) are

not taken according to a specific experimental design, which means using these pictures

could result in an unbalanced representation – e.g., spatial or temporal gaps in

distributions (e.g., Mt. Rainier meadow pictures predominantly may be from around

visitor centers, weather permitting times and might have weekend effects). However, one

can deal with skewness introduced by source images by statistically controlling for any

spatiotemporal bias in pictures taken (Breckheimer et al. 2020).

Finally, we believe that the floral detection combined with ML approaches could

allow for quick assessment of ecological patterns, which could be important in a changing

world. For example, it might be valuable to know about the distribution of a rare species

threatened by climate change and / or the co-occurrence of wildflowers that compete for

pollinators whose phenology shift at different rates or due to climate driven changes in

species composition (Taheri et al. 2021). Documenting species presence-absence using

any of our methods as applied to existing (crowd-sourced) pictures could address

questions such as these and provide a preliminary assessment of patterns before

designing new (and costly) experiments (Elliott and Davies 2019).

4.4   Future directions

Although our flower detection algorithm performed well, there are likely improvements

that could be made to detect a greater number of wildflowers and better distinguish

between similar looking flowers. For example, we have not looked at the fluorescence

signal but hypothesize that shape and color are being learnt during feature extraction, and

as future directions, incorporation of additional information (e.g., thermal, or infrared

bands) might help in the presented algorithms.

Embedding the trained models in community science apps (Mobile Applications)

could also be one of the end uses of these algorithms. In other domains, apps have

increased community involvement, e.g., the bird identification app, Merlin (Nugent 2018)

and the plant identification functionality of iNaturalist. Use in apps both provides the

public with a way to identify flowers and a way to gather data.

Combining photo data with drone and remotely sensed products could potentially

offer opportunities to measure things in remote areas and on larger spatial scales.

Improvement of algorithms that can use additional bands (produced in hyperspectral or

multispectral imagery) could be promising and might detect more kinds of flowers and

distinguish between flowers that are more similar (Wieme et al. 2022).

5. CONCLUSIONS

We explored three deep learning techniques to detect and count flowering plants in alpine

meadows. The two-stage detector we used (Mask R-CNN) performed better than the

single-stage detectors (YOLO and RetinaNet), with the first stage cropping flowers and

filtering out the background; and the second stage classifying and detecting the flowers.

However, all techniques gave us detailed information on flowering occurrences in

meadows, allowing us to calculate floral richness as well as flower abundance. These

approaches were also able to capture known ecological patterns; namely that flowering

varies temporally (i.e., phenology) and by elevation. While our trained models exhibit

promise for application in areas dominated by our focal species from Mt. Rainier National Park, caution should be exercised considering potential variations in local ecological dynamics. Looking forward, further investigations into the suitability and transferability of these models to subalpine regions across the broader Pacific Northwest (such as Washington, Oregon, and British Columbia) will be essential for robust and regionally informed study efforts. In the future, including a larger catalog of species for training could allow for coverage from other areas, for example Rocky Mountain meadows or Swiss Alpine meadows. More generally, these approaches could aid in climate change studies, because they can be used in a non-invasive, low-investment way to rapidly get information on spatiotemporal flowering patterns.

Acknowledgements

We want to acknowledge support of the University of Washington eScience Institute and Hille Ris Lambers lab in developing the ideas and supporting discussions. Authors want to acknowledge assistance from John H., Sapti S., Tommy L., and Richard L. Cristea was supported by the NSF award OAC-2117834 and the University of Washington eScience Institute. We also acknowledge support from Microsoft AI for Earth and Microsoft Azure collegial support. This study was carried out on the ancestral lands of the people of the Nisqually, Puyallup, Squaxin Island, Muckleshoot, Yakama, and Cowlitz Nations.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Authors' contributions

A.J. and J.H.R.L. conceived the ideas; A.J., J.H.R.L., N.C., A.T. designed the methodology; A.J. and MeadoWatch volunteers collected the data; A.J. analyzed the data; A.J., E.J.T. and J.H.R.L. led the writing of the manuscript. All authors contributed critically to the drafts and gave final approval for publication.

Data Availability Statement

All code will be made available on public GitHub repositories and will be archived in Zenodo.

570
References

Abdolali, F., J. Kapur, J. L. Jaremko, M. Noga, A. R. Hareendranathan, and K. Punithakumar. 2020. Automated thyroid nodule detection from ultrasound imaging using deep convolutional neural networks. Computers in Biology and Medicine 122:103871.

Aide, T. M., C. Corrada-Bravo, M. Campos-Cerqueira, C. Milan, G. Vega, and R. Alvarez. 2013. Real-time bioacoustics monitoring and automated species identification. PeerJ 1:e103.

Anderson, C. B. 2018. Biodiversity monitoring, earth observations and the ecology of scale. Ecology Letters 21:1572–1585.

Anthelme, F., and O. Dangles. 2012. Plant–plant interactions in tropical alpine environments. Perspectives in Plant Ecology, Evolution and Systematics 14:363–372.

Antonelli, A., W. D. Kissling, S. G. A. Flantua, M. A. Bermúdez, A. Mulch, A. N. Muellner-Riehl, H. Kreft, H. P. Linder, C. Badgley, J. Fjeldså, S. A. Fritz, C. Rahbek, F. Herman,

H. Hooghiemstra, and C. Hoorn. 2018. Geological and climatic influences on mountain biodiversity. Nature Geoscience 11:718–725.

Breckheimer, I. K., E. J. Theobald, N. C. Cristea, A. K. Wilson, J. D. Lundquist, R. M. Rochefort, and J. HilleRisLambers. 2020. Crowd-sourced data reveal social–ecological mismatches in phenology driven by climate. Frontiers in Ecology and the Environment 18:76–82.

Cardinale, B. J., J. P. Wright, M. W. Cadotte, I. T. Carroll, A. Hector, D. S. Srivastava, M. Loreau, and J. J. Weis. 2007. Impacts of plant diversity on biomass production increase through time because of species complementarity. Proceedings of the National Academy of Sciences 104:18123–18128.

Chen, Y., W. S. Lee, H. Gan, N. Peres, C. Fraisse, Y. Zhang, and Y. He. 2019. Strawberry Yield Prediction Based on a Deep Neural Network Using High-Resolution Aerial Orthoimages. Remote Sensing 11:1584.

Chiang, C.-Y., C. Barnes, P. Angelov, and R. Jiang. 2020. Deep learning-based automated forest health diagnosis from aerial images. IEEE Access 8:144064–144076.

Clark, M. L., L. Salas, S. Baligar, C. A. Quinn, R. L. Snyder, D. Leland, W. Schackwitz, S. J. Goetz, and S. Newsam. 2023. The effect of soundscape composition on bird vocalization classification in a citizen science biodiversity monitoring project. Ecological Informatics 75:102065.

Dias, P. A., A. Tabb, and H. Medeiros. 2018. Apple flower detection using deep convolutional networks. Computers in Industry 99:17–28.

Dreyer, L. L., K. J. Esler, and J. Zietsman. 2006. Flowering phenology of South African

Oxalis—possible indicator of climate change? South African Journal of Botany 72:150–

156.

Duan, H., X. Xue, T. Wang, W. Kang, J. Liao, and S. Liu. 2021. Spatial and Temporal

Differences in Alpine Meadow, Alpine Steppe and All Vegetation of the Qinghai-Tibetan

Plateau and Their Responses to Climate Change. Remote Sensing 13:669.

Elliott, T. L., and T. J. Davies. 2019. A comparison of phylogenetic and species beta diversity

measures describing vegetation assemblages along an elevation gradient. Journal of

Vegetation Science 30:98–107.

Fabbro, T., and C. Körner. 2004. Altitudinal differences in flower traits and reproductive

allocation. Flora - Morphology, Distribution, Functional Ecology of Plants 199:70–81.

Fierer, N., C. M. McCain, P. Meir, M. Zimmermann, J. M. Rapp, M. R. Silman, and R. Knight.

2011. Microbes do not follow the elevational diversity patterns of plants and animals.

Ecology 92:797–804.

Gallagher, R. V., L. Hughes, and M. R. Leishman. 2009. Phenological trends among Australian

alpine species: using herbarium records to identify climate-change indicators. Australian

Journal of Botany 57:1.

Ganjurjav, H., Q. Gao, E. S. Gornish, M. W. Schwartz, Y. Liang, X. Cao, W. Zhang, Y. Zhang,

W. Li, Y. Wan, Y. Li, L. Danjiu, H. Guo, and E. Lin. 2016. Differential response of

alpine steppe and alpine meadow to climate warming in the central Qinghai–Tibetan

Plateau. Agricultural and Forest Meteorology 223:233–240.

Gottfried, M., H. Pauli, A. Futschik, M. Akhalkatsi, P. Barančok, J. L. Benito Alonso, G. Coldea,

J. Dick, B. Erschbamer, M. R. Fernández Calzado, G. Kazakis, J. Krajči, P. Larsson, M.

628  Mallaun, O. Michelsen, D. Moiseev, P. Moiseev, U. Molau, A. Merzouki, L. Nagy, G.

629  Nakhutsrishvili, B. Pedersen, G. Pelino, M. Puscas, G. Rossi, A. Stanisci, J.-P. Theurillat,

630  M. Tomaselli, L. Villar, P. Vittoz, I. Vogiatzakis, and G. Grabherr. 2012. Continent-wide

631  response of mountain vegetation to climate change. Nature Climate Change 2:111–115.

632 Guo, Q., D. A. Kelt, Z. Sun, H. Liu, L. Hu, H. Ren, and J. Wen. 2013. Global variation in

633  elevational diversity patterns. Scientific Reports 3:3007.

634 Hao, Z., L. Lin, C. J. Post, E. A. Mikhailova, M. Li, Y. Chen, K. Yu, and J. Liu. 2021.

635  Automated tree-crown and height detection in a young forest plantation using mask

636  region-based convolutional neural network (Mask R-CNN). ISPRS Journal of

637  Photogrammetry and Remote Sensing 178:112–123.

638 He, K., G. Gkioxari, P. Dollár, and R. Girshick. 2017. Mask r-cnn. Pages 2961–2969

639  Proceedings of the IEEE international conference on computer vision.

640 Hik, D. S., C. J. McColl, and R. Boonstra. 2001. Why are Arctic ground squirrels more stressed

641  in the boreal forest than in alpine meadows? Écoscience 8:275–288.

642 Hille Ris Lambers, J., A. F. Cannistra, A. John, E. Lia, R. D. Manzanedo, M. Sethi, J. Sevigny,

643  E. J. Theobald, and J. K. Waugh. 2021. Climate change impacts on natural icons: Do

644  phenological shifts threaten the relationship between peak wildflowers and visitor

645  satisfaction? Climate Change Ecology 2:100008.

646 Hopping, K. A., A. K. Knapp, T. Dorji, and J. A. Klein. 2018. Warming and land use change

647  concurrently erode ecosystem services in Tibet. Global Change Biology 24:5534–5548.

648 Iglovikov, V., and A. Shvets. 2018. TernausNet: U-Net with VGG11 Encoder Pre-Trained on

649  ImageNet for Image Segmentation. arXiv:1801.05746 [cs].

Inouye, D. W. 2008. EFFECTS OF CLIMATE CHANGE ON PHENOLOGY, FROST

DAMAGE, AND FLORAL ABUNDANCE OF MONTANE WILDFLOWERS. Ecology

89:353–362.

Inouye, D. W., F. Saavedra, and W. Lee-Yang. 2003. Environmental influences on the

phenology and abundance of flowering by Androsace septentrionalis (Primulaceae).

American Journal of Botany.

Jia, W., Y. Tian, R. Luo, Z. Zhang, J. Lian, and Y. Zheng. 2020. Detection and segmentation of

overlapped fruits based on optimized mask R-CNN application in apple harvesting robot.

Computers and Electronics in Agriculture 172:105380.

Jiang, S., Y. Liu, J. Luo, M. Qin, N. C. Johnson, M. Öpik, M. Vasar, Y. Chai, X. Zhou, L. Mao,

G. Du, L. An, and H. Feng. 2018. Dynamics of arbuscular mycorrhizal fungal community

structure and functioning along a nitrogen enrichment gradient in an alpine meadow

ecosystem. New Phytologist 220:1222–1235.

Jiang, Y., C. Li, R. Xu, S. Sun, J. S. Robertson, and A. H. Paterson. 2020. DeepFlower: a deep

learning-based approach to characterize flowering patterns of cotton plants in the field.

Plant Methods 16:156.

Kattenborn, T., J. Leitloff, F. Schiefer, and S. Hinz. 2021. Review on Convolutional Neural

Networks (CNN) in vegetation remote sensing. ISPRS Journal of Photogrammetry and

Remote Sensing 173:24–49.

Khalighifar, A., R. M. Brown, J. Goyes Vallejos, and A. T. Peterson. 2021. Deep learning

improves acoustic biodiversity monitoring and new candidate forest frog species

identification (genus Platymantis) in the Philippines. Biodiversity and Conservation

30:643–657.

Klein, G., M. Rebetez, C. Rixen, and Y. Vitasse. 2018. Unchanged risk of frost exposure for

674   subalpine and alpine plants after snowmelt in Switzerland despite climate warming.

675   International Journal of Biometeorology 62:1755–1762.

676   Kudo, G., and S. Suzuki. 1999. Flowering phenology of alpine plant communities along gradient

677   of snowmelt timing. Polar Bioscience 12:100–113.

678   Li, Y., and F. Ren. 2019. Light-weight retinanet for object detection. arXiv preprint

679   arXiv:1905.10011.

680   Lin, T.-Y., P. Goyal, R. Girshick, K. He, and P. Dollár. 2017. Focal loss for dense object

681   detection. Pages 2980–2988 Proceedings of the IEEE international conference on

682   computer vision.

683   Machefer, M., F. Lemarchand, V. Bonnefond, A. Hitchins, and P. Sidiropoulos. 2020. Mask R-

684   CNN Refitting Strategy for Plant Counting and Sizing in UAV Imagery. Remote Sensing

685   12:3015.

686   Manzanedo, R. D., A. John, M. L. Sethi, E. J. Theobald, B. Brosi, J. Jenkins, A. Kloss-Schmidt,

687   E. Lia, A. Schiffer, J. Sevigny, A. Wilson, Y. Yogev, and J. Hille Ris Lambers. 2022.

688   MeadoWatch: a long-term community-science database of wildflower phenology in

689   Mount Rainier National Park. Scientific Data 9:151.

690   Marini, L., M. Scotton, S. Klimek, J. Isselstein, and A. Pecile. 2007. Effects of local factors on

691   plant species richness and composition of Alpine meadows. Agriculture, Ecosystems &

692   Environment 119:281–288.

693   Miller-Rushing, A. J., D. W. Inouye, and R. B. Primack. 2008. How well do first flowering dates

694   measure plant responses to climate change? The effects of population size and sampling

695   frequency. Journal of Ecology 96:1289–1296.

Moyroud, E., and B. J. Glover. 2017. The Evolution of Diverse Floral Morphologies. Current

Biology 27:R941–R951.

Nugent, J. 2018. Citizen Science: Birds, Binoculars, and Biodiversity. Science Scope 041.

Ogilvie, J. E., S. R. Griffin, Z. J. Gezon, B. D. Inouye, N. Underwood, D. W. Inouye, and R. E.

Irwin. 2017. Interannual bumble bee abundance is driven by indirect climate effects on

floral resource phenology. Ecology Letters 20:1507–1515.

Osherov, E., and M. Lindenbaum. 2017. Increasing CNN Robustness to Occlusions by Reducing

Filter Support. Pages 550–561 2017 IEEE International Conference on Computer Vision

(ICCV). IEEE, Venice.

Phan, S., D. Torrejon, J. Furseth, E. Mee, and C. Luscombe. 2023. Exploiting weak supervision

to facilitate segmentation, classification, and analysis of microplastics (<100 μm) using

Raman microspectroscopy images. Science of The Total Environment 886:163786.

Picek, L., M. Šulc, Y. Patel, and J. Matas. 2022. Plant recognition by AI: Deep neural nets,

transformers, and kNN in deep embeddings. Frontiers in Plant Science 13:787527.

Prasetyo, E., N. Suciati, and C. Fatichah. 2020. A Comparison of YOLO and Mask R-CNN for

Segmenting Head and Tail of Fish. Pages 1–6 2020 4th International Conference on

Informatics and Computational Sciences (ICICoS).

Prevéy, J. S. 2020. Climate Change: Flowering Time May Be Shifting in Surprising Ways.

Current Biology 30:R112–R114.

Rawal, D. S., S. Kasel, M. R. Keatley, and C. R. Nitschke. 2015. Herbarium records identify

sensitivity of flowering phenology of eucalypts to climate: Implications for species

response to climate change: Herbarium Records Identify Phenological Shift. Austral

Ecology 40:117–125.

Redmon, J., S. Divvala, R. Girshick, and A. Farhadi. 2016. You Only Look Once: Unified, Real-

Time Object Detection. arXiv:1506.02640 [cs].

Redmon, J., and A. Farhadi. 2017. YOLO9000: Better, Faster, Stronger. Pages 6517–6525 2017

IEEE Conference on Computer Vision and Pattern Recognition (CVPR). IEEE,

Honolulu, HI.

Redmon, J., and A. Farhadi. 2018. YOLOv3: An Incremental Improvement. arXiv:1804.02767

[cs].

Ren, S., K. He, R. Girshick, and J. Sun. 2017. Faster R-CNN: Towards Real-Time Object

Detection with Region Proposal Networks. IEEE Transactions on Pattern Analysis and

Machine Intelligence 39:1137–1149.

Ronneberger, O., P. Fischer, and T. Brox. 2015. U-net: Convolutional networks for biomedical

image segmentation. Pages 234–241 International Conference on Medical image

computing and computer-assisted intervention. Springer.

Roslin, T., L. Antão, M. Hällfors, E. Meyke, C. Lo, G. Tikhonov, M. del M. Delgado, E.

Gurarie, M. Abadonova, O. Abduraimov, O. Adrianova, T. Akimova, M. Akkiev, A.

Ananin, E. Andreeva, N. Andriychuk, M. Antipin, K. Arzamascev, S. Babina, M.

Babushkin, O. Bakin, A. Barabancova, I. Basilskaja, N. Belova, N. Belyaeva, T.

Bespalova, E. Bisikalova, A. Bobretsov, V. Bobrov, V. Bobrovskyi, E. Bochkareva, G.

Bogdanov, V. Bolshakov, S. Bondarchuk, E. Bukharova, A. Butunina, Y. Buyvolov, A.

Buyvolova, Y. Bykov, E. Chakhireva, O. Chashchina, N. Cherenkova, S. Chistjakov, S.

Chuhontseva, E. A. Davydov, V. Demchenko, E. Diadicheva, A. Dobrolyubov, L.

Dostoyevskaya, S. Drovnina, Z. Drozdova, A. Dubanaev, Y. Dubrovsky, S. Elsukov, L.

Epova, O. Ermakova, O. S. Ermakova, E. Ershkova, A. Esengeldenova, O. Evstigneev, I.

Fedchenko, V. Fedotova, T. Filatova, S. Gashev, A. Gavrilov, I. Gaydysh, D. Golovcov,

N. Goncharova, E. Gorbunova, T. Gordeeva, V. Grishchenko, L. Gromyko, V.

Hohryakov, A. Hritankov, E. Ignatenko, S. Igosheva, U. Ivanova, N. Ivanova, Y.

Kalinkin, E. Kaygorodova, F. Kazansky, D. Kiseleva, A. Knorre, L. Kolpashikov, E.

Korobov, H. Korolyova, N. Korotkikh, G. Kosenkov, S. Kossenko, E. Kotlugalyamova,

E. Kozlovsky, V. Kozsheechkin, A. Kozurak, I. Kozyr, A. Krasnopevtseva, S. Kruglikov,

O. Kuberskaya, A. Kudryavtsev, E. Kulebyakina, Y. Kulsha, M. Kupriyanova, M.

Kurbanbagamaev, A. Kutenkov, N. Kutenkova, N. Kuyantseva, A. Kuznetsov, E. Larin,

P. Lebedev, K. Litvinov, N. Luzhkova, A. Mahmudov, L. Makovkina, V. Mamontov, S.

Mayorova, I. Megalinskaja, A. Meydus, A. Minin, O. Mitrofanov, M. Motruk, A.

Myslenkov, N. Nasonova, N. Nemtseva, I. Nesterova, T. Nezdoliy, T. Niroda, T.

Novikova, D. Panicheva, A. Pavlov, K. Pavlova, S. Podolski, N. Polikarpova, T.

Polyanskaya, I. Pospelov, E. Pospelova, I. Prokhorov, I. Prokosheva, L. Puchnina, I.

Putrashyk, J. Raiskaya, Y. Rozhkov, O. Rozhkova, M. Rudenko, I. Rybnikova, S.

Rykova, M. Sahnevich, A. Samoylov, V. Sanko, I. Sapelnikova, S. Sazonov, Z.

Selyunina, K. Shalaeva, M. Shashkov, A. Shcherbakov, V. Shevchyk, S. Shubin, E.

Shujskaja, R. Sibgatullin, N. Sikkila, E. Sitnikova, A. Sivkov, N. Skok, S. Skorokhodova,

E. Smirnova, G. Sokolova, V. Sopin, Y. Spasovski, S. Stepanov, V. Stratiy, V.

Strekalovskaya, A. Sukhov, G. Suleymanova, L. Sultangareeva, V. Teleganova, V.

Teplov, V. Teplova, T. Tertitsa, V. Timoshkin, D. Tirski, A. Tolmachev, A. Tomilin, L.

Tselishcheva, M. Turgunov, Y. Tyukh, P. Van, V. Van, A. Vasin, A. Vasina, A. Vekliuk,

L. Vetchinnikova, V. Vinogradov, N. Volodchenkov, I. Voloshina, T. Xoliqov, E.

Yablonovska-Grishchenko, V. Yakovlev, M. Yakovleva, O. Yantser, Y. Yarema, A.

Zahvatov, V. Zakharov, N. Zelenetskiy, A. Zheltukhin, T. Zubina, J. Kurhinen, and O.

Ovaskainen. 2021. Phenological shifts of abiotic events, producers and consumers across

a continent. Nature Climate Change 11:241–248.

Rui, J., J. Li, S. Wang, J. An, W. Liu, Q. Lin, Y. Yang, Z. He, and X. Li. 2015. Responses of

Bacterial Communities to Simulated Climate Changes in Alpine Meadow Soil of the

Qinghai-Tibet Plateau. Applied and Environmental Microbiology 81:6070–6077.

Russell, B. C., A. Torralba, K. P. Murphy, and W. T. Freeman. 2008. LabelMe: A Database and

Web-Based Tool for Image Annotation. International Journal of Computer Vision

77:157–173.

Safaldin, M., N. Zaghden, and M. Mejdoub. 2023. Moving Object Detection Based on Enhanced

Yolo-V2 Model. Pages 1–8 2023 5th International Congress on Human-Computer

Interaction, Optimization and Robotic Applications (HORA). IEEE.

Santos Pereira, L. F., S. Barbon, N. A. Valous, and D. F. Barbin. 2018. Predicting the ripening of

papaya fruit with digital imaging and random forests. Computers and Electronics in

Agriculture 145:76–82.

Sargent, R. D., and D. D. Ackerly. 2008. Plant–pollinator interactions and the assembly of plant

communities. Trends in Ecology & Evolution 23:123–130.

Shannon, C. E. 1948. A mathematical theory of communication. The Bell system technical

journal 27:379–423.

Sponsler, D. B., F. Requier, K. Kallnik, A. Classen, F. Maihoff, J. Sieger, and I. Steffan-

Dewenter. 2022. Contrasting patterns of richness, abundance, and turnover in mountain

bumble bees and their floral hosts. Ecology 103:e3712.

Steinbauer, M. J., J.-A. Grytnes, G. Jurasinski, A. Kulonen, J. Lenoir, H. Pauli, C. Rixen, M.

Winkler, M. Bardy-Durchhalter, E. Barni, A. D. Bjorkman, F. T. Breiner, S. Burg, P.

Czortek, M. A. Dawes, A. Delimat, S. Dullinger, B. Erschbamer, V. A. Felde, O.

Fernández-Arberas, K. F. Fossheim, D. Gómez-García, D. Georges, E. T. Grindrud, S.

Haider, S. V. Haugum, H. Henriksen, M. J. Herreros, B. Jaroszewicz, F. Jaroszynska, R.

Kanka, J. Kapfer, K. Klanderud, I. Kühn, A. Lamprecht, M. Matteodo, U. M. di Cella, S.

Normand, A. Odland, S. L. Olsen, S. Palacio, M. Petey, V. Piscová, B. Sedlakova, K.

Steinbauer, V. Stöckli, J.-C. Svenning, G. Teppa, J.-P. Theurillat, P. Vittoz, S. J. Woodin,

N. E. Zimmermann, and S. Wipf. 2018. Accelerated increase in plant species richness on

mountain summits is linked to warming. Nature 556:231–234.

Stöcklin, J., P. Kuss, and A. R. Pluess. 2009. Genetic diversity, phenotypic variation and local

adaptation in the alpine landscape: case studies with alpine plant species. Botanica

Helvetica 119:125–133.

Suh, S., Y. Park, K. Ko, S. Yang, J. Ahn, J.-K. Shin, and S. Kim. 2021. Weighted Mask R-CNN

for Improving Adjacent Boundary Segmentation. Journal of Sensors 2021:1–8.

Sumit, S. S., J. Watada, A. Roy, and D. Rambli. 2020. In object detection deep learning methods,

YOLO shows supremum to Mask R-CNN. Journal of Physics: Conference Series

1529:042086.

Taheri, S., B. Naimi, C. Rahbek, and M. B. Araújo. 2021. Improvements in reports of species

redistribution under climate change are required. Science Advances 7:eabe1110.

Theobald, E. J., I. Breckheimer, and J. HilleRisLambers. 2017. Climate drives phenological

reassembly of a mountain wildflower meadow community. Ecology 98:2799–2812.

Tian, Y., G. Yang, Z. Wang, E. Li, and Z. Liang. 2020. Instance segmentation of apple flowers

using the improved mask R–CNN model. Biosystems Engineering 193:264–278.

Tu, S., Y. Xue, C. Zheng, Y. Qi, H. Wan, and L. Mao. 2018. Detection of passion fruits and

maturity classification using Red-Green-Blue Depth images. Biosystems Engineering

175:156–167.

Vorkauf, M., A. Kahmen, C. Körner, and E. Hiltbrunner. 2021. Flowering phenology in alpine

grassland strongly responds to shifts in snowmelt but weakly to summer drought. Alpine

Botany 131:73–88.

Wan, P., A. Toudeshki, H. Tan, and R. Ehsani. 2018. A methodology for fresh tomato maturity

detection using computer vision. Computers and Electronics in Agriculture 146:43–50.

Wang, X. (Annie), J. Tang, and M. Whitty. 2021. DeepPhenology: Estimation of apple flower

phenology distributions based on deep learning. Computers and Electronics in

Agriculture 185:106123.

Wieme, J., K. Mollazade, I. Malounas, M. Zude-Sasse, M. Zhao, A. Gowen, D. Argyropoulos, S.

Fountas, and J. Van Beek. 2022. Application of hyperspectral imaging systems and

artificial intelligence for quality assessment of fruit, vegetables and mushrooms: A

review. Biosystems Engineering 222:156–176.

Wilson, A., K. Bacher, I. Breckheimer, J. Lundquist, R. Rochefort, E. Theobald, L. Whiteaker,

and J. HilleRisLambers. 2017. Monitoring wildflower phenology using traditional

science, citizen science, and crowd sourcing. Park Science 33:17–26.

Wu, Y., A. Kirillov, F. Massa, W.-Y. Lo, and R. Girshick. 2019. Detectron2.

Xu, M., S. Yoon, A. Fuentes, and D. S. Park. 2023. A Comprehensive Survey of Image

Augmentation Techniques for Deep Learning. Pattern Recognition 137:109347.

Yongheng, G. 2007. Grazing intensity impacts on carbon sequestration in an alpine meadow on

the eastern Tibet.

Yu, Y., K. Zhang, L. Yang, and D. Zhang. 2019. Fruit detection for strawberry harvesting robot

in non-structural environment based on Mask-RCNN. Computers and Electronics in

Agriculture 163:104846.

Zhou, H., W. Ouyang, J. Cheng, X. Wang, and H. Li. 2019. Deep Continuous Conditional

Random Fields With Asymmetric Inter-Object Constraints for Online Multi-Object

Tracking. IEEE Transactions on Circuits and Systems for Video Technology 29:1011–

1022.

6. APPENDIX A

Mask R-CNN

We used Mask R-CNN (Mask Region based CNN), a deep learning algorithm that

generates bounding boxes and segmentation for each instance of an object (here a flower)

in an image (Fig. A.1). It is classified as a two-pass network; where the first part is

dedicated to feature learning, while the second part is for detection and segmentation. In

the end, the network identifies individual objects in an image and localizes each of them

with a bounding box (Machefer et al. 2020) along with predicting pixel level segmentation

masks. Instance segmentation in Mask R-CNN comprises of object detection and

semantic segmentation. Mask R-CNN is an improvement upon Faster R-CNN (Ren et al.

2017) which used region proposal and classification networks to detect region of interest

(RoI) and bounding boxes. Mask R-CNN supports various backbones as feature pyramid

network (FPN) in order to produce feature maps of the input image. Region proposal

network (RPN) is then applied atop feature maps that returns object proposals with

corresponding objectness scores. Mask R-CNN identifies the anchor boxes based on the

IoU, which is a critical step in the first part of the network (RPN). Objectness tells how

good the model did to predict an object regardless of class. A RoIAlign layer is then

applied to filter the proposals; the predecessor (Faster R-CNN (Ren et al., 2017)) used RoI

pooling which made it slower. The key feature is the use of RoI Align instead of RoiPool

(quantization free layer). The proposals are then passed to a fully connected layer that

returns class and bounding boxes for the objects. Simultaneously, the mask branch that

is a fully convolutional network (FCN) works on each RoI, and outputs a segmented mask

at a pixel level. Mask R-CNN uses class loss, box loss and mask loss for scoring the model.

In our study Mask R-CNN was configured with ResNext 101 backbone.

Figure A.1: High level architecture of the Mask R-CNN algorithm.

YOLO

YOLO stands for "You Only Look Once" that uses regression to achieve object detection and localization (Redmon et al. 2016). YOLO is a one stage method that works by regressing the bounding boxes and class probabilities and is known for its accuracy and high speed (Fig. A.2). YOLO has constantly evolved by releasing significant iterations with changes to its network structure resulting in better speed, consistent classification accuracy and recognizing smaller targets (Redmon et al. 2016, Redmon and Farhadi 2017, 2018). The YOLO method divides each input into regions or grids (S by S where S =8 for e.g.). For each grid it predicts B bounding boxes with confidence scores along with class conditional probabilities C. Confidence score translates to accuracy of the predicted object(s) in a grid. YOLO employs non-maximum suppression (NMS) similarly to Mask R-CNN if there are multiple candidate bounding boxes for the same target object. The loss function has three parts: objectness, box and classification. Objectness like in Mask R-CNN is a measure of how well the model did to predict an object irrespective of the class, Box tells how close the predicted box aligns with the actual box regardless of class and Classification loss tells how well each grid is classified with respect to its class probabilities. The YOLO algorithm penalizes the model's inability to draw the bounding box correctly and where it missed estimating it completely. YOLO v5 was the last publicly available version by the original authors, and the recent release (unofficially known as Yolo v5) is now maintained by Ultralytics (GitHub handle - https://github.com/ultralytics/yolov5)

Figure A.2: High level architecture of the YOLO algorithm.

Retina-Net

RetinaNet is a three-part learner where the convolution layer learns from the architecture

of the images (Lin et al. 2017, Li and Ren 2019). The architecture uses a feature pyramid

network (FPN) backbone on the top of ResNet architecture to generate a rich, multi-scale

convolutional feature pyramid (Fig. A.3). Two subnetworks are attached to it next –

classification subnet and regression subnet. The first subnetwork is for classifying anchor

boxes, while the second subnetwork is for regressing anchor boxes to the ground-truth

boxes. The complex head and body of the network involves convolutions which reduces

the image to a deep low-resolution image. The network is tuned by using the regression

loss and the classification loss. Both metrics look at overall objectness and classification

ability of the model. Regression loss is calculated by using smooth L1 whereas

classification is evaluated by Focal loss.

Figure A.3: High level architecture of the RetinaNet algorithm.
910

7. APPENDIX B

Predictions by RetinaNet and YOLO

Figure B.1: Predictions from the RetinaNet method showing the fluency of the model; the
percent next to the predicted class represents model confidence. (A and B) Meadow in
peak flowering with predictions (B) corresponding original tile. (C and D) Another
meadow in peak flowering with predictions (D) corresponding to the original tile.

Figure B.2: Predictions from the YOLO method showing the fluency of the model; the percent next to the predicted class represents model confidence. (A and B) Meadow in peak flowering with predictions (B) corresponding original tile. (C and D) Another meadow in peak flowering with predictions (D) corresponding to the original tile.

8. APPENDIX C

Hyperparameters for the three methods

927
928 Table C.1: Results of anchor box variation with RetinaNet. Used batch size of 10 and ran
929 it for 250 epochs (iterations).

Anchor Box Size	AP - Bounding Box
[4, 8, 16, 32, 64]	6.99
[8, 16, 32, 64, 128]	9.05
[16, 32, 64, 128, 256]	13.41
[32, 64, 128, 256,512]	8.22

930
931

932 **9.** APPENDIX D

933 Correlation between richness as inferred by Mask R-CNN, YOLO and RetinaNet for
934 August 2020 dataset.
935

937   Figure D.1: Correlation between richness (number of unique flowers) as calculated by
938   Mask R-CNN, YOLO and RetinaNet for all the collections from August 2020. Significant
939   association is found between YOLO and Mask R-CNN