Transcriptomes Often Resemble Biological Aging Upon Perturbation

Thomas Stoeger[1,2,3]

[1] Feinberg School of Medicine in the Division of Pulmonary and Critical Care Medicine, Northwestern University, Chicago, IL, USA

[2] The Potocsnak Longevity Institute, Northwestern University, Chicago, IL, USA

[3] Simpson Querrey Lung Institute for Translational Science, Chicago, IL, USA

Correspondence: thomas.stoeger@northwestern.edu

Summary

Aging is deeply interwoven with cellular and physiological biology, but the extent is not known.
Here, I demonstrate that the Gene Length-dependent Transcription Decline (GLTD) which marks
age-dependent transcriptional change is not specific to aging. Instead, it follows perturbations in
5,231 of 11,959 public gene expression profiles. My analysis suggests the hypothesis that during
the passage of time, exposure to myriad perturbations change transcriptomes in a way that
would be described as "aging".

Keywords: Aging, transcriptomics, damage, convergence, transcription, gene length-dependent
transcription decline

14
15 **Main Text**
16
Conditions that promote health tend to delay phenotypic manifestations of aging, whereas the
inverse tends to hold for conditions that worsen health[1]. However, it is not known to what
extent perturbations of cellular and organismal physiology yield phenotypes resembling those
encountered during aging[2].

To address this question, I drew from the recent insight that transcriptomic changes in humans
and mice with age can be approximated by a negative correlation between gene length and
transcriptomic change[3-7], termed Gene Length-dependent Transcription Decline (GLTD) [*in press,*
Trends in Genetics[8]]. Unlike most molecular and physiological markers of age[9], GLTD can be
retrospectively measured in a wide set of biomedical studies, as transcriptomics are frequently
reported in a wide vary of biomedical settings. I begin my investigation with resources that
aggregate studies.

EBI Gene Expression Atlas (EBI-GXA) gathers and processes high-quality transcriptomic studies[10],
which reflect areas of interest to individual investigators (10 of 10 randomly sampled studies,
Extended Data Table 1). I considered those 7,108 (4,077 human and 3,031 mouse) differential
gene expression profiles that report the relative fold-changes of transcripts of perturbed
samples relative to control samples for at least 10,000 genes. Because GLTD in aging originates
in a loss of productive transcription over the gene body[7], I quantified gene length through the
median length of primary transcripts, omitting non-coding regulatory regions. As before[5], I
quantified length-correlated change through the Spearman correlation between gene length
and relative fold-changes of transcripts of protein-coding genes using Kendall and Stuart's
method for P values[11].

Though EBI-GXA does not directly report age, it reproduces insights into a model of accelerated
aging[12] in which GLTD has been firmly established[3,7]. Specifically, mice with mutations in the
DNA excision repair protein *DNA excision repair protein ERCC-1* (ERCC1) gene show a significant
negative correlation (Figure 1A, Spearman correlation: -0.14, P value = 2.4E-68). Further, the 7[th]
most negative correlation in EBI-GXA derives from UV-irradiated mice with mutations in the
related *DNA repair protein complementing XP-A cells* (XPA) gene and ERCC6 genes (Figure 1B;
Spearman correlation: -0.35, P value of "~ 0" as not precisely computable). Complete results are
provided as Extended Data Table 2.

In 44% of all profiles, I found that perturbations often resemble biological aging, as they often
yield a negative correlation (=GLTD) (Figure 1C, at a significance cutoff of P value = 0.01). These
perturbations were not confined to any single domain of biology (Figure 1C, Extended Data
Table 2). Importantly, perturbations preferentially caused length-correlated changes in the
direction that is dominant in aging[3-7], as more profiles showed a negative correlation (=GLTD)
than positive correlation (Figure 1C, Extended Data Figure 2, e.g.: Fisher's Exact P value of 1.76E-
5 when for length correlations considering a P value threshold of 0.01).

Similarly, the median and mean effect size of length-correlated change both were negative
(P~"0" via bootstrapping, and P=2.06E-11 via T-test, respectively). While effect sizes were small
(-0.0077 and -0.0074, respectively), the negative effect sizes include all perturbations in EBI-GXA
independent of the significance of length-correlated change, consistent with the finding that
age-dependent molecular change often[13,14] (but not always[14,15]) appears to be a gradual process
in which individual steps are not readily discernable from each other.

As in aging[5], GLTD following perturbations was largely independent of transcriptional regulation
of known age-associated pathways, as 40% of profiles still displayed GLTD when excluding 7,691
of 19,163 human and 8,212 of 21,167 mouse protein-coding genes that have been annotated by
the Gene Ontology Consortium[16] to participate in any of 27 processes regulated by age[17], some
as broad as "metabolism", "inflammation", "synaptic transmission", and "stress" (Extended Data
Figure 2).

While length-correlated changes in gene expression profiles have been suspected to have
technical origins in RNA-sequencing (RNA-seq)[18], negative correlations (=GLTD) are also more
numerous than positive correlations in human and mouse microarray data (Extended Data
Figure 3). While there is no discernable preference for positive or negative correlations in RNA-
seq data of humans or mice (Extended Data Figure 4), this result may be attributable to lower
statistical power (n=486 and n=371 versus n=2,788 and n=1,988 profiles) or increased technical

variability. Length-correlated changes in RNA-seq studies have been suspected to possibly
originate from length-dependent technical artifacts[18]. However, the preferential directionality of
the correlations favors biological origins, and this manuscript will subsequently confirm this
directionally through a variety of experimental methods and datasets, and observations on
nascent transcripts.

Another resource, GPSAdb[19], that exclusively contains RNA-seq data following loss-of-function
perturbations of individual genes, confirms that GLTD is common after perturbations. GPSAdb
aggregates a total of 2,713 profiles mapping to unambiguous gene identifiers, corresponding to
1,249 genes. Among these profiles, 45% show GLTD at a P value of 0.01 (Figure 1E, Extended
Data Table 3). Again, there are more negative than positive correlations (Figure 1E and Extended
Data Figure 5, e.g.: Fisher's Exact P value of 0.004 when for length correlations considering a P
value threshold of 0.01), and the median and mean effect size of length-correlated change are
negative (Figure 1F, bootstrapped P value 0.0002; Figure 1F, P value of 1.17E-6, T-test).

Closer inspection of GPSAdb hints toward further complexity. Loss-of-function by RNA
interferences (n=1,844) or genetic knockout (n=487) yielded median length correlations that are
negative, but CRISPR interferences (CRISPRi) (n=73) yielded median length correlations that are
positive (+0.09, bootstrapped P value of ~0 as not precisely computable). While the lower
number of perturbations with CRISPRi may have increased the chance for uncontrolled
covariates, the reasons the observed increase are presently unexplained.

Avoiding potential confounding factors of data aggregation and more comprehensively
inspecting the loss-of-function of genes, I next reanalyze one recent perturb-seq study[20] that
combined perturbations in a single experiment rather than after parallel experiments. The
authors applied CRISPRi constructs against all expressed genes in K562 cells, identified 1,958
genes to cause a strong transcriptional phenotype, and provide bioinformatically processed data
for those perturbations.

42% of profiles showed GLTD at a P value of 0.01 (Figure 2A, Extended Data Table 4). Opposing
the observations of CRISPRi in GPSAdb, more perturbations yielded a negative correlation
(=GLTD) rather than positive correlation (Figure 2B and Extended Data Figure 7, e.g.: Fisher's
Exact P value of 6.3E-38 when for length correlations considering a P value threshold of 0.01).
The median and mean effect sizes of length-correlated change again were negative (Figure 2B,
bootstrapped P value of ~0 as not precisely computable; and P value of 8.31E-79, T-test,
respectively).

Superimposing length-correlated change onto a map created by the authors of this datasets to
depict global dependencies of cellular transcriptomes on the function of individual genes
identified length-correlated change as a latent factor aligning with the first two dimensions of
the map (Spearman correlations of -0.40 and +0.30; P values of 1.2E-77 and 2.0E-42,
respectively). This alignment does not reflect the length of the genes whose function is lost
(Extended Data Figure 8A), is not affected by excluding differentially expressed genes annotated
the for any of 27 processes associated with aging[17] (Extended Data Figure 8B), and holds when
independently mapping genes through Principal Components (Extended Data Figure 8C).
Resembling observations in aging in which known transcription factors are less informative than
gene length[5], length-correlated change better aligns with the map than the maximal alignment
observed for any of 1,268 transcription factors (Extended Data Table 5 with maximal effect sizes

114 of correlation of +0.11 and +0.13, respectively). Gene Ontology Enrichment analysis confirmed
115 the preceding annotation that accompanied the map[20] (Figure 2C), as GLTD is most enriched
116 among genes that contribute to transcription or translation (Extended Data Table 6).

117 To provide initial mechanistic clues, I turn toward a recent parallel perturbation experiment[21]
118 that succeeded in biochemically tracing nascent transcripts to infer synthesis and degradation
119 rates in HEK293 cells upon CRISPRi of 180 genes. I found changes in synthesis and degradation
120 rate to be predominantly negatively correlated with gene length (Figure 2D,E; Extended Data
121 Tables 7 and 8; Extended Data Figure 9). Curiously, effects on length-correlated transcript
122 synthesis appeared particularly pronounced after loss-of-function of genes involved in
123 translation (Extended Data Figure 10). Overall, loss-of-function of these 180 genes resembled
124 GLTD in aging by originating during the transcript synthesis[7]. It is not known if there is a length-
125 correlated compensation of transcript stability in aging or if modulation of proteostasis[22] affects
126 GLTD in aging.

127 My findings align well with wear-and-tear-based theories of biological aging, especially
128 imperfection-driven non-random damage[23] and DNA damage[3,24], but cannot exclude additional
129 explanations and covariates or extend beyond the transcriptome. The main contribution of my
130 present analysis is the demonstration that the primary[5] transcriptional phenotype encountered
131 during aging is a frequent mode of transcriptional change following perturbations. These
132 transcriptional data suggest the hypothesis that "aging" is not a specific biological process but
133 rather that during the passage of time, exposure to myriad perturbations accumulate in a
134 manner that would be described as "aging". Alternatively, the transcriptional phenotype of
135 older individuals may reflect a higher susceptibility of their transcriptomes toward change
136 following perturbations.

138
139 **Acknowledgments**
140
141 I thank Zihan Xu and Junyue Cao for sharing intermediate data behind their publication. I thank
142 Ben D. Singer and M. Jamal Jenkins for feedback on the manuscript.

144 **Author contributions**

145 T.S. conceptualized the study, curated the data, performed formal analysis, visualized the
146 findings, and wrote the manuscript.

148 **Declaration of interests**

149 The author declares no competing interests.

151
152

Materials and Methods

Length Correlations

The primary transcript lengths for humans and mice were obtained from BioMart
(http://useast.ensembl.org/biomart/martview), specifically from GRCh38.p13, and GRCm39,
respectively. For each gene, the median length across all primary transcripts was calculated, and
genes marked as protein-coding by NCBI's reference file
(https://ftp.ncbi.nlm.nih.gov/gene/DATA/GENE_INFO/All_Data.gene_info.gz) were selected. For
mice and humans, only genes with an unambiguous 1:1 mapping between Entrez and Ensembl
gene identifiers were retained. Length correlations were defined as the Spearman correlation
between the lengths of primary transcripts and fold-changes of transcripts and computed using
SciPy 1.2.1. P-values of Spearman correlations were obtain by Kendall and Stuart's method for P-
values of Spearman correlations [11].

Bootstrapping

Bootstrapping was done with 100,000 iterations.

Reanalysis of EBI-GXA

Expression profiles were downloaded from
http://ftp.ebi.ac.uk/pub/databases/microarray/data/atlas/experiments/ . To account for some
experiments reporting few genes, experiments reporting less than 10,000 protein-coding genes
with unambiguous 1:1 mapping between Entrez and Ensembl were excluded from the analysis.

Reanalysis of GPSAdb

Expression profiles were downloaded from https://www.gpsadb.com using their bulk-download
option. Length correlations were computed over the transcripts of genes that were not directly
targeted by a given perturbation.

Exclusion of pathways associated with aging

Gene Ontology annotations from NCBI were used,
https://ftp.ncbi.nlm.nih.gov/gene/DATA/gene2go.gz. Genes were excluded if "go_term"
matched case-insensitively the following regular expression, which represents the list of age-
associated pathways compiled by Stegeman and Weake[17]:

dna repair|inflamm|calcium-mediated signaling|\btranslation|insulin|response to heat|heat
shock|cell cycle|protein catabolic process|\bliver|MAPK cascade|mitochondrial
matrix|potassium|ATPase|glucose|apoptosis|retinoic|matrix|oxidative stress|RNA
processing|cell differentiation|\bT cell activation|cellular matrix|metabolic process|axon
guidance|synaptic trans|stress

Reanalysis of Perturb-seq

The fold-changes and coordinates of Figure 2 of Replogle and Saunders et al.[20] were obtained
from the author-provided files at https://doi.org/10.25452/figshare.plus.21632564.v1. . Length
correlations were computed over the transcripts of genes that were not directly targeted by a
given perturbation. PCA analysis was preceded by an imputation of missing values according to
the median fold-change observed for a gene, and z-scoring across of a gene over all
perturbations.

Transcription factor analysis

Transcription factor binding sites were obtained from GTRD[25], version 21,
http://gtrd.biouml.org:8888/downloads/current/intervals/target_genes/, and the included
"genes promoter[-1000,+100]" used to map transcription factors toward genes.

Gene Ontology enrichment analysis

Gene Ontology annotations from NCBI were used,
https://ftp.ncbi.nlm.nih.gov/gene/DATA/gene2go.gz. Enrichment was done in Python via
Fisher's Exact test, implemented in scipy.stats.

Reanalysis of Xu et al.
204
205 Per-gene resolved intermediate data behind Xu et al.[21] was requested from, and kindly shared by,
206 the authors, Zihan Xu and Junyue Cao. Length correlations were computed over the transcripts
207 of genes that were not directly targeted by a given perturbation.

208
209 **Data and code availability**

210 Code will be made public upon acceptance via GitHub, and is meanwhile attached for reviewers
211 as a zip file.

212
213
214
215

216 **Figures**
217

218
Figure 1. Gene Length-dependent Transcription Decline (GLTD) is frequent in public
transcriptomic data. (**A, B**) Transcriptomic fold changes of genes (grey dots). **(C)** Fraction of gene
expression profiles with length-correlated change in EBI-GXA[10] at different *P* value thresholds,
determined by Kendall and Stuart's method[11]. Dashed vertical line indicates *P* value of 0.01. **(D)**
Histogram of all observed length-correlated changes in EBI-GXA, including non-significant ones.
P value of median inferred from bootstrapping and *P* value of mean from two-sided one group T-
test. n.c. is non computable as close to 0. (**E, F**) as (A,B) but for GPSAdb[19].
226

227
Figure 2. Gene Length-dependent Transcription Decline (GLTD) is frequent following the loss of
genes. **(A)** Fraction of gene expression profiles with length-correlated change in Replogle et al.[20]
at different *P* value thresholds, determined by Kendall and Stuart's method[11]. Dashed vertical line
indicates *P* value of 0.01. **(B)** Histogram of all observed length-correlated changes in in Replogle
et al., including non-significant ones. *P* value of median inferred from bootstrapping and *P* value
of mean from two-sided one group T-test. n.c. is non computable as close to 0. **(C)** Length-
correlated change superimposed on map by Replogle et al., a Uniform Manifold Approximation
and Projection (UMAP), of genes according to the effect of their loss-of-function toward the
transcriptome. **(D)** As (B) but for synthesis rates contained in Xu et al.[21]. **(E)** As (B) but for
synthesis rates contained in Xu et al.[21].

239
240
241 **Extended Data Figure 1.** *P* value of two-sided Fisher's Exact test for hypothesis that at a given
242 significance threshold of length-correlated change (*P* value of length correlation) within EBI-GXA
243 there was the same amount of negative and positive correlations.

244
245

246
247
Extended Data Figure 2. Gene Length-dependent Transcription Decline (GLTD) persists in EBI-
GXA when excluding genes annotated for aging-related pathways. **(A)** Fraction of gene
expression profiles with length-correlated change in EBI-GXA[10] at different *P* value thresholds,
determined by Kendall and Stuart's method[11]. Dashed vertical line indicates *P* value of 0.01. **(B)**
Histogram of all observed length-correlated changes in EBI-GXA, including non-significant ones.
P value of median inferred from bootstrapping and *P* value of mean from two-sided one group T-
test. n.c. is non computable as close to 0. **(C)** *P* value of two-sided Fisher's Exact test for
hypothesis that at a given significance threshold of length-correlated change (*P* value of length
correlation) within EBI-GXA there was the same amount of negative and positive correlations.

258
259
260
261
Extended Data Figure 3. Gene Length-dependent Transcription Decline (GLTD) persists in EBI-
GXA when only considering experiments annotated to have been performed by microarrays. **(A)**
Fraction of gene expression profiles with length-correlated change for human studies at different
P value thresholds, determined by Kendall and Stuart's method[11]. Dashed vertical line indicates *P*
value of 0.01. **(B)** As (A) but for mouse studies. **(C)** Histogram of all observed length-correlated
changes in human studies, including non-significant ones. *P* value of median inferred from
bootstrapping and *P* value of mean from two-sided one group T-test. n.c. is non computable as
close to 0. **(D)** As (C) but for mouse studies. **(E)** *P* value of two-sided Fisher's Exact test for
hypothesis that at a given significance threshold of length-correlated change (*P* value of length
correlation) within human studies there was the same amount of negative and positive
correlations. **(F)** As (E) but for mouse studies.

273
274
275
276

277
278
279
Extended Data Figure 4. Gene Length-dependent Transcription Decline (GLTD) persists in EBI-
GXA when only considering experiments annotated to have been performed by RNA-sequencing
(RNA-Seq). **(A)** Fraction of gene expression profiles with length-correlated change for human
studies at different P value thresholds, determined by Kendall and Stuart's method[11]. Dashed
vertical line indicates P value of 0.01. **(B)** As (A) but for mouse studies. **(C)** Histogram of all
observed length-correlated changes in human studies, including non-significant ones. P value of
median inferred from bootstrapping and P value of mean from two-sided one group T-test. n.c. is
non computable as close to 0. **(D)** As (C) but for mouse studies. **(E)** P value of two-sided Fisher's
Exact test for hypothesis that at a given significance threshold of length-correlated change (P
value of length correlation) within human studies there was the same amount of negative and
positive correlations. **(F)** As (E) but for mouse studies.
291
292

293
294
295 **Extended Data Figure 5.** *P* value of two-sided Fisher's Exact test for hypothesis that at a given
296 significance threshold of length-correlated change (*P* value of length correlation) within GPSAdb
297 there was the same amount of negative and positive correlations.
298

299
300 **Extended Data Figure 6.** Length correlations within GPSAdb according to method used for loss-
301 of-function. Notches indicate bootstrapped 95% confidence interval of median. Box contains 25-
302 75% range of observe length correlations. n indicates number of genes. n.c. indicates that not
303 precisely computable.
304
305

306
307 **Extended Data Figure 7.** *P* value of two-sided Fisher's Exact test for hypothesis that at a given
308 significance threshold of length-correlated change (*P* value of length correlation) within Replogle
309 et al.[20] there was the same amount of negative and positive correlations.

Alignment with UMAP1:
Spearman correlation +0.01
P value: 0.52

Alignment with UMAP2:
Spearman correlation -0.13
P value: 2.9E-8

Alignment with UMAP1:
Spearman correlation -0.40
P value: 1.21E-77

Alignment with UMAP2:
Spearman correlation +0.30
P value: 1.97E-42

Alignment with PCA1:
Spearman correlation -0.40
P value: 5.61E-78

Alignment with PCA2:
Spearman correlation -0.40
P value: 5.4E-76

Extended Data Figure 8. The alignment of length-correlated change with the map of loss-of-function effects by Replogle et al.[20] is robust toward alternative analysis. **(A)** Superimposition of gene-length of targeted genes (dots). **(B)** Superimposition of length-correlated change computed without genes annotated for pathways associated with aging. **(C)** Alternative map of loss-of-function using Principal Component Analysis (PCA).

320
321
Extended Data Figure 9. Loss-of-function causes length-correlated change of transcript
synthesis and degradation when re-analyzing Xu et al.[21]. **(A)** Histogram of all observed length-
correlated changes in transcript synthesis Xu et al., including non-significant ones. *P* value of
median inferred from bootstrapping and *P* value of mean from two-sided one group T-test. n.c. is
non computable as close to 0. **(B)** As (A) but for degradation rates. **(C)** *P* value of two-sided
Fisher's Exact test for hypothesis that at a given significance threshold of length-correlated
change (*P* value of length correlation) within transcriptome synthesis data of Xu et al. there was
the same amount of negative and positive correlations. **(D)** Same as (C) but for degradation
rates.

331
332
333 **Extended Data Figure 10.** Length correlations within Xu et al.[21] according to process annotations
334 by them. Notches indicate bootstrapped 95% confidence interval of median. Box contains 25-75%
335 range of observe length correlations. n indicates number of genes.
336

Extended Data Table 1. Manual inspection of 10 randomly selected gene expression profiles in EBI-GXA.

Extended Data Table 2. Length-correlated changes in EBI-GXA. Rho is Spearman correlation, pval the *P* value for correlation obtained by Kendall and Stuart's method[11]. n is number of protein-coding genes.

Extended Data Table 3. Length-correlated changes in GPSAdb. Rho is Spearman correlation, pval the *P* value for correlation obtained by Kendall and Stuart's method[11]. n is number of protein-coding genes.

Extended Data Table 4. Length-correlated changes in Replogle et al.[20]. Rho is Spearman correlation, pval the *P* value for correlation obtained by Kendall and Stuart's method[11]. umap_x is first UMAP dimension, and umap_y is the second UMAP dimension.

Extended Data Table 5. Associations of transcription factors with first and second UMAP dimension (axis). Correlations obtained via point-biserial correlations and Spearman correlations.

Extended Data Table 6. Gene Ontology analysis for genes with GLTD in Replogle et al.[20].

Extended Data Table 7. Length-correlated changes for synthesis in Xu et al.[21]. Rho is Spearman correlation, pval the *P* value for correlation obtained by Kendall and Stuart's method[11]. gene_class is annotation of process provided by authors.

Extended Data Table 8. Length-correlated changes for degradation in Xu et al.[21]. Rho is Spearman correlation, pval the *P* value for correlation obtained by Kendall and Stuart's method[11]. gene_class is annotation of process provided by authors.

References

1   Kennedy, B. K. *et al.* Geroscience: linking aging to chronic disease. *Cell* **159**, 709-713 (2014). https://doi.org:10.1016/j.cell.2014.10.039

2   Freund, A. Untangling Aging Using Dynamic, Organism-Level Phenotypic Networks. *Cell Syst* **8**, 172-181 (2019). https://doi.org:10.1016/j.cels.2019.02.005

3   Vermeij, W. P. *et al.* Restricted diet delays accelerated ageing and genomic stress in DNA-repair-deficient mice. *Nature* **537**, 427-431 (2016). https://doi.org:10.1038/nature19329

4   Hall, H. *et al.* Transcriptome profiling of aging Drosophila photoreceptors reveals gene expression trends that correlate with visual senescence. *BMC Genomics* **18**, 894 (2017). https://doi.org:10.1186/s12864-017-4304-3

5   Stoeger, T. *et al.* Aging is associated with a systemic length-associated transcriptome imbalance. *Nature Aging* **2**, 1191-1206 (2022). https://doi.org:10.1038/s43587-022-00317-6

6   Ibañez-Solé, O., Barrio, I. & Izeta, A. Age or lifestyle-induced accumulation of
genotoxicity is associated with a length-dependent decrease in gene expression.
iScience, 106368 (2023). https://doi.org:https://doi.org/10.1016/j.isci.2023.106368
7   Gyenis, A. *et al.* Genome-wide RNA polymerase stalling shapes the transcriptome during
aging. *Nat Genet* **55**, 268-279 (2023). https://doi.org:10.1038/s41588-022-01279-6
8   Soheili-Nezhad, S., Ibáñez-Solé, O., Izeta, A., Hoeijmakers, J. H. & Stoeger, T. Time is
ticking faster for long genes in aging. *Trends in Genetics* (in press).
9   Hartmann, A. *et al.* Ranking Biomarkers of Aging by Citation Profiling and Effort Scoring.
Front Genet **12**, 686320 (2021). https://doi.org:10.3389/fgene.2021.686320
10   Papatheodorou, I. *et al.* Expression Atlas: gene and protein expression across multiple
studies and organisms. *Nucleic Acids Res* **46**, D246-D251 (2018).
https://doi.org:10.1093/nar/gkx1158
11   Kendall, M. G. & Stuart, A. Inference and Relationship. *The Advanced Theory of Statistics*
2 (1973).
12   Niedernhofer, L. J. *et al.* A new progeroid syndrome reveals that genotoxic stress
suppresses the somatotroph axis. *Nature* **444**, 1038-1043 (2006).
https://doi.org:10.1038/nature05456
13   Cellerino, A. & Ori, A. What have we learned on aging from omics studies? *Semin Cell*
Dev Biol **70**, 177-189 (2017). https://doi.org:10.1016/j.semcdb.2017.06.012
14   Schaum, N. *et al.* Ageing hallmarks exhibit organ-specific temporal signatures. *Nature*
583, 596-602 (2020). https://doi.org:10.1038/s41586-020-2499-y
15   Poganik, J. R. *et al.* Biological age is increased by stress and restored upon recovery. *Cell*
Metabolism **35**, 807-+ (2023). https://doi.org:10.1016/j.cmet.2023.03.015
16   Gene Ontology, C. Gene Ontology Consortium: going forward. *Nucleic Acids Res* **43**,
D1049-1056 (2015). https://doi.org:10.1093/nar/gku1179
17   Stegeman, R. & Weake, V. M. Transcriptional Signatures of Aging. *J Mol Biol* **429**, 2427-
2437 (2017). https://doi.org:10.1016/j.jmb.2017.06.019
18   Mandelboum, S., Manber, Z., Elroy-Stein, O. & Elkon, R. Recurrent functional
misinterpretation of RNA-seq data caused by sample-specific gene length bias. *PLoS Biol*
17, e3000481 (2019). https://doi.org:10.1371/journal.pbio.3000481
19   Guo, S. *et al.* GPSAdb: a comprehensive web resource for interactive exploration of
genetic perturbation RNA-seq datasets. *Nucleic Acids Res* **51**, D964-D968 (2023).
https://doi.org:10.1093/nar/gkac1066
20   Replogle, J. M. *et al.* Mapping information-rich genotype-phenotype landscapes with
genome-scale Perturb-seq. *Cell* **185**, 2559-2575 e2528 (2022).
https://doi.org:10.1016/j.cell.2022.05.013
21   Xu, Z., Sziraki, A., Lee, J., Zhou, W. & Cao, J. Dissecting key regulators of transcriptome
kinetics through scalable single-cell RNA profiling of pooled CRISPR screens. *Nature*
Biotechnology (2023). https://doi.org:10.1038/s41587-023-01948-9
22   Labbadia, J. & Morimoto, R. I. The biology of proteostasis in aging and disease. *Annu Rev*
Biochem **84**, 435-464 (2015). https://doi.org:10.1146/annurev-biochem-060614-033955
23   Gladyshev, V. N. The origin of aging: imperfectness-driven non-random damage defines
the aging process and control of lifespan. *Trends Genet* **29**, 506-512 (2013).
https://doi.org:10.1016/j.tig.2013.05.004

24   Schumacher, B., Pothof, J., Vijg, J. & Hoeijmakers, J. H. J. The central role of DNA damage
in the ageing process. *Nature* **592**, 695-703 (2021). https://doi.org:10.1038/s41586-021-
03307-7
25   Yevshin, I., Sharipov, R., Kolmykov, S., Kondrakhin, Y. & Kolpakov, F. GTRD: a database
on gene transcription regulation-2019 update. *Nucleic Acids Res.* **47**, D100-D105 (2019).
https://doi.org:10.1093/nar/gky1128