Changes in large-scale neural networks under stress are linked to affective reactivity to stress in real life

Authors:

Rayyan Tutunji[1*], Martin Krentz[1], Nikos Kogias[1], Lycia de Voogd[1,2], Florian Krause[1], Eliana Vassena[1,2], Erno J. Hermans[1]

Affiliations:

1. Donders Institute for Brain, Cognition and Behaviour. Radboud University Medical Center. Nijmegen, The Netherlands.
2. Behavioural Science Institute. Radboud University. Nijmegen, The Netherlands.

*Corresponding Author:

Rayyan Tutunji, rayyan.tutunji@donders.ru.nl

Abstract

Controlled laboratory stress induction procedures are very effective in inducing physiological and subjective stress. However, whether such stress responses are representative for stress reactivity in real life is not clear. Using a combined within-subject functional MRI laboratory stress and ecological momentary assessment stress paradigm, we investigated dynamic shifts in large-scale neural network configurations under stress and how these relate to affective reactivity to stress in real life. Laboratory stress induction resulted in significantly increased cortisol levels, and shifts in task-driven neural activity. Namely, increased salience network (SN) activation in an oddball task and decreased default mode network activity in a memory retrieval task. Crucially, individuals showing increased SN reactivity specifically in the early phase of the acute stress response also expressed increased affective reactivity in real life. Our findings provide (correlational) evidence that real-life affective stress reactivity is driven primarily by vigilant attentional reorientation mechanisms associated with SN.

Introduction

Acute stress triggers a cascade of time-dependent processes that result in dynamic shifts of large-scale brain network configurations[1,2]. These processes are driven by distinct actions of stress-sensitive hormones and neuromodulators in the acute phase and in the recovery phase of the stress response[3,4]. While these dynamics have been studied extensively in experimental models in both animals and humans, it is unknown how interindividual differences in stress-induced perturbations of large-scale networks are reflected in interindividual differences in reactivity to acute stressors in real life. This variability is critical for understanding the role of networks/stress responses in healthy adaptation to stress and psychopathology, especially given the involvement of alterations in large-scale networks in both stress and psychopathology[5–8].

Research has implicated three core large-scale brain networks - the salience (SN), executive control (ECN), and default mode networks (DMN) - in both reactivity to, and recovery from a stressor through the effects of stress-related hormones and neurotransmitters [1,6]. The early reactivity phase of the stress response is thought to be driven by actions of catecholamines and corticosteroids[4,9]. In this phase, activation of the locus coeruleus results in tonically elevated release of norepinephrine[10]. Increased LC activity and norepinephrine have also been associated with overall increased salience network activity[3,11,12]. This coincides with specific environmental demands during acute stress, with an increased need for threat vigilance[13]. At the same time, studies have also shown ECN and DMN suppression, associated with impaired working memory and memory retrieval under acute stress, functions supported by these two networks, respectively[14–16].

Interestingly, these processes may be reversed through slow, gene transcription-dependent corticosteroid effects in the later recovery stage of the stress response, starting 1-2 hours following stress[4]. Following the administration of hydrocortisone, within a time window in which genomic actions can be expected, a decrease in SN-related regions is seen[17,18]. Delayed effects of hydrocortisone in a time window that is consistent with genomic mechanisms have also been linked to upregulation of regions associated with the ECN, as well as with improved working memory performance[19]. Additionally, delayed effects of stress hormones have been shown to result in improved DMN linked memory retrieval[20]. Thus, growing evidence points to a reversal of overall changes in network balance under stress, with a decrease in SN, and increase in DMN and ECN related regions driven by later, genomic effects of corticosteroids. Arguably, this reversal in the late stage of the stress response serves an adaptive function by promoting higher-order cognitive functions required for a return to homeostasis and an

optimal preparation for future stressors. While many studies have shown the shifts within these large-scale networks over the course of either the early or late stress response, the temporal dynamics have thus far been only inferred . Within-subject, time dependent shifts in these large-scale networks over the full duration of the stress response – including both early nongenomic and late genomic effects – have not been thoroughly investigated.

While laboratory findings are important in establishing a mechanistic understanding of the neural stress response, a critical question is how individual dynamics of neural stress reactivity relate to healthy and functional responses to stress in real life. Real-life stress is often studied using approaches such as Ecological Momentary Assessments [EMA, also known as experience sampling methods or ESM[21]. These methods leverage repeated assessments in day-to-day lives of participants to derive measures of stress reactivity and sensitivity[22–24]. While such studies have created the opportunity to quantify stress reactivity in daily life, linking these observations to the lab has been a more difficult endeavor. Current attempts to derive daily life stress measures are limited by study designs that are unable to disentangle measures of exposure to stressors from measures of the consequences of said exposure. One measure of reactivity that can be adapted from resilience research to EMA studies is the residualization-based stressor reactivity measure[25]. In resilience research, this measure is derived by regressing the change in mental health onto a measure of exposure to life stressors. The residuals of this regression then indicate changes in mental health that are not explained by the amount of stress exposure[26–28]. Here, we adapted this measure for EMA research to quantify individual affective reactivity to acute stressors in real life at a time scale that is comparable to laboratory studies on acute stress[29]. As outcome measure, we used changes in positive affect, which has directly been linked to resilience [30].

In this study, we used a within-subject cross-sectional design (n=83) to investigate the effects of an established psychosocial laboratory stressor on the dynamics of large-scale networks under stress and task demand (see Figure 1). We investigated both the initial reactivity (i.e., early phase) and recovery phases (i.e., late phase) of this response in a novel functional MRI paradigm combining three tasks that have each been shown to preferentially recruit one of the three networks involved in stress (SN, ECN, DMN). We then compared temporal dynamics of stress to a control scan using a matched, non-stressful procedure to determine within-person changes in large-scale network balance across the phases of the stress response. Next, we investigated the relationship between this balance and real-life affective reactivity to stress using a residualization-based stressor reactivity score. We expected to see increased SN, and decreased ECN and DMN activity during the early phase of the stress response, with a reversal of

this balance in the later recovery phase (i.e., decreased SN, and increased ECN and DMN). This hypothesis is based on a previously published working model[1]. We also expected greater SN activation and ECN and DMN suppression to be associated with increased real-life stress reactivity, with the opposite effects in the recovery phase of the neural response to stress (i.e., smaller decreases or recovery in SN and increases in ECN and DMN).

Figure 1. Study Design. A) Stress and control weeks were counterbalanced for order, followed by stress and control MRI scans that were also counterbalanced between subjects. B) Typical scan day for a participant. C) Three tasks performed during MRI task blocks including a standard 2-Back, facial oddball, and associative retrieval task.

Methods

Participants

Eighty-three students in the first year of the bachelor's program for biomedical science and medicine were recruited for this study. Participants were healthy, right-handed, Dutch speaking volunteers with no history of psychiatric or neurological illness at the time of recruitment. Participants first completed two counterbalanced weeks of stress assessments in daily life (i.e., Ecological Momentary Assessments, EMA), one with an ecological exam stressor and the other without. At the end of each of the weeks, participants completed a series of computer tasks and a questionnaire battery. This was followed by two counterbalanced fMRI sessions, one with a stress induction procedure using a modified version of the

socially evaluated cold pressors (SECPT, Schwabe & Schächinger, 2018), and the other with a matched control task. All procedures were approved by the local ethics committee (Figure 1).

Five participants withdrew prior to completion of all scanning sessions. Due to issues that occurred during scanning, an additional three participants were also excluded. Reasons included incorrect stimuli presentation during scanning (1), scanner related malfunctions (1), and incidental findings (1). Finally, due to the COVID-19 outbreak an additional five participants were unable to complete either one or both scan sessions and were thus excluded from the MRI portion of the study, bringing the total number of participants to 70 (f=43 (61.4%)).

Real-life affective reactivity

Participants completed repeated EMA surveys (or beeps) delivered to their phones six times a day for two separate weeks: One during a high-stakes examination period (i.e., a stress week), and the other outside these period (i.e., control week). Surveys assessed stress levels as predictors and affect as an outcome. Stress was assessed using question regarding event-related stress relating to the most stressful experienced event, activity-related stress relating to the activity participants were engaged in when answering the surveys, and social-related stress relating to social context in which participants were in at the time of the beep. Affect items were collected for positive and negative mood. Additionally, ambulatory data was collected from wrist worn devices measuring aspects of physiological arousal. At the end of each of these weeks, participants filled in a questionnaire battery. Within the scope of the current paper, only subjective stress and positive affect measures are used from the EMA data. Full details and results of the EMA weeks are reported in previous work, and the full questionnaire set can be found in the associated GitHub directory[31].

Laboratory stress and the neural response

We investigated laboratory stress using a multi-day fMRI paradigm. Participants took part in three MRI scan days: One structural scan day, and two functional scan days (stress and control days, order counterbalanced between participants). Structural scans were used to reduce scanner related apprehension in scanning naïve participants. Those with prior scanning experience were only scheduled for the two functional scan days, and structural scans were appended to the end of their first fMRI session. Participants were asked to be onsite two-hours prior to the functional scans between 10:00 and 16:00 to allow cortisol levels to return to baseline prior to testing, and to account for diurnal cortisol fluctuations. During this time, participants practiced the tasks they would later perform in the scanner

(reported in the fMRI Task Section, Figure 1C). Following the rest period, participants were escorted to the MRI scanner, where skin conductance electrodes and a PPG heart rate sensor were attached to their left hand, and respiration belt was attached below the chest. The stress sessions included a modified version of the Socially Evaluated Cold Pressor Test (SECPT), and the other a matched control protocol[61]. During the SECPT, participants had their foot immersed in cold water while laying on the scanner bed and were then asked to perform a difficult mental arithmetic task. The control procedure was matched for time with room temperature water, and a simple arithmetic task. Saliva samples were collected right before the start of the SECPT/Control procedures (T=-3min), and then again at T=14min, T=42min, T=87min and T=160min (Full protocol and operating procedures in the supplementary material text). Participants were not informed which fMRI session would occur first, nor what the details of the stress procedure entailed beyond having their foot placed in water.

fMRI Tasks

Participants performed a series of fMRI tasks in two phases immediately following the SECPT/Control procedure. The first phase examining stress reactivity (i.e., the early phase) followed the SECPT/control procedure and consisted of three task runs of 11.5 minutes each, with two interleaved resting state runs of five minutes in between (Figure 1B) for a total of approximately 50 minutes including saliva sampling following the first resting state run. Participants were then taken out of the scanner for a short 20-minute break before continuing with the second half of scanning (hereafter the late phase). A total of 21 different task blocks were presented during each run, and a total of 21 blocks of each task were shown in each phase of the scanning session. Task blocks consisted of 27 seconds, followed by 6 seconds of rest before the next task. During this time, participants were shown a shape on the screen that indicated what the next task would be. Task order was pseudorandomized to maximize the number of times participants had to switch from one task to another. Tasks were selected based on previous evidence implicating the involvement of one of the three networks of interest (i.e., SN, ECN, DMN) as follows:

Oddball. A facial oddball paradigm was selected for Salience Network recruitment based on previous research showing robust SN activity during the presentation of an oddball stimulus[36]. Participants were shown a stream of faces with one standard neutral face being shown most of the time, and a novel oddball face with an emotional expression being shown 16% of the time. Participants also performed a facial recognition task outside the scanner following each phase (i.e., early and late), with 40 oddball targets shown in the scanner, and 20 lures. Faces were selected from four databases: Amsterdam Face

Database, Radboud Face Database, Karolinska Directed Emotional Faces, and the Chicago face database[62–65].

2-Back. Working memory tasks are often used to recruit the Executive Control Network[66]. To this end, we selected a 2-back task. Participants were presented with a stream of numbers and were instructed to press a button when the most recent number they had seen was the same as the number two places back (e.g., 3,1,3). A total of 15 trials per block were presented. Individual trials lasted for 1.8 seconds, with a stimulus being present for approximately 400 milliseconds[15].

Associative Retrieval. The default mode network has been shown to be active during internal thought processes, and memory retrieval tasks. To this end, we used an adapted version of an associative retrieval task[50]. On the scan day, but prior to the scan, participants performed two encoding sessions spaced 45 minutes apart to simulate the time in between the two scan phases (i.e., early and late encoding) during that day. During the encoding sessions, participants viewed a series of images that were negatively or positively valent which moved to one of four corners of the screen. Participants were told that they would be tested on the object-location association in the scanner. During scanning, participants engaged in a retrieval task where they were presented with the images from the encoding session and asked to indicate the location of the image using a button box.

fMRI Data Acquisition

Data was acquired on site using 3T MAGNETON Prisma and PrismaFit MR scanners (Siemens AG, Healthcare Sector, Erlangen, Germany) using a product 32-channel head coil. Participants were scanned on the same scanner for both of their functional sessions. T1-weighted images were acquired in the sagittal orientation using a 3D MPRAGE sequence with the following parameters: TR/TI/TE = 2300/11000/3ms, 8° flip angle, FOV 256 mm × 216 mm × 176 mm and a 1 mm isotropic resolution. Parallel imaging (iPAT = 2) was used to accelerate the acquisition resulting in an acquisition time of 5 minutes and 21 sec. Functional images were acquired with a multiband (3), multiecho (3) sequence with the following specifications: TR/TE1-3 1500/13.4/34.8/56.2ms, 75° flip angle, FOV 84 mm × 84 mm × 64 mm and a 2.5 mm isotropic resolution, multiband factor=3. Parallel imaging (iPAT = 2) was used to accelerate the acquisition.

EMA Data Analysis

We adopted a residual-based stress score to derive a measure of real-life affective reactivity to stress [25]. First, two general linear mixed effects models were used to estimate the effect of stress exposure (i.e.,

control or exam week) on stress reports (model 1) and positive affect (model 2). For model 1, an aggregate measure of stress was calculated from the total of the event, activity, and social stress scales. When participants reported stress levels above their individual means, surveys were labelled as "More Stress". Responses below the individual mean were labeled as "Less Stress". This was done to reduce affect-congruency effects and was modeled using a binomial family. Model 2 used positive affect as a continuous variable. In both models, we added a random effect of subject, and a random slope and intercept for week type effects. Random effects of these models would therefore indicate the subject level impact of stress on subjective positive affect. Random effects were then modeled against each other using linear regression to derive the overall change in positive affect relative to experienced stress during the stress week. Residuals from these models were used as a residual-based affective stress reactivity measure. In order to make interpretation easier, inverse scores are presented where increased scores represent increased affective reactivity in real life. This score was used in following fMRI models.

fMRI Arousal Measures

We next established the validity of our laboratory stress induction procedure by examining the effects of stress induction via the SECPT on biophysiological measures of salivary cortisol and heart rate (IBI and RMSSD) using mixed effects models with subject as a random effect and session (i.e., stress or control), time, and scan order as fixed effects. Sex was controlled for as a fixed effect. Interactions were modeled for all three of these predictors. Session was modeled with a random slope and intercept. Distributions of the data were examined, and model families were chosen according to the optimal fit as determined by residual normality and the Akaike Information Criterion (AIC). Follow-up models were carried out using the "emmeans" package. The difference in the area under the curve with respect to increase (AUCi) for the cortisol response was calculated per subject using the first four samples to derive individual measures of cortisol stress reactivity ($AUCi_{stress} - AUCi_{control}$).

fMRI Data Analysis

Standard fMRI processing steps were carried out including weighted echo recombination, registration, distortion correction, slice time correction, motion correction using ICA-AROMA, and finally high pass filtering (full details of acquisition protocol and preprocessing reported in SM Text 4). First level fixed effects models were then constructed at the run level. For the 2-Back task, we modelled the occurrence of a 2-Back target and the non-targets as events and the contrast 2-Back>non-target used to model ECN activity. For the oddball task, oddball and standard trials were each modelled as events. The contrast used to measure SN activity was the Oddball>Standard trials contrast. Finally, for the associative retrieval

task, remembered and forgotten trials were each modeled, and the contrast Remembered>Forgotten was used to model DMN activity.

Second level analysis was performed for each subject using a 2x2 fixed effect design, with session (control vs stress) and phase (early: runs 1-3 and late: runs 4-6) as fixed effects resulting in four main EVs: Control Early, Control Late, Stress Early, Stress Late. Contrasts were modelled for mean task related activity across the four EVs. Z-statistic images were thresholded using non-parametric cluster-based thresholding at Z>3.1 and a corrected cluster significance threshold of p=0.05. Previously established templates were used for ROI analyses, with a single mask created from all subdivisions of the SN, ECN, and DMN [67]. Parameter estimates were extracted from threshold z-statistic images of specific target task-network combinations for each of the four levels (Control Early, Control Late, Stress Early, Stress Late).

We first investigated the main effects with three-way interactions for session (stress/control), phase (early/late), and network (SN/ECN/DMN). We next investigate the specific hypothesis regarding temporal shifts as a function of stress in each network individually, with additional interaction terms modeled for cortisol stress reactivity (Stress*Phase*AUCi) and the real-life affective reactivity score (Stress*Phase*Real-LifeAffectivity). A maximal fitting approach was used in which subject was modeled as a random effect, and correlated random intercepts and slopes modeled for all other fixed effects of interest. Since AUCi and the real-life stress score only have one level per subject, no random slopes or intercepts were added for these terms. Scan order was also modeled as a fixed effect with no random slope and intercept to control for potential differences due to when sessions occurred.

fMRI Task Data Analysis

Task-related MRI measures were analyzed using mixed effects models, with individual trials being modelled for each of the three tasks. Trials with reaction times lower than 200ms were removed prior to analysis. For the 2-back, reaction time, error count, and a combined score (LIASES score) were analyzed[68]. For the oddball task, the average reaction time per trial and the oddball facial recognition (as measured by dprime, d') and the number of hits and misses were analyzed. For the associative retrieval task, reaction times and recall percentages were analyzed. In all models, session (i.e., stress or control MRI) were modelled as fixed effects with an interaction term for phase (i.e., early or late). A random effect was modelled for subject, with a correlated random intercept and slope modelled for session and phase effects. The same models used for the reaction time were used for the out-of-scanner oddball recognition task. We additionally ran separate models with an interaction term modelled for neural

activity in the targeted ROI's to examine the relationship between task performance and neural responses, with random slopes and intercepts also modelled for ROI activity.

Results

Deriving a residual-based stress score

In order to first establish the validity of our real-life stress paradigm, we first verified whether exam weeks were more stressful than control weeks without exams using EMA. During these weeks, students received repeated assessments (i.e., beeps) probing stress and affect. We used a measure of stress combining three types of stress: Event-related, activity-related, and social stress. Positive affect was measured using a 4 item questionnaire adapted from previous EMA studies (See [31] for full results and details). Participants showed a significant increase in the numbers of beeps during the exam week in which stress was above their overall average (Odds Ratio=-1.67, SE=0.14, t-stat=6.033, p<0.001, Figure 2A) in addition to lower overall positive affect (β=-0.77, SE=0.11, t-stat=-6.80, p<0.001, Figure 2B) compared to the control week.

To derive subject-level estimates of the effects of exam weeks on mood and stress levels, random effects were extracted from the mixed effects models. Implementing this approach, as opposed to a simple average and correlation measure, accounts for within subject variation, thus providing better estimates of within-subject effects[32]. A significant negative within-subject association between experiences of increased stress and reduced positive affect was found, indicating individuals with increased perceived stress had greater reductions in positive affect (β=-0.84, SE=0.17, t-stat=-5.09, p<0.001, Figure 2C). By extracting the residuals from this analysis, a measure can be derived of the deviation from the expected change in positive affect relative to the amount of reported stress exposure. That is, residuals above the line indicate a lower impact of stress exposure on affect, and vice versa for those below the line. The inverse of these residuals can thus be used as a residual-based affective reactivity score to estimate the within-subject reactivity controlling for potential differences in individual levels of exposure to stress (Figure 2C).

As a final step, we wanted to investigated whether inter-individual differences in stress reactivity were related to personality or trait characteristics that are linked to psychopathology. This was used to determine whether our derived measure of real-life reactivity corresponds to measures that are relevant for mental health. Therefore, we correlated this measure with state anxiety (STAI), depression symptoms (BDI) and scales of the NEO-FFI. Within the NEO-FFI, there were trend-level associations between

neuroticism and the residual-based score (r=0.21, p=0.054). This indicates that participants who experience greater changes in mood relative to the same level of perceived stress as their peers also exhibit personality traits that are associated with increased risk for psychopathology.

Figure 2. Residual Based Regressor. The averaged effects of examination weeks on (A) subjective stress levels and (B) positive affect, with individual-colored lines shown for the modeled random subject level effects. (C) These random effects (RE) were extracted from each model and then modeled against each other to derive the relative change in affect with respect to change in stress exposure. Blue dots above the line indicate less stress reactivity, red dots under the regression lines indicate more stress reactivity.

Increased physiological measures in response to SECPT

We next investigated whether our laboratory stressor induced changes in physiological arousal. This was done to establish the effectiveness of our laboratory stress induction procedure. Results of the mixed model showed a significant stress (stress, control) by time (1-5 samples) effect on salivary cortisol levels (log transformed estimate, β=1.00, SE=0.006, t-stat=2.44, p=0.015, Figure 3A). There was an additional significant effect of sex (β=-0.16, SE=0.08, t-stat=-2.06, p=0.040), with the difference being driven by attenuated cortisol responsivity in hormonal contraceptive users specifically (β=0.26, SE=0.10, t-stat=2.69, p=0.007). Follow-up tests showed significantly lower cortisol in the stress session relative to the control session immediately following the stress induction procedure following Tukey correction (T=9 minutes, β=0.173, SE=0.084, p=0.039), but expectedly higher cortisol at the third and fourth samples (T=39 minutes, β =-0.296, SE=0.84, p<0.001, and T=85 minutes, β=-0.165, SE=, p=0.0497). No significant

differences in cortisol were observed in the last sample (T=160 minutes, SE=-0.101, t-stat=0.084, p=0.229). We used the difference in the area under the curve with respect to increase (AUCi) in the early phase as a base measure of cortisol stress reactivity in later models. To this end, a paired-sample t-test showed that AUCi of salivary cortisol was significantly higher in the stress session compared to the control session when controlling for scan order effects (M_{diff}=103.86, t=3.975 p<0.001).

Figure 3. Physiological changes in response to SECPT. A) Salivary cortisol response showing increased cortisol in response to the SECPT compared to the control procedure. B) Average heart rate (beats per minute) and C) average heart rate variability (RMSSD, ms) per scanner run separated by whether participants had the control or stress scan first. Participants who had the stress session first exhibited increased heart rate and decreased heart rate variability (RMSSD) during the stress session. Error bars = SEM.

We examined further physiological reactivity to stress by looking at average heart rate and heart rate variability during the scanner runs (starting at 18, 30, 41, 53, 60 minutes post stress in the early phase, and 98, 111-, 117-, 128-, and 135-minutes post stress in the late phase). Within average heart rate, there was a main effect of time (β=-0.11, SE=0.04, t-stat=-2.90, p=0.004), a stress by scan order interaction (β=-2.51, SE=0.81, t-stat=-3.08, p=0.002), and a time by scan order interaction (β=-0.12, SE=0.04, t-stat=-3.20, p=0.001). A three-way interaction between stress, time, and scan order was also present for the effects of stress on average heart rate (BPM, β=0.20, SE=0.05, t-stat=3.83, p<0.001). Overall, participants who had the stress scan first showed increased heart rate during the stress session, with no differences being seen in those with the control scan first. Additionally, over time heart rate in the stress session

remained higher than in the control session only in those who had the stress scan first (Full model results in SM text 1).

There was also a significant main effect of time on heart rate variability (RMSSD, β=-0.13E-3, SE=0.052E-3, t-stat=-2.494, p=0.013), and significant interactions between stress and time (β=0.223E-3, SE=0.076e-03, t-stat=2.933, p=0.003) and stress and scan order (β=1.894E-3, SE=0.798E-3, t-stat=-2.373, p=0.018) during scanning. Follow-up tests indicated that participants who had the stress scan first showed reduced heart rate variability in the stress session (β=-0.003, SE=0.001, t-stat=-3.022, p=0.003, Figure 3). No significant effects were seen in participants who had the control session first. Additionally, over time, there appeared to be a significant increase in heart rate variability in the control, but not the stress session (See SM Text 1 for full results). Together these results indicate increased arousal in response to the stress induction procedure.

Task-Network Activations in Salience and Executive Control Networks

Before testing our main hypothesis with regards to temporal shifts in network activation under stress, we first verified whether the oddball, 2-back, and memory encoding task induced the expected network activation. With regards to the fMRI task-related activity, we first investigated network (i.e., Salience, Executive Control, and Default Mode networks) based task-related activation in the contrasts of interest using mixed-effects models on the subject-level fMRI estimates. Contrasts for network activity were modelled using only regressors from the specific tasks used for each network. SN activity was extracted from the SN mask for the contrast oddball > standard trials, ECN activity from the 2-Back > no-back trials contrast from the ECN mask, and DMN activity was from the remembered > forgotten trials from the DMN mask.

We found a significant main effect of network (F(2,76.9)=170.87, P<0.001), indicating differences in how well our tasks recruited the different networks and significant stress by phase (F(1,620)=5.488, P=0.019) and stress by network effects (F(2,620)=3.898, P=0.021), indicating that stress impacts neural activity differently between the networks. As our hypothesis pertains to specific changes within networks over time, further models were fit for each of the SN, ECN, and DMN separately. There was expected significant SN-related activity in the Oddball>Standard contrast (β=1.11, SE=0.06, t-stat=18.21, p<0.001, Figure 4A) and significant activation in the ECN to the 2Back>non-target trials (β=0.57, SE=0.07, t-

stat=8.73, p<0.001, Figure 4B). Contrary to expectations, we saw a significant overall decrease in the DMN for the Remembered>Forgotten contrast (β=-0.35, SE=0.05, t-stat=-7.45, p<0.001, Figure 4C).

Figure 4. Main Task Effects. Main task effects across all task runs for each of the contrasts for the ECN/2-back task (A, MNI-coordinates (XYZ mm): 27 33 29), SN/oddball Task (B, MNI-coordinates (XYZ mm): 41 91 35), and DMN/associative retrieval task (C, MNI-coordinates (XYZ mm): 41 91 35). Increased activity seen within expected regions of interest as well as some overlap in the SN and ECN contrasts, as well as an unexpected decrease in the DMN contrast. Results are whole-brain corrected with a cluster forming threshold of Z>3.1 and whole-brain cluster significance level of P<0.05.

Stress results in shifts in SN and DMN

We next tested our main hypotheses regarding temporal shifts under stress in the networks across time in each network separately. Within the SN, there was a significant stress by phase interaction effect (β=0.22, SE=0.07, t-stat=02.99, p=0.003), with increased SN activity in the early phase of the stress response relative to the control condition (β=0.35, SE=0.13, t-stat=2.77, p=0.007), but not in the late phase (β=-0.10, SE=0.13, t-stat=0.77, p=0.444). Within the SN, there was an additional reduction of activity in the late stress session compared to the early stress session (β=-0.44, SE=0.15, t-stat=-2.99,

p=0.003). Within the DMN, there was an overall decrease in activity in the stress session compared to the control session across both phases (β=0.11, SE=0.04, t-stat=2.40, p=0.016). Furthermore, there was a significant main effect of scan order on DMN, with greater suppression seen in participants who had the stress scan first (β=-0.11, SE=0.05, t-stat=-2.24, p=0.025). No significant stress effects were seen in ECN activity (Main effect of stress: β=-0.02, SE=0.09, t-stat=-0.18, p=0.858, Figure 5A).

SN stress reactivity is related to affective reactivity in real life

We finally investigate the association between shifts in the large-scale networks under stress and measures of laboratory and real-life affective reactivity (the AUCi difference in cortisol, and the residual-based reactivity scores respectively) using mixed effects models with network as a factor. Contrary to expectations, no significant relationship between cortisol stress reactivity and any of the large-scale networks were seen even when controlling for sex (analysis results can be found in the SM). There was however a significant three-way interaction between stress, phase, and real-life affective reactivity within the SN, but not the ECN and DMN, indicating that affective reactivity in real life was linked to changes in the stress response in SN over time (β=-0.23, SE=0.10, t-stat=-2.17, p=0.030). Follow-up tests showed that effects were driven by changes in the early phase of the stress response (β=0.71, SE=0.35, t-stat=-2.05, p=0.045) as opposed to the late phase of stress recovery(β=0.18, SE=0.35, t-stat=0.50, p=0.622), indicating that stress reactivity in daily life was associated with increased SN reactivity to stress in the early phase, but not the later recovery phase of the neural stress response (Figure 5B).

Figure 5. Effects of stress on large scale networks over time and real-life stress reactivity. A) Stress-Control differences in each of the networks in both the early and late phases of the stress response showing overall suppression of DMN under stress, and an increase in SN activity in early phase of stress reactivity. B) Increased SN reactivity to stress is related to increased real-life stress reactivity. Error bars = SEM.

Stress related SN activity enhances vigilance

We finally investigated associations between changes in networks under stress and performance measures from each of the corresponding tasks. This was done to investigate the impact of stress not just on brain activity, but also on task performance associated with the networks. In the oddball task, participants reacted significantly faster to the presentation of an oddball stimulus in the stress compared to the control session (β=0.97, SE=0.01, t-stat=-3.09, p=0.002). There was an additional main effect of time, with average reaction times being slower in the late phase of the MRI scan (β=0.98, SE=0.01, t-stat=-4.09, p<0.001). The addition of ROI measures for SN activity resulted in non-significant results. An exploratory mediation analysis was run to examine whether this was due to significant stress effects in SN under stress. Mediation models were run with simple slopes and subject as a random effect. SN activity significantly mediated the relationship between stress and reaction times during the

presentation of oddball stimuli (β=0.0361, 95% CI=[-0.819, 0.00], p=0.03). This indicates that increased SN activity under stress enhances reaction times in a vigilance oriented task.

Participants also performed worse during on the oddball facial recognition outside of the scanner in the stress session, as measured by d-prime (β=-0.42, SE=0.06, t-stat=7.55, p<0.001, Figure 6A). This was driven by both fewer hits (β=-3.29, SE=0.67, t-stat=4.87, p<0.001, Figure 6B), and more false alarms (β=1.68, SE=0.42, t-stat=4.04, p<0.001, Figure 6C). Interestingly, there was also a significant interaction effect between SN activity and session on false alarms across participants (β=1.05, SE=0.48, t-stat=2.18, p=0.030, Figure 6D). Post-hoc tests looking at control-stress showed a significant difference between the slopes (β=-1.05, SE=0.5 t-stat=-2.098, p=0.0371) with increased SN activity being associated with fewer false alarms in control session (slope=-0.588) and more false alarms in the stress session (slope=0.461).

Figure 6. Oddball Task Measures. D-prime (A), Hits (B), and False alarms (C) differed significantly between the stress and control

sessions, with lower d-prime under stress driven by both fewer hits,

and more false alarms. SN (salience network) activity was also

related to false alarms differently in the stress and control session

(D).

A one sample, two-sided t-test against the expected chance level (25%) in the DMN memory retrieval task showed that participants performed significantly above chance levels (Mean accuracy=61.5%, t-stat=255.65, p<0.001). There were no significant stress effects, or phase effects in the reaction times, nor in the ability to recall images. There was however a significant effect of valence. Participants responded faster (β=-0.0018, SE=0.0009, t-stat=-2.063, p=0.0392) and with more accuracy (β=0.027, SE=0.004, t-stat=7.266, p<0.001) to images that were negatively valanced compared to neutral ones. There were no significant differences between the control and stress session in the 2Back/ECN task when looking at the reaction times, proportion of errors, nor the LIASES score (Null results in SM text 1).

Discussion

In this study, we aimed to tested how brain activity changes during the course of the stress response, and whether changes in large-scale neural networks under stress were associated with real-life stress reactivity. Laboratory stress induced a network-dependent and time-dependent shift in BOLD activity, with an early increase and a later decrease in responsiveness of the salience network (SN), alongside increased suppression of the default mode network (DMN) throughout. Importantly, increased early SN reactivity to stress was associated with increased affective reactivity to stress in real life as measured with ecological momentary assessment (EMA). These results show how neural networks dynamically change in response to (laboratory induced) stress. Crucially, these inter-individual neural networks responses are related to affective reactivity to real-life stressors.

Affective reactivity to daily life stress was linked to SN reactivity to salient stimuli in the early, catecholaminergic dominated, phase of the stress response. By contrast, although we found a return back to baseline, we found no association with SN activity during the late phase of the stress response, which is thought to be dominated by genomically driven effects of glucocorticoids. Interestingly, our affective reactivity measure corresponds to in-the-moment stress, which falls into a time scale similar to that of the early phase of the acute stress response. Our results are in line with the two previous studies that connected real-life affective dynamics to neural measures. One study was able to link negative affect inertia to increased responses to social feedback in the insula – a core node of the SN[33]. Another study found a link between stress-induced dopaminergic activity using PET to psychotic reactivity to stress in real life[34]. The usage of PET imaging however does not allow for investigations of temporal shifts as a result of hormonal changes relating to stress response. Utilizing fMRI, we were able to link real-life measures to the full scope of the stress response, including both the reactivity and recovery phases.

In our data, stress related SN activation mediated decreased reaction times to unexpected oddball stimuli in the stress session, indicating heightened vigilance supported by SN activity. Previous evidence has linked the SN to vigilance and attentional reorienting mechanisms[35,36] which are enhanced by stress[37]. As suggested by our findings, in real-life, this may translate to increased vigilance in stressful situations, resulting in greater affective reactivity to stressors. This mechanism may possibly be related to increased attention to threats or negative events, and may be at the core of attentional bias to negative or stressful events typically observed in anxiety and depression[38].

The fMRI results additionally confirm the hypothesis based on the model proposed by Hermans and colleagues (2014) that the early stress response is characterized by a shift towards increased SN activity in a task-based setting[1]. This is in line with findings of increased functional connectivity in the SN in response to stress[7,39,40], with the process thought to be driven by locus coeruleus (LC) norepinephrine release[3,12]. Norepinephrine release is accompanied by activation of the sympathetic autonomic nervous system, resulting in increased heart rate which often corresponds with reduced heart rate variability, as also found in our study[15]. Finally, we found a return to baseline in SN activity in the later recovery stage of the stress response where corticosteroids have been shown to be involved[1,17]. This indicates that the initial shift in stress systems is later reversed when recovering from a stressor.

We found suppressed DMN activity during the early phase of the acute stress response, which is in line with previous studies[16,41]. More surprisingly, this suppression persisted in the recovery phase, two hours after the onset of the stressor. The DMN is implicated in memory retrieval functions, as well as in self-referential processing, rumination[42–45]. Yet no impairment of retrieval functions as a function of stress was seen. The reduction in DMN may instead indicate decreased internal attention, which is the cost of increased SN driven external attention in the early phase[43,46]. Additionally, no links between the DMN and real-life stress were established. While previous evidence has linked the DMN to depressive rumination and negative affect in daily life[45], our findings do not support the role of DMN driven rumination as a mechanism of affective reactivity to stress.

Worth noting is that the associative retrieval task did not elicit increased DMN activity relative to a fixation baseline, in contrast with previous evidence[42,46–48]. Our specific paradigm may not allow for sufficient DMN engagement given the traditional view of the DMN as a task negative network[49]. Another possible explanation may come from a deviation from the original protocol which used only neutral images[50]. Our design utilized both negative and neutral images. Negative images are often used by conditioning and stress research to elicit reactive stress responses[51,52]. Given the higher accuracy and faster reaction times to negatively valent images, it may be that valence-related mechanisms overshadowed previously reported task effects, resulting in suppression of the DMN in our contrast. Despite that however, we were still able to see stress effects on DMN in our study.

Based on previous literature we expected to see decreased ECN activity in the early reactivity phase of the stress response, and an increase in the aftermath of the stressor[1,15]. This latter process is putatively driven by genomic effects of cortisol. The lack of ECN suppression in the early phase may be due to coactivation of the ECN and SN, where increased SN activity results in a multi-tasking state of task

switching[40,53]. Such task switching would result in poorer performance on ECN tasks, though this was not the case in our data. It may instead be that arousal levels in our study population were insufficient to show this effect in our study, in addition to the lack of ECN upregulation in the later phase. Previous work has found administration of corticosteroids enhances ECN activity within the time frame of genomic corticosteroid effects following stress exposure[19,54]. Such studies used high doses of corticosteroids (peak measures of salivary cortisol at around 40-130 nmol/L) that may not be comparable to those resulting from the stress induction paradigm (around 5.6 nmol/L in our study). Higher doses in pharmacological studies, as well as individual variations in responses to stress induction may explain the lack of suppression. Finally, we found a significant sex difference in cortisol responses that was driven by blunted responses in males relative to hormonal contraceptive users. While important to investigate, such effects of contraceptive use and sex are beyond the scope and power of the current study, and thus we refrain from further interpretation.

In addition to neural findings, we also demonstrate the successful utilization of a dynamic EMA-based residualization method on a daily affect measure (Ref your preprint?). This measure is novel in capturing inter-individual differences in positive affect changes in response to in-the-moment stress. We focus on positive affect as our previous work has shown that stress seems to have a bigger impact on positive rather than negative affect[31]. This may be due to respondents showing greater variability when answering positive affect items, which is reflected in the skewed distribution of negative affect. Additionally, previous work has linked self-reported momentary resilience directly to positive affect in daily life[30]. By correlating this measure to neuroticism, we also show that it partially relates to an established personality trait that is linked to psychopathology[55].

While our results on stress dynamics in the lab and real life are novel, their long-term relationship to resilience is still an open question. Some studies have shown affective responses in daily life to be linked not only to momentary mood and psychiatric symptom expression, but also future mental health outcomes[56–58]. Thus, our measure may capture these same mechanisms at work in a shorter time frame. Some prospective studies have also investigated how long-term resilience is related to brain activity of large-scale networks. These studies have shown increased SN activation to be predictive of later mental health outcomes as well[39,59]. Together, these findings suggest that the mechanism leading to poorer mental health outcomes could be related to stress-induced enhancement of vigilance related processing that has a long-term impact.

In conclusion, our study demonstrates that neural stress reactivity is associated with stress reactivity in real life. These findings demonstrate ecological validity of neuroscientific research in the context of stress. Furthermore, partly in line with our hypothesis, we demonstrate a change in large scale networks under stress, with increased SN reactivity immediately follow threat that returns to baseline during stress recovery. Importantly, our results indicate that SN-related attentional mechanisms following stress are linked to affective reactivity in daily life. Meaning, individuals that show enhanced SN activity immediately following a laboratory stressor also show enhanced reactivity to stressful events in real-life contexts outside of the laboratory. Mechanisms such as increased vigilance under stress may have implications for our understanding of how stress related disorders develop. It may be that increased vigilance to threat can lead to impaired recovery from stress in the long-term, resulting in poorer mental health outcomes. Indeed, recent studies have indicated this hypervigilance may be risk markers for development of stress related disorders [60]. Future extension of these findings in prospective studies and clinical populations may help in uncovering the real-life consequences of neural underpinnings of these disorders.

References

1. Hermans, E. J., Henckens, M. J. A. G., Joëls, M. & Fernández, G. Dynamic adaptation of large-scale brain networks in response to acute stressors. *Trends Neurosci* **37**, 304–314 (2014).

2. Schwabe, L., Hermans, E. J., Joëls, M. & Roozendaal, B. Mechanisms of memory under stress. *Neuron* **110**, 1450–1467 (2022).

3. Hermans, E. J. *et al.* Stress-related noradrenergic activity prompts large-scale neural network reconfiguration. *Science (1979)* **334**, 1151–1153 (2011).

4. Joëls, M., Sarabdjitsingh, R. A. & Karst, H. Unraveling the Time Domains of Corticosteroid Hormone Influences on Brain Activity: Rapid, Slow, and Chronic Modes. *Pharmacol Rev* **64**, 901–938 (2012).

5. Hermans, E. J. & Fernández, G. Heterogeneity of cognitive-neurobiological determinants of resilience. *Behavioral and Brain Sciences* **38**, e103 (2015).

6. Menon, V. Large-scale brain networks and psychopathology: a unifying triple network model. *Trends Cogn Sci* **15**, 483–506 (2011).

7. van Leeuwen, J. M. C. *et al.* Disrupted upregulation of salience network connectivity during acute stress in siblings of schizophrenia patients. *Psychol Med* **51**, 1038–1048 (2021).

8. Kühnel, A. *et al.* Spatiotemporal Dynamics of Stress-Induced Network Reconfigurations Reflect Negative Affectivity. *Biol Psychiatry* (2022) doi:10.1016/j.biopsych.2022.01.008.

9. Joëls, M. & Baram, T. Z. The neuro-symphony of stress. *Nat Rev Neurosci* **10**, 459–466 (2009).

10. Aston-Jones, G. & Cohen, J. D. AN INTEGRATIVE THEORY OF LOCUS COERULEUS-NOREPINEPHRINE FUNCTION: Adaptive Gain and Optimal Performance. *Annu Rev Neurosci* **28**, 403–450 (2005).

11. van Marle, H. J. F., Hermans, E. J., Qin, S. & Fernández, G. From Specificity to Sensitivity: How Acute Stress Affects Amygdala Processing of Biologically Salient Stimuli. *Biol Psychiatry* **66**, 649–655 (2009).

12. Zerbi, V. *et al.* Rapid Reconfiguration of the Functional Connectome after Chemogenetic Locus Coeruleus Activation. *Neuron* **103**, 702-718.e5 (2019).

13. Henckens, M. J. A. G. *et al.* Interindividual differences in stress sensitivity: Basal and stress-induced cortisol levels differentially predict neural vigilance processing under stress. *Soc Cogn Affect Neurosci* **11**, 663–673 (2016).

14. Quaedflieg, C. W. E. M. & Schwabe, L. Memory dynamics under stress. *Memory* **26**, 364–376 (2018).

15. Qin, S., Hermans, E. J., van Marle, H. J. F., Luo, J. & Fernández, G. Acute Psychological Stress Reduces Working Memory-Related Activity in the Dorsolateral Prefrontal Cortex. *Biol Psychiatry* **66**, 25–32 (2009).

16. Zhang, W. *et al.* Acute stress alters the 'default' brain processing. *Neuroimage* **189**, 870–877 (2019).

17. Henckens, M. J. A. G., van Wingen, G. A., Joëls, M. & Fernández, G. Time-dependent effects of cortisol on selective attention and emotional interference: a functional MRI study. *Front Integr Neurosci* **6**, (2012).

18. Henckens, M. J. A. G., van Wingen, G. A., Joels, M. & Fernandez, G. Time-Dependent Effects of Corticosteroids on Human Amygdala Processing. *Journal of Neuroscience* **30**, 12725–12732 (2010).

19. Henckens, M. J. A. G., van Wingen, G. A., Joëls, M. & Fernández, G. Time-dependent corticosteroid modulation of prefrontal working memory processing. *Proceedings of the National Academy of Sciences* **108**, 5801–5806 (2011).

20. van Ast, V. A., Cornelisse, S., Meeter, M., Joëls, M. & Kindt, M. Time-Dependent Effects of Cortisol on the Contextualization of Emotional Memories. *Biol Psychiatry* **74**, 809–816 (2013).

21. Shiffman, S., Stone, A. A. & Hufford, M. R. Ecological Momentary Assessment. *Annu Rev Clin Psychol* **4**, 478–481 (2008).

22. Myin-Germeys, I. *et al.* Emotional reactivity to daily life stress in psychosis and affective disorder. *Schizophr Res* **53**, 236–237 (2002).

23. Myin-Germeys, I., Delespaul, P. & van Os, J. Behavioral sensitization to daily life stress in psychosis. *Psychol Med* **35**, 733–741 (2005).

24. Jacobs, N. *et al.* A momentary assessment study of the relationship between affective and adrenocortical stress responses in daily life. *Biol Psychol* **74**, 60–66 (2007).

25. van Harmelen, A.-L. *et al.* Adolescent friendships predict later resilient functioning across psychosocial domains in a healthy community cohort. *Psychol Med* **47**, 2312–2322 (2017).

26. Ioannidis, K., Askelund, A. D., Kievit, R. A. & van Harmelen, A.-L. The complex neurobiology of resilient functioning after childhood maltreatment. *BMC Med* **18**, 32 (2020).

27. Amstadter, A. B., Myers, J. M. & Kendler, K. S. Psychiatric resilience: longitudinal twin study. *British Journal of Psychiatry* **205**, 275–280 (2014).

28. Cahill, S., Hager, R. & Chandola, T. The validity of the residuals approach to measuring resilience to adverse childhood experiences. *Child Adolesc Psychiatry Ment Health* **16**, 18 (2022).

29. Sin, N. L., Graham-Engeland, J. E., Ong, A. D. & Almeida, D. M. Affective reactivity to daily stressors is associated with elevated inflammation. *Health Psychology* **34**, 1154–1165 (2015).

30. Hoorelbeke, K., van den Bergh, N., Wichers, M. & Koster, E. H. W. Between vulnerability and resilience: A network analysis of fluctuations in cognitive risk and protective factors following remission from depression. *Behaviour Research and Therapy* **116**, 1–9 (2019).

31. Tutunji, R. *et al.* Using wearable biosensors and ecological momentary assessments for the detection of prolonged stress in real life. *bioRxiv* (2021) doi:10.1101/2021.06.29.450360.

32. Rouder, J. N. & Haaf, J. M. A psychometrics of individual differences in experimental tasks. *Psychon Bull Rev* **26**, 452–467 (2019).

33. Provenzano, J. *et al.* Different Aspects of the Neural Response to Socio-Emotional Events Are Related to Instability and Inertia of Emotional Experience in Daily Life: An fMRI-ESM Study. *Front Hum Neurosci* **12**, 1–10 (2018).

34. Hernaus, D. *et al.* Psychotic reactivity to daily life stress and the dopamine system: A study combining experience sampling and [^{18}F]fallypride positron emission tomography. *J Abnorm Psychol* **124**, 27–37 (2015).

35. Krebs, R. M., Fias, W., Achten, E. & Boehler, C. N. Picture novelty attenuates semantic interference and modulates concomitant neural activity in the anterior cingulate cortex and the locus coeruleus. *Neuroimage* **74**, 179–187 (2013).

36. Krebs, R. M., Park, H. R. P., Bombeke, K. & Boehler, C. N. Modulation of locus coeruleus activity by novel oddball stimuli. *Brain Imaging Behav* **12**, 577–584 (2018).

37. Schwabe, L. & Wolf, O. T. Emotional Modulation of the Attentional Blink: Is There an Effect of Stress? *Emotion* **10**, 283–288 (2010).

38. Ellenbogen, M. A. & Schwartzman, A. E. Selective attention and avoidance on a pictorial cueing task during stress in clinically anxious and depressed participants. *Behaviour Research and Therapy* **47**, 128–138 (2009).

39. Zhang, W. *et al.* Acute-stress-induced change in salience network coupling prospectively predicts post-trauma symptom development. *Transl Psychiatry* **12**, 63 (2022).

40. Young, C. B. *et al.* Dynamic Shifts in Large-Scale Brain Network Balance As a Function of Arousal. *The Journal of Neuroscience* **37**, 281–290 (2017).

41. van Oort, J. *et al.* Absence of default mode downregulation in response to a mild psychological stressor marks stress-vulnerability across diverse psychiatric disorders. *Neuroimage Clin* **25**, 102176 (2020).

42. Spreng, R. N., Mar, R. A. & Kim, A. S. N. The Common Neural Basis of Autobiographical Memory, Prospection, Navigation, Theory of Mind, and the Default Mode: A Quantitative Meta-analysis. *J Cogn Neurosci* **21**, 489–510 (2009).

43. Davey, C. G., Pujol, J. & Harrison, B. J. Mapping the self in the brain's default mode network. *Neuroimage* **132**, 390–397 (2016).

44. Philippi, C. L., Tranel, D., Duff, M. & Rudrauf, D. Damage to the default mode network disrupts autobiographical memory retrieval. *Soc Cogn Affect Neurosci* **10**, 318–326 (2013).

45. Provenzano, J., Fossati, P., Dejonckheere, E., Verduyn, P. & Kuppens, P. Inflexibly sustained negative affect and rumination independently link default mode network efficiency to subclinical depressive symptoms. *J Affect Disord* **293**, 347–354 (2021).

46. Spreng, R. N. & Grady, C. L. Patterns of Brain Activity Supporting Autobiographical Memory, Prospection, and Theory of Mind, and Their Relationship to the Default Mode Network. *J Cogn Neurosci* **22**, 1112–1123 (2010).

47. Sestieri, C., Corbetta, M., Romani, G. L. & Shulman, G. L. Episodic Memory Retrieval, Parietal Cortex, and the Default Mode Network: Functional and Topographic Analyses. *Journal of Neuroscience* **31**, 4407–4420 (2011).

48. Harrison, B. J. *et al.* Consistency and functional specialization in the default mode brain network. *Proc Natl Acad Sci U S A* **105**, 9781–9786 (2008).

49. Raichle, M. E. *et al.* A default mode of brain function. *Proc Natl Acad Sci U S A* **98**, 676–82 (2001).

50. Wagner, I. C., van Buuren, M., Bovy, L. & Fernandez, G. Parallel Engagement of Regions Associated with Encoding and Later Retrieval Forms Durable Memories. *Journal of Neuroscience* **36**, 7985–7995 (2016).

51. Takai, N. *et al.* Effect of psychological stress on the salivary cortisol and amylase levels in healthy young adults. *Arch Oral Biol* **49**, 963–968 (2004).

52. Suzuki, A., Poon, L., Papadopoulos, A. S., Kumari, V. & Cleare, A. J. Long term effects of childhood trauma on cortisol stress reactivity in adulthood and relationship to the occurrence of depression. *Psychoneuroendocrinology* **50**, 289–299 (2014).

53. Sridharan, D., Levitin, D. J. & Menon, V. A critical role for the right fronto-insular cortex in switching between central-executive and default-mode networks. *Networks* **105**, 12569–12574 (2008).

54. Oei, N. Y. L., Tollenaar, M. S., Spinhoven, P. & Elzinga, B. M. Hydrocortisone reduces emotional distracter interference in working memory. *Psychoneuroendocrinology* **34**, 1284–1293 (2009).

55. Kotov, R., Gamez, W., Schmidt, F. & Watson, D. Linking "big" personality traits to anxiety, depressive, and substance use disorders: A meta-analysis. *Psychol Bull* **136**, 768–821 (2010).

56. Charles, S. T., Piazza, J. R., Mogle, J., Sliwinski, M. J. & Almeida, D. M. The Wear and Tear of Daily Stressors on Mental Health. *Psychol Sci* **24**, 733–741 (2013).

57. Leger, K. A., Charles, S. T. & Almeida, D. M. Let It Go: Lingering Negative Affect in Response to Daily Stressors Is Associated With Physical Health Years Later. *Psychol Sci* **29**, 1283–1290 (2018).

58. Vaessen, T. *et al.* Is sensitivity to daily stress predictive of onset or persistence of psychopathology? *European Psychiatry* **45**, 167–173 (2017).

59.   Grueschow, M. *et al.* Real-world stress resilience is associated with the responsivity of the locus coeruleus. *Nat Commun* **12**, 2275 (2021).

60.   Kredlow, A. M., Fenster, R. J., Laurent, E. S., Ressler, K. J. & Phelps, E. A. Prefrontal cortex, amygdala, and threat processing: implications for PTSD. *Neuropsychopharmacology* **47**, 247–259 (2022).

61.   Schwabe, L. & Schächinger, H. Ten years of research with the Socially Evaluated Cold Pressor Test: Data from the past and guidelines for the future. *Psychoneuroendocrinology* **92**, 155–161 (2018).

62.   van der Schalk, J., Hawk, S. T., Fischer, A. H. & Doosje, B. Moving faces, looking places: Validation of the Amsterdam Dynamic Facial Expression Set (ADFES). *Emotion* **11**, (2011).

63.   Langner, O. *et al.* Presentation and validation of the Radboud Faces Database. *Cogn Emot* **24**, (2010).

64.   Lundqvist, E., Flykt, D. A. & Öhman, A. The Karolinska Directed Emotional Faces - KDEF. Preprint at (1998).

65.   Ma, D. S., Correll, J. & Wittenbrink, B. The Chicago face database: A free stimulus set of faces and norming data. *Behav Res Methods* **47**, (2015).

66.   Owen, A. M., McMillan, K. M., Laird, A. R. & Bullmore, E. N-back working memory paradigm: A meta-analysis of normative functional neuroimaging studies. *Hum Brain Mapp* **25**, 46–59 (2005).

67.   Shirer, W. R., Ryali, S., Rykhlevskaia, E., Menon, V. & Greicius, M. D. Decoding Subject-Driven Cognitive States with Whole-Brain Connectivity Patterns. *Cerebral Cortex* **22**, 158–165 (2012).

68.   Vandierendonck, A. A comparison of methods to combine speed and accuracy measures of performance: A rejoinder on the binning procedure. *Behav Res Methods* **49**, 653–673 (2017).

69.   Poser, B. A., Versluis, M. J., Hoogduin, J. M. & Norris, D. G. BOLD contrast sensitivity enhancement and artifact reduction with multiecho EPI: Parallel-acquired inhomogeneity-desensitized fMRI. *Magn Reson Med* **55**, (2006).

70.   Jenkinson, M., Bannister, P., Brady, M. & Smith, S. Improved optimization for the robust and accurate linear registration and motion correction of brain images. *Neuroimage* **17**, (2002).

71.   Jenkinson, M. & Smith, S. A global optimisation method for robust affine registration of brain images. *Med Image Anal* **5**, (2001).

72.   Pruim, R. H. R. *et al.* ICA-AROMA: A robust ICA-based strategy for removing motion artifacts from fMRI data. *Neuroimage* **112**, (2015).

73.   Smith, S. M. Fast robust automated brain extraction. *Hum Brain Mapp* **17**, (2002).

74. Worsley, K. J. Statistical analysis of activation images. in *Functional Magnetic Resonance Imaging* (Oxford University Press, 2001). doi:10.1093/acprof:oso/9780192630711.003.0014.

75. Woolrich, M. W., Ripley, B. D., Brady, M. & Smith, S. M. Temporal Autocorrelation in Univariate Linear Modeling of FMRI Data. *Neuroimage* **14**, (2001).

Acknowledgments

This project was funded by the European Research Council grant ERC-CoG-682591/STRESNET.

Author Contributions

Study design: RT, MK, FK, LdV, EJH

Data Collection: RT, MK, NK

Data processing: RT

Data Analysis: RT, EV, EJH

Writing: RT, EV, EJH

Editing and Final approval: All authors

Funding: EJH