# Mapping the blood vasculature in an intact human kidney using hierarchical phase-contrast tomography

**Running title: Analysis of the vasculature in an intact human kidney**

Shahrokh Rahmani [1,2] , Daniyal J Jafree [3, 4, 5], Peter D Lee [1], Paul Tafforeau [6],Joseph Brunet[1,6] Sonal Nandanwar[1], Joseph Jacob [7,8], Alexandre Bellier [9], Maximilian Ackermann [10,11, 12] Danny D Jonigk [12,13], Rebecca J Shipley [1], David A Long [3,5], Claire L Walsh [1]

1. Department of Mechanical Engineering, University College London, London, UK, WC1E 6BT.
2. National Heart & Lung Institute, Faculty of Medicine, Imperial College London, London, United Kingdom
3. Developmental Biology and Cancer Research & Teaching Department, UCL Great Ormond Street Institute of Child Health, University College London, London, UK, WC1N 1EH.
4. UCL MB/PhD Programme, Faculty of Medical Science, University College London, London, UK, WC1E 6BT.
5. UCL Centre of Kidney and Bladder Health, UCL London UK
6. European Synchrotron Radiation Facility, Grenoble, France, 38043.
7. Satsuma Lab, Centre for Medical Image Computing, UCL, London, UK.
8. Lungs for Living Research Centre, UCL, London, UK.
9. Department of Anatomy (LADAF), Grenoble Alpes University, Grenoble, France, 38058.
10. Institute of Anatomy, University Medical Center of the Johannes Gutenberg University Mainz, Mainz, Germany.
11. Institute of Pathology and Department of Molecular Pathology, Helios University Clinic Wuppertal, University of Witten-Herdecke, Wuppertal, Germany.
12. Institute of Pathology, RWTH Aachen Medical University, Aachen, Germany.
13. German Center for Lung Research (DZL), Biomedical Research in Endstage and Obstructive Lung Disease Hannover (BREATH), Hannover, Germany.

**Correspondence:** Dr. Claire L. Walsh, Mechanical Engineering, University College London, London, UK, WC1E 6BT

Email: c.walsh.11@ucl.ac.uk

**ABSTRACT**

The architecture of the kidney vasculature is essential for its function. Although structural profiling of the intact rodent kidney vasculature has been performed, it is challenging to map vascular architecture of larger human organs. We hypothesised that hierarchical phase-contrast tomography (HiP-CT) would enable quantitative analysis of the entire human kidney vasculature. Combining label-free HiP-CT imaging of an intact kidney from a 63-year-old male with topology network analysis, we quantitated vasculature architecture in the human kidney down to the scale of arterioles. Although human and rat kidney vascular topologies are comparable, vascular radius decreases at a significantly faster rate in humans as vessels branch from artery towards the cortex. At branching points of large vessels, radii are theoretically optimised to minimise flow resistance, an observation not found for smaller arterioles. Structural differences in the vasculature were found in different spatial zones of the kidney reflecting their unique functional roles. Overall, this represents the first time the entire arterial vasculature of a human kidney has been mapped providing essential inputs for computational models of kidney vascular flow and synthetic vascular architectures, with implications for understanding how the structure of individual blood vessels collectively scales to facilitate organ function.

**INTRODUCTION**

The vascular system of the kidney is highly specialised, serving multiple functions

including delivery of oxygen and nutrients to the organ's parenchyma whilst also

facilitating plasma ultrafiltration and solute reabsorption. Despite only comprising

approximately 1% of body weight, the kidney receives up to 20% of cardiac output

[1]. Blood enters the kidney through the kidney artery which branches from the

abdominal aorta and enters the kidney hilum. Once within the kidney, the kidney

arteries divide hierarchically, first into segmental arteries which pass through the

kidney pelvis, then branching into interlobar branches which pass through columns

between the pyramids of the kidney medulla. At the distal end of the kidney columns,

interlobular arteries branch into arcuate arteries that arch around the outer surface of

the kidney pyramids. From these, the interlobular vessels branch and penetrate into

the surrounding kidney cortex before finally terminating at efferent arterioles [1]. This

complex network perfuses specialised glomerular capillaries for plasma ultrafiltration,

before peritubular capillaries located in the cortex, and vasa recta located in the

medullary pyramids facilitate dynamic solute exchange. Thereafter, venous return

follows the arterial supply out of the organ [2].

Structural and molecular changes to the kidney vasculature are a common

feature of kidney pathologies including multiple aetiologies of chronic kidney disease

(CKD) and transplant rejection in both animal models and patients [3] Therefore,

studying kidney vascular patterning has implications for understanding the basis of

kidney function in health and disease, and also aids surgical planning for tumour

resection, nephrectomy and transplantation. Vascular geometries also have a central

role to play in computational models which underpin the creation of digital twins, such

as through the creation of synthetic data [4], and flow modelling [4], [5], [6], [7], [8],

which are playing an increasing role in biomedical research.

Vascular imaging of the kidney has advanced following technological

innovations in micro-computed tomography ($\mu$CT) [9], [10], [11], magnetic resonance

imaging (MRI) [12], ultrasound [13], lightsheet microscopy [14], [15] and

photoacoustic imaging [16], [17]. These techniques have been used to generate

quantitative analyses of vascular network geometry in intact kidneys of model

organisms, particularly rodents where kidney diameter reaches up to 12 mm [18].

Comparatively human kidneys, with a diameter of approximately 5 cm [19] are far

more challenging to image at high resolution whilst still intact. Corrosion casting of

human kidneys has highlighted vascular heterogeneity and generated intricate 3D

casts (down to 100 $\mu$m) but has provided limited quantitative or accessible digitized

geometries of the vascular network [20]. Optical clearing and lightsheet microscopy
have been used to quantify portions of the human kidney vascular network [21]
however, it has not yet been possible to capture the intact vascular network of the
human kidney without physical sectioning the tissue, beyond approximately six vessel
divisions [22].  MRI has been used to quantify larger vessels both *in vivo* and post
mortem [23], [24], but lacks the resolution capable of imaging small vessels and
arterioles [23].
Due to these limitations, human kidney vascular network analysis is often
predominantly focused on the very large, first three branches of the arterial tree [23],
or on only small portions of the network [25]. Where multiscale modelling has been
performed, parameters from rodent kidneys are assumed to be representative of
human kidney vascular networks [4], [5], [8]. The semi-quantitative studies of human
kidney vascular casts have shown large anatomical variation in even the segmental
artery (first or second branch after the kidney artery) patterns [22], while smaller
vessels such as arcuate arteries, interlobular arteries and afferent or efferent
arterioles have not been assessed quantitatively at the organ scale.
Here, we show how a recently developed X-ray based non-destructive
imaging technique - hierarchical phase-contrast tomography (HiP-CT), can be used to
map and quantify the arterial vascular network of an intact human kidney down to the
arteriolar level for the first time. HiP-CT is a technique which leverages the European
Synchrotron Radiation Facility's (ESRF) Extremely Brilliant Source (EBS); a high-
energy fourth generation synchrotron source, to image intact human organs at
unprecedented scale and resolution. Previously we have demonstrated the feasibility
of applying HiP-CT to profile the human glomerular morphology and number across
cubic centimetres of intact human kidney [26].  We now extend the use of this
technology to extract and quantify the arterial network of an intact human kidney
across multiple length scales without using antibodies, dyes or contrast agents.
Within the human kidney, we delineated the extent and morphology of the
vasculature, down to afferent and efferent arterioles, quantifying variation in vascular
morphology within the context of vascular ordering schemes. This enabled
quantitative comparison between human and previously published rodent kidney
vascular networks, the latter of which has been used as inputs for biophysical
modelling of human vascular flow [4], [5], [6], [7], [8]. We further demonstrate regional
heterogeneity in the context of the anatomical compartments of the kidney. Such
variations highlight the link between regional structure and function, re-enforcing the

importance of quantitative analyses for understanding and modelling regional micro-
environments within the human kidney.
126
**METHODS**
**Sample preparation**
An intact human kidney was obtained from a 63-year-old male (cause of death:
pancreatic cancer) who consented to body donation to the Laboratoire d'Anatomie
des Alpes Françaises before death. Post-mortem study was conducted according to
Quality Appraisal for Cadaveric Studies scale recommendations [27]. The body was
embalmed by injecting 4500 mL of 1.15% formalin in lanolin followed by 1.44%
formalin into the right carotid artery, before storage at 3.6°C. During evisceration of
the right kidney, vessels were exposed, and surrounding fat and connective tissue
removed. The kidney was post-fixed in 4% neutral-buffered formaldehyde at room
temperature for one week. The kidney was then dehydrated through an ethanol
gradient over 9 days to a final equilibrium of 70% [27]. Each solution was four-fold
greater than the volume of the organ and during dehydration, the solution was
degassed using a diaphragm vacuum pump (Vacuubrand, MV2, 1.9m$^3$/h) to remove
excess dissolved gas. The dehydrated kidney was transferred to a polyethylene
terephthalate jar where it was physically stabilised using a crushed agar-agar ethanol
mixture, and then imaged [26], [27].
144
**Scanning, image acquisition and reconstruction**
Imaging was performed on the BM05 beamline at the ESRF following the HiP-CT
protocol [26], [27]. Initially the whole kidney was imaged at 25 μm per voxel (isotropic
edge length). Volumes of interest within the same kidney were also imaged at 6.5
and 2.6 μm per voxel. [26] Tomographic reconstruction was performed [26], [27],
using the PyHST2 software [28]. Briefly, a filtered back-projection algorithm with
single-distance phase retrieval coupled to an unsharp mask filter was applied to the
collected radiographs. Reconstruction parameters are provided in image metadata.
The reconstructed volumes were binned (averaged) to 50, 13, and 5.2 μm per voxel
respectively to reduce computational load for subsequent image segmentation and
quantification (**see Figure S1**). All reconstructed image volumes and metadata can
be accessed at human-organ-atlas.esrf.eu, table for direct DOI links for each dataset
is provide in **Table S1**
158
**Image filtering, enhancement, and segmentation**

Prior to semi-automated segmentation, images were filtered to enhance blood vessel

contrast using Amira v2021.1 software. A 3D median filter (iterations =2 and 26

neighbourhood analysis) was used to reduce image noise and image normalisation

was performed using background detection correction (Amira v2021.1; default

parameter settings). Semi-automated segmentation of the arterial networks was

performed in Amira v2021.1 using a manual region growing tool where the user

selects an initial voxel within the vasculature along with set intensity and contrast

thresholds. Any voxel within the connected neighbourhood of the initially selected

voxel that has an intensity and contrast within the thresholds are added to the region.

The annotator continues this process in an iterative fashion selecting seed points

altering the thresholds to expanding the region, (Method shown in **Supplementary**

**Video 3)**.  Once the primary annotator believes they have filled the interior of all

vessels the data is passed to a second annotator who repeats the process. A third

annotator (referred to as the proof-reader) will then quantitatively review the labels.

The data the proof-reader is presented with are randomised regions of 2D slices of

the data. They then count the number of vessels cross-sections present in the slice,

recoding the true positive and false negative number of vessel cross-sections that

have been segmented. The proof-reader returns the data to the initial two annotators

and the whole process repeats iteratively until the proof-reader does not find any

false negatives. This method was applied to segment the kidney arterial network from

the intact human kidney from the imaging data at 50 µm per voxel, and portions of the

same network in the 13 and 5.2 µm per voxel datasets.

A second approach to quantitative validation of the segmentation was

performed using smaller segmented regions of the 13 µm per voxel dataset. Here, the

higher resolution volume of interest at 13 µm per voxel was rigidly registered to the

whole organ volume with affine registration toolkit (Amira-Avizo) (See

**Supplementary Methods §1.1 and Tables S2 & S3**). Overlapping portions of the 13

µm voxel segmentations and 50 µm per voxel datasets were extracted and the 50 µm

per voxel datasets was up-sampled to the resolution of the 13 µm voxel dataset. An

overlap measure known as topological precision and recall score following Paetzold et

al. [29], was applied (see **Supplementary Information §2.1** & **Figure S3)**

**Visualization and skeletonization**

To quantify branching metrics of the human kidney vasculature, the segmented 3D

vascular network at 50 µm per voxel was skeletonized using the centreline tree

algorithm in Amira-Avizo. The choice of skeletonization algorithm and the

parameterising of the algorithm were optimised by utilising the super-metric approach

outline by Walsh & Berg et al. [7] (tube parameters: slope = 4 and zeroval = 10, see

**Supplementary Methods § 2.2 and Figure S4** for parameter selection method). The

resulting spatial graph describes the vessel network in terms of 'nodes', 'points',

'segments', and 'sub-segments'. A segment is defined as being between a start and

end node; which correspond to either a branching point leading into another segment

branch or a terminal end where no further branches were detectable. Between the

start and terminal node of each segment lie sub-segments with 'points', marking the

start and end of each sub-segment. Each sub-segment has an associated radius and

length (**Figure S5A**). A multiscale smoothing approach was applied to the larger

vessels (those of Strahler generation greater than 5), through a weighted smoothing

algorithm, and corrections for the radius of collapsed vessels (see **Supplementary**

**Methods § 2.3** for details). The final spatial graph was manually proofread to mitigate

errors in node locations and remove spurious branches in large collapsed segments

of vessels.

**Morphological analysis**

Topological/morphological metrics of the network were calculated from the spatial

graph as follows codes are provided at

https://github.com/HiPCTProject/Skeleton_analysis:

(i)  branching angle, calculated as either (a) the angle between the two child segments

from a common parent segment or (b) the angle between a child segment and its

parent segment. In both cases the vector for the segment of parent and child were

calculated between the start node and end node (i.e. ignoring vascular tortuosity);

(ii)  tortuosity defined as the Euclidean distance between start and end node of a

segment divided by the sum of all subsegment lengths;

(iii) radius calculated per segment as either, the mean of all subsegment radii, or for

larger vessels that had fully collapsed (See **Supplementary Methods § 2.3** for

details), as the equivalent radius for the perimeter of vessel cross-section in the

binary image;

(iv) length defined as the sum of all subsegment lengths;

(v)  inter-vessel distance calculated by two approaches to facilitate different analyses.

Firstly, using the segmentation binary image the distance of every non-vessel voxel

from its nearest vessel voxel was calculated *via* a 3D distance transform (ImageJ)

applied to the binary vessel segmentation. Secondly, using the skeleton form, the

Euclidean distance between the midpoint of every segment to its nearest-

neighbouring segment midpoint was calculated.

Additionally, we also assessed vessel generation or order using two methods. Firstly,

using the centripetal system known as Strahler ordering system [30], [31], [32], where

the most distal segments are assigned as the first order, if two segments with the

same order intersect, the resulting segment has order one greater. Alternatively, if

two segments with different orders intersect, the higher order of the two is given to

the resulting segment, **(Figure S5B).** Secondly, we took a centrifugal or 'topological'

approach, starting with most proximal artery as generation one, at each branching

node the generation is increased, this approach has been utilised by e.g. Pries and

Secomb [33] **(Figure S5C)**.

From the ordering analyses we assessed the branching ratio ($\gamma$) defined as the

antilog of the reciprocal for the linear fit to the plot of Stahler order (*O*) against the

logarithm of the number of segments (*N*) in each order:

$$N = N_0 e^{-\frac{O}{\gamma}} \qquad (1)$$

We also examined whether our data followed Murray's Law which states that the

cube of the parent segment radius should be equal to the sum of the cubed child

segment radii:

$$R^3_{parent} = \sum_{i=2}^{i=n} R_i^3 \qquad (2)$$

The radius of the arterial network in the human kidney obtained from this study were

compared to those of the rat kidney taken from[11] scanned with 20 and 4 µm voxels

using a micro-filling approach.

**Kidney Compartment Segmentation**

Segmentation of the compartments within the human kidney, including cortex,

medulla, inter-medullary pillars and hilum, was performed in Dragonfly (version:

2021.3) using a 2D convolutional neural network (CNN). The final hyperparameters of

the CNN are given in **Table S4**. Correction of the CNN output was manually

performed in Amira (v2021.1) to provide the final compartment delineation. These

compartments were used to group and then analyse vascular network parameters.

**Statistical analysis**

Statistical comparisons of vascular network morphology between human and rat

kidneys[11]  was performed in GraphPad Prism (version: 10.1.2). For all statistical

tests, a *p* value of less than 0.05 was considered statistically significant. Radius

against Strahler Order were normalised to the 9th Strahler Order (the largest vessel

for which the human data contained complete vessel segments). Log of radius

against Strahler generation for each of the human and rat datasets was plotted

facilitating a linear least squares regression analysis. A sum of squares $F$ test was

performed with the null hypothesis that a single set of global parameters for slope

and intercept would fit vessel radius or vessel length for both the rat and human

cases. For Murray's law the same sum of squares $F$ test was performed with the null

hypothesis that Murray's law would fit the human data; in this analysis automated

outlier detection was performed with the Graphpad ROUT method where Q is the

desired maximum false discovery rate, Q = 0.05%.

**RESULTS**

**HiP-CT can visualise the arterial vascular network in the intact human kidney**

**down to efferent and afferent arterioles.**

Using HiP-CT [26], [27] we imaged the whole intact kidney obtained from a 63-year-

old male organ donor in a hierarchical fashion. We initially performed an overview

scan of the entire kidney at 25 µm per voxel, followed by selecting and imaging

representative volumes of interest at 6.5 µm per voxel and 2.6 µm per voxel (**Figure**

**1A**). As these image volumes are inherently aligned, expert annotation was  applied

to the image volumes taken at each resolution to produced a multi-scale

segmentation of the arterial network (**Figure 1B and Supplementary Video 1**). From

the segmented data we were able to identify all known anatomical subdivisions of the

kidney arterial system (**Figure 1C**), down to arterioles that terminate in the

specialised plasma ultrafiltration units of the kidney: the glomerulus. The segmental

pattern of anterior, posterior, superior and inferior territories supplying the kidney

parenchyma were clearly delineated. Each vascular territory (**Figure 1D and**

**Supplementary Video 2**) had a corresponding kidney arterial branch originating from

the hilum, which bifurcated before hierarchical branching towards the cortical

parenchyma.

**Figure 1. Multi-level segmentation of the human kidney arterial network. A)** *Overview of the hierarchical image volumes that can be acquired with Hierarchical Phase-Contrast Tomography (HiP-CT). Brown, cyan and yellow volumes show the whole organ acquired at 25 µm, sub-volume acquired at 6 µm and sub-volume acquired at 2.6 µm, respectively, in the intact human kidney.* **B i-iv)** *showing the vascular segmentation performed across the three resolutions of HiP-CT data enabling the whole organ (***Bi***) through to glomeruli (red arrows ***Biv***) to be visualised and segmented.* **C)** *Diagram of the anatomical organisation of the human kidney arterial network.* **D)** *The vascular territories of the kidney imaged in this study.*

**An error-bounded image processing pipeline to reproducibly quantify the**

**arterial network in human organs.**

We next sought to quantitate the arterial network in a reliable and reproducible manner.

As we have previously shown that quantitative features of vascular networks are

influenced heavily by the image processing pipeline [7], we developed an image

processing pipeline (**Figure 2**), involving reduction of the initial image to a skeleton or

spatial graph representation of the arterial network. The graph representation

comprises a set of nodes; 3D locations where vessels meet or end, and the

connections between these nodes, defined as 'segments' (see **Figure S5A and Figure**

**4A**). Our pipeline, which is overviewed in **Figure 2**, comprises 8 steps which are fully

detailed in **Supplementary Methods §2**, and enables the generation of a spatial graph

from segmented HiP-CT with quantification of error at the segmentation and

skeletonization stages.

In brief, the pipeline included utilising the aligned higher resolution HiP-CT

volumes to provide quantification of the segmentation accuracy (**Figure 2, Step 1**).

Secondly, we applied three different skeletonization algorithms and utilised a formal

metric (the recently developed skeleton super-metric [7]) to assess and optimise the

skeletonisation process **(Figure 2 Step 2)**. The skeleton was then corrected (**Figure 2**

**Steps 3 – 8**) to overcome the two primary challenges of HiP-CT data. The multiscale

nature of the vasculature captured was corrected with a multi-scale smoothing

approach; **Figure 2 Steps 3 and 4.** Collapsed vessels were identified from outliers in

radial distributions (**Figure 2 Steps 5),** followed by automated extraction of vessel

cross-section and radius correction using the collapsed vessel perimeter **Figure 2**

**Steps 6-8.**

The result of our novel pipeline was the generation the first open-source spatial

graph of the human kidney arterial vasculature in its entirety. We were able to identify

97% of vessels <50 μm radius across the whole intact human kidney, with an imaging

resolution of 50 μm per voxel. The network consisted of 10,193 nodes, 376,603 points

and 10190 segments, or vessels. The total network volume was $1.68 \times 10^{12}$ μm$^3$ its was

length of $2.3 \times 10^7$ μm. This spatial graph, which is provided as spatial graphs in

**Supplementary data** captures all the morphological features and connectivity of the

human kidney arterial vasculature, which was then used for downstream analyses as

described below.

*Figure 2. Pipeline for the extraction and correction of the vascular network skeletonization.* *Step 1* *Segmentation is performed with quantitative validation using a higher resolution volume of interest,* *Step 2* *Skeletonization is optimised by comparison of skeletonization algorithms and optimisation of skeleton super-metric, the super-metric is a projection of the distance vector between the reconstructed skeleton sand the segmented image, onto a weighted space, it contains 5 contributing terms: network volume (Vol.), connected components (CC), Euler Number, Centerline sensitivity (cl sens.), Bifurcation DICE (BB DICE).* *Step 3* *An initial Strahler Order (O) calculation is made on the skeletonized network.* *Step 4* *Using the ordering from Step 3 the network can be split into larger calibre (O ≥ 5) and smaller calibre vessels (O > 5) , the larger calibre vessel can then be smoothed as shown in insets, orange arrows show the points where smoothing has noticeably acted on regions of larger vessels.* *Step 5* *Strahler order vs Mean radius is plotted for every segment (blue circles); outliers (red circles) are identified as segments with a radius below the 90% percentile for their Order.* *Step 6* *The segments identified as outliers are visualised and collapse status is manually confirmed.* *Step 7* *For vessels which are confirmed as collapsed, planes which are normal to the centreline of the vessel (indicated by orange arrows) are created at every point along the centreline and the 2D image for each plane is extracted (lower panels). From these 2D planes the collapsed vessel is identified (red cross) and the perimeter (yellow dashed line) is extracted.* *Step 8* *The perimeter is used to calculate an equivalent radius and assigned as the new radius of the segment.*

## Multi-scale generational and ordering analysis of the arteriolar vasculature in the human kidney.

Having produced a reproducible spatial graph of the human arterial vasculature of the

kidney, we then performed topological generation [33] and Strahler Ordering [30],

[31], [32] analyses. This resulted in nine Strahler Orders (**Figure 3A**) and twenty-five

topological generations (**Figure 3B**). As the main artery supplying the kidney was cut

during autopsy we can infer that 10 Strahler order, 27 topological generations, can be

imaged over the entire intact human kidney with HiP-CT.

Leveraging the hierarchical capability of HiP-CT we imaged regions of the

cortex at 2.6 µm per voxel and from these data, we segmented down to the afferent

arteriolar level, evidence by the presence of glomeruli at the terminal ends of these

vessels **(Figure 3Ei-3Eiii)**. Interestingly, we were also able to segment glomeruli from

non-terminal arteries, (the artery from which the arterioles branch have additional

bifurcations downstream, which are not arterioles) (**Figure 3Eii black arrow and**

**Figure S7**). This supports recent findings, [34] in the rat kidney, which demonstrated

the existence of non-terminal branch arterioles and their contribution to the

synchronicity of blood flow in the kidney [5], [35].

*Figure 3. Ordering and branching ratio analyses. Rendering of the vascular network with vessels coloured according to **A)** Strahler order and **B)** Topological generation. **C)** Plot showing the number of vessels per Strahler order with fit for the log plot to calculate branching ratio. **D)** Strahler order against cumulative vascular volume fraction. **Ei)** Strahler ordering down to efferent and afferent arterioles, using the 2.6 µm/voxel image dataset; **Eii)** showing the same region segmented at 5.2 µm/voxel (binned data); and **Eiii)** the same small region from the whole kidney overview (50 µm/voxel). Black arrows in **Ei** and **Eiii** indicate the same vessel in both cases with Strahler order of 1 or 3 respectively indicated by the colour cases. The black arrow in **Eii** indicates a bifurcation downstream of the arterioles that can be segmented, which is not an arteriole itself, i.e. this demonstrates the presence of non-terminal arterioles in the human kidney.*

Glomeruli branching from non-terminal arterioles  (see arrow in **Figure 3Ei** compared

to **3Eiii**) prevents the application approaches used by Nordlestten et al. [11]  to

estimate the Strahler Order of every terminal node in the whole arterial tree (extracted

from the 50 μm/voxel data) relative to glomeruli. However, our high resolution data

shows there are at least 12 Strahler orders (an additional two) between afferent

arterioles and the kidney artery. The estimate of 12 Strahler Orders between afferent

arterioles and the kidney artery can also be supported by considering the estimates

of total glomeruli number and number of terminal nodes in the 50 μm/voxel data (**See**

**Supplementary Methods §3** and **Figure S7**). Given the scale of the arterial tree

captured at our 50 μm per voxel dataset, we used this to perform further quantitative

analysis.

By plotting the number of segments within each Strahler order (**Figure 3C**) we

determine the branching ratio arterial vessels in the human kidney to be 2.921. This

value is similar to that of the human pulmonary arterial tree (3.0 [32]) and to that of

the rat kidney (2.85 [11]). To give spatial context to our data, we mapped Strahler

orders to known anatomical subdivisions of the human arterial tree, including

interlobar, arcuate and interlobular arteries. Strahler orders 7-9 ($n = 25$ segments;

mean radius = 929 ± 477 μm) mapped to the branches of the kidney artery entering

the kidney hilum. Orders 5-6 comprised interlobar arteries ($n = 219$ segments; mean

radius = 417 ± 247 μm), and orders 2-4 arcuate arteries ($n = 4841$ segments; mean

radius = 78 ± 45 μm). Interlobular arteries fell within orders 1-3 ($n = 9430$ segments;

mean radius = 55 ± 23 μm). We further plotted the cumulative volume of the kidney

vascular network is plotted in **Figure 3D,** finding that over 1/5 of the volume of the

network lies within Strahler orders 1-4, corresponding to segments from interlobular

arteries to arcuate arteries.

**Analysis of vascular network metrics in the human kidney reveals limit of**

**Murray's law and concordance with a rodent model organism**

Vascular network geometric properties, such as, diameter, length and branching

angles, are an important means of quantitatively comparison of vascular networks in

health or disease [36], [37]. To address this, we have extracted and reported the

metrics for the human kidney vasculature. We grouped our data according to Strahler

order (**Figure 4**, **Table 1**) to enable quantitative comparison to rat and other human

organ data, and also provide raw data for each segment in **Supplementary data** as

inputs for modelling applications.

As Strahler order increases, in the human kidney vascular network there is a

reduction in the ratio of vessel length:diameter (**Figure 4B**), whilst the mean radius

(**Figure 4E**) and inter-vessel distance increase (**Figure 4F**). Tortuosity does not vary

significantly with Strahler order (**Figure 4D**); with most segments having tortuosity

close to 1 implying limited deviation from a straight path. The above findings are

largely consistent with anticipated trends for healthy tissue i.e. where a vasculature

network is assumed to be a fractal structure, with branching pattern driven by

optimised delivery of blood to the whole organ. Interestingly for Strahler orders 8-6

the mean branching angle is approximately 150º, which decreases slightly to 130º for

Strahler orders 3-1 (**Figure 4C**), the latter being the predicted optimal theoretical

branching angle for vascular growth that is volume constrained [38].

**Table 1. Human kidney vascular branching metrics by Strahler**

**generation (means with standard deviation are shown).**

| Strahler order | Segments | Rad μm | Length μmx10³ | Tort. | LDR | Vol. x10⁸ μm³ | Branching Angle º | IVD μmx10³ |
|---|---|---|---|---|---|---|---|---|
| 1 | 5105 | 45±5 | 2.6±1.7 | 1.2±0.20 | 57.4±36 | 0.18±0.22 | 129±28 | 1.1±0.4 |
| 2 | 3030 | 56±15 | 1.8±1.4 | 1.1±0.16 | 32.3±26 | 0.21±0.46 | 128±29 | 0.9±0.4 |
| 3 | 1295 | 95±37 | 1.8±1.4 | 1.1±0.14 | 20.2±18 | 6.7±10.3 | 128±29 | 1.1±0.4 |
| 4 | 516 | 165±60 | 2.3±1.9 | 1.1±0.13 | 14.6±12 | 25.5±34.9 | 135±29 | 1.4±0.5 |
| 5 | 150 | 294±110 | 3.3±2.6 | 1.1±0.05 | 12.2±9.4 | 11.3±17.1 | 142±25 | 1.8±0.7 |
| 6 | 69 | 684±250 | 4.3±3.1 | 1.1±0.06 | 6.5±4.5 | 84.6±104 | 149±21 | 2.9±0.7 |
| 7 | 20 | 839±251 | 7.2±4.9 | 1.1±0.05 | 8.5±4.8 | 223±302 | 148±24 | 6.1±1.5 |
| 8 | 4 | 877±188 | 8.1±4.7 | 1.1±0.07 | 9.6±6.3 | 227±159 | 141±6 | 4.7±1.5 |
| 9 | 1 | 2923 | 669 | 1.0 | 0.2 | 180 | - | - |

**Figure 4. Metrics of the adult human kidney arterial network. A)** *Schematic diagram of how the metrics in B-E are calculated.* ***B)*** *The length:diameter ratio.* ***C)*** *The branching angle between the child and parent segments.* ***D)*** *The tortuosity of segments,* ***E)*** *their radius, and* ***F)*** *the inter-vessel distance as measured between the mid-point of each segment.*

As simulation of kidney haemodynamics has previously been performed using

micro-CT data from the rat kidney, we used our high-resolution segmentations to

align our network with those derived the previously published rat dataset [11]. In

doing so, we were able to relate normalised vessel metrics from each species at

corresponding Strahler orders. The increase in vessel radius with Strahler order

followed a similar trend between human and rat kidney (**Figure 5A**); The only

parameter that showed a considerable difference was radius, as human and rat vary

significantly different based on a fit of log(radius) to Strahler order (**Figure 5B)**,

($p < 0.0001$ Sum-of-F test $F$ (DFn, DFd) = 700.6 (2, 12)), with human vessel radii

increasing with Strahler Order more quickly than the rat. To provide physiological

relevance to this discrepancy, we compared our radius data to the Murray's Law; a

theoretical relationship between the radii of parent and child vessels derived from

considering an optimisation of energy between blood flow through a network and

diffusion into tissue with fixed metabolic demands [35], [38], [39]. The application of

Murray's Law has been previously supported by data from e.g. Nordlestten et al.

which shows a deviation from Murry's law by ~1% for the rat kidney [11].

When a best fit linear regression is applied to our data (red line **Figure 5C**),

the slope 0.63 and intercept $4 \times 10^{-6}$ values are with an $R^2 = 0.68$, applying a Extra-

sum-of-F test, $F$ (DFn, DFd) = 5474 (1, 3949) between our data and a theoretical

Murray's law we found a significant difference $P < 0.0001$ between Murray's law and

our data fit (**Figure 5C**). The deviation from Murray's law is more pronounced for

smaller vessels (**Figure 5C** inset). and is consistent with the previously calculated

branching ratios, thus inferring deviation from a symmetric network as arterial vessels

undergo hierarchal branching towards the cortex of the organ.

***Figure 5. Comparison to rodent data and Murray's theoretical law of energy balance**. **A)** Normalised radius against Strahler Order for our data and for the rat data of Nordsletten et al.* [11]*, **B)** for the same data but plotted for log(Radius), a similar pattern is seen but significant statistical difference is found between the best fit for the two datasets. **C)** Our data plotted to compare to Murray's Law, plotted showing each Strahler order in a different colour. The best fit line for the data is shown in red with the theoretical Murray's law in dashed black. The inset shows how the human data differs from Murray's law predominantly for the smaller vessels.*

**Regional heterogeneity within the kidney creates local microenvironments that enable specialised kidney functions.**

Having demonstrated deviation in Murray's law at lower Strahler orders, corresponding, we then sought to resolve regional heterogeneity in the human kidney vasculature. This regional heterogeneity is postulated to reflect different functions corresponding to the anatomical zone of the organ. For example, the kidney medulla possesses low oxygen tension, generating hypoxia that is inherent to the medulla's urinary concentration mechanisms. A longstanding hypothesis, supported by blood oxygenation level-dependent MRI studies, [40] is that vascular rarefaction in CKD results in hypoxia

within the kidney cortex, stimulating neighbouring cells into a pro-fibrotic phenotype

and manifesting in loss of organ function [3]. Although the regional heterogeneity of

vascular patterning is likely to be fundamental for local microenvironments, such as the

generation of physiological hypoxia or susceptibility to pathological hypoxia, it has not

been quantitatively explored in the human kidney vasculature.

**Table 2. Human kidney vascular branching metrics by zone**

| | Cortex | Medulla | Hilum | Inter-medullary pillars | Organ |
|---|---|---|---|---|---|
| **Volume of tissue x $10^{13}$ $\mu m^3$ (% of total)** | 8.70 (63.7%) | 3.21 (23.5%) | 1.18 (8.66%) | 0.57 (4.14%) | 13.7 (100%) |
| **Number of segments* (% of total)** | 6141 (60.27%) | 554 (5.4%) | 151 (1.5%) | 727 (7.1%) | 10190 (100%) |
| **Mean segment length, $\mu m$ ± STD** | 1999±1374 | 1493±1113 | 3993±3568 | 1720±1386 | 2260±1720 |
| **Mean segment radius, $\mu m$ ± STD** | 48±12.6 | 95±49 | 496±335 | 136±80 | 71±87 |
| **Mean inter-vessel distances, $x10^3$ $\mu m$ ± STD** | 1.10±0.677 | 1.55±0.881 | 1.55±1.312 | 0.664±0.543 | 1.2±0.833 |
| **Mean segment volume, $x10^8$ $\mu m^3$ ± STD** | 0.148±0.116 | 0.623±1.12 | 69.0±147 | 1.73±4.48 | 1.65± 20.6 |
| **Mean segment tortuosity ± STD** | 1.14±0.17 | 1.08±0.13 | 1.08±0.1 | 1.1±0.12 | 1.15±0.18 |
| *segments that crossed over two regions were excluded. | | | | | |

**Figure 6. Analysis of zonal heterogeneity in vascular branching metrics within the human kidney. (Ai)** *3D reconstruction vasculature colour according to anatomical compartment within the human kidney cortex (green), medulla (yellow) and hilum (pink), inter-medullar pillars (dark blue). (**Aii**) Showing the 3D surface masks of the same regions. (**Aiii**) Inter-vessel distances are plotted against the total number of vessel voxels for each kidney compartment. (**Bi**) Visual heatmap of inter-vessel distance for the entire human kidney, where pink represents the largest inter-vessel distance (> 4.5 mm) and white (0 mm) the smallest. (**Bii**) A digital zoomed region within cortex and medulla. (**Biii**) The 2D slice of the associated HiP-CT raw image with the compartments overlaid.*

467
468
469
470

Leveraging the contrast-free approach of HiP-CT imaging, we were able to

segment the kidney into known anatomical compartments including hilum, medulla,

intramedullary kidney columns and cortex (**Figure 6A**). The total tissue volume of

each zone in addition to the number of vessels, length, radius and volume of

segmented vessels within each zone were quantified (**Table. 2**). Most of the tissue

volume of the human kidney was measured to be occupied by the cortex (63.7%) as

compared with the medulla (23.5%) or hilum (8.7%) or intermedullary pillars (4.1%).

The number of segments for the vascular network within each compartment followed

this trend. As a proxy for kidney tissue oxygenation, we quantified (**Figure 6Ai-Aiii**)

and mapped (**Figure 6Bi-Biii**) the inter-vessel distance, reflecting the extravascular

distance across which oxygen and solutes diffuse, compartmentalised by hilum,

medulla, cortex, and inter-medullary pillars. Mean inter-vessel distances for these

compartments **(Table. 2)** demonstrates the medulla having the highest inter-vessel

distance which follows the anticipated distributions of hypoxia. Notably, there were

large portions within the medulla where inter-vessel distance was > 4.5 mm **(Figure**

**6Bi)**; in line with the known hypoxic characteristic of the medulla. Whilst the cortex

has one of smallest inter-vessel distance, it has a large standard deviation and the

heatmap in **Figure 6Bi and 6Bii** shows small areas with inter-vessel distance >

4.5mm are also found, predominantly towards the kidney capsule.

**DISCUSSION**

Owing to the limited volume of tissue imageable using pre-clinical modalities such as

micro-CT and lightsheet microscopy, and insufficient resolution of technologies

routinely used in clinical practice such a CT and MRI, it had previously been impossible

to capture the entire vascular network of the intact adult human kidney. Using

synchrotron-based HiP-CT we were able to segment and quantify the human kidney

arterial network from kidney artery to interlobular arteries without the need for

exogenous contrast agents. With this method, we demonstrated that vessels mapping

from interlobar to interlobular arteries occupy approximately 1/5[th] of the vascular

volume of the human kidney. By imaging regions of interest in the intact kidney at

higher resolution and aligning with low resolution data, we further demonstrate that,

akin to rat [34] and varying from the traditional hierarchy of the kidney vasculature

observed in textbooks, that the glomeruli in humans can originate from non-terminal

arterioles. Although there existed similarity in topology between human and rat kidney,

there was a significant difference in the change in radius with Strahler order between

organisms. As vessels branch towards the cortex and radius decreases in the human

kidney vasculature, they also do not adhere to Murray's theoretical law of energy
balance. Finally, we demonstrate vascular volume fractions and inter vessel distances
vary between anatomical zones of the kidney, corresponding to regionally specialised
functions and known physiological gradients in local oxygen tension.

The human kidney vasculature is exquisitely specialised to meet the
physiological demands of the kidney. Underpinning this specialisation is the cellular
and molecular heterogeneity of endothelial beds with the renal vasculature [41] of
which we are gaining an increasing understanding due to the advent of improved
techniques such as cellular transcriptomics. The rapid advances in our understanding
of cellular and molecular heterogeneity of the kidney vasculature has not been matched
by structural insights, likely as a consequence of limitations in imaging technologies.
We have overcome these limitations using HiP-CT, and capture the 3D vascular
architecture of an entire human kidney at twenty-fold greater resolution than
conventional hospital CT scanners (400 μm per voxel), instead comparable to that of
light microscopy (1-8 μm per voxel) yet on a volume many orders of magnitude larger
than that of a kidney punch biopsy. The balance between imaging volume and
resolution afforded by HiP-CT thus bridges the scale between local cellular architecture
and global tissue structure, providing quantitative vascular branching metrics from an
intact human organ for the first time. Testament to this balance, we report up to 10
Strahler orders or 27 topological generations for the human kidney vasculature. These
exceed previous *in vivo MRI* studies, which report up to six topological orders [23]
Despite being taken from the lowest resolution HiP-CT scan, these metrics also
exceeds studies on cadaveric cast and dye injections which report up to arterial
branches corresponding to Strahler orders 7-9) [22], [42]. We demonstrate
concordance in topological layout between the human kidney vasculature and that of
the rat, the latter of which has been key for inputs to generate  biophysical models of
kidney haemodynamics [5], [11] However, we demonstrate deviations in the
progressive decrease in magnitude of vessel radii between human and rat kidney
vasculature, and non-adherence to a symmetrical network that would be consistent
with Murray's law. This discrepancy could potentially influence the prior simulations of
haemodynamics, oxygenation or drug delivery [8], [43]; and generation of synthetic
vessel trees for *in silico* experiments [44], [45]. This divergence between theoretical
prediction and our human data may be accounted for by regional heterogeneity in
architecture, corroborated by our segmentation of hilar, medullary, intra-medullary and
cortical zones of HiP-CT images from the same kidney. The increased inter-vessel
distance observed within the medullary compared to other kidney anatomical zones

corresponds to the known decreased oxygen tension within this region of the kidney.

Thus, structural heterogeneity could impart varying metabolic demands and generate

local microenvironments within the kidney [46], [47], which may underpin the unique

cellular and molecular adaptations of specialised endothelia across the kidney vascular

network [41].

The quantitative analysis pipeline performed in this paper serves multiple

purposes. Firstly, it surmounts one of the limitations of HiP-CT, in the size of the

datasets generated. Whereas the whole kidney dataset amounts to ~500GB,

necessitating considerable computational power and data storage facilities,

segmentation followed by skeletonization allows the whole kidney vasculature dataset

to be represented in a spatial graph comprising only KB of data. The spatial graph,

which is provided as **Supplementary Data**, is readily quantifiable. Whereas prior

simulations of kidney haemodynamics and perfusion have relied on seminal micro-CT

studies performed in rat, we provide, for the first time, a complete map of the kidney

arterial network in its entirety. The segmentation is accurate, with 97% of vessels of

<50 µm radius captured across the intact human kidney, and thus provides vital inputs

for future biophysical modelling frameworks of kidney physiology. It also serves as a

reference dataset to study kidney diseases, in which vascular rarefaction is a

pathophysiological hallmark[48]. The pipeline could be used to generate vascular maps

from multiple kidneys or other human organs, potentially giving rise to spatial 'atlases'

of human organ vasculature across healthy and pathological contexts. Beyond these,

our openly available dataset has immediate practical applications, such as providing

inputs for bioprinting for tissue engineering of artificial kidneys [49] or planning surgical

resection of kidney tumours whilst preserving kidney function. [50] These datasets can

also be used as a tool for medical education and training, as well as for the creation

and advancement of surgical methods.

There are several limitations of this work, including the low throughput of the

segmentation of this type of data. Here we present the complete analysis from a single

kidney, as a framework for future studies to study further kidneys in health and disease,

or other intact human organs. The accuracy of the segmentation, however, lays a

foundation for tools such as machine learning methods for automated segmentation of

blood vasculature from imaging data,[17], [51], [52]. Although the resolution of the

organ-wide HiP-CT scan far exceeds conventional *in vivo* and *ex vivo* imaging

measures, it still cannot achieve arteriole resolution across the whole organ.

Nevertheless, improvements of the ESRF beamline (BM18) have already been

developed, and will extend the resolution limit for whole organs down to 8um, whilst

increasing the speed of scanning. Although the physical access to HiP-CT via the

synchrotron is limited, we have released all our data thus far through the Human Organ

Atlas portal (https://human-organ-atlas.esrf.eu/), for open-access download and use of

the data for biomedical researchers.

In summary, we have achieved quantitative mapping of the entire arterial

network of an intact human kidney for the first time: a vital step towards understanding

how physical properties of the kidney vasculature relate to cellular and molecular

heterogeneity, whilst generating key inputs for future biophysical modelling of kidney

vascular physiology. Ultimately, we envisage that mapping of microstructural detail will

become routine at the scale of the whole kidney, providing a means to link cellular

events with organ physiology and pathology.

**FUNDING**

This [publication, dataset, software etc.] has been made possible in part by grants

DAF2020-225394 and 2022-316777 (DOI /10.37921/331542rbsqvn) from the Chan

Zuckerberg Initiative DAF, an advised fund of Silicon Valley Community Foundation,

and grant CZIF2021-006424 from the Chan Zuckerberg Initiative Foundation, the

MRC (MR/R025673/1), the RAEng (CiET1819-10), and ESRF beamtimes (md1252 &

md1290). DALs laboratory is supported by a Wellcome Trust Investigator Award

(220895/Z/20/Z) and the NIHR Biomedical Research Centre at Great Ormond Street

Hospital for Children NHS Foundation Trust and University College London.

**ACKNOWLEDGEMENTS**

**AUTHOR CONTRIBUTIONS**

P.D.L, R.J.S., and, P.T. conceptualized the project and designed experiments; P.T.

designed and built instrumentation for HiP-CT imaging; P.T. J.B and C.L.W. designed

and implemented tomographic reconstruction areas; S.R. , J.B, S.N and C.L.W.

designed, managed and performed image analysis; S.R. and C.L.W. modelled,

quantified and provided results; S.R., C.L.W., R.J.S., D.A.L., D.J.J., and P.D.L,

provided results interpretation and discussion; S.R., C.L.W., D.J.J., and D.A. L. wrote

the paper; D.J.J. and D.A.L. provided medical interpretation for kidney analysis. All

authors assisted in reviewing and revising the manuscript.

**DATA SHARING STATEMENT**

The image data that form the basis of the study findings are freely available at the

ESRF data repository (https://human-organ-atlas.esrf.eu). Additionally, the spatial

graph data of the kidney arterial network, along with the computed morphological

parameters, can be accessed in Supplementary data which will be released following

peer review.

## REFERENCES

[1]   G. Molema and W. C. Aird, 'Vascular heterogeneity in the kidney', in *Seminars in nephrology*, Elsevier, 2012, pp. 145–155.

[2]   S. J. Dumas *et al.*, 'Phenotypic diversity and metabolic specialization of renal endothelial cells', *Nat Rev Nephrol*, vol. 17, no. 7, pp. 441–464, 2021.

[3]   D. A. Long, J. T. Norman, and L. G. Fine, 'Restoring the renal microvasculature to treat chronic kidney disease', *Nat Rev Nephrol*, vol. 8, no. 4, pp. 244–250, 2012.

[4]   L. F. M. Cury, G. D. Maso Talou, M. Younes-Ibrahim, and P. J. Blanco, 'Parallel generation of extensive vascular networks with application to an archetypal human kidney model', *R Soc Open Sci*, vol. 8, no. 12, Dec. 2021, doi: 10.1098/rsos.210973.

[5]   D. J. Marsh, D. D. Postnov, O. V Sosnovtseva, and N.-H. Holstein-Rathlou, 'The nephron-arterial network and its interactions', *American Journal of Physiology-Renal Physiology*, vol. 316, no. 5, pp. F769–F784, 2019.

[6]   A. d'Esposito *et al.*, 'Computational fluid dynamics with imaging of cleared tissue and of in vivo perfusion predicts drug uptake and treatment responses in tumours', *Nat Biomed Eng*, vol. 2, no. 10, pp. 773–787, 2018, doi: 10.1038/s41551-018-0306-y.

[7]   C. L. Walsh, M. Berg, H. West, N. A. Holroyd, S. Walker-Samuel, and R. J. Shipley, 'Reconstructing microvascular network skeletons from 3D images: what is the ground truth?', *Comput Biol Med*, vol. 171, p. 108140, 2024.

[8]   P. Xu *et al.*, 'A hybrid approach to full-scale reconstruction of renal arterial network', *Sci Rep*, vol. 13, no. 1, p. 7569, 2023.

[9]   D. J. Marsh, D. D. Postnov, D. J. Rowland, A. S. Wexler, O. V Sosnovtseva, and N.-H. Holstein-Rathlou, 'Architecture of the rat nephron-arterial network: analysis with micro-computed tomography', *American Journal of Physiology-Renal Physiology*, vol. 313, no. 2, pp. F351–F360, 2017.

[10]  D. S. Perrien *et al.*, 'Novel methods for microCT-based analyses of vasculature in the renal cortex reveal a loss of perfusable arterioles and glomeruli in eNOS-/-mice', *BMC Nephrol*, vol. 17, no. 1, pp. 1–10, 2016.

[11]  D. A. Nordsletten, S. Blackett, M. D. Bentley, E. L. Ritman, and N. P. Smith, 'Structural morphology of renal vasculature', *Am J Physiol Heart Circ Physiol*, vol. 291, no. 1, 2006, doi: 10.1152/ajpheart.00814.2005.

[12]  N. Parvin, J. R. Charlton, E. J. Baldelomar, J. J. Derakhshan, and K. M. Bennett, 'Mapping vascular and glomerular pathology in a rabbit model of neonatal acute kidney injury using MRI', *Anat Rec*, vol. 303, no. 10, pp. 2716–2728, 2020.

[13]  J. Foiret, H. Zhang, T. Ilovitsh, L. Mahakian, S. Tam, and K. W. Ferrara, 'Ultrasound localization microscopy to image and assess microvasculature in a rat kidney', *Sci Rep*, vol. 7, no. 1, p. 13662, 2017.

[14]  J. Huang *et al.*, 'A cationic near infrared fluorescent agent and ethyl-cinnamate tissue clearing protocol for vascular staining and imaging', *Sci Rep*, vol. 9, no. 1, p. 521, 2019.

[15]  A. Klingberg *et al.*, 'Fully automated evaluation of total glomerular number and capillary tuft size in nephritic kidneys using lightsheet microscopy', *Journal of the American Society of Nephrology*, vol. 28, no. 2, pp. 452–459, 2017.

[16]  O. Ogunlade *et al.*, 'In vivo three-dimensional photoacoustic imaging of the renal vasculature in preclinical rodent models', *American Journal of Physiology-Renal Physiology*, vol. 314, no. 6, pp. F1145–F1153, 2018.

[17]  W. Zheng *et al.*, 'Deep Learning Enhanced Volumetric Photoacoustic Imaging of Vasculature in Human', *Advanced Science*, vol. 10, no. 29, p. 2301277, Oct. 2023, doi: https://doi.org/10.1002/advs.202301277.

| 670 | [18] | S. Balıkçı Dorotea, T. Banzato, L. Bellini, B. Contiero, and A. Zotti, 'Kidney |
| 671 | | Measures in the Domestic Rat: A Radiographic Study and a Comparison to |
| 672 | | Ultrasonographic Reference Values', *J Exot Pet Med*, vol. 25, no. 2, pp. 157–162, |
| 673 | | 2016, doi: https://doi.org/10.1053/j.jepm.2016.03.011. |
| 674 | [19] | M. J. Musa and A. Abukonna, 'Sonographic measurement of renal size in normal high |
| 675 | | altitude populations', *J Radiat Res Appl Sci*, vol. 10, no. 3, pp. 178–182, 2017, doi: |
| 676 | | https://doi.org/10.1016/j.jrras.2017.04.004. |
| 677 | [20] | O. K. Zenin, E. S. Kafarov, O. A. Beshulya, L. A. Udochkina, and H. M. Bataev, |
| 678 | | 'Quantitative Anatomy of the Intrainganic Arterial Kidney', in *International* |
| 679 | | *Conference on Health and Well-Being in Modern Society (ICHW 2019)*, Atlantis Press, |
| 680 | | 2019, pp. 129–132. |
| 681 | [21] | S. Zhao *et al.*, 'Cellular and molecular probing of intact human organs', *Cell*, vol. 180, |
| 682 | | no. 4, pp. 796–812, 2020. |
| 683 | [22] | I. U. Vagabov, E. S. Kafarov, O. K. Zenin, T. S. Dokaeva, and K. M. Bataev, |
| 684 | | 'Segmental Arteries as Sources of Formation of Arterial Segments of Human Kidney', |
| 685 | | in *The International Conference "Health and wellbeing in modern society"(ICHW* |
| 686 | | *2020)*, Atlantis Press, 2020, pp. 341–346. |
| 687 | [23] | L. Timms *et al.*, 'Ferumoxytol-enhanced ultrashort TE MRA and quantitative |
| 688 | | morphometry of the human kidney vasculature', *Abdominal Radiology*, vol. 46, pp. |
| 689 | | 3288–3300, 2021. |
| 690 | [24] | J. R. Charlton *et al.*, 'Image analysis techniques to map pyramids, pyramid structure, |
| 691 | | glomerular distribution, and pathology in the intact human kidney from 3-D MRI', |
| 692 | | *American Journal of Physiology-Renal Physiology*, vol. 321, no. 3, pp. F293–F304, |
| 693 | | 2021. |
| 694 | [25] | D. J. Jafree *et al.*, 'Three-dimensional imaging and single-cell transcriptomics of the |
| 695 | | human kidney implicate perturbation of lymphatics in alloimmunity', *bioRxiv*, p. |
| 696 | | 2022.10.28.514222, Jan. 2022, doi: 10.1101/2022.10.28.514222. |
| 697 | [26] | C. L. Walsh *et al.*, 'Imaging intact human organs with local resolution of cellular |
| 698 | | structures using hierarchical phase-contrast tomography', *Nat Methods*, vol. 18, no. 12, |
| 699 | | pp. 1532–1541, 2021. |
| 700 | [27] | J. Brunet *et al.*, 'Preparation of large biological samples for high-resolution, |
| 701 | | hierarchical, multi-modal imaging', *bioRxiv*, 2022. |
| 702 | [28] | A. Mirone, E. Brun, E. Gouillart, P. Tafforeau, and J. Kieffer, 'The PyHST2 hybrid |
| 703 | | distributed code for high speed tomographic reconstruction with iterative |
| 704 | | reconstruction and a priori knowledge capabilities', *Nucl Instrum Methods Phys Res B*, |
| 705 | | vol. 324, pp. 41–48, 2014. |
| 706 | [29] | J. C. Paetzold *et al.*, 'clDice—A novel connectivity-preserving loss function for vessel |
| 707 | | segmentation', in *Medical Imaging Meets NeurIPS 2019 Workshop*, 2019. |
| 708 | [30] | G. S. Kassab, K. Imoto, F. C. White, C. A. Rider, Y.-C. Fung, and C. M. Bloor, |
| 709 | | 'Coronary arterial tree remodeling in right ventricular hypertrophy', *American Journal* |
| 710 | | *of Physiology-Heart and Circulatory Physiology*, vol. 265, no. 1, pp. H366–H375, |
| 711 | | 1993. |
| 712 | [31] | A. N. Strahler, 'Quantitative analysis of watershed geomorphology', *Eos, Transactions* |
| 713 | | *American Geophysical Union*, vol. 38, no. 6, pp. 913–920, Dec. 1957, doi: |
| 714 | | https://doi.org/10.1029/TR038i006p00913. |
| 715 | [32] | K. Horsfield, 'Morphometry of the small pulmonary arteries in man.', *Circ Res*, vol. |
| 716 | | 42, no. 5, pp. 593–597, 1978. |
| 717 | [33] | A. R. Pries and T. W. Secomb, 'Blood Flow in Microvascular Networks', |
| 718 | | *Microcirculation*, pp. 3–36, 2008, doi: 10.1016/B978-0-12-374530-9.00001-2. |

[34] D. J. Marsh, D. D. Postnov, D. J. Rowland, A. S. Wexler, O. V Sosnovtseva, and N.-
H. Holstein-Rathlou, 'Architecture of the rat nephron-arterial network: analysis with
micro-computed tomography', *Am J Physiol Renal Physiol*, vol. 313, no. 2, pp. F351–
F360, Aug. 2017, doi: 10.1152/ajprenal.00092.2017.
[35] D. D. Postnov *et al.*, 'Modeling of Kidney Hemodynamics: Probability-Based
Topology of an Arterial Network', *PLoS Comput Biol*, vol. 12, no. 7, pp. e1004922–
e1004922, Jul. 2016, doi: 10.1371/journal.pcbi.1004922.
[36] C. O'Connor, E. Brady, Y. Zheng, E. Moore, and K. R. Stevens, 'Engineering the
multiscale complexity of vascular networks', *Nat Rev Mater*, vol. 7, no. 9, pp. 702–
716, 2022.
[37] J. Ehling *et al.*, 'Quantitative micro-computed tomography imaging of vascular
dysfunction in progressive kidney diseases', *Journal of the American Society of
Nephrology*, vol. 27, no. 2, pp. 520–532, 2016.
[38] E. Tekin, D. Hunt, M. G. Newberry, and V. M. Savage, 'Do Vascular Networks
Branch Optimally or Randomly across Spatial Scales?', *PLoS Comput Biol*, vol. 12,
no. 11, pp. e1005223–e1005223, Nov. 2016, doi: 10.1371/journal.pcbi.1005223.
[39] W. Deng and K. Tsubota, 'Numerical simulation of the vascular structure dependence
of blood flow in the kidney', *Med Eng Phys*, vol. 104, p. 103809, 2022.
[40] M. Pruijm *et al.*, 'Renal blood oxygenation level-dependent magnetic resonance
imaging to measure renal tissue oxygenation: a statement paper and systematic
review', *Nephrology Dialysis Transplantation*, vol. 33, no. suppl_2, pp. ii22–ii28,
2018.
[41] S. J. Dumas *et al.*, 'Phenotypic diversity and metabolic specialization of renal
endothelial cells', *Nat Rev Nephrol*, vol. 17, no. 7, pp. 441–464, 2021, doi:
10.1038/s41581-021-00411-9.
[42] M. M. Shoja *et al.*, 'Peri-hilar branching patterns and morphologies of the renal artery:
a review and anatomical study', *Surgical and Radiologic Anatomy*, vol. 30, pp. 375–
382, 2008.
[43] A. d'Esposito *et al.*, 'Computational fluid dynamics with imaging of cleared tissue and
of in vivo perfusion predicts drug uptake and treatment responses in tumours', *Nat
Biomed Eng*, vol. 2, no. 10, pp. 773–787, 2018.
[44] L. F. M. Cury, G. D. Maso Talou, M. Younes-Ibrahim, and P. J. Blanco, 'Parallel
generation of extensive vascular networks with application to an archetypal human
kidney model', *R Soc Open Sci*, vol. 8, no. 12, p. 210973, 2021.
[45] E. Brown *et al.*, 'Physics-informed deep generative learning for quantitative
assessment of the retina', *bioRxiv*, p. 2023.07.10.548427, Jan. 2023, doi:
10.1101/2023.07.10.548427.
[46] A. B. Brummer *et al.*, 'Branching principles of animal and plant networks identified by
combining extensive data, machine learning and modelling', *J R Soc Interface*, vol. 18,
no. 174, p. 20200624, 2021.
[47] E. Tekin, D. Hunt, M. G. Newberry, and V. M. Savage, 'Do vascular networks branch
optimally or randomly across spatial scales?', *PLoS Comput Biol*, vol. 12, no. 11, p.
e1005223, 2016.
[48] D. A. Long, J. T. Norman, and L. G. Fine, 'Restoring the renal microvasculature to
treat chronic kidney disease', *Nat Rev Nephrol*, vol. 8, no. 4, pp. 244–250, 2012, doi:
10.1038/nrneph.2011.219.
[49] C. G. Lebedenko and I. A. Banerjee, 'Enhancing Kidney vasculature in tissue
engineering—Current trends and approaches: A Review', *Biomimetics*, vol. 6, no. 2, p.
40, 2021.

[50]   J. J. Hsieh *et al.*, 'Renal cell carcinoma', *Nat Rev Dis Primers*, vol. 3, no. 1, pp. 1–19,
769           2017.
770    [51]   C. Kirst *et al.*, 'Mapping the fine-scale organization and plasticity of the brain
771           vasculature', *Cell*, vol. 180, no. 4, pp. 780–795, 2020.
772    [52]   M. I. Todorov *et al.*, 'Machine learning analysis of whole mouse brain vasculature',
773           *Nat Methods*, vol. 17, no. 4, pp. 442–449, 2020.
774