Fungal melanin suppresses airway epithelial chemokine secretion through blockade of calcium fluxing

Jennifer L. Reedy[1,2,#], Kirstine Nolling Jensen[1,2,#], Arianne J. Crossen[1], Kyle J. Basham[1], Rebecca A. Ward[1], Christopher M. Reardon[1], Hannah Brown Harding[1,2], Olivia W. Hepworth[1,2], Patricia Simaku[1], Geneva N. Kwaku[1], Kazuya Tone[3,4], Janet A. Willment[3,5], Delyth M. Reid[3], Mark H.T. Stappers[3,5], Gordon D. Brown[3,5], Jayaraj Rajagopal[6,7,8,9], Jatin M. Vyas[1,2,*]

[1] Department of Medicine, Division of Infectious Diseases, Massachusetts General Hospital, Boston, MA, 02114, USA

[2] Harvard Medical School, Boston, MA, 02115, USA

[3] Aberdeen Fungal Group, University of Aberdeen, Institute of Medical Sciences, Foresterhill, Aberdeen, United Kingdom

[4] Division of Respiratory Diseases, Department of Internal Medicine, The Jikei University School of Medicine, Tokyo, Japan

[5] MRC Centre for Medical Mycology, University of Exeter, Exeter, United Kingdom

[6] Center for Regenerative Medicine, Massachusetts General Hospital, Boston, MA, 02114, USA

[7] Division of Pulmonary and Critical Care Medicine, Department of Medicine, Massachusetts General Hospital, Boston, MA, 02114, USA

[8] Harvard Stem Cell Institute, Cambridge, MA, 02138, USA

23   [9] Klarman Cell Observatory, Broad Institute of Massachusetts Institute of Technology

24   and Harvard, Cambridge, MA, 02142, USA

25   *Correspondence: jvyas@mgh.harvard.edu

26   [#]Co-first authors

Summary

Respiratory infections caused by the human fungal pathogen *Aspergillus fumigatus* are a major

cause of mortality for immunocompromised patients. Exposure to these pathogens occurs

through inhalation, although the role of the respiratory epithelium in disease pathogenesis has

not been fully defined. Employing a primary human airway epithelial model, we demonstrate that

fungal melanins potently block the post-translational secretion of the chemokines CXCL1 and

CXCL8 independent of transcription or the requirement of melanin to be phagocytosed, leading

to a significant reduction in neutrophil recruitment to the apical airway both *in vitro* and *in vivo*.

Aspergillus-derived melanin, a major constituent of the fungal cell wall, dampened airway

epithelial chemokine secretion in response to fungi, bacteria, and exogenous cytokines.

Furthermore, melanin muted pathogen-mediated calcium fluxing and hindered actin filamentation.

Taken together, our results reveal a critical role for melanin interaction with airway epithelium in

shaping the host response to fungal and bacterial pathogens.

Keywords: airway epithelium, *Aspergillus*, melanin, innate immunity, host-pathogen interaction,

fungal immunology

Introduction

Aspergillus fumigatus is the most prominent respiratory fungal pathogen and causes a spectrum of clinical manifestations ranging from allergic disease to severe invasive infections. Despite the ubiquity of this fungal pathogen in the environment, only ~10% of immunocompromised patients (*e.g.,* neutropenic or allogenic bone marrow transplant recipients) develop invasive aspergillosis (IA), indicating that other factors play a significant role in determining the true risk for this infection[1]. In addition to infection associated with the immunocompromised state, there is an increased risk of IA following pulmonary viral infections (*e.g.,* influenza, SARS-CoV-2) and in those with underlying lung disease (*e.g.,* asthma, cystic fibrosis [CF], chronic obstructive pulmonary disease [COPD])[2-5]. The mortality from *Aspergillus*-related pulmonary disease remains unacceptably high (>50%), coinciding with elevated rates of *Aspergillus* multidrug resistance[6-8]. Despite this, the mechanistic interactions governing the invasion of *Aspergillus* at its first point of contact with the host, the airway epithelium, remains poorly understood.

In immune cells, activation of pattern recognition receptors (PRRs), such as the C-type lectin receptor (CLRs), Toll-like receptors (TLRs), NOD-like receptors (NLRs), and Rig-I-like receptors, mediate fungal recognition of cell wall components and subsequent host responses[9]. The cell wall of *Aspergillus* is primarily composed of polysaccharides, including galactosaminogalactan, galactomannan, β-1,3 glucan, β-1,4 glucan, and chitin[10]. Most humans are exposed to *Aspergillus* through the inhalation of conidia, the reproductive propagules of the fungus, which range from 2-3 μm in size and are easily aerosolized in the environment[11,12]. In addition to the typical cell wall components, *Aspergillus* conidia are surrounded by a hydrophobic rodlet and 1,8-dihydroxy naphthalene (DHN) melanin layers[10,12]. Once inhaled by the host, conidia swell and shed these outer layers, exposing the carbohydrate matrix of the cell wall. In immune cells, fungal melanin possesses the remarkable capacity to absorb reactive oxygen species (ROS) and blunt pro-inflammatory cascades, serving as a key virulence factor for many

pathogens, including *Aspergillus*[13-19]. Amelanotic strains of clinically relevant fungal pathogens

lose their capacity to mount successful infections in mammalian host model systems, indicating

that melanin plays a key and non-redundant role in virulence. Indeed, the presence of melanized

Aspergillus leads to blockage of phagosome biogenesis[20] and removal of the melanin in the

phagosome is required to reprogram macrophage metabolism, promoting an antifungal

response[21].

Clinical data and current literature indicate that several immune system components are

critical for the recognition and swift clearance of fungal pathogens. Still, the rules governing the

inflammatory responses to fungi in the lungs are poorly understood. Past studies have focused

on innate and adaptive immune cells but have often overlooked a key player in respiratory

infections: the airway epithelium - the primary point of contact for inhaled conidia[22-25]. The

conducting airways are sites of *Aspergillus* colonization and invasion[22-25]. Fortunately, *in vitro*

differentiation of primary airway basal cells recapitulates the true diversity of conducting airway

epithelium with all of the relevant common (basal, ciliated, club, goblet) and rare (ionocytes, tuft,

neuroendocrine) cell types in pseudostratified layers, enabling mechanistic studies *ex vivo*[26].

Additionally, these cells form polarized barriers and enable the investigation of basolateral and

apical epithelial responses when grown at air-liquid interface (ALI)[27,28]. While the primary human

airway epithelial cell (hAEC) ALI model may not fully recapitulate the *in vivo* environment (*e.g.,*

lacks resident immune cells, endothelial cells, fibroblasts), this model offers the advantages of a

highly pliable system to conduct mechanistic studies without compromising relevant human

physiology[28-30] and allow us to pinpoint mechanisms specific to the interaction of the epithelium

and inhaled conidia where the microbe first encounters host defense. In addition to utilizing this

system, we also use the airway epithelial cell line H292, a mucoepitheliod non-small cell lung

carcinoma cell line, which has previously been used extensively in modeling airway epithelial

responses to bacterial pathogens. While this cell line does not capture the cellular diversity of the

airway epithelium, it is well-suited to mechanistic studies due to cellular uniformity. We previously

demonstrated a role for melanin in immune response to *Aspergillus fumigatus* orchestrated by

airway epithelium using both primary hAEC and H292 epithelial models[28]. We observed that

airway epithelium infected with *A. fumigatus* lacking DHN melanin promoted greater

transepithelial migration of neutrophils than wildtype conidia. The mechanism by which melanin

modulates epithelial-mediated neutrophil transmigration remains unknown.

In the present study, we leveraged primary hAEC to demonstrate that purified fungal

melanin at the apical surface of epithelial cells blocked migration of neutrophils and blunted the

apical secretion of chemoattractant chemokines. We determined that DHN melanin suppression

of neutrophil transmigration and chemokines occured in response to both fungal and bacterial

pathogens. Furthermore, L-DOPA melanin from *Cryptococcus neoformans* and other

commercially available synthetic melanins inhibited CXCL8 (also known as IL-8) secretion,

showing these effects were not unique to DHN melanin, but rather is a preserved property of

fungal melanins. We determined that fungal melanin abolished the apical-basolateral CXCL8

gradient as a mechanism to block neutrophil recruitment. We demonstrated that melanin

suppressed secretion, but not translation or transcription, of CXCL8 in airway epithelium in a

phagocytosis-independent manner. Lastly, we unveiled melanin exerts effects through dampened

calcium fluxing and disrupted actin filamentation in airway epithelium. Together, these data

indicated that fungal melanin inhibits epithelial-mediated inflammation through blocking calcium

signaling and the secretion of critical pro-inflammatory chemokines.

Results

Aspergillus melanin inhibits transepithelial migration of primary neutrophils

We previously demonstrated that hAEC induce neutrophil transmigration upon stimulation with *A.*

fumigatus wildtype conidia[28]. However, induction of neutrophil transmigration required at least 4h

of infection with *A. fumigatus* in contrast to the bacterial pathogen, *Pseudomonas aeruginosa*,

which induced robust neutrophil transmigration within 1h[28]. Furthermore, *Aspergillus* conidia that

lacked the ability to produce DHN melanin *(ΔpksP)* induced neutrophil transmigration within 1h of

infection similarly to *P. aeruginosa*, suggesting that melanin was blocking hAEC responses to

resting conidia[28]. Although prior studies identified MelLec as a receptor for DHN melanin in

endothelial cells and myeloid cells[31-33], primary hAECs and H292 cell line lacked this receptor

(**Supplementary Figure 1**), suggesting that the effects of fungal melanin is exerted through

alternative mechanisms. Thus, we hypothesized that melanin could act via two mechanisms to

block neutrophil migration: melanin could actively downregulate pro-inflammatory responses or

serve as a passive barrier to prevent host access to immunostimulatory epitopes on the fungal

cell wall. To test these hypotheses, we performed mixing studies with *A. fumigatus* melanin ghosts

and amelanotic *ΔpksP* conidia. Generation of melanin ghosts is a well-described technique in

which melanized conidia are biochemically and enzymatically degraded to strip all cellular

components except a rigid shell of purified melanin resembling the original size and shape of the

conidia[20,34-38]. To ascertain whether melanin coating of conidia was required for its inhibitory

function, we stimulated primary hAEC with *ΔpksP* conidia alone or in combination with melanin

ghosts at the apical side at a ratio of 5:1 melanin ghosts to *ΔpksP* conidia and then measured

neutrophil recruitment across the epithelium (basolateral→apical) (**Figure 1a**). Stimulation of

hAEC with *ΔpksP* conidia mixed with melanin ghosts dampened neutrophil transmigration when

compared to *ΔpksP* alone (**Figure 1b**). As expected, melanin ghosts alone did not stimulate

neutrophil transmigration.

We next examined if this response seen in primary hAEC extended to the human lung

epithelial cell line, H292, which is routinely used to dissect airway responsiveness to bacterial

pathogens[39-41]. While these cells do not recapitulate the cellular diversity of the primary airway

epithelium, they are well suited to mechanistic studies given their homogeneity, thus we wished

to determine if our phenotype could translate to this model system. We observed that infection

with *ΔpksP* conidia alone induced more neutrophil recruitment across H292 monolayers

compared with *ΔpksP* conidia in the presence of melanin ghosts (**Figure 1c**) recapitulating the

primary hAECs findings. Since the melanin in these experiments was applied as discrete particles

unable to coat the surface of the *ΔpksP* conidia, these results indicate a model where melanin

actively downregulates inflammatory responses rather than acting as a passive barrier on the

surface of the conidia.

Aspergillus *melanin inhibits secretion of airway epithelial-derived CXCL8 and*

CXCL1

To understand the mechanism of how neutrophil transmigration was silenced, we performed a

Luminex assay using primary human airway epithelium derived from three healthy human donors.

Primary hAEC was differentiated at ALI for 16 days prior to stimulation with either media alone

(HBSS), *P. aeruginosa* (positive control), three wildtype strains of *A. fumigatus* (CEA10, Af293,

B5233), or the amelanotic *A. fumigatus ΔpksP* strain (derived from B5233). CXCL8 and CXCL1

(also known as GROα) are two neutrophil chemoattractants produced by the airway epithelium

and were the most abundant chemokines specifically induced on the apical side by the mutant

ΔpksP A. fumigatus strain, but not by wildtype strains (**Figure 2a-b, Supplementary Figure 2**).

We confirmed these differences in cytokine expression in H292 cells using both the *ΔpksP* and

ΔpksP complemented strains (**Figure 2c-d**) to validate the use of these cells for mechanistic

studies.

To determine if melanin acted through a blockade of epithelial chemokine secretion, we

performed mixing studies in which we stimulated with a fixed concentration of *ΔpksP* conidia

combined with graded amounts of melanin ghosts. The mixing assay revealed that the addition

of melanin ghosts suppressed both CXCL8 and CXCL1 chemokine apical secretion in a dose-

dependent manner (**Figure 2e-f**). Melanin ghosts alone applied at the highest concentration did

not induce chemokine secretion. To confirm that these results were not due to absorption by

melanin ghosts of secreted CXCL8, we examined neutrophil migration to the apical compartment

in epithelium exposed to melanin ghosts in the presence and absence of exogenous CXCL8.

Indeed, despite the presence of melanin ghosts, exposure to CXCL8 triggered robust neutrophil

migration (**Figure 2g**). These data suggest that melanin ghosts block the secretion of CXCL8 and

CXCL1 rather than adsorb extracellular chemokines.

The inhibitory effects of Aspergillus melanin extend to other pro-inflammatory

stimuli

Thus far, our experiments suggested that melanin was not acting as a passive protective shield,

but directly downregulated pro-inflammatory responses. We next sought to determine if fungal

melanin anti-inflammatory actions were specific to fungal pathogens or if they represented a more

general host-microbe interaction. *P. aeruginosa* is an important bacterial respiratory pathogen

frequently found in co-infections with *A. fumigatus* in patients with CF. Therefore, we examined

whether *Aspergillus* melanin ghosts could dampen *P. aeruginosa*-induced airway epithelial

inflammation. We infected H292 monolayers with *P. aeruginosa* alone or in combination with

melanin ghosts for 6h then quantified chemokine secretion. The addition of melanin ghosts

suppressed CXCL8 (**Figure 3a**) and CXCL1 (**Supplementary Figure 3a**) release in response to

P. aeruginosa in a dose-dependent manner*.* To ensure that the melanin ghosts were not

interfering with the ability of *P. aeruginosa* to bind to the epithelium, we confirmed that *P.*

aeruginosa binding or internalization by the epithelial cells was not altered in the presence of

melanin ghosts (**Supplementary Figure 3b**), suggesting that the difference in chemokine

secretion was not due to difference in the ability of *P. aeruginosa* to access the cells. Additionally,

we performed an LDH cell viability assay that demonstrated no difference in cell viability in the

presence of *Pseudomonas* with or without melanin (**Supplementary Figure 3c**).

TNFα is a known inducer of secretion of CXCL8 by airway epithelial cells, therefore we

hypothesized that melanin ghosts also block cytokine-induced CXCL8 in airway epithelial cells.

We stimulated H292 epithelium with recombinant human TNFα for 6h in the presence of absence

of melanin ghosts then quantified CXCL8 secretion in supernatants. The combination of melanin

ghosts and TNFα stimulation of epithelial cells failed to secrete CXCL8 compared to stimulation

with TNFα alone (**Figure 3b**).

Melanin blocks* Pseudomonas-*induced transepithelial migration of neutrophils

Since co-stimulation of airway epithelium with *P. aeruginosa* and melanin ghosts resulted

in a potent decrease of CXCL8 secretion, we examined the role of melanin ghosts in airway

epithelial-induced neutrophils transmigration in response to *P. aeruginosa*. While CXCL8 is a

known neutrophil chemoattractant that plays a role in epithelial-induced neutrophil

transmigration[42,43], recruitment of neutrophils across the epithelium has been linked to other

secreted compounds, such as hepoxilin A3, in response to *P. aeruginosa* [44-47]. We stimulated both

primary hAEC and H292 with *P. aeruginosa* alone or in combination with melanin ghosts and

quantified neutrophil transmigration using a myeloperoxidase assay. The addition of melanin

ghosts decreased *P. aeruginosa*-induced neutrophil transmigration (**Figure 3c**), demonstrating

that melanin blocked all mediators required for neutrophil recruitment to *Pseudomonas*.

To determine if blunting of neutrophil influx occurs in an *in vivo* model of infection, we

infected mice with *P. aeruginosa* alone or in combination with melanin ghosts. As predicted, *P.*

aeruginosa induced potent neutrophil recruitment into the airways of mice; however the number

of neutrophils in the bronchoalveolar lavage fluid was significantly decreased in the presence of

melanin (**Figure 3d**). Together, these data indicated that modulation of pro-inflammatory

processes by melanin ghosts extends to recruitment of neutrophils to the site of infection *in vivo*.

Structurally distinct melanins from diverse fungal sources block airway epithelial

CXCL8 expression

Melanins are darkly pigmented molecules created through the polymerization of phenol or indole precursors forming polymers of high molecular weight[48,49]. Despite the ubiquity of melanin pigments across kingdoms (*e.g.,* animals, fungi, insects, helminths), the structure of melanin remains poorly defined. *Aspergillus* and dematiaceous molds produces DHN melanin catalyzed by the polyketide synthase pathway[14-16,50]. However, some fungal pathogens contain melanin derived from L-tyrosine or L-dihydroxyphenylalanine (L-DOPA) produced through the action of tyrosinase or laccase, respectively[34,35,51-56]. One example of laccase-derived melanin occurs in *C. neoformans*, an environmental yeast that causes human infections including pneumonia and meningitis. L-DOPA-based melanins are structurally distinct from *Aspergillus* DHN melanin. This structural difference impacts host responses as demonstrated by the recognition of DHN, but not L-DOPA, melanin by the CLR, MelLec or Clec1A, identified on endothelial and immune cells[31-33]. To determine if the ability to inhibit the appearance of epithelial-derived CXCL8 in the supernatant was specific to *Aspergillus* DHN melanin, we utilized both *C. neoformans* melanin ghosts and a synthetic tyrosine-based melanin. As strong inducers of epithelial CXCL8, *Pseudomonas* and amelanotic *ΔpksP A. fumigatus* were used to stimulate H292 epithelium in the presence or absence of *C. neoformans* melanin ghosts. Similar to *Aspergillus* melanin ghosts, *Cryptococcus* melanin ghosts suppressed CXCL8 and CXCL1 in the supernatant (**Figure 4a, Supplementary Figure 4**).

In addition to our *C. neoformans* melanin ghosts, we also tested epithelial responses to a synthetic source of tyrosine-derived melanin. We previously created fungal-like particles (FLP) using purified fungal carbohydrates which associated with amine-coated polystyrene 3 μM microspheres using either chemical conjugation or adsorption[57,58]. Using this technique, we coated beads to create FLP with synthetic L-DOPA and β-1,3 glucan. β-1,3-glucan is a pro-

inflammatory carbohydrate found in most fungal cell walls and was added as an additional control

to demonstrate that any effect was due to the melanin and not a result of any coating applied to

the microspheres. We stimulated H292 monolayers with *P. aeruginosa* either alone or in

combination with unmodified FLP (naked beads), L-DOPA melanin-coated FLP, or β-1,3-glucan-

coated FLP. We found that combined treatment with *P. aeruginosa* and L-DOPA-coated FLP led

to a reduction in CXCL8 secretion, while no difference in CXCL8 secretion was observed when

P. aeruginosa was combined with either unmodified FLP or β-1,3-glucan-coated FLP (**Figure 4b**).

Taken together, these results demonstrate that the immunomodulatory effect of melanin is not

specific to *Aspergillus* melanin, but also occurs in response to *C. neoformans* L-DOPA based

melanin and a synthetic tyrosine-based melanin.

Melanin abolishes the transepithelial gradient of CXCL8 required for neutrophil transmigration

Since epithelial cells are polarized, they secrete effector molecules selectively to either the apical

or basolateral compartments or bidirectionally. Neutrophil transmigration to the apical surface (*i.e.*

airway), requires the creation of a chemokine gradient established by differential secretion at the

apical and basolateral surfaces[59-61]. To understand whether apical stimulation with melanin

affected only apical chemokine secretion or both the apical and basolateral compartments, we

stimulated H292 monolayers grown on Transwells with *P. aeruginosa* in the presence or absence

of melanin ghosts and collected supernatants from both the apical and basolateral compartments.

H292 were chosen to model this interactions because basolateral chemokine secretion by hAECs,

even in the presence of *P. aeruginosa* was below the limit of detection (*data not shown*).

Stimulation with *P. aeruginosa* induced higher release of CXCL8 on the apical surface of the

epithelium when compared to the basolateral surface (**Figure 5a**). The combination of *P.*

aeruginosa with *Aspergillus* melanin ghosts significantly decreased both apical and basolateral

epithelium CXCL8 secretion when compared with *Pseudomonas* alone. Furthermore, this co-

stimulation effectively abolished the apical-basolateral chemokine gradient generated by *P.*

aeruginosa.

Melanin induces durable impact on CXCL8 secretion

Thus far, our experiments examined the impact of melanin ghosts for short durations (6h or less).

Thus, we sought to understand whether melanin stimulation of epithelium produced a lasting

effect at the epithelium or was a transient suppression of chemokine release. Melanin ghosts

were durable and were not degraded in our cell culture systems as shown in previous studies

utilizing melanin ghosts[19,62,63]. We stimulated H292 epithelium with *P. aeruginosa* or TNFα in

combination with melanin ghosts for 24h. After 24h stimulation, there was a persistent reduction

of apical CXCL8 secretion by epithelial cells co-stimulated melanin ghosts compared with either

P. aeruginosa or TNFα alone (**Figure 5b**), supporting the hypothesis that melanin silenced the

airway epithelium to prevent inflammation, not simply delaying the initiation of response.

Melanin suppression of chemokine secretion is independent of phagocytosis of

melanin or de novo transcription

To determine the mechanism by which melanin blocked CXCL8 secretion by airway epithelial

cells, we first examined whether phagocytosis of melanin ghosts was required. H292 epithelium

were pre-treated with cytochalasin D, an inhibitor of actin polymerization that effectively blocks

phagocytosis, then stimulated with either TNFα alone or in combination with *Aspergillus* melanin

ghosts. Melanin ghosts inhibited TNFα–induced CXCL8 secretion both in the presence and

absence of cytochalasin D, suggesting that phagocytosis of melanin ghosts was not required for

its anti-inflammatory action (**Figure 6a**). Cytochalasin D alone increased basal levels of secretion

compared to media alone as shown previously in H292 cells[64].

We next investigated if melanin-induced transcription was required for CXCL8

suppression using the transcription inhibitor, actinomycin D. Following pre-treatments with vehicle

control or actinomycin D, epithelial cells were incubated with TNFα, TNFα plus *Aspergillus*

melanin ghosts, or melanin ghosts alone. TNFα induction of CXCL8 required CXCL8

transcription[65], therefore as expected addition of actinomycin D blocked TNFα−induced CXCL8

secretion (**Figure 6b**). Airway epithelial cells produce basal levels of CXCL8 as demonstrated by

the media only controls where there was low, but detectable concentrations of CXCL8. Melanin

ghosts suppressed basal levels of CXCL8 secretion by H292 cells. Actinomycin D-mediated

inhibition of transcription did not alter melanin ghosts-induced suppression of basal CXCL8

(**Figure 6b**). Together, these data revealed that the suppression of CXCL8 secretion was

independent of phagocytosis and *de novo* gene transcription.

Melanin exerts a post-translational blockade on CXCL8 secretion

To understand further how melanin acted at a cellular level, we sought to illuminate the

mechanism by which melanin interfered with CXCL8 secretion by epithelial cells. Therefore, we

performed paired experiments, in which we stimulated both primary hAEC cultures

(**Supplementary Figure 5a-b**) or H292 monolayers (**Figure 6c-d**) with media alone, wildtype *A.*

fumigatus conidia, *ΔpksP* conidia alone or in combination with melanin ghosts, *P. aeruginosa*

alone or in combination with melanin ghosts, and melanin ghosts alone. CXCL8 transcription was

induced by all stimuli compared with media treatment alone and no significant difference in

transcript levels between our pro-inflammatory stimuli (*ΔpksP* or *P. aeruginosa)* either alone or in

combination with melanin, demonstrating that melanin ghosts did not block transcription of CXCL8

(**Figure 6c, Supplementary Figure 5a**). Immunoblotting of epithelial cell lysates revealed

increased intracellular CXCL8 in cells treated with melanin ghosts in combination with a pro-

inflammatory stimulant (*i.e., ΔpksP* conidia or *P. aeruginosa*) compared to cells treated with the

pro-inflammatory stimulant alone (**Figure 6d, Supplementary Figure 5b**). Interestingly, we

observed an increase of intracellular CXCL8 protein within cells treated with melanin ghosts,

which is in direct contrast to the decrease in extracellular release of CXCL8 demonstrated by

ELISA. These data demonstrated that the difference in extracellular CXCL8 protein in epithelial

supernatants was not due to difference in either transcription or translation of the CXCL8 protein.

Taken together our results revealed that melanin interfered with CXCL8 signaling by airway

epithelium through a post-translational blockade of chemokine secretion.

328　*Fungal melanin blocked calcium fluxing in epithelial cells*

Given our data suggesting melanin exerts its effects through a transcription/translation-

independent mechanism, we hypothesized that fungal melanin mediated chemokine reductions

through perturbation of an early signaling pathway- calcium fluxing. To examine if fungal melanin

modulated host responses through calcium signaling, we measured calcium fluxing over time in

airway epithelium stimulated by ionomycin and calcium chloride or media alone in the presence

and absence of melanin ghosts by plate assay (**Figure 7a**) and confocal microscopy (**Figure 7b**).

As expected, treatment with ionomycin and calcium chloride induced robust calcium fluxing, with

a peak around 10 minutes. Melanin potently muted calcium fluxing to similar levels as

unstimulated cells (media alone). Together, these data suggest that fungal melanin modulates

epithelial host responses through calcium signaling.

340　*Melanin disrupts actin structure in airway epithelial cells*

Our results demonstrate chemokine secretion, but not production, is blocked by fungal melanin.

To examine whether actin polymerization, which can be mediated by calcium, changes in airway

epithelium in the presence of *A. fumigatus* melanin ghosts, we imaged phalloidin-stained epithelial

cells pre-treated with *P. aeruginosa* or media alone with or without melanin ghosts. Phalloidin

enables visualization of actin filaments (*i.e.,* F actin). In the presence of fungal melanin, epithelial

cells showed dampened levels of phalloidin staining (**Figure 8a**). Using the same experimental

groups, we next measured the amount of F and G actin by immunoblotting. We added latrunculin

A (an inhibitor of actin polymerization), phalloidin (positive F actin control), and DMSO vehicle as

additional controls. These results confirmed reduced F actin levels (**Figure 8b**), indicating that the

fungal melanin disrupts actin polymerization resulting in diminished chemokine secretion in

response to inflammatory stimulus.

Discussion

Humans are exposed to most fungal pathogens, including *A. fumigatus*, through inhalation of fungal

particles including conidia, yeasts, or hyphal fragments. Upon inhalation, *Aspergillus* conidia are

deposited throughout the host airways, however we lack a detailed understanding of the role that

respiratory epithelium plays in directing the outcome of host-pathogen interaction towards clearance,

invasion, or allergic disease. Here, we utilized primary, well differentiated hAEC at ALI and the

monomorphic H292 cell line to demonstrate that fungal melanin actively downregulates airway

epithelial-driven chemokine responses. Our results demonstrate that *Aspergillus* melanin potently

blocks epithelial-driven chemokines via a post-translational inhibition of secretion by muting calcium

fluxing and disruption of actin polymerization, resulting in the ablation of the chemokine gradient that

silences neutrophil transmigration.

The primary hAEC model is powerful model as it recapitulates the pseudostratified

structure and cellular diversity of conducting airway epithelium, enabling study of rare populations

and the interplay of epithelial cells in disease[26]. Most epithelial cell types cannot be grown in pure

cultures, limiting our understanding of the role of specific epithelial cell subtypes in the

pathogenesis of disease. While this system has clear advantages over oncogenic and

immortalized cell lines, we also confirmed our phenotype in the the cell line, H292, a

mucoepitheliod non-small cell lung carcinoma cell line, as this reagent has previously been used

extensively in modeling airway epithelial responses to bacterial pathogens[27,39,40,66-68]. While this

cell line does not capture the cellular diversity of the airway epithelium, it is well-suited to

mechanistic studies due to uniformity of cellular responses and ability to grow large-scale cultures

without a dependence on patient sample quality. Importantly, our results demonstrate that

melanin effects both primary hAEC and H292 epithelial similarly, indicating that H292 cells can

serve as a relevant model system for mechanistic dissection of melanin sensing and response

pathways.

Prior studies identified an impact on the innate immune sensing of fungal pathogens by

fungal melanin. Melanin reprogramed the metabolome of macrophages and prevented LC3-

associated phagocytosis of *Aspergillus* conidia through sequestration of calcium preventing

calcineurin activation[21,37,69,70]. Additionally, melanin is a known inhibitor of ROS production by

phagocytes[12-17], which is an important mechanism of pathogen killing by macrophages and

neutrophils. Since the airways are the first point of contact in invasive aspergillosis, we sought to

determine the biological role melanin played in innate immune responses. Here, we depict a

critical role for fungal melanin in decreasing airway epithelial pro-inflammatory chemokine

secretion via a post-translational mechanism, independent of both phagocytosis or *de novo*

transcription. A recent study suggested depletion of extracellular CXCL10 and CCL20

chemokines by DHN melanin, but not IL-1α or CXCL8[19]. This study indicates that *Aspergillus*

melanin binds to some pro-inflammatory cytokines in the extracellular space. While we

demonstrate a reduction in CXCL8 and CXCL1 in our epithelial model, our data suggest that this

is not largely in part to adsorption by melanin. The mechanisms responsible for recognition of

fungal melanin by epithelial cells remains unknown and warrant further studies. MelLec (also

known as Clec1A) is a CLR that recognizes *Aspergillus* DHN melanin, but not L-DOPA melanin

of *C. neoformans*[31,69]. Both *A. fumigatus* DHN melanin and *C. neoformans* L-DOPA melanin block

airway epithelial-derived inflammation suggesting a universal melanin effect at the epithelial

surface. Given the specificity of MelLec for only DHN melanin, and the expression of human

MelLec is restricted to endothelial cells and innate cells (**Supplementary Figure 1**)[31], the

respiratory epithelial response to melanin is likely independent of MelLec. Our experiments

suggest that the phagocytosis of melanin particles is not necessary, since cytochalasin D does

not block the ability of melanin ghosts to suppress epithelial CXCL8 secretion, suggesting that

melanin is interacting with a cell surface receptor or a soluble factor released by epithelial cells.

The observations that multiple forms of melanin retain the capacity to blunt the chemokine

secretion by airway epithelial cells suggests that human pathogens exploit this pathway to

establish successful infection. Indeed, multiple organisms possess melanin including *A.*

fumigatus, C. neoformans, and *Bordetella pertussis*[16,34,71]. Our understanding of melanin is

evolving from a structural scaffold that provides protection from the harsh environment it inhabits

to a multi-purpose molecule that disables the immune response at multiple steps. Our finding that

Aspergillus melanin inhibits *P. aeruginosa*-induced airway epithelial chemokine secretion

indicates that *Aspergillus* could play a role in dampening inflammation during co-infections

common in CF patients. Colonization by *Aspergillus* could enhance the virulence of *P. aeruginosa*

in patients by blocking early innate immune cell recruitment to sites of *P. aeruginosa* infection.

Our results indicate the suppression of early signaling cascades induced by fungal

melanin, specifically calcium signaling. Calcium contributes through key signaling mechanisms in

host immunity, such as TLR/CLR signaling, ROS production, and cytokine induction[72,73]. In innate

immune cells, calcium has been linked to CXCL8 secretion in response to thapsigargin. Indeed,

intracellular calcium stores in neutrophils are required and depletion of this calcium was not

sufficient to trigger CXCL8 secretion[74]. Although this phenomenon has not been expanded to

airway epithelial cells, thapsigargin increases CXCL8 in immortalized airway epithelium[75]. Prior

studies have linked *Aspergillus* melanin to sequestration of calcium inside the phagosome to block

LC3-associated phagocytosis in monocytes[70], yet whether melanin impacted calcium-mediated

secretion of cytokines and chemokines was not reported. Our data suggest that sequestration of

calcium extends beyond inhibition of phagocytosis to include dampened chemokine secretion.

We propose that the melanin blunts calcium fluxing resulting in perturbation of actin

polymerization and subsequent impairment in the release of CXCL1 and CXCL8. It is important

to note that changes in actin polymerization may not be the sole contributor of calcium-mediated

signaling in airway epithelial antifungal defense.

While we focused primarily on CXCL8 for this work, fungal melanin also inhibited the

secretion of CXCL1 (**Figure 2b, Supplementary Figure 3a, and Supplementary Figure 4**).

While CXCL8 and CXCL1 are known neutrophil chemoattractants that plays a role in epithelial-

induced neutrophil transmigration other secreted compounds are also required for *P. aeruginosa*

dependent neutrophil transmigration including hepoxilin A3[44-47]. Melanin ghosts dampened

neutrophil transmigration in response to *P.aeruginosa,* indicating that the effect of fungal melanin

may have a broad impact on airway epithelial cell secretion. Both CXCL8 and CXCL1 are secreted

through the trans-Golgi network. The intracellular accumulation of CXCL8 protein with little

detectable extracellular CXCL8 suggests that melanin may interfere with trans-Golgi secretion of

CXCL8. Future studies will interrogate both traditional trans-Golgi secretomal pathways, as well

as alternative secretion systems, such as IL-1β which is secreted via a gasdermin dependent

mechanism, to determine the extent of the melanin secretional blockade.

Fungal melanins act as an anti-inflammatory agent at the respiratory epithelium, blocking

post-translational secretion of chemokines to recruit host immune cells. *Aspergillus* melanin is

required to establish invasive disease as the *ΔpksP* is avirulent in animal models of infection. The

mechanism underpinning the virulence defect of the *ΔpksP* strain has long been attributed to

melanin blockade of phagosomal maturation and ROS production by innate immune cells, critical

requirements for host response to *Aspergillus*. However, our work highlights an additional role for

melanin in the host-pathogen interaction, as an anti-inflammatory that prevents airway epithelial

chemokine secretion thereby decreasing early recruitment of neutrophils to sites of infection.

While there is a clear benefit to the pathogen in preventing inflammatory cell recruitment to the

airways, there is potential benefit to the host in dampening host response to melanin at the airway

epithelium. If melanin fails to silence inflammation in some patients this could lead to chronic

inflammatory conditions including chronic pulmonary aspergillosis and allergic diseases such as

allergic bronchopulmonary aspergillosis and fungal asthma.

Acknowledgements

We would like to thank Drs. Bryan Hurley and Hongmei Mou for insightful discussions regarding

our work. Additionally, we would like to thank Dr. George Chamilos for providing melanin ghosts

to test in parallel with those created in our own lab. J.L.R was supported by NIH/NIAID grant

1K08AI14755. J.M.V is supported by R01AI150181, R01AI136529, and R21AI152499. GDB

acknowledges funding from the MRC Centre for Medical Mycology at the University of Exeter

(MR/N006364/2 and MR/V033417/1) and the Wellcome Trust (102705, 217163).

Author Contributions

Conceptualization – Jennifer Reedy, Kirstine Nolling Jensen, and Jatin Vyas. *Investigation* –

Jennifer Reedy, Kirstine Nolling Jensen, Arianne Crossen, Kyle J. Basham, Chris Reardon,

Hannah Brown Harding, Olivia W. Hepworth, Patricia Simaku, Geneva N. Kwaku, Kazuya Tone,

Janet A. Willment, Delyth M. Reid, Mark H.T. Stappers. *Writing* – Jennifer Reedy, Kirstine Nolling

Jensen, Rebecca Ward, Arianne Crossen, Jatin Vyas. *Manuscript review and revision* – Arianne

Crossen, Kyle Basham, Hannah Brown Harding, Chris Reardon, Olivia W. Hepworth, Patricia

Simaku, Geneva N. Kwaku, Kazuya Tone, Janet A. Willment, Delyth M. Reid, Mark H.T. Stappers,

Gordon D. Brown. *Resources* – Jennifer Reedy, Jayaraj Rajagopal, Gordon D. Brown Jatin Vyas.

Competing Interests

472 The authors declare no competing interests.

474 **Data Availability**

475 The data that support the findings of this study are available from the corresponding author upon

476 reasonable request.

Figure Legends

Figure 1. *Aspergillus* melanin inhibits transepithelial migration of neutrophils. (a) Schematic of neutrophil migration assay. Neutrophil migration across primary human airway epithelial cells (hAECs) **(b)** or H292 cells **(c)** measured by apical MPO assay following infection (2h hAECs; 1h H292) with $1 \times 10^7/cm^2$ wildtype B5233 or *ΔpksP A. fumigatus* conidia in the presence or absence of purified *A. fumigatus* melanin ghosts (MG; $5 \times 10^7/cm^2$). Negative control of media (HBSS) or MG alone were included. Data represents the number of neutrophils as determined by standard curve. Data are represented as mean ± SEM. N=4-6; One-Way ANOVA with Tukey's multiple comparisons test; **$p<0.01$, ***$p<0.001$, ****$p<0.0001$.

Figure 2. *Aspergillus* melanin inhibits secretion of airway epithelial-derived CXCL8 and CXCL1. CXCL8 and CXCL1 in apical cell supernatants measured by Luminex **(a-b)** and ELISA **(c-f)**. **(a-b)** Primary hAECs following 4h stimulation by media alone (HBSS), *P. aeruginosa* PAO1 strain (PSA), three different wildtype (WT) *A. fumigatus* strains ($1 \times 10^7/cm^2$) or *ΔpksP A. fumigatus* conidia ($1 \times 10^7/cm^2$). N=3. See also Figure S2 **(c-d)** H292 cells were stimulated for 4h with HBSS, two WT *A. fumigatus* strains, *ΔpksP* conidia, or *ΔpksP* complemented strain ($1 \times 10^7/cm^2$). N=4. **(e-f)** H292 were stimulated for 4h with HBSS, WT B5233 conidia ($1 \times 10^7/cm^2$), *ΔpksP* conidia ($1 \times 10^7/cm^2$), *ΔpksP* conidia mixed with increasing concentrations of *Aspergillus* melanin ghosts (MG; $0.1x = 1 \times 10^6/cm^2$; $1x = 1 \times 10^7/cm^2$; $10x = 1 \times 10^8/cm^2$), or MG alone ($1 \times 10^8/cm^2$). N=3. **(g)** Neutrophil migration to the apical compartment across H292 epithelium pretreated for 1h with MG ($1 \times 10^8/cm^2$) in the absence and presence of exogenous CXCL8 (3,000 pg/mL). N=3. Data are represented as mean ± SEM. One-Way ANOVA with Tukey's multiple comparisons test; *$p<0.05$, **$p<0.01$, ***$p<0.001$, ****$p<0.0001$ compared to *ΔpksP* alone (a-f) or MG+CXCL8 (g).

Figure 3. The inhibitory effects of *Aspergillus* melanin extended to other pro-inflammatory stimuli. (a) H292 epithelium infected with media alone (HBSS; negative control), *P. aeruginosa* PAO1 strain (PSA) in the absence and presence of increasing *A. fumigatus* melanin ghosts (MG; $0.2x = 2 \times 10^6/cm^2$; $1x = 1 \times 10^7/cm^2$; $5x = 5 \times 10^7/cm^2$) or MG alone for 6h. Apical CXCL8 secretion in the supernatant was measured by ELISA. n=3. See also Figure S3. **(b)** Apical CXCL8 secretion was measured by ELISA following a 6h stimulation of H292 cells with HBSS, PSA ± MG, TNFα (100 ng/mL) ± MG, or *Aspergillus* MG alone ($5 \times 10^7/cm^2$). n=3. **(c)** Neutrophil migration across H292 epithelium in response to PSA alone or in the presence of MG or MG alone ($5 \times 10^7/cm^2$) was measured by apical MPO assay. Data represents the number of neutrophils as determined by the standard curve. n=6. **(d)** 6-12-week-old C57BL/6 mice were infected via oropharyngeal aspiration with PSA (5.85×10^5) with or without *Aspergillus* MG (5.85×10^6) for 4h. The number of neutrophils in the bronchoalveolar lavage fluid was measured by flow cytometry. n=5-6. Data are represented as mean ± SEM. One-Way ANOVA with Tukey's multiple comparisons test; *p<0.05, **p<0.01, ***p<0.001, compared to stimulation alone (*i.e.,* PSA or TNFα alone).

Figure 4. Structurally distinct melanins from diverse fungal sources block airway epithelial CXCL8. (a) H292 cells were stimulated for 6h with wildtype *A. fumigatus* (B5233; $1 \times 10^7/cm^2$), *ΔpksP* conidia ($1 \times 10^7/cm^2$) with or without *C. neoformans* H99 melanin ghosts (*Cn*-MG) ($5 \times 10^7/cm^2$), *P. aeruginosa* PAO1 strain (PSA) with or without *Cn*-MG, or *Cn*-MG alone. CXCL8 apical secretion was measured by ELISA. See also Figure S4. **(b)** H292 epithelium was infected with PSA in the absence or presence of synthetic L-DOPA-coated FLP ($5 \times 10^7/cm^2$), β-1,3-glucan-coated FLP (BG; $5 \times 10^7/cm^2$), or naked FLP ($5 \times 10^7/cm^2$) for 6h, then apical CXCL8 secretion was measured by ELISA. Media or coated beads alone were included as negative controls. Data are represented as mean ± SEM. n=3; One-Way ANOVA with Dunnett's multiple comparisons test; *p<0.05, **p<0.01, ****p<0.0001 compared to stimulation alone (*i.e., ΔpksP* or PSA alone).

Figure 5. Melanin abolished the transepithelial gradient of CXCL8 required for neutrophil transmigration, with a durable impact on CXCL8 secretion. (a) CXCL8 expression in the media from the apical (A) and basolateral (BL) sides of H292 cells was measured by ELISA. H292 cells were infected for 6h with *P. aeruginosa* PAO1 strain (PSA) and *Aspergillus* melanin ghosts (MG; $5 \times 10^7/cm^2$) together or alone. Media alone (HBSS) was used as a negative control. n=6; ~~one~~ two-way ANOVA with Sidak's multiple comparisons test. ****p<0.0001. **(b)** An ELISA assessed apical CXCL8 levels following a 24h stimulation with PSA or TNFα (100 ng/mL) in the absence or presence of MG ($5 \times 10^7/cm^2$). n=3; one-way ANOVA with Tukey's multiple comparison test. Data are represented as mean ± SEM. ***p<0.001, ****p<0.0001.

Figure 6. Melanin suppression of chemokine secretion is independent of phagocytosis of melanin or *de novo* transcription, but exerts a post-translational blockade on CXCL8 secretion. (a) H292 cells were incubated with DMSO vehicle control or cytochalasin D (CytoD; 20μM) for 30 min, then stimulated with TNFα (100 ng/mL) and/or *Aspergillus* melanin ghosts (MG; $2 \times 10^7/cm^2$) for 6h. Apical CXCL8 levels were measured by ELISA. n=3. **(b)** H292 epithelium was exposed to Actinomycin D (1μg/mL), then stimulated as described in A. CXCL8 apical secretion was measured using an ELISA assay. One-Way ANOVA with Tukey's multiple comparisons test; **p<0.01; ****p<0.0001. n=3. **(c-d)** CXCL8 expression in H292 cells lysates was measured by **(c)** qPCR for RNA and **(d)** Western blot for intracellular protein, respectively, following 6h stimulation with *A. fumigatus* wildtype (WT; B5233 strain; $1 \times 10^7/cm^2$), *ΔpksP* conidia ($1 \times 10^7/cm^2$) ± MG, *P. aeruginosa* PAO1 strain (PSA) ± MG, or MG alone ($5 \times 10^7/cm^2$). For Western blots, GAPDH was used as a loading control. See also Figure S5. n=2. Data are represented as mean ± SEM.

Figure 7. Epithelial calcium fluxing dampened by fungal melanin. (a) H292 cells were stimulated with media (HBSS) alone or melanin ghosts for 30 min followed by a 1 h incubation with ionomycin (2 μM) and calcium chloride (1.3 mM) for 2h. Calcium fluxing measured by plate

reader every 15 sec. Cell stimulated with media in the presence or absence of melanin ghosts

produced virtually identical values. **(b)** Representative DIC (top panel) and confocal (bottom

panel) images of H292 calcium fluxing by fluo-4 in the presence or absence of melanin ghosts

when treated with media alone (HBSS without calcium or magnesium) or ionomycin and calcium.

Images from 10 min timepoint. Scale bar = 10 µm.

Figure 8. Melanin disrupts actin structure in airway epithelial cells. (a) Representative

images of phalloidin (green) and DAPI (blue) stained H292 cells stimulated with media alone

(HBSS), *P. aeruginosa* PAO1 strain (PSA; $1x10^7/cm^2$) ± MG, or MG alone ($5x10^7/cm^2$) for 6h.

Scale bar = 10 µm. **(b)** Representative of F actin (top blot), G actin (1:20 dilution; middle blot),

and total actin (bottom blot) measured by Western blot from three independent experiments. H292

cells were stimulated with medial alone (HBSS), *P. aeruginosa* PAO1 strain (PSA; $1x10^7/cm^2$) ±

MG, or MG alone ($5x10^7/cm^2$), DMSO vehicle control, or latrunculin A (LatA) for 6h. Phalloidin

was additionally used as a positive control. **(c)** Quantification of F/G actin ration from (b).

Materials and Methods

H292 culture, upright and inverted Transwell culture

Human NCI-H292 pulmonary mucoepidermoid carcinoma cell line (H292) (American Type Cell Collection) was grown as previously described[28,47,66,67]. H292 cells were cultured in complete RPMI medium (cRMPI) (RPMI 1640, L-glutamine, 10% heat-inactivated fetal bovine serum [FBS] (Gibco) and 1% penicillin/streptomycin) at 37°C in the presence of 5% CO_2. For upright Transwell culture, the apical compartments of Costar 6.5 mm or 12mm Transwell® inserts with 0.4 µm pore polyester membrane (Corning) were treated with a 60% Ethanol, 1:1000 rat tail collagen I (Corning) mixture and left open in tissue culture hood for at least 4h to allow for evaporation. H292 cells were then liberated with 0.25% Trypsin-EDTA (Thermo Fisher Scientific) and resuspended in cRPMI. The cell suspension was seeded onto the apical surface of the Transwells and cRPMI was added to the basolateral compartment. For inverted Transwell culture for use in neutrophil migration assays, Costar 6.5 mm Transwell® inserts with 3.0 µm pore polyester membranes (Corning) were inverted and treated with collagen. H292 cells in suspension were applied to the surface of the inverted Transwells to form a small tension bubble and incubated overnight. The following day, Transwells were placed upright in a Costar culture plate (Corning) and both the apical and basolateral compartments were filled with cRPMI. All transwell cultures were incubated at 37°C in the presence of 5% CO_2 and used on day 7.

Primary human airway epithelial basal cell proliferation and upright and inverted ALI Transwell culture

Primary hAEC were grown as previously described[28,40,76-78]. Basal cells were cultured in small-airway epithelial cell medium (SAGM) (PromoCell) supplemented with 5 µM Y-27632 (Tocris), 1 µM A-83-01 (Tocris), 0.2 µM DMH-1(Selleck Chemicals), and 0.5 µM CHIR99021 (Tocris) and

1% penicillin/streptomycin on laminin-enriched 804G-conditioned media coated plates. For

differentiation of basal cells into a fully pseudostratified epithelium at ALI, the apical compartments

of Costar 6.5 mm or 12mm Transwell® inserts with 0.4 µm pore polyester membrane (Corning)

were coated with 804G-conditioned media for at least 4 hours. Upon removal, a basal cell

suspension in SAGM was applied to the apical compartment, and SAGM was added to the

basolateral compartment and the transwells were incubated overnight. The following day, the

SAGM was replaced with PneumaCult-ALI medium (StemCell) combined 1:1 with DMEM/F-12

(Gibco) and incubated overnight. The media in the apical compartment was then removed to

establish ALI. ALI hAEC were incubated for 16-21 days wherein the apical compartment is kept

dry and the ALI media in the basolateral compartment is refreshed frequently.

For inverted ALI culture for use in neutrophil migration assays, Costar 6.5 mm Transwell®

inserts with 3.0 µm pore polyester membranes (Corning) were inverted and treated with 804G-

conditioned media. Basal cells in suspension in SAGM were applied to the surface of the inverted

Transwells to form a small tension bubble and incubated overnight. The following day, Transwells

were placed upright in a Costar culture plate (Corning) and both the apical and basolateral

compartments were filled with SAGM. The following day, SAGM was replaced with PneumaCult-

ALI medium combined 1:1 with DMEM/F-12 and incubated overnight. The media in the apical

compartment was then removed to establish air-liquid interface. The inverted ALI were incubated

for 16-21 days wherein the apical compartment is kept dry and the media in the basolateral

compartment is refreshed frequently. All ALI cultures were incubated at 37°C in the presence of

5% CO_2. Prior to use, the transepithelial electrical resistance (TEER) was measured on the day

of experiment to ensure that all epithelium were intact.

Fungal and bacterial culture

Aspergillus fumigatus strains *Af293, CEA10, B5233, ΔpksP,* and *ΔpksP* + pksP were used. The

B5233, ΔpksP, and complemented *ΔpksP* + pksP strains were gifted by K.J. Kwon-Chung

(National Institutes of Health; NIH). The *Af293* and *CEA10* strains were gifted by R.A. Cramer,

Darmouth College. All *A. fumigatus* strains were grown at 37°C for 3 – 5 days on glucose minimal

media slants (GMM). Conidia were harvested by applying a sterile solution of deionized water

(Milli-Q, Millipore Sigma, Burlington, MA) containing 0.01% Tween 20 to each slant followed by

gentle surface agitation to liberate conidia using a sterile cotton-tipped swab. The solution was

passed through a 40 μm cell strainer to separate conidia from hyphal fragments. Conidia were

then washed three times with sterile PBS and counted on a LUNA™ automated cell counter

(Logos Biosystems). Conidia were used either on the day of harvest or stored at 4°C overnight for

use the following day. For *in vitro* experiments, conidia were applied to the apical surface. For

animal experiments, conidia were used on the day of harvest.

P. aeruginosa strain PAO1 (ATCC) was grown overnight in 4 mL of LB liquid media at

37°C with shaking. Cultures were inoculated from frozen stocks of PAO1 stored at -80°C. After

shaking overnight, 1 mL of overnight culture was centrifuged at 16,000 g x 5 minutes and washed

three times in HBSS. After the final wash the *P. aeruginosa* was resuspended in 600 μl of HBSS.

For *in vitro* stimulations, the *P. aeruginosa* was used at a 1:100 dilution. For quantification of

bacterial number for *in vivo* experiments, the OD600 of a 1:10 dilution of bacteria in HBSS was

obtained and bacteria were diluted appropriately. For *in vitro* experiments, *P. aeruginosa* was

applied to the apical surface.

Neutrophil isolation

Neutrophils were isolated from healthy volunteer donors in accordance with Massachusetts

General Hospital IRB-approved protocol (2015P000818) as previously described[28,67]. Briefly, 50

mL of blood was drawn via venipuncture into a syringe containing acid citrate-dextrose. The blood

was split between three 50 mL conical tubes and layered on top of a Ficoll-Paque solution, then

centrifuged at 1,500 g for 20 minutes at room temperature to obtain a buffy coat layer. Plasma

and mononuclear cells were removed by aspiration. Red blood cells (RBC) were removed using

2% gelatin sedimentation technique followed by a wash with RBC lysis buffer. The remaining

neutrophils were washed and resuspended in HBSS without calcium or magnesium (HBSS-) to a

concentration of 5 x 10^7 neutrophils/mL.

Neutrophil transmigration experiments

The neutrophil transepithelial migration experiments were performed as previously described

(**Figure 1a**)[28,40,41,67]. H292 cell monolayers or primary hAEC were grown on the underside of 3.0

μm pore size Transwell filters (Corning Life Sciences) to enable neutrophil migration through the

Transwell. Mature epithelium (at least 16 days at ALI for primary hAECs or day 7 monolayers of

H292 cells) in 12 well format were used for all experiments. On the day of the experiment,

Transwells were equilibrated in HBSS for at least 30 minutes prior to stimulation. Transwells were

then inverted into a sterile 15 cm cell culture plate and infected on the apical surface. Cells were

stimulated with media only (HBSS), *P. aeruginosa* (PAO1), *A. fumigatus* conidia, melanin ghosts,

or melanin ghost plus 3,000 pg/mL exogenous CXCL8 (#208-IL; R&D Systems). Infections were

for 1 h on H292 cell monolayers and for 2 h on primary hAEC. After stimulation, the apical surface

was washed and transwells were inverted into fresh 24 well culture plates containing HBSS alone

or HBSS containing 100 nM *N*-formyl-methionyl-leucyl-phenylalanine (fMLP; Sigma; uninfected

positive control). Neutrophils were applied to the basolateral compartment and transmigration was

allowed to proceed for 2h (H292 cells) and 4h (primary hAECs) at 37°C with 5% CO_2. After the

migration period, the Transwells were discarded and the migrated neutrophils were quantified

using a myeloperoxidase (MPO) assay. The migrated neutrophils in the apical compartment were

lysed with 0.5% Triton X-100 and neutrophil MPO activity was quantified using a colorimetric

assay using citrate buffered 2,2'-azino-bis(3-ethylbenxothiazoline-6-sulfonic acid) diammonium salt (ABTS) solution (Sigma-Aldrich). After 10 min of incubation, the optical density at 405 nm was read using a spectrophotometer plate reader. In addition, to the experimental samples, a standard curve was prepared for each experiment was used to determine the number of migrated neutrophils based on the OD_{405} value. Data are displayed as the number of transmigrated neutrophils, extrapolated from the standard curve.

Melanin ghost preparation

Melanin ghosts were prepared using previously published methods[20,34-38]. In addition to melanin ghosts created in our lab using the following protocol, we also tested melanin ghosts created in the laboratory of Dr. George Chamilos with similar results to our own. Briefly, *A. fumigatus* strain B5233 was grown for 3 days in T75 flasks (CELLTREAT) on GMM at 30°C. *C. neoformans* strain H99 (gift from Arturo Casadevall) was grown in liquid L-dopa media and incubatefd at 30°C for 10-14 days. Flasks were seeded from frozen stocks maintained at −80°C. To harvest fungal organisms, a sterile solution of deionized water (Milli-Q; Millipore Sigma) containing 0.01% Tween 20 (Sigma-Aldrich) was added to each flask, and spores were liberated using gentle surface agitation with a sterile swab. The spore solution was passed through a 40-μm cell strainer to separate hyphal debris. To remove the fungal cell wall, spores were washed three times with sterile PBS, then resuspended in 10mg/mL lysing enzymes from *Trichoderma harzianum* which has since been discontinued and replaced with 5mg/mL lyticase from *Arthrobacter luteus* (Sigma-Aldrich) dissolved in a 1 M Sorbitol (Sigma-Aldrich) / 0.1 M Sodium Citrate (Sigma-Aldrich) solution in deionized water. The solution was incubated in a 30°C water bath for 24h. To denature proteins, the spore solution was then washed twice with sterile PBS and resuspended in a 4 M guanidine thiocyanate (Sigma-Aldrich) solution and incubated at room temperature for 18h with nutation. To hydrolyze proteins, spores were then washed twice with sterile PBS and resuspended in a 1.0 mg/mL Proteinase K (Roche Laboratories) in 10mM Tris-HCl (Sigma-

Aldrich), 1mM $CaCl_2$ (Sigma-Aldrich), and 0.5% Sodium Dodecyl Sulfate (Sigma-Aldrich) pH 7.8,

solution in distilled water. The suspension was incubated in a 65°C water bath for 4h, then washed

three times with sterile PBS and resuspended in a 0.9% NaCl solution. A Folch lipid extraction

was then performed wherein the suspension was transferred to a separatory funnel along with

methanol (Thermo Fisher Scientific) and chloroform (Midland Scientific) and mixed vigorously.

This was repeated twice, with the organic layer being drained each time and the soluble melanin-

containing interfacial layer maintained. For the hydrolysis of residual acid-labile constituents, 6M

HCl (Thermo Fisher Scientific) was added to the flask and boiled for 1 hour, then combined with

equal volume sterile PBS to neutralize pH. The melanin ghost suspension was then washed with

sterile PBS and placed in 0.22 µm centrifugal filter units (Merck Millipore Ltd). The melanin ghosts

were then washed 3 times and resuspended in deionized water. The suspension was frozen

overnight at -80°C and lyophilized for two days. The melanin ghosts were finally resuspended in

sterile PBS, sonicated, and counted on a LUNA automated cell counter (Logos Biosystems).

Fungal-like particle (FLP) creation

We have previously created FLP using purified fungal carbohydrates which we can associated

with amine-coated polystyrene 3 µM microspheres (Polysciences) using either chemical

conjugation or adsorption[57,58]. To create melanin-coated particles, synthetic melanin (Sigma) was

sonicated robustly in DMSO to create a fine suspension that was then incubated with amine-

coated polystyrene microspheres for 1h at room temperature with shaking. The microspheres

were washed copiously with PBS to remove all traces of DMSO and free melanin particles. All of

the microspheres obtained a brown coloration consistent with melanin association and there was

no free melanin visible in the flow through from the FLP.

ELISA

Mature epithelium (at least 16 days at ALI for primary hAECs or day 7 monolayers of H292 cells)

in either 12 or 24 well format were used for all experiments. On the day of the experiment, growth

media was removed and cells were equilibrated for at least 30 min in HBSS. HBSS was removed

and replaced with fresh HBSS alone or containing the appropriate stimulants. Since the primary

hAEC are pseudostratified epithelium, the concentration of applied stimuli was based upon the

surface area of the epithelium rather than an MOI calculation. After application of stimuli, cell

culture plates were spun gently at 250 g x 5 min to bring pathogens in contact with epithelial

surface. Stimulations were allowed to proceed for the desired time (4h – 24h). Supernatants were

harvest and centrifuged through Captiva 96 well 0.2 uM filter plate (Agilent Technologies) at 625

g for 10 min. Supernatants were frozen at -80°C until use. Unless otherwise noted, supernatants

measured were from the apical side. ELISAs were performed using either R&D Duoset kits (R&D

Systems) or Biolegend LegendPlex assays (Legendplex™ Custom Panel (Human CXCL8, IL-8))

according to the manufacturers instructions. For the R&D Duoset kits the ELISA were read using

an i3X Spectrophotometer (Molecular Devices, LLC). For the LegendPlex assays, a BD

FACSCelesta™ Flow Cytometer (BD) was used for data acquisition. The resultant flow cytometry

files were then analyzed using the LEGENDplex™ Data Analysis Software Suite. Results were

analyzed using GraphPad PRISM9 software (GraphPad Software). All reported ELISAs were

done in both technical and biological triplicate.

For cytochalasin D experiments, cytochalasin D (Sigma-Aldrich) was added to HBSS

media at a final concentration of 20 μM. Epithelial cells were incubated for 30 min at 37°C prior to

stimulation. At that time the media was exchanged for fresh HBSS containing the indicated

stimulants in addition to 20 μM cytochalasin D. To inhibit transcription, actinomycin D (Sigma-

Aldrich) dissolved in DMSO was added to HBSS at a final concentration of 1μg/mL. An equal

volume of DMSO in HBSS was added to the control wells. Epithelial cells were then incubated for

1h at 37°C prior to stimulation.

In vivo *neutrophil recruitment experiment*

C57BL/6 mice of 6 -12 weeks of age were obtained from Jackson Laboratories and housed in the

Massachusetts General Hospital specific pathogen-free animal facilities for at least one week

prior to infection. *Pseudomonas* strain PAO1 was diluted and quantified using OD600. Inoculums

were prepared consisting of HBSS alone, 5.8×10^5 CFU PAO1 alone, 5.8×10^6 melanin ghosts

alone, or a 5.8×10^5 CFU PAO1 combined with 5.8×10^6 melanin ghosts. Each inoculum was

prepared in 50 μl total volume in HBSS. Mice were infected via oropharyngeal aspiration of the

inoculum under isoflurane anesthesia. Infection was allowed to proceed for 4 hours, at which time

mice were euthanized using CO_2. Mice were exsanguinated, and the trachea was exposed and

cannulated with a 22G catheter (Exel Safelet Catheter) that was secured in place using suture

string. The lungs were flushed with a total of 2.3 ml of PBS containing 5% fetal bovine serum. The

collected bronchoalveolar lavage fluid was centrifuged at 300 g x 5 min to pellet cells. The pelleted

cells were resuspended in FACS buffer (PBS + 1% bovine serum albumin + 2 mM EDTA + 0.1%

sodium azide) containing 1:100 dilution of Fc Block and a 1:250 dilution of Zombie Violet dye

(423113; Biolegend). After incubation at 4°C for 15 minutes, cells were centrifuged and then

resuspended in FACS buffer containing an antibody cocktail consistent of anti-CD11b AF488

(101217; Biolegend), anti-CD45 APC/Cy7 (157618; Biolegend), anti-F4/80 AF647 (123122;

Biolegend), anti-SiglecF PE (155506; Biolegend), and anti-Gr-1 BV650 (108442; Biolegend).

After incubation at 4°C for 20 min cells were washed with FACS buffer and then fixed in 2%

paraformaldehyde for 10 min at room temperature. Cells were washed and then resuspended in

FACs buffer and stored at 4°C (cells were stored no longer than overnight prior to flow cytometry).

Splenocytes for a single mouse were prepared in parallel to use as controls for flow cytometry

gating. Counting beads were added prior to flow and calibration beads were prepared using each

of the fluorophores. Samples were run on a a BD FACSCelesta™ Flow Cytometer, neutrophils

were identified as CD45$^+$F4/80$^-$SiglecF$^-$CD11b$^+$Gr-1$^+$ cells, and analysis was performed using

FlowJo 10 software (BD). Five to six mice were used per experimental group. All animal studies

were conducted under protocols approved by the Institutional Animal Care and Use Committee

Subcommittee on Research Animal Care at Massachusetts General Hospital.

RNA extraction and qRT-PCR

Epithelium was grown on 0.4 μm pore-size Transwells until mature (at least 16 days hAEC or 7

days for H292 cells). On the day of stimulation, media was removed and cells were equilibrated

in HBSS for at least 30 min prior to stimulation with either media alone (HBSS), *Pseudomonas*

aeruginosa (PAO1), *A. fumigatus* conidia or melanin ghosts alone or in combination. Conidia were

applied at 1 x 10^7 conidia per cm^2 and melanin ghosts were applied at 5X concentration at 5 x 10^7

conidia per cm^2. After 6h of stimulation, supernatants were harvested for ELISA. At least 4

Transwells of epithelium were stimulated per condition, one well of epithelium was used for

Western blot analysis and three wells were each harvested for RNA. For RNA prep, the epithelium

was washed with ice-cold HBSS and then RNA was isolated using the QIAGEN RNeasy Mini Kit.

After harvest of cell lysates in QIAGEN RLT buffer containing beta-mercaptoethanol, cell lysates

were transferred to QIAGEN QIAshredder columns to homogenize the cells, the rest of the RNA

extraction was performed according to RNeasy mini kit instructions. RNA was quantified using

nanodrop. cDNA was created using the SuperScript IV VILO kit (Thermo Fisher Scientific)

according to manufacturers instructions. qRT-PCR was performed on cDNA using the Taqman$^\circledR$

Fast Advanced Master Mix (Thermo Fisher Scientific) and Taqman$^\circledR$ Gene Expression assay

probes CXCL8 labelled with FAM (#4331182;Hs00174103_m1; Thermo Fisher Scientific) and

GAPDH labelled with VIC-MGB(#448480;Hs99999905_m1; Thermo Fisher Scientific). qRT-PCR

reactions were run using the Applied Biosystems™ 7500 Fast Real-Time PCR System per kit

instructions and analyzed using the ΔΔCt method. All qRT-PCR experiments were performed in

at least biological duplicates.

Western blot

Epithelium were prepared as described in "RNA extraction and qRT-PCR". After 6h of stimulation

the epithelium was placed on ice and lysed with 1% NP40 lysis buffer containing protease

inhibitor. Following denaturation with NuPAGE LDS Sample Buffer (Thermo Fisher Scientific)

proteins were resolved by SDS-PAGE on NuPAGE gels (NuPAGE gels, Thermo Fisher

Scientific). Proteins were then transferred to methanol-activated PDVF membrane (Perkin Elmer)

using transfer buffer (0.025M Tris, 0.192 M glycine, 20% methanol) and electrophoretic transfer

at 100V for 1h. The membranes were blocked for 1h at room temperature in 5% Milk in PBS-

0.01% Tween 20 (PBST), and then incubated with anti-CXCL8 antibody (#99407; Cell Signaling

Technologies) overnight at 4^{o}C. The G-actin/F-actin assay biochem kit (#BK037; Cytoskeleton,

Inc.) was used for F/G actin fractionation experiments. Latrunculin A at a concentration of 10 µM

and 1:100 phalloidin were used as positive controls for G actin and F actin, respectively. Phalloidin

was added post lysate-collection as described in manufacture instructions for the kit. Membranes

were washed and incubated for 1h at room temperature with secondary HRP conjugated antibody

(#P0447; DAKO). After washing, membranes were visualized using Western Lightning Plus ECL

chemiluminescent substrate (Perkin Elmer, Waltham, MA) and developed using Kodak BioMax

XAR film (Millipore Sigma). Films were then scanned and processed using Adobe Photoshop

2021. Any contrast adjustments were applied evenly to the entire image and adheres to the

standards set forth by the scientific community[79]. All reported western blots were repeated in at

least biological duplicate.

Calcium flux plate assay

H292 cells were seeded into a 96-well plate at a density of 2×10^4 cells/well and incubated

overnight at 37°C with 5% CO_2 and 95% humidity. The cells were washed in serum-free cRPMI

medium and treated with 5 nM Fluo-4 calcium indicator dye (Invitrogen, Thermo Fisher Scientific,

Massachusetts, USA) with Pluronic (Invitrogen) for 1h at 37°C with 5% CO_2 and 95% humidity.

Subsequently, the cells were washed with HBSS without calcium and magnesium (14175-095;

Gibco) and pre-treated with *A. fumigatus* melanin ghosts or media (HBSS) alone for 30 min at

room temperature. Melanin ghosts were removed by washing and cells were stimulated with 2µM

ionomycin and 1.3 mM calcium chloride for 2h. Data was collected during the stimulation with 30

second intervals using a SpectraMax i3x Plate Reader (Molecular Devices, California, USA).

Confocal imaging

For calcium flux assay, H292 cells were plated in Nunc Lab-Tek 8-chambered coverglass

(Thermo Fisher Scientific, Rochester, NY) and allowed to adhere. Cells were grown in HBSS

without calcium and magnesium. Fluo-4 in DMSO was added to cells for 1 h at 37°C in the dark.

Cells were washed and then stimulated with 2µM ionomycin and 1.3 mM calcium chloride in the

presence or absence of melanin ghosts. Cells were imaged every minute for 15 minutes (as

described below).

For phalloidin staining, H292 cells were plated in Nunc Lab-Tek 8-chambered coverglass (Thermo

Fisher Scientific, Rochester, NY) and allowed to adhere. Cells were stimulated with for 6 h,

washed with cold PBS, incubated with glyoxal solution for 1 h at room temperature, and quenched

with 50mM NH_4Cl at room temperature. Cells were permeabilized with Triton X-100 for 4 min at

room temperature. Non-specific binding in cells was prevented by incubating with blocking buffer

(5% gelatin from bovine skin [Sigma Aldrich]) for 15 min. Cells were incubated overnight at 4°C

with phalloidin iFlur 488 conjugate antibodies (Caymen Chemical) in dilution buffer (PBS, 1%

BSA). Following three PBS washes, the chambered coverglass was then mounted with

Vectashield Antifade Mounting Medium (Vector Laboratories, Burlingame, CA) and Prolong

antifade with DAPI (Invitrogen).

Images were captured on a Nikon Ti-Eclipse inverted microscope equipped with a CSU-X1

confocal spinning-disk head (Yokogawa, Sugarland, TX) and Coherent 4W continuous-wave

laser (Coherent, Santa Clara, CA) excited the sample. A high-magnification, high-numerical

aperture objective (Nikon, 100x, 1.49 numerical aperture, oil immersion) was used. Images were

obtained using an EM-CCD camera (Hamamatsu, Bridgewater, NJ). Image acquisition was

performed using MetaMorph software version 7.10.5.476 (Molecular Devices, Downingtown, PA).

Raw images were then cropped using Adobe Photoshop CS5 and assembled in Adobe Illustrator,

version CS4 (Adobe Systems, San Jose, CA).

Luminex assay

24-well Transwells were prepared using basal epithelial cells from three healthy human donors

and differentiated for 16 days at ALI. On the day of stimulation, ALI media was removed and

epithelium were equilibrated in HBSS media for 1.5 h prior to stimulation. The HBSS was removed

and replaced with fresh HBSS media on the basolateral side, and 200 μL of HBSS media

containing 1 x 10^7/cm^2 wild type conidia (strains *CEA10*, *Af293*, *B5233*), *ΔpksP* conidia or *P.*

aeruginosa strain PAO1. After 4h, the apical and basolateral media was harvested and put

through a Captiva 96 well 0.2 μm filter plate at 3000 rpm for 10 min. Supernatants were frozen at

-80°C. Data were processed using the Luminex analytical software. MILLIPLEX Map Human

cytokine/chemokine Magnetic Bead Panel – Premixed 41 Plex (Millipore Sigma), run on a

MAGPIX® xMAP instrument according to manufacturer's instructions. Analysis was performed

using GraphPad Prism.

Pseudomonas *epithelial association assay*

H292 epithelium were infected with PAO1 alone or in combination with 5×10^7 *Aspergillus* melanin ghosts and incubated for 4h. Wells were washed three times with HBSS to remove any nonadherent bacteria. To lyse epithelial cells, 1% Triton X-100 was added to each well, and plates were incubated with gentle agitation for 1h at 4°C. The lysate was collected, vortexed, and 1:10 serial dilutions were made in HBSS. 100 μL of each dilution was plated onto LB agar plates containing ampicillin and spread using glass beads. The plates were incubated at 37°C overnight and colony forming units were counted the following day.

LDH viability assay

The LDH viability assay was performed using Invitrogen CyQUANT LDH Cytotoxicity Assay (Fisher Scientific) according to the manufacturers instructions. After 6h infection of H292 cells, a 1:10 dilution of 1% Triton X-100 was added to control well containing unstimulated epithelium to lyse the cells and incubated for 10 min at 37°C 5% CO_2. 50 μL of supernatant from each experimental condition (all experimental conditions were performed in triplicate) was transferred to a 96 well plate and mixed with 50 μL of the kit Reaction mixture and allowed to incubated for 30 min at room temperature in the dark. Stop solution was then added to each sample and absorbance at 490 and 680 was measured using an i3X Spectrophotometer (Molecular Devices, LLC). LDH activity and % cytotoxicity were calculated according to assay kit instructions.

REFERENCES

1   Thompson, G. R., 3rd & Young, J. H. *Aspergillus* infections. *N Engl J Med* **385**, 1496-1509 (2021). https://doi.org:10.1056/NEJMra2027424

2   Huang, L. *et al.* Invasive pulmonary aspergillosis in patients with influenza infection: A retrospective study and review of the literature. *Clin Respir J* **13**, 202-211 (2019). https://doi.org:10.1111/crj.12995

3   Rawson, T. M. *et al.* Bacterial and fungal coinfection in individuals with coronavirus: a
rapid review to support COVID-19 antimicrobial prescribing. *Clin Infect Dis* **71**, 2459-
2468 (2020). https://doi.org:10.1093/cid/ciaa530

4   Rayens, E., Norris, K. A. & Cordero, J. F. Mortality trends in risk conditions and invasive
mycotic disease in the United States, 1999-2018. *Clin Infect Dis* **74**, 309-318 (2021).
https://doi.org:10.1093/cid/ciab336

5   Takazono, T. & Sheppard, D. C. *Aspergillus* in chronic lung disease: modeling what
goes on in the airways. *Med Mycol* **55**, 39-47 (2017).
https://doi.org:10.1093/mmy/myw117

6   Bueid, A. *et al.* Azole antifungal resistance in *Aspergillus fumigatus*: 2008 and 2009. *J*
Antimicrob Chemother **65**, 2116-2118 (2010). https://doi.org:10.1093/jac/dkq279

7   Iversen, M. *et al. Aspergillus* infection in lung transplant patients: incidence and
prognosis. *Eur J Clin Microbiol Infect Dis* **26**, 879-886 (2007).
https://doi.org:10.1007/s10096-007-0376-3

8   Lin, S. J., Schranz, J. & Teutsch, S. M. Aspergillosis case-fatality rate: systematic review
of the literature. *Clin Infect Dis* **32**, 358-366 (2001). https://doi.org:10.1086/318483

9   Ward, R. A. & Vyas, J. M. The first line of defense: effector pathways of anti-fungal
innate immunity. *Curr Opin Microbiol* **58**, 160-165 (2020).
https://doi.org:10.1016/j.mib.2020.10.003

10   Latgé, J. P., Beauvais, A. & Chamilos, G. The Cell Wall of the Human Fungal Pathogen
A*spergillus fumigatus*: Biosynthesis, Organization, Immune Response, and Virulence.
Annu Rev Microbiol **71**, 99-116 (2017). https://doi.org:10.1146/annurev-micro-030117-
020406

11   Cho, S.-H., Seo, S.-C., Schmechel, D., Grinshpun, S. A. & Reponen, T. Aerodynamic
characteristics and respiratory deposition of fungal fragments. *Atmospheric Environment*
39, 5454-5465 (2005). https://doi.org:https://doi.org/10.1016/j.atmosenv.2005.05.042

12   Valsecchi, I. *et al.* The puzzling construction of the conidial outer layer of *Aspergillus*
fumigatus. Cell Microbiol **21**, e12994 (2019). https://doi.org:10.1111/cmi.12994

13   Wang, Y., Aisen, P. & Casadevall, A. *Cryptococcus neoformans* melanin and virulence:
mechanism of action. *Infect Immun* **63**, 3131-3136 (1995).
https://doi.org:10.1128/iai.63.8.3131-3136.1995

14   Jahn, B. *et al.* Isolation and characterization of a pigmentless-conidium mutant of
Aspergillus fumigatus with altered conidial surface and reduced virulence. *Infect Immun*
65, 5110-5117 (1997). https://doi.org:10.1128/iai.65.12.5110-5117.1997

15   Langfelder, K. *et al.* Identification of a polyketide synthase gene (pksP) of *Aspergillus*
fumigatus involved in conidial pigment biosynthesis and virulence. *Med Microbiol*
Immunol **187**, 79-89 (1998).

16 Tsai, H. F., Chang, Y. C., Washburn, R. G., Wheeler, M. H. & Kwon-Chung, K. J. The
developmentally regulated *alb1* gene of *Aspergillus fumigatus*: its role in modulation of
conidial morphology and virulence. *J Bacteriol* **180**, 3031-3038 (1998).

17 Sugareva, V. *et al.* Characterisation of the laccase-encoding gene abr2 of the
dihydroxynaphthalene-like melanin gene cluster of *Aspergillus fumigatus*. *Arch Microbiol*
186, 345-355 (2006). https://doi.org:10.1007/s00203-006-0144-2

18 Chai, L. Y. *et al. Aspergillus fumigatus* conidial melanin modulates host cytokine
response. *Immunobiology* **215**, 915-920 (2010).
https://doi.org:10.1016/j.imbio.2009.10.002

19 Graf, K. T., Liu, H., Filler, S. G. & Bruno, V. M. Depletion of extracellular chemokines by
Aspergillus melanin. *mBio* **14**, e0019423 (2023). https://doi.org:10.1128/mbio.00194-23

20 Akoumianaki, T. *et al. Aspergillus* cell wall melanin blocks LC3-associated phagocytosis
to promote pathogenicity. *Cell Host Microbe* **19**, 79-90 (2016).
https://doi.org:10.1016/j.chom.2015.12.002

21 Goncalves, S. M. *et al.* Phagosomal removal of fungal melanin reprograms macrophage
metabolism to promote antifungal immunity. *Nat Commun* **11**, 2282 (2020).
https://doi.org:10.1038/s41467-020-16120-z

22 Kradin, R. L. & Mark, E. J. The pathology of pulmonary disorders due to *Aspergillus spp*.
Arch Pathol Lab Med **132**, 606-614 (2008). https://doi.org:10.5858/2008-132-606-
TPOPDD

23 Latge, J. P. The pathobiology of *Aspergillus fumigatus*. *Trends Microbiol* **9**, 382-389
(2001). https://doi.org:10.1016/s0966-842x(01)02104-7

24 Lee, J. Y. *et al. Aspergillus* Tracheobronchitis and Influenza A Co-infection in a Patient
with AIDS and Neutropenia. *Infect Chemother* **46**, 209-215 (2014).
https://doi.org:10.3947/ic.2014.46.3.209

25 Nyga, R. *et al.* Invasive Tracheobronchial Aspergillosis in Critically Ill Patients with
Severe Influenza. A Clinical Trial. *Am J Respir Crit Care Med* **202**, 708-716 (2020).
https://doi.org:10.1164/rccm.201910-1931OC

26 Montoro, D. T. *et al.* A revised airway epithelial hierarchy includes CFTR-expressing
ionocytes. *Nature* **560**, 319-324 (2018). https://doi.org:10.1038/s41586-018-0393-7

27 Yonker, L. M. *et al.* Neutrophil-Derived Cytosolic PLA2α Contributes to Bacterial-Induced
Neutrophil Transepithelial Migration. *J Immunol* **199**, 2873-2884 (2017).
https://doi.org:10.4049/jimmunol.1700539

28 Feldman, M. B. *et al. Aspergillus fumigatus* Cell Wall Promotes Apical Airway Epithelial
Recruitment of Human Neutrophils. *Infect Immun* **88** (2020).
https://doi.org:10.1128/IAI.00813-19

29 Feldman, M. B., Wood, M., Lapey, A. & Mou, H. SMAD Signaling Restricts Mucous Cell
Differentiation in Human Airway Epithelium. *Am J Respir Cell Mol Biol* **61**, 322-331
(2019). https://doi.org:10.1165/rcmb.2018-0326OC

30 Yonker, L. M. *et al.* Neutrophil dysfunction in cystic fibrosis. *J Cyst Fibros* **20**, 1062-1071
(2021). https://doi.org:10.1016/j.jcf.2021.01.012

31 Stappers, M. H. T. *et al.* Recognition of DHN-melanin by a C-type lectin receptor is
required for immunity to *Aspergillus*. *Nature* **555**, 382-386 (2018).
https://doi.org:10.1038/nature25974

32 Lopez Robles, M. D. *et al.* Cell-surface C-type lectin-like receptor CLEC-1 dampens
dendritic cell activation and downstream Th17 responses. *Blood Adv* **1**, 557-568 (2017).
https://doi.org:10.1182/bloodadvances.2016002360

33 Colonna, M., Samaridis, J. & Angman, L. Molecular characterization of two novel C-type
lectin-like receptors, one of which is selectively expressed in human dendritic cells. *Eur J*
Immunol **30**, 697-704 (2000). https://doi.org:10.1002/1521-4141(200002)30:2<697::Aid-
immu697>3.0.Co;2-m

34 Wang, Y., Aisen, P. & Casadevall, A. Melanin, melanin "ghosts," and melanin
composition in *Cryptococcus neoformans*. *Infect Immun* **64**, 2420-2424 (1996).
https://doi.org:10.1128/iai.64.7.2420-2424.1996

35 Rosas, A. L. *et al.* Synthesis of polymerized melanin by *Cryptococcus neoformans* in
infected rodents. *Infect Immun* **68**, 2845-2853 (2000).
https://doi.org:10.1128/IAI.68.5.2845-2853.2000

36 Folch, J., Lees, M. & Sloane Stanley, G. H. A simple method for the isolation and
purification of total lipides from animal tissues. *J Biol Chem* **226**, 497-509 (1957).

37 Chamilos, G. *et al.* Melanin targets LC3-associated phagocytosis (LAP): A novel
pathogenetic mechanism in fungal disease. *Autophagy* **12**, 888-889 (2016).
https://doi.org:10.1080/15548627.2016.1157242

38 Camacho, E. *et al.* N-acetylglucosamine affects Cryptococcus neoformans cell-wall
composition and melanin architecture. *Microbiology (Reading)* **163**, 1540-1556 (2017).
https://doi.org:10.1099/mic.0.000552

39 Pazos, M. A. *et al. Pseudomonas aeruginosa* ExoU augments neutrophil transepithelial
migration. *PLoS Pathog* **13**, e1006548 (2017).
https://doi.org:10.1371/journal.ppat.1006548

40 Yonker, L. M. *et al.* Development of a Primary Human Co-Culture Model of Inflamed
Airway Mucosa. *Sci Rep* **7**, 8182 (2017). https://doi.org:10.1038/s41598-017-08567-w

41 Yonker, L. M. *et al.* Neutrophil-Derived Cytosolic PLA2alpha Contributes to Bacterial-
Induced Neutrophil Transepithelial Migration. *J Immunol* **199**, 2873-2884 (2017).
https://doi.org:10.4049/jimmunol.1700539

42   Das, S. T. *et al.* Monomeric and dimeric CXCL8 are both essential for *in vivo* neutrophil
recruitment. *PLoS One* **5**, e11754 (2010). https://doi.org:10.1371/journal.pone.0011754

43   Long, M. B. & Chalmers, J. D. Treating Neutrophilic Inflammation in Airways Diseases.
Arch Bronconeumol **58**, 463-465 (2022). https://doi.org:10.1016/j.arbres.2021.11.003

44   Tamang, D. L. *et al.* Hepoxilin A(3) facilitates neutrophilic breach of lipoxygenase-
expressing airway epithelial barriers. *J Immunol* **189**, 4960-4969 (2012).
https://doi.org:10.4049/jimmunol.1201922

45   Pazos, M. A. *et al.* Distinct cellular sources of hepoxilin A3 and leukotriene B4 are used
to coordinate bacterial-induced neutrophil transepithelial migration. *J Immunol* **194**,
1304-1315 (2015). https://doi.org:10.4049/jimmunol.1402489

46   Mrsny, R. J. *et al.* Identification of hepoxilin A3 in inflammatory events: a required role in
neutrophil migration across intestinal epithelia. *Proc Natl Acad Sci U S A* **101**, 7421-
7426 (2004). https://doi.org:10.1073/pnas.0400832101

47   Kubala, S. A., Patil, S. U., Shreffler, W. G. & Hurley, B. P. Pathogen induced chemo-
attractant hepoxilin A3 drives neutrophils, but not eosinophils across epithelial barriers.
Prostaglandins Other Lipid Mediat **108**, 1-8 (2014).
https://doi.org:10.1016/j.prostaglandins.2013.11.001

48   Riley, P. A. Melanin. *Int J Biochem Cell Biol* **29**, 1235-1239 (1997).
https://doi.org:10.1016/s1357-2725(97)00013-7

49   White, L. P. Melanin: a naturally occurring cation exchange material. *Nature* **182**, 1427-
1428 (1958). https://doi.org:10.1038/1821427a0

50   Wheeler, M. H. & Bell, A. A. Melanins and their importance in pathogenic fungi. *Curr Top
Med Mycol* **2**, 338-387 (1988). https://doi.org:10.1007/978-1-4612-3730-3_10

51   Rhodes, J. C., Polacheck, I. & Kwon-Chung, K. J. Phenoloxidase activity and virulence
in isogenic strains of Cryptococcus neoformans. *Infect Immun* **36**, 1175-1184 (1982).
https://doi.org:10.1128/iai.36.3.1175-1184.1982

52   Kwon-Chung, K. J., Tom, W. K. & Costa, J. L. Utilization of indole compounds by
Cryptococcus neoformans to produce a melanin-like pigment. *J Clin Microbiol* **18**, 1419-
1421 (1983). https://doi.org:10.1128/jcm.18.6.1419-1421.1983

53   Salas, S. D., Bennett, J. E., Kwon-Chung, K. J., Perfect, J. R. & Williamson, P. R. Effect
of the laccase gene CNLAC1, on virulence of Cryptococcus neoformans. *J Exp Med*
184, 377-386 (1996). https://doi.org:10.1084/jem.184.2.377

54   Tsai, H. F., Washburn, R. G., Chang, Y. C. & Kwon-Chung, K. J. Aspergillus fumigatus
arp1 modulates conidial pigmentation and complement deposition. *Mol Microbiol* **26**,
175-183 (1997). https://doi.org:10.1046/j.1365-2958.1997.5681921.x

55   Eisenman, H. C. *et al.* Microstructure of cell wall-associated melanin in the human
pathogenic fungus Cryptococcus neoformans. *Biochemistry* **44**, 3683-3693 (2005).
https://doi.org:10.1021/bi047731m

Eisenman, H. C. *et al. Cryptococcus neoformans* laccase catalyses melanin synthesis from both D- and L-DOPA. *Microbiology (Reading)* **153**, 3954-3962 (2007). https://doi.org:10.1099/mic.0.2007/011049-0

Tam, J. M., Mansour, M. K., Khan, N. S., Yoder, N. C. & Vyas, J. M. Use of fungal derived polysaccharide-conjugated particles to probe Dectin-1 responses in innate immunity. *Integr Biol (Camb)* **4**, 220-227 (2012). https://doi.org:10.1039/c2ib00089j

Reedy, J. L. *et al.* The C-Type Lectin Receptor Dectin-2 Is a Receptor for *Aspergillus fumigatus* Galactomannan. *mBio* **14**, e0318422 (2023). https://doi.org:10.1128/mbio.03184-22

Chow, A. W. *et al.* Polarized secretion of interleukin (IL)-6 and IL-8 by human airway epithelia 16HBE14o- cells in response to cationic polypeptide challenge. *PLoS One* **5**, e12091 (2010). https://doi.org:10.1371/journal.pone.0012091

Pillai, D. K. *et al.* Directional secretomes reflect polarity-specific functions in an in vitro model of human bronchial epithelium. *Am J Respir Cell Mol Biol* **50**, 292-300 (2014). https://doi.org:10.1165/rcmb.2013-0188OC

Skronska-Wasek, W., Durlanik, S., Garnett, J. P. & Pflanz, S. Polarized cytokine release from airway epithelium differentially influences macrophage phenotype. *Mol Immunol* **132**, 142-149 (2021). https://doi.org:10.1016/j.molimm.2021.01.029

Thywißen, A. *et al.* Conidial dihydroxynaphthalene melanin of the human pathogenic fungus *Aspergillus fumigatus* interferes with the host endocytosis pathway. *Front Microbiol* **2**, 96 (2011). https://doi.org:10.3389/fmicb.2011.00096

Amin, S., Thywissen, A., Heinekamp, T., Saluz, H. P. & Brakhage, A. A. Melanin dependent survival of Apergillus fumigatus conidia in lung epithelial cells. *Int J Med Microbiol* **304**, 626-636 (2014). https://doi.org:10.1016/j.ijmm.2014.04.009

van den Berg, A. *et al.* Cytoskeletal architecture differentially controls post-transcriptional processing of IL-6 and IL-8 mRNA in airway epithelial-like cells. *Exp Cell Res* **312**, 1496-1506 (2006). https://doi.org:10.1016/j.yexcr.2006.01.010

Cambier, S., Gouwy, M. & Proost, P. The chemokines CXCL8 and CXCL12: molecular and functional properties, role in disease and efforts towards pharmacological intervention. *Cellular & Molecular Immunology* **20**, 217-251 (2023). https://doi.org:10.1038/s41423-023-00974-6

Hurley, B. P., Siccardi, D., Mrsny, R. J. & McCormick, B. A. Polymorphonuclear cell transmigration induced by *Pseudomonas aeruginosa* requires the eicosanoid hepoxilin A3. *J Immunol* **173**, 5712-5720 (2004). https://doi.org:10.4049/jimmunol.173.9.5712

Kusek, M. E., Pazos, M. A., Pirzai, W. & Hurley, B. P. *In vitro* coculture assay to assess pathogen induced neutrophil trans-epithelial migration. *J Vis Exp*, e50823 (2014). https://doi.org:10.3791/50823

68 Crossen, A. J. *et al.* Human Airway Epithelium Responses to Invasive Fungal Infections:
A Critical Partner in Innate Immunity. *J Fungi (Basel)* **9** (2022).
https://doi.org:10.3390/jof9010040

69 Chamilos, G. & Carvalho, A. *Aspergillus fumigatus* DHN-Melanin. *Curr Top Microbiol*
Immunol **425**, 17-28 (2020). https://doi.org:10.1007/82_2020_205

70 Kyrmizi, I. *et al.* Calcium sequestration by fungal melanin inhibits calcium-calmodulin
signalling to prevent LC3-associated phagocytosis. *Nat Microbiol* **3**, 791-803 (2018).
https://doi.org:10.1038/s41564-018-0167-x

71 Hiramatsu, Y., Nishida, T., Nugraha, D. K., Sugihara, F. & Horiguchi, Y. Melanin
Produced by Bordetella parapertussis Confers a Survival Advantage to the Bacterium
during Host Infection. *mSphere* **6**, e0081921 (2021).
https://doi.org:10.1128/mSphere.00819-21

72 Hann, J., Bueb, J. L., Tolle, F. & Bréchard, S. Calcium signaling and regulation of
neutrophil functions: Still a long way to go. *J Leukoc Biol* **107**, 285-297 (2020).
https://doi.org:10.1002/jlb.3ru0719-241r

73 Kong, F., You, H., Zheng, K., Tang, R. & Zheng, C. The crosstalk between pattern-
recognition receptor signaling and calcium signaling. *Int J Biol Macromol* **192**, 745-756
(2021). https://doi.org:10.1016/j.ijbiomac.2021.10.014

74 Kuhns, D. B., Young, H. A., Gallin, E. K. & Gallin, J. I. Ca2+-dependent production and
release of IL-8 in human neutrophils. *J Immunol* **161**, 4332-4339 (1998).

75 Waters, V. *et al.* The effect of cyclosporin A on airway cell proinflammatory signaling and
pneumonia. *Am J Respir Cell Mol Biol* **33**, 138-144 (2005).
https://doi.org:10.1165/rcmb.2005-0005OC

76 Levardon, H., Yonker, L. M., Hurley, B. P. & Mou, H. Expansion of Airway Basal Cells
and Generation of Polarized Epithelium. *Bio Protoc* **8** (2018).
https://doi.org:10.21769/BioProtoc.2877

77 Mou, H. *et al.* Dual SMAD Signaling Inhibition Enables Long-Term Expansion of Diverse
Epithelial Basal Cells. *Cell Stem Cell* **19**, 217-231 (2016).
https://doi.org:10.1016/j.stem.2016.05.012

78 Mou, H. *et al.* Airway basal stem cells generate distinct subpopulations of PNECs. *Cell*
Rep **35**, 109011 (2021). https://doi.org:10.1016/j.celrep.2021.109011

79 Rossner, M. & Yamada, K. M. What's in a picture? The temptation of image
manipulation. *J Cell Biol* **166**, 11-15 (2004). https://doi.org:10.1083/jcb.200406019

1110

a Stimulus

MPO Assay from Apical Compartment

b Primary hAECs

c H292 Cells

a

CXCL8 (pg/ml)

HBSS
PSA
PSA + MG
MG

A   BL   A   BL   A   BL   A   BL

****

b

24 Hours

CXCL8 (pg/ml)

HBSS   PSA   PSA +MG   TNFα   TNFα +MG   MG

***

****