Competence for transcellular infection in the root cortex involves a post-

replicative, cell-cycle exit decision in *Medicago truncatula*.

Morgane Batzenschlager[1,*], Beatrice Lace[1], Ning Zhang[1,2], Chao Su[1], Sabrina Egli[1], Pascal Krohn[1,3],

Jule Salfeld[1], Franck Anicet Ditengou[1,4], Thomas Laux[1,2], and Thomas Ott[1,3,*].

[1] University of Freiburg, Faculty of Biology, 79104 Freiburg, Germany

[2] CIBSS – Centre of Integrative Biological Signalling Studies, University of Freiburg, Freiburg,

Germany

[3] present address: Max Planck Institute for Plant Breeding Research, Department for Plant-Microbe

Interactions, 50829 Cologne, Germany

[4] present address: Freiburg University Medical Centre, Institute for Disease Modelling and Targeted

Medicine (IMITATE 79106 , Freiburg, Germany

*Corresponding authors:

morgane.batzenschlager@biologie.uni-freiburg.de; thomas.ott@biologie.uni-freiburg.de

ABSTRACT

During root nodule symbiosis (RNS), cell-division activity is re-initiated and sustained in the root

cortex to create a hospitable cellular niche. Such temporary and spatially confined site is required to

render host cells compatible with the intracellular progression of rhizobia. Although it has been

suggested that early infection events might involve a pre-mitotic cell-cycle arrest, this process has

not been dissected with cellular resolution. Here, we show that a dual-colour Medicago histone

reporter robustly identifies cells with different mitotic or endoreduplication activities in the root

cortex. By imaging deep root tissues, we found that a confined trajectory of cortical cells that are

transcellularly passed by infection threads are in a stage of the cell-cycle that is distinct from directly

adjacent cells. Distinctive features of infected cells include nuclear widening and large-scale

chromatin rearrangements consistent with a cell-cycle exit prior to differentiation. Using a

combination of fluorescent reporters demarcating cell-cycle phase progression, we confirmed that a

reduced proliferation potential and modulating the G2/M transition, a process possibly controlled by

the NF-YA1 transcription factor, mark the success of rhizobial delivery to nodule cells.

INTRODUCTION

Cell-cycle regulation is a key pathway to maintain beneficial and sustainable symbiotic associations

that are of primary importance in both marine and terrestrial ecosystems. One common theme to

stabilize facultative associations is to achieve a mutual control of cell proliferation and differentiation

dynamics in both the host and the symbionts (Kondorosi & Kondorosi, 2004; Russo & Genre, 2021;

Rivera & Davies, 2021). Prime examples are flowering plants from a single phylogenetic clade,

comprising four orders (Fabales, Fagales, Cucurbitales, Rosales) with legumes being the most

prominent ones, that evolved the unique ability to accommodate soil-borne rhizobia intracellularly

(Parniske, 2018). Within this symbiosis, rhizobia fix atmospheric nitrogen and deliver ammonium to

the host in exchange for plant photosynthates. The conversion of atmospheric dinitrogen into a usable

form is highly energy-demanding and requires a low-oxygen environment, which is provided within

nodules, specialized lateral organs formed on the roots that are densely populated by rhizobia. A

widely adopted mechanism to access the developing nodule cells is the formation of cortical infection

threads, transcellular conduits formed upon the invagination of the host plasma membrane and

hijacking the cellular machinery of infected root cortical cells (Timmers et al., 1999; Su et al., 2023).

Although our molecular and cellular understanding of the infection mechanisms is expanding

(Tsyganova et al., 2021; Monroy-Morales et al., 2022; Gao et al., 2022), our knowledge on how these

processes are interconnected with the control of cell-cycle progression in cells hosting intracellular

infection threads has remained fragmentary.

Plants engaging into arbuscular mycorrhizal (AM) or root nodule symbioses (RNS) have recruited a common range of cell-cycle processes (Foucher & Kondorosi, 2000; Russo & Genre, 2021). In *M. truncatula* (hereafter, Medicago) the perception of and colonization by the AM fungus result in occasional cell divisions in the inner cortical layers (Russo et al., 2019) and in diffuse endoreduplication events as intraradical hyphae spread throughout the root cortex (Carotenuto et al., 2019a; Carotenuto et al., 2019b). During RNS, the perception of rhizobia results in the onset of a signalling cascade that triggers an outward gradient of cell-cycle reactivation and cell divisions. Although this is initiated in the pericycle, cell-cycle reactivation occurs throughout the inner cortical (C) layers C5, C4 and the middle C3 layer in Medicago (Timmers et al., 1999; Xiao et al., 2014). The successful establishment of RNS further involves the tight coordination of the infection and organogenesis programmes (Guan et al., 2013; Madsen et al., 2010). It requires the progression of cortical infection threads through the C3 layer prior to subsequent periclinal divisions that produce a persistent nodule meristem (Xiao et al., 2014). The resulting indeterminate nodules will later form well-defined developmental zones, with actively dividing meristematic cells giving rise to post-meristematic, differentiating central tissues of increasing ploidy (Vinardell et al., 2003; Nagymihály et al., 2017). While those cells can be intracellularly colonized, the infection competence of nodule-like structures as being induced by exogenous application of cytokinin (Gauthier-Coles et al., 2019), constitutive activation of common symbiotic components (Gleason et al., 2006; Ried et al., 2014; Singh et al., 2014) or the expression of meristematic factors (Dong et al., 2021) may be massively reduced in many of these spontaneous nodules (Hayashi et al., 2010; Liu et al., 2022). The lack of infectibility cannot be overcome even when applying bacterial strains which enter plant cells from the intercellular space (Liang et al., 2019). This clearly demonstrates the ultimate requirement of cells being in a so far uncharacterized state of infection competence.

Transcriptome profiling of Medicago root hairs, treated with rhizobial Nod factors or undergoing rhizobial colonization, uncovered changes in several cell-cycle related genes (Breakspear et al., 2014; Liu et al., 2019a). This perception of Nod factors results in the increased transcription of

D-type cyclins, genes known to respond to extrinsic signals and to promote cell-cycle progression

from the first GAP (G1) phase to the DNA-synthesis (S) phase (Soni et al., 1995; Dewitte et al.,

2003). Subsequently, the initiation of infection threads coincides with the highest expression of genes

required for DNA replication, modification and repair, strongly suggesting that the epidermal

infection programme involves cell cycle re-entry and progression to a post-replicative phase

(Breakspear et al., 2014). A repression of the endocycle was further proposed to occur since infected

root hairs accumulate transcripts of the *OSD1/UVI4* gene (Breakspear et al., 2014), whose activity

prevents an unscheduled increase in ploidy by inhibiting the anaphase-promoting complex/cyclosome

(Iwata et al., 2011). However, much less is known about the transcriptional signature of infected cells

along the cortical trajectory of transcellularly progressing infection threads. Such ability to guide

rhizobia *via* cortical infection threads is, however, thought to be an important innovation since it has

been evolutionary maintained in the vast majority of nodulating species as it, most likely, maximizes

the host control over bacterial delivery (Parniske, 2018; Cathebras et al., 2022 – pre-print).

A central regulator for controlling infection and organogenesis is the transcription factor

Nodule Inception (*NIN*). Co-opted from nitrate response circuits, NIN acts as a master coordinator of

infection, nodule organogenesis and nodule number (Schauser et al., 1999; Marsh et al., 2007;

Fournier et al., 2015; Soyano et al., 2014; Cathebras et al., 2022 – pre-print). It promotes the

expression of several growth targets (Liu et al., 2019a) including *NF-YA1* (Soyano et al., 2013; Feng

et al., 2021), a subunit of the conserved heterotrimeric Nuclear Factor-Y (NF-Y) transcription factor.

A-, B- and C-type subunits of the NF-Y complex act as specialized modules, interacting with other

transcription factors and recruiting chromatin remodelling enzymes to adjust cell specification to

environmental or developmental needs (Baudin et al., 2015; Zanetti et al., 2016; Myers & Holt, 2018).

In vertebrates, the CCAAT-binding NF-Y transcription factor targets genes involved in all major

activities executed in G2 and mitosis (M; Bolognese et al., 1999; Linhart et al., 2005) and is an

important regulator of the G2/M transition (Manni et al., 2001). In legumes, the rhizobium-induced

NF-YA1 subunit functions in cortical infection thread progression, in the establishment and

maintenance of the nodule meristem and in the early differentiation of nodule cells (Combier et al.,

2006; Laporte et al., 2014; Xiao et al., 2014; Shrestha et al., 2021; Schiessl et al., 2023 – pre-print).

Yet, whether the Medicago NF-YA1 protein regulates the G2/M transition is not known.

The hypothesis that intracellular infection at least partially relies on cell-cycle control is

further supported by structural re-arrangements in cortical cells anticipating transcellular IT

progression. These include the formation of pre-infection threads (PITs), which are transvacuolar,

cytoplasmic bridges comparable to those occurring in pre-mitotic cells (van Brussel et al., 1992;

Timmers et al., 1999). *In situ* detection of marker transcripts revealed that outer cortical cells recruit

histone H4 used during DNA replication but do not express a B-type, mitotic cyclin gene – implying

that activated target cells stop in the G2 phase of the cell cycle (Yang et al., 1994). G2-phase cell-

cycle arrests have been repeatedly described in eukaryotic cells as appropriate gates for executing

cell fate and patterning decisions (Meserve & Duronio, 2017; Meyer et al., 2017), for wound healing

(Cosolo et al., 2019) or for host infection during pathogenic and biotrophic interactions (Wildermuth

et al., 2017; Bardetti et al., 2019). In plants, transiently or sustainably reducing the potential to enter

mitosis is an appropriate response when cells must, for instance, cope with genotoxic stresses (Chen

et al., 2017; Takahashi et al., 2019) or switch to another cell-cycle program, such as the meiotic

(Hernandez-Lagana et al., 2021) or the endoreduplication cycle (De Veylder et al., 2011; Wildermuth

et al., 2017). Modulating the G2/M transition combines well-described transcriptional and post-

translational mechanisms, leading to the repression of genes necessary to execute cell division

(Berckmans & De Veylder, 2009; Kobayashi et al., 2015), controlling the activity of cyclin-dependent

kinases (CDK; Shimotohno et al., 2021) and the stability of transcriptional repressors and mitotic

inducers (Fülöp et al., 2005; Chen et al., 2017). Although previous reports have suggested such cell-

cycle patterns to occur in roots during rhizobial infections (Yang et al., 1994; Breakspear et al., 2014),

this has never been spatially resolved at cellular level.

Here, we employed a combination of cell-cycle reporters to resolve functional steps of cell-

cycle progression on the cortical infection thread trajectory with cellular and subcellular resolution.

We demonstrate that the first generation of Medicago cells successfully internalizing rhizobia shows unique features, including a marked decrease in their histone H3.1 content and a reduced competence for both cell division and chromosome segregation. We also show that tetraploid (4n) cells reaching an 8C DNA content are optimal for infection by rhizobia. Our results reveal that, upon transcellular infection thread passage, cortical cells undergo a last GAP phase after DNA replication and exit to differentiation. Furthermore, we present first evidence that the symbiotically-induced NF-YA1 subunit, by controlling mitotic entry in a heterologous system, holds potential to contribute to this process.

RESULTS

A reduced proliferative potential typifies cells supporting cortical infection thread progression

In *A. thaliana* (hereafter, Arabidopsis), the incorporation and eviction dynamics of histone H3 variants have been used to identify cells with different cell division and reprogramming potentials, during organ patterning (Otero et al., 2016) or when acquiring reproductive competence (Lagana & Autran, 2020). To test whether gaining a stage of intracellular infection competence involves a similar modification of the cell-division potential and associated chromatin reorganization, we searched for Medicago genes coding for the replicative histone variant H3.1 and the replacement variant H3.3 (Figure 1–figure supplement 1A; Probst et al., 2020). The encoded Medicago H3.1 protein is identical to the one present in *L. japonicus* and *H. vulgare* and, as in other flowering plants, differs from H3.3 by only four residues (Figure 1–figure supplement 1B; Shi et al., 2011; Lu et al., 2018). To check that these Medicago *H3* genes encode for *bona fide* H3.1 and H3.3 counterparts, native promoter-genomic fusions with fluorescent tags were expressed in *A. rhizogenes*–induced transgenic roots, formed on plants that were inoculated with the compatible symbiont *S. meliloti*. We adopted a fixation, clearing and cell-wall counter-staining procedure (Ursache et al., 2018) that enabled us to perform detailed imaging deep inside the multi-layered root cortical tissues. Medicago H3 fusion proteins fully recapitulated the patterns observed in Arabidopsis roots (*i.e.,* patchy pattern of H3.1

and constitutive presence of H3.3; Figure 1–figure supplement 1C–E) and showed the expected localization in euchromatin (diffuse labelling) and heterochromatic regions (subnuclear foci; Figure 1–figure supplement 1C–E; Ingouff et al., 2010; Shi et al., 2011; Otero et al., 2016). A low H3.1 content, typical for cells with a reduced proliferative potential, also enabled us to identify putative quiescent centre (QC) cells within the open root apical meristem of Medicago (Figure 1–figure supplement 1E; Rost, 2011; Otero et al., 2016; Xiao et al., 2019; Ghahremani & MacLean, 2020). Preferentially incorporated during DNA replication, the canonical histone H3.1 variant controls the maintenance of genome integrity (Davarinejad et al., 2022) and epigenetic inheritance (Jiang & Berger, 2017) and is consequently maintained at high levels in cells engaged in recurrent cell division or endoreduplication cycles (Figure 1A). Once cells exit the cell cycle, H3.1 is evicted and likely replaced by H3.3 to mediate cellular differentiation in various Arabidopsis cell types (Otero et al., 2016). In Medicago, increased levels of *H3.1* transcripts were coherently detected at an early stage of nodule primordia development (Figure 1– figure supplement 1F; Schiessl et al., 2019) and mRNAs accumulate only in meristematic (ZI) and infection zones of mature nodules (ZIId and ZIIp; Roux et al., 2014), while *H3.3* transcription stayed high in the differentiation zone (ZIII; Figure 1– figure supplement 1G). We verified a corresponding differential accumulation of H3.1 and H3.3 tagged proteins at cellular resolution in nodule sections (Figure 1B). The double histone reporter proved also useful to distinguish actively endoreduplicating cells prior to fungal colonization (Carotenuto et al., 2019; Carotenuto et al., 2019a) from fully differentiated, arbuscule-containing cells (Figure 1C). Since the selected Medicago histone variants recapitulated the expected transcriptional and localization patterns in well-defined developmental zones, we considered them as functional reporters of H3.1 and H3.3. In addition, the H3.1/H3.3 balance enables to track cortical cells re-entering the cell cycle and adapting their responses to both bacterial and fungal symbionts.

We then focused on the H3.1 nuclear content at early interaction stages between rhizobia and the host plant (8 days post inoculation [dpi]; Figure 2A–B). In accordance with RNA sequencing data (Figure 1– figure supplement 1F; Schiessl et al., 2019), we observed the replicative H3.1 variant

accumulating in regularly dividing cortical cells forming the nodule primordium (Figure 2A, left panel). The H3.1-eGFP fusion protein was also visible in cells penetrated but not fully passed by an infection thread (Figure 2A, close up). This is consistent with earlier observations that cells preparing for intracellular infection pass through S-phase (Yang et al., 1994). By contrast, host cells sustaining cortical infection thread progression in the middle and inner cortex (C3 and C4) up to the first recipient cells in the nodule primordium (C4/5-derived; Figure 2B) exhibited enlarged nuclei (Figure 2C) with a significantly decreased histone H3.1 content compared to their direct neighbouring cells (Figure 2D). Such pattern was also observed using another *H3.1* gene expressed in root tissues (*H3.1 (2)*; Figure 2–figure supplement 1A–E). Nuclear enlargement associated with a massive eviction of H3.1 typically identifies cells having completed a DNA replication round, but pausing in the following GAP phase for cell-cycle switches or exit decisions (Figure 2E; Otero et al., 2016; Lagana & Autran, 2020; Probst et al., 2020). Together, our data strongly suggest that cortical cells along the infection thread trajectory, different to their direct neighbours, exhibit a reduced proliferation potential and presumably exit the cell cycle (Figure 2 B–E). The observed nuclear enlargement and large-scale chromatin rearrangements are consistent with an increased transcriptional activity (Knaack et al., 2022) concomitant with sustained infection thread progression (Breakspear et al., 2014; Liu et al., 2019a). This proves the long-standing hypothesis that cortical cells competent for transcellular infection suspend their course in G2 (Yang et al., 1994; Timmers et al., 1999) in Medicago.

Exiting the cell cycle on the cortical infection thread trajectory aligns infection and organogenesis

Next, we tested whether a cell-cycle exit decision and associated remodelling of the histone H3 composition requires the tight coupling of infection and nodule organogenesis as observed under wild-type (WT) conditions. Thus, we compared the H3.1/H3.3 patterns in WT roots to those of two mutants in which nodule organogenesis is either abolished (*daphne-like*; Liu et al., 2019b) or where nodules form but cortical infection threads are impaired in their progression (*nf-ya1-1*; Laporte et al.,

2014). In the WT situation, the majority of cortical cells (90%) enabling transcellular passage until

reaching the nodule primordium showed a reduced proliferation potential (low H3.1 content, n = 41;

Figure 3A and Figure 3–figure supplement 1A). A higher H3.1 content was occasionally observed in

the last infected cell in the absence of nodule primordium organogenesis (NOD [-]; Figure 3–figure

supplement 1B). Absence of inner cortical cell division is also a characteristic feature of the *daphne-*

like Medicago mutant (Figure 3–figure supplement 1C), which lacks a remote cis-regulatory region

in the *NIN* promoter perceiving cytokinin signals (Liu et al., 2019b). This non-nodulating mutant still

produces WT-like root hair infections occasionally reaching outer cortical layers. In this context, the

percentage of infected cortical cells visibly accumulating H3.1 in the euchromatin increased from

10% in the WT to almost 41% in the *daphne-like* mutant (n = 44; Figure 3A and Figure 3–figure

supplement 1D). This indicates that outer cortical cells hosting an infection thread lack an extensive

chromatin reorganization in the absence of sustained nodule organogenesis.

As a second mutant, we chose a *nf-ya1* null allele, producing smaller nodules that are additionally

delayed and impaired in their development (Laporte et al., 2014; Xiao et al., 2014). Nodule primordia

of the *nf-ya1-1* mutant showed reduced cell layers, with a lower frequency of cell divisions in the

inner cortex (C4/5) and only few anticlinal divisions in the middle layer (C3; Figure 3–figure

supplement 1E-G; Xiao et al., 2014; Schiessl et al., 2023 – pre-print). Cortical infection threads

appeared abnormally bulbous (Figure 3A and Figure 3–figure supplement 1E-F) and showed signs

of early abortion (Figure 3–figure supplement 1G; Laporte et al., 2014; Schiessl et al., 2023 – pre-

print). We observed that, even though the majority (76%) of crossed cortical cells were presumably

arrested in a GAP phase (low H3.1 content, n = 54; Figure 3–figure supplement 1F), such pausing

was not observed in about 24% of the inspected events. Here, infection threads were either blocked

or delayed in the outer cortex and the replicative H3.1 variant was retained (Figure 3A and Figure 3–

figure supplement 1G). Although the last infected cortical cells showed a significant nuclear

enlargement in all three genetic backgrounds investigated (Figure 3B), failure to coordinate

organogenesis and infection programs in the mutants resulted in an increased occurrence of infected

cells accumulating more H3.1 than their direct neighbours (Figure 3C). Together, these data reveal

that an appropriate reduction in cell-cycle activity and associated chromatin changes in infected cells

coincide with an efficient progression of cortical infection threads.

Cortical cells supporting transcellular infection control their commitment to cell division

In Medicago and other plants producing indeterminate nodules, a local control of mitotic activity

could be especially important for rhizobia to timely cross the middle cortex, where sustained cellular

proliferation gives rise to the nodule meristem (Xiao et al., 2014). To investigate in greater detail

whether cells competent for transcellular infection are selectively kept away from transiting to

mitosis, we first used a triple fluorescent sensor enabling to track all cell-cycle phases *in planta*.

Originally developed for Arabidopsis, the plant cell-cycle indicator (PlaCCI) combines a component

of the pre-replication complex (CDT1a), the replicative histone variant (H3.1) and the N-terminal

domain of a B-type cyclin (N-CYCB1;1) as G1-, S- and late G2-phase reporters, respectively (Figure

4A; Desvoyes et al., 2020; Echevarria et al., 2021). The N-terminal domain of CYCB1;1 contains a

destruction box (D-box) ensuring a rapid turnover of the protein after one mitotic cell cycle (Colón-

Carmona et al., 1999). These cell-cycle markers are expressed at early stages of organ development

but disappear in terminally differentiated cells (Desvoyes et al., 2020). When expressing the PlaCCI

construct in transgenic Medicago roots, all three markers were successfully visualized in reactivated

cortical cells upon nodule primordium formation (Figure 4B–C). Cortical cells crossed by an infection

thread kept low but detectable amounts of the Arabidopsis H3.1 variant, indicating that they transited

through S-phase (Figure 4C). However, these cells stopped before accumulating substantial levels of

N-CYCB1;1, in contrast to their direct neighbours. We also confirmed latter observations when using

an Arabidopsis CYCB1;2-based fluorescent reporter (Figure 4D; Trolet et al., 2019), labelling only

cells that transit through G2/M (Figure 4–figure supplement 1A; Weimer et al., 2016). Together, these

results verified a selective decrease in mitotic competence occurring on the cortical infection thread

trajectory (Yang et al., 1994) as rhizobia pass the middle cortical layer (C3) in Medicago roots.

Importantly, we did not observe the G1-phase marker accumulating in cells undergoing infection (Figure 4C), favouring the hypothesis of a cell-cycle arrest and exit decision (Figure 2E) rather than a rapid switch to another endoreduplication round (Desvoyes et al., 2019 – pre-print).

We next tested whether transcriptional reporters of a target gene necessary to execute cell division get selectively repressed on the path of a cortical infection. The late G2/M gene *KNOLLE* codes for a syntaxin protein specialized in angiosperm cytokinesis (Park et al., 2018) and is oppositely controlled when cells do (Haga et al., 2007) or shall not (Takahashi et al., 2019) commit to mitosis. The transcription of *KNOLLE* and other late cell-cycle genes (*e.g.*, *CYCB1;2*) in Arabidopsis is primarily regulated by mitosis-specific activator (MSA) cis-elements present in their promoters that are targeted by activator- and repressor-type three Myb repeats (MYB3R) transcription factors (Haga et al., 2007; Kobayashi et al., 2015). Mutations in all three MSA core sequences present in the Medicago *KNOLLE* promoter significantly reduced the expression of a β-glucuronidase (*GUS*) reporter gene in *N. benthamiana* leaf cells in the presence of a hyperactive form of a MYB3R transcriptional activator (NtmybA2Δ630; Figure 4–figure supplement 1B; Araki et al., 2004), suggesting that the Medicago *KNOLLE* gene can be used as a *bona fide* late G2/M readout. Using the pKNOLLE::NLS-3xVenus construct we found that the activation of the *KNOLLE* promoter was lower on the cortical infection thread trajectory in comparison to neighbouring, proliferating cells from nodule primordia (Figure 4E). An equivalent construct, ensuring the timely elimination of fluorescent reporters after one canonical cell cycle (pKNOLLE::D-box-3xVenus-NLS; Figure 4–figure supplement 1C), corroborated a selective control of the G2/M transition in cells undergoing transcellular infection. Combining our observations of enlarged nuclei (Figure 2C), the absence of a G1 marker and the reduced H3.1 levels and expression of mitotic genes (Figure 2D and Figure 4B–E), we conclude that the initial cortical cells along the infection thread trajectory conduct a single cell-cycle round and stay in a post-replicative (*i.e.*, G2) phase as long as they are crossed.

We further took advantage of the fact that the PlaCCI sensor provides a full view on cell-cycle phase progression in root tissues, in Arabidopsis (Desvoyes et al., 2020) and Medicago (this study).

CDT1a- and CYCB1;1-associated signals in the root tip enabled us to identify 2C (*i.e.*, G1) and 4C

(*i.e.*, G2) nuclei, respectively (Figure 4-figure supplement 2A) – where 2C and 4C chromatin-values

correspond to the DNA content of a basic diploid genome. Measuring the area of nuclei exhibiting

the CYCB1;1-YFP signal in cortical layers of the root apical meristem (root tip) or the nodule

primordium revealed that they were on average 1.57 or 1.63 times larger than the ones producing

CDT1a-CFP, respectively (Figure 4-figure supplement 2B). This is in agreement with an increased

DNA content as cells transit from the G1- to the late G2-phase. Surprisingly, proliferating nodule

cells contain significantly wider nuclei than their diploid counterparts in the proliferation zone of the

root apical meristem (Figure 4-figure supplement 2B), suggesting a higher ploidy level early on.

These results lead to the intriguing possibility that reactivated inner cortical cells in Medicago enter

the mitotic cell cycle with an increased amount of genetic material.

Proliferation of endopolyploid cortical cells is a distinctive feature of Medicago root nodules

Our data suggest that the dividing cortical cells contributing to nodule primordium formation are

polyploid – *i.e.*, that they contain more than the two genome copies of a diploid (2n) cell, where "n"

refers to the number of separate chromosomes. To assess this further, we identified and used the

centromere-specific histone H3 variant (CENH3) from Medicago as a marker of individual

chromosomes. Such fluorescently-tagged CENH3 proteins have already been successfully applied

for chromosome quantification *in planta* (Lermontova et al., 2006; De Storme et al., 2016). Aligning

the Medicago CENH3 amino acid sequence (G8A083) with characterized CENH3 sequences from

L. japonicus (Tek et al., 2014), Arabidopsis (Talbert et al., 2002), *H. vulgare* (Sanei et al., 2011) and

O. sativa (Hirsh et al., 2009) revealed the presence of canonical CENH3 features (Figure 5–figure

supplement 1), including a variable N-terminal tail and the centromere targeting domain (CATD;

Karimi-Ashtiyani et al., 2015). When expressing a mCitrine-CENH3 fusion protein under the control

of its own promoter (pCENH3::mCitrine-CENH3), CENH3 labelled up to 16 nuclear foci and

localized exclusively at the presumed centromeres during mitosis (Figure 5A and B) in diploid root

tip cells. As the signal was rather weak, we scored centromeric foci in roots ectopically expressing

the protein (pUbi::mCitrine-CENH3; Figure 5A). Here, a reliable signal was observed in an increased

proportion of nuclei (82%) displaying not more than 14 to 16 mCitrine-CENH3 foci (versus only

55% when *CENH3* was expressed under its endogenous promoter; n = 55; Figure 5B) – reflecting the

base ploidy in *M. truncatula* with 2n = 16. We therefore used the pUbi::mCitrine-CENH3 reporter

construct for our further analyses.

In a first set of experiments, we used CENH3 to discriminate between polyploid cells

undergoing endomitosis or endocycling. In endomitosis, entire chromosome replication and partial

mitotic progression result in a doubling of the chromosome / centromeric foci number (Iwata et al.,

2011; De Storme et al., 2016). By contrast, endocycling cells undergo repeated rounds of DNA

synthesis without segregating the newly replicated parts (Edgar et al., 2014). Here, copied

chromosome arms do not condense and spatial separation of sister kinetochores does not occur, so

that the number of compact CENH3 signals corresponds to the initial number of chromosomes (up to

8C; Figure 5–figure supplement 2A; Lermontova et al., 2006). As the occurrence of endocycling cells

has been previously reported in tissues infected by AM fungi, we first tested the CENH3-reporter in

roots inoculated with *R. irregularis*. Indeed, cortical cells entered by the fungus displayed large nuclei

with no more than 16 labelled foci (Figure 5–figure supplement 2B), supporting their endocycling

nature (Carotenuto et al., 2019a; Lermontova et al., 2006). Up to 16 CENH3 foci were also observed

in diploid cortical cells from the root tip and pericycle cells from lateral root primordia (Figure 5–

figure supplement 2C). However, the majority of nodule cells exhibited between 26 and 32

fluorescent centromeric foci (75,5% in nodule primordia from stage IV to VI, n = 106; Figure 5–

figure supplement 2B–C) and a visibly increased number of chromosomes on mitotic figures (Figure

5–figure supplement 2B). These observations strongly support previously published data proposing

that proliferating cortical cells giving rise to Medicago root nodules are tetraploid (4n = 32; Wipf &

Cooper, 1938; Torrey & Barrios, 1969).

We next tested whether endopolyploid cells are already present at an earlier stage of nodule

development, where cell division events are confined to the pericycle, the inner cortical cells (C4,

C5) and occasionally occur in the third (C3) and second (C2) cortical layers (Figure 5C; Xiao et al.,

2014). Indeed, cells originating only from cortical cell divisions (C2 to C5) showed an increased

number of chromosomes (Figure 5D-E) compared to recently divided cells originating from the

pericycle (Figure 5E). Notably, cells performing anticlinal divisions in the middle cortex (Figure 5D,

left panel) that will later be crossed by the infection thread (Figure 5E, violet triangle) were also

polyploid. From these observations, we concluded that a key feature of cortical cells in Medicago

nodules is their ability to segregate a higher number of chromosome copies - corresponding to 8C in

a tetraploid cell entering mitosis.

Infected cells within the nodule primordium selectively halt their preparations for mitosis

In a next set of experiments, we studied to which extent the cells positioned on the infection thread

trajectory (C3 to C4/5) prepare for mitosis. For this, we made use of the pre-mitotic loading

characteristics of CENH3 to the centromeres, which mainly occurs during the G2 phase and prior to

the spatial separation of sister kinetochores (Figure 6A; Lermontova et al., 2006, 2007, 2011). In

proliferating populations of the root tip (Figure 6–figure supplement 1A) and nodule primordia

(Figure 6A-B), late G2 cells contained large nuclei where the majority of sister kinetochores was split

and appeared as twin foci. We further noticed that centromere separation in Medicago is a gradual

rather than a simultaneous process, as root tip cells with comparable nuclear areas often exhibited an

intermediate number of fluorescent doublets (Figure 6–figure supplement 1A) of similar appearance

and length on a single focal plane (Figure 6–figure supplement 1B–D). To address the question to

which extent cortical cells that are competent for infection progress through the G2 phase and commit

to nuclear division, we assessed the gradual formation of sister centromeres. Although the nuclear

size of infected C3 cells suggested an 8C DNA content, as found in late G2-phase cells of the nodule

tissue (Figure 6B, C), none of these infected cells showed the typical CENH3 twin foci formed when

sister kinetochores split (n = 13; Figure 6B and D). Such control over mitotic commitment was tightly maintained in the majority of infected, C4/5-derived cells (75,5%; n = 53; Figure 6D) which did not display any visible sign of centromere separation across image stacks (Figure 6B), despite a presumably duplicated genomic DNA content (Figure 6C). A smaller proportion of the recently infected cells (20,7%) partially prepared for chromosome segregation and showed up to 3 fluorescent doublets (Figure 6D) of comparable lengths to those measured in G2-phase progressing cells (Figure 6E), whereas only two cells (3,8%) with internalized bacteria got closer to nuclear and likely to cell division (Figure 6B and D). Together, by combining several marker strategies based on histone H3 dynamics and on *bona fide* mitotic reporters, we conclude that most middle (C3) and inner (C4/5-derived) cortical cells competent for intracellular infection enter a post-replicative, G2-type phase without proceeding towards M-phase. Endopolyploid cortical cells hosting infection threads have presumably gained the ability to execute a differentiation program rather than entering cell division at the end of G2.

The symbiosis-specific NF-YA1 subunit holds potential to modulate the G2/M transition

Our data suggest that cell-cycle progression and bacterial infection are highly coordinated, although the precise molecular players able to redirect cell-cycle decisions while promoting cortical infections remain to be identified. Prominent candidates are NF-YA1 proteins that were evolutionary recruited to RNS where they sustain both organogenesis and intracellular infection in a range of nodulating legume and non-legume species (Combier et al., 2006; Laporte et al., 2014; Xiao et al., 2014; Hossain et al., 2016; Soyano et al., 2019; Bu et al., 2020; Shrestha et al., 2021). NF-YA1 plays an unambiguous role as a positive regulator of cell divisions (Soyano et al., 2013) but was also recently shown to function in the early differentiation and specification of nodule cells in Medicago (Schiessl et al., 2023 – pre-print). This prompted us to assess whether NF-YA1 could act as a switch at mitotic entry. To test this, we used WT NF-YA1 and a mutated variant lacking a functional DNA-binding domain (mutDBD) as eGFP fusion proteins. As expected, both variants localized to the nucleus in *N.*

benthamiana epidermal cells (Figure 7A; Laporte et al., 2014). To foster cell cycle re-entry in such differentiated cells, we ectopically expressed the Arabidopsis D-type cyclin *CYCD3;1* in mature leaves as described in the recently developed 'cell division-enabled leaf system' (CDELS; Xu et al., 2020). As these re-activated cells commit to mitosis and perform cytokinesis (Figure 7B), they also enable the activation of promoters containing MSA cis-elements (Xu et al., 2020) as found in the Medicago *KNOLLE* promoter (Figure 4–figure supplement 1B). We confirmed this by expressing eGFP-KNOLLE from its endogenous promoter in the absence and presence of Arabidopsis CYCD3;1. The KNOLLE fusion protein was only detected when co-expressed with CYCD3;1 using immunoblot analyses (Figure 7–figure supplement 1A) and labelled newly formed cell-division planes (Figure 7–figure supplement 1B). This is consistent with previous reports showing that the syntaxin KNOLLE accumulates only in dividing cells. We also confirmed the activation of the Medicago KNOLLE promoter (pKNOLLE::GUS) in CDEL samples ectopically expressing CYCD3;1 using a fluorometric GUS assay (Figure 7C). Next, we tested the impact of NF-YA1 on the activity of promoters driving cell cycle-dependent gene transcription in the CDEL samples. While NF-YA1 did not alter *GUS* expression when driven from the histone *H4* promoter (pH4::GUS) that is active during DNA synthesis (Figure 7C-D; Yang et al., 1994), it significantly repressed pKNOLLE::GUS activation (Figure 7E). This effect was fully abolished when using the mutated NF-YA1$^{\text{mutDBD}}$ variant (Figure 7E). Interestingly, the repressive effect of NF-YA1 was stronger compared to the one observed when using a MYB3R transcriptional repressor from Medicago (Figure 7–figure supplement 2A–B) known to halt the expression of late G2/M genes (Haga et al., 2007; Takahashi et al., 2019). These results suggest that the NF-YA1 DNA-binding subunit prevents an unscheduled entry into mitosis in de-differentiating leaf cells, despite the action of ectopically-expressed CYCD3;1.

We then tested whether such effect could be observed at the cellular level in the CDEL system. For this, we ectopically co-expressed non-fluorescently-tagged CYCD3;1 and NF-YA1 and additionally transformed these leaves with fluorescently tagged Medicago H3.3 and KNOLLE

proteins, expressed from their native regulators (Figure 7–figure supplement 1B). While the constitutive presence of NF-YA1 did not prevent cell division events *per se*, a significantly lower proportion of transformed cells produced detectable levels of KNOLLE (Figure 7–figure supplement 1C), indicating that less pavement cells engaged in mitosis at the investigated experimental timepoint. Cells visibly accumulating the KNOLLE protein still progressed through M-phase in the presence of NF-YA1 (Figure 7–figure supplement 1B), although a smaller proportion of them fully completed cell division at 64 hours when compared to the control samples (Figure 7–figure supplement 1D). From these data we concluded that NF-YA1 is able to control mitotic entry in de-differentiating, endoreduplicated cells at least in the CDEL system. Since symbiotically infected middle and inner primordium layers, where *NF-YA1* expression was found to be confined in (Laporte et al., 2014; Liu et al., 2019b), are comprised of endopolyploid cells (Figure 5C–E), we hypothesize that the Medicago NF-YA1 protein might similarly modulate cell-cycle progression during symbiotic infections of legume roots by rhizobia.

DISCUSSION

Failure to timely coordinate infection and organogenesis programs in the root cortex results in the formation of poorly colonized nodules that are unable to support the plant's demand to obtain fixed nitrogen (Guan et al., 2013; Laporte et al., 2014; Schiessl et al., 2023–pre-print). Here, we sought to understand at cellular resolution, whether cell-cycle progression and cellular reprogramming are altered in a spatially confined array of host cells that gain a time-limited stage of competence for transcellular infection.

Our data support a model where dividing cells contributing to nodule formation, including target cells for cortical infection, are polyploid and contain 4 copies (C) of each chromosome (Figure 5D–E). In Medicago, these 4C cells either pre-exist in the root cortex (Cebolla et al., 1999; Vinardell et al., 2003; Carotenuto et al., 2019a; Carotenuto et al., 2019b) or arise from a single endoreduplication round prior to their possible entry into cell division (Libbenga & Torrey, 1973).

As competent cells are crossed by a cortical infection thread or fully internalize bacteria and despite being embedded into a proliferating tissue, they reach an 8C DNA content (Figure 6C and Figure 8) but without committing to a subsequent cell division (Figure 4C–E). Rather, cells supporting transcellular infections presumably use a cell-cycle GAP phase after DNA replication (*i.e.,* G2) and extend such phase to remodel their histone H3 composition (Figure 2A–B and D). Finally, this differentiation process and concomitant transcellular passage are accomplished within a single, last cell-cycle round (Figure 2E; Otero et al., 2016). Here, and similar to infected root hairs (Breakspear et al., 2014), a rapid switch to another endocycle is likely prevented as both G1- and S-phase markers are kept low (Figure 4C). Taken together, these events control a cell-cycle progression state where molecular players operate to ensure a successful transmission of rhizobia to the first nodule cells.

Modulating endopolyploidy during rhizobial infections

In line with findings from Arabidopsis reporting that a downregulation of *H3.1* gene expression and a massive eviction process contribute to various differentiation programs in the male and female gametes (Ingouff et al., 2010), the root meristem, the embryo (Otero et al., 2016) and the female germline precursor (Lagana & Autran, 2020), we show that the genetic program enabling intracellular infections of Medicago roots accesses the same mechanism (Figure 2A–B and Figure 2–figure supplement 1C–E). Reciprocally, our data indicate that high H3.1 levels tend to be maintained in infected cells with blocked or aborted infections (Figure 3A and C and Figure 3–figure supplement 1A–D). Furthermore, the nuclear enlargement in infected cells (Figure 2C) indicates profound transcriptional reprogramming and activation of new gene regulatory networks as previously shown to occur in LCO-treated (Knaack et al., 2022) or young symbiotic cells (*i.e.*, infected root hairs and 8C cells from mature nodules; Breakspear et al., 2014; Nagymihály et al., 2017; Liu et al., 2019a). We expanded the latter view by showing that tetraploid cortical cells crossed by an infection thread also reach a presumable 8C DNA content (Figure 6C). As reported for growing cell types in Arabidopsis (Bhosale et al., 2018), such alteration in endopolyploidy might support, among others,

changes in cell wall compositions, which have also recently been demonstrated to occur specifically around infected cells in Medicago (Su et al., 2023; Gaudioso-Pedraza et al., 2018). In the determinate nodulator *L. japonicus*, polyploidization of early nodule cells might be controlled by *VAGRANT INFECTION THREAD 1* (*VAG1*), a putative component of DNA topoisomerase VI (topo VI) complex, as cortical cells with reduced ploidy levels cannot be infected in the *vag1* mutant (Suzaki et al., 2014).

Interestingly, but different to infections by mycorrhizal fungi where repetitive rounds of endoreduplication occur (Figure 1C; Russo & Genre, 2021) and support fungal colonization (Carotenuto et al., 2019a; Carotenuto et al., 2019b), markedly reduced histone H3.1 contents in cells maintaining rhizobial infections in the root cortex up to the first recipient cells in nodule primordia indicate reduced endocycling rates (Figure 2D-E; Otero et al., 2016). Such attenuation of cell-cycle progression together with decreased transcript levels of mitotic cyclins were shown to occur in budding yeast and Arabidopsis (Negishi et al., 2016; Gigli-Bisceglia et al., 2018) following the activation of a cell-wall integrity (CWI) checkpoint. More recently, Ma and colleagues proposed a scenario where the CWI kinase THESEUS1 (THE1), with a related gene being induced in infected root hairs (Breakspear et al., 2014; Liu et al., 2019a), triggers cell wall stiffening as a response to endoreduplication attenuation (Ma et al., 2022).

A G2-phase arrest and attenuated endoreduplication as a plausible context for cortical infections

Reaching and staying in a pre-mitotic stage has been proposed to facilitate the long-distance progression of infection threads in legume species forming so-called cortical, pre-infection threads (PITs) (Yang et al., 1994; Timmers et al., 1999; Van Spronsen et al., 2001; Foucher & Kondorosi, 2000). These structures resemble the onset of a cell division plane that requires the formation of a cortical division zone (Smertenko et al., 2018), which is established from late S- through G2 phases (Costa et al., 2016; Schaefer et al., 2017). Interestingly, this cell-cycle progression window coincides

with the one we observed in infected cortical cells (Figure 2A–E and Figure 4A-C). In addition, the existence of pre-mitotic and G2-arrested cortical cells on the infection thread trajectory is in line with a recent report showing that a rhizobium-induced mitotic module, comprising an α-Aurora kinase gene and its MYB3R transcriptional activator, supports endosymbiotic infection in Medicago (Gao et al., 2022). Interestingly though, overexpression of the activator-type *MYB3R1* gene resulted in the formation of abnormally bulbous or branched infection threads (Gao et al., 2022). These data emphasize the need to balance the activity of transcriptional regulators driving mitosis and cytokinesis in infected cells.

In summary, we hypothesize that endocycle suppression after a round of DNA replication (Breakspear et al., 2014) and exiting the cell-cycle from the G2-phase might therefore create a permissive, yet controlled state for cortical infection thread progression where polarization and cell-wall remodelling factors specialized for intracellular infection (Lace et al., 2023; Su et al., 2023) act in a defined cell-cycle window to sustain infection thread growth and guidance across cortical layers.

Endopolyploidization following cytokinin perception: a legume innovation?

Using the centromeric histone H3 (CENH3) fluorescent reporter enabled us to assess chromosome numbers, the progression through the G2 phase (Figure 6–figure supplement 1A–B; Lermontova et al., 2006) and to clearly identify diploid and polyploid cells (Figure 5 B and E; De Storme et al., 2016). Combining CENH3 patterns with the histone H3.3 marker showed that somatic cortical cells giving rise to indeterminate Medicago root nodules are mainly tetraploid, which is consistent with previous observations (Wipf & Cooper, 1938; Torrey & Barrios, 1969). Furthermore, our imaging data indicate that dividing endopolyploid cells are homogenously maintained in the nodule tissue (Figure 5–figure supplement 2B–C). This suggests that tetraploidization is an important feature for nodule fitness and infectability as it may enable processes such as the required integration of many contextual stimuli (*e.g.* bacterial signals) (Doyle & Coate, 2019; Russo & Genre, 2021), the coordination of controlled organogenesis and a greater cellular adaptability to hypoxic environments

as also suggested for human tumour cells (Yant & Bomblies, 2015). Furthermore, our findings point

towards tetraploidization of cortical cells being a hallmark for nodule identity as pericycle cells in

nodule or lateral root primordia remain strictly diploid (Figure 5E and Figure 5–figure supplement

2C). Interestingly, when dissected and cultured in suspension, cortical cells from the susceptible zone

of pea roots entered cell divisions as tetraploid cells only when cytokinin (kinetin) was applied in

addition to auxin (2,4-D; Libbenga & Torrey, 1973). This is particularly interesting since a specific

cortical response to cytokinin, an auxin/cytokinin interplay and subsequent cellular proliferations in

the root cortex appear to be a legume innovation (Guan et al., 2013; Gauthier-Coles et al., 2019).

NF-YA1, possible intermediary between intracellular infection and cell-cycle control

While this and other reports have shed light onto cell-cycle patterns during rhizobial infections in

roots and nodules, genetic factors coordinating cell-cycle progression and intracellular infections

remained enigmatic. Using the heterologous CDEL system (Xu et al., 2020) has enabled us to test the

impact of at least one of the potential candidates, the symbiotic transcription factor NF-YA1, on cell

cycle-dependent activities in a de-differentiating cellular population. Expression of NF-YA1 in this

system did not prevent cells from passing through S-phase (Figure 7D). But, while the activated cells

(as indicated by the induction of the KNOLLE reporter) still engage in mitosis and eventually divide

(Figure 7–figure supplement 1B), significantly fewer transformed cells promoted the expression of

this G2/M marker at our experimental timepoints (Figure 7E and Figure 7–figure supplement 1C)

These results strongly suggest that, in the CDEL system, NF-YA1 negatively regulates the entry into

mitosis despite the presence of CYCD3;1. The latter has been shown to promote mitotic cycles and

to restrain differentiation and endoreduplication when accumulated in shoot tissues (Schnittger et al.,

2002; Dewitte et al., 2003). Our observations are also consistent with findings in mammalian cells,

where the NF-Y trimer mediates the repression of key mitotic genes in complex with the WT p53

protein (Imbriano et al., 2005; Di Agostino et al., 2006) during a G2 arrest induced by genotoxic

stress (Manni et al., 2001). A link to DNA damage control, potentially arising when endoreduplicated

pavement cells are pushed back into a chromosome segregation cycle, seems likely in the CDEL

system. However, whether NF-YA1 similarly monitors cell division activity in polyploid nodule

primordium cells (Figure 5E) and controls the G2/M transition along the infection thread trajectory

and in intracellularly colonized nodules (Figure 6B) are aspects that need to be addressed

independently.

In summary, we propose that a local control of cell-cycle activity and a cell-cycle exit decision along

the infection thread trajectory provide a window of cellular competence for cortical infections to

progress down to the nodule primordium niche in Medicago. We demonstrated that the cell division

and differentiation potentials are controlled in a spatially confined cell-file in the root cortex. We also

show that post-replicative, tetraploid cells reaching an 8C DNA content are optimal for intracellular

infection. Consequently, it is tempting to hypothesize that exactly these cells comprise a highly

specific transcriptional and regulatory landscape that ultimately defines the molecular susceptibility

state for intracellular infection competence.

MATERIALS AND METHODS

Plant materials and growth conditions

The Medicago wild-type (WT) ecotype A17, cultivar Jemalong was used in this study. The *daphne-*

like (FN8113) and *nf-ya1-1*mutant lines have been described previously (J. Liu et al., 2019; Laporte

et al., 2014). Seeds from the *daphne-like* mutant were kindly provided by Rene Geurts (Laboratory

of Molecular Biology, Wageningen University, The Netherlands).

Seeds were scarified by covering them with sulfuric acid (H_2SO_4) 96% for 8 minutes and washed 6

times with sterile tap water. After surface sterilization with a bleach solution (1.2% sodium

hypochlorite, NaClO and 0.1% sodium dodecyl sulfate, SDS) for 1 min, the 6 washing steps were

repeated and seeds were sown on 1% (w/v) agar plates. Seeds were then stratified at 4 °C for 5 days

in the dark before being germinated at 22 °C in the dark for 16 h.

Hairy root transformation

Composite *Medicago* plants were generated with transgenic hairy roots as previously described

(Boisson-Dernier et al., 2001). *Agrobacterium rhizogenes* ARquaI cells carrying the desired binary

vector were cultured in liquid LB medium containing appropriate antibiotics at 28 °C, shaking

overnight. Three hundred μl of the culture (OD 600 nm: 0.5-0.7) were plated on solid LB medium

with corresponding antibiotics and incubated at 28 °C for 2 days. Plants were prepared by removing

the seed coat from germinated seedlings and about 5 mm of the root tip region was removed before

the cut site was dipped into the solid *A. rhizogenes* culture. Up to 25 inoculated seedlings were placed

on solid Fahräeus medium supplemented with 0.5 mM NH_4NO_3. Transformed seedlings were then

grown vertically in a climate chamber at 21° C for 3 days in the dark. After another 4 days at 21 °C

(16 h light/8 h dark, 35 μmol m^{-2} s^{-1} light intensity), with only the roots protected from illumination,

composite plants were transferred to fresh solid Fahräeus plates (0.5 mM NH_4NO_3) and grown for 10

days at 22 °C following the same photoperiod. Transgenic roots were visualized based on the

fluorescence emitted by nuclear markers (*M.t.* pH3.3::H3.3-mCherry; *A.t.* pH3.1::H3.1-mCherry;

pLjUbi::NLS-2xmCherry) using an AXIO Zoom.V16 stereomicroscope (Zeiss). Un-transformed

roots were removed and the composite plants were transferred in pots (7x7x8 cm) containing a

mixture of quartz sand and vermiculite (1:1 v/v) equilibrated with 25 ml of a ¼ Hoagland solution

(0.1 mM KNH3). After 4 days of acclimation (16/8 h light/dark photoperiod, 24 °C temperature, 60%

humidity and 70-90 μmol m-2 s-1 light intensity), plants were subjected to rhizobial inoculation.

For this, *S. meliloti* (Sm2011) strains producing mCherry or GFP were grown on solid TY medium

containing 10 μg/ml tetracycline for 2 days at 28 °C. A single colony was used as an inoculum and

bacteria were grown in liquid TY medium containing the antibiotic at 28 °C, shaking for 2 days. Cells

were centrifuged for 8 min at 2205 g, washed once with liquid Fahräeus medium and finally

resuspended into the same medium to reach an OD 600 nm of 0.0015. Five ml of this bacterial

suspension were applied to each composite plant. Pots were watered twice a week with tap water and

fertilized with a ¼ Hoagland solution (0.1mM KNH3) once a week. Transgenic roots were harvested

at different timepoints (8 to 14 days) for tissue fixation, clearing and deep tissue imaging by confocal

microscopy.

Construct design

For expressing the canonical (*H3.1*) and replacement (*H3.3*) histone genes of *M. truncatula*

(Medicago), genomic fragments containing the promoters (*H3.1*: 2623 bp; *H3.1* (2): 2580 bp; *H3.3*:

2739 bp) and open reading frames (ORF) except the termination codon were used. The pH4::GUS

reporter used in quantitative fluorometric assays comprised a 2080 bp fragment of the promoter.

Expression of the centromeric histone (*CENH3*) gene (coding sequence) was driven by a tentative

promoter comprising 1000 bp upstream of the start codon. Transcriptional KNOLLE reporters were

driven by a 2325 bp-long fragment of the *KNOLLE* promoter. The core pentamer of all three mitosis-

specific activator (MSA) cis-elements (AACGG) within the *KNOLLE* promoter was mutated by

changing two bases (ATTGG) according to Ito et al., 2001. The *KNOLLE* genomic sequence was

used to build the translational reporter (pKNOLLE::eGFP-KNOLLE). For Medicago NF-YA1, the

genomic sequence was used and the NF-YA1[mutDBD] variant was created by replacing the conserved

RGC amino acid residues (positions 213-215) by an AAA triplet according to Laloum et al., 2014.

For the repressor-type *MYB3R* transcription factor, the coding sequence (CDS) was used. All

Medicago sequence information was obtained from the Mt4.0v1 genome through Phytozome.

The *A. thaliana* (Arabidopsis) pCYCB1;2::N-CYCB1;2-NLS-3xVenus reporter construct was

designed according to Trolet et al., 2019 and included a genomic fragment (AT5G06150) of 1929 bp,

comprising 1090 bp of promoter sequence, the 5' untranslated region and 839 bp from the ORF

including the first intron. Addition of 2 bases (GG for Gly) at the 3' end of this genomic fragment

enabled the production of the N-terminal CYCB1;2 domain (including the destruction box [D-box]

motif) fused in-frame to a triple-Venus nuclear reporter (Binder et al., 2014; Nadzieja et al., 2019).

The CYCB1;1 N-terminal (D-box) domain was produced from a genomic fragment (AT4G37490) of

713 bp including the first 2 introns (nucleotides [nts] 41-308 and 488-584) and 129 bp of the third

exon (nts 585-713). This module was fused in-frame to a triple-Venus nuclear FP to generate a

destabilized version of the fluorescent reporter. The CYCD3;1 (AT4G34160) CDS was placed under

control of the cauliflower mosaic virus (CaMV) 35S promoter (426 bp) and fused in frame to a 3xHA

module to build the p35S::CYCD3;1-HA construct. Arabidopsis sequence data can be found on

Phytozome (*Arabidopsis thaliana* TAIR10 annotation version). To produce the C-terminally

truncated, hyperactive version of the *N. tabacum* MybA2 transcriptional activator (pLjUbi::eGFP-

NtMybA2ΔC), the CDS (GenBank accession: AB056123.1) coding for the first 630 amino acid

residues was used (Araki et al., 2004). The GoldenBraid PlaCCI transformation plasmid was kindly

provided by Bénédicte Desvoyes and Crisanto Gutierrez (Centro de Biologia Molecular Severo

Ochoa, CSIC-UAM, Madrid, Spain).

Construct production using the Golden Gate cloning system

The Golden Gate modular cloning (MoClo) system was used to generate binary plasmids for

transgene expression in plant cells (Weber et al., 2011). Promoter, genomic and coding sequences

were synthetized and cloned into pMS-RQ (GeneArt, Thermo Fisher Scientific, Regensburg,

Germany). Level 0 (L0) functional modules used in this study are held for distribution in the ENSA

project collection (https://www.ensa.ac.uk/). Details of the L1 expression cassettes and L2 binary

vector composition, including used promoters, genetically encoded fluorescent proteins (stable or

short-lived) or epitope tags and the accession numbers for requesting constructs, are provided in a

supplementary file (Table S1). One-pot reactions using the Type IIS restriction enzymes BsaI-HFv2

(R3733S) and BpiI / BbsI-HF (R3539S) and the T4 DNA ligase (M0202S), all from New England

BioLabs (Frankfurt am Main, Germany), were assembled according to Binder et al., 2014. Cut-

ligation reactions were cycled 50 times between 37 °C for 2 min and 25 °C for 5 min, followed by 5

min at 50 °C and 5 min at 80 °C. Five µl of the reaction were transformed into *E. coli* Stbl3 (for L1

and L2 plasmids carrying histone H3.1 or H3.3 expression cassettes) or TOP10.

Root colonization by *R. irregularis*

Following excision of un-transformed roots, composite Medicago plants were transferred to chive (*A.*

schoenoprasum) nurse pot (9 x 9 cm) systems containing *R. irregularis* (DAOM 197198, sterile spore

inoculum from Agronutrition, France) set up according to Demchenko et al., 2004 (4 plants x pot).

Each pot was covered with a transparent plastic bag (20 x 30 cm) for two days to keep plants in

moisturized conditions. Plants were grown in a controlled environment chamber at 24 °C with a 16/8

hour light/dark photoperiod, with each pot being watered every second or third day with 50 ml

deionized water and fertilized once per week with 50 ml ¼ Hoagland medium. After 7 to 15 days of

co-culturing, composite plants were harvested and excised transgenic roots were subjected to fixation,

clearing and Calcofluor White staining prior to imaging.

Clearing and staining of the root material

The presence of all expected fluorescent reporters was verified in the meristematic region using an

AXIO Zoom.V16 stereomicroscope (Zeiss) before further analyzing the transgenic roots. Fixation,

clearing and cell-wall counter-staining procedures were performed according to Ursache et al., 2018.

Briefly, transformed roots were isolated and fixed in a freshly prepared 4% paraformaldehyde (PFA)

solution in 1x phosphate-buffered saline (PBS), infiltrated under vacuum for 15 min and then gently

agitated for 60-120 min at room temperature in the dark. After 2 washes in 1x PBS, roots were

transferred to the clearing (ClearSee) solution and incubated 7 to 10 days with gentle shaking,

changing the ClearSee solution every 2 days. Prior to imaging, roots were stained with 0.1%

Calcofluor White in ClearSee for 1 h, before being rinsed once and washed in the clearing solution

for 30 min. Calcofluor White was imaged using a 405 nm laser diode excitation source and a 425–

475 nm detection window.

Nodule sections

To assess the localization pattern of H3.1 and H3.3 histone variants across different nodule zones,

composite plants were generated by hairy root transformation as described above. Nodules were

harvested 35 to 40 days after inoculation in open pots and fixed in 4% (w/v) PFA in 1x PBS, under

vacuum treatment for 15 min and then gently agitated for 45 min at room temperature. The fixed

samples were embedded in 6% (w/v) low melting temperature agarose (Biozym) and longitudinal

sections (70 μm thickness) were obtained using a VT1000S vibratome microtome (Leica Biosystems,

Mannheim, Germany).

Fluorescent reporter imaging by Confocal Laser-Scanning Microscopy (CLSM)

Imaging of plant material (*i.e.,* nodule semi-thin sections and cleared root samples) was performed

using a TCS SP8 confocal microscope (Leica Microsystems, Mannheim, Germany) controlled by the

LAS X v3 software. Images were acquired with a 20x/0.75 or a 40x/1.10 (HC PL APO CS2) water

immersion objective. Genetically encoded FPs were excited with an argon laser or a White Light

Laser (WLL) source. Emitted photons were collected using hybrid detectors (HyDs). eCFP was

excited with an argon laser line at 458 nm and the emission was detected at 470-510 nm. All other

FPs were excited with WLL excitation (ex) lines and emissions (em) were detected according to the

following settings: eGFP = 488 nm (ex) / 500-550 nm (em); YFP = 515 nm (ex) / 525-586 nm (em);

mCitrine and Venus = 515 nm (ex) / 525-550 (em); mCherry: 561 or 587 nm (ex) / 600-650 or 620-

650 nm (em).

Image analysis and quantification of nuclear fluorescent signals

All images were processed and analyzed using the Fiji/ImageJ open-source software (Schindelin et al., 2012). Image files were used to create figure panels through the FigureJ plugin (Mutterer & Zinck, 2013). If not indicated differently, 2D maximum intensity projections of z-stacks are presented in the figures. All quantitative measurements were performed on single focal planes and single channels from un-processed z-stacks, acquired from fully transformed Medicago roots. The Fire lookup table (LUT) was applied to facilitate the visualization of low-intensity pixels. Nuclear areas were measured at the equatorial plane using the freehand selection tool, on channels displaying the signal from chromatin-associated proteins (*i.e.*, H3.1 or H3.3-FP fusions). H3.1-eGFP levels were quantified in the nucleus of infected cells (IC) and of directly adjacent, neighbouring cells (NC) present on the same z-stack in the same cortical layer. For this, an outline was manually drawn around each nucleus at the equatorial plane and area, integrated density and mean gray values were measured. Three oval selections were drawn adjacent to all measured nuclei to be used for normalization against background fluorescence. The corrected total nuclear fluorescence (CTNF) = integrated density – (area of selected nucleus × mean fluorescence of background readings) was calculated according to McCloy et al., 2014 using Excel. The number of mCitrine-CENH3 signals in individual nuclei and mitotic figures was manually determined across image stacks using the multi-point tool. The length of CENH3 fluorescent doublets was measured between the 2 most distant pixels using the straight-line tool.

Transient expression in *N. benthamiana* leaves

Four to 5-weeks old tobacco (*N. benthamiana*) plants were transiently transformed with *A. rhizogenes* (ARqua1) cells carrying the construct of interest. The first, second and third fully expanded true leaves were used for infiltration experiments using needle-less syringes. Agrobacteria were resuspended in a MES/KOH infiltration buffer (10 mM, pH 5.6) supplemented with 10 mM $MgCl_2$ and 150 μM acetosyringone. Bacterial cultures used for co-transformations were diluted to a final OD

600 nm of 0.1 for each construct. *A. tumefaciens* (GV3101) cells producing the p19 silencing

suppressor (Silhavy et al., 2002), were systematically included at a final OD 600 nm of 0.05. Upon

infiltration, plants were further grown for 64 to 72 h before harvesting leaf samples with a biopsy

puncher (4 mm diameter) in the distal half of fully infiltrated leaves. Four to 5 leaf discs were either

mounted in sterile tap water and imaged with a confocal microscope, or 10 leaf disks were frozen in

liquid nitrogen prior to *in vitro* transactivation assays.

Cell division-enabled leaf system (CDELS)

To foster differentiated pavement cells to enter the cell division cycle, non-fluorescently-tagged

Arabidopsis CYCD3;1 was expressed under control of the CaMV 35S promoter by Agrobacterium

infiltration (Xu et al., 2020). The p35S::CYCD3;1-HA construct was co-transformed with binary

vectors allowing native expression of cell cycle regulated-reporters (pH4::GUS, pKNOLLE::GUS,

pKNOLLE::eGFP-KNOLLE) and constitutive expression of transcription factors (NF-YA1, NF-

YA1^{mutDBD}) from Medicago. All plasmids used in the experiments involving the CDEL system are

listed in Table S1. Transformed leaves homogenously expressing nuclear transformation markers

included on the corresponding binary vectors were selected using an AXIO Imager.M2m ApoTome.2

light microscope (Zeiss). The activity of HA-tagged CYCD3;1 was verified by the presence of at

least one mitotic figure with chromosomes labeled by H3.3-mCherry before harvesting.

Immunoblot analyses

To detect proteins transiently produced in the CDEL samples, 3 discs from 64 h post-infiltration

leaves were frozen in liquid nitrogen and ground to powder using a Tissue Lyser MM300 (Retsch).

The powder was resuspended in 120 μl of 2x loading buffer and heated for 8 min at 95 °C. Equal

volumes of samples were loaded and separated by 10% SDS-polyacrylamide gel electrophoresis

(PAGE) at constant voltage (80V). Proteins were transferred to a PVDF membrane (0.2 μm pore size)

using the Trans-Blot® Turbo™ semi-dry transfer system (7 min mixed MW MIDI program) (Bio-

Rad). After blocking in a 2.5% (w/v) milk/TBST solution for 1 h at RT, transferred membranes were

hybridized overnight at 4 °C with the following primary antibodies: anti-GFP (632381, Takara)

(1/5000); anti-DsRed (632496, Takara) (1/5000); peroxidase-conjugated anti-HA (clone 3F10,

12013819001, Roche) (1/2000); anti-FLAG M2 (F3165, Sigma-Aldrich) (1/2000). Membranes were

washed and incubated 1 h at RT with peroxidase-conjugated secondary antibodies (anti-mouse, A

4416; anti-rabbit, A 6154; Sigma-Aldrich) (1/2000). Immunoblotted proteins were detected using the

Clarity Western ECL Substrate (Bio-Rad). Digital images were acquired using an ECL ChemoCam

Imager (Intas Science imaging) with exposures stopped before saturation.

Transactivation and quantitative fluorometric GUS assays

N. benthamiana leaf samples used for GUS fluorometric assays were harvested 72 h post-infiltration.

The same number of plasmids was used for all co-transformation experiments. A construct driving

the accumulation of a nuclear FP (proH3.1::NLS:3xVenus) was included in the absence of exogenous

transcription factors. Ten leaf discs per transformed leaf were frozen in liquid nitrogen inside a Safe

Seal Micro tube containing a stainless-steel bead (5 mm diameter). Samples were ground twice for

30 s (28/s frequency) using a Tissue Lyser MM300 (Retsch) and the powder was resuspended in 400

µl of cold GUS extraction buffer (50 mM NaHPO$_4$, pH 7.0, 1 mM EDTA, 10 mM DTT, 0.05% [v/v]

Triton X-100 and 1x Protease Inhibitor Cocktail [Roche]). Crude leaf extracts were incubated for 2 h

at 4 °C on a turning wheel and centrifuged at 16.900 g for 5 min at 4 °C. Ten µl of the supernatant

were used to determine protein concentrations by the dye-binding Bradford assay (Bio-Rad). Twenty-

five µg of total proteins in a final volume of 100 µl were pipetted into a white 96 well plate (Greiner

Bio-One) before the addition of 200 µl of GUS reaction buffer (50 mM NaHPO$_4$, pH 7.0, 1 mM

EDTA, 1% [v/v] Triton X-100 and 0.5 mM 4-Methylumbelliferyl-β-D-glucuronide [4-MUG] [Carl

Roth]). After 10 min of incubation at 37 °C in the dark, the fluorescence emitted by GUS-mediated

hydrolysis of 4-MUG into 4-Methylumbelliferone (4-MU) was measured using a microplate reader

(POLARstar Omega, BMG LABTECH). Fluorescence was recorded at 360 nm excitation and 460

nm emission filter (gain 500, 20 flashes per well) every 5 min over a time period of 130–180 min.

Samples were run in technical triplicates and blank measurements (100 µl extraction buffer + 200 µl

reaction buffer) were performed to correct for any nonenzymatic hydrolysis of 4-MUG. Corrected

GUS activities (4-MU fluorescence units [FU] min^{-1}) were calculated as: $A_{GUS} = \Delta F / \Delta t = (F1-F0) /$

Δt where ΔF is the difference in fluorescence intensities measured in the linear range of the

fluorescence curves over a 15 min period of time (Δt). Values were normalized to 1 µg of protein

content by using the protein concentration determined via Bradford assay. Two to 5 independent

experiments were performed with at least 3 leaves from 2 different plants analyzed.

Data visualization and statistical analysis

Alignment of multiple protein sequences was performed using Jalview (version 2.11.2.6) (Procter et

al., 2021). Statistical analyses and generation of figures were performed using the GraphPad Prism

software (version 9.5.1) (GraphPad Software Inc.). Normal distribution of data was tested using the

Shapiro-Wilk and Kolmogorov-Smirnov normality tests. A Brown-Forsythe and Welch ANOVA test

followed by Dunnett's multiple T3 comparisons or an unpaired t-test with Welch's correction were

applied as parametric tests. Kruskal-Wallis followed by Dunn's multiple comparisons or Mann-

Whitney tests were applied as non-parametric tests. Boxes in the box-and-whiskers plots extend from

the 25th to 75th percentiles and whiskers range from the smallest to the largest values. Middle

horizontal lines are plotted at the median. Sample size n, statistical tests and significance levels are

provided in the figure legends. Source data and results of the statistical analyses are provided as

supplement files associated to the corresponding figure.

***M. truncatula* gene identifiers**

Accession numbers for the *M. truncatula* (Medicago) genes used in this study are provided from the

following genome portals, Phytozome (Mt4.0v1 version; https://phytozome-next.jgi.doe.gov/;

Goodstein et al., 2012) and / Medicago truncatula A17 r5.0 (A17 r5.1.9 version;

https://medicago.toulouse.inra.fr/MtrunA17r5.0-ANR/; Pecrix et al., 2018), respectively: *H3.1* (1)

(Medtr8g092720 / MtrunA17_Chr8g0383781); *H3.1* (2) (Medtr8g103245 /

MtrunA17_Chr8g0390361); *H3.3* (Medtr4g097175 / MtrunA17_Chr4g0054161); *H4*

(Medtr7g099610 / MtrunA17_Chr7g0262791); *KNOLLE* (Medtr5g012010 /

MtrunA17_Chr5g0398891); *CENH3* (Medtr8g027840 / MtrunA17_Chr8g0347231); repressor-type

MYB3R (Medtr7g061330 / MtrunA17_Chr7g0238631); *NF-YA1* (Medtr1g056530 /

MtrunA17_Chr1g0177091).

DATA AVAILABILITY STATEMENT

All data generated and analyzed during this study are included in the manuscript and supporting files.

All sequences have been obtained from publicly available repositories and indicated in the Materials

and Methods section.

MATERIALS AVAILABILITY STATEMENT

All material generated within this study can be obtained from the corresponding authors upon request.

Plasmids obtained from other laboratories as indicated in the respective sections need to be requested

from the original producer.

ACKNOWLEDGEMENTS

We would like to thank the entire team for the constant and fruitful input into the project and

especially Eija Schulze and Carmen Schubert for their excellent experimental support. Furthermore,

we greatly appreciated the technical help of Zenglin Li and Philipp Schwenk (University of Freiburg,

Germany) on GUS fluorometric assays. Jean Keller (LRSV Toulouse, France now Max-Planck

Institute for Chemical Ecology, Jena, Germany) conducted a comprehensive phylogeny on MYB3R,

MEME and FIMO searches on MSA motifs. Katharina Schiessl (University of Cambridge, UK)

provided the *nf-ya1-1* seeds and fruitful comments throughout the project. We also thank the staff of

the Life Imaging Center (LIC) in the Hilde Mangold House (HMH) of the Albert-Ludwigs-University

of Freiburg for the help with their confocal microscopy resources, and the excellent support in image

recording. The microscopes are operated by the Microscopy and Image Analysis Platform (MIAP)

and the Life Imaging Center (LIC), Freiburg.

FUNDING

Engineering Nitrogen Symbiosis for Africa (ENSA) project currently supported through a grant to

the University of Cambridge by the Bill & Melinda Gates Foundation (OPP1172165) and UK

government's Department for International Development (DFID) (TO)

Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) 431626755 (TO)

DFG under Germany's Excellence Strategy grant CIBSS – EXC-2189 – Project ID 39093984 (TO)

China Scholarship Council (CSC) grant 201708080016 (CS)

DFG project number 414136422 (CLSM; TO), DFG project number 426849454 (TEM; TO)

AUTHOR CONTRIBUTIONS

Conceptualization: MB, TL, TO; Investigation: MB, BL, NZ, CS, SE, PK, JS, FAD; Writing –

Original Draft: MB, TO; Writing –Review & Editing: MB, BL, NZ, CS, SE, PK, JS, FAD, TL, TO;

Supervision: MB, TL, TO; Project administration: TO; Funding Acquisition: TL, TO

REFERENCES

Araki, S., Ito, M., Soyano, T., Nishihama, R., & Machida, Y. (2004). Mitotic cyclins stimulate the

activity of c-Myb-like factors for transactivation of G2/M phase-specific genes in tobacco. *J

Biol Chem*, *279*(31), 32979-32988. https://doi.org/10.1074/jbc.M403171200

Bardetti, P., Castanheira, S. M., Valerius, O., Braus, G. H., & Perez-Martin, J. (2019). Cytoplasmic

retention and degradation of a mitotic inducer enable plant infection by a pathogenic fungus.

Elife, *8*. https://doi.org/10.7554/eLife.48943

Baudin, M., Laloum, T., Lepage, A., Ripodas, C., Ariel, F., Frances, L., Crespi, M., Gamas, P.,
Blanco, F. A., Zanetti, M. E., de Carvalho-Niebel, F., & Niebel, A. (2015). A Phylogenetically
Conserved Group of Nuclear Factor-Y Transcription Factors Interact to Control Nodulation
in Legumes. *Plant Physiol*, *169*(4), 2761-2773. https://doi.org/10.1104/pp.15.01144

Berckmans, B., & De Veylder, L. (2009). Transcriptional control of the cell cycle. *Curr Opin Plant*
Biol, *12*(5), 599-605. https://doi.org/10.1016/j.pbi.2009.07.005

Bhosale, R., Boudolf, V., Cuevas, F., Lu, R., Eekhout, T., Hu, Z., Van Isterdael, G., Lambert, G. M.,
Xu, F., Nowack, M. K., Smith, R. S., Vercauteren, I., De Rycke, R., Storme, V., Beeckman,
T., Larkin, J. C., Kremer, A., Hofte, H., Galbraith, D. W., … De Veylder, L. (2018). A
Spatiotemporal DNA Endoploidy Map of the Arabidopsis Root Reveals Roles for the
Endocycle in Root Development and Stress Adaptation. *Plant Cell*, *30*(10), 2330-2351.
https://doi.org/10.1105/tpc.17.00983

Binder, A., Lambert, J., Morbitzer, R., Popp, C., Ott, T., Lahaye, T., & Parniske, M. (2014). A
modular plasmid assembly kit for multigene expression, gene silencing and silencing rescue
in plants. *PLoS One*, *9*(2), e88218. https://doi.org/10.1371/journal.pone.0088218

Boisson-Dernier, A., Chabaud, M., Garcia, F., Becard, G., Rosenberg, C., & Barker, D. G. (2001).
Agrobacterium rhizogenes-transformed roots of Medicago truncatula for the study of
nitrogen-fixing and endomycorrhizal symbiotic associations. *Mol Plant Microbe Interact*,
14(6), 695-700. https://doi.org/10.1094/MPMI.2001.14.6.695

Bolognese, F., Wasner, M., Dohna, C. L., Gurtner, A., Ronchi, A., Muller, H., Manni, I., Mossner,
J., Piaggio, G., Mantovani, R., & Engeland, K. (1999). The cyclin B2 promoter depends on
NF-Y, a trimer whose CCAAT-binding activity is cell-cycle regulated. *Oncogene*, *18*(10),
1845-1853. https://doi.org/10.1038/sj.onc.1202494

Breakspear, A., Liu, C., Roy, S., Stacey, N., Rogers, C., Trick, M., Morieri, G., Mysore, K. S., Wen,
J., Oldroyd, G. E., Downie, J. A., & Murray, J. D. (2014). The root hair « infectome » of
Medicago truncatula uncovers changes in cell cycle genes and reveals a requirement for Auxin
signaling in rhizobial infection. *Plant Cell*, *26*(12), 4680-4701.
https://doi.org/10.1105/tpc.114.133496

Bu, F., Rutten, L., Roswanjaya, Y. P., Kulikova, O., Rodriguez-Franco, M., Ott, T., Bisseling, T.,
van Zeijl, A., & Geurts, R. (2020). Mutant analysis in the nonlegume Parasponia andersonii
identifies NIN and NF-YA1 transcription factors as a core genetic network in nitrogen-fixing
nodule symbioses. *New Phytol*, *226*(2), 541-554. https://doi.org/10.1111/nph.16386

Carotenuto, G., Sciascia, I., Oddi, L., Volpe, V., & Genre, A. (2019b). Size matters : Three methods
for estimating nuclear size in mycorrhizal roots of Medicago truncatula by image analysis.
BMC Plant Biol, *19*(1), 180. https://doi.org/10.1186/s12870-019-1791-1

Carotenuto, G., Volpe, V., Russo, G., Politi, M., Sciascia, I., de Almeida-Engler, J., & Genre, A.
(2019a). Local endoreduplication as a feature of intracellular fungal accommodation in
arbuscular mycorrhizas. *New Phytol*, *223*(1), 430-446. https://doi.org/10.1111/nph.15763

Carrere, S., Verdier, J., & Gamas, P. (2021). MtExpress, a Comprehensive and Curated RNAseq-
based Gene Expression Atlas for the Model Legume Medicago truncatula. *Plant Cell Physiol*,
62(9), 1494-1500. https://doi.org/10.1093/pcp/pcab110

Cathebras, C., Gong, X., Andrade, R. E., Vondenhoff, K., Keller, J., Delaux, P.-M., Hayashi, M.,
Griesmann, M., & Parniske, M. (2022). A novel cis-element enabled bacterial uptake by plant
cells (p. 2022.03.28.486070). *bioRxiv*. https://doi.org/10.1101/2022.03.28.486070

Cebolla, A., Vinardell, J. M., Kiss, E., Olah, B., Roudier, F., Kondorosi, A., & Kondorosi, E. (1999).
The mitotic inhibitor ccs52 is required for endoreduplication and ploidy-dependent cell
enlargement in plants. *EMBO J*, *18*(16), 4476-4484.
https://doi.org/10.1093/emboj/18.16.4476

Chen, P., Takatsuka, H., Takahashi, N., Kurata, R., Fukao, Y., Kobayashi, K., Ito, M., & Umeda, M.
(2017). Arabidopsis R1R2R3-Myb proteins are essential for inhibiting cell division in
response to DNA damage. *Nat Commun*, *8*(1), Article 1. https://doi.org/10.1038/s41467-017-
00676-4

Colón-Carmona, A., You, R., Haimovitch-Gal, T., & Doerner, P. (1999). Technical advance : Spatio-
temporal analysis of mitotic activity with a labile cyclin-GUS fusion protein. *Plant J*, *20*(4),
503-508. https://doi.org/10.1046/j.1365-313x.1999.00620.x

Combier, J. P., Frugier, F., de Billy, F., Boualem, A., El-Yahyaoui, F., Moreau, S., Vernie, T., Ott,
T., Gamas, P., Crespi, M., & Niebel, A. (2006). MtHAP2-1 is a key transcriptional regulator
of symbiotic nodule development regulated by microRNA169 in Medicago truncatula. *Genes
Dev*, *20*(22), 3084-3088. https://doi.org/10.1101/gad.402806

Cosolo, A., Jaiswal, J., Csordas, G., Grass, I., Uhlirova, M., & Classen, A. K. (2019). JNK-dependent
cell cycle stalling in G2 promotes survival and senescence-like phenotypes in tissue stress.
Elife, *8*. https://doi.org/10.7554/eLife.41036

Costa, S. (2017). Are division plane determination and cell-cycle progression coordinated? *New
Phytol*, *213*(1), 16-21. https://doi.org/10.1111/nph.14261

Davarinejad, H., Huang, Y. C., Mermaz, B., LeBlanc, C., Poulet, A., Thomson, G., Joly, V., Munoz,
M., Arvanitis-Vigneault, A., Valsakumar, D., Villarino, G., Ross, A., Rotstein, B. H., Alarcon,

E. I., Brunzelle, J. S., Voigt, P., Dong, J., Couture, J. F., & Jacob, Y. (2022). The histone H3.1 variant regulates TONSOKU-mediated DNA repair during replication. *Science*, *375*(6586), 1281-1286. https://doi.org/10.1126/science.abm5320

De Storme, N., Keceli, B. N., Zamariola, L., Angenon, G., & Geelen, D. (2016). CENH3-GFP: a visual marker for gametophytic and somatic ploidy determination in Arabidopsis thaliana. *BMC Plant Biol*, *16*, 1. https://doi.org/10.1186/s12870-015-0700-5

De Veylder, L., Larkin, J. C., & Schnittger, A. (2011). Molecular control and function of endoreplication in development and physiology. *Trends Plant Sci*, *16*(11), 624-634. https://doi.org/10.1016/j.tplants.2011.07.001

Desvoyes, B., Arana-Echarri, A., Barea, M. D., & Gutierrez, C. (2020). A comprehensive fluorescent sensor for spatiotemporal cell cycle analysis in Arabidopsis. *Nat Plants*, *6*(11), 1330-1334. https://doi.org/10.1038/s41477-020-00770-4

Desvoyes, B., Noir, S., Masoud, K., López, M. I., Genschik, P., & Gutierrez, C. (2019). FBL17 targets CDT1a for degradation in early S-phase to prevent Arabidopsis genome instability (p. 774109). *bioRxiv*. https://doi.org/10.1101/774109

Dewitte, W., Riou-Khamlichi, C., Scofield, S., Healy, J. M. S., Jacqmard, A., Kilby, N. J., & Murray, J. A. H. (2003). Altered Cell Cycle Distribution, Hyperplasia, and Inhibited Differentiation in Arabidopsis Caused by the D-Type Cyclin CYCD3. *Plant Cell*, *15*(1), 79-92. https://doi.org/10.1105/tpc.004838

Di Agostino, S., Strano, S., Emiliozzi, V., Zerbini, V., Mottolese, M., Sacchi, A., Blandino, G., & Piaggio, G. (2006). Gain of function of mutant p53 : The mutant p53/NF-Y protein complex reveals an aberrant transcriptional mechanism of cell cycle regulation. *Cancer Cell*, *10*(3), 191-202. https://doi.org/10.1016/j.ccr.2006.08.013

Dong, W., Zhu, Y., Chang, H., Wang, C., Yang, J., Shi, J., Gao, J., Yang, W., Lan, L., Wang, Y., Zhang, X., Dai, H., Miao, Y., Xu, L., He, Z., Song, C., Wu, S., Wang, D., Yu, N., & Wang, E. (2021). An SHR-SCR module specifies legume cortical cell fate to enable nodulation. *Nature*, *589*(7843), 586-590. https://doi.org/10.1038/s41586-020-3016-z

Doyle, J. J., & Coate, J. E. (2019). Polyploidy, the Nucleotype, and Novelty : The Impact of Genome Doubling on the Biology of the Cell. *Int J Plant Sci*, *180*(1), 1-52. https://doi.org/10.1086/700636

Draizen, E. J., Shaytan, A. K., Marino-Ramirez, L., Talbert, P. B., Landsman, D., & Panchenko, A. R. (2016). HistoneDB 2.0 : A histone database with variants—An integrated resource to explore histones and their variants. *Database (Oxford)*, *2016*. https://doi.org/10.1093/database/baw014

Echevarría, C., Gutierrez, C., & Desvoyes, B. (2021). Tools for Assessing Cell-Cycle Progression in Plants. *Plant and Cell Physiology*, *62*(8), 1231-1238. https://doi.org/10.1093/pcp/pcab066

Edgar, B. A., Zielke, N., & Gutierrez, C. (2014). Endocycles : A recurrent evolutionary innovation for post-mitotic cell growth. *Nat Rev Mol Cell Biol*, *15*(3), 197-210. https://doi.org/10.1038/nrm3756

Feng, J., Lee, T., Schiessl, K., & Oldroyd, G. E. D. (2021). Processing of NODULE INCEPTION controls the transition to nitrogen fixation in root nodules. *Science*, *374*(6567), 629-632. https://doi.org/10.1126/science.abg2804

Foucher, F., & Kondorosi, E. (2000). Cell cycle regulation in the course of nodule organogenesis in Medicago. *Plant Mol Biol*, *43*(5-6), 773-786. https://doi.org/10.1023/a:1006405029600

Fournier, J., Teillet, A., Chabaud, M., Ivanov, S., Genre, A., Limpens, E., de Carvalho-Niebel, F., & Barker, D. G. (2015). Remodeling of the infection chamber before infection thread formation reveals a two-step mechanism for rhizobial entry into the host legume root hair. *Plant Physiol*, *167*(4), 1233-1242. https://doi.org/10.1104/pp.114.253302

Fulop, K., Tarayre, S., Kelemen, Z., Horvath, G., Kevei, Z., Nikovics, K., Bako, L., Brown, S., Kondorosi, A., & Kondorosi, E. (2005). Arabidopsis anaphase-promoting complexes : Multiple activators and wide range of substrates might keep APC perpetually busy. *Cell Cycle*, *4*(8), 1084-1092.

Gao, J. P., Jiang, S., Su, Y., Xu, P., Wang, J., Liang, W., Liu, C. W., & Murray, J. D. (2022). Intracellular infection by symbiotic bacteria requires the mitotic kinase AURORA1. *Proc Natl Acad Sci U S A*, *119*(43), e2202606119. https://doi.org/10.1073/pnas.2202606119

Gaudioso-Pedraza, R., Beck, M., Frances, L., Kirk, P., Ripodas, C., Niebel, A., Oldroyd, G. E. D., Benitez-Alfonso, Y., & de Carvalho-Niebel, F. (2018). Callose-Regulated Symplastic Communication Coordinates Symbiotic Root Nodule Development. *Curr Biol*, *28*(22), 3562-3577 e6. https://doi.org/10.1016/j.cub.2018.09.031

Gauthier-Coles, C., White, R. G., & Mathesius, U. (2018). Nodulating Legumes Are Distinguished by a Sensitivity to Cytokinin in the Root Cortex Leading to Pseudonodule Development. *Front Plant Sci*, *9*, 1901. https://doi.org/10.3389/fpls.2018.01901

Gigli-Bisceglia, N., Engelsdorf, T., Strnad, M., Vaahtera, L., Khan, G. A., Yamoune, A., Alipanah, L., Novak, O., Persson, S., Hejatko, J., & Hamann, T. (2018). Cell wall integrity modulates Arabidopsis thaliana cell cycle gene expression in a cytokinin- and nitrate reductase-dependent manner. *Development*, *145*(19). https://doi.org/10.1242/dev.166678

Gleason, C., Chaudhuri, S., Yang, T., Munoz, A., Poovaiah, B. W., & Oldroyd, G. E. (2006).

Nodulation independent of rhizobia induced by a calcium-activated kinase lacking

autoinhibition. *Nature*, *441*(7097), 1149-1152. https://doi.org/10.1038/nature04812

Goodstein, D. M., Shu, S., Howson, R., Neupane, R., Hayes, R. D., Fazo, J., Mitros, T., Dirks, W.,

Hellsten, U., Putnam, N., & Rokhsar, D. S. (2012). Phytozome : A comparative platform for

green plant genomics. *Nucleic Acids Res*, *40*(Database issue), D1178-86.

https://doi.org/10.1093/nar/gkr944

Guan, D., Stacey, N., Liu, C., Wen, J., Mysore, K. S., Torres-Jerez, I., Vernié, T., Tadege, M., Zhou,

C., Wang, Z., Udvardi, M. K., Oldroyd, G. E. D., & Murray, J. D. (2013). Rhizobial Infection

Is Associated with the Development of Peripheral Vasculature in Nodules of Medicago

truncatula1[W][OA]. *Plant Physiol*, *162*(1), 107-115. https://doi.org/10.1104/pp.113.215111

Haga, N., Kato, K., Murase, M., Araki, S., Kubo, M., Demura, T., Suzuki, K., Muller, I., Voss, U.,

Jurgens, G., & Ito, M. (2007). R1R2R3-Myb proteins positively regulate cytokinesis through

activation of KNOLLE transcription in Arabidopsis thaliana. *Development*, *134*(6),

1101-1110. https://doi.org/10.1242/dev.02801

Hayashi, T., Banba, M., Shimoda, Y., Kouchi, H., Hayashi, M., & Imaizumi-Anraku, H. (2010). A

dominant function of CCaMK in intracellular accommodation of bacterial and fungal

endosymbionts. *Plant J*, *63*(1), 141-154. https://doi.org/10.1111/j.1365-313X.2010.04228.x

Hernandez-Lagana, E., & Autran, D. (2020). H3.1 Eviction Marks Female Germline Precursors in

Arabidopsis. *Plants*, *9*(10), Article 10. https://doi.org/10.3390/plants9101322

Hernandez-Lagana, E., Mosca, G., Mendocilla-Sato, E., Pires, N., Frey, A., Giraldo-Fonseca, A.,

Michaud, C., Grossniklaus, U., Hamant, O., Godin, C., Boudaoud, A., Grimanelli, D., Autran,

D., & Baroux, C. (2021). Organ geometry channels reproductive cell fate in the Arabidopsis

ovule primordium. *Elife*, *10*. https://doi.org/10.7554/eLife.66031

Hirsch, C. D., Wu, Y., Yan, H., & Jiang, J. (2009). Lineage-specific adaptive evolution of the

centromeric protein CENH3 in diploid and allotetraploid Oryza species. *Mol Biol Evol*,

26(12), 2877-2885. https://doi.org/10.1093/molbev/msp208

Hossain, M. S., Shrestha, A., Zhong, S., Miri, M., Austin, R. S., Sato, S., Ross, L., Huebert, T.,

Tromas, A., Torres-Jerez, I., Tang, Y., Udvardi, M., Murray, J. D., & Szczyglowski, K.

(2016). Lotus japonicus NF-YA1 Plays an Essential Role During Nodule Differentiation and

Targets Members of the SHI/STY Gene Family. *Mol Plant Microbe Interact*, *29*(12), 950-964.

https://doi.org/10.1094/MPMI-10-16-0206-R

Imbriano, C., Gurtner, A., Cocchiarella, F., Di Agostino, S., Basile, V., Gostissa, M., Dobbelstein,

M., Del Sal, G., Piaggio, G., & Mantovani, R. (2005). Direct p53 transcriptional repression :

In vivo analysis of CCAAT-containing G2/M promoters. *Mol Cell Biol*, *25*(9), 3737-3751.
https://doi.org/10.1128/MCB.25.9.3737-3751.2005

Ingouff, M., Rademacher, S., Holec, S., Soljic, L., Xin, N., Readshaw, A., Foo, S. H., Lahouze, B.,
Sprunck, S., & Berger, F. (2010). Zygotic resetting of the HISTONE 3 variant repertoire
participates in epigenetic reprogramming in Arabidopsis. *Curr Biol*, *20*(23), 2137-2143.
https://doi.org/10.1016/j.cub.2010.11.012

Ito, M., Araki, S., Matsunaga, S., Itoh, T., Nishihama, R., Machida, Y., Doonan, J. H., & Watanabe,
A. (2001). G2/M-phase-specific transcription during the plant cell cycle is mediated by c-
Myb-like transcription factors. *Plant Cell*, *13*(8), 1891-1905.
https://doi.org/10.1105/tpc.010102

Iwata, E., Ikeda, S., Matsunaga, S., Kurata, M., Yoshioka, Y., Criqui, M. C., Genschik, P., & Ito, M.
(2011). GIGAS CELL1, a novel negative regulator of the anaphase-promoting
complex/cyclosome, is required for proper mitotic progression and cell fate determination in
Arabidopsis. *Plant Cell*, *23*(12), 4382-4393. https://doi.org/10.1105/tpc.111.092049

Jiang, D., & Berger, F. (2017). DNA replication–coupled histone modification maintains Polycomb
gene silencing in plants. *Science*, *357*(6356), 1146-1149.
https://doi.org/10.1126/science.aan4965

Karimi-Ashtiyani, R., Ishii, T., Niessen, M., Stein, N., Heckmann, S., Gurushidze, M., Banaei-
Moghaddam, A. M., Fuchs, J., Schubert, V., Koch, K., Weiss, O., Demidov, D., Schmidt, K.,
Kumlehn, J., & Houben, A. (2015). Point mutation impairs centromeric CENH3 loading and
induces haploid plants. *Proc Natl Acad Sci U S A*, *112*(36), 11211-11216.
https://doi.org/10.1073/pnas.1504333112

Knaack, S. A., Conde, D., Chakraborty, S., Balmant, K. M., Irving, T. B., Maia, L. G. S., Triozzi, P.
M., Dervinis, C., Pereira, W. J., Maeda, J., Schmidt, H. W., Ane, J. M., Kirst, M., & Roy, S.
(2022). Temporal change in chromatin accessibility predicts regulators of nodulation in
Medicago truncatula. *BMC Biol*, *20*(1), 252. https://doi.org/10.1186/s12915-022-01450-9

Kobayashi, K., Suzuki, T., Iwata, E., Nakamichi, N., Suzuki, T., Chen, P., Ohtani, M., Ishida, T.,
Hosoya, H., Müller, S., Leviczky, T., Pettkó-Szandtner, A., Darula, Z., Iwamoto, A., Nomoto,
M., Tada, Y., Higashiyama, T., Demura, T., Doonan, J. H., … Ito, M. (2015). Transcriptional
repression by MYB3R proteins regulates plant organ growth. *The EMBO Journal*, *34*(15),
1992-2007. https://doi.org/10.15252/embj.201490899

Kondorosi, E., & Kondorosi, A. (2004). Endoreduplication and activation of the anaphase-promoting
complex during symbiotic cell development. *FEBS Lett*, *567*(1), 152-157.
https://doi.org/10.1016/j.febslet.2004.04.075

Lace, B., Su, C., Invernot Perez, D., Rodriguez-Franco, M., Vernie, T., Batzenschlager, M., Egli, S.,
Liu, C. W., & Ott, T. (2023). RPG acts as a central determinant for infectosome formation
and cellular polarization during intracellular rhizobial infections. *Elife*, *12*.
https://doi.org/10.7554/eLife.80741

Laloum, T., Baudin, M., Frances, L., Lepage, A., Billault-Penneteau, B., Cerri, M. R., Ariel, F.,
Jardinaud, M.-F., Gamas, P., de Carvalho-Niebel, F., & Niebel, A. (2014). Two CCAAT-box-
binding transcription factors redundantly regulate early steps of the legume-rhizobia
endosymbiosis. *Plant J*, *79*(5), 757-768. https://doi.org/10.1111/tpj.12587

Laporte, P., Lepage, A., Fournier, J., Catrice, O., Moreau, S., Jardinaud, M. F., Mun, J. H., Larrainzar,
E., Cook, D. R., Gamas, P., & Niebel, A. (2014). The CCAAT box-binding transcription
factor NF-YA1 controls rhizobial infection. *J Exp Bot*, *65*(2), 481-494.
https://doi.org/10.1093/jxb/ert392

Lermontova, I., Rutten, T., & Schubert, I. (2011). Deposition, turnover, and release of CENH3 at
Arabidopsis centromeres. *Chromosoma*, *120*(6), 633-640. https://doi.org/10.1007/s00412-
011-0338-5

Lermontova, I., Schubert, V., Bornke, F., Macas, J., & Schubert, I. (2007). Arabidopsis CBF5
interacts with the H/ACA snoRNP assembly factor NAF1. *Plant Mol Biol*, *65*(5), 615-626.
https://doi.org/10.1007/s11103-007-9226-z

Lermontova, I., Schubert, V., Fuchs, J., Klatte, S., Macas, J., & Schubert, I. (2006). Loading of
Arabidopsis centromeric histone CENH3 occurs mainly during G2 and requires the presence
of the histone fold domain. *Plant Cell*, *18*(10), 2443-2451.
https://doi.org/10.1105/tpc.106.043174

Liang, J., Klingl, A., Lin, Y.-Y., Boul, E., Thomas-Oates, J., & Marín, M. (2019). A subcompatible
rhizobium strain reveals infection duality in Lotus. *J Exp Bot*, *70*(6), 1903-1913.
https://doi.org/10.1093/jxb/erz057

Libbenga, K. R., & Torrey, J. G. (1973). Hormone-Induced Endoreduplication Prior to Mitosis in
Cultured Pea Root Cortex Cells. *Am J Bot*, *60*(4), 293-299. https://doi.org/10.2307/2441195

Linhart, C., Elkon, R., Shiloh, Y., & Shamir, R. (2005). Deciphering transcriptional regulatory
elements that encode specific cell cycle phasing by comparative genomics analysis. *Cell*
Cycle, *4*(12), 1788-1797. https://doi.org/10.4161/cc.4.12.2173

Liu, C. W., Breakspear, A., Stacey, N., Findlay, K., Nakashima, J., Ramakrishnan, K., Liu, M., Xie,
F., Endre, G., de Carvalho-Niebel, F., Oldroyd, G. E. D., Udvardi, M. K., Fournier, J., &
Murray, J. D. (2019a). A protein complex required for polar growth of rhizobial infection
threads. *Nat Commun*, *10*(1), 2848. https://doi.org/10.1038/s41467-019-10029-y

Liu, H., Lin, J.-S., Luo, Z., Sun, J., Huang, X., Yang, Y., Xu, J., Wang, Y.-F., Zhang, P., Oldroyd, G.
E. D., & Xie, F. (2022). Constitutive activation of a nuclear-localized calcium channel
complex in Medicago truncatula. *Proc Natl Acad Sci U S A*, *119*(34), e2205920119.
https://doi.org/10.1073/pnas.2205920119

Liu, J., Rutten, L., Limpens, E., van der Molen, T., van Velzen, R., Chen, R., Chen, Y., Geurts, R.,
Kohlen, W., Kulikova, O., & Bisseling, T. (2019b). A Remote cis-Regulatory Region Is
Required for NIN Expression in the Pericycle to Initiate Nodule Primordium Formation in
Medicago truncatula. *Plant Cell*, *31*(1), 68-83. https://doi.org/10.1105/tpc.18.00478

Lu, L., Chen, X., Qian, S., & Zhong, X. (2018). The plant-specific histone residue Phe41 is important
for genome-wide H3.1 distribution. *Nat Commun*, *9*(1), Article 1.
https://doi.org/10.1038/s41467-018-02976-9

Ma, Y., Jonsson, K., Aryal, B., De Veylder, L., Hamant, O., & Bhalerao, R. P. (2022).
Endoreplication mediates cell size control via mechanochemical signaling from cell wall. *Sci*
Adv, *8*(49), eabq2047. https://doi.org/10.1126/sciadv.abq2047

Madsen, L. H., Tirichine, L., Jurkiewicz, A., Sullivan, J. T., Heckmann, A. B., Bek, A. S., Ronson,
C. W., James, E. K., & Stougaard, J. (2010). The molecular network governing nodule
organogenesis and infection in the model legume Lotus japonicus. *Nat Commun*, *1*(1), 10.
https://doi.org/10.1038/ncomms1009

Manni, I., Mazzaro, G., Gurtner, A., Mantovani, R., Haugwitz, U., Krause, K., Engeland, K., Sacchi,
A., Soddu, S., & Piaggio, G. (2001). NF-Y mediates the transcriptional inhibition of the cyclin
B1, cyclin B2, and cdc25C promoters upon induced G2 arrest. *J Biol Chem*, *276*(8),
5570-5576. https://doi.org/10.1074/jbc.M006052200

Marsh, J. F., Rakocevic, A., Mitra, R. M., Brocard, L., Sun, J., Eschstruth, A., Long, S. R., Schultze,
M., Ratet, P., & Oldroyd, G. E. (2007). Medicago truncatula NIN is essential for rhizobial-
independent nodule organogenesis induced by autoactive calcium/calmodulin-dependent
protein kinase. *Plant Physiol*, *144*(1), 324-335. https://doi.org/10.1104/pp.106.093021

McCloy, R. A., Rogers, S., Caldon, C. E., Lorca, T., Castro, A., & Burgess, A. (2014). Partial
inhibition of Cdk1 in G 2 phase overrides the SAC and decouples mitotic events. *Cell Cycle*,
13(9), 1400-1412. https://doi.org/10.4161/cc.28401

Meserve, J. H., & Duronio, R. J. (2017). A population of G2-arrested cells are selected as sensory
organ precursors for the interommatidial bristles of the Drosophila eye. *Dev Biol*, *430*(2),
374-384. https://doi.org/10.1016/j.ydbio.2017.06.023

Meyer, H. M., Teles, J., Formosa-Jordan, P., Refahi, Y., San-Bento, R., Ingram, G., Jönsson, H.,
Locke, J. C. W., & Roeder, A. H. K. (2017). Fluctuations of the transcription factor ATML1

generate the pattern of giant cells in the Arabidopsis sepal. *eLife*, *6*, e19131. https://doi.org/10.7554/eLife.19131

Monroy-Morales, E., Davila-Delgado, R., Ayala-Guzman, E., Gamboa-deBuen, A., & Sanchez-Lopez, R. (2022). Visualization of the Crossroads between a Nascent Infection Thread and the First Cell Division Event in Phaseolus vulgaris Nodulation. *Int J Mol Sci*, *23*(9). https://doi.org/10.3390/ijms23095267

Mutterer, J., & Zinck, E. (2013). Quick-and-clean article figures with FigureJ. *J Microsc*, *252*(1), 89-91. https://doi.org/10.1111/jmi.12069

Myers, Z. A., & Holt, B. F. (2018). NUCLEAR FACTOR-Y: still complex after all these years? *Curr Opin Plant Biol*, *45*(Pt A), 96-102. https://doi.org/10.1016/j.pbi.2018.05.015

Nadzieja, M., Stougaard, J., & Reid, D. (2019). A Toolkit for High Resolution Imaging of Cell Division and Phytohormone Signaling in Legume Roots and Root Nodules. *Front Plant Sci*, *10*, 1000. https://doi.org/10.3389/fpls.2019.01000

Nagymihaly, M., Veluchamy, A., Gyorgypal, Z., Ariel, F., Jegu, T., Benhamed, M., Szucs, A., Kereszt, A., Mergaert, P., & Kondorosi, E. (2017). Ploidy-dependent changes in the epigenome of symbiotic cells correlate with specific patterns of gene expression. *Proc Natl Acad Sci U S A*, *114*(17), 4543-4548. https://doi.org/10.1073/pnas.1704211114

Negishi, T., Veis, J., Hollenstein, D., Sekiya, M., Ammerer, G., & Ohya, Y. (2016). The Late S-Phase Transcription Factor Hcm1 Is Regulated through Phosphorylation by the Cell Wall Integrity Checkpoint. *Mol Cell Biol*, *36*(6), 941-953. https://doi.org/10.1128/MCB.00952-15

Otero, S., Desvoyes, B., Peiro, R., & Gutierrez, C. (2016). Histone H3 Dynamics Reveal Domains with Distinct Proliferation Potential in the Arabidopsis Root. *Plant Cell*, *28*(6), 1361-1371. https://doi.org/10.1105/tpc.15.01003

Park, M., Krause, C., Karnahl, M., Reichardt, I., El Kasmi, F., Mayer, U., Stierhof, Y. D., Hiller, U., Strompen, G., Bayer, M., Kientz, M., Sato, M. H., Nishimura, M. T., Dangl, J. L., Sanderfoot, A. A., & Jurgens, G. (2018). Concerted Action of Evolutionarily Ancient and Novel SNARE Complexes in Flowering-Plant Cytokinesis. *Dev Cell*, *44*(4), 500-511 e4. https://doi.org/10.1016/j.devcel.2017.12.027

Parniske, M. (2018). Uptake of bacteria into living plant cells, the unifying and distinct feature of the nitrogen-fixing root nodule symbiosis. *Curr Opin Plant Biol*, *44*, 164-174. https://doi.org/10.1016/j.pbi.2018.05.016

Pecrix, Y., Staton, S. E., Sallet, E., Lelandais-Briere, C., Moreau, S., Carrere, S., Blein, T., Jardinaud, M. F., Latrasse, D., Zouine, M., Zahm, M., Kreplak, J., Mayjonade, B., Satge, C., Perez, M., Cauet, S., Marande, W., Chantry-Darmon, C., Lopez-Roques, C., … Gamas, P. (2018).

Whole-genome landscape of Medicago truncatula symbiotic genes. *Nat Plants*, *4*(12), 1017-1025. https://doi.org/10.1038/s41477-018-0286-7

Probst, A. V., Desvoyes, B., & Gutierrez, C. (2020). Similar yet critically different : The distribution, dynamics and function of histone variants. *J Exp Bot*, *71*(17), 5191-5204. https://doi.org/10.1093/jxb/eraa230

Procter, J. B., Carstairs, G. M., Soares, B., Mourao, K., Ofoegbu, T. C., Barton, D., Lui, L., Menard, A., Sherstnev, N., Roldan-Martinez, D., Duce, S., Martin, D. M. A., & Barton, G. J. (2021). Alignment of Biological Sequences with Jalview. *Methods Mol Biol*, *2231*, 203-224. https://doi.org/10.1007/978-1-0716-1036-7_13

Ried, M. K., Antolin-Llovera, M., & Parniske, M. (2014). Spontaneous symbiotic reprogramming of plant roots triggered by receptor-like kinases. *Elife*, *3*. https://doi.org/10.7554/eLife.03891

Rivera, H. E., & Davies, S. W. (2021). Symbiosis maintenance in the facultative coral, Oculina arbuscula, relies on nitrogen cycling, cell cycle modulation, and immunity. *Sci Rep*, *11*(1), 21226. https://doi.org/10.1038/s41598-021-00697-6

Rost, T. L. (2011). The organization of roots of dicotyledonous plants and the positions of control points. *Ann Bot*, *107*(7), 1213-1222. https://doi.org/10.1093/aob/mcq229

Roux, B., Rodde, N., Jardinaud, M. F., Timmers, T., Sauviac, L., Cottret, L., Carrere, S., Sallet, E., Courcelle, E., Moreau, S., Debelle, F., Capela, D., de Carvalho-Niebel, F., Gouzy, J., Bruand, C., & Gamas, P. (2014). An integrated analysis of plant and bacterial gene expression in symbiotic root nodules using laser-capture microdissection coupled to RNA sequencing. *Plant J*, *77*(6), 817-837. https://doi.org/10.1111/tpj.12442

Russo, G., Carotenuto, G., Fiorilli, V., Volpe, V., Chiapello, M., Van Damme, D., & Genre, A. (2019). Ectopic activation of cortical cell division during the accommodation of arbuscular mycorrhizal fungi. *New Phytol*, *221*(2), 1036-1048. https://doi.org/10.1111/nph.15398

Russo, G., & Genre, A. (2021). Divide and Be Conquered-Cell Cycle Reactivation in Arbuscular Mycorrhizal Symbiosis. *Front Plant Sci*, *12*, 753265. https://doi.org/10.3389/fpls.2021.753265

Sanei, M., Pickering, R., Kumke, K., Nasuda, S., & Houben, A. (2011). Loss of centromeric histone H3 (CENH3) from centromeres precedes uniparental chromosome elimination in interspecific barley hybrids. *Proc Natl Acad Sci U S A*, *108*(33), E498-505. https://doi.org/10.1073/pnas.1103190108

Schaefer, E., Belcram, K., Uyttewaal, M., Duroc, Y., Goussot, M., Legland, D., Laruelle, E., de Tauzia-Moreau, M. L., Pastuglia, M., & Bouchez, D. (2017). The preprophase band of

1186  microtubules controls the robustness of division orientation in plants. *Science*, *356*(6334),
1187    186-189. https://doi.org/10.1126/science.aal3016

1188 Schauser, L., Roussis, A., Stiller, J., & Stougaard, J. (1999). A plant regulator controlling
1189    development of symbiotic root nodules. *Nature*, *402*(6758), 191-195.
1190    https://doi.org/10.1038/46058

1191 Schiessl, K., Lee, T., Orvosova, M., Bueno-Batista, M., Stuer, N., Bailey, P. C., Mysore, K. S., Wen,
1192    J., & Oldroyd, G. E. D. (2023). *Light sensitive short hypocotyl (LSH) confer symbiotic nodule*
1193    *identity in the legume Medicago truncatula* (p. 2023.02.12.528179). bioRxiv.
1194    https://doi.org/10.1101/2023.02.12.528179

1195 Schiessl, K., Lilley, J. L. S., Lee, T., Tamvakis, I., Kohlen, W., Bailey, P. C., Thomas, A., Luptak, J.,
1196    Ramakrishnan, K., Carpenter, M. D., Mysore, K. S., Wen, J., Ahnert, S., Grieneisen, V. A.,
1197    & Oldroyd, G. E. D. (2019). NODULE INCEPTION Recruits the Lateral Root
1198    Developmental Program for Symbiotic Nodule Organogenesis in Medicago truncatula. *Curr*
1199    *Biol*, *29*(21), 3657-3668 e5. https://doi.org/10.1016/j.cub.2019.09.005

1200 Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S.,
1201    Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J. Y., White, D. J., Hartenstein, V., Eliceiri,
1202    K., Tomancak, P., & Cardona, A. (2012). Fiji : An open-source platform for biological-image
1203    analysis. *Nat Methods*, *9*(7), 676-682. https://doi.org/10.1038/nmeth.2019

1204 Schnittger, A., Schöbinger, U., Bouyer, D., Weinl, C., Stierhof, Y.-D., & Hülskamp, M. (2002).
1205    Ectopic D-type cyclin expression induces not only DNA replication but also cell division in
1206    Arabidopsis trichomes. *Proc Natl Acad Sci U S A*, *99*(9), 6410-6415.
1207    https://doi.org/10.1073/pnas.092657299

1208 Shi, L., Wang, J., Hong, F., Spector, D. L., & Fang, Y. (2011). Four amino acids guide the assembly
1209    or disassembly of Arabidopsis histone H3.3-containing nucleosomes. *Proc Natl Acad Sci U S*
1210    *A*, *108*(26), 10574-10578. https://doi.org/10.1073/pnas.1017882108

1211 Shimotohno, A., Aki, S. S., Takahashi, N., & Umeda, M. (2021). Regulation of the Plant Cell Cycle
1212    in Response to Hormones and the Environment. *Annu Rev Plant Biol*, *72*, 273-296.
1213    https://doi.org/10.1146/annurev-arplant-080720-103739

1214 Shrestha, A., Zhong, S., Therrien, J., Huebert, T., Sato, S., Mun, T., Andersen, S. U., Stougaard, J.,
1215    Lepage, A., Niebel, A., Ross, L., & Szczyglowski, K. (2021). Lotus japonicus Nuclear Factor
1216    YA1, a nodule emergence stage-specific regulator of auxin signalling. *New Phytol*, *229*(3),
1217    1535-1552. https://doi.org/10.1111/nph.16950

1218 Silhavy, D., Molnar, A., Lucioli, A., Szittya, G., Hornyik, C., Tavazza, M., & Burgyan, J. (2002). A
1219   viral protein suppresses RNA silencing and binds silencing-generated, 21- to 25-nucleotide
1220   double-stranded RNAs. *EMBO J*, *21*(12), 3070-3080. https://doi.org/10.1093/emboj/cdf312

1221 Singh, S., Katzer, K., Lambert, J., Cerri, M., & Parniske, M. (2014). CYCLOPS, a DNA-binding
1222   transcriptional activator, orchestrates symbiotic root nodule development. *Cell Host Microbe*,
1223   *15*(2), 139-152. https://doi.org/10.1016/j.chom.2014.01.011

1224 Smertenko, A., Assaad, F., Baluška, F., Bezanilla, M., Buschmann, H., Drakakaki, G., Hauser, M.-
1225   T., Janson, M., Mineyuki, Y., Moore, I., Müller, S., Murata, T., Otegui, M. S., Panteris, E.,
1226   Rasmussen, C., Schmit, A.-C., Šamaj, J., Samuels, L., Staehelin, L. A., … Žárský, V. (2017).
1227   Plant Cytokinesis : Terminology for Structures and Processes. *Trends Cell Biol*, *27*(12),
1228   885-894. https://doi.org/10.1016/j.tcb.2017.08.008

1229 Soni, R., Carmichael, J. P., Shah, Z. H., & Murray, J. A. (1995). A family of cyclin D homologs from
1230   plants differentially controlled by growth regulators and containing the conserved
1231   retinoblastoma protein interaction motif. *Plant Cell*, *7*(1), 85-103.
1232   https://doi.org/10.1105/tpc.7.1.85

1233 Soyano, T., Hirakawa, H., Sato, S., Hayashi, M., & Kawaguchi, M. (2014). Nodule Inception creates
1234   a long-distance negative feedback loop involved in homeostatic regulation of nodule organ
1235   production. *Proc Natl Acad Sci U S A*, *111*(40), 14607-14612.
1236   https://doi.org/10.1073/pnas.1412716111

1237 Soyano, T., Kouchi, H., Hirota, A., & Hayashi, M. (2013). Nodule inception directly targets NF-Y
1238   subunit genes to regulate essential processes of root nodule development in Lotus japonicus.
1239   *PLoS Genet*, *9*(3), e1003352. https://doi.org/10.1371/journal.pgen.1003352

1240 Su, C., Zhang, G., Rodriguez-Franco, M., Hinnenberg, R., Wietschorke, J., Liang, P., Yang, W.,
1241   Uhler, L., Li, X., & Ott, T. (2023). Transcellular progression of infection threads in Medicago
1242   truncatula roots is associated with locally confined cell wall modifications. *Curr Biol*, *33*(3),
1243   533-542 e5. https://doi.org/10.1016/j.cub.2022.12.051

1244 Suzaki, T., Ito, M., Yoro, E., Sato, S., Hirakawa, H., Takeda, N., & Kawaguchi, M. (2014).
1245   Endoreduplication-mediated initiation of symbiotic organ development in Lotus japonicus.
1246   *Development*, *141*(12), 2441-2445. https://doi.org/10.1242/dev.107946

1247 Takahashi, N., Ogita, N., Takahashi, T., Taniguchi, S., Tanaka, M., Seki, M., & Umeda, M. (2019).
1248   A regulatory module controlling stress-induced cell cycle arrest in Arabidopsis. *Elife*, *8*.
1249   https://doi.org/10.7554/eLife.43944

Talbert, P. B., Masuelli, R., Tyagi, A. P., Comai, L., & Henikoff, S. (2002). Centromeric localization

and adaptive evolution of an Arabidopsis histone H3 variant. *Plant Cell*, *14*(5), 1053-1066.

https://doi.org/10.1105/tpc.010425

Tek, A. L., Kashihara, K., Murata, M., & Nagaki, K. (2014). Identification of the centromere-specific

histone H3 variant in Lotus japonicus. *Gene*, *538*(1), 8-11.

https://doi.org/10.1016/j.gene.2014.01.034

Timmers, A. C., Auriac, M. C., & Truchet, G. (1999). Refined analysis of early symbiotic steps of

the Rhizobium-Medicago interaction in relationship with microtubular cytoskeleton

rearrangements. *Development*, *126*(16), 3617-3628. https://doi.org/10.1242/dev.126.16.3617

Torrey, J. G., & Barrios, S. (1969). Cytological Studies on Rhizobial Nodule Initiation in Pisum.

Caryologia, *22*(1), 47-61. https://doi.org/10.1080/00087114.1969.10796324

Trolet, A., Baldrich, P., Criqui, M. C., Dubois, M., Clavel, M., Meyers, B. C., & Genschik, P. (2019).

Cell Cycle-Dependent Regulation and Function of ARGONAUTE1 in Plants. *Plant Cell*,

31(8), 1734-1750. https://doi.org/10.1105/tpc.19.00069

Tsyganova, A. V., Brewin, N. J., & Tsyganov, V. E. (2021). Structure and Development of the

Legume-Rhizobial Symbiotic Interface in Infection Threads. *Cells*, *10*(5).

https://doi.org/10.3390/cells10051050

Ursache, R., Andersen, T. G., Marhavy, P., & Geldner, N. (2018). A protocol for combining

fluorescent proteins with histological stains for diverse cell wall components. *Plant J*, *93*(2),

399-412. https://doi.org/10.1111/tpj.13784

van Brussel, A. A., Bakhuizen, R., van Spronsen, P. C., Spaink, H. P., Tak, T., Lugtenberg, B. J., &

Kijne, J. W. (1992). Induction of pre-infection thread structures in the leguminous host plant

by mitogenic lipo-oligosaccharides of Rhizobium. *Science*, *257*(5066), 70-72.

https://doi.org/10.1126/science.257.5066.70

van Spronsen, P. C., Gronlund, M., Pacios Bras, C., Spaink, H. P., & Kijne, J. W. (2001). Cell

biological changes of outer cortical root cells in early determinate nodulation. *Mol Plant*

Microbe Interact, *14*(7), 839-847. https://doi.org/10.1094/MPMI.2001.14.7.839

Vinardell, J. M., Fedorova, E., Cebolla, A., Kevei, Z., Horvath, G., Kelemen, Z., Tarayre, S., Roudier,

F., Mergaert, P., Kondorosi, A., & Kondorosi, E. (2003). Endoreduplication mediated by the

anaphase-promoting complex activator CCS52A is required for symbiotic cell differentiation

in Medicago truncatula nodules. *Plant Cell*, *15*(9), 2093-2105.

https://doi.org/10.1105/tpc.014373

Weber, E., Engler, C., Gruetzner, R., Werner, S., & Marillonnet, S. (2011). A modular cloning system for standardized assembly of multigene constructs. *PloS One*, *6*(2), e16765. https://doi.org/10.1371/journal.pone.0016765

Weimer, A. K., Biedermann, S., Harashima, H., Roodbarkelari, F., Takahashi, N., Foreman, J., Guan, Y., Pochon, G., Heese, M., Van Damme, D., Sugimoto, K., Koncz, C., Doerner, P., Umeda, M., & Schnittger, A. (2016). The plant-specific CDKB1-CYCB1 complex mediates homologous recombination repair in Arabidopsis. *EMBO J*, *35*(19), 2068-2086. https://doi.org/10.15252/embj.201593083

Wildermuth, M. C., Steinwand, M. A., McRae, A. G., Jaenisch, J., & Chandran, D. (2017). Adapted Biotroph Manipulation of Plant Cell Ploidy. *Annu Rev Phytopathol*, *55*, 537-564. https://doi.org/10.1146/annurev-phyto-080516-035458

Wipf, L., & Cooper, D. C. (1938). Chromosome Numbers in Nodules and Roots of Red Clover, Common Vetch and Garden Pea. *Proc Natl Acad Sci U S A*, *24*(2), 87-91. https://doi.org/10.1073/pnas.24.2.87

Xiao, T. T., Schilderink, S., Moling, S., Deinum, E. E., Kondorosi, E., Franssen, H., Kulikova, O., Niebel, A., & Bisseling, T. (2014). Fate map of Medicago truncatula root nodules. *Development*, *141*(18), 3517-3528. https://doi.org/10.1242/dev.110775

Xiao, T. T., van Velzen, R., Kulikova, O., Franken, C., & Bisseling, T. (2019). Lateral root formation involving cell division in both pericycle, cortex and endodermis is a common and ancestral trait in seed plants. *Development*, *146*(20), dev182592. https://doi.org/10.1242/dev.182592

Xu, J., Lee, Y. J., & Liu, B. (2020). Establishment of a mitotic model system by transient expression of the D-type cyclin in differentiated leaf cells of tobacco (Nicotiana benthamiana). *New Phytol*, *226*(4), 1213-1220. https://doi.org/10.1111/nph.16309

Yang, W. C., de Blank, C., Meskiene, I., Hirt, H., Bakker, J., van Kammen, A., Franssen, H., & Bisseling, T. (1994). Rhizobium nod factors reactivate the cell cycle during infection and nodule primordium formation, but the cycle is only completed in primordium formation. *Plant Cell*, *6*(10), 1415-1426. https://doi.org/10.1105/tpc.6.10.1415

Yant, L., & Bomblies, K. (2015). Genome management and mismanagement—Cell-level opportunities and challenges of whole-genome duplication. *Genes Dev*, *29*(23), 2405-2419. https://doi.org/10.1101/gad.271072.115

Zanetti, M. E., Ripodas, C., & Niebel, A. (2017). Plant NF-Y transcription factors : Key players in plant-microbe interactions, root development and adaptation to stress. *Biochim Biophys Acta Gene Regul Mech*, *1860*(5), 645-654. https://doi.org/10.1016/j.bbagrm.2016.11.007

FIGURES

Figure 1. Medicago histone H3.1 patterns reveal sustained mitotic and endocycling activities in a symbiotic context.
(**A**) Schematic representation of histone H3.3-mCherry and H3.1-eGFP distribution (horizontal bars) and fluorescence intensity (grey saturation) throughout the different cell-cycle phases (adapted from Echevarria et al., 2021). H3.1 is predominantly expressed during S-phase and incorporated during DNA replication in proliferating (G1, S, G2, M) and endocycling (G1, S, G2) cells. H3.3 is constitutively produced. Nuclei of increasing size and DNA content are coloured according to their H3.3 (pink) and H3.1 (green) content in decondensed chromatin (G1, S, G2) or condensed chromosomes (M). (**B**) Confocal images of nodule sections isolated from WT transgenic roots inoculated with mCherry-producing *S. meliloti* (35 to 40 dpi), showing H3.1-eGFP (green) and H3.3-mCherry (magenta) localization across characteristic developmental zones. H3.1 accumulates in the meristematic zone (ZI) enriched in proliferating cells (late-anaphase chromosomes indicated by a filled arrowhead) and in regions with high endoreduplication activity (ZIId and ZIIp, distal and proximal parts of the infection zone) where rhizobia are released into membrane-bound compartments called symbiosomes. H3.1 is extensively replaced by H3.3 in the fixation zone (ZIII) where host cells and rhizobia complete their differentiation process. Images are maximum intensity projections except the top-right panel (single focal plane). Scale bars: left panel = 50 μm; right panels = 10 μm. Transformation experiments were repeated 3 times with a total of 9 nodules from 6 composite plants showing similar results. (**C**) Confocal images of whole-mount WT roots colonized by *R. irregularis* (15 dpi). Plant and fungal cell walls were stained with Calcofluor white (grayscale). H3.1 was enriched in chromocenters (heterochromatin foci in nuclei indicated by stars) and kept at high levels in the euchromatin (diffuse labeling in nuclei pointed by filled arrowheads) from neighbouring (upper panel) and early-arbusculated cells (lower panel) of the inner cortical tissue. The empty arrowhead points to a nucleus with a low H3.1 content in a fully-arbusculated, differentiated cell. Images are maximum intensity projections. Scale bars: 50 μm. Two independent transformation experiments were performed with 3 to 5 composite plants analyzed per replica.

Figure 2. Individual reprogramming for infection includes large-scale chromatin rearrangements.
(**A-B**) Confocal images of whole-mount WT roots expressing the pH3.1::H3.1-eGFP / pH3.3::H3.3-mCherry reporter and
inoculated with mCherry-producing *S. meliloti* (8 dpi). Images are maximum intensity projections (eGFP: green;
mCherry: magenta; Calcofluor white cell-wall staining: grayscale). (**A**) Left panel: the solid-line frame indicates the outer
and middle cortical regions of an early nodule primordium shown in the close up. Right panel: dotted-line regions of
interest (ROIs) indicate nuclei of infected cells (IC) fully passed by a cortical infection thread in the first (C1) and second
(C2) cortical layers. NC: neighbouring cell. The filled arrowhead points to the nucleus of a host cell from the middle
cortical layer (C3), which is penetrated by the progressing infection thread. Scale bars: 20 μm. (**B**) Left panel: dotted-line
ROIs indicate nuclei of infected cells (IC) being passed (C4) or recently passed (C3) by a cortical infection thread. Right
panel: dotted-line ROIs indicate nuclei of infected cells in the inner cortex (C4/5) of a nodule primordium with several
cell layers. Scale bars: 20 μm. (**C-D**) Quantification of the nuclear area (**C**) and corrected total H3.1-eGFP nuclear
fluorescence (**D**) at the equatorial plane in couples of neighbouring (NC) and infected cells (IC) from the same cortical
layer (C2 to C4, n = 27; see Materials and methods for more details). Roots from 2 to 7 composite plants with visible
signs of inner cortical cell division (*i.e.*, showing high H3.1-eGFP signal) were analyzed from 3 independent
transformation experiments. All data points are shown and crosses indicate sample means. Differences were statistically
significant (P values < 0.0001) using an unpaired t-test with Welch's correction (**C**) or a Mann-Whitney test (**D**). (**E**)
Schematic representation of histone H3.3-mCherry and H3.1-eGFP distribution (horizontal bars) and fluorescence
intensity (grey saturation) throughout the different cell-cycle phases. Nuclei of increasing size and DNA content are
coloured according to their merged H3.3 (magenta) and H3.1 (green) content in decondensed chromatin (G1, S, G2). This
diagram illustrates cells exiting the cell cycle within proliferating or endoreduplicating populations, which proceed to a
massive H3.1 eviction following their last DNA replication round (Otero et al., 2016).

Figure 3. H3.1 eviction coincides with cortical infection thread progression.
(**A**) Confocal images of whole-mount transgenic roots co-expressing H3.1-eGFP and H3.3-mCherry in three different genotypes: WT (11 dpi), *daphne-like* (14 dpi) and *nf-ya1-1* (12 dpi), inoculated with mCherry-producing *S. meliloti*. WT and *nf-ya1-1* plants initiate nodule formation (NOD[+]) whereas the *daphne-like* mutant is non-nodulating (NOD[-]). Numbers indicate the frequencies of observation of a low (WT) or high (*daphne-like*, *nf-ya1-1*) H3.1-eGFP content in the last infected cortical cell. Corresponding schematic representations below each panel depict cellular proliferation or endocycling activities indicated by nuclear H3.1 levels (low: magenta; high: green) on the cortical infection thread trajectory (C1 to C3; diffuse magenta or green colouring) and in inner cortical layers (C4 to C5). Rhizobia inside infection threads are depicted in red. Confocal images are maximum intensity projections and show merged fluorescent channels (eGFP: green; mCherry: magenta; Calcofluor white cell-wall staining: grayscale in the middle and right panels). Scale bars: 20 μm. IC: infected cell. Ep: epidermis. (**B-C**) Quantification of the nuclear area (**B**) and the relative corrected total H3.1-eGFP nuclear fluorescence (IC/NC) (**C**) at the equatorial plane in couples of neighbouring (NC) and infected cells (IC) from the same cortical layer in WT (C1 to C4), *daphne-like* (C1 to C2) and *nf-ya1-1* (C1 to C4) genetic backgrounds (n = 43, 19 and 39 nuclei, respectively). Different letters indicate statistically significant differences according to a Kruskal-Wallis test followed by Dunn's multiple comparisons test. (**B**) All data points are shown and crosses indicate sample means. (**C**) All data points are shown with magenta or green indicating relative H3.1-eGFP nuclear signals (IC/NC) below 1,5 or above 1,6, respectively. Horizontal bars indicate sample means with 95% confidence interval. Ten to 21 composite plants (8-14 dpi) from 2 (*daphne-like*, *nf-ya1-1*) to 7 (WT) independent transformation experiments were analyzed.

Figure 4. A tight control over host cells' mitotic commitment enables passage of the future nodule meristem.
(**A**) Schematic representation of Arabidopsis CDT1a-eCFP, H3.1-mCherry and N-CYCB1;1-YFP distribution (horizontal bars) and fluorescence intensity (grey saturation) throughout the different cell-cycle phases (adapted from Echevarria et al., 2021; see also Desvoyes et al., 2020). CDT1a-CFP accumulates in G1 and is rapidly degraded during the G1/S transition (blunt end bar). H3.1 is predominantly expressed during S-phase and incorporated during DNA replication. N-CYCB1;1-YFP is present in late G2 and mitotic cells and is completely degraded in anaphase (blunt end bar). Nuclei of increasing size and DNA content are coloured according to their CDT1a (sky blue), H3.1 (vermillion) and N-CYCB1;1 (yellow) content in/around decondensed chromatin (G1, S, G2) or condensed chromosomes (M). H3.1 levels decrease in differentiating cells (dotted-line box). (**B-C**) Confocal images of whole-mount Medicago WT roots expressing the Arabidopsis PlaCCI reporter in nodule primordia at 7 dpi with mCherry- (**B**) or GFP-producing *S. meliloti* (**C**). Images are maximum intensity projections and show merged fluorescent channels (eCFP and GFP: cyan; mCherry: red; YFP: yellow; Calcofluor white cell-wall staining: grayscale). (**B**) The filled arrowhead points to a cell of the C3 layer penetrated by an infection thread. Cell-cycle phases of non-infected neighbouring cells are indicated. (**C**) Left panel: the dotted-line ROI indicates the nucleus of an infected cell (IC) from the C3 layer being passed by a cortical infection thread. Numbers indicate the frequencies of observation of the absence of CDT1a- or N-CYCB1;1-associated signals in the nucleus of the last infected cortical cell. Right panels: fluorescence intensity profiles of CDT1a-, H3.1- and N-CYCB1.1-associated signals along the cyan and green transects shown in (**C**). Scale bars: 20 μm. (**D**) Left panels: confocal images of a whole-mount WT root co-expressing a destabilized triple-Venus nuclear reporter driven by the Arabidopsis *CYCB1;2* promoter (pCYCB1;2::N-CYCB1;2-NLS-3xVenus) and a nuclear-localized tandem-mCherry as a transformation marker, 12 dpi with mCherry-producing *S. meliloti*. Numbers indicate the frequencies of observation of the absence of the triple-Venus reporter signal in the nucleus of the last infected cortical cell. Two independent transformation experiments were performed with 3 to 4 composite plants analyzed per replica. (**E**) Left panels: confocal images of a whole-mount WT root co-expressing a transcriptional reporter of Medicago *KNOLLE* driving a nuclear-localized triple-Venus (pKNOLLE::NLS-3xVenus) together with a nuclear-localized tandem-mCherry as a transformation marker, 12 dpi with mCherry-producing *S. meliloti*. Numbers indicate the frequencies of observation of nodule primordia where the triple-Venus reporter signal is kept comparably low on the cortical infection thread trajectory. A total of 12 composite plants from 2 independent transformation experiments were analyzed. (**D-E**) Images are maximum intensity projections and show merged fluorescent channels (Venus: green; mCherry: magenta; Calcofluor white cell-wall staining: grayscale in

the upper panels). The Venus channel is shown in Green Fire Blue when isolated in lower panels, with blue or yellow indicating low or high fluorescence levels, respectively. The dotted-line ROIs indicate nuclei of infected cells (IC) from the C3 layer being passed by a cortical infection thread. The fluorescence intensity profiles of pCYCB1;2 and pKNOLLE reporter-associated signals along the green transects are shown in the corresponding right panels. NC: neighbouring cell. NOD: nodule primordium. Scale bars: 20 μm.

1426
1427
1428
Figure 5. Dividing cortical cells initiating Medicago nodule primordium formation are tetraploid.
(**A**) Confocal images of whole-mount WT transgenic roots expressing mCitrine-CENH3 under the control of native (*M.t.*
pCENH3; left and middle panels) or constitutive promoters (pLjUbi; right panel). Simultaneous expression of H3.3-
mCherry enables the recognition of condensed (star) and segregating chromosomes (middle panel). mCitrine-CENH3
labels the centromeric region of individual chromosomes (filled arrowheads). The number of CENH3-labeled foci
determined across image stacks is indicated in yellow. Images are maximum intensity projections and show merged
fluorescent channels (mCitrine: Green Fire Blue; mCherry and Calcofluor white cell-wall staining: grayscale). In the
Green Fire Blue colour scheme, blue or yellow indicate low or high fluorescence levels, respectively (min. to max. = 1-
140 in the middle panel). Scale bars: left and right panels = 10 μm; middle panel = 2 μm. (**B**) Quantification of the number
of centromeric signals in transgenic root tips expressing mCitrine-CENH3 under native (pCENH3) or constitutive (pUbi)
promoters (n = 55 nuclei). All data points are shown and are from 2 (pUbi) to 3 (pCENH3) independent experiments with
7 composite plants analyzed per construct. Horizontal bars indicate sample means with 95% confidence interval.
Differences were statistically significant (P value = 0.0006) using a Mann-Whitney test. (**C**) Schematic representation of
an early nodule primordium where anticlinal cell divisions occurred in the pericycle (P; dark violet) and cortical cells (C2
to C5; light violet). Rhizobia inside the infection thread are depicted in red. Ep: epidermis. En: endodermis. (**D**) Maximum
intensity projections of an early nodule primodium at the developmental stage represented in (**C**), formed in a WT root
expressing the pLjUbi::mCitrine-CENH3 / pH3.3::H3.3-mCherry construct at 7 dpi with mCherry-producing *S. meliloti*.
Dotted-line ROIs indicate the contours of divided cells in the inner (C4) and middle (C3) cortical layers. The number of
CENH3-labeled foci, determined across image stacks and indicated in yellow are: 11 and 16 in undivided, infected cells
(IC) from the outer cortical layers (C1; left panel and C2, right panel) and 25 to 28 in divided, uninfected cells from the
C3 (filled arrowhead; left panel) and C4 layers (right panel). Images show merged fluorescent channels (mCitrine: Green
Fire Blue; mCherry and Calcofluor white cell-wall staining: grayscale). In the Green Fire Blue colour scheme, blue or
yellow indicate low or high fluorescence levels, respectively. Scale bars: 20 μm. (**E**) Quantification of the number of
centromeric signals in early nodule primordia at the developmental stage depicted in (**C**), formed in inoculated WT roots
(7 to 12 dpi) constitutively expressing mCitrine-CENH3. The number (n) of analyzed nuclei in each cell layer is indicated
on the top. E: endodermis. P: pericycle. All data points are shown with black or violet symbols indicating undivided or
divided cells, respectively. NI: non-infected cell (discs). I: infected cell (triangles). Horizontal dotted lines are positioned
at y = 16 and y = 32, corresponding to diploid (2n = 16) or tetraploid (4n = 32) cellular states in Medicago. Data are from
2 independent experiments with 7 nodule primordia from 5 composite plants analyzed.

Figure 6. Primary infected nodule primordium cells do not reach full competence for chromosome segregation.
(**A**) Schematic representation of CENH3 deposition at centromeres (blue dots) throughout the different pre-mitotic (G1, S, G2 early and G2 late) cell-cycle phases in plants. Nuclei of increasing size and DNA content are depicted as ovals. The dotted-line frame indicates the timing of CENH3 loading at replicated centromeres, occurring during G2 before sister kinetochore (blue doublets) split in preparation to mitosis (Lermontova et al., 2007). Right panel: confocal image of a late G2-phase cell observed in a nodule (NOD) in an inoculated WT transgenic root (12 dpi) constitutively expressing mCitrine-CENH3. The majority of fluorescent signals appear as doublets corresponding to sister kinetochores. Scale bar: 5 μm. (**B**) Confocal images of whole-mount WT roots expressing the pLjUbi::mCitrine-CENH3 / pH3.3::H3.3-mCherry construct and inoculated with mCherry-producing *S. meliloti* (7 to 12 dpi). Filled arrowheads point to nuclei of infected nodule primordium cells. Dotted-line ROIs indicate mCitrine-CENH3 doublets appearing as twin spots on the same focal plane. Neighbouring cells in late G2 or undergoing mitosis (M) are indicated. The presence of CENH3-labeled twin spots was assessed across image stacks in the nuclei of infected cells from the middle (C3) and inner (C4/5) cortical layers and colour-coded as follows: grey = no doublet (upper panels); light blue = 1 to 3 doublets (lower left panel); dark violet = 11 or more doublets (lower right panel). Scale bars: 5 μm. (**A-B**) Images are maximum intensity projections (mCitrine: Green Fire Blue; mCherry and Calcofluor white cell-wall staining: grayscale). In the Green Fire Blue colour scheme, blue or yellow indicate low or high fluorescence levels, respectively. (**C**) Quantification of the nuclear area at the equatorial plane in cells being passed by a cortical infection thread (infected cells) in nodule primordia formed by WT transgenic roots inoculated with *S. meliloti* (7 to 12 dpi). The number (n) of analyzed nuclei in each cell layer (C3 to C4/5) is indicated on the top. The 8C chromatin-value is given for uninfected nodule primordium cells in late G2 showing close to 32 mCitrine-CENH3 doublets as identified in (**A**) and (**B**). All data points are shown and crosses indicate sample means. Differences were not statistically significant according to a Kruskal-Wallis test followed by Dunn's multiple comparisons test. At least 9 composite plants from 2 independent experiments were analyzed. (**D**) Quantification of infected cells in C3 (n = 13) and C4/5 (n = 53) cortical layers in nodule primordia, showing no doublet (grey), 1 to 3 doublets (light blue) or more than 11 doublets (dark violet) labelled by mCitrine-CENH3. Nine composite plants from 2 independent experiments were analyzed. (**E**) Quantification of the length of mCitrine-CENH3 doublets appearing as twin spots on the same focal plane in infected cells (IC) from the C4/5 cortical layer (n = 11), early G2 (n = 12) and late G2-

phase cells (n = 27) in nodule primordia. n = 21 (IC $^{C4/5}$), 28 (G2^{early}), 94 (G2^{late}) doublets. Horizontal bars indicate sample means with 95% confidence interval. Differences were not statistically significant according to a Kruskal-Wallis test followed by Dunn's multiple comparisons test. All data points are shown and are from 6 to 10 composite plants from 2 independent transformation experiments.

Figure 7. NF-YA1 specifically reduces the expression of a G2/M gene in a cell-division enabled system.
(**A**) Live-cell confocal images of *N. benthamiana* epidermal cells ectopically expressing eGFP-NF-YA1 variants: WT (left panel) or impaired in DNA recognition (mutDBD; right panel). Both fusion proteins accumulate in the nucleus and were used in fluorometric β-glucuronidase (GUS) assays. Images are maximum intensity projections and show merged fluorescent (eGFP: green) and bright field (BF) channels. Scale bars: 10 μm. (**B**) Schematic representation of the cell division-enabled leaf system (CDELS) on the scale of epidermal pavement cells in *N. benthamiana*. Differentiated cells in mature leaves (left panel) re-enter the canonical cell cycle upon ectopic expression of the Arabidopsis D-type cyclin CYCD3;1 (right panel). Nearly all re-activated pavement cells (diffuse yellow colouring) have completed cytokinesis after 3 days (filled arrowheads). See also Figure 7 – figure supplement 2B. Scale bars: 20 μm. (**C**) Left panel: schematic representation of the CDEL system on the scale of the canonical cell cycle. Ectopically-expressed CYCD3;1 targets and activates cyclin-dependent kinases (CDK, not shown), fostering the G1/S transition. Progression throughout the different cell-cycle phases is accompanied by the sequential activation of DNA synthesis (S) and mitotic genes (M). Histone *H4* and *KNOLLE* promoter-reporters were selected as readouts for G1/S and late G2/M transcriptional waves, respectively. Right panel: fluorometric GUS assay in tobacco leaf cells ectopically expressing CYCD3;1 and the Medicago pKNOLLE::GUS (pKNOLLE) reporter construct. Fluorescence curves (GUS-mediated hydrolysis of 4-MUG) over time are shown for 3 biological replicates. Error bars indicate standard deviation. Data are from 1 of 4 independent transient transformations. (**D**) Activity of GUS driven by the histone *H4* promoter (pH4) in the absence or presence of ectopically-expressed eGFP-NF-YA1 (NF-YA1) in CDEL samples (+ CYCD3;1). All data points are shown and crosses indicate sample means. Differences were not statistically significant (P value = 0.3323) according to an unpaired t test with Welch's correction. Data are from 3 independent transformation experiments. n = 12 ([-] NF-YA1) and 12 ([+] NF-YA1) biological samples. (**E**) Activity of GUS driven by the *KNOLLE* promoter (pKNOLLE) in CDEL samples (+ CYCD3;1) in the absence or presence of ectopically-expressed eGFP-NF-YA1 variants with WT (NF-YA1) or mutated DNA-binding domain (NF-YA1^{mutDBD}). All data points are shown and crosses indicate sample means. Statistically significant differences ([-] NF-YA1 versus [+] NF-YA1; P value < 0.0001) or non-significant differences ([-] NF-YA1 versus [+] NF-YA1^{mutDBD}; P value = 0.7861) are based on a Kruskal-Wallis test followed by Dunn's multiple comparisons test. Data are from 4 to 5 independent transformation experiments. n = 18 ([-] NF-YA1), 16 ([+] NF-YA1), 16 ([+] NF-YA1^{mutDBD}) biological samples.

1524
1525
1526
Figure 8. Model illustrating characteristic cellular traits on the cortical infection thread trajectory.
(**A**) Cortical cells competent for sustained transcellular infection thread progression are characterized by a reduced
proliferation or endoreduplication activity (pink to green two-colour gradient). By the time they are crossed and ultimately
internalize bacteria, such tetraploid cells (this study, Torrey and Barrios, 1969) reach a post-replicative, 8C ploidy stage
but contrary to direct neighbour cells, they do not commit to a subsequent cell division (red to white two-colour gradient).
(**B**) Rather, cells supporting transcellular infection presumably enter a prolonged G2-phase (this study, Yang et al., 1994)
during which they remodel their histone H3 composition as part of their cellular reprogramming for infection. This
differentiation process and concurrent transcellular passage are accomplished within a last cell-cycle round. We propose
that cellular factors controlling the G2/M transition and eventually the exit from the canonical cell cycle (dashed arrow)
are prominent candidates to support intracellular infection competence.
1537
1538