Caenorhabditis elegans **SEL-5/AAK1 regulates cell migration and cell outgrowth independently of its kinase activity**

Filip Knop[1], Apolena Zounarová[1], Vojtěch Šabata[1], Teije Corneel Middelkoop[2], Marie Macůrková[1,*]

[1] Department of Cell Biology, Faculty of Science, Charles University, Vinicna 7, 128 00 Prague 2, Czech Republic

[2] Institute of Molecular Genetics of the Czech Academy of Sciences, Videnska 1083, 142 20 Prague 4, Czech Republic

* Corresponding author

Keywords: SEL-5/AAK1, Wnt signalling, DPY-23/AP2M1, excretory cell, retromer

Abstract

During *Caenorhabditis elegans* development multiple cells migrate long distances or extend processes to reach their final position and/or attain proper shape. Wnt signalling pathway stands out as one of the major coordinators of cell migration or cell outgrowth along the anterior-posterior body axis. Wnt signalling outcome is fine-tuned by various mechanisms including endocytosis. In this study, we show that SEL-5, the *C. elegans* orthologue of mammalian AP2-associated kinase AAK1, acts together with the retromer complex as a positive regulator of EGL-20/Wnt signalling during the migration of QL neuroblast daughter cells. At the same time, SEL-5 in cooperation with the retromer complex is also required during excretory canal cell outgrowth. Importantly, SEL-5 kinase activity is not required for its role in either neuronal migration or excretory cell outgrowth and neither of these processes is dependent on DPY-23/AP2M1 phosphorylation. We further establish that Wnt proteins CWN-1 and CWN-2 together with Frizzled receptor CFZ-2 positively regulate excretory cell outgrowth, while LIN-44/Wnt and LIN-17/Frizzled together generate a stop signal inhibiting its extension.

Introduction

During the development of *Caenorhabditis elegans* several cells, both neuronal and non-neuronal, migrate in a well-defined and invariant manner along the anterior-posterior (A-P) body axis. Other cells do not migrate but rather extend their processes along the same axis. In both instances, the directionality of migration and growth is controlled by a set of guidance cues provided by the surrounding tissue and received by the corresponding signalling apparatus in the migrating or growing cell (Silhankova and Korswagen, 2007; Sundaram and Buechner, 2016; Sherwood and Plastino, 2018; Hutter, 2019). Gradients of Wnt proteins and a panel of their receptors are the key determinants of the A-P guidance in *C.elegans*. Wnts can act either as attractants, repellents or permissive cues for a given cell and the cell response depends on a given combination of Wnts, Frizzleds and other components of the Wnt pathway expressed in and around that particular cell (Silhankova and Korswagen, 2007; Ackley, 2014; Middelkoop and Korswagen, 2014; Rella et al., 2016).

Endocytosis is an important regulatory mechanism of Wnt signalling applied in both Wnt-producing and Wnt-receiving cells. On the way out of the producing cell, Wnts are accompanied by a Wnt sorting receptor Wntless (WLS) (Banziger et al., 2006; Bartscherer et al., 2006; Goodman et al., 2006). Once WLS is relieved of its cargo at the plasma membrane, it is internalized by clathrin- and adaptor protein complex 2 (AP2)-dependent endocytosis (Pan et al., 2008; Port et al., 2008; Yang et al., 2008). Internalized WLS is subsequently recycled by a multisubunit retromer complex to the Golgi apparatus (Belenkaya et al., 2008; Franch-Marro et al., 2008; Pan et al., 2008; Port et al., 2008; Yang et al., 2008) and further to the endoplasmic reticulum (Yu et al., 2014). Disruption of AP2 or clathrin function results in decreased Wnt secretion due to improper WLS recycling (Pan et al., 2008; Port et al., 2008; Yang et al., 2008). In Wnt-receiving cells, the activity of the Wnt-receptor complex, the signalosome, can be promoted by endocytosis, while at the same time, endocytosis can also

attenuate Wnt signalling by regulating the membrane availability of Wnt receptors (Albrecht

et al., 2021; Colozza and Koo, 2021; Wu et al., 2021).

AP2-associated kinase 1 (AAK1) belongs to the Ark/Prk or Numb-associated (NAK) family

of serine/threonine protein kinases (Smythe and Ayskough, 2003; Sorrell et al., 2016).

Members of this family play an important role in regulating endocytosis in yeast, *Drosophila*

or mammals (Smythe and Ayskough, 2003; Peng et al., 2009). Human AAK1 has been shown

to phosphorylate AP2M1, the µ2 subunit of the AP2 adaptor complex (Conner and Schmid,

2002). Phosphorylation of µ2 at Thr156 by AAK1 enhances the binding affinity of AP2 to

sorting signals of endocytosed proteins (Ricotta et al., 2002). AAK1 has also been shown to

phosphorylate the endocytic adaptor and negative regulator of Notch signalling Numb and

affect its subcellular localization (Sorensen and Conner, 2008). AAK1 can also interact with a

membrane-tethered activated form of the Notch receptor and promote its internalization

(Gupta-Rossi et al., 2011).

AAK1 knock-down upregulated Wnt signalling in a mouse embryonic stem cell-based kinase

and phosphatase siRNA screen (Groenendyk and Michalak, 2011). A recent study in human

cells confirmed this observation and offered a mechanism. By phosphorylating AP2 AAK1

promotes endocytosis of LRP6 and thus shuts down the signalling (Agajanian et al., 2018). In

C. elegans, AAK1 orthologue SEL-5 has been genetically implicated in Notch signalling as a

mutation in *sel-5* can suppress the constitutively active *lin-12*/Notch mutants. Suppression is

however observed only in the *lin-12* allele activating the membrane-anchored Notch and not

when the pathway is activated by expression of the intracellular domain (Fares and

Greenwald, 1999). This hints that SEL-5 could regulate endocytosis in a similar manner as its

mammalian counterpart. So far, no link has been made between SEL-5 and *C. elegans* Wnt

signalling regulation.

In this study, we set to analyse the potential role of the SEL-5 kinase in Wnt signalling

regulation and its contribution to DPY-23/AP2M1 phosphorylation in *C. elegans*. We show

that SEL-5 acts in collaboration with the retromer complex as a positive regulator of EGL-

20/Wnt signalling during the migration of QL neuroblast descendants. We further

demonstrate that although SEL-5 participates in DPY-23 phosphorylation, the observed role

of SEL-5 in neuroblast migration is independent of its kinase activity and the phosphorylation

of DPY-23 is dispensable for Wnt-dependent migration of QL neuroblast descendants. SEL-5

together with the retromer is further required during the active phase of excretory cell canal

extension, again independently of its kinase activity. Finally, we show that Wnt proteins

CWN-1 and CWN-2 together with CFZ-2/Frizzled promote excretory canal outgrowth while

LIN-44/Wnt and LIN-17/Frizzled together define the stopping point for canal extension.

Results

Loss of SEL-5 affects migration of QL neuroblast descendants

Q cell descendant (Q.d) migration is dependent on EGL-20/Wnt signalling. While on the right

side of the L1 larva QR divides and QR.d migrate in the default anterior direction, on the left

side QL.d respond to the EGL-20 signal produced in several cells around the rectum by

expressing homeotic gene *mab-5* and migrate to the posterior. In the absence of active EGL-

20 signalling *mab-5* is not expressed and QL.d migrate to the anterior (Salser and Kenyon,

1992) (Fig. 1A). To visualize Q.d migration we utilized *Pmec-7::gfp* expressing transgene

muIs32 that is, apart from other cells, active in AVM and PVM touch neurons, which are QR

and QL descendants, respectively (Ch'ng et al., 2003). We performed RNA interference

(RNAi) against *sel-5* in wild type background and observed no effect on QL.d migration (Fig.

1B). To increase the sensitivity of the RNAi approach, we repeated the RNAi experiment

using *vps-29(tm1320)* mutant strain. This strain harbours a null mutation in the VPS-29

retromer subunit, displays only weakly penetrant QL.d migration defect (Coudreuse et al., 2006) and has been used previously as a sensitized background to uncover genes required for the Q.d migration (Harterink et al., 2011b; Lorenowicz et al., 2014). RNAi against *sel-5* significantly increased the QL.d migration defect of *vps-29(tm1320)* (Fig. 1B, Figure 1 – figure supplement 1). To verify the results obtained with RNAi we assessed the QL.d migration in *sel-5(ok363)* and *sel-5(ok149)* mutants. Similarly to RNAi results, single *sel-5* mutation did not affect QL.d migration while *sel-5 vps-29* double mutants displayed almost fully penetrant defect (Fig. 1C, D).

Loss of *vps-29* affects EGL-20/Wnt signalling at the level of Wnt production (Yang et al., 2008). We further tested whether loss of *sel-5* could enhance partially penetrant defects in QL.d migration in mutants acting at a different step of the EGL-20/Wnt pathway. Loss of *sel-5* strongly increased the weak QL.d migration defect of the *lin-17*/Frizzled mutants but did not affect the already more penetrant QL.d defect of *mig-1*/Frizzled mutant animals (Fig. 1D). We next asked whether loss of *sel-5* in the *vps-29* background affects other Wnt-dependent processes such as the polarization of the ALM and PLM neurons, CAN neuron migration or T cell polarity. Polarization of ALM and PLM is regulated by several Wnt proteins (Pan et al., 2006; Prasad and Clark, 2006; Hilliard and Bargmann, 2006), CAN neuron migration is governed predominantly by CWN-2 with minor contribution from CWN-1 and EGL-20 (Zinovyeva and Forrester, 2005; Zinovyeva et al., 2008) and T cell polarity depends on the LIN-44/Wnt signal (Herman et al., 1995). We observed weakly penetrant ALM polarization defects in both *sel-5(ok363) vps-29* and *sel-5(ok149) vps-29* double mutants, a very mild PLM polarization defect was detected in a strain carrying the *ok149* allele (Table 1). Mildly penetrant anterior displacement of the CAN neuron was observed in the double mutants, while about 17% of double mutant animals carrying the *ok149* allele did not correctly form the phasmid sensillum which indicates defects in T cell polarization (Table 1). Apart from

these Wnt-dependent phenotypes, we also observed reduced brood size in the double mutants

compared to controls (Table 1), defective tail formation (Fig. 1E) and a progressive egg-

laying defect leading to a bag-of-worms phenotype.

The penetrance of the observed phenotypes was not the same in the two *sel-5* alleles. This

was striking as both alleles carry deletions hitting the kinase domain of SEL-5 protein. We

were able to amplify shorter than wild type transcripts from both *sel-5(ok363)* and *sel-*

5(ok149) animals using primers in exon 1 and exon 17 (Fig. 1F). Sequencing of the *ok363*

transcript revealed that amino acids 1 to 243 of the wild type protein followed by extra 32

amino acids would be translated due to a frameshift and a premature stop codon. Such protein

encompasses a larger part of the SEL-5 kinase domain including the active site. In agreement

with data published by Fares and Greenwald (Fares and Greenwald, 1999), sequencing of the

ok149 transcript revealed that it can give rise to a protein containing the first 153 amino acids

of SEL-5 followed by 21 extra amino acids and is thus devoid of the active site (Fig. 1F). If

such truncated proteins are present in the mutants, either one or both may possess some

residual activity that is responsible for the observed phenotypic differences.

SEL-5 is required in EGL-20/Wnt producing cells to direct QL.d migration

The asymmetric migration of the Q neuroblast is regulated both cell-autonomously and non-

autonomously. To analyse the tissue-specific requirement of SEL-5 in QL.d migration we

expressed full-length SEL-5 tagged with GFP at its N-terminus under the control of different

tissue-specific promoters in *sel-5 vps-29* background and assayed the QL.d migration.

Expression of SEL-5 from *egl-20* promoter (Coudreuse et al., 2006) resulted in a significant

rescue of the QL.d migration defect of *sel-5 vps-29* double mutants albeit not to the level of

vps-29 single mutant (Fig. 2A, Figure 2 – figure supplement 1). No rescue was obtained when

SEL-5 was expressed from the *wrt-2* promoter that is active in the seam cells and the Q cell

(Aspöck et al., 1999; Middelkoop et al., 2012). These data suggest that *sel-5* is required in the

EGL-20/Wnt-producing cells to control QL.d migration. However, a similar level of rescue as

with *egl-20* promoter was obtained also when SEL-5 was expressed from *hlh-1* promoter

specific for the body wall muscle cells (Krause et al., 1990, Harfe et al., 1998). When SEL-5

was expressed from *egl-20* and *hlh-1* promoter at the same time, the rescue almost reached the

background frequency of the QL.d defect, caused by *vps-29* alone (Fig. 1D, 2A, Figure 2 –

figure supplement 1). Apart from the rectal epithelial cells, *egl-20* expression was also

detected in posterior ventral body wall muscle quadrants VL23 and VR24 (Harterink et al.,

2011a), it is thus possible that *hlh-1* promoter-driven expression of SEL-5 in these muscle

quadrants is responsible for the observed rescue and only the combined expression from all

EGL-20 producing cells is sufficient to drive a full rescue. Alternatively, the observed rescue

pattern could indicate that SEL-5 is required both in the Wnt-producing and in the muscle

cells.

To confirm that the tissues identified above in the rescue experiments are tissues with a

genuine *sel-5* expression we set to determine the endogenous *sel-5* expression pattern which

has not been previously analysed. The only published expression data (Fares and Greenwald,

1999) were based on expression from *sel-12* instead of *sel-5* promoter. We generated GFP

knock-in strains by inserting GFP into the *sel-5* genomic sequence using the SEC cassette

approach (Dickinson et al., 2015). We obtained two strains, one expressing a transcriptional

P*sel-5::gfp* reporter and one, after excision of the SEC cassette, an N-terminally tagged

GFP::SEL-5 under the control of its own promoter (P*sel-5::gfp::sel-5*). Microscopic analyses

of both strains revealed that *sel-5* is expressed broadly, but at a low level judged by the GFP

signal intensity. The most prominent expression was observed in the gonad from the

beginning of its development (Fig. 2B). Expression was further observed in the developing

oocytes, in the vulva, in epidermal cells and most notably, in the rectal epithelial cells that are

known to produce EGL-20/Wnt (Whangbo and Kenyon, 1999) (Fig. 2B, C). We could not

detect *sel-5* expression in the muscles, although we cannot exclude that a low level of

expression, below the detection limit using the endogenous locus-tagging approach, is

present. Interestingly oocytes and epidermal seam cells expressing the GFP::SEL-5 fusion

protein revealed its distinct subcellular localization. Protein localized to punctate structures

located close to the cell surface (Fig. 2C). This pattern resembles the subcellular distribution

of human AAK1 (Conner and Schmid, 2002) hinting that SEL-5 could be involved in

regulating intracellular transport similar to AAK1.

SEL-5 does not affect MIG-14 endocytosis

SEL-5 orthologue AAK1 has been implicated in endocytosis regulation (Conner and Schmid,

2002; Ricotta et al., 2002). The obvious candidate that could be affected by SEL-5-dependent

endocytosis in Wnt-producing cells is the Wnt cargo receptor MIG-14/Wls. Human AAK1

can phosphorylate AP2 subunit μ2 (AP2M1) (Conner and Schmid, 2002; Ricotta et al., 2002)

and thus increase AP2 affinity to cargo molecules (Ricotta et al., 2002). It is conceivable that

SEL-5 could regulate MIG-14/Wls trafficking at the level of AP2-dependent endocytosis.

The *C. elegans* μ2 subunit of the AP2 complex, DPY-23 (also known as APM-2), has been

shown to participate in MIG-14 internalization (Pan et al., 2008; Yang et al., 2008). We first

tested whether loss of *sel-5* expression has any effect on the level of phosphorylation of DPY-

23. A phospho-specific antibody recognizing the phosphorylated threonine T160 (T156 in

mammalian AP2M1) of DPY-23 (Hollopeter et al., 2014) revealed a decrease in the level of

phosphorylated endogenous DPY-23 in *sel-5* mutants compared to wild type animals (Fig.

3A). Similarly, T160 phosphorylation was reduced on overexpressed GFP-tagged DPY-23

(Fig. 3B). Next, we assessed MIG-14 levels and localization in *sel-5 vps-29* mutants using a

transgene expressing a functional MIG-14::GFP protein (Lorenowicz et al., 2014). However,

as shown in Fig. 3C and D (Fig. 3C, D), levels of MIG-14::GFP did not change significantly

in *sel-5 vps-29* compared to *vps-29* single mutant. Levels of MIG-14::GFP in *sel-5* single

mutants were variable yet on average comparable to the wild type. We therefore concluded

that the significant increase in QL.d migration defect in *sel-5 vps-29* compared to *vps-29*

cannot be attributed to changes in MIG-14/Wls levels. Importantly, MIG-14::GFP did not re-

localize to the plasma membrane in either *sel-5* or *sel-5 vps-29* mutants, in striking contrast to

the MIG-14::GFP behaviour in animals treated with *dpy-23* RNAi (Figure 3 – figure

supplement 1). This rules out the possibility that the Wnt-related phenotypes in *sel-5 vps-29*

mutants arise from a defect in MIG-14 internalization.

Our tissue-specific rescue experiments revealed that for a full rescue, simultaneous *sel-5*

expression is necessary from both *egl-20* and *hlh-1* promoters. SEL-5 could therefore regulate

endocytosis along the EGL-20 transport route, thus shaping the EGL-20 gradient. To test this

hypothesis, we endogenously tagged EGL-20 with GFP and assessed Wnt gradient formation

in control and mutant backgrounds. This approach has been used previously and revealed that

extracellular EGL-20 can be detected in the form of distinct puncta that most likely represent

EGL-20 bound to Frizzled (Pani and Goldstein, 2018). We detected EGL-20 puncta in all

strains tested, albeit with different frequencies (Figure 3 – figure supplement 2). While the

number of EGL-20 puncta in *sel-5* mutants was not significantly different from the one in

control animals, in *sel-5 vps-29* mutants the number of puncta was substantially reduced.

However, statistical evaluation did not reveal significant differences between *vps-29*, *sel-5*

vps-29 and *vps-35*; although a decreasing trend in puncta number was visible (Figure 3 –

figure supplement 2). In *vps-35*, the EGL-20 gradient was shown to be highly reduced or

absent (Coudreuse et al., 2006; Harterink et al., 2011b) and was used for comparison. We also

compared the signal intensity in the EGL-20 spots among the different strains but no

significant differences were detected (Figure 3 – figure supplement 2). Our data thus do not

support the notion that *sel-5* significantly affects EGL-20 gradient formation.

SEL-5-associated phenotypes are independent of DPY-23 phosphorylation

Although endocytic cargo other than MIG-14 may be affected in *sel-5 vps-29* mutants, we

could not exclude the possibility that the observed phenotypes are a consequence of defects in

other mechanisms unrelated to DPY-23 phosphorylation. To test this hypothesis we repeated

the rescue experiment presented in Fig. 2A now with SEL-5 carrying either K75A or D178A

point mutations. Position D178 corresponds to the D176 of human AAK1 and is part of the

conserved HRD motif in the catalytic loop, K75 corresponds to K74 in AAK1 and is

predicted to affect ATP binding (Figure 4 – figure supplement 1). In AAK1 both mutations

abolished its ability to phosphorylate AP2M1 (Conner and Schmid, 2003). When expressed

under the control of *egl-20* promoter, the mutant SEL-5 proteins could still rescue the QL

migration phenotype (Fig. 4A, Figure 4 – figure supplement 2). This indicates that SEL-5

kinase activity is not responsible for the phenotypes observed in *sel-5 vps-29* mutants. To test

this further, we next asked whether the phosphorylation at the T160 position of DPY-23 is

necessary for QL.d migration. To this end, we utilized *dpy-23(mew25)* allele harbouring

T160A point mutation (G. Hollopeter and G. Beacham, unpublished). Characterization of

dpy-23(mew25) animals revealed that homozygous mutants are viable; they are not dumpy

and look superficially wild type. This indicates that the mutants do not suffer from a gross

endocytosis defect even though no DPY-23 phosphorylation was detected in *dpy-23(mew25)*

animals (Fig. 4B). In compliance with previous findings using *dpy-23* alleles harbouring

mutations changing the T160 amino acid (Hollopeter et al., 2014, Partlow et al., 2019),

mew25 is able to rescue the "jowls" phenotype of *fcho-1(ox477)* mutants (Fig. 4C). FCHO-1

is a member of the muniscin family of proteins and a proposed allosteric activator of AP2

(Hollopeter et al., 2014). We then asked whether complete loss of DPY-23 phosphorylation

would lead to a QL.d migration defect. This implies that *mew25* produces a protein with

altered function. Animals carrying *dpy-23(mew25)* alone or in combination with *sel-5(ok149)*

did not show any defects in QL.d migration (Fig. 4D). Variable increase in QL.d migration

defects was observed in *vps-29;dpy-23(mew25)* animals, but the penetrance of the defect

never reached levels comparable to those in *sel-5 vps-29* mutants (Fig. 1D, 4D). Furthermore,

MIG-14::GFP did not re-localize to the plasma membrane in *dpy-23(mew25)* similar to the

sel-5 mutants and unlike in *dpy-23* RNAi animals (Fig. 4E). Together these observations

strongly suggest that the role of SEL-5 in regulation of QL.d migration is not dependent on its

kinase activity and moreover, that DPY-23 phosphorylation at T160 is not a major regulatory

event in this process.

The outgrowth of excretory cell canals is impaired in *sel-5 vps-29* mutants

Apart from the QL.d migration defect in the *sel-5 vps-29* double mutants, in some animals, we

unexpectedly noticed a severe shortening of the posterior canals of the excretory cell that

prompted us to analyse this phenotype in more detail. The excretory cell (also called the

excretory canal cell) is a large H-shaped cell required for osmoregulation (Buechner et al.,

1999; Liegeois et al., 2007). The excretory canal cell body is located near the posterior bulb

of the pharynx and four excretory canals emanate from the cell body – two short ones directed

to the anterior and two posterior canals extending to the rectum (Fig. 5A). To assess the

morphology of the excretory cell we expressed GFP under the control of *pgp-12* promoter

which is active exclusively in the excretory cell (Zhao et al., 2004) and analysed the length of

the canals in late L4 or early adult animals. While the posterior canal length in *sel-5* or *vps-29*

single mutants was indistinguishable from the wild type controls, in more than 60 % of *sel-5*

vps-29 mutants the posterior canals stopped at various positions anterior to the rectum (Fig.

5A, B). A similar effect, albeit with lower penetrance, was observed when assessing the

anterior canal length (Fig. 5C). Interestingly, the posterior canal on the right side of the

animal was significantly more affected than its counterpart on the left side in *sel-5 vps-29*

double mutants carrying the *ok149* allele (Fig. 5D). Similar trend was observed in *sel-*

5(ok363) vps-29 animals, although the difference observed there did not reach statistical

significance (Fig. 5D).

The active growth of the canals starts during embryogenesis and continues during L1, after

that the canals passively grow with the growing animal (Fujita et al., 2003). Comparison of

canal lengths at several time points within a 24-hour interval after hatching revealed that in

sel-5 vps-29 double mutants, the posterior canals of the excretory cell are shorter already at

the time of hatching and the growth defect prevails into the larval development (Fig. 5E).

Excretory canal shortening has not previously been reported for any of the retromer

components. We therefore tested whether this phenotype is specific for the *vps-29* retromer

subunit or whether the whole retromer complex is required together with *sel-5* for proper

excretory canal extension. We performed RNAi against *vps-35*, *vps-26*, *snx-1* and *snx-3* in

sel-5(ok149) background and assessed the posterior canal length. Except for *snx-1* RNAi,

significant shortening was observed with RNAi against all other retromer subunits, with *vps-*

35 showing the strongest phenotype (Fig. 5F). These data confirm that the whole retromer

complex acts together with *sel-5* to control excretory canal outgrowth.

We next tested in which tissue SEL-5 activity is required for excretory cell outgrowth. We

expressed SEL-5 from either *pgp-12* (excretory cell), *hlh-1* (body wall muscle cells) or *col-10*

(hypodermis) promoters in *sel-5 vps-29* double mutants and checked for any rescue. As

shown in Fig. 5G, expression of SEL-5 from *pgp-12* promoter almost fully rescued the

excretory cell shortening while SEL-5 expression from the muscle-specific *hlh-1* promoter

did not affect the excretory cell phenotype of the *sel-5 vps-29* double mutants. As in the case

of the QL.d migration defect, the excretory canal length was efficiently rescued also by the

SEL-5D178A variant expressed from *pgp-12* promoter arguing that SEL-5 kinase activity is

not required for canal extension (Fig. 5G). Expression of VPS-29 in the excretory cell

partially rescued the canal shortening suggesting that SEL-5 and the retromer complex act

together within the same cell to regulate canal length. Interestingly, the expression of SEL-5

from *col-10* promoter which is active in the hypodermis (Spencer et al., 2001) also resulted in

a significant rescue of the excretory canal shortening. This suggests that SEL-5 activity in

both the excretory cell and in the hypodermis contributes to the regulation of the canal

outgrowth.

To support our finding that SEL-5 role in excretory canal outgrowth is independent of DPY-

23 phosphorylation, we analysed posterior canal length in *dpy-23(mew25)* mutant animals. No

canal shortening was observed in animals carrying *dpy-23(mew25)* alone (Fig. 5H), while

only very mild shortening was occasionally observed in either *sel-5; dpy-23(mew25)* or *vps-*

29; dpy-23(mew25). These observations are consistent with a role of SEL-5 other than DPY-

23 phosphorylation at T160.

Wnt-dependent signalling is required to establish proper excretory cell canal length

We were interested in whether the observed shortening of the excretory canals in *sel-5 vps-29*

mutants could be a consequence of crosstalk with the Wnt signalling as in the case of QL.d

migration. *lin-17/Frizzled* mutants have been reported to show an overgrowth of the posterior

excretory cell canals past the rectum into the tip of the tail (Hedgecock et al., 1987) while loss

of *axl-1*, one of the two *C. elegans* Axin orthologues, resulted in ectopic branching of the

posterior excretory canal but no effect on the canal length was observed. The ectopic

branching in *axl-1* mutants could be rescued by simultaneous loss of *bar-1*/β-catenin or *pop-*

1/Tcf expression (Oosterveen et al., 2007). Wnt signalling thus seems to play a role in

excretory canal growth, but so far, the other Wnt pathway components and their mechanism

of action remained unknown. We therefore tested whether mutations in other Wnt pathway

components could affect excretory cell growth. Among the four Wnts tested (EGL-20, LIN-

44, CWN-1, CWN-2), only mutants in *lin-44*/Wnt exhibited almost fully penetrant posterior

excretory canal overgrowth phenotype, as observed in *lin-17*/Frizzled mutants (Fig. 6A, B).

Mild canal shortening was observed in *cwn-2*/Wnt and *cfz-2*/Frizzled mutants, while a very

weak overgrowth phenotype could be detected in *egl-20*/Wnt mutants (Fig. 6B). No change in

excretory canal length was displayed by *cwn-1*/Wnt or *mig-1*/Frizzled mutants. Interestingly,

highly penetrant canal overgrowth was also observed in *mig-14*/Wls mutants, while loss of

mig-5/Dishevelled or *dsh-1*/Dishevelled resulted in partially penetrant overgrowth phenotype

(Fig. 6B). We next tested various combinations of Wnt and Frizzled mutants and found out

that simultaneous loss of *lin-44* and *cwn-1* or *lin-44* and *cwn-2* expression resulted in partial

rescue of the canal overgrowth, same effect was observed after simultaneous loss of *lin-17*

and *cfz-2*. Surprisingly, the loss of *egl-20* and *mig-1* also partially rescued the canal

overgrowth in *lin-17* even though neither of them displayed canal shortening on its own,

while at the same time *mig-1* suppressed canal shortening in *cfz-2*. Most prominent canal

shortening was observed in either *cwn-1; cfz-2* or *cwn-2; cfz-2* double mutants. Apart from

shortening, three instead of two posterior canal branches were detected in 11% of *cwn-1; cfz-

2* and 15% of *cwn-2; cfz-2* double mutants (Fig. 6D). These data suggest that at least two

Wnt-dependent pathways are acting during the extension of the excretory canal; one involving

lin-44, *lin-17*, *dsh-1* and *mig-5* that is responsible for determining the stopping point for the

growing canal, and one involving *cwn-1*, *cwn-2* and *cfz-2* that contributes to the canal growth.

The most striking phenotype is the overgrowth of the posterior excretory canal in *lin-17*, *lin-

44* and *mig-14* mutants. Interestingly when we measured the length of the excretory canal

within the 24-hour interval after hatching, no difference in canal length was detected between

wild type controls and *lin-17*, *lin-44* and *mig-14* mutants (Fig. 6C). This suggests that the

initial growth is not affected in the mutants but rather that these mutants miss a signal

determining the final position of the canal in later larval stages. Simultaneous loss of *lin-17*

and *sel-5* led to a partial rescue of the *lin-17* phenotype (Fig. 6B). Similar effect was observed

in *lin-44; sel-5* double mutants (Fig. 6B). Analysis of a *lin-17; sel-5 vps-29* triple mutant was

not possible as such combination was lethal. These data indicate that *sel-5* is required for the

initial growth of the excretory canals and the LIN-44/Wnt signal acting through LIN-

17/Frizzled receptor is required at a later stage to terminate excretory canal extension.

Discussion

The NAK family kinase AAK1 has recently been added to the long list of Wnt signalling

regulators (Agajanian et al., 2019), acting at the level of LRP6 endocytosis. In the present

study, we probed the role of *C. elegans* AAK1 orthologue SEL-5 in Wnt signalling and we

uncovered its requirement for Wnt-dependent QL neuroblast daughter cell migration. We also

unveiled that SEL-5 is required during the outgrowth of the excretory canal cell prior to a Wnt

dependent stop signal. Importantly, in both instances the role of SEL-5 is independent of its

kinase activity, defining an alternative mode of action for this kinase.

The role of SEL-5 has not been extensively studied in *C. elegans*, the only available reports

uncovered the role of *sel-5* as a genetic suppressor of a constitutively active *lin-12*/Notch

mutation (Tax et al., 1997; Fares and Greenwald, 1999). Information about the protein

function was missing. SEL-5 shows the highest similarity to the mammalian Numb-associated

kinase family members AAK1 and BMP2K (Shaye and Greenwald 2011; Sorrell et al., 2016).

Both AAK1 and BMP2K were shown to phosphorylate the μ2 subunit of the AP2 adaptor at

position T156 and thus regulate the clathrin-mediated endocytosis (Conner and Schmid, 2002;

Ricotta et al., 2002; Ramesh et al., 2021). Our findings uncovered some similarities between

SEL-5 and AAK1 or BMP2K. Firstly, we show that the level of DPY-23/AP2M1

phosphorylation in *sel-5* mutants is decreased compared to controls. Secondly, the subcellular

localization of SEL-5 to punctate structures proximal to the plasma membrane in oocytes and

in epidermal seam cells resembles the subcellular distribution of human AAK1 (Conner and

Schmid, 2002). These data support the notion that SEL-5 has a function analogous to its

mammalian counterparts.

Given the established role of AAK1 in mammalian endocytosis regulation (Kadlecova et al.,

2017; Agajanian et al., 2019; Wrobel et al., 2019), it was striking to see that loss of *sel-5*

expression alone had no obvious developmental consequences. However, even in mammalian

cells complete loss of AP2M1 T156 phosphorylation does not result in endocytosis block but

only in a reduction of its efficiency (Motley et al., 2006; Wrobel et al., 2019). We did not

observe complete loss but only a partial reduction of DPY-23 phosphorylation in *sel-5*

mutants. This implies that SEL-5 is not the only kinase responsible for DPY-23

phosphorylation. It also suggests that endocytosis is not inhibited but could be only partially

compromised. In such case, the level of ongoing endocytosis might be sufficient to support

proper developmental decisions such as the direction of Q neuroblast migration. We argued

that in combination with an additional insult to the intracellular trafficking machinery, in this

case, the retromer complex, the suboptimal function of endocytosis would manifest itself.

Indeed, *sel-5 vps-29* double mutants display two highly penetrant phenotypes – the QL.d

migration defect and the shortening of the excretory cell canals. Strikingly, neither of these

phenotypes can be attributed to a decrease in endocytic efficiency due to a lack of DPY-23

phosphorylation. Firstly, both phenotypes can be rescued by a tissue-specific expression of a

kinase-inactive SEL-5 mutant protein. Secondly, *vps-29; dpy-23(mew25)* mutants expressing

only non-phosphorylatable DPY-23 do not phenocopy *sel-5 vps-29* mutants. Alternative

mechanisms of SEL-5 function thus have to be considered. It is possible that either SEL-5

affects endocytosis by a mechanism not dependent on DPY-23 phosphorylation or that SEL-5

regulates QL.d migration and excretory cell canal extension by endocytosis-independent

mechanisms. Our findings about MIG-14/Wls behaviour, the most obvious endocytic cargo

candidate for QL.d migration, argue against the first possibility; most notably the fact that

MIG-14 does not re-localize to the plasma membrane in the absence of SEL-5. However, the

effect of SEL-5 on a different endocytic cargo cannot be excluded. For example, SEL-5 could

regulate the turnover of guidance receptors at the plasma membrane of the growing tip in case

of the excretory canal extension. In *Drosophila*, loss of Numb-associated kinase (Nak) results

in the absence or shortening of higher-order dendrites and Nak has been shown to promote

endocytosis of Neuroglian/L1-CAM during dendrite extension and branching (Yang et al.,

2011). In the *C. elegans* excretory cell, PAT-3/β1-integrin is required for canal outgrowth

(Hedgecock et al., 1987). Integrins constantly cycle between the plasma membrane and

intracellular compartments using various trafficking routes (Moreno-Layseca et al., 2019) and

could thus serve as a putative SEL-5 cargo.

A possible mechanism for how SEL-5 could affect endocytosis independently of DPY-23

phosphorylation follows from the finding, that SEL-5 physically interacted with REPS-1

protein in a yeast-two-hybrid assay (Tsushima et al., 2013). REPS-1 is an orthologue of

mammalian REPS1 (RALBP1-associated Eps Domain-containing 1), an Eps homology-

containing protein known to interact with RALBP1 (RalA-binding protein 1) (Yamaguchi et

al., 1997). RALBP1 is an effector of Ral GTPase that can interact with the μ2 subunit of AP2

and regulate endocytosis (Jullien-Flores et al., 2000). Importantly, human AAK1 interacted

with both RALBP1 and REPS1 in several high-throughput mass spectrometry analyses

(Huttlin et al., 2017; Huttlin et al., 2021; Cho et al., 2022; Golkowski et al., 2023). SEL-5 thus

could regulate the AP2 complex indirectly via interaction with the REPS1-RALBP1 complex.

Searching for a possible endocytosis-independent mechanism of SEL-5 activity is

complicated by the fact that while in the case of QL.d migration SEL-5 function is cell-

nonautonomous, cell-autonomous expression of SEL-5 is sufficient to rescue the *sel-5 vps-29*

excretory canal extension defect. This suggests that more than one mechanism might exist.

One possibility common for both phenotypes comes from recent studies in mammalian cells.

BMP2K has been shown to localize to the early secretory compartment and to regulate COPII

coat assemblies (Cendrowski et al., 2020). SEL-5 thus could affect Wnt secretion in case of

QL.d migration or new guidance receptor delivery to the plasma membrane in case of the

excretory canals. EGL-20 gradient analyses did not detect significant differences between

wild type and *sel-*5 animals, but also not between *vps-29* and *sel-5 vps-29* in the number of

EGL-20 puncta despite the almost 8-fold difference in the QL.d migration phenotype

penetrance between these two mutants. This would argue against a SEL-5 role in secretion.

However, it is still possible that even a small decrease in secretion efficiency in *vps-29* could

substantially decrease the frequency of reaching the EGL-20 concentration threshold required

to initiate QL migration and such a decrease may not be captured by our assay. A direct

correlation between the amount of EGL-20 and the QL.d phenotype would be necessary to

answer this question. More experiments will thus be required to establish how SEL-5 exerts

its function. The biggest question to answer will be how it does so independently of its kinase

activity.

Apart from the role of SEL-5 and the retromer complex in excretory cell canal extension, we

have also uncovered a significant contribution of Wnt pathway components in defining

excretory canal length. The role of Wnt signalling in the outgrowth of the excretory canals has

been suggested (Sundaram and Buechner, 2016), however, the only evidence came from the

early observation that canals overgrow in *lin-17* mutants (Hedgecock et al., 1987) and that

erroneous canal branching is observed in *axl-1*/Axin-like mutants (Oosterveen et al., 2007).

More recently, it has been shown that loss of PLR-1, a transmembrane E3 ubiquitin ligase,

results in excretory canal shortening (Bhat et al., 2015). PLR-1 is responsible for the

internalization of Frizzled receptors from the cell surface (Moffat et al., 2014). However, a

direct link between PLR-1 function in the excretory cell and Wnt signalling has not been

established. Our analysis of single Wnt and Frizzled mutants revealed that while loss of *cwn-2*

or *cfz-2* expression resulted in a very mild shortening of the excretory canal, loss of *lin-44* or

lin-17 led to profound canal overgrowth (summarized in Fig. 7A). These findings suggested

that two independent Wnt pathways could be employed to establish proper excretory canal

length – one promoting canal extension and one generating the stop signal for growth

termination. Further analyses of double mutants and other Wnt signalling components

revealed that the extension-promoting pathway includes *cwn-1* in addition to *cwn-2* and *cfz-2*,

while the stop-signal pathway encompasses *lin-44*, *lin-17*, *dsh-1*, *mig-5* and *mig-14*. A similar

repulsive role of LIN-44/LIN-17 complex has been described in the case of a posterior axon

of *C. elegans* GABAergic DD6 motor neuron (Maro et al., 2009) or PLM, ALN and PLN

neurons (Zheng et al., 2015). Loss of *lin-44* or *lin-17* expression promoted outgrowth of the

posterior neurites of these neurons implicating that in wild type animals, LIN-44 serves as a

repulsive cue. On the other hand, *cwn-2* and *cfz-2* were shown to positively regulate the

posterior neurite outgrowth of RMED/V neurons with *cwn-2* acting as an attractive cue (Song

et al., 2010). The role of two other Wnt signalling components, *egl-20* and *mig-1*, is less clear.

No effect (*mig-1*) or only very mild overgrowth defect (*egl-20*) is observed in single mutants.

However, both *egl-20* and *mig-1* significantly rescue the overgrowth phenotype of *lin-17*

mutants, while at the same time, *mig-1* can suppress the shortening of canals in *cfz-2* mutants.

EGL-20-producing cells are localized around the rectum (Whangbo et al., 1999; Harterink et

al., 2011a), exactly where the excretory canals stop, while LIN-44 is expressed more

posteriorly (Herman et al., 1995; Harterink et al., 2011a). A possible explanation could thus

be that while LIN-44 provides a general posterior repulsive signal, EGL-20 fine-tunes the

exact stopping position of the growing canal. The role of different Wnts and Frizzleds in

excretory canal outgrowth is summarized in Fig. 7B.

Further investigation will be required to decipher the exact way how SEL-5 and the retromer

crosstalk with Wnt signalling during excretory cell outgrowth. It is clear though that more

than one mechanism is likely involved. First, *sel-5 vps-29* mutants display canal shortening

similarly to *cwn-1; cfz-2* or *cwn-2; cfz-2* suggesting a positive regulatory role. Mutants in *lin-*

17 and *lin-44* display canal overgrowth, yet *sel-5* is partially able to suppress this phenotype.

This would imply a negative regulatory role of *sel-5* and be in agreement with the role of

AAK1 in Wnt pathway regulation (Agajanian et al., 2019). However, *sel-5* and *vps-29* are

required already during the initial larval outgrowth while the LIN-44/LIN-17 signal is

required later. The observed rescue might thus also be explained by a delayed growth of the

canal and not by a direct impact of *sel-5* and *vps-29* on LIN-44 or LIN-17 levels or

localization.

Taken together, we have uncovered cooperation between *C. elegans* orthologue of AAK1

kinase, SEL-5, and the retromer complex in regulating cell migration and cell outgrowth, and

we have also outlined the engagement of Wnt pathway components in defining the length of

the excretory cell canals. The surprising independence of SEL-5 function on its kinase activity

opens a new path for investigation with high relevance also for mammalian biology. AAK1

has been proposed as a potential drug target for treating various neurological disorders and

preventing viral entry, where the kinase activity is the target for inhibition (Martinez-Gualda

et al., 2021; Xin et al., 2023). In light of our findings, the non-enzymatic activities of AAK1

should also be taken into account.

Materials and methods

Caenorhabditis elegans strains and culture

Standard methods of cultivation, manipulation and genetics of *C. elegans* were used as described previously (Brenner 1974). Bristol N2 strain was used as wild type, *Escherichia coli* strain OP50 was used as a food source. Other strains, extrachromosomal and integrated arrays used in this study were:

LGI: *pry-1(mu38)* (Maloof et al, 1999), *lin-17(n671)* (Brenner 1974), *lin-44(n1792)* (Herman and Horvitz, 1994), *mig-1(e1787)* (Brenner 1974)

LGII: *cwn-1(ok546), dsh-1(ok1445), mig-14(mu71)* (Harris et al., 1996), *muIs32 [Pmec-7::gfp; lin-15(+)]* (Ch'ng et al, 2003), *huSi2 [Pmig-14::mig-14::gfp], mig-5(cp385[mNG-GLO^AID::mig-5])* (Heppert et al., 2018)

LGIII: *sel-5(ok363), sel-5(ok149), vps-29(tm1320), sIs10089* (McKay et al., 2003)

LGIV: *egl-20(n585)* (Harris et al., 1996), *cwn-2(ok895)*

LGV: *muIs35 [Pmec-7::gfp; lin-15(+)]* (Ch'ng et al, 2003), *cfz-2(ok1201)*

LGX: *dpy-23(mew25)*(G. Hollopeter and G. Beacham, unpublished), *kyIs4 [Pceh-23::unc-76::gfp + lin-15(+)]* (Zallen et al., 1999)

Molecular biology, germline transformation and RNA interference

Total RNA was isolated using the TRIzol™ reagent (cat.# 15596026, ThermoFisher Scientific) from a mixed stage population of *C. elegans* N2 strain collected from a single 9 cm NGM plate. cDNA was transcribed from total RNA using SuperScript™ III Reverse Transcriptase (cat.# 18080093, ThermoFisher Scientific) and an oligo(dT) primer. Full-length *sel-5* (isoform a), *vps-29* (isoform a) or *dpy-23* (isoform b) cDNA was PCR-amplified using gene-specific primers. PCR products were cloned into pJet1.2/blunt vector (cat.# K1232, ThermoFisher Scientific) and sequenced. Same procedure was applied to clone truncated *sel-5*

cDNA from *sel-5(ok363)* and *sel-5(ok149)* mutant animals. Two- or three-fragment Gibson

assembly (Gibson et al., 2009) was employed to construct tissue-specific gene expression

vectors. For that, 524 bp of *col-10* promoter sequence, 600 bp of *eft-3* promoter sequence,

4419 bp of *egl-20* promoter sequence, 2896 bp of *hlh-1* promoter sequence, 3410 bp of *pgp-*

12 promoter sequence and 1629 bp of *wrt-2* promoter sequence were PCR-amplified from N2

genomic DNA, cDNAs were amplified from the corresponding pJet1.2 vectors and pPD95.81

(a gift from Andrew Fire (Addgene plasmid # 1497 ; http://n2t.net/addgene:1497 ;

RRID:Addgene_1497)) was a source of backbone and GFP sequences. Fragments were

assembled in the desired combinations and correct assembly was verified by sequencing.

SEL-5 was tagged with GFP at its N-terminus, VPS-29 and DPY-23 were tagged at their C-

termini. Point mutations were introduced into *sel-5* and *dpy-23* sequences using the

QuikChange Site-directed Mutagenesis kit (Agilent). Transcriptional reporter P*pgp-12::gfp*

was created by inserting 3410 bp of *pgp-12* promoter upstream of GFP in pPD95.81. P*egl-*

20::mCherry::PH was constructed by Gibson assembly using P*wrt-2::gfp::PH* (Wildwater et

al., 2011) and pCFJ104 (Frøkjaer-Jensen et al., 2008) (a gift from Erik Jorgensen (Addgene

plasmid # 19328 ; http://n2t.net/addgene:19328 ; RRID:Addgene_19328)) as PCR templates

to amplify PH domain and mCherry, respectively. To create extrachromosomal arrays

constructs were microinjected into distal gonads of young adults using inverted microscope

Leica DMi8 equipped with DIC filters and microinjection system InjectMan® 4 and

FemtoJet® 4i (Eppendorf). Microinjection mixtures contained 10 ng/µL of the plasmid of

interest, 5 ng/µL P*myo-2::tdTomato* as a co-injection marker and 135 ng/µL pBluescript as a

carrier DNA.

CRISPR/Cas9 SEC knock-in method (Dickinson et al., 2015) was used to generate GFP

tagged endogenous *sel-5* locus and performed virtually as described (Dickinson et al., 2015

and http://wormcas9hr.weebly.com/protocols.html). 647 bp upstream of *sel-5* ATG (5'

homology arm) and 558 bp *sel-5* genomic sequence starting from ATG (3' homology arm)

were PCR-amplified from N2 genomic DNA with primers harbouring overhangs for Gibson

assembly with the pDD282 vector (Dickinson et al., 2013) (a gift from Bob Goldstein

(Addgene plasmid # 66823; http://n2t.net/addgene:66823; RRID:Addgene_66823)) and

assembled with pDD282 digested with ClaI and SpeI. Sequence

GCTGAAAAGCCCTAGAGGCA was inserted into pDD162 vector for sgRNA expression

(Dickinson et al., 2013) (pDD162 was a gift from Bob Goldstein (Addgene plasmid # 47549;

http://n2t.net/addgene:47549 ; RRID:Addgene_47549)). Injection was performed as described

above, injection mix contained 10 ng/μl pDD282 with homology arms, 50 ng/μl pDD162 with

corresponding sgRNA sequence, 5 ng/μl P*myo-2::tdTomato*. Line expressing P*sel-5::gfp* was

first established, several L1 animals were then exposed to a heat shock for 4 hours at 34°C to

excise the selection SEC cassette and create a P*sel-5::gfp::sel-5* knock-in line. The EGL-

20::GFP knock-in strain was generated as described (Pani and Goldstein, 2018) with one

modification – the pDD282 vector was used for cloning of the homology arms.

RNA interference experiments were conducted by feeding using bacterial strains from the

Ahringer library (Kamath et al. 2003) except for *dpy-23*, which came from the Vidal library

(Rual et al., 2004). L4 larvae were transferred to RNAi plates with the desired bacterial clones

and the effect of RNAi was assessed in the next generation after 3-4 days at 20°C.

Protein isolation and Western blotting

For MIG-14::GFP detection, gravid hermaphrodites were subjected to hypochlorite treatment

and released embryos were left to hatch overnight in M9 buffer. Larvae were collected and

washed twice in M9 buffer. Pelleted larvae were then resuspended in TX-114 buffer (25 mM

Tris–HCl pH 7.5, 150 mM NaCl, 0.5 mM CaCl2, 1% TX-114 and cOmplete® protease

inhibitors (Roche)), snap frozen and then ground in liquid nitrogen. Thawed lysates were

centrifuged at 20,800 g for 30 minutes at 4°C, mixed with Laemmli sample buffer, separated

on 8% SDS–PAGE and transferred onto Amersham™ Protran® Premium Western blotting

nitrocellulose membrane. GFP was detected with monoclonal anti-GFP antibody (cat.#

11814460001, Roche, RRID: AB_390913) and equal loading was assessed by staining with

monoclonal anti α-tubulin antibody (cat. # T9026, Sigma Aldrich, RRID: AB_477593).

Secondary goat anti-mouse HRP-conjugated antibody (cat.# 115-035-146, Jackson Immuno-

Research Laboratories, RRID: AB_2307392) or goat anti-rabbit HRP-conjugated antibody

(cat.# 111-035-045, Jackson Immuno-Research Laboratories, RRID: AB_2337938) and

WesternBright ECL HRP substrate (cat.# K-12045-D20, Advansta) were used to visualize the

signal. Images of membranes were taken on ImageQuant (LAS4000) and Fiji Gels plug-in

was used for subsequent densitometric analysis. For endogenous phosphorylated DPY-23 and

DPY-23::GFP detection, 200 L4 larvae or young adults positive for DPY-23::GFP were

collected into M9 buffer for each analysed strain. Animals were washed once with M9 buffer

and two times with M9 buffer with 0.001% Triton X-100. 33 μL of animal pellet in

M9/0.001% Triton X-100 was mixed with 33 μL of 4X Laemmli buffer, 4 μL of 20x

PhosStop® phosphatase inhibitors (Roche), 4 μL of 20x cOmplete® protease inhibitors

(Roche) and 1.6 μL of 1 M DTT. Samples were snap frozen in liquid nitrogen, thawed on ice,

and sonicated with 2x 10 pulses at 0.8 amplitude, 0.85 duty cycle with UP50H ultrasound

processor (Hielscher). Samples were heated to 99 °C for 5 minutes and centrifuged at 20,000

g for 10 minutes. 20 μL of each sample were loaded on 9% SDS-PAGE, separated and further

processed as above using anti-GFP and anti-phosphoAP2M1 (cat. # ab109397, Abcam,

RRID: AB_10866362) antibodies.

Caenorhabditis elegans **phenotypes, microscopy and statistical analyses**

Final position of QL.paa (PVM – QL.d phenotype), polarity of ALM and PLM neurons was

assessed in L4 larvae carrying transgene *muIs32* or *muIs35*. PVM migration was scored as

defective when PVM was located anteriorly to the posterior edge of the vulva. ALM polarity

was scored as defective when bipolar or reversed neurites were observed. CAN neurons were

visualized with *kyIs4* transgene and their position relative to V3 seam cell was scored. CAN

positioned anterior to V3 was scored as displaced. For DiI staining, L3-L4 well-fed animals

were washed from a plate with M9 buffer and incubated for three hours in 10 μg/ml DiI

solution (D282, ThermoFisher Scientific, dissolved to 2 mg/ml stock solution in

dimethylformamide). Animals were then washed three times in M9 buffer and directly

observed. Excretory cell canals lengths were scored in L4 or young adult animals carrying

integrated P*pgp-12::gfp* transgene (*sIs10089*) or P*pgp-12::gfp* expressed from an

extrachromosomal array (*mamEx11 or mamEx29*). Posterior canals were scored as wild type

when they reached region between the inner edge of posterior gonadal turn and the rectum.

Shortened canal phenotype was graded according to the region within worm body the canal

reached and following landmarks were used for each category: 0 – excretory canal missing

entirely or reaching the inner edge of anterior gonadal turn, 1 – canal reaching anterior

spermatheca, 2 – canal reaching vulva, 3 – canal reaching posterior spermatheca, 4 – canal

reaching inner edge of posterior gonadal turn, 5 – canal reaching rectum (wild type). Canals

which overgrew the rectum and reached the tail tip were scored as 6. Anterior excretory cell

canals were measured and their length normalized to measured distance between the posterior

edge of the pharynx and the nose tip. Larval excretory cell posterior canals were measured at

four developmental stages (hatch, 4, 8 and 24 h after hatching) and normalized to animal body

length. For microscopy imaging worms were anesthetised either with 10 mM sodium azide or

1 mM levamisol and mounted on 3% agarose pads. Images were taken on inverted

microscope Leica DM6. For confocal imaging Zeiss LSM880 confocal microscope was used.

The EGL-20/Wnt gradient was visualized in L2/L3 animals using a spinning disc microscopy

setup, based on a Nikon Eclipse Ti2 equipped with a Yokogawa CSU-W1 scan head, 50 um

pinhole disc, 488 nm laser, Teledyne Photometrics PRIME BSI camera, 60X W NA1.2

objective and operated with NIS elements. For each animal a z-stack of 31 slices, dz=1μm

was acquired. Imaging was done at room temperature (22-23°C). Images were processed with

Fiji image processing package (Schindelin et al., 2012). For EGL-20 gradient quantification, a

maximum projection of 7 consecutive z-sections where the rectal epithelial cells were visible

was made for each sample. A curved line of 150 μm length and 3.9 μm width was fitted

through the ventral side of each animal using the rectum as the starting point and the selected

region was straightened using the "Straighten" function in Fiji. To count the number of EGL-

20 puncta in these selections, images were blinded and the puncta were manually counted.

Counting was repeated three times and the values for each image were averaged. These

averaged values were then plotted and used for statistical evaluation. For mean pixel intensity

calculations a gaussian blur (sigma=3) was applied to the straightened images. Subsequently,

a single segmentation threshold was determined from a representative wild type image by

using the 'graythresh' inbuilt function in Matlab (which uses the Otsu method). This threshold

was multiplied by 1.7 (a factor that was determined manually) in order to generate a mask that

segments only bright EGL-20 spots. Using the derived threshold a binary mask was made for

all images. Using this mask, the mean pixel intensity of all pixels inside the segmented EGL-

20 spots was computed. Statistical analyses were performed either in GraphPad Prism or in

Excel equipped with the Real Statistics Resource Pack software (Release 7.6). Copyright

(2013 – 2021) Charles Zaiontz, www.real-statistics.com.

Materials availability

Strains and plasmids generated in this study are available upon reasonable request from academic researchers.

Acknowledgements

We thank Gunther Hollopeter and Gwendolyn Beacham (Cornell University, USA) for sharing the *dpy-23(mew25)* allele. We thank Jan Mašek for critically reading the manuscript and Gunther Hollopeter for helpful discussion. Some strains were provided by the Caenorhabditis Genetics Center (CGC) which is supported by the National Institutes of Health - Office of Research Infrastructure Programs (P40 OD010440). This work was funded by Charles University Grant Agency grant 1446218/2018 to F.K., by Czech Science Foundation grant 16-17966Y to M.M., by Lumina quaeruntur grant LQ200522301 of the Czech Academy of Sciences to T.C.M. and by Charles University programme SVV 260559. Microscopy was performed in the Vinicna Microscopy Core Facility co-financed by the Czech-BioImaging large RI project LM2023050. The authors acknowledge the Imaging Methods Core Facility at BIOCEV, institution supported by the MEYS CR (LM2023050 Czech-BioImaging), for their support and assistance on the spinning disc imaging done in this work. Computational resources were supplied by the project "e-Infrastruktura CZ" (e-INFRA LM2018140) provided within the program Projects of Large Research, Development and Innovations Infrastructures.

References

Ackley, B. D. (2014). Wnt-signaling and planar cell polarity genes regulate axon guidance along the anteroposterior axis in C. elegans. *Developmental Neurobiology*, *74*(8), 781–796. https://doi.org/10.1002/dneu.22146

Agajanian, M. J., Walker, M. P., Axtman, A. D., Ruela-de-Sousa, R. R., Serafin, D. S., Rabinowitz, A. D., Graham, D. M., Ryan, M. B., Tamir, T., Nakamichi, Y., Zuercher, W. J., & Major, M. B. (2019). WNT Activates the AAK1 Kinase to Promote Clathrin-Mediated Endocytosis of LRP6 and Establish a Negative Feedback Loop. *Cell Reports*, *26*(1), 79-93.e8. https://doi.org/10.1016/j.celrep.2018.12.023

Albrecht, L. V., Tejeda-Muñoz, N., & de Robertis, E. M. (2021). Cell Biology of Canonical Wnt Signaling. *Annual Review of Cell and Developmental Biology*, *37*, 369-389. https://doi.org/10.1146/annurev-cellbio-120319-023657

Aspöck, G., Kagoshima, H., Niklaus, G., & Bürglin, T. R. (1999). Caenorhabditis elegans has scores of hedgehog-related genes: Sequence and expression analysis. *Genome Research*, *9*(10), 909–923. https://doi.org/10.1101/gr.9.10.909

Bänziger, C., Soldini, D., Schütt, C., Zipperlen, P., Hausmann, G., & Basler, K. (2006). Wntless, a Conserved Membrane Protein Dedicated to the Secretion of Wnt Proteins from Signaling Cells. *Cell*, *125*(3), 509–522. https://doi.org/10.1016/j.cell.2006.02.049

Bartscherer, K., Pelte, N., Ingelfinger, D., & Boutros, M. (2006). Secretion of Wnt Ligands Requires Evi, a Conserved Transmembrane Protein. *Cell*, *125*(3), 523–533. https://doi.org/10.1016/j.cell.2006.04.009

Belenkaya, T. Y., Wu, Y., Tang, X., Zhou, B., Cheng, L., Sharma, Y. V., Yan, D., Selva, E. M., & Lin, X. (2008). The Retromer Complex Influences Wnt Secretion by Recycling Wntless from Endosomes to the Trans-Golgi Network. *Developmental Cell*, *14*(1), 120–131. https://doi.org/10.1016/j.devcel.2007.12.003

Bhat, J. M., Pan, J., & Hutter, H. (2015). PLR-1, a putative E3 ubiquitin ligase, controls cell polarity and axonal extensions in C. elegans. *Developmental Biology*, *398*(1), 44–56. https://doi.org/10.1016/j.ydbio.2014.11.008

Brenner, S. (1974). The genetics of Caenorhabditis elegans. *Genetics*, 77(1), 71-94. https://doi.org/10.1093/genetics/77.1.71

Buechner, M., Hall, D. H., Bhatt, H., & Hedgecock, E. M. (1999). Cystic canal mutants in Caenorhabditis elegans are defective in the apical membrane domain of the renal (excretory) cell. *Developmental Biology*, *214*(1), 227–241. https://doi.org/10.1006/dbio.1999.9398

Cendrowski, J., Kaczmarek, M., Mazur, M., Kuzmicz-Kowalska, K., Jastrzebski, K., Brewinska-Olchowik, M., Kominek, A., Piwocka, K., & Miaczynska, M. (2020). Splicing variation of bmp2k balances abundance of copii assemblies and autophagic degradation in erythroid cells. *ELife*, *9*, 1–26. https://doi.org/10.7554/ELIFE.58504

Ch'ng, Q., Williams, L., Lie, Y. S., Sym, M., Whangbo, J., & Kenyon, C. (2003). Identification of genes that regulate a left-right asymmetric neuronal migration in Caenorhabditis elegans. *Genetics*, *164*(4), 1355–1367. https://doi.org/10.1093/genetics/164.4.1355

Cho, N. H., Cheveralls, K. C., Brunner, A. D., Kim, K., Michaelis, A. C., Raghavan, P., Kobayashi, H., Savy, L., Li, J. Y., Canaj, H., Kim, J.Y.S., Stewart, E. M., Gnann, C., McCarthy, F., Cabrera, J. P., Brunetti, R. M., Chhun, B. B., Dingle, G., Hein, M.Y., Huang, B., Mehta, S. B., Weissman,

J. S., Gómez-Sjöberg, R., Itzha, D. N., Royer, L. A., Mann, M., & Leonetti, M. D. (2022) OpenCell: Endogenous tagging for the cartography of human cellular organization. *Science*, *375*(6585):eabi6983. https://doi.org/10.1126/science.abi6983

Colozza, G., & Koo, B.-K. (2021). Ub and Dub of RNF43/ZNRF3 in the WNT signalling pathway. *EMBO Reports*, *22*(5). https://doi.org/10.15252/embr.202152970

Conner, S. D., & Schmid, S. L. (2002). Identification of an adaptor-associated kinase, AAK1, as a regulator of clathrin-mediated endocytosis. *Journal of Cell Biology*, *156*(5), 921–929. https://doi.org/10.1083/jcb.200108123

Conner S.D., & Schmid S.L. (2003). Differential requirements for AP-2 in clathrin-mediated endocytosis. *Journal of Cell Biology*, *162*(5), 773–779, https://doi.org/10.1083/jcb.200304069

Coudreuse, D. Y. M., Roël, G., Betist, M. C., Destrée, O., & Korswagen, H. C. (2006). Wnt gradient formation requires retromer function in Wnt-producing cells. *Science*, *312*(5775), 921–924. https://doi.org/10.1126/science.1124856

Dickinson, D. J., Ward, J. D., Reiner, D. J., & Goldstein, B. (2013). Engineering the Caenorhabditis elegans genome using Cas9-triggered homologous recombination. *Nature Methods*, *10*(10), 1028-1034. https://doi.org/10.1038/nmeth.2641

Dickinson, D. J., Pani, A. M., Heppert, J. K., Higgins, C. D., & Goldstein, B. (2015). Streamlined genome engineering with a self-excising drug selection cassette. *Genetics*, *200*(4), 1035–1049. https://doi.org/10.1534/genetics.115.178335

Fares, H., & Greenwald, I. (1999). SEL-5, a serine/threonine kinase that facilitates lin-12 activity in Caenorhabditis elegans. *Genetics*, *153*(4), 1641–1654. https://doi.org/10.1093/genetics/153.4.1641

Franch-Marro, X., Wendler, F., Guidato, S., Griffith, J., Baena-Lopez, A., Itasaki, N., Maurice, M. M., & Vincent, J.-P. (2008). Wingless secretion requires endosome-to-Golgi retrieval of Wntless/Evi/ Sprinter by the retromer complex. *Nature Cell Biology*, *10*(2), 170–177. https://doi.org/10.1038/ncb1678

Frøkjaer-Jensen, C., Davis, M.W., Hopkins, C.E., Newman, B.J., Thummel, J.M., Olesen, S.P., Grunnet, M., & Jorgensen, E.M. (2008). Single-copy insertion of transgenes in Caenorhabditis elegans. *Nature Genetics*, *40*(11), 1375-83. https://doi: 10.1038/ng.248

Fujita, M., Hawkinson, D., King, K. V., Hall, D. H., Sakamoto, H., & Buechner, M. (2003). The role of the ELAV homologue EXC-7 in the development of the Caenorhabditis elegans excretory canals. *Developmental Biology*, *256*(2), 290–301. https://doi.org/10.1016/S0012-1606(03)00040-X

Gibson, D. G., Young, L., Chuang, R.-Y., Venter, J. C., Hutchison, C. A., & Smith, H. O. (2009). Enzymatic assembly of DNA molecules up to several hundred kilobases. *Nature Methods*, *6*(5), 343–345. https://doi.org/10.1038/nmeth.1318

Golkowski, M., Lius, A., Sapre, T., Lau, H. T., Moreno, T., Maly, D.J, & Ong, S. E. (2023). Multiplexed kinase interactome profiling quantifies cellular network activity and plasticity. *Molecular Cell*, 83(5), 803-818.e8. https://doi.org/10.1016/j.molcel.2023.01.015

Goodman, R. M., Thombre, S., Firtina, Z., Gray, D., Betts, D., Roebuck, J., Spana, E. P., & Selva, E. M. (2006). Sprinter: A novel transmembrane protein required for Wg secretion and signaling. *Development*, *133*(24), 4901–4911. https://doi.org/10.1242/dev.02674

Groenendyk, J., & Michalak, M. (2011). A Genome-Wide siRNA Screen Identifies Novel Phospho-enzymes Affecting Wnt/β-Catenin Signaling in Mouse Embryonic Stem Cells. *Stem Cell Reviews and Reports*, *7*(4), 910–926. https://doi.org/10.1007/s12015-011-9265-3

Gupta-Rossi, N., Ortica, S., Meas-Yedid, V., Heuss, S., Moretti, J., Olivo-Marin, J.-C., & Israël, A. (2011). The adaptor-associated kinase 1, AAK1, is a positive regulator of the notch pathway. *Journal of Biological Chemistry*, *286*(21), 18720–18730. https://doi.org/10.1074/jbc.M110.190769

Harfe, B. D., Branda, C. S., Krause, M., Stern, M. J., & Fire, A. (1998). MyoD and the specification of muscle and non-muscle fates during postembryonic development of the C. elegans mesoderm. *Development*, *125*(13), 2479–2488. https://doi.org/10.1242/dev.125.13.2479

Harris, J., Honigberg, L., Robinson, N., & Kenyon, C. (1996). Neuronal cell migration in C. elegans: Regulation of Hox gene expression and cell position. *Development*, *122*(10), 3117–3131. https://doi.org/10.1242/dev.122.10.3117

Harterink, M., Kim, D. H., Middelkoop, T. C., Doan, T. D., van Oudenaarden, A., & Korswagen, H. C. (2011). Neuroblast migration along the anteroposterior axis of C. elegans is controlled by opposing gradients of Wnts and a secreted Frizzled-related protein. *Development*, *138*(14), 2915–2924. https://doi.org/10.1242/dev.064733

Harterink, M., Port, F., Lorenowicz, M.J., McGough, I.J., Silhankova, M., Betist, M.C., van Weering, J.R.T., van Heesbeen, R.G.H.P., Middelkoop, T.C., Basler, K., Cullen, P.J., & Korswagen, H.C. (2011). A SNX3-dependent retromer pathway mediates retrograde transport of the Wnt sorting receptor Wntless and is required for Wnt secretion. *Nature Cell Biology*, *13*(8), 914-923, https://doi.org/10.1038/ncb2281

Hedgecock, E. M., Culotti, J. G., Hall, D. H., & Stern, B. D. (1987). Genetics of cell and axon migrations in Caenorhabditis elegans. *Development*, *100*(3), 365–382. https://doi.org/10.1242/dev.100.3.365

Heppert, J. K., Pani, A. M., Roberts, A. M., Dickinson, D. J., & Goldstein, B. (2018). A CRISPR tagging-based screen reveals localized players in wnt-directed asymmetric cell division. *Genetics*, *208*(3), 1147–1164. https://doi.org/10.1534/genetics.117.300487

Herman, M. A., & Horvitz, H.R. (1994). The Caenorhabditis elegans gene lin-44 controls the polarity of asymmetric cell divisions. *Development*, *120*(5), 1035–1047. https://doi.org/10.1242/dev.120.5.1035

Herman, M. A., Vassilieva, L. L., Horvitz, H. R., Shaw, J. E., & Herman, R. K. (1995). The C. elegans gene lin-44, which controls the polarity of certain asymmetric cell divisions, encodes a Wnt protein and acts cell nonautonomously. *Cell*, *83*(1), 101–110. https://doi.org/10.1016/0092-8674(95)90238-4

Hilliard, M. A., & Bargmann, C. I. (2006). Wnt signals and Frizzled activity orient anterior-posterior axon outgrowth in C. elegans. *Developmental Cell*, *10*(3), 379–390. https://doi.org/10.1016/j.devcel.2006.01.013

Hollopeter, G., Lange, J. J., Zhang, Y., Vu, T. N., Gu, M., Ailion, M., Lambie, E. J., Slaughter, B. D., Unruh, J. R., Florens, L., Florens, L., & Jorgensen, E. M. (2014). The membrane-associated proteins FCHo and SGIP are allosteric activators of the AP2 clathrin adaptor complex. *ELife*, *3*. https://doi.org/10.7554/eLife.03648

Hutter, H. (2019). Formation of longitudinal axon pathways in Caenorhabditis elegans. *Seminars in Cell and Developmental Biology*, *85*, 60–70. https://doi.org/10.1016/j.semcdb.2017.11.015

Huttlin, E. L., Bruckner, R. J., Navarrete-Perea, J., Cannon, J. R., Baltier, K., Gebreab, F., Gygi, M. P., Thornock, A., Zarraga, G., Tam, S., Harper, J. W., & Gygi, S. P. (2021). Dual proteome-scale networks reveal cell-specific remodeling of the human interactome. *Cell*, *184*(11), 3022-3040.e28. https://doi.org/10.1016/j.cell.2021.04.011

Huttlin, E. L., Bruckner, R. J., Paulo, J. A., Cannon, J. R., Ting, L., Baltier, K., Colby, G., Gebreab, F., Gygi, M. P., Parzen, H., Gygi, S. P., & Wade Harper, J. (2017). Architecture of the human interactome defines protein communities and disease networks. *Nature*, *545*(7655), 505–509. https://doi.org/10.1038/nature22366

Jullien-Flores, V., Mahe, Y., Mirey, G., Leprince, C., Meunier-Bisceuil, B., Sorkin, A., & Camonis, J. H. (2000). RLIP76, an effector of the GTPase Ral, interactas with the AP2 complex: Involvement of the Ral pathway in receptor endocytosis. *Journal of Cell Science*, *113*(16), 2837–2844. https://doi.org/10.1242/jcs.113.16.2837

Kadlecova, Z., Spielman, S. J., Loerke, D., Mohanakrishnan, A., Reed, D. K., & Schmid, S. L. (2017). Regulation of clathrin-mediated endocytosis by hierarchical allosteric activation of AP2. *Journal of Cell Biology*, *216*(1), 167–179. https://doi.org/10.1083/jcb.201608071

Kamath, R. S., & Ahringer, J. (2003). Genome-wide RNAi screening in Caenorhabditis elegans. *Methods*, *30*(4), 313–321. https://doi.org/10.1016/S1046-2023(03)00050-1

Krause, M., Fire, A., Harrison, S. W., Priess, J., & Weintraub, H. (1990). CeMyoD accumulation defines the body wall muscle cell fate during C. elegans embryogenesis. *Cell*, *63*(5), 907–919. https://doi.org/10.1016/0092-8674(90)90494-Y

Liégeois, S., Benedetto, A., Michaux, G., Belliard, G., & Labouesse, M. (2007). Genes required for osmoregulation and apical secretion in Caenorhabditis elegans. *Genetics*, *175*(2), 709–724. https://doi.org/10.1534/genetics.106.066035

Lorenowicz, M. J., Macurkova, M., Harterink, M., Middelkoop, T. C., de Groot, R., Betist, M. C., & Korswagen, H. C. (2014). Inhibition of late endosomal maturation restores Wnt secretion in Caenorhabditis elegans vps-29 retromer mutants. *Cellular Signalling*, *26*(1), 19–31. https://doi.org/10.1016/j.cellsig.2013.09.013

Maloof, J. N., Whangbo, J., Harris, J. M., Jongeward, G. D., & Kenyon, C. (1999). A Wnt signaling pathway controls Hox gene expression and neuroblast migration in C. elegans. *Development*, *126*(1), 37–49. https://doi.org/10.1242/dev.126.1.37

Maro, G. S., Klassen, M. P., & Shen, K. (2009). A β-catenin-dependent Wnt pathway mediates anteroposterior axon guidance in C. elegans motor neurons. *PLoS ONE*, *4*(3). https://doi.org/10.1371/journal.pone.0004690

Martinez-Gualda, B., Graus, M., Camps, A., Vanhulle, E., Saul, S., Azari, S., Nhu Tran, D. H., Vangeel, L., Chiu, W., Neyts, J., Vermeire, K., & de Jonghe, S. (2022). Synthesis and evaluation of 3-alkynyl-5-aryl-7-aza-indoles as broad-spectrum antiviral agents. *Frontiers in Chemistry*, *10*. https://doi.org/10.3389/fchem.2022.1058229

McKay, S. J., Johnsen, R., Khattra, J., Asano, J., Baillie, D. L., Chan, S., Dube, N., Fang, L., Goszczynski, B., Ha, E., Zhao, Z., & Moerman, D. G. (2003). Gene expression profiling of cells, tissues, and developmental stages of the nematode C. elegans. In *Cold Spring Harbor Symposia on Quantitative Biology* (Vol. 68). https://doi.org/10.1101/sqb.2003.68.159

Middelkoop, T. C., & Korswagen, H. C. (2014). Development and migration of the C. elegans Q neuroblasts and their descendants. *WormBook : The Online Review of C. elegans Biology*, 1–23. https://doi.org/10.1895/wormbook.1.173.1

Middelkoop, T. C., Williams, L., Yang, P.-T., Luchtenberg, J., Betist, M. C., Ji, N., van Oudenaarden, A., Kenyon, C., & Korswagen, H. C. (2012). The thrombospondin repeat containing protein MIG-21 controls a left-right asymmetric Wnt signaling response in migrating C. elegans neuroblasts. *Developmental Biology*, *361*(2), 338–348. https://doi.org/10.1016/j.ydbio.2011.10.029

Moffat, L. L., Robinson, R. E., Bakoulis, A., & Clark, S. G. (2014). The conserved transmembrane RING finger protein PLR-1 downregulates Wnt signaling by reducing Frizzled, Ror and Ryk cell-surface levels in C. elegans. *Development (Cambridge)*, *141*(3), 617–628. https://doi.org/10.1242/dev.101600

Moreno-Layseca, P., Icha, J., Hamidi, H., & Ivaska, J. (2019). Integrin trafficking in cells and tissues. *Nature Cell Biology*, *21*(2), 122–132. https://doi.org/10.1038/s41556-018-0223-z

Motley, A. M., Berg, N., Taylor, M. J., Sahlender, D. A., Hirst, J., Owen, D. J., & Robinson, M. S. (2006). Functional analysis of AP-2 α and μ2 subunits. *Molecular Biology of the Cell*, *17*(12), 5298–5308. https://doi.org/10.1091/mbc.E06-05-0452

Oosterveen, T., Coudreuse, D. Y. M., Yang, P.-T., Fraser, E., Bergsma, J., Dale, T. C., & Korswagen, H. C. (2007). Two functionally distinct Axin-like proteins regulate canonical Wnt signaling in C. elegans. *Developmental Biology*, *308*(2), 438–448. https://doi.org/10.1016/j.ydbio.2007.05.043

Pan, C.-L., Baum, P. D., Gu, M., Jorgensen, E. M., Clark, S. G., & Garriga, G. (2008). C. elegans AP-2 and Retromer Control Wnt Signaling by Regulating MIG-14/Wntless. *Developmental Cell*, *14*(1), 132–139. https://doi.org/10.1016/j.devcel.2007.12.001

Pan, C.-L., Howell, J. E., Clark, S. G., Hilliard, M., Cordes, S., Bargmann, C. I., & Garriga, G. (2006). Multiple Wnts and Frizzled receptors regulate anteriorly directed cell and growth cone migrations in Caenorhabditis elegans. *Developmental Cell*, *10*(3), 367–377. https://doi.org/10.1016/j.devcel.2006.02.010

Pani, A.M., & Goldstein, B. (2018). Direct visualization of a native Wnt in vivo reveals that a long-range Wnt gradient forms by extracellular dispersal. *eLife*, *7*:e38325. https://doi.org/10.7554/eLife.38325

Partlow, E.A., Baker, R.W., Beacham, G.M., Chappie, J.S., Leschziner, A.E., & Hollopeter, G. (2019). A structural mechanism for phosphorylation-dependent inactivation of the AP2 complex. *eLife*, 8:e50003, https://doi.org/10.7554/eLife.50003

Peng, Y.-H., Yang, W.-K., Lin, W.-H., Lai, T.-T., & Chien, C.-T. (2009). Nak regulates Dlg basal localization in Drosophila salivary gland cells. *Biochemical and Biophysical Research Communications*, *382*(1), 108–113. https://doi.org/10.1016/j.bbrc.2009.02.139

Port, F., Kuster, M., Herr, P., Furger, E., Bänziger, C., Hausmann, G., & Basler, K. (2008). Wingless secretion promotes and requires retromer-dependent cycling of Wntless. *Nature Cell Biology*, *10*(2), 178–185. https://doi.org/10.1038/ncb1687

Prasad, B. C., & Clark, S. G. (2006). Wnt signaling establishes anteroposterior neuronal polarity and requires retromer in C. elegans. *Development*, *133*(9), 1757–1766. https://doi.org/10.1242/dev.02357

Ramesh, S. T., Navyasree, K. V., Sah, S., Ashok, A. B., Qathoon, N., Mohanty, S., Swain, R. K., & Umasankar, P. K. (2021). BMP2K phosphorylates AP-2 and regulates clathrin-mediated endocytosis. *Traffic*, *22*(11), 377–396. https://doi.org/10.1111/tra.12814

Rella, L., Fernandes Póvoa, E. E., & Korswagen, H. C. (2016). The Caenorhabditis elegans Q neuroblasts: A powerful system to study cell migration at single-cell resolution in vivo. *Genesis*, *54*(4), 198–211. https://doi.org/10.1002/dvg.22931

Ricotta, D., Conner, S. D., Schmid, S. L., von Figura, K., & Höning, S. (2002). Phosphorylation of the AP2 μ subunit by AAK1 mediates high affinity binding to membrane protein sorting signals. *Journal of Cell Biology*, *156*(5), 791–795. https://doi.org/10.1083/jcb.200111068

Rual, J.F., Ceron, J., Koreth, J., Hao, T., Nicot, A.S., Hirozane-Kishikawa, T., Vandenhaute, J., Orkin, S.H., Hill, D.E., van den Heuvel, S., & Vidal, M. (2004). Toward improving Caenorhabditis elegans phenome mapping with an ORFeome-based RNAi library. *Genome Research*, *14*(10B), 2162-8. https://doi: 10.1101/gr.2505604

Salser, S. J., & Kenyon, C. (1992). Activation of a C. elegans Antennapedia homologue in migrating cells controls their direction of migration. *Nature*, *355*(6357), 255–258. https://doi.org/10.1038/355255a0

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., Tomancak, P., & Cardona, A. (2012). Fiji: An open-source platform for biological-image analysis. *Nature Methods*, *9*(7), 676–682. https://doi.org/10.1038/nmeth.2019

Shaye, D. D., & Greenwald, I. (2011). Ortholist: A compendium of C. elegans genes with human orthologs. *PLoS ONE*, *6*(5):e20085. https://doi.org/10.1371/journal.pone.0020085

Sherwood, D. R., & Plastino, J. (2018). Invading, leading and navigating cells in caenorhabditis elegans: Insights into cell movement in vivo. *Genetics*, *208*(1), 53–78. https://doi.org/10.1534/genetics.117.300082

Silhankova, M., & Korswagen, H. C. (2007). Migration of neuronal cells along the anterior-posterior body axis of C. elegans: Wnts are in control. *Current Opinion in Genetics and Development*, *17*(4), 320–325. https://doi.org/10.1016/j.gde.2007.05.007

Smythe, E., & Ayscough, K. R. (2003). The Ark1/Prk1 family of protein kinases. Regulators of endocytosis and the actin cytoskeleton. *EMBO Reports*, *4*(3), 246–251. https://doi.org/10.1038/sj.embor.embor776

Song, S., Zhang, B., Sun, H., Li, X., Xiang, Y., Liu, Z., Huang, X., & Ding, M. (2010). A Wnt-Frz/Ror-Dsh pathway regulates neurite outgrowth in Caenorhabditis elegans. *PLoS Genetics*, *6*(8):e1001056, https://doi.org/10.1371/journal.pgen.1001056

Sorensen, E. B., & Conner, S. D. (2008). AAK1 regulates Numb function at an early step in clathrin-mediated endocytosis. *Traffic*, *9*(10), 1791–1800. https://doi.org/10.1111/j.1600-0854.2008.00790.x

Sorrell, F. J., Szklarz, M., Abdul Azeez, K. R., Elkins, J. M., & Knapp, S. (2016). Family-wide Structural Analysis of Human Numb-Associated Protein Kinases. *Structure*, *24*(3), 401–411. https://doi.org/10.1016/j.str.2015.12.015

Spencer, A. G., Orita, S., Malone, C. J., & Han, M. (2001). A RHO GTPase-mediated pathway is required during P cell migration in Caenorhabditis elegans. *Proceedings of the National Academy of Sciences of the United States of America*, *98*(23), 13132–13137. https://doi.org/10.1073/pnas.241504098

Sundaram, M. V., & Buechner, M. (2016). The caenorhabditis elegans excretory system: A model for tubulogenesis, cell fate specification, and plasticity. *Genetics*, *203*(1), 35–63. https://doi.org/10.1534/genetics.116.189357

Tax, F. E., Thomas, J. H., Ferguson, E. L., & Horvitz, H. R. (1997). Identification and characterization of genes that interact with lin-12 in Caenorhabditis elegans. *Genetics*, *147*(4), 1675–1695. https://doi.org/10.1093/genetics/147.4.1675

Tsushima, H., Malabarba, M. G., Confalonieri, S., Senic-Matuglia, F., Verhoef, L. G. G. C., Bartocci, C., D'Ario, G., Cocito, A., di Fiore, P. P., & Salcini, A. E. (2013). A Snapshot of the Physical and Functional Wiring of the Eps15 Homology Domain Network in the Nematode. *PLoS ONE*, *8*(2). https://doi.org/10.1371/journal.pone.0056383

Whangbo, J., & Kenyon, C. (1999). A Wnt signaling system that specifies two patterns of cell migration in C. elegans. *Molecular Cell*, *4*(5), 851–858. https://doi.org/10.1016/S1097-2765(00)80394-9

Wildwater, M., Sander, N., de Vreede, G., & van den Heuvel, S. (2011). Cell shape and Wnt signaling redundantly control the division axis of C. elegans epithelial stem cells. *Development*, *138*(20), 4375-85. https://doi: 10.1242/dev.066431

Wrobel, A. G., Kadlecova, Z., Kamenicky, J., Yang, J.-C., Herrmann, T., Kelly, B. T., McCoy, A. J., Evans, P. R., Martin, S., Müller, S., Honing, S., & Owen, D. J. (2019). Temporal Ordering in Endocytic Clathrin-Coated Vesicle Formation via AP2 Phosphorylation. *Developmental Cell*, *50*(4), 494-508.e11. https://doi.org/10.1016/j.devcel.2019.07.017

Wu, Y. C., Chiang, Y. C., Chou, S. H., & Pan, C. L. (2021). Wnt signalling and endocytosis: Mechanisms, controversies and implications for stress responses. *Biology of the Cell*, *113*(2), 95-106. https://doi.org/10.1111/boc.202000099.

Xin, X., Wang, Y., Zhang, L., Zhang, D., Sha, L., Zhu, Z., Huang, X., Mao, W., & Zhang, J. (2023). Development and therapeutic potential of adaptor-associated kinase 1 inhibitors in human multifaceted diseases. *European Journal of Medicinal Chemistry*, 248:115102. https://doi.org/10.1016/j.ejmech.2023.115102

Yamaguchi, A., Urano, T., Goi, T., & Feig, L. A. (1997). An Eps homology (EH) domain protein that binds to the Ral-GTPase target, RalBP1. *Journal of Biological Chemistry*, *272*(50), 31230–31234. https://doi.org/10.1074/JBC.272.50.31230

Yang, P.-T., Lorenowicz, M. J., Silhankova, M., Coudreuse, D. Y. M., Betist, M. C., & Korswagen, H. C. (2008). Wnt Signaling Requires Retromer-Dependent Recycling of MIG-14/Wntless in Wnt-Producing Cells. *Developmental Cell*, *14*(1), 140–147. https://doi.org/10.1016/j.devcel.2007.12.004

Yang, W.-K., Peng, Y.-H., Li, H., Lin, H.-C., Lin, Y.-C., Lai, T.-T., Suo, H., Wang, C.-H., Lin, W.-H., Ou, C.-Y., Chang, H., & Chien, C.-T. (2011). Nak regulates localization of clathrin sites in higher-order dendrites to promote local dendrite growth. *Neuron*, *72*(2), 285–299. https://doi.org/10.1016/j.neuron.2011.08.028

Yu, J., Chia, J., Canning, C. A., Jones, C. M., Bard, F. A., & Virshup, D. M. (2014). WLS Retrograde transport to the endoplasmic reticulum during Wnt secretion. *Developmental Cell*, *29*(3), 277–291. https://doi.org/10.1016/j.devcel.2014.03.016

Zallen, J. A., Kirch, S. A., & Bargmann, C. I. (1999). Genes required for axon pathfinding and extension in the C. elegans nerve ring. *Development*, *126*(16), 3679–3692. https://doi.org/10.1242/dev.126.16.3679

Zhao, Z., Sheps, J. A., Ling, V., Fang, L. L., & Baillie, D. L. (2004). Expression analysis of ABC transporters reveals differential functions of tandemly duplicated genes in Caenorhabditis elegans. *Journal of Molecular Biology*, *344*(2), 409–417. https://doi.org/10.1016/j.jmb.2004.09.052

Zheng, C., Diaz-Cuadros, M., & Chalfie, M. (2015). Dishevelled attenuates the repelling activity of Wnt signaling during neurite outgrowth in Caenorhabditis elegans. *Proceedings of the National Academy of Sciences of the United States of America*, *112*(43), 13243–13248. https://doi.org/10.1073/pnas.1518686112

Zinovyeva, A. Y., & Forrester, W. C. (2005). The C. elegans Frizzled CFZ-2 is required for cell migration and interacts with multiple Wnt signaling pathways. *Developmental Biology*, *285*(2), 447–461. https://doi.org/10.1016/j.ydbio.2005.07.014

Zinovyeva, A. Y., Yamamoto, Y., Sawa, H., & Forrester, W. C. (2008). Complex network of Wnt signaling regulates neuronal migrations during Caenorhabditis elegans development. *Genetics*, *179*(3), 1357–1371. https://doi.org/10.1534/genetics.108.090290

Table 1. Phenotypes of *sel-5* and *vps-29* single and double mutants

Genotype	ALM (%)[a]	PLM (%)[b]	CAN (%)[c]	Dye filling (%)[d]	Fecundity (n)[e]
Wild type	0.0	0.0	0.6	0.5	280 ± 41
sel-5(ok149)	0.0	0.0	8.9	4.7	236 ± 24
sel-5(ok363)	0.0	0,0	10.9	1.0	249 ± 37
vps-29(tm1320)	0.0	0.0	2.9	4.3	221 ± 28
sel-5(ok149) vps-29	29.9	9.1	17.7	17.4	132 ± 40
sel-5(ok363) vps-29	7.2	2.8	16.3	7.1	159 ± 23

[a] % of animals with axons of one or both ALM reversed/bipolar, n > 60

[b] % of animals with axons of one or both PLM neurons reversed, n > 60

[c] % of CAN neurons located anteriorly of V3 seam cell, n > 40

[d] % of animals with one or both phasmid sensilla not dyed with DiI, n > 40

[e] average number of progeny from 5 hermaphrodites, ± s.d.

Figure 1. Loss of *sel-5* potentiates QL migration defect in retromer and Wnt pathway mutants. A) QL neuroblast lineage and a cartoon indicating the position of terminally differentiated neurons (depicted with empty circles) (adapted from Rella et al., 2016). Aberrant position of PVM neuron, as observed in QL.d migration defect, is highlighted in magenta. B) RNAi against *sel-5* increases the penetrance of *vps-29* QL.d migration defect when compared to the control (L4440) RNAi. No defect is observed in wild type background. C) PVM position in L2 larvae of control (transgene only) and *sel-5 vps-29* double mutant animals. PVM position indicated with white arrowhead, neurons visualized by expression of P*mec-7::gfp* transgene *muIs32*, scale bar represents 100 μm. D) Mutation in *sel-5* results in increased penetrance of QL.d migration defect of *vps-29* and *lin-17*, but not *mig-1* mutants. E) Examples of mild and severe alteration of tail tip morphology in *sel-5 vps-29* double mutants. F) Shortened transcripts are produced from the *sel-5* locus in both *ok363* and *ok149* alleles. Potential protein products resulting from these transcripts are depicted showing the impact of the truncation on the kinase domain (dark grey box). Extra amino acids resulting from a frameshift and thus not present in the wild type protein are also depicted (light grey box). Position of the active site is indicated. For B) and D) results are shown as % of WT and QL.d animals, Fisher's exact test was performed to assess the difference between the samples, Bonferroni correction for multiple testing was applied. For B) a representative RNAi experiment is shown, n > 55 per condition, * p-value < 0.02, additional experiments provided in Figure 1 – figure supplement 1, for D) data combined from three independent experiments are shown, n > 150 animals in total for each strain, ** p-value < 0.002, *** p-value < 0.0002, ns – not significant.

Figure 1 – figure supplement 1. Additional examples of RNAi against *sel-5* in wild type and *vps-29* background. Results are shown as % of WT and QL.d animals, Fisher's exact test was performed to assess the difference between the samples, Bonferroni correction for multiple testing was applied, n > 55 per condition, *** p-value < 0.0002, ns - not significant.

Figure 2. SEL-5 is expressed in multiple tissues and is required cell-nonautonomously for QL.d migration. A) Transgenic rescue of the QL.d migration defect. *sel-5* was expressed under the control of various promoters from an extrachromosomal array in *sel-5 vps-29; muIs32* background and the effect of such expression on QL.d migration was quantified. Comparison was made between animals carrying the transgene and their siblings which have lost the transgenic array. Expression of each transgene in the expected tissue is shown on the right. Results are shown as % of WT and QL.d animals, data are combined from three independent experiments, n > 100 animals in total for each condition. Fisher's exact test was performed to assess the difference between the samples, Bonferroni correction for multiple testing was applied, *** p-value < 0.0001, ns – not significant. Results from additional independent transgenic strains are provided as Figure 2 – figure supplement 1. B) GFP expression driven by an endogenous *sel-5* promoter in the gonad of L1 and adult animals. C) Expression of GFP::SEL-5 fusion protein driven by endogenous *sel-5* promoter in various larval and adult tissues. Localized GFP::SEL-5 expression is indicated by white arrowheads, white arrows point to autofluorescent signal from gut granules. Scale bar represents 20 μm in A) – C).

Figure 2 – figure supplement 1. Transgenic rescue of the QL.d migration defect. Several independent transgenic strains for each rescue construct were tested in *sel-5 vps-29; muIs32* background and the effect of such expression on QL.d migration was quantified. Comparison was made between animals carrying the transgene and their siblings which have lost the transgenic array. Results are shown as % of WT and QL.d animals, data are combined from three independent experiments, n > 100 animals in total for each condition. Fisher's exact test was performed to assess the difference between the samples, Bonferroni correction for multiple testing was applied, ** p-value < 0.001, *** p-value < 0.0001, ns – not significant. Lines used in Figure 2A are labelled in bold.

Figure 3. SEL-5 alters the phosphorylation status of DPY-23 but does not affect MIG-14 levels. A) Level of DPY-23 phosphorylation at position T160 is reduced in *sel-5* mutant animals. Phosphorylation was detected by Western blot analysis in lysates from a population of L4/young adults of indicated strains using phospho-specific antibody against human AP2M1. Band intensities were normalized to an alpha tubulin loading control and compared to the control sample (sample/control ratio indicated below each lane). Representative Western blot example is shown on the left, relative quantification of four independent experiments is shown on the right. B) Level of DPY-23::GFP phosphorylation is reduced in *sel-5* mutants. Phosphorylation at T160 of GFP tagged DPY-23 was detected by Western blot analysis in lysates from a population of L4/young adults using phospho-specific antibody against human AP2M1. Strains expressing either no GFP fusion protein or a GFP tagged DPY-23T160A mutant protein were included as controls. Band intensities were normalized to a GFP signal intensity and compared to the control sample (DPY-23::GFP in wild type background), sample/control ratio is indicated below each lane. Representative Western blot example is shown on the left, relative quantification of three independent experiments is shown on the right. C) L2/L3 animals expressing MIG-14::GFP from the *huSi2* transgene were imaged using confocal microscope. Posterior part of the body with Wnt expressing cells is shown. Images are maximum pojections of 4 consecutive z-sections taken at xxx interval. Anterior to the left, dorsal up, scale bar represents 20 μm. D) Western blot analysis of MIG-14::GFP levels expressed from a *huSi2*

(continued) transgene in various mutant backgrounds. MIG-14::GFP was detected in lysates from synchronized populations of L1 larvae of the indicated strains. Band intensities were normalized to an alpha tubulin loading control and compared to the control sample (sample/control ratio indicated below each lane). Representative Western blot example is shown on the left, relative quantification of five independent experiments is shown on the right. In A, B and D error bars represent mean ± s.d., statistical significance was assessed by unpaired two-tailed Student's t-test for samples with unequal variance, * p-value < 0.05, ** p-value < 0.01, *** p-value < 0.001, ns – not significant.

Pmig-14::mig-14::gfp	Pegl-20::mCherry::PH	merge	
ctrl			
sel-5			
vps-29			
sel-5 vps-29			
dpy-23 RNAi			

Figure 3 – figure supplement 1. MIG-14 does not re-localize to the plasma membrane in *sel-5* or *sel-5 vps-29* mutants. L2/L3 animals expressing MIG-14::GFP from the *huSi2* transgene together with mCherry::PH driven by *egl-20* promoter were imaged using confocal microscope. Posterior part of the body with Wnt expressing cells is shown. Images are single z-sections. Anterior to the left, dorsal up, scale bar represents 20 μm.

Figure 3 – figure supplement 2. EGL-20 gradient formation. A) Endogenously tagged EGL-20::GFP was visualized in various backgrounds. Images were acquired with spinning disc microscopy using L2/L3 animals and represent maximum projections of 7 consecutive z-sections captured at 1 μm interval. A distance of 150 μm from the rectum of each animal is indicated by a yellow line. B) A curved line of 150 μm length (measured from the rectum) and 3.9 μm width was fitted with the ventral side of each animal presented in A) and the selected region was straightened using the "Straighten" function in Fiji for easier comparison. C) Number of EGL-20 particles in various mutant backgrounds. Identical selection as in B) was applied to all samples and the number of clearly visible puncta was manually counted in blinded images. Counting was repeated three times for each sample, the number of puncta was averaged and plotted as a single data point. D) Mean signal intensity in EGL-20 particles was measured in the same selections as in C). For C) and D) Wilcoxon rank sum test was performed to assess statistical significance. Bonferroni correction for multiple testing was applied. * p-value < 0.005,** p-value < 0.001, error bars represent 95% confidence interval, only statistically significant differences shown in C), no statistically significant differences revealed in D).

Figure 4 – SEL-5 kinase activity and DPY-23 phoshorylation are not required for QL.d migration. A) Transgenic rescue of the QL.d migration defect with kinase-inactive SEL-5. D178A or K75A SEL-5 mutant protein was expressed from *egl-20* promoter in *sel-5 vps-29; muIs32* mutant background from an extrachromosomal array and the QL.d migration defect was quantified in animals carrying the transgene and their siblings which have lost the transgenic array. Results are shown as % of WT and QL.d animals, data are combined from three independent experiments, n > 100 animals in total for each condition. Fisher's exact test was performed to assess the difference between the samples, Bonferroni correction for multiple testing was applied, *** p-value < 0.0001. Results from additional independent transgenic strains are provided as Figure 4 – figure supplement 2. B) DPY-23 phosphorylation at position T160 is absent in *dpy-23(mew25)* mutant animals. Phosphorylation was detected by Western blot analysis in lysates from a population of L4/young adults of indicated strains using phospho-specific antibody against human AP2M1. Band intensities were normalized to an alpha tubulin loading control. C) *dpy-23(mew25)* rescues the „jowls" phenotype (white arrow) of *fcho-1(ox477)* animals. Heads of adult animals are shown, anterior to the left, scale bar represents 10 μm. D) The absence of DPY-23 T160 phosphorylation does not result in QL.d migration defect. The presence of *dpy-23(mew25)* allele carrying T160A substitution either alone or in combination with *sel-5* or *vps-29* does not significantly contribute to the QL.d migration defect. Results are shown as % of WT and QL.d animals, data combined from at least three independent experiments are shown, n > 180 animals in total for each strain. Fisher's exact test was performed to assess the difference between the samples, ** p-value < 0.01. E) MIG-14::GFP does not relocalize to the plasma membrane in *dpy-23(mew25)* animals. L3 animals expressing MIG-14::GFP from the *huSi2* transgene were imaged using confocal microscope. Posterior part of the body with Wnt expressing cells is shown. Images are maximum projections of 5 consecutive z-sections taken at 0.3 μm interval. Anterior to the left, dorsal up, scale bar represents 20 μm.

```
hAAK1      46  VTVDEVLAEGGFAIVFLVRTSNGMKCALKRMFV-NNEHDLQVCKREIQIMRDLSGHKNIVGYIDSSINNVSSGDVWEVLI  124
SEL-5      47  VTIEKQIAEGGFAIVYVASDRKNNKFALKRQFTKDNEKQLEACCREHSFLKQCIGHKNIVEFVDSYTNCLGNG-IWECML  125
               **:::  :*********::.    :.  *  **** *. :**::*:.* **  .:::: ****** ::** * :..* :** ::

hAAK1     125  LMDFCRGGQVVNLMNQRLQT--GFTENEVLQIFCDTCEAVARLHQCKTPIIHRDLKVENILL--HDRGHYVLCDFGSATN  200
SEL-5     126  LTEYHQ-KNVLQLMNERISQNQYLTNDEILSIFTDLCEAVSFIHNRPQPIIHRDLKVENVLISSHKPPHYVLCDFGSATT  204
               * :: :  :*::***:*:.      :*::*:*.** * ****: :*:    ***********:*:  *.  ***********.

hAAK1     201  KFQNPQTEGVNAVEDEIKKYTTLSYRAPEMVNLYSGKIITTKADIWALGCLLYKLCYFTLPFGESQVAICDGNFTIPDNS  280
SEL-5     205  QILSVEKYGVEYVKSEVERNTTMCYRSPEMIDFYSGLEIGLKSDIWALGVLLYRLCFFCVPFEESPLAIQSVNYQFPSVP  284
               ::  . :.  **: *:.*::: **:.**:**::*** *  *:****** ***:**:* :** ** :** . *: :*. .

hAAK1     281  RYSQDMHCLIRYMLEPDPDKRPDIYQVSYFSFKLL  315
SEL-5     285  NIPDEIKVLIYMLLDIDVNRRPSIYQTSVLAFE--  317
               . .:::: **  :*: * ::**.***.* ::*:
```

Figure 4 - figure supplement 1. Sequence alignment of the kinase domain of SEL-5 and human AAK1. Residues mutated in *C. elegans* rescue constructs highlighted in grey. Black rectangle marks the conserved HRD motif.

Figure 4 - figure supplement 2. Trangenic rescue of QL.d migraton defect by kinase-inactive SEL-5. Additional independent transgenic strains for each rescue construct were tested in *sel-5 vps-29; muIs32* background and the effect of such expression on QL.d migration was quantified. Comparison was made between animals carrying the transgene and their siblings which have lost the transgenic array. Results are shown as % of WT and QL.d animals, data are combined from three independent experiments, n > 170 animals in total for each condition. Fisher's exact test was performed to assess the difference between the samples, Bonferroni correction for multiple testing was applied, *** p-value < 0.0001. Lines used in Figure 4A labelled in bold.

Figure 5. *sel-5* cooperates with the retromer complex to regulate the length of excretory cell canals.
A) Posterior canals of the excretory cell are significantly shortened in *sel-5 vps-29* mutants. Excretory cell was visualized by P*pgp-12::gfp* expression from a *mamEx11* transgene. Scale bar represents 200 µm. B) Quantification of posterior canal outgrowth defects. The outgrowth of the posterior canal of the

(continued) excretory cell was quantified by dividing the region between the posterior bulb of the pharynx to the tip of the tail into seven segments. Percentage of canal arms terminating in each segment is indicated. Segment scoring 50% or higher was highlighted in dark blue, segment scoring 15 – 49% was highlighted light blue for easier orientation. All strains contained the *mamEx11* transgene to visualize the excretory cell. Statistical significance of differences between the strains was analysed using Fisher's exact test for 2x3 table. For the test, data from segments 0 – 4 were pooled into one category ("shorter"), data from segments 5 and 6 were used as the two other categories ("normal" and "longer", respectively). Comparison was made either to the control strain or to the more severe single mutant in the case of double mutant strains. Bonferroni correction for multiple testing was applied. C) Anterior canals of the excretory cell are shortened in *sel-5 vps-29* double mutants. Excretory cell was visualized by P*pgp-12::gfp* expression from a *mamEx29* transgene. Scale bar represents 100 μm. D) Posterior canals on the right side of the animals are more severely affected compared to their left counterparts. The length of the posterior canal on each side of the animal was measured and normalized to the length of the whole body of the animal. Paired t-test was used to assess the significance of the difference between the two sides, *** p-value < 0.001, ns – not significant. E) Posterior excretory canals are shorter already at the time of hatching in *sel-5 vps-29* mutants. The dynamics of posterior canal outgrowth was assessed by measuring the canal length and normalizing it to the total body length at hatching and at three time points during early larval development. Results are presented as mean + s.d. of at least 30 canals for each condition. Unpaired two-tailed Student's t-test was performed to assess the difference between the samples, * p-value < 0.05, ** p-value < 0.01. F) Loss of either *vps-35*, *vps-26* or *snx-3* retromer component expression induces posterior canal shortening in *sel-5* mutants. Canal outgrowth was scored as in 4B. G) Both cell-autonomous and non-autonomous expression of *sel-5* rescues excretory canal shortening in *sel-5 vps-29* mutants. SEL-5 was expressed from an extrachromosomal array under the control of *pgp-12*, *hlh-1* or *col-10* promoters. Canal outgrowth was scored as in 4B and comparison was made between animals carrying the array and their siblings which have lost the transgenic array. All strains contained also the *mamEx29* extrachromosomal array to visualize the excretory canal. H) T160 phosphorylation of DPY-23 is not required for posterior excretory canal outgrowth. Canal outgrowth was scored as in 4B in strains containing *dpy-23(mew25)* allele carrying T160A substitution. Comparison was made between the double mutants and either *sel-5* or *vps-29* single mutant. For F) – H) † – differences not significant, ‡ – same data as in B).

B)

	0	1	2	3	4	5	6	N	p-value	versus
ctrl	0	0	0	0	1	99	0	248		
egl-20(n585)	0	0	0	0	1	93	6	169	0.0007	*ctrl*
cwn-1(ok546)	0	0	0	0	0	98	2	230	0.1328†	*ctrl*
cwn-2(ok895)	0	1	0	0	6	91	2	176	0.0008	*ctrl*
lin-44(n1792)	0	0	0	0	0	6	94	194	<0.0001	*ctrl*
mig-1(e1787)	0	0	0	0	0	100	0	180	0.5115†	*ctrl*
lin-17(n671)	0	0	0	0	0	14	86	175	<0.0001	*ctrl*
cfz-2(ok1201)	0	0	0	1	12	85	2	177	<0.0001	*ctrl*
mig-14(mu71)	0	0	0	0	3	14	83	182	<0.0001	*ctrl*
pry-1(mu38)	0	0	0	0	3	95	2	187	0.0939†	*ctrl*
dsh-1(ok1445)	0	0	0	0	0	84	16	232	<0.0001	*ctrl*
mig-5(cp385)‡	0	0	0	0	3	65	32	182	<0.0001	*ctrl*
lin-44; egl-20	0	0	0	0	0	10	90	180	0.1262†	*lin-44*
lin-44; cwn-1	0	0	0	0	2	43	55	204	<0.0001	*lin-44*
lin-44; cwn-2	0	0	0	1	2	24	73	181	<0.0001	*lin-44*
cwn-1; egl-20	0	0	0	1	1	96	2	180	0.2201†	*egl-20*
cwn-1; cwn-2	0	1	1	2	4	91	1	179	0.4533†	*cwn-2*
mig-1; egl-20	0	0	0	0	0	91	9	180	0.1998†	*egl-20*
cwn-1; cfz-2	0	0	0	7	29	64	0	178	<0.0001	*cfz-2*
cwn-2; cfz-2	1	1	1	7	19	70	1	189	0.0008	*cfz-2*
lin-17; egl-20	0	0	0	2	11	53	34	178	<0.0001	*lin-17*
mig-1 lin-17	0	0	1	1	9	57	32	180	<0.0001	*lin-17*
mig-1; cfz-2	0	0	0	0	3	97	0	178	<0.0001	*cfz-2*
lin-17; cfz-2	0	1	1	1	8	61	28	180	<0.0001	*lin-17*
lin-17; sel-5(ok149)	0	0	0	0	0	35	65	180	<0.0001	*lin-17*
lin-44; sel-5(ok149)	0	0	0	0	7	43	50	128	<0.0001	*lin-44*

Figure 6. Wnt pathway components determine the length of the posterior excretory canal. A) Posterior canals of the excretory cell overgrow into the tip of the tail in *lin-17* mutants. Excretory cell was visualized by *Ppgp-12::gfp* expression from the *sIs10089* transgene. Boxed areas are magnified in the bottom row, scale bare represents 200 μm. B) Quantification of posterior canal outgrowth in Wnt pathway mutants. Canal outgrowth was scored as in Fig. 5B and comparison was made either to the control strain (*sIs10089* only) or to the more severe single mutant in the case of double mutant strains. All strains contained the *sIs10089* transgene to visualize excretory canals except for *lin-17; sel-5* and *lin-44; sel-5* which contained *mamEx11*, and *mig-5*, which contained *mamEx29*. † – differences not significant. ‡ Full genotype was *mig-5(cp385[mNG-GLO^AID::mig-5]); mamEx29; mamEx34[eft-3p::TIR::mRuby; myo2p::tdTomato]* and animals were grown on NGM plates with 1 mM auxin from L1 till L4/young adult stage. Only animals containing the *mamEx34* array were used for analysis. C) Posterior excretory canals display normal length during early larval development. The dynamics of posterior canal outgrowth in *lin-44*, *lin-17* and *mig-14* mutants was assessed by measuring the canal length and normalizing it to the total body length at hatching and at three time points during early larval development. Results are presented as mean + s.d. of at least 30 measured canals. Comparison between samples assessed by unpaired two-tailed Student's t-test did not reveal any significant differences. D) Three branches of the posterior canal detected in *cwn-1; cfz-2* animals. Excretory canal visualized by the *sIs10089* transgene, scale bar represents 200 μm.

Figure 7. Model summarizing the inputs acting in the excretory cell outgrowth. A) Three possible outcomes of the canal outgrowth depending on the genetic background. B) Model summarizing the role of Wnt pathway components, *sel-5* and the retromer in the outgrowth of the excretory cell posterior canals. Only players discussed in this work are depicted. First, the active outgrowth of the posterior canals is controlled by *sel-5*, retromer and the Wnt pathway consisting of *cwn-1*, *cwn-2* and *cfz-2*. It remains to be determined whether these players act sequentially, in parallel or as components of a single pathway. After reaching the proper length, further canal extension is inhibited by a second Wnt signalling pathway governed by *lin-44*, *lin-17*, *dsh-1* and *mig-5*. Last, *egl-20* and *mig-1* might fine-tune the precise stopping position of the growing canal.