Comammox **and Unknown Candidate AOB Contribute to Nitrite Accumulation in an**

Integrated A-B stage process that Incorporates Side-stream EBPR (S2EBPR)

Yuan Yan[a,#], Jangho Lee[a,#], IL Han[a], Zijian Wang[a], Guangyu Li[a], Kester McCullough[a,b],

Stephanie Klaus[b], Da Kang[a,c], DongQi Wang[a], Anand Patel[a], Jim McQuarrie[d], Beverley M.

Stinson[e], Christine deBarbadillo[f], Paul Dombrowski[g], Charles Bott[b,*], April Z. Gu[a,*]

[a] School of Civil and Environmental Engineering, Cornell University, Ithaca, United States

[b] Hampton Roads Sanitation District, Virginia Beach, United States

[c] Department of Environmental Engineering, College of Environmental & Resource Sciences,

Zhejiang University, China

[d] Denver Metro Wastewater Reclamation District, Denver, Colorado, United States

[e] AECOM, Laurel, Maryland

[f] District of Columbia Water and Sewer Authority, Washington, District of Columbia

[g] Woodard & Curran, Inc, Enfield, Connecticut, United States

[#] Joint first authors

[*] Corresponding authors:

April Z. Gu. aprilgu@cornell.edu, Charles Bott, cbott@hrsd.com

Abstract

A novel integrated pilot-scale A-stage high rate activated sludge, B-stage short-cut biological

nitrogen removal and side-stream enhanced biological phosphorus removal (A/B-shortcut N-

S2EBPR) process for treating municipal wastewater was demonstrated with the aim to achieve

simultaneous and carbon- and energy-efficient N and P removal. In this studied period, an

average of 7.62 ± 2.17 mg-N/L nitrite accumulation was achieved through atypical partial

nitrification without canonical known NOB out-selection. Network analysis confirms the central

hub of microbial community as Nitrospira, which was one to two orders of magnitude higher

than canonical aerobic oxidizing bacteria (AOB) in a B-stage nitrification tank. The contribution

of comammox Nitrospira as AOB was evidenced by the increased amoB/nxr ratio and higher

ammonia oxidation activity. Furthermore, oligotyping analysis of Nitrospira revealed two

dominant sub-clusters (microdiveristy) within the Nitrospira. The relative abundance of

oligotype II, which is phylogenetically close to Nitrospira_midas_s_31566, exhibited a positive

correlation with nitrite accumulation in the same operational period, suggesting its role as

comammox Nitrospira. Additionally, the phylogenetic investigation suggested that heterotrophic

organisms from the family Comamonadacea and the order Rhodocyclaceae embedding ammonia

monooxygenase and hydroxylamine oxidase may function as heterotrophic nitrifiers. This is the

first study that elucidated the impact of integrating the S2EBPR on nitrifying populations with

implications on short-cut N removal. The unique conditions in the side-stream reactor, such as

low ORP, favorable VFA concentrations and composition, seemed to exert different selective

forces on nitrifying populations from those in conventional biological nutrient removal

processes. The results provide new insights for integrating EBPR with short-cut N removal

process for mainstream wastewater treatment.

1 Introduction

The adsorption/bio-oxidation (A-B) process is a promising engineering application towards carbon capture, energy recovery, and efficient nutrient removal. It diverts organic carbon to anaerobic digestion and thus manipulates the C/N in B-stage reactors through carbon flow redirection (Ren and Pagilla, 2022). The first stage (A-stage) is designed for high-rate carbon capture and removal using high-rate activated sludge (HRAS), which is widely operated with a low hydraulic retention time (HRT), a low sludge retention time (SRT) and a low dissolved oxygen (DO) concentration less than 0.5 mg L^{-1} (Boehnke et al., 1997). After HRAS, the wastewater COD/N ratio could be lowered to 0.67 (Lotti et al., 2015), which is suitable for second stage (B-stage) nutrient (mostly nitrogen), like short-cut nitrogen removal processes, to achieve "carbon-free" nitrogen removals (Sancho et al., 2019).

Partial nitrification with anammox (PN/A) process has emerged to be one of the promising short-cut N removal system, which can save 100% carbon, 60% aeration and reduce 80% sludge production (Jetten et al., 1997). It consists of two step processes where aerobic ammonia-oxidizing bacteria (AOB) oxidize half of the ammonia to nitrite and, then anaerobic ammonia-oxidizing bacteria (anammox) bacteria convert ammonia and nitrite into dinitrogen. The out-selection of nitrite oxidizing bacteria (NOB) in B-stage is crucial for PN/A process and it has been successfully demonstrated for side-stream high-strength wastewater (Lackner et al., 2014). A number of operational strategies have been suggested to enable side-stream NOB out-selection and consequently nitrite accumulation, including high temperature, low DO, low SRT, high pH, and high free nitrous acid (FNA) and/or free ammonia (FA) concentration (Hellinga et al., 1998; Joss et al., 2009; Van Dongen et al., 2001; Xu et al., 2012). However, these strategies were not readily applicable in mainstream treatment since the inhibitory levels of 0.02 mg/L FNA and 1

mg/L FA is unavailable with only 50 mg-N/L ammonia at neutral pH in the mainstream (Wang

et al., 2017). Also, it has been shown that NOB can adapt to high concentration of FA (An et al.,

2021) and low DO (Liu and Wang, 2013), which makes it harder to be out-selected in

mainstream process. Therefore, maintaining long-term stable NOB out-selection still remains a

challenge for main-stream wastewater treatments and the underlined mechanisms need to be

explored.

Most PN/A studies focus only on nitrogen removal, however, phosphorus co-occur with nitrogen

and is also a critical nutrient that needs to be treated in wastewater (Rittmann et al., 2011).

Enhanced biological phosphorus removal (EBPR) process has been widely applied worldwide in

full-scale facilities. Its performance and stability strongly rely on the activity of phosphorus

accumulating organisms (PAOs) with favorable influent rbCOD/TP ratio >15 (Gu et al., 2008).

Therefore, EBPR is considered incompatible with B-stage autotrophic nitrogen removal due to

the insufficiency of readily biodegradable carbon in A-stage effluent (B-stage influent). The

implementation of a side-stream anaerobic biological sludge hydrolysis and fermentation reactor

called side-stream EBPR (S2EBPR) can be an alternative option to address the carbon shortage

challenge of coupling EBPR with B-stage nitrogen removal process. S2EBPR takes advantage of

the volatile fatty acids (VFAs) generated via hydrolysis/fermentation of side-stream return

activated sludge (RAS) and mixed liquor, and thus overcome the reliance on the influent

carbon (Barnard et al., 2017; Onnis-Hayden et al., 2020; Gu et al., 2018; Wang et al., 2019).

However, the influence of implementing S2EBPR on the PN/A removal system, especially on

the nitrite accumulation remains unknown. We hypothesize that the incorporation of S2EBPR

would affect the nitrifying populations because of the extended anaerobic conditions in the side-

steam reactor with low oxidation reduction potential (ORP <300 mV) and high VFAs. In

addition, PAOs and the surplus of sludge fermentate from the S2EBPR reactor to the down-

stream B-stage would compete oxygen with nitrifiers (Carvalheira et al., 2014).

Recently, our team reported a successful incorporation of S2EBPR into a A-B stage process to

achieve simultaneous nitrogen and phosphorus removal for main-stream treatment (McCullough

et al., 2023). This pilot-scale system combined A-stage HRAS, B-stage short-cut biological

nitrogen removal process, and S2EBPR, referring as A/B-shortcut N-S2EBPR herein, provides

us a good opportunity to explore the interacting impacts of PAOs and nitrogen removal related

microorganisms. In this study, we focused on the comprehensive examination of the nitrite

accumulation and AOB and NOB population dynamics in B-stage processes that generated

nitrite with intermittent aeration AvN (ammonia versus NO_x^--N (nitrite+nitrate) control strategy.

Process performance, batch activity tests were performed to investigate the underlying processes

involved in the nitrite accumulation. 16S rRNA gene amplification sequencing and quantitative

PCR were applied to reveal the microbial community composition and dynamics with emphasis

on nitrification-relevant populations. To our knowledge, this is the first study that revealed the

influence of implementing S2EBPR on B-stage nitrification population and, provided new

insights on the distinctive nitrifying populations that contributed to nitrite accumulation in this

A/B-shortcut N-S2EBPR process that is different from those typically observed in previous

WWTPs. The results pointed out a possible new strategy in achieving nitritation to enable

anammox for main-stream municipal WWTPs.

2   Material methods

2.1   Pilot reactor setup and Operation conditions

The pilot plant operated by Hampton Roads Sanitation District (HRSD) was located at

Chesapeake-Elizabeth WWTP in Virginia Beach, VA, USA. It consisted of an A-stage HRAS

process followed by the B-stage process, a parallel S2EBPR reactor and a final stage anammox

polishing tank (Figure S1). A portion of A-stage wasted activated sludge (WAS) was split into a

combined fermenter and thickener. The B-stage consisted of four CSTRs bioreactors in series,

each with independent aeration control via AvN control (Regmi et al., 2014). In the studied

period, B-stage reactors were switched from clarifier wasting to aeration tank wasting (hydraulic

wasting) and recycling. A portion of B-stage RAS and the A-stage WAS fermentate were added

to the side-stream reactor for PAOs selection. The detailed parameters of A-B process have been

introduced elsewhere (McCullough et al., 2023).

Chemical analysis of soluble COD, OP, total ammonia nitrogen (NH_4^+-N + NH_3-N), NO_2^--N,

and NO_3^--N, volatile fatty acids (VFA) including acetic, propionic, butyric, isobutyric, valeric,

isovaleric, and caproic acids were summarized in Text S3.

2.2   AOB and NOB activity analysis

AOB and NOB activities were analysis weekly on site. About 4L mixed liquor suspended solids

(MLSS) were withdrawn from the last B-stage aerobic tank and pre-aerated for 30 min to oxidize

the residual carbon. Ammonium chloride was spiked to reach 25 mg-N/L final concentration.

DO and pH were maintained around 3-5 mg/L and 7, respectively. The samples were taken every

15 minutes for a total of 1 hour. After filtered by 0.45 μm filter, the filtrate was used to analyze

the NH_4^+-N, NO_2^--N, and NO_3^--N concentrations using HACH TNT kits (HACH Loveland,

CO). AOB activity was calculated as the sum of nitrite and nitrate production rates, and NOB

activity was calculated as the nitrate production rates by linear regression.

2.3   Quantification of nitrifying candidates

Quantitative polymerase chain reaction (qPCR) was performed on a CFX96 real-time PCR

detection system (Bio-Rad, USA) to determine the concentrations of functional genes including

amoA, *nxrB*, *amoB* and 16S. The amoA-1F/2R primer for the canonical ammonia

monooxygenase (AMO) enzymes in AOB (Rotthauwe et al., 1997), and the *nxrB* 169F/638R

primer for the beta subunit of nitrite oxidoreductase (NXR) (Pester et al., 2014) were used to

target canonical AOB and NOB. The *amoB* 148F/485R primer were used to target the beta

subunit of AMO in *Nitrospira* that can do complete ammonia oxidation (comammox) (Cotto et

al., 2020). All functional genes were then normalized by 16S rRNA with primer set 341f/534r

(He et al., 2007). Calibration curves were obtained using 10-fold serial dilutions of known

concentrations of cloned DNA samples that were prepared following the instruction from TOPO

TA cloning kit (Thermo Fisher, USA) in our laboratory. The R^2 of standard curves were above

0.99 and the amplification efficiency was around 95%. The detailed qPCR reaction conditions

were summarized in Table S1.

2.4   Bioinformatics and statistical analysis

2.4.1   Microbial population analysis

Sludge samples from B-stage were monthly collected for DNA extraction. Genomic DNA was

extracted using Instagene matrix (Kim et al., 2013) and then used for both 16S ribosomal RNA

(16S rRNA) sequencing and functional genes quantification.

The V4 region of the 16S rRNA genes were amplified using primers 515F-Y (Parada et al.,

2016) and 926R (Quince et al., 2011). Detailed amplification process is listed in text S2. To

minimize the inflation of rare species in the community analysis, we removed the operational

taxonomic units (OTUs) that are lower than 0.01%. The filtered-OUT matrix was used to

perform statistical analyses in R-4.2.3. The matrix was also used to predict the bacterial

metagenomes using Phylogenetic Investigation of Communities by Reconstruction of

Unobserved States 2 (PICRUSt2) pipeline (Douglas et al., 2020) with implemented tools EPA-ng

(Barbera et al., 2019), gappa (Czech et al., 2020), and castor (Louca and Doebeli, 2018). The

network analysis of microbial community and the correlation between the environmental

variables and the most abundant 50 taxa were done through microbiomeSeq package (Ssekagiri

et al., 2017; Torondel et al., 2016). All bioinformatic figures were generated with ggplot2

package (Villanueva and Chen, 2019, p. 2).

2.4.2   Oligotyping analysis

Oligotyping analysis was performed according to (Eren et al., 2013; Srinivasan et al., 2021) to

reveal the sub-clusters (i.e. microdiveristy) within the dominant nitrifying population. A total of

65,24316S rRNA gene amplicon sequences classified as *Nitrospira* were extracted and formatted

as required by the oligotyping pipeline. After quality filtering, 18 nucleotide positions were

selected for entropy decomposition (102, 120, 134, 135, 149, 151, 160, 181, 197, 199, 210, 219,

240, 262, 265, 375, 488, 491), which were used to generate a total of 205 raw oligotypes. After

denoising and elimination based on the -s (oligotypes that occurred in at least 3 samples) and -M

(a minimum count of 30 (~0.1%)) parameters, 2 dominant oligotypes, presumably representing

micro-diversity of *Nitrospira* sub-groups, were finally identified. The representative sequence of

the 2 dominant oligotypes were BLAST'd against a custom database built from MiDAS 4.8.1 for

top hits at 99% identity threshold (Dueholm et al., 2022).

3   Results and Discussion

3.1   Simultaneous nitrite accumulation and efficient P removal in A/B-shortcut N-

S2EBPR process

The pilot was designed to perform nitrogen and phosphorus removal simultaneously. To

overcome the shortage of influent VFA and enable bio-P removal in main-stream treatment line,

a side-stream reactor was implemented to facilitate bio-P removal through receiving

supplemental carbon from A-stage sludge fermenter (Figure S1). With the increase in the A-

stage WAS fermentate supplement flow and RAS slit ratio to the S2EBPR reactor (Figure S2),

the nitrite accumulation was observed over a period of approximately 150 days (Figure 1).

During this period, the effluent nitrite concentration raised to a maximum level of 6.55 mg-N/L

with an average concentration of 2.65 ± 1.89 mg-N/L (Figure 2A). The average nitrite

accumulating ratio (NAR) was 49%, suggesting that a significant nitrite accumulation

occurred. Figure 2A shows the N mass balance in the system and changes of the various N

species along the B-stage reactors in series. As ammonia decreased along the B-stage zones, the

nitrite began to accumulate from 3.9 g/d to 20.7 g/d (Figure 2A). Previous research indicated that

nitrite can inhibit PAO activity by increasing maintenance energy and disturbing the TCA cycle

(Zhou et al., 2011). Nevertheless, our results indicated an efficient P removal (77%-92%) could

be achieved concurrently with nitrite accumulation (0.2 ± 0.05 - 2.7 ± 1.9 mg-N/L) during this

same period (Text S1, Figure S2).

3.2   Nitrite accumulation due to partial nitrification

Since nitrite can be accumulated either through partial nitrification (PN) or partial denitrification (PDN) pathways, both AOB/NOB activities were evaluated over long-term on site to identify the dominant mechanisms involved in the nitrite accumulation (Figure 1). Examination of the N species during the aerobic/anoxic cycle showed that nitrite only accumulated in aeration periods while it was consumed after aeration was off (Figure 2B), suggesting that nitrite accumulation was likely due to PN rather than PDN. This was further supported by the AOB and NOB activities assessment. During the study period, the average AOB and NOB activities were 1.9 ± 0.9 and 1.1 ± 1.0 mgN/gSS/h, respectively (Figure 1). AOB activities were higher than the NOB activities during the period of study, consistent with the nitrite accumulation via PN (Figure 1). This evidence led to our conclusion that the higher nitrite accumulation in this study period is likely attributed to, at least mostly, PN.

3.3   Unknown AOB that contributed to the High AOB activity

3.3.1   Low and negligible known AOB, NOB abundance

Temporal Microbial communities via 16S rRNA gene sequencing analysis revealed the microbial community structure dynamics and patterns during the study period (Figure 3). The average relative abundance of known AOB, including *Nitrosomonas* and *Nitrosomonadaceae* were remarkably low at 0.07% and 0.2%, respectively. The only known NOB is *Nitrospira*, has a relative abundance ranging from 1.7-6%. Their abundances were depicted in Figure 1. to show the variation trends. In our system, the relative abundances of both known AOB and NOB are significantly lower than that typically reported in mainstream PN/A systems (0.23-4.4% and 0.2-53%) (summarized in Table S2). Particularly interesting, the relative abundance of *Nitrospira*

was much higher than the known AOB, which did not align with the higher ammonia oxidation

activity compared to nitrite oxidation activity.

To further verify the abundances of nitrifiers in the sludge, the *amoA* genes in AOB and the *nxr*

genes in NOB were quantified and normalized to 16S rRNA genes (Figure 3B). The average

nxr/16S ratio was $8.3\text{x}10^{-3}$ in this study period. Generally, *Nitrospira* genome harbor 1-6 copies

of *nxr* (Palomo et al., 2018; Pester et al., 2014), while AOB in betaproteobacteria contains 2-3

copies of *amoA* (Norton et al., 2002). Considering the copy difference, the abundance of *nxrB*

was still one order of magnitude higher than *amoA,* suggesting a higher known *NOB* abundance

than known AOB abundance (Figure 3B). Both 16S sequencing and functional gene

quantification confirmed that the abundance of known AOB (*Nitrosomonas* and

Nitrosomonadaceae) was lower than NOB (*Nitrospira*) throughout the treatment periods, yet the

system still achieved PN. In addition, the correlation analysis demonstrates that the microbial

community changes have significant impact on the effluent nitrite accumulation (Figure S5).

This result implies either the low abundance of nitrification communities played a significant

role or some group of organisms carried out ammonia oxidation yet was unknown to us.

Theoretically, the numerical ratio of AOB to NOB in a balanced nitrifying system should be

higher than 2:1 (Winkler et al., 2012). However, the current investigation shows a disparity

between the rates of nitrite and ammonia oxidation, and the abundance of nitrifying biomass.

Similar observations have been reported before (Wang et al., 2015; Fitzgerald et al., 2015; Zhang

et al., 2018; Gao et al., 2021). Wang et al. (2015) observed NOB:AOB ratios from 9:1 to 5000:1

in five activated sludge reactors through qPCR. Zhang et al. (2018) achieved stable nitrite

accumulation with higher abundance of *Nitrospira* comparing to AOB. Several scenarios can

potentially explain the higher relative abundance of known NOB but lower nitrite oxidation

activity in our system: 1) there might be a group of *Nitrospira* who can perform not only nitrite

oxidation, but also ammonia oxidation, such as complete ammonia oxidizer (comammox); 2)

underestimation of AOB in the system due to unknown AOB; and 3) some substances in

wastewater could have more severe inhibitory effects on NOB than AOB.

3.3.2   Comammox *Nitrospira* may contribute to nitrite accumulation.

Roots et al., (2019) reported that the abundance of Comammox *Nitrospira* was much higher than

both AOB and ammonia oxidizing archaea (AOA) in their in nitrification reactor (Roots et al.,

2019), challenging the commonly held assumption that comammox play a minor role in

wastewater treatment bioreactors. Among the versatile *Nitrospira* genus, the comammox

Nitrospira encode a phylogenetically distinct *amo* gene from the canonical AOB and AOA,

allowing them to fully oxidize ammonia to nitrate (Daims et al., 2016; Koch et al., 2015). High

presence of comammox was reported in nitrite accumulating condition, indicating a potential

correlation between the comammox and nitrite accumulation, given the higher ammonia

oxidation rate than the nitrite oxidation rate in comammox *Nitrospira* (Shao and Wu, 2021). A

recent isotope labelling test has proven that comammox species (*N. nitrosa* and *N. nitrificans*)

provide nitrite for anammox bacteria (*Brocadia*) (van Kessel et al., 2015). The pure cultured *N.*

inopinata was recorded to accumulate nitrite during complete nitrification (Kits et al., 2017).

Given the function versatility of comammox species and the high abundance of *Nitrospira*, it's

possible that a portion of the comammox *Nitrospira* in our system functioned as AOB, leading to

the accumulation of nitrite.

The presence of comammox *Nitrospira* in our system was determined by quantifying the *amoB*

gene that was specifically found in comammox (Cotto et al., 2020; Roots et al., 2019) (Figure

3B). Results indicated an occurrence of higher abundance of comammox *Nitrospira* and nitrite

accumulation. In addition, *amoB* was present and remained much more stable than both *nxrB* and

amoA (Figure 3B). The average of *amoB* ratio was 7.8×10^{-4} per 16S gene copies. The *amoB*/*nxrB*

ratio was deployed to indicate the relative abundance of comammox *Nitrospira* to the total

Nitrospira. Results clearly indicate that comammox *Nitrospira* abundance increased from 1% to

13.6% during nitrite accumulation period. Considering that *Nitrospira* genome contains 1-5

times more copies of *nxr* (1-2 copies in most comammox genomes but 4 copies in *N. nitrificans*

and 5 copis in *N. mosocoviensis*) than *amo* (single copy) (Palomo et al., 2018; Pester et al.,

2014), the relative abundance of comammox *Nitrospira* to the total NOB ranges from 13.6% to

68% in our system.

The increasing abundance of comammox *Nitrospira* motivated us to employ the oligotyping

analysis to further elucidate the micro-diversity of *Nitrospira*. This analysis identified 3

dominant oligotypes of *Nitrospira* and revealed their changes in microbial communities over

time (Figure 4B).

Though the oligotype 1 consistently dominated the *Nitrospira* ($> 71\%$), oligotype II gradually

increased in tandem with the increasing nitrite accumulation. The similar increasing trends of

amoB/*nxrB* and the relative abundance of oligotype II indicates a positive relationship between

the comammox and the oligotype II *Nitrospira*. Note that comammox *Nitrospira* do not form

a monophyletic group in 16S rRNA gene sequence or *nxrB*-based phylogenetic analyses, but are

instead interspersed with strict nitrite-oxidizing *Nitrospira* (Daims et al., 2015; van Kessel et al.,

2015). Different comammox *Nitrospira* encode *amo* orthologs that dissimilar to each other, with

only ~60% amino acids identity between members of the different clades (Daims et al., 2015),

making it difficult to design a uniform comammox qPCR primers (Beach and Noguera, 2019;

Cotto et al., 2020). The *amoB* primer set used in this study was specifically optimized for clade

A comammox (Cotto et al., 2020). However, it is possible that clade B comammox also played a

role in our S2EBPR-PN/A processes, but current primers did not capture the whole comammox

community. Nevertheless, oligotyping and qPCR analysis indicated that a group of strict nitrite-

oxidizing *Nitrospira* decreased, while another group of *Nitrospira*, which might belong to

comammox and can oxidize ammonia increased, partially contributing to the nitrite accumulation

($r^2 = 0.60$, Figure 4B). BLASTing the two major oligotypes of *Nitrospira* again MiDAS4

database, indicating that all sequences in oligotype I were identified as *Nitrospira defluvii*

(Figure 4C). By summarizing the individual taxonomic identifications of all sequences in

oligotype II subcluster, oligotype II is closer to Nitrospira_midas_s_31566 than

Nitrospira_midas_s_64422. Considering the positive correlation of the nitrite accumulation and

oligotype II of *Nitrospira*, it is possible that Nitrospira_midas_s_31566 is a candidate

comammox *Nitrospira*.

3.3.3   Atypical ammonia oxidation bacteria

Fitzgerald et al. (2015) had proposed that the abundance of ammonia oxidizing organisms may

be underestimated, particularly in low DO condition (Fitzgerald et al., 2015). Despite the

presence of complete nitrification to nitrate, no known AOB were identified in their low DO

nitrification systems. To address this issue, they did pure culture and found 5 isolates that had not

previously been recognized as AOB but can oxidize NH_4 under both autotrophic and

heterotrophic conditions. They proposed that genus *Pseudomonas* and *Rhodococcus*, and family

Xanthomonadaceae and Sphingomonas are involved in heterotrophic nitrification under low-DO

conditions.

To further explore the potential ammonia oxidizers in our samples, the functional potential of

bacterial assemblages was predicted with PICRUSt2 in both enzymes and pathways. The

dominated enzyme genes are related to carbohydrate metabolism (e.g., glycolysis/gluconeogesis

metabolism), amino acid metabolism, DNA polymerase, and electron transport, which is

conceivable that all those functions are vital for microorganisms. Then we specifically aimed at

genes that can oxidize ammonia into nitrite. However, only three OTUs have the potential to

hold the methane/ammonia monooxygenase subunits (AMO, EC:1.14.18.3 and 1.14.99.39) that

oxidize ammonia into hydroxylamine, and they all belong to unclassified genus under order

Rhodocyclaceae. Another 11 OTUs belonging to Comamonadaceae_unclassified have the

potential to hold hydroxylamine oxidase (HAO, EC: 1.7.2.6) that can further oxidize

hydroxylamine to nitrite (Table S3 and Figure S6). Indeed, AMO and HAO were purified from

heterotrophic denitrifiers back in 1996 (Moir et al., 1996; Wehrfritz et al., 1996), and

heterotrophic nitrification has drawn a lot attention nowadays. 134 isolates were identified at the

genus/species level, and members from family Comamonadaceae and order Rhodocyclaceae

were reported to be aerobic denitrifiers (Duan et al., 2022).

In addition, the co-occurrence microbial patterns during nitrite accumulation periods revealed

that genus *Nitrospira*, midas_g_894, and midas_g_5535 are the main hubs for this specific

microbiome structure (Figure 3C). The latter two belong to the family Saprospiraceae and

Rhodocyclaceae, respectively, and the species in those families have all been shown to play a

role in autotrophic/heterotrophic nitrification (Duan et al., 2022; Mehrani et al., 2022).

Though the relative abundance of OTUs that are predicted to hold *hao* only represents a small

fraction of the entire Comamonadaceae_unclassified group (Figure S6), the prediction guides us

to look into the function played by Comamonadaceae. Though most heterotrophic nitrifiers can

oxidize ammonia to N_2O or N_2 by a full nitrification and denitrification pathway, some of them

are not capable of aerobic denitrification and ends with nitrite (Duan et al., 2022). A test with

enriched Comamonadaceae (~96%) showed ammonium oxidation potential under aerobic

conditions, with nitrite as the main product (Bao and Li, 2017), indicating the importance of

Comamonadaceae in PN. Some research also indicates that they contribute to both nitrogen and

phosphorus removal and can function as PAO (Ge et al., 2015). Our unique coupling of A-B-

stage with S2EBPR created unprecedented selection forces and conditions for both P and N

removal relevant organisms in this system, which would conceivably lead to enrichment of

organisms with versatile functions different from those in conventional EBPR or PDN systems

for main-stream treatment. In the current study, the relative abundance of Comamonadaceae

family remained stable at unusually high relative abundance level. (13%) in the study period

(Figure 3A) and they might contribute to nitrite accumulation.

It's important to note that the predicted functional potential was based on the current database,

which only includes the sequences of known AMO. Considering the fact that the *amo* genes in

comammox *Nitrospira* are phylogenetically distinct from previously identified AMOs (van

Kessel et al., 2015), it is possible that some other groups of bacteria are oxidizing ammonia but

cannot be captured by the current database.

3.4   Possible factors affecting nitrite accumulation

Another possible scenario to explain nitrite accumulation is that NOB activities were more

severely inhibited than AOB. Multiple strategies have been postulated to suppress NOB in PN

reactors, including low DO, transient anoxic condition, FNA inhibition, FA inhibition, and VFA

composition (Eilersen et al., 1994; Jiang et al., 2019; Park et al., 2010; Wang et al., 2014). The

FNA concentrations in both B-stage influent and the mixing zone (B-stage influent + flow from

Side-stream reactor) are nearly zero, which did not provide inhibition pressure on NOB. The DO

ranged between 0.2-1.8 mg/L in 4 intermittently aerated CSTRs, in line with the DO range that is

suitable for PN (Cao et al., 2017; Regmi et al., 2014). However, more and more evidence show

that DO regulation is not reliable to suppress NOB alone (Park et al., 2010).

Compared to the conventional B-stage nutrient removal system, the coupling of S2EBPR with

PN created unique environmental selection conditions differ from those in conventional EBPR or

short-cut N system without EBPR. A split of RAS will flow through S2EBPR, where the unique

feature of S2EBPR system exerted different condition for nitrifying populations since they were

exposed to extended longer anaerobic time (HRT of anaerobic time is 3.65 hrs.) with lower ORP

(-383 ± 17 mv) in comparison to conventional nutrient removal processes where AOB and NOB

generally experience shorter anoxic conditions (1-2 hrs) with shallow ORP level (~-150 mv). In

addition, high level (126.7 ± 72.1 mg/L as COD) and more complex VFA (Figure 5B and Table

S4, containing acetate, propionate, butyric, valeric acids) were added in side-stream reactor.

Indeed, the sCOD concentrations into SBRP highly covariates with the nitrite accumulation in B-

stage (Figure 5A). The sCOD mass in S2EBPR in this Period is 288 mg/L. When the carbon

mass in S2EBRP decreased at the end of the study period, the nitrite accumulation also declined.

This trend cannot be observed in B-stage reactor, indicating that side stream EBPR reactor is the

place where NOB was inhibited. Spearman correlation analysis indicated that the VFA

concentration into S2EBPR abundance is negatively correlated with the abundance of *Nitrospira*

but positively correlated with the abundance of midas_g_887, which belongs to family

Comamonadaceae (Figure 5E). This influence is more profound when the VFA mass flow into

S2EBPR is high (> 90 g/d). The unique conditions in the S2EBPR, including the low ORP

environment and the VFAs concentration and composition, may play roles for the nitrite

accumulation and they are further discussed in the following sections.

3.4.1   VFAs Inhibit NOB activity and alter NOB metabolic pathway:

In 1994, Eilersen et al. reported that nitrite oxidation is more easily to be inhibited by VFA than

ammonia oxidation, especially acetic, propionic, and butyric acids (Table S5). Especially, the

butyric acid exerted strong inhibition on NOB activity (IC_{50} = 399 mg/L) comparing to acetic

(IC_{50} = 3707 mg/L) and propionic (IC_{50} = 3082 mg/L). Following research confirmed that NOB

activity was more susceptible to VFA inhibition comparing to AOB and the inhibition level is

proportional to the VFA molecular weight (Delgado et al., 2004; Gomez et al., 2000). All those

researchers came to this conclusion since the nitrate production decreased when they added

VFAs into NOB consortia. However, a later study showed that VFA (1500 mg/L) addition did

not influence the CO_2 fixation that is proportional to nitrite oxidation in NOB consortia

(*Nitrospira* spp.), indicating other metabolic pathway was activated by VFAs rather than activity

inhibition (Oguz et al., 2006). Though some *Nitrobacter* and *Nitrospira* benefit from the

presence of simple organic compounds, e.g., pyruvate, formate, urea (Daims et al., 2016; Palomo

et al., 2018), only comammox *Nitrospira* was reported to utilize fatty acids (acetate and

propionate) (Xu et al., 2022), and even have potential to synthesize and degrade poly(3-

hydroxybutyrate) (Palomo et al., 2018; Yang et al., 2020).

Comparing to the conventional WWTP that rely on mono external carbon source, such as

methanol or acetate, our system's A-stage fermentate contained a higher and stable proportion

(58.9%) of high molecular size VFAs, including propionic, butyric, iso-butyric and isovaleric

acids (Figure 5B). A-stage fermentate addition in S2EBPR may alter comammox *Nitrospira*

towards organic matter utilization pathway and down regulate the nitrite oxidation enzymes.

Notably, nitrite accumulation was not observed until the VFA mass flow was increased higher

than 66 g/d (Figure 5C). The activity tests also indicated that higher residual VFA concentration

(> 30 mg/L) in B-stage resulted lower NOB activity (in terms of nitrate production rates) ($p <$

0.01, Figure 5D), supporting the proposed role of residual VFAs in alternating the NOB activity

(Figure 5E). All those results indicated that implementation of S2EBPR with VFAs supplement

seem to favor the proliferation of versatile *Nitrospira* over canonical *Nitrospira*. This speculation

was supported by quick increase of the relative abundance of comammox *Nitrospira* over the

total *Nitrospira* community (indicated as *amoB*/*nxrB*) (Figure 4), although underlying

mechanisms need further investigation.

3.4.2 Longer exposure to anaerobic conditions with lower ORP impacts NOB:

In our system, S2EBPR induced an extended anaerobic phase (~3.7 hours) where AOB and NOB

have to survive/maintain their activities for a longer time than typical duration in conventional

WWTPs. Some research reported that the endogenous decay rate of AOB ($0.020\ \mathrm{d}^{-1}$) was slower

than that of NOB ($0.035\ \mathrm{d}^{-1}$) in anoxic condition (Cui et al., 2019), this decay difference was

also supported by the FISH staining analysis Morgenroth et al., (2000). However, other results

showed identical decay rates of AOB and NOB under both anoxic and anaerobic conditions

(Munz et al., 2011; Salem et al., 2006), The differences may have been due to wastewater

characteristics, operation parameters, and the composition of the microbial communities and the

related survival strategies (Geets et al., 2006; Hao et al., 2010). The ORP in S2EBPR was $-383 \pm$

17 mV, which is significantly below the +100 to +350 mV in normal nitrification tank, and +50

to -50 mV in denitrification tank (Environmental, 2008). Generally, the endogenous decay rates

of nitrifiers decreases with the decreasing ORP condition, as $b_{\text{aerobic}} \gg b_{\text{anoxic}} > b_{\text{anaerobic}}$ (Geets et

al., 2006; Munz et al., 2011; Ruiz-Martínez et al., 2018; Salem et al., 2006). We assume that the

actual decay rates in our system would be even lower than $0.02\ \mathrm{d}^{-1}$. Especially, the abundance of

Nitrospira was much higher than AOB throughout the experiment periods, indicating that the

decay differences between AOB/NOB did not play a significant role in NOB out-selection.

Other than low ORP, NOB is prone to be sensitive to the aerobic/anoxic perturbation. The nitrite

oxidation enzyme reactivation time of NOB is longer than AOB when they are exposed to

aerobic condition again (Gilbert et al., 2014; Kornaros et al., 2010). This characteristic has been

applied as intermittent aeration strategy in PN tank to suppress the NOB activity (Kornaros et al.,

2010). With more RAS split into S2EBPR (Figure S2), more AOB and NOB population would

undergo anaerobic and VFA exposure. With less versability, AOB may slow down their

metabolic activity while NOB may seek for other alternative pathway to maintain their energy.

This explains why they need a longer (enzyme) reactivation time (lag phase) to change back to

autotrophic nitrite oxidation in intermittent aeration systems (Gilbert et al., 2014) and thus lead

to nitrite accumulation.

4   Conclusions

Nitrite accumulation was observed along with phosphorus removal in a pilot-scale A/B-shortcut

N-S2EBPR process treating municipal wastewater.

(1) Nitrite accumulation was achieved by PN in A/B-shortcut N-S2EBPR system without

434   canonical NOB out-selection, which was the higher ammonia oxidation activity than the

435   nitrite oxidation rate.

436   (2) *Nitrospira* dominated and was the center hub of the autotrophic nitrification community

437   throughout the operation periods with one to two orders of magnitude higher than AOB,

438   as indicated by 16s rDNA phylogenetic analysis and network analysis.

439  (3) The contribution of comammox *Nitrospira* oligotype II (Nitrospira_midas_s_31566) in

440    PN was suggested by the increased *amoB*/*nxr* ratio, the significant positive correlation

441    between the abundance of *Nitrospira* Oligotype II and the ammonia oxidation activity.

442  (4) The integration of S2EBPR into the A/B-shortcut N-S2EBPR seemed to exert distinct

443    selective pressure on nitrifying community from those in conventional BNR systems.

444  (5) Phylogenetic investigation suggested that ammonia oxidation enzymes (AMO and HAO)

445    were embedded in heterotrophic organisms that belong to the family Comamonadaceae

446    and order Rhodocyclaceae. These organisms have potential to function as heterotrophic

447    nitrifiers in response to elevated VFA addition.

5 Acknowledgement

449 This study was funded by the Water Research Foundation (WRF, Grant No. 4901). We

450 gratefully acknowledge the financial support and scientific input from the WRF project team,

451 particularly from Jim McQuarrie, Issac Avila, Dan Freedman, Rudy Maltos, Blair Wisdom

452 (Denver Metro Wastewater Reclamation District), Beverley M. Stinson (AECOM), Christine

453 deBarbadillo, Haydee De Clippeleir (District of Columbia Water and Sewer Authority), Paul

454 Dombrowski (Woodard & Curran), Daniel R. Dair and Chandler H. Johnson (World Water

455 Works) and James Barnard (Black & Veatch). Special thanks are given to the generous support

456 of operators and staff of (Hampton Roads Sanitation District) for their assistance with sample

457 and data collection.

6   Reference

An, Z., Kent, T.R., Sun, Y., Bott, C.B., Wang, Z.-W., 2021. Free ammonia resistance of nitrite-oxidizing bacteria developed in aerobic granular sludge cultivated in continuous upflow airlift reactors performing partial nitritation. Water Environment Research 93, 421–432.

Bao, P., Li, G.-X., 2017. Sulfur-Driven Iron Reduction Coupled to Anaerobic Ammonium Oxidation. Environ. Sci. Technol. 51, 6691–6698.

Barbera, P., Kozlov, A.M., Czech, L., Morel, B., Darriba, D., Flouri, T., Stamatakis, A., 2019. EPA-ng: Massively Parallel Evolutionary Placement of Genetic Sequences. Systematic Biology 68, 365–369.

Barnard, J.L., Dunlap, P., Steichen, M., 2017. Rethinking the Mechanisms of Biological Phosphorus Removal. Water Environment Research 89, 2043–2054.

Beach, N.K., Noguera, D.R., 2019. Design and Assessment of Species-Level qPCR Primers Targeting Comammox. Frontiers in Microbiology 10.

Boehnke, B., Diering, B., Zuckut, S.W., 1997. Cost-effective wastewater treatment process for removal of organics and nutrients. Water Engineering and Management 144, 18–21.

Cao, Y., van Loosdrecht, M.C.M., Daigger, G.T., 2017. Mainstream partial nitration–anammox in municipal wastewater treatment: status, bottlenecks, and further studies. Appl Microbiol Biotechnol 101, 1365–1383.

Carvalheira, M., Oehmen, A., Carvalho, G., Eusébio, M., Reis, M.A.M., 2014. The impact of aeration on the competition between polyphosphate accumulating organisms and glycogen accumulating organisms. Water Research 66, 296–307.

Cotto, I., Dai, Z., Huo, L., Anderson, C.L., Vilardi, K.J., Ijaz, U., Khunjar, W., Wilson, C., De Clippeleir, H., Gilmore, K., Bailey, E., Pinto, A.J., 2020. Long solids retention times and attached growth phase favor prevalence of comammox bacteria in nitrogen removal systems. Water Research 169, 115268.

Cui, H., Zhang, L., Zhang, Q., Li, X., Peng, Y., 2019. Stable partial nitrification of domestic sewage achieved through activated sludge on exposure to nitrite. Bioresource Technology 278, 435–439.

Czech, L., Barbera, P., Stamatakis, A., 2020. Genesis and Gappa: processing, analyzing and visualizing phylogenetic (placement) data. Bioinformatics 36, 3263–3265.

Daims, H., Lebedeva, E.V., Pjevac, P., Han, P., Herbold, C., Albertsen, M., Jehmlich, N., Palatinszky, M., Vierheilig, J., Bulaev, A., Kirkegaard, R.H., von Bergen, M., Rattei, T., Bendinger, B., Nielsen, P.H., Wagner, M., 2015. Complete nitrification by Nitrospira bacteria. Nature 528, 504–509.

Daims, H., Lücker, S., Wagner, M., 2016. A New Perspective on Microbes Formerly Known as Nitrite-Oxidizing Bacteria. Trends Microbiol 24, 699–712.

Delgado, A., Aspé, E., Martí, M.C., Roeckel, M., 2004. The Effect of Volatile Fatty Acids on the Nitrification of a Saline Effluent. Environmental Technology 25, 413–422.

Douglas, G.M., Maffei, V.J., Zaneveld, J.R., Yurgel, S.N., Brown, J.R., Taylor, C.M., Huttenhower, C., Langille, M.G.I., 2020. PICRUSt2 for prediction of metagenome functions. Nat Biotechnol 38, 685–688.

Duan, S., Zhang, Y., Zheng, S., 2022. Heterotrophic nitrifying bacteria in wastewater biological nitrogen removal systems: A review. Critical Reviews in Environmental Science and Technology 52, 2302–2338.

Dueholm, M.K.D., Nierychlo, M., Andersen, K.S., Rudkjøbing, V., Knutsson, S., Albertsen, M.,
Nielsen, P.H., 2022. MiDAS 4: A global catalogue of full-length 16S rRNA gene
sequences and taxonomy for studies of bacterial communities in wastewater treatment
plants. Nature communications 13, 1–15.
Eilersen, A.M., Henze, M., Kløft, L., 1994. Effect of Volatile Fatty Acids and Trimethylamine
on Nitrification in Activated Sludge. Water Research 28, 1329–1336.
Environmental, Y.S.I., 2008. ORP Management in wastewater as an indicator of process
efficiency. YSI, Yellow Springs, OH http://www. ysi. com/media/pdfs/A567-ORP-
Management-in-Wastewater-as-an-Indicator-of-Process-Efficiency. pdf (accessed on
15.08. 13).
Eren, A.M., Maignien, L., Sul, W.J., Murphy, L.G., Grim, S.L., Morrison, H.G., Sogin, M.L.,
2013. Oligotyping: differentiating between closely related microbial taxa using 16S
ʀRNA gene data. Methods Ecol Evol 4, 1111–1119.
Fitzgerald, C.M., Camejo, P., Oshlag, J.Z., Noguera, D.R., 2015. Ammonia-oxidizing microbial
communities in reactors with efficient nitrification at low-dissolved oxygen. Water
Research 70, 38–51.
Gao, R., Peng, Y., Li, J., Zhang, Q., Li, X., Deng, L., Li, W., Kao, C., 2021. Improving
performance and efficiency of partial anammox by coupling partial nitrification and
partial denitrification (PN/A-PD/A) to treat municipal sewage in a step-feed reactor.
Bioresource Technology 341, 125804.
Ge, H., Batstone, D.J., Keller, J., 2015. Biological phosphorus removal from abattoir wastewater
at very short sludge ages mediated by novel PAO clade Comamonadaceae. Water
Research 69, 173–182.
Geets, J., Boon, N., Verstraete, W., 2006. Strategies of aerobic ammonia-oxidizing bacteria for
coping with nutrient and oxygen fluctuations. FEMS Microbiology Ecology 58, 1–13.
Gilbert, E.M., Agrawal, S., Brunner, F., Schwartz, T., Horn, H., Lackner, S., 2014. Response of
Different Nitrospira Species To Anoxic Periods Depends on Operational DO. Environ.
Sci. Technol. 48, 2934–2941.
Gomez, J., Mendez, R., Lema, J.M., 2000. Kinetic study of addition of volatile organic
compounds to a nitrifying sludge. Appl Biochem Biotechnol 87, 189–202.
Gu, A.Z., Saunders, A., Neethling, J.B., Stensel, H.D., Blackall, L.L., 2008. Functionally
Relevant Microorganisms to Enhanced Biological Phosphorus Removal Performance at
Full-Scale Wastewater Treatment Plants in the United States. Water Environment
Research 80, 688–698.
Gu, A.Z., Tooker, N.B., Onnis-Hayden, A., Wang, D., Li, G., Srinivasan, V., Takács, I., 2018.
Investigation of the Mechanisms for Optimization and Design of a Side-Stream EBPR
Process as a Sustainable Approach for Achieving Stable and Efficient Phosphorus
Removal. by Water Environmental Research Foundation (WERF) and International
Water Association (IWA) Publishing.
Hao, X.-D., Wang, Q.-L., Zhu, J.-Y., Van Loosdrecht, M.C.M., 2010. Microbiological
Endogenous Processes in Biological Wastewater Treatment Systems. Critical Reviews in
Environmental Science and Technology 40, 239–265.
He, S., Gall, D.L., McMahon, K.D., 2007. "Candidatus Accumulibacter" Population Structure in
Enhanced Biological Phosphorus Removal Sludges as Revealed by Polyphosphate
Kinase Genes. Applied and Environmental Microbiology 73, 5865–5874.

Hellinga, C., Schellen, A., Mulder, J.W., van Loosdrecht, M.C.M. v, Heijnen, J.J., 1998. The
SHARON process: an innovative method for nitrogen removal from ammonium-rich
waste water. Water science and technology 37, 135–142.
Jetten, M.S.M., Horn, S.J., van Loosdrecht, M.C.M., 1997. Towards a more sustainable
municipal wastewater treatment system. Water Science and Technology, Sustainable
Sanitation 35, 171–180.
Jiang, C., Xu, S., Wang, R., Feng, S., Zhou, S., Wu, Shimin, Zeng, X., Wu, Shanghua, Bai, Z.,
Zhuang, G., Zhuang, X., 2019. Achieving efficient nitrogen removal from real sewage
via nitrite pathway in a continuous nitrogen removal process by combining free nitrous
acid sludge treatment and DO control. Water Research 161, 590–600.
Joss, A., Salzgeber, D., Eugster, J., König, R., Rottermann, K., Burger, S., Fabijan, P., Leumann,
S., Mohn, J., Siegrist, H.R., 2009. Full-scale nitrogen removal from digester liquid with
partial nitration and anammox in one SBR. Environmental Science and Technology 43,
5301–5306.
Kits, K.D., Sedlacek, C.J., Lebedeva, E.V., Han, P., Bulaev, A., Pjevac, P., Daebeler, A.,
Romano, S., Albertsen, M., Stein, L.Y., Daims, H., Wagner, M., 2017. Kinetic analysis of
a complete nitrifier reveals an oligotrophic lifestyle. Nature 549, 269–272.
Koch, H., Lücker, S., Albertsen, M., Kitzinger, K., Herbold, C., Spieck, E., Nielsen, P.H.,
Wagner, M., Daims, H., 2015. Expanded metabolic versatility of ubiquitous nitrite-
oxidizing bacteria from the genus Nitrospira. PNAS 112, 11371–11376.
Kornaros, M., Dokianakis, S.N., Lyberatos, G., 2010. Partial Nitrification/Denitrification Can Be
Attributed to the Slow Response of Nitrite Oxidizing Bacteria to Periodic Anoxic
Disturbances. Environ. Sci. Technol. 44, 7245–7253.
Lackner, S., Gilbert, E.M., Vlaeminck, S.E., Joss, A., Horn, H., van Loosdrecht, M.C.M., 2014.
Full-scale partial nitration/anammox experiences – An application survey. Water
Research 55, 292–303.
Liu, G., Wang, J., 2013. Long-Term Low DO Enriches and Shifts Nitrifier Community in
Activated Sludge. Environ. Sci. Technol. 47, 5109–5117.
Lotti, T., Kleerebezem, R., Hu, Z., Kartal, B., De Kreuk, M.K., Van Erp Taalman Kip, C., Kruit,
J., Hendrickx, T.L.G., Van Loosdrecht, M.C.M., 2015. Pilot-scale evaluation of
anammox-based mainstream nitrogen removal from municipal wastewater.
Environmental Technology (United Kingdom) 36, 1167–1177.
Louca, S., Doebeli, M., 2018. Efficient comparative phylogenetics on large trees. Bioinformatics
34, 1053–1055.
McCullough, K., Klaus, S., Wilson, C., Vanrolleghem, P.A., Gu, A.Z., Bott, C.B., 2023.
Sidestream bio-P and mainstream anammox in a BNR process with upstream carbon
capture.
Mehrani, M.-J., Sobotka, D., Kowal, P., Guo, J., Mąkinia, J., 2022. New insights into modeling
two-step nitrification in activated sludge systems – The effects of initial biomass
concentrations, comammox and heterotrophic activities. Science of The Total
Environment 848, 157628.
Moir, J.W.B., Crossman, L.C., Spiro, S., Richardson, D.J., 1996. The purification of ammonia
monooxygenase from Paracoccus denitrficans. FEBS Letters 387, 71–74.
Morgenroth, E., Obermayer, A., Arnold, E., Brühl, A., Wagner, M., Wilderer, P.A., 2000. Effect
of long-term idle periods on the performance of sequencing batch reactors. Water Science
and Technology 41, 105–113.

Munz, G., Lubello, C., Oleszkiewicz, J.A., 2011. Factors affecting the growth rates of
ammonium and nitrite oxidizing bacteria. Chemosphere 83, 720–725.
Norton, J.M., Alzerreca, J.J., Suwa, Y., Klotz, M.G., 2002. Diversity of ammonia
monooxygenase operon in autotrophic ammonia-oxidizing bacteria. Arch Microbiol 177,
139–149.
Oguz, M.T., Robinson, K.G., Layton, A.C., Sayler, G.S., 2006. Volatile fatty acid impacts on
nitrite oxidation and carbon dioxide fixation in activated sludge. Water Research 40,
665–674.
Onnis-Hayden, A., Srinivasan, V., Tooker, N.B., Li, G., Wang, D., Barnard, J.L., Bott, C.,
Dombrowski, P., Schauer, P., Menniti, A., Shaw, A., Stinson, B., Stevens, G., Dunlap, P.,
Takács, I., McQuarrie, J., Phillips, H., Lambrecht, A., Analla, H., Russell, A., Gu, A.Z.,
2020. Survey of full-scale sidestream enhanced biological phosphorus removal
(S2EBPR) systems and comparison with conventional EBPRs in North America: Process
stability, kinetics, and microbial populations. Water Environment Research 92, 403–417.
Palomo, A., Pedersen, A.G., Fowler, S.J., Dechesne, A., Sicheritz-Pontén, T., Smets, B.F., 2018.
Comparative genomics sheds light on niche differentiation and the evolutionary history
of comammox Nitrospira. ISME J 12, 1779–1793.
Parada, A.E., Needham, D.M., Fuhrman, J.A., 2016. Every base matters: assessing small subunit
rRNA primers for marine microbiomes with mock communities, time series and global
field samples. Environmental microbiology 18, 1403–1414.
Park, S., Bae, W., Rittmann, B.E., 2010. Operational Boundaries for Nitrite Accumulation in
Nitrification Based on Minimum/Maximum Substrate Concentrations That Include
Effects of Oxygen Limitation, pH, and Free Ammonia and Free Nitrous Acid Inhibition.
Environ. Sci. Technol. 44, 335–342.
Pester, M., Maixner, F., Berry, D., Rattei, T., Koch, H., Lücker, S., Nowka, B., Richter, A.,
Spieck, E., Lebedeva, E., Loy, A., Wagner, M., Daims, H., 2014. NxrB encoding the beta
subunit of nitrite oxidoreductase as functional and phylogenetic marker for nitrite-
oxidizing Nitrospira. Environmental Microbiology 16, 3055–3071.
Quince, C., Lanzen, A., Davenport, R.J., Turnbaugh, P.J., 2011. Removing noise from
pyrosequenced amplicons. BMC bioinformatics 12, 38.
Regmi, P., Miller, M.W., Holgate, B., Bunce, R., Park, H., Chandran, K., Wett, B., Murthy, S.,
Bott, C.B., 2014. Control of aeration, aerobic SRT and COD input for mainstream
nitritation/denitritation. Water Research 57, 162–171.
Ren, Z. (Jason), Pagilla, K. (Eds.), 2022. Pathways to Water Sector Decarbonization, Carbon
Capture and Utilization. IWA Publishing.
Rittmann, B.E., Mayer, B., Westerhoff, P., Edwards, M., 2011. Capturing the lost phosphorus.
Chemosphere 84, 846–853.
Roots, P., Wang, Y., Rosenthal, A.F., Griffin, J.S., Sabba, F., Petrovich, M., Yang, F., Kozak,
J.A., Zhang, H., Wells, G.F., 2019. Comammox Nitrospira are the dominant ammonia
oxidizers in a mainstream low dissolved oxygen nitrification reactor. Water Research
157, 396–405.
Rotthauwe, J.H., Witzel, K.P., Liesack, W., 1997. The ammonia monooxygenase structural gene
amoA as a functional marker: molecular fine-scale analysis of natural ammonia-oxidizing
populations. Applied and Environmental Microbiology 63, 4704–4712.

Ruiz-Martínez, A., Claros, J., Serralta, J., Bouzas, A., Ferrer, J., 2018. Modeling the decay of nitrite oxidizing bacteria under different reduction potential conditions. Process Biochemistry 71, 159–165.

Salem, S., Moussa, M. s., van Loosdrecht, M. c. m., 2006. Determination of the decay rate of nitrifying bacteria. Biotechnology and Bioengineering 94, 252–262.

Sancho, I., Lopez-Palau, S., Arespacochaga, N., Cortina, J.L., 2019. New concepts on carbon redirection in wastewater treatment plants: A review. Science of The Total Environment 647, 1373–1384.

Shao, Y.-H., Wu, J.-H., 2021. Comammox Nitrospira Species Dominate in an Efficient Partial Nitrification–Anammox Bioreactor for Treating Ammonium at Low Loadings. Environ. Sci. Technol. 55, 2087–2098.

Srinivasan, V.N., Li, G., Wang, D., Tooker, N.B., Dai, Z., Onnis-Hayden, A., Bott, C., Dombrowski, P., Schauer, P., Pinto, A., Gu, A.Z., 2021. Oligotyping and metagenomics reveal distinct Candidatus Accumulibacter communities in side-stream versus conventional full-scale enhanced biological phosphorus removal (EBPR) systems. Water Research 206, 117725.

Ssekagiri, A., Sloan, W., Zeeshan Ijaz, U., 2017. microbiomeSeq: An R package for analysis of microbial communities in an environmental context. ISCB Africa ASBCB Conference.

Torondel, B., Ensink, J.H.J., Gundogdu, O., Ijaz, U.Z., Parkhill, J., Abdelahi, F., Nguyen, V.-A., Sudgen, S., Gibson, W., Walker, A.W., Quince, C., 2016. Assessment of the influence of intrinsic environmental and geographical factors on the bacterial ecology of pit latrines. Microbial Biotechnology 9, 209–223.

Van Dongen, U., Jetten, M.S.M., Loosdrecht, M.C.M. Van, 2001. Ammonium Rich Wastewater. Growth Lakeland 44, 153–160.

van Kessel, M.A.H.J., Speth, D.R., Albertsen, M., Nielsen, P.H., Op den Camp, H.J.M., Kartal, B., Jetten, M.S.M., Lücker, S., 2015. Complete nitrification by a single microorganism. Nature 528, 555–559.

Villanueva, R.A.M., Chen, Z.J., 2019. ggplot2: Elegant Graphics for Data Analysis (2nd ed.). Measurement: Interdisciplinary Research and Perspectives 17, 160–167.

Wang, D., Tooker, N.B., Srinivasan, V., Li, G., Fernandez, L.A., Schauer, P., Menniti, A., Maher, C., Bott, C.B., Dombrowski, P., Barnard, J.L., Onnis-Hayden, A., Gu, A.Z., 2019. Side-stream enhanced biological phosphorus removal (S2EBPR) process improves system performance - A full-scale comparative study. Water Research 167, 115109.

Wang, Q., Duan, H., Wei, W., Ni, B.-J., Laloo, A., Yuan, Z., 2017. Achieving Stable Mainstream Nitrogen Removal via the Nitrite Pathway by Sludge Treatment Using Free Ammonia. Environ. Sci. Technol. 51, 9800–9807.

Wang, Q., Ye, L., Jiang, G., Hu, S., Yuan, Z., 2014. Side-stream sludge treatment using free nitrous acid selectively eliminates nitrite oxidizing bacteria and achieves the nitrite pathway. Water Research 55, 245–255.

Wang, Shanyun, Peng, Y., Ma, B., Wang, Shuying, Zhu, G., 2015. Anaerobic ammonium oxidation in traditional municipal wastewater treatment plants with low-strength ammonium loading: Widespread but overlooked. Water Research 84, 66–75.

Wehrfritz, J.-M., Carter, J.P., Spiro, S., Richardson, D.J., 1996. Hydroxylamine oxidation in heterotrophic nitrate-reducing soil bacteria and purification of a hydroxylamine-cytochromec oxidoreductase from aPseudomonas species. Arch. Microbiol. 166, 421–424.

Winkler, M.K.H., Bassin, J.P., Kleerebezem, R., Sorokin, D.Y., van Loosdrecht, M.C.M., 2012.
Unravelling the reasons for disproportion in the ratio of AOB and NOB in aerobic
granular sludge. Appl Microbiol Biotechnol 94, 1657–1666.
Xu, G., Xu, X., Yang, F., Liu, S., Gao, Y., 2012. Partial nitrification adjusted by hydroxylamine
in aerobic granules under high DO and ambient temperature and subsequent Anammox
for low C/N wastewater treatment. Chemical Engineering Journal 213, 338–345.
Xu, S., Chai, W., Xiao, R., Smets, B.F., Palomo, A., Lu, H., 2022. Survival strategy of
comammox bacteria in a wastewater nutrient removal system with sludge fermentation
liquid as additional carbon source. Science of The Total Environment 802, 149862.
Yang, Y., Daims, H., Liu, Y., Herbold, C.W., Pjevac, P., Lin, J.-G., Li, M., Gu, J.-D., 2020.
Activity and Metabolic Versatility of Complete Ammonia Oxidizers in Full-Scale
Wastewater Treatment Systems. mBio 11, e03175-19.
Zhang, T., Wang, B., Li, X., Zhang, Q., Wu, L., He, Y., Peng, Y., 2018. Achieving partial
nitrification in a continuous post-denitrification reactor treating low C/N sewage.
Chemical Engineering Journal 335, 330–337.
Zhou, Y., Oehmen, A., Lim, M., Vadivelu, V., Ng, W.J., 2011. The role of nitrite and free
nitrous acid (FNA) in wastewater treatment plants. Water Research 45, 4672–4682.
700

Figure 1 AOB and NOB activities, nitrite accumulation, and the relative abundance of AOB and

NOB throughout the operation period for this study.

Figure 2 (A) Nitrogen mass balance of the pilot plant in study period (units are in g/d; values with the percentage sigh represent the removal rate, values of TIN, ammonia, nitrite, and nitrate are colored with black, green, red, and blue, respectively The B-stage effluent will go into the following anammox reactor; (B) Intermittent aeration profile taken on Day 252, showing nitrite accumulating during aerobic periods and being denitrified during anoxic periods. Reproduced from McCullough et al. (2023).

Figure 3 (A) Relative abundance of microbial community during nitrite accumulation period at genus level by 16S rRNA sequencing. (B) Quantitative nitrifying genes, *amoA* indicating ordinary proteobacteria AOB, *amoB* indicating comammox *Nitrospira*, and *nxr* indicating NOB *Nitrospira*, normalized by 16S rRNA genes. (C) Network analysis on OTU level with Spearman associations; The size of the nodes is proportional to its own total degree. The width of the edges is proportional to the correlation between the two nodes to which it corresponds. Positive and

negative correlations between taxa (nodes) are indicated by blue and red color of the edges

respectively. Detailed network analysis figure can be found in Figure S5.

Figure 4 (A) Oligotypes (3 total) of 16S rRNA gene amplicon sequences classified as genus *Nitrospira* throughout the investigation period; (B) Positive correlation between the relative abundance of oligotype II (CCGGCTTAGTGGGTATTA) and the nitrite concentration; (C) Blast results of taxonomy identification of oligotypes I and II against MiDAS4 database.

Figure 5 (A) Soluble COD concentrations in SBPR, B-stage and nitrite accumulation; (B) VFA

speciation in A-stage fermentate; (C) VFA mass flow into SBPR and nitrite accumulation; (D)

VFA concentration in SBPR and AOB/NOB activities; (E) The Spearman correlation between the most abundant 50 taxa and numerical environmental variables. The Spearman correlation test is performed and associated p-values are adjusted for multiple testing.