Comammox and Unknown Candidate AOBs Contribute to Nitrite Accumulation in An

Integrated A-B stage process that Incorporates Side-stream EBPR (S2EBPR)

Yuan Yan[a,#], Jangho Lee[a,#], IL Han[a], Zijian Wang[a], Kester McCullough[a,b], Stephanie Klaus[b], Da

Kang[a,c], DongQi Wang[a], Anand Patel[a], Guangyu Li[a], Jim McQuarrie[d], Beverley M. Stinson[e],

Christine deBarbadillo[f], Paul Dombrowski[g], Charles Bott[b, *], April Gu[a,*]

[a] School of Civil and Environmental Engineering, Cornell University, Ithaca, United States

[b] Hampton Roads Sanitation District, Virginia Beach, United States

[c] Department of Environmental Engineering, College of Environmental & Resource Sciences,

Zhejiang University, China

[d] Denver Metro Wastewater Reclamation District, Denver, Colorado, United States

[e] AECOM, Laurel, Maryland

[f] DC Water and Sewer Authority, Washington, District of Columbia

[g] Woodard & Curran, Inc, Enfield, Connecticut, United States

[#] Joint first authors

[*] Corresponding authors

Graphical abstract

Highlights

• Nitrite accumulation was achieved in HRAS-P(d)N-S2EBPR system via partial nitrification

with *Nitrospira*/AOB >10.

• The positive correlation between *Nitrospira* oligo type 2 and nitrite accumulation suggested the

contribution of comammox *Nitrospira* in partial nitrification.

• The integration of S2EBPR and elevating fermentate VFA encourages the development of

heterotrophic nitrifiers from family Comamonadaceae and order Rhodocyclaceae.

Abstract

In a pilot-scale A-B process integrated with side-stream enhanced biological phosphorus removal (EBPR) for treating municipal wastewater, average of 7.62 ± 2.17 mg-N/L nitrite accumulation with nitrite accumulation ratio of 49% was achieved by partial nitrification, which was verified through relative higher ammonia oxidation activity (1.94 ± 0.89 mgN/gSS/h) than nitrite oxidation rate (1.06 ± 0.99 mgN/gSS/h). *Nitrospira* dominated the ammonia oxidizing community in a B-stage nitrification tank throughout the operation periods with one to two orders of magnitude higher than canonical aerobic oxidizing bacteria (AOB). The higher ammonia oxidation activity along with increased *amoB*/*nxr* ratio in nitrite accumulation period suggested the contribution of comammox *Nitrospira* as AOB. Phylogenetic investigation suggested that AMO and HAO were embedded in heterotrophic organisms that belong to the family Comamonadacea and the order Rhodocyclaceae. These organisms may function as heterotrophic nitrification. The integration of S2EBPR provides PAOs suitable environment, such as low ORP and favorable VFA concentrations and composition, but also exerts additional selective pressure on NOB activity than AOB, which may be beneficial for main-stream partial nitrification.

1 Introduction

The adsorption/bio-oxidation (A/B) process is a promising engineering innovation towards carbon capture, energy recovery, and efficient nutrient removal. It diverts organic carbon to anaerobic digestion and thus manipulates the C/N in B-stage reactors through carbon flow redirection (Ren and Pagilla, 2022). The first stage (A-stage) is designed for high-rate carbon capture and removal using high-rate activated sludge (HRAS), and the second stage (B-stage) is for nutrient (mostly nitrogen) removal (Sancho et al., 2019). HRAS is widely operated with a low hydraulic retention time (HRT), a low sludge retention time (SRT) and a low dissolved oxygen (DO) concentration less than 0.5 mg L^{-1} (Boehnke et al., 1997). After HRAS, the wastewater COD/N ratio could be lowered to 0.67 (Lotti et al., 2015), which is suitable for B-stage short-cut nitrogen removal processes to achieve "carbon-free" autotrophic nitrogen removals.

Partial nitrification with anammox (PN/A) process has emerged to be one of the promising nitrogen removal processes, which can save 100% carbon, 60% aeration and reduce 80% sludge production (Jetten et al., 1997). It consists of two step processes where aerobic ammonia-oxidizing bacteria (AOB) oxidize half of the ammonia to nitrite and, then anaerobic ammonia-oxidizing bacteria (anammox) bacteria convert ammonia and nitrite into dinitrogen. Nitrite accumulation through partial nitrification in B-stage via the out-selection of nitrite oxidizing bacteria (NOB) is crucial for PN/A and it has been successfully demonstrated for side-stream high-strength wastewater (Lackner et al., 2014). A number of operational strategies have been suggested to enable side-stream NOB out-selection and consequently nitrite accumulation, including high temperature, low DO, low SRT, high pH, and high free nitrous acid (FNA) and/or free ammonia (FA) concentration (Hellinga et al., 1998; Joss et al., 2009; Van Dongen et al.,

2001; Xu et al., 2012). However, these strategies were not readily applicable in mainstream

treatment since the inhibitory levels of 0.02 mg/L FNA and 1 mg/L FA is unavailable with ca. 50

mg-N/L ammonia at neutral pH in the mainstream (Wang et al., 2017). Also, it has been shown

that NOB can adapt to high concentration of FA (An et al., 2021) and low DO (Liu and Wang,

2013), which makes it harder to be out-selected in mainstream process. Therefore, maintaining

long-term stable NOB out-selection still remains a challenge for main-stream wastewater

treatments and the underlined mechanisms need to be explored.

Most PN/A studies focus only on nitrogen removal, however, phosphorus co-occur with nitrogen

and is also a critical nutrient that needs to be treated in wastewater (Rittmann et al., 2011).

Enhanced biological phosphorus removal (EBPR) process has been widely applied worldwide in

full-scale facilities. Its performance and stability strongly rely on the activity of phosphorus

accumulating organisms (PAOs) with favorable influent rbCOD/TP ratio >15 (Gu et al., 2008).

Therefore, EBPR is considered incompatible with B-stage autotrophic nitrogen removal due to

the insufficiency of readily biodegradable carbon in A-stage effluent (B-stage influent). The

implementation of a side-stream anaerobic biological sludge hydrolysis and fermentation reactor

called side-stream EBPR (S2EBPR) can be an alternative option to address the carbon shortage

challenge of coupling EBPR with B-stage nitrogen removal process. S2EBPR takes advantage of

the volatile fatty acids (VFAs) generated via hydrolysis/fermentation of side-stream return

activated sludge (RAS) and mixed liquor, and thus overcome the reliance on the influent

carbon (Vollertsen et al., 2006; Barnard et al., 2017; Onnis-Hayden et al., 2020; Gu et al., 2018;

Wang et al., 2019). However, the influence of implementing S2EBPR on the PN/A removal

system, especially on the nitrite accumulation remains unknown. We hypothesize that the

incorporation of S2EBPR would affect the nitrifying populations because of the extended

anaerobic conditions in the side-steam reactor with low oxidation reduction potential (ORP,

<300 mV) and high VFAs. In addition, PAOs and the surplus of sludge fermentate from the

S2EBPR reactor to the down-stream B-stage would compete oxygen with nitrifiers (Carvalheira

et al., 2014).

Recently, our team has reported a successful incorporation of S2EBPR into a A-B stage process

with downstream Annamox polishing to achieve simultaneous short-cut N removal and

biological P removal for main-stream treatment (Klaus, 2019; McCullough et al., 2020). This

pilot-scale HRAS-P(d)N-S2EBPR-Anammox system provides us good opportunity to explore

the interacting impacts of PAOs and nitrogen removal related microorganisms. In this study, we

focused on the comprehensive examination of the nitrite accumulation and AOB and NOB

population dynamics in B-stage processes that generated nitrite with intermittent aeration AvN

(ammonia versus NO_x^--N (nitrite+nitrate) control strategy. Process performance, batch activity tests

were performed to investigate the underlying processes involved in the nitrite accumulation. 16S

rRNA gene amplification sequencing and quantitative PCR were applied to reveal the microbial

community composition and dynamics with emphasis on nitrification-relevant populations. To

our knowledge, this is the first study that revealed the influence of implementing S2EBPR on B-

stage nitrification population and, provided new insights on the distinctive nitrifying populations

that contributed to nitrite accumulation in this HRAS-P(d)N-S2EBPR process that is different

from those typically observed in previous WWTPs. The results pointed out a possible new

strategy in achieving nitritation to enable annamox for main-stream municipal WW treatment.

2   Material methods

2.1   Pilot reactor setup and Operation conditions

The pilot plant operated by Hampton Roads Sanitation District (HRSD) is located at

Chesapeake-Elizabeth WWTP in Virginia Beach, VA, USA. It mainly consisted of an A-stage

HRAS process followed by the B-stage process, a parallel S2EBPR reactor and a final stage

Annamox polishing tank (Figure S1). The detailed parameters of A-B process have been

introduced elsewhere (Klaus et al., 2020; Printz, 2019). The A-stage HRAS consisted of three

bioreactors in series (V_{total} = 511 L) followed by an intermediate clarifier with effluent going to

the B-stage process. A portion of A-stage wasted activated sludge (WAS) was splited into a

combined fermenter and thickener. The B-stage consisted of four CSTRs bioreactors in series

(V_{total} = 606 L) each with independent aeration control via Ammonia versus NOx (nitrite +

nitrate) (AvN) control (Regmi et al., 2014) and mechanical mixing. The B-stage system was

operated in a fully intermittent configuration in which all CSTRs were intermittently aerated for

the same interval and duration, and it was maintained at a 5-hour HRT using a flow-paced (2.0 L

min^{-1}) influent progressive cavity pump (Seepex BW2, Bottrop, Germany) with a variable

frequency drive (VFD) and magnetic flow meter (Rosemount 8705, Houston, TX). In the

studied period, B-stage reactors were switched from clarifier wasting to aeration tank wasting

(hydraulic wasting) and recycling. A portion of B-stage RAS and the A-stage WAS fermentate

were added to the side-stream reactor for simultaneous VFA production via hydrolysis and

fermentation, and for PAOs selection. The HRT of side-stream reactor varies depending on the

percentage of B-stage RAS sent to the reactor, but it maintained around 4 hours during most of

the operation period in this study (Figure S2). The SRT of the S2EBPR reactor was estimated as

$SRT_{total} - SRT_{B-stage,}$ resulting in an average SRT of 6 days.

2.2   Chemical analysis: sCOD, VFA, NH4, NO2, NO3, PO4

Total and soluble COD, OP, total ammonia nitrogen (NH_4^+-N + NH_3-N), NO_2^--N, and NO_3^--N

were measured with HACH TNTplus kits (Loveland, Colorado) and a HACH DR2800

spectrophotometer (Loveland, CO). Nutrient and soluble COD samples were filtered through

0.45 μm and 1.5 μm filters respectively. The sCOD measurement includes readily biodegradable

and colloidal COD fractions. Daily pH and temperature readings of the reactors were recorded

using a handheld pH and temperature meter (Beckman Coulter, Brea, CA). Readings for DO

were recorded using a handheld DO sensor (Insite Model 10, LA, USA).

Concentrations of acetic, propionic, butyric, isobutyric, valeric, isovaleric, and caproic acids

were determined by GC-FID as previously described in (Kinyua et al., 2017). VFA

concentrations were determined from a 0.5-100 mg/L calibration curve prepared from

commercial standards (Absolute Standards).

2.3   AOB and NOB activity analysis

AOB and NOB activities were analysis weekly on site. About 4L mixed liquor suspended solids

(MLSS) were withdrawn from the last B-stage aerobic tank and pre-aerated for 30 min to oxidize

the residue carbon. Ammonium chloride was spiked to reach 25 mg-N/L final concentration. DO

and pH were maintained around 3-5 mg/L and 7, respectively. The samples were taken every 15

minutes for a total of 1 hour. After filtered by 0.45 μm filter, the filtrate was used to analyze the

NH_4^+-N, NO_2^--N, and NO_3^--N concentrations using HACH TNT kits (HACH Loveland, CO).

AOB activity was calculated as the sum of nitrite and nitrate production rates, and NOB activity

was calculated as the nitrate production rates by linear regression.

2.4   Quantification of nitrifying candidates

Quantitative polymerase chain reaction (qPCR) was performed on a CFX96 real-time PCR

detection system (Bio-Rad, USA) to determine the concentrations of functional genes including

amoA, *nxrB*, *amoB* and 16S. The amoA-1F/2R primer for the canonical ammonia

monooxygenase (AMO) enzymes in AOB (Rotthauwe et al., 1997), and the *nxrB* 169F/638R

primer for the beta subunit of nitrite oxidoreductase (NXR) (Pester et al., 2014) were used to

target canonical AOB and NOB. The *amoB* 148F/485R primer were used to target the beta

subunit of AMO in *Nitrospira* that can do complete ammonia oxidation (comammox) (Cotto et

al., 2020). All functional genes were then normalized by 16S rRNA with primer set 341f/534r

(He et al., 2007). Calibration curves were obtained using 10-fold serial dilutions of known

concentrations of cloned DNA samples that were prepared following the instruction from TOPO

TA cloning kit (Thermo Fisher, USA) in our laboratory. The R^2 of standard curves were above

0.99 and the amplification efficiency was around 95%. The detailed qPCR reaction conditions

were summarized in Table S1.

2.5   Bioinformatics and statistical analysis

2.5.1   Microbial population analysis

Sludge samples in effluent from B-stage were monthly collected for DNA extraction. Genomic

DNA was extracted using Instagene matrix (Kim et al., 2013) and then used for both 16S

ribosomal RNA (16S rRNA) sequencing and functional genes quantification.

The V4 region of the 16S rRNA genes were amplified using primers 515F-Y (Parada et al.,

2016) and 926R (Quince et al., 2011). Each primer contained a unique six base index sequence

for sample multiplexing, as well as Illumina flow cell binding and sequencing sites (Bartram et

al., 2011). The 25 μl PCR mix contained 1× ThermoPol buffer, 0.2 μM forward primer, 0.2 μM

reverse primer, 200 μM dNTPs, 15 μg bovine serum albumin, 0.625 U *Taq* DNA polymerase

(New England Biolabs, USA), and 10 ng of template. The PCR was prepared in triplicate as

follows: 95°C for 3 min, 35 cycles of 95°C for 30 sec, 50°C for 30 sec, 68°C for 1 min, and a

final extension of 68°C for 7 min. Equal quantities of 16S rRNA gene amplicons were pooled,

excised from an agarose gel, and purified using the Wizard SV Gel and PCR Clean-Up System

(Promega, WI, USA). A 3 pM library containing 10% PhiX Control v3 (Illumina, CA, USA) was

sequenced on a MiSeq (Illumina, CA, USA) using a 2 x 250 cycle MiSeq Reagent Kit v2.

Paired-end reads were assembled using the paired-end assembler for Illumina sequences

(PANDAseq version 2.8) (Masella et al., 2012). The raw paired-end reads were assembled for

each sample and processed in Mothur v. 1.39.4 following the MiSeq standard operating

procedure (SOP) (Kozich et al., 2013). High-quality reads were obtained after quality control and

chimera screening and then clustered at a 97% similarity to obtain the operational taxonomic

units (OTUs).

To minimize the inflation of rare species in the community analysis, we removed OTUs that are

lower than 0.01%. The filtered-OUT matrix was used to perform statistical analyses in R-4.2.3.

The matrix was also used to predict the bacterial metagenomes using Phylogenetic Investigation

of Communities by Reconstruction of Unobserved States 2 (PICRUSt2) pipeline (Douglas et al.,

2020) with implemented tools EPA-ng (Barbera et al., 2019), gappa (Czech et al., 2020), and

castor (Louca and Doebeli, 2018). The co-occurrence patterns of microbial community and the

correlation between the environmental variables and the most abundant 50 taxa were done

through microbiomeSeq package (Ssekagiri et al., 2017; Torondel et al., 2016). All bioinformatic

figures were generated with ggplot2 package (Villanueva and Chen, 2019, p. 2).

2.5.2   Oligotyping analysis

Oligotyping analysis was performed according to (Eren et al., 2013; Srinivasan et al., 2021) to

reveal the sub-clusters (i.e. microdiveristy) within the dominant nitrifying population. A total of

65,24316S rRNA gene amplicon sequences classified as *Nitrospira* were extracted and formatted

as required by the oligotyping pipeline. After quality filtering, 18 nucleotide positions were

selected for entropy decomposition (102, 120, 134, 135, 149, 151, 160, 181, 197, 199, 210, 219,

240, 262, 265, 375, 488, 491), which were used to generate a total of 205 raw oligotypes. After

denoising and elimination based on the -s (oligotypes that occurred in at least 3 samples) and -M

(a minimum count of 30 (~0.1%)) parameters, 3 dominant oligotypes, presumably representing

micro-diversity of *Nitrospira* sub-groups, were finally identified. The representative sequence of

the 3 dominant oligotypes were retrieved and blasted in the NCBI's nt database

(http://www.ncbi.nlm.nih.gov/).

3   Results and Discussion

3.1   Simultaneous nitrite accumulation and efficient P removal in A-B-stage-S2EBPR

process

The HRAS-P(d)N-S2EBPR treatment line was designed to perform nitrogen and phosphorus

removal simultaneously. To overcome the shortage of influent VFA and enable bio-P removal in

main-stream treatment line, a side-stream reactor was implemented to facilitate bio-P removal

through receiving supplemental carbon from A-stage sludge fermenter (Figure S1). With the

increase in the A-stage WAS fermentate supplement flow and RAS slit ratio to the S2EBPR

reactor (Figure S2), the nitrite accumulation was observed over a period of approximately 150

days (Figure 1). During this period, the effluent nitrite concentration raised to a maximum level

of 6.55 mg-N/L with an average concentration of 2.65 ± 1.89 mg-N/L (Figure 1). The average

nitrite accumulating ratio (NAR) was 49%, suggesting that a significant nitrite accumulation

occurred. Figure 2A shows the N mass balance in the system and changes of the various N

species along the B-stage reactors in series. As ammonia decreased along the B-stage zones, the

nitrite began to accumulate from 3.9 g/d to 20.7 g/d (Figure 2A). Previous research indicated that

nitrite can inhibit PAO activity by increasing maintenance energy and disturbing the TCA cycle

(Zhou et al., 2011). Nevertheless, our results indicated an efficient P removal (77%-92%) could

be achieved concurrently with nitrite accumulation (0.2 ± 0.05 - 2.7 ± 1.9 mg-N/L) during this

same period (Supplementary text in SI).

Figure 1 AOB and NOB activities, nitrite accumulation, and the relative abundance of AOB and

NOB throughout the operation period for this study.

Figure 2 (A) Nitrogen mass balance of the pilot plant in study period (units are in g/d; values

with the percentage sigh represent the removal rate, values of TIN, ammonia, nitrite, and nitrate

are colored with black, green, red, and blue, respectively The B-stage effluent will go into the

following anammox reactor; (B) Intermittent aeration profile taken on Day 522, showing nitrite

accumulating during aerobic periods and being denitrified during anoxic periods. Reproduced

from McCullough et al., (2020).

3.2 Nitrite accumulation due to partial nitrification

Since nitrite can be accumulated either through partial nitrification or partial denitrification

pathways, both AOB/NOB activities were evaluated over long-term on site to identify the

dominant mechanisms involved in the nitrite accumulation (Figure 1). Examination of the N

species during the aerobic/anoxic cycle showed that nitrite only accumulated in aeration periods

while it was consumed after aeration was off (Figure 2B), suggesting that nitrite accumulation

was likely due to partial nitrification (PN) rather than partial denitrification. This was further

supported by the AOB and NOB activities assessment. During the study period, the average

AOB and NOB activities were 1.9 ± 0.9 and 1.1 ± 1.0 mgN/gSS/h, respectively (Figure 1). AOB

activities were higher than the NOB activities during the period of study, consistent with the

nitrite accumulation via partial nitrification (Figure 1). This evidence led to our conclusion that

the higher nitrite accumulation in this study period is likely attributed to, at least mostly, partial

nitrification.

3.3   Unknown AOB that contributed to the High AOB activity

3.3.1   Low and negligible known AOB, NOB abundance

Temporal Microbial communities via 16S rRNA gene sequencing analysis revealed the

microbial community structure dynamics and patterns during the study period (Figure 3). The

average relative abundance of known AOB, including *Nitrosomonas* and *Nitrosomonadaceae*

were remarkably low at 0.07% and 0.2%, respectively. The only known NOB is *Nitrospira*, has a

relative abundance ranging from 1.7-6%. Their abundances were depicted in Figure 1 to show

the variation trends. In our system, the relative abundances of both known AOB and NOB are

significantly lower than that typically reported in mainstream PN/A systems (0.23-4.4% and 0.2-

53%) (summarized in Table S2). Particularly interesting, the relative abundance of *Nitrospira*

was much higher than the known AOB, which did not align with the higher ammonia oxidation

activity compared to nitrite oxidation activity.

Figure 3 (A) Relative abundance of microbial community during nitrite accumulation period at

genus level by 16S rRNA sequencing. (B) Quantitative nitrifying genes, *amoA* indicating

ordinary proteobacteria AOB, *amoB* indicating comammox *Nitrospira*, and *nxr* indicating NOB

Nitrospira, normalized by 16S rRNA genes. (C) Network analysis on OTU level with Spearman

associations; The size of the nodes is proportional to its own total degree. The width of the edges

is proportional to the correlation between the two nodes to which it corresponds. Positive and

negative correlations between taxa (nodes) are indicated by blue and red color of the edges

respectively.

To further verify the abundances of nitrifiers in the sludge, the *amoA* genes in AOB and the *nxr*

genes in NOB were quantified and normalized to 16S rRNA genes (Figure 3B). The average

nxr/16S ratio was 8.3×10^{-3} in this study period. Generally, *Nitrospira* genome harbor 1-6 copies

of *nxr* (Palomo et al., 2018; Pester et al., 2014), while AOB in betaproteobacteria contains 2-3

copies of *amoA* (Norton et al., 2002). Considering the copy difference, the abundance of *nxrB*

was still one order of magnitude higher than *amoA,* suggesting a higher known *NOB* abundance

than known AOB abundance (Figure 3B). Both 16S sequencing and functional gene

quantification confirmed that the abundance of known AOB (*Nitrosomonas* and

Nitrosomonadaceae) was lower than NOB (*Nitrospira*) throughout the treatment periods, yet the

system still achieved partial nitrification.

Theoretically, the numerical ratio of AOB to NOB in a balanced nitrifying system should be

higher than 2:1 (Arciero et al., 1991; Winkler et al., 2012). However, the current investigation

shows a disparity between the rates of nitrite and ammonia oxidation, and the abundance of

nitrifying biomass. Similar observations have been reported before (Wang et al., 2015; Fitzgerald

et al., 2015; Zhang et al., 2018; Gao et al., 2021). Wang et al. (2015) observed NOB:AOB ratios

from 9:1 to 5000:1 in five activated sludge reactors through qPCR. Zhang et al. (2018) achieved

stable nitrite accumulation with higher abundance of *Nitrospira* comparing to AOB. Several

scenarios can potentially explain the higher relative abundance of known NOBs but lower nitrite

oxidation activity in our system: 1) there might be a group of *Nitrospira* who can perform not

only nitrite oxidation, but also ammonia oxidation, such as complete ammonia oxidizer

(comammox); 2) underestimation of AOB in the system due to unknown AOBs; and 3) some

substances in wastewater could have more severe inhibitory effects on NOB than AOB.

3.3.2   Comammox *Nitrospira* may contribute to nitrite accumulation.

Roots et al., (2019) reported that the abundance of Comammox *Nitrospira* was much higher than

both AOB and ammonia oxidizing archaea (AOA) in their in nitrification reactor (Roots et al.,

2019), challenging the commonly held assumption that comammox play a minor role in

wastewater treatment bioreactors. Among the versatile *Nitrospira* genus, the comammox

Nitrospira encode a phylogenetically distinct *amo* gene from the canonical AOB and AOA,

allowing them to fully oxidize ammonia to nitrate (Daims et al., 2016; Koch et al., 2015). High

presence of comammox was reported in nitrite accumulating condition, indicating a potential

correlation between the comammox and nitrite accumulation, given the higher ammonia

oxidation rate than the nitrite oxidation rate in comammox *Nitrospira* (Shao and Wu, 2021). A

recent isotope labelling test has proven that comammox species (*N. nitrosa* and *N. nitrificans*)

provide nitrite for anammox bacteria (*Brocadia*) (van Kessel et al., 2015). The pure cultured *N.*

inopinata was recorded to accumulate nitrite during complete nitrification (Kits et al., 2017).

Given the function versatility of comammox species and the high abundance of *Nitrospira*, it's

possible that a portion of the comammox *Nitrospira* in our system functioned as AOB, leading to

the accumulation of nitrite.

The presence of comammox *Nitrospira* in our system was determined by quantifying the *amoB*

gene that was specifically found in comammox (Cotto et al., 2020; Roots et al., 2019) (Figure

3B). Results indicated an occurrence of higher abundance of comammox *Nitrospira* and nitrite

accumulation. In addition, *amoB* was present and remained much more stable than both *nxrB* and

amoA (Figure 3B). The average of *amoB* ratio was 7.8×10^{-4} per 16S gene copies. The *amoB*/*nxrB*

ratio was deployed to indicate the relative abundance of comammox *Nitrospira* to the total

Nitrospira. Results clearly indicate that comammox *Nitrospira* abundance increased from 1% to

13.6% during nitrite accumulation period. Considering that *Nitrospira* genome contains 1-5

times more copies of *nxr* (1-2 copies in most comammox genomes but 4 copies in *N. nitrificans*

and 5 copis in *N. mosocoviensis*) than *amo* (single copy) (Palomo et al., 2018; Pester et al.,

2014), the relative abundance of comammox *Nitrospira* to the total NOB ranges from 13.6% to

68% in our system.

The increasing abundance of comammox *Nitrospira* motivated us to employ the oligotyping

analysis to further elucidate the micro-diversity of *Nitrospira*. This analysis identified 3

dominant oligotypes of *Nitrospira* and revealed their changes in microbial communities over

time (Figure 4A).

Figure 4 A) Oligotypes (3 total) of 16S rRNA gene amplicon sequences classified as genus

Nitrospira throughout the investigation period; 2) Positive correlation between the relative

abundance of Oligotype 2 (CCGGCTTAGTGGGTATTA) and the nitrite concentration.

Though the oligotype 1 consistently dominated the *Nitrospira* (> 71%), Oligotype 2 gradually

increased in tandem with the increasing nitrite accumulation. The similar increasing trends of

amoB/*nxrB* and the relative abundance of oligotype II indicates a positive relationship between

the comammox and the oligotype II *Nitrospira*. Note that comammox *Nitrospira* do not form

a monophyletic group in 16S rRNA gene sequence or *nxrB*-based phylogenetic analyses, but are

instead interspersed with strict nitrite-oxidizing *Nitrospira* (Daims et al., 2015; van Kessel et al.,

2015). Different comammox *Nitrospira* encode *amo* orthologs that dissimilar to each other, with

only ~60% amino acids identity between members of the different clades (Daims et al., 2015),

making it difficult to design a uniform comammox qPCR primers (Beach and Noguera, 2019;

Cotto et al., 2020). The *amoB* primer set used in this study was specifically optimized for clade

A comammox (Cotto et al., 2020). However, it is possible that clade B comammox also played a

role in our S2EBPR-PN/A processes, but current primers did not capture the whole comammox

community. Nevertheless, oligotyping and qPCR analysis indicated that a group of strict nitrite-

oxidizing *Nitrospira* decreased, while another group of *Nitrospira*, which might belong to

comammox and can oxidize ammonia increased ($r^2 = 0.60$, Figure 4B), partially contributing to

the nitrite accumulation. We also retrieved the representative sequence for the oligotypes

TTTTTCCGCGATAGGGAC (1) and CCGGCTTAGTGGGTATTA (2) and aligned them in the

nt database. However, the majority of the top hits found in NCBI's nt database are labelled as

"uncultured bacteria", indicating that the *Nitrospira* detected in our system is phylogenetic far

from the records of known reference sequences in *Nitrospira* genera. The genome information of

the *Nitrospira* in S2EBPR system needs further investigation.

3.3.3   Atypical ammonia oxidation bacteria

Fitzgerald et al. (2015) had proposed that the abundance of ammonia oxidizing organisms may

be underestimated, particularly in low DO condition (Fitzgerald et al., 2015). Despite the

presence of complete nitrification to nitrate, no known AOB were identified in their low DO

nitrification systems. To address this issue, they did pure culture and found 5 isolates that had not

previously been recognized as AOB but can oxidize NH_4 under both autotrophic and

heterotrophic conditions. They proposed that genus *Pseudomonas* and *Rhodococcus*, and family

Xanthomonadaceae and Sphingomonas are involved in heterotrophic nitrification under low-DO

conditions.

To further explore the potential ammonia oxidizers in our samples, the functional potential of

bacterial assemblages was predicted with PICRUSt2 in both enzymes and pathways. The

dominated enzyme genes are related to carbohydrate metabolism (e.g., glycolysis/gluconeogesis

metabolism), amino acid metabolism, DNA polymerase, and electron transport, which is

conceivable that all those functions are vital for microorganisms. Then we specifically aimed at

genes that can oxidize ammonia into nitrite. However, only three OTUs have the potential to

hold the methane/ammonia monooxygenase subunits (AMO, EC:1.14.18.3 and 1.14.99.39) that

oxidize ammonia into hydroxylamine, and they all belong to unclassified genus under order

Rhodocyclaceae. Another 11 OTUs belonging to Comamonadaceae_unclassified have the

potential to hold hydroxylamine oxidase (HAO, EC: 1.7.2.6) that can further oxidize

hydroxylamine to nitrite (Table S3 and Figure S4). Indeed, AMO and HAO were purified from

heterotrophic denitrifiers back in 1996 (Moir et al., 1996; Wehrfritz et al., 1996), and

heterotrophic nitrification has drawn a lot attention nowadays. 134 isolates were identified at the

genus/species level, and members from family Comamonadaceae and order Rhodocyclaceae

were reported to be aerobic denitrifiers (Duan et al., 2022).

In addition, the co-occurrence microbial patterns during nitrite accumulation periods revealed

that genus Nitrospira, midas_g_894, and midas_g_5535 are the main hubs for this specific

microbiome structure (Figure 3C). The latter two belong to the family Saprospiraceae and

Rhodocyclaceae, respectively, and the species in those families have all been shown to play a

role in autotrophic/heterotrophic nitrification (Duan et al., 2022; Mehrani et al., 2022).

Though the relative abundance of OTUs that are predicted to hold *hao* only represents a small

fraction of the entire Comamonadaceae_unclassified group (Figure S4), the prediction guides us

to look into the function played by Comamonadaceae. Though most heterotrophic nitrifiers can

oxidize ammonia to N_2O or N_2 by a full nitrification and denitrification pathway, some of them

are not capable of aerobic denitrification and ends with nitrite (Duan et al., 2022). A test with

enriched Comamonadaceae (~96%) showed ammonium oxidation potential under aerobic

conditions, with nitrite as the main product (Bao and Li, 2017), indicating the importance of

Comamonadaceae in partial nitrification. Some research also indicates that they contribute to

both nitrogen and phosphorus removal and can function as PAO (Ge et al., 2015). Our unique

coupling of A-B-stage with S2EBPR created unprecedented selection forces and conditions for

both P and N removal relevant organisms in this system, which would conceivably lead to

enrichment of organisms with versatile functions different from those in conventional EBPR or

PDN systems for main-stream treatment. In the current study, the relative abundance of

Comamonadaceae family remained stable at unusually high relative abundance level. (13%) in

the study period (Figure 3A) and they might contribute to nitrite accumulation.

It's important to note that the predicted functional potential was based on the current database,

which only includes the sequences of known AMO. Considering the fact that the *amo* genes in

comammox *Nitrospira* are phylogenetically distinct from previously identified AMOs (van

Kessel et al., 2015), it is possible that some other groups of bacteria are oxidizing ammonia but

cannot be captured by the current database.

3.4 Possible factors on nitrite accumulation

Another possible scenario to explain nitrite accumulation is that NOB activities were more

severely inhibited than that on AOB. Multiple strategies have been postulated to suppress NOB

in partial nitrification reactors, including low DO, transient anoxic condition, FNA inhibition, FA

inhibition, and VFA composition (Eilersen et al., 1994; Jiang et al., 2019; Park et al., 2010;

Wang et al., 2014). The average DO is 1.1 ± 0.9, 1.8 ± 0.7, 1.3 ± 0.5, 0.76 ± 0.51 mg/L in 4

CSTRs, respectively, in line with the DO range that is suitable for partial nitrification (Cao et al.,

2017; Regmi et al., 2014). However, more and more evidence show that DO regulation is not

reliable to suppress NOB alone (Park et al., 2010). Also, the FNA concentrations in both B-stage

influent and the mixing zone (B-stage influent + flow from Side-stream reactor) are nearly zero,

which did not provide inhibition pressure on NOB.

Compared to the conventional B-stage nutrient removal system, the coupling of S2EBPR with

PN created unique environmental selection conditions differ from those in conventional EBPR or

short-cut N system without EBPR. A split of RAS will flow through S2EBPR, where the unique

feature of S2EBPR system exerted different condition for nitrifying populations since they were

exposed to extended longer anaerobic time (HRT of anaerobic time is 3.65 hrs.) with lower ORP

$(-383 \pm 17$ mv) in comparison to conventional nutrient removal processes where AOB and NOB

generally experience shorter anoxic conditions (1-2 hrs) with shallow ORP level ($\sim$-150 mv). In

addition, high level (126.7 ± 72.1 mg/L as COD) and more complex VFA (Figure 5B and Table

S4, containing acetate, propionate, butyric, valeric acids) were added in side-stream reactor.

Indeed, the sCOD concentrations into SBRP highly covariates with the nitrite accumulation in B-

stage (Figure 5A). The sCOD mass in S2EBPR in this Period is 288 mg/L. When the carbon

mass in S2EBRP decreased at the end of the study period, the nitrite accumulation also declined.

This trend cannot be observed in B-stage reactor, indicating that side stream EBPR reactor is the

place where NOB was inhibited. Spearman correlation analysis indicated that the VFA

concentration into S2EBPR abundance is negatively correlated with the abundance of *Nitrospira*

and this influence is more profound when the VFA mass flow into S2EBPR is high (> 90 g/d)

(Figure 5E). The unique conditions in the S2EBPR, including the low ORP environment and the

VFAs concentration and composition, may play roles for the nitrite accumulation and they are

further discussed in the following sections.

Figure 5 (A) Soluble COD concentrations in SBPR, B-stage and nitrite accumulation; (B) VFA

speciation in A-stage fermentate; (C) VFA mass flow into SBPR and nitrite accumulation; (D)

VFA concentration in SBPR and AOB/NOB activities; (E) The Spearman correlation between

the most abundant 50 taxa and numerical environmental variables. The Spearman correlation test

is performed and associated p-values are adjusted for multiple testing.

3.4.1   VFA Inhibit NOB activity and alter NOB metabolic pathway:

Back to 1994, Eilersen et al. has reported that nitrite oxidation is more easily to be inhibited by

VFA than ammonia oxidation, especially acetic, propionic, and butyric acids (Table S5).

Especially, the butyric acid exerted strong inhibition on NOB activity (IC_{50} = 399 mg/L)

comparing to acetic (IC_{50} = 3707 mg/L) and propionic (IC_{50} = 3082 mg/L). Following research

confirmed that NOB activity was more susceptible to VFA inhibition comparing to AOB and the

inhibition level is proportional to the VFA molecular weight (Delgado et al., 2004; Gomez et al.,

2000). All those researchers came to this conclusion since the nitrate production decreased when

they added VFAs into NOB consortia. However, a later study showed that VFA (1500 mg/L)

addition did not influence the CO_2 fixation that is proportional to nitrite oxidation in NOB

consortia (*Nitrospira* spp.), indicating other metabolic pathway was activated by VFAs rather

than activity inhibition (Oguz et al., 2006). Though some *Nitrobacter* and *Nitrospira* benefit

from the presence of simple organic compounds, e.g., pyruvate, formate, urea (Daims et al.,

2016; Palomo et al., 2018), only comammox *Nitrospira* was reported to utilize fatty acids

(acetate and propionate) (Xu et al., 2022), and even have potential to synthesize and degrade

poly(3-hydroxybutyrate) (Palomo et al., 2018; Yang et al., 2020).

Comparing to the conventional WWTP that rely on mono external carbon source, such as

methanol or acetate, our system's A-stage fermentate contained a higher and stable proportion

(58.9%) of high molecular size VFAs, including propionic, butyric, iso-butyric and isovaleric

acids (Figure 5B). A-stage fermentate addition in S2EBPR may alter comammox *Nitrospira*

towards organic matter utilization pathway and down regulate the nitrite oxidation enzymes.

Notably, nitrite accumulation was not observed until the VFA mass flow was increased higher

than 66 g/d (Figure 5C). The activity tests also indicated that higher residual VFA concentration

(> 30 mg/L) in B-stage resulted lower NOB activity (in terms of nitrate production rates) ($p <$

0.01, Figure 5D), confirming the role of residual VFAs in alternating the NOB activity (Figure

5E). All those results indicated that implementation of S2EBPR with VFAs supplement seem to

favor the proliferation of versatile *Nitrospira* over canonical *Nitrospira*. This speculation was

supported by quick increase of the relative abundance of comammox *Nitrospira* over the total

Nitrospira community (indicated as *amoB*/*nxrB*) (Figure 4).

3.4.2   Longer exposure to anaerobic conditions with lower ORP impact NOB:

In our system, S2EBPR induced an extended anaerobic phase (~3.7 hours) where AOB and NOB

have to survive/maintain their activities for a longer time than typical duration in conventional

WWTPs. Some research reported that the endogenous decay rate of AOB (0.020 d^{-1}) was slower

than that of NOB (0.035 d^{-1}) in anoxic condition (Cui et al., 2019), this decay difference was

also supported by the FISH staining analysis Morgenroth et al., (2000). However, other results

showed identical decay rates of AOB and NOB under both anoxic and anaerobic conditions

(Munz et al., 2011; Salem et al., 2006), The differences may have been due to wastewater

characteristics, operation parameters, and the composition of the microbial communities and the

related survival strategies (Geets et al., 2006; Hao et al., 2010). The ORP in S2EBPR was $-383 \pm$

17 mV, which is significantly below the +100 to +350 mV in normal nitrification tank, and +50

to -50 mV in denitrification tank (Environmental, 2008). Generally, the endogenous decay rates

of nitrifiers decreases with the decreasing ORP condition, as $b_{aerobic} \gg b_{anoxic} > b_{anaerobic}$ (Geets et

al., 2006; Munz et al., 2011; Ruiz-Martínez et al., 2018; Salem et al., 2006). We assume that the

actual decay rates in our system would be even lower than 0.02 d^{-1}. Especially, the abundance of

Nitrospira was much higher than AOB throughout the experiment periods, indicating that the

decay differences between AOB/NOB did not play a significant role in NOB out-selection.

Other than low ORP, NOB is prone to be sensitive to the aerobic/anoxic perturbation. The nitrite

oxidation enzyme reactivation time of NOB is longer than AOB when they are exposed to

aerobic condition again (Gilbert et al., 2014; Kornaros et al., 2010). This characteristic has been

applied as intermittent aeration strategy in partial nitrification tank to suppress the NOB activity

(Kornaros et al., 2010). With more RAS split into S2EBPR (Figure S2), more AOB and NOB

population would undergo anaerobic and VFA exposure. With less versability, AOB may slow

down their metabolic activity while NOB may seek for other alternative pathway to maintain

their energy. This explains why they need a longer (enzyme) reactivation time (lag phase) to

change back to autotrophic nitrite oxidation in intermittent aeration systems (Gilbert et al., 2014)

and thus lead to nitrite accumulation.

4   Conclusions

Nitrite accumulation was observed along with good bio-P removal in a pilot-scale S2EBPR

implemented A-B stage process treating municipal wastewater.

(1) Average of 7.62 ± 2.17 mg-N/L nitrite accumulation with nitrite accumulation ratio of

49% was achieved by partial nitrification, which was verified through relative higher

ammonia oxidation activity (1.94 ± 0.89 mgN/gSS/h) than nitrite oxidation rate (1.06 ±

0.99 mgN/gSS/h).

(2) The integration of S2EBPR provides PAOs with a suitable environment, such as low

ORP and high VFA concentrations, but also inhibits NOB activity and provides

additional selective pressure nitrifying community.

(3) *Nitrospira* dominated the autotrophic nitrification community throughout the operation

periods with one to two orders of magnitude higher than AOB. The positive correlation

between *Nitrospira* oligo type 2 and nitrite accumulation suggested the contribution of

comammox *Nitrospira* as AOB.

(4) Phylogenetic investigation suggested that AMO and HAO were embedded in

heterotrophic organisms that belong to the family Comamonadaceae and order

Rhodocyclaceae. These organisms may function as heterotrophic nitrifiers in response to

elevated VFA addition.

5   Acknowledgement

This study was funded by the Water Environment Research Foundation (WERF, Grant No.

4901). We gratefully acknowledge the financial support and scientific discussion from Charles

B. Bott (HRSD), Jim McQuarrie (Denver Metro), Beverley M. Stinson (AECOM), Christine

deBarbadillo (DC Water), Paul Dombrowski (Woodard & Curran) and James Barnard (Black &

Veatch). Special thanks are given to the generous support of operators and staff of HRSD for

their assistance with sample and data collection.

6   Reference

An, Z., Kent, T.R., Sun, Y., Bott, C.B., Wang, Z.-W., 2021. Free ammonia resistance of nitrite-oxidizing bacteria developed in aerobic granular sludge cultivated in continuous upflow airlift reactors performing partial nitritation. Water Environment Research 93, 421–432. https://doi.org/10/gq26r9

Arciero, D.M., Balny, C., Hooper, A.B., 1991. Spectroscopic and rapid kinetic studies of reduction of cytochrome c554 by hydroxylamine oxidoreductase from Nitrosomonas europaea. Biochemistry 30, 11466–11472. https://doi.org/10/fhqgdt

Bao, P., Li, G.-X., 2017. Sulfur-Driven Iron Reduction Coupled to Anaerobic Ammonium Oxidation. Environ. Sci. Technol. 51, 6691–6698. https://doi.org/10/gbmh7c

Barbera, P., Kozlov, A.M., Czech, L., Morel, B., Darriba, D., Flouri, T., Stamatakis, A., 2019. EPA-ng: Massively Parallel Evolutionary Placement of Genetic Sequences. Systematic Biology 68, 365–369. https://doi.org/10/gd458h

Barnard, J.L., Dunlap, P., Steichen, M., 2017. Rethinking the Mechanisms of Biological Phosphorus Removal. Water Environment Research 89, 2043–2054. https://doi.org/10.2175/106143017x15051465919010

Bartram, A.K., Lynch, M.D.J., Stearns, J.C., Moreno-Hagelsieb, G., Neufeld, J.D., 2011. Generation of multi-million 16S rRNA gene libraries from complex microbial communities by assembling paired-end Illumina reads. Applied and environmental microbiology 77, 3846–3852.

Beach, N.K., Noguera, D.R., 2019. Design and Assessment of Species-Level qPCR Primers Targeting Comammox. Frontiers in Microbiology 10.

Boehnke, B., Diering, B., Zuckut, S.W., 1997. Cost-effective wastewater treatment process for removal of organics and nutrients. Water Engineering and Management 144, 18–21.

Cao, Y., van Loosdrecht, M.C.M., Daigger, G.T., 2017. Mainstream partial nitration–anammox in municipal wastewater treatment: status, bottlenecks, and further studies. Appl Microbiol Biotechnol 101, 1365–1383. https://doi.org/10/ghbhhp

Carvalheira, M., Oehmen, A., Carvalho, G., Eusébio, M., Reis, M.A.M., 2014. The impact of aeration on the competition between polyphosphate accumulating organisms and glycogen accumulating organisms. Water Research 66, 296–307. https://doi.org/10/f6p578

Cotto, I., Dai, Z., Huo, L., Anderson, C.L., Vilardi, K.J., Ijaz, U., Khunjar, W., Wilson, C., De Clippeleir, H., Gilmore, K., Bailey, E., Pinto, A.J., 2020. Long solids retention times and attached growth phase favor prevalence of comammox bacteria in nitrogen removal systems. Water Research 169, 115268. https://doi.org/10/gqxst8

Cui, H., Zhang, L., Zhang, Q., Li, X., Peng, Y., 2019. Stable partial nitrification of domestic sewage achieved through activated sludge on exposure to nitrite. Bioresource Technology 278, 435–439. https://doi.org/10/grv5t4

Czech, L., Barbera, P., Stamatakis, A., 2020. Genesis and Gappa: processing, analyzing and
visualizing phylogenetic (placement) data. Bioinformatics 36, 3263–3265.
https://doi.org/10/ghkddf

Daims, H., Lebedeva, E.V., Pjevac, P., Han, P., Herbold, C., Albertsen, M., Jehmlich, N.,
Palatinszky, M., Vierheilig, J., Bulaev, A., Kirkegaard, R.H., von Bergen, M., Rattei, T.,
Bendinger, B., Nielsen, P.H., Wagner, M., 2015. Complete nitrification by Nitrospira
bacteria. Nature 528, 504–509. https://doi.org/10/f3s3d3

Daims, H., Lücker, S., Wagner, M., 2016. A New Perspective on Microbes Formerly Known as
Nitrite-Oxidizing Bacteria. Trends Microbiol 24, 699–712. https://doi.org/10/f82hxv

Delgado, A., Aspé, E., Martí, M.C., Roeckel, M., 2004. The Effect of Volatile Fatty Acids on the
Nitrification of a Saline Effluent. Environmental Technology 25, 413–422.
https://doi.org/10/c43jcn

Douglas, G.M., Maffei, V.J., Zaneveld, J.R., Yurgel, S.N., Brown, J.R., Taylor, C.M.,
Huttenhower, C., Langille, M.G.I., 2020. PICRUSt2 for prediction of metagenome
functions. Nat Biotechnol 38, 685–688. https://doi.org/10/gg5v22

Duan, S., Zhang, Y., Zheng, S., 2022. Heterotrophic nitrifying bacteria in wastewater biological
nitrogen removal systems: A review. Critical Reviews in Environmental Science and
Technology 52, 2302–2338. https://doi.org/10/gkzvwc

Eilersen, A.M., Henze, M., Kløft, L., 1994. Effect of Volatile Fatty Acids and Trimethylamine
on Nitrification in Activated Sludge. Water Research 28, 1329–1336.
https://doi.org/10.1016/0043-1354(94)90298-4

Environmental, Y.S.I., 2008. ORP Management in wastewater as an indicator of process
efficiency. YSI, Yellow Springs, OH http://www. ysi. com/media/pdfs/A567-ORP-
Management-in-Wastewater-as-an-Indicator-of-Process-Efficiency. pdf (accessed on
15.08. 13).

Eren, A.M., Maignien, L., Sul, W.J., Murphy, L.G., Grim, S.L., Morrison, H.G., Sogin, M.L.,
2013. Oligotyping: differentiating between closely related microbial taxa using 16S
rRNA gene data. Methods Ecol Evol 4, 1111–1119. https://doi.org/10/f5jsk9

Fitzgerald, C.M., Camejo, P., Oshlag, J.Z., Noguera, D.R., 2015. Ammonia-oxidizing microbial
communities in reactors with efficient nitrification at low-dissolved oxygen. Water
Research 70, 38–51. https://doi.org/10/f64qdj

Gao, R., Peng, Y., Li, J., Zhang, Q., Li, X., Deng, L., Li, W., Kao, C., 2021. Improving
performance and efficiency of partial anammox by coupling partial nitrification and
partial denitrification (PN/A-PD/A) to treat municipal sewage in a step-feed reactor.
Bioresource Technology 341, 125804. https://doi.org/10/gqmxsm

Ge, H., Batstone, D.J., Keller, J., 2015. Biological phosphorus removal from abattoir wastewater
at very short sludge ages mediated by novel PAO clade Comamonadaceae. Water
Research 69, 173–182. https://doi.org/10.1016/j.watres.2014.11.026

Geets, J., Boon, N., Verstraete, W., 2006. Strategies of aerobic ammonia-oxidizing bacteria for coping with nutrient and oxygen fluctuations. FEMS Microbiology Ecology 58, 1–13. https://doi.org/10/b8hws5

Gilbert, E.M., Agrawal, S., Brunner, F., Schwartz, T., Horn, H., Lackner, S., 2014. Response of Different Nitrospira Species To Anoxic Periods Depends on Operational DO. Environ. Sci. Technol. 48, 2934–2941. https://doi.org/10/f5zq9f

Gomez, J., Mendez, R., Lema, J.M., 2000. Kinetic study of addition of volatile organic compounds to a nitrifying sludge. Appl Biochem Biotechnol 87, 189–202. https://doi.org/10/dq46rz

Gu, A.Z., Saunders, A., Neethling, J.B., Stensel, H.D., Blackall, L.L., 2008. Functionally Relevant Microorganisms to Enhanced Biological Phosphorus Removal Performance at Full-Scale Wastewater Treatment Plants in the United States. Water Environment Research 80, 688–698. https://doi.org/10.2175/106143008x276741

Gu, A.Z., Tooker, N.B., Onnis-Hayden, A., Wang, D., Li, G., Srinivasan, V., Takács, I., 2018. Investigation of the Mechanisms for Optimization and Design of a Side-Stream EBPR Process as a Sustainable Approach for Achieving Stable and Efficient Phosphorus Removal. by Water Environmental Research Foundation (WERF) and International Water Association (IWA) Publishing.

Hao, X.-D., Wang, Q.-L., Zhu, J.-Y., Van Loosdrecht, M.C.M., 2010. Microbiological Endogenous Processes in Biological Wastewater Treatment Systems. Critical Reviews in Environmental Science and Technology 40, 239–265. https://doi.org/10/csmtq8

He, S., Gall, D.L., McMahon, K.D., 2007. "Candidatus Accumulibacter" Population Structure in Enhanced Biological Phosphorus Removal Sludges as Revealed by Polyphosphate Kinase Genes. Applied and Environmental Microbiology 73, 5865–5874. https://doi.org/10/dcfkfh

Hellinga, C., Schellen, A., Mulder, J.W., van Loosdrecht, M.C.M. v, Heijnen, J.J., 1998. The SHARON process: an innovative method for nitrogen removal from ammonium-rich waste water. Water science and technology 37, 135–142.

Jetten, M.S.M., Horn, S.J., van Loosdrecht, M.C.M., 1997. Towards a more sustainable municipal wastewater treatment system. Water Science and Technology, Sustainable Sanitation 35, 171–180. https://doi.org/10/bm4qhb

Jiang, C., Xu, S., Wang, R., Feng, S., Zhou, S., Wu, Shimin, Zeng, X., Wu, Shanghua, Bai, Z., Zhuang, G., Zhuang, X., 2019. Achieving efficient nitrogen removal from real sewage via nitrite pathway in a continuous nitrogen removal process by combining free nitrous acid sludge treatment and DO control. Water Research 161, 590–600. https://doi.org/10/gph6pq

Joss, A., Salzgeber, D., Eugster, J., König, R., Rottermann, K., Burger, S., Fabijan, P., Leumann, S., Mohn, J., Siegrist, H.R., 2009. Full-scale nitrogen removal from digester liquid with partial nitration and anammox in one SBR. Environmental Science and Technology 43, 5301–5306. https://doi.org/10.1021/es900107w

Kinyua, M.N., Miller, M.W., Wett, B., Murthy, S., Chandran, K., Bott, C.B., 2017. Polyhydroxyalkanoates, triacylglycerides and glycogen in a high rate activated sludge A-stage system. Chemical Engineering Journal 316, 350–360. https://doi.org/10/f94pm3

Kits, K.D., Sedlacek, C.J., Lebedeva, E.V., Han, P., Bulaev, A., Pjevac, P., Daebeler, A., Romano, S., Albertsen, M., Stein, L.Y., Daims, H., Wagner, M., 2017. Kinetic analysis of a complete nitrifier reveals an oligotrophic lifestyle. Nature 549, 269–272. https://doi.org/10/gbs6kc

Klaus, S.A., 2019. Intensification of Biological Nutrient Removal Processes (Doctor of Philosophy). Virginia Polytechnic Institute and State University, Blacksburt, Virginia.

Klaus, S.A., Sadowski, M.S., Kinyua, M.N., Miller, M.W., Regmi, P., Wett, B., De Clippeleir, H., Chandran, K., Bott, C.B., 2020. Effect of influent carbon fractionation and reactor configuration on mainstream nitrogen removal and NOB out-selection. Environmental Science: Water Research and Technology 6, 691–701. https://doi.org/10.1039/c9ew00873j

Koch, H., Lücker, S., Albertsen, M., Kitzinger, K., Herbold, C., Spieck, E., Nielsen, P.H., Wagner, M., Daims, H., 2015. Expanded metabolic versatility of ubiquitous nitrite-oxidizing bacteria from the genus Nitrospira. PNAS 112, 11371–11376. https://doi.org/10/f3sxtj

Kornaros, M., Dokianakis, S.N., Lyberatos, G., 2010. Partial Nitrification/Denitrification Can Be Attributed to the Slow Response of Nitrite Oxidizing Bacteria to Periodic Anoxic Disturbances. Environ. Sci. Technol. 44, 7245–7253. https://doi.org/10/dwnr6j

Kozich, J.J., Westcott, S.L., Baxter, N.T., Highlander, S.K., Schloss, P.D., 2013. Development of a dual-index sequencing strategy and curation pipeline for analyzing amplicon sequence data on the MiSeq Illumina sequencing platform. Applied and environmental microbiology 79, 5112–5120. https://doi.org/10/f46m6z

Lackner, S., Gilbert, E.M., Vlaeminck, S.E., Joss, A., Horn, H., van Loosdrecht, M.C.M., 2014. Full-scale partial nitration/anammox experiences – An application survey. Water Research 55, 292–303. https://doi.org/10/f52dp8

Liu, G., Wang, J., 2013. Long-Term Low DO Enriches and Shifts Nitrifier Community in Activated Sludge. Environ. Sci. Technol. 47, 5109–5117. https://doi.org/10/f4w4gz

Lotti, T., Kleerebezem, R., Hu, Z., Kartal, B., De Kreuk, M.K., Van Erp Taalman Kip, C., Kruit, J., Hendrickx, T.L.G., Van Loosdrecht, M.C.M., 2015. Pilot-scale evaluation of anammox-based mainstream nitrogen removal from municipal wastewater. Environmental Technology (United Kingdom) 36, 1167–1177. https://doi.org/10.1080/09593330.2014.982722

Louca, S., Doebeli, M., 2018. Efficient comparative phylogenetics on large trees. Bioinformatics 34, 1053–1055. https://doi.org/10/gc6265

Masella, A.P., Bartram, A.K., Truszkowski, J.M., Brown, D.G., Neufeld, J.D., 2012. PANDAseq: paired-end assembler for illumina sequences. BMC bioinformatics 13, 31.

McCullough, K., Klaus, S., Schoepflin, S., Wilson, C., Gu, A., Han, I., Clippeleir, H. de, Bott,
C., 2020. Simultaneous N and P Removal with Anammox and Sidestream BioP.
Presented at the WEFTEC 2020, Water Environment Federation.

Mehrani, M.-J., Sobotka, D., Kowal, P., Guo, J., Mąkinia, J., 2022. New insights into modeling
two-step nitrification in activated sludge systems – The effects of initial biomass
concentrations, comammox and heterotrophic activities. Science of The Total
Environment 848, 157628. https://doi.org/10/grz33r

Moir, J.W.B., Crossman, L.C., Spiro, S., Richardson, D.J., 1996. The purification of ammonia
monooxygenase from Paracoccus denitrficans. FEBS Letters 387, 71–74.
https://doi.org/10/bwc7p3

Morgenroth, E., Obermayer, A., Arnold, E., Brühl, A., Wagner, M., Wilderer, P.A., 2000. Effect
of long-term idle periods on the performance of sequencing batch reactors. Water Science
and Technology 41, 105–113. https://doi.org/10/gqzrpc

Munz, G., Lubello, C., Oleszkiewicz, J.A., 2011. Factors affecting the growth rates of
ammonium and nitrite oxidizing bacteria. Chemosphere 83, 720–725.
https://doi.org/10/c579xz

Norton, J.M., Alzerreca, J.J., Suwa, Y., Klotz, M.G., 2002. Diversity of ammonia
monooxygenase operon in autotrophic ammonia-oxidizing bacteria. Arch Microbiol 177,
139–149. https://doi.org/10/cp85t6

Oguz, M.T., Robinson, K.G., Layton, A.C., Sayler, G.S., 2006. Volatile fatty acid impacts on
nitrite oxidation and carbon dioxide fixation in activated sludge. Water Research 40,
665–674. https://doi.org/10/cpfbkj

Onnis-Hayden, A., Srinivasan, V., Tooker, N.B., Li, G., Wang, D., Barnard, J.L., Bott, C.,
Dombrowski, P., Schauer, P., Menniti, A., Shaw, A., Stinson, B., Stevens, G., Dunlap, P.,
Takács, I., McQuarrie, J., Phillips, H., Lambrecht, A., Analla, H., Russell, A., Gu, A.Z.,
2020. Survey of full-scale sidestream enhanced biological phosphorus removal
(S2EBPR) systems and comparison with conventional EBPRs in North America: Process
stability, kinetics, and microbial populations. Water Environment Research 92, 403–417.
https://doi.org/10/gn4whz

Palomo, A., Pedersen, A.G., Fowler, S.J., Dechesne, A., Sicheritz-Pontén, T., Smets, B.F., 2018.
Comparative genomics sheds light on niche differentiation and the evolutionary history
of comammox Nitrospira. ISME J 12, 1779–1793. https://doi.org/10/gdb9m2

Parada, A.E., Needham, D.M., Fuhrman, J.A., 2016. Every base matters: assessing small subunit
rRNA primers for marine microbiomes with mock communities, time series and global
field samples. Environmental microbiology 18, 1403–1414.

Park, S., Bae, W., Rittmann, B.E., 2010. Operational Boundaries for Nitrite Accumulation in
Nitrification Based on Minimum/Maximum Substrate Concentrations That Include
Effects of Oxygen Limitation, pH, and Free Ammonia and Free Nitrous Acid Inhibition.
Environ. Sci. Technol. 44, 335–342. https://doi.org/10/c5vtq7

Pester, M., Maixner, F., Berry, D., Rattei, T., Koch, H., Lücker, S., Nowka, B., Richter, A.,
Spieck, E., Lebedeva, E., Loy, A., Wagner, M., Daims, H., 2014. NxrB encoding the beta
subunit of nitrite oxidoreductase as functional and phylogenetic marker for nitrite-
oxidizing Nitrospira. Environmental Microbiology 16, 3055–3071.
https://doi.org/10/f3src8

Printz, K.E., 2019. The Investigation of Nitrite Accumulation and Biological Phosphorus
Removal in an Intermittently Aerated Process Combining Shortcut Nitrogen Removal
and Sidestream Biological Phosphorus Removal (Thesis). Virginia Tech.

Quince, C., Lanzen, A., Davenport, R.J., Turnbaugh, P.J., 2011. Removing noise from
pyrosequenced amplicons. BMC bioinformatics 12, 38.

Regmi, P., Miller, M.W., Holgate, B., Bunce, R., Park, H., Chandran, K., Wett, B., Murthy, S.,
Bott, C.B., 2014. Control of aeration, aerobic SRT and COD input for mainstream
nitritation/denitritation. Water Research 57, 162–171.
https://doi.org/10.1016/j.watres.2014.03.035

Ren, Z. (Jason), Pagilla, K. (Eds.), 2022. Pathways to Water Sector Decarbonization, Carbon
Capture and Utilization. IWA Publishing. https://doi.org/10.2166/9781789061796

Rittmann, B.E., Mayer, B., Westerhoff, P., Edwards, M., 2011. Capturing the lost phosphorus.
Chemosphere 84, 846–853. https://doi.org/10.1016/j.chemosphere.2011.02.001

Roots, P., Wang, Y., Rosenthal, A.F., Griffin, J.S., Sabba, F., Petrovich, M., Yang, F., Kozak,
J.A., Zhang, H., Wells, G.F., 2019. Comammox Nitrospira are the dominant ammonia
oxidizers in a mainstream low dissolved oxygen nitrification reactor. Water Research
157, 396–405. https://doi.org/10/gg6rhv

Rotthauwe, J.H., Witzel, K.P., Liesack, W., 1997. The ammonia monooxygenase structural gene
amoA as a functional marker: molecular fine-scale analysis of natural ammonia-oxidizing
populations. Applied and Environmental Microbiology 63, 4704–4712.
https://doi.org/10/ghzcgs

Ruiz-Martínez, A., Claros, J., Serralta, J., Bouzas, A., Ferrer, J., 2018. Modeling the decay of
nitrite oxidizing bacteria under different reduction potential conditions. Process
Biochemistry 71, 159–165. https://doi.org/10/gd9fwp

Salem, S., Moussa, M. s., van Loosdrecht, M. c. m., 2006. Determination of the decay rate of
nitrifying bacteria. Biotechnology and Bioengineering 94, 252–262.
https://doi.org/10/fc5zz7

Sancho, I., Lopez-Palau, S., Arespacochaga, N., Cortina, J.L., 2019. New concepts on carbon
redirection in wastewater treatment plants: A review. Science of the Total Environment
647, 1373–1384. https://doi.org/10.1016/j.scitotenv.2018.08.070

Shao, Y.-H., Wu, J.-H., 2021. Comammox Nitrospira Species Dominate in an Efficient Partial
Nitrification–Anammox Bioreactor for Treating Ammonium at Low Loadings. Environ.
Sci. Technol. 55, 2087–2098. https://doi.org/10/gqmxwm

Srinivasan, V.N., Li, G., Wang, D., Tooker, N.B., Dai, Z., Onnis-Hayden, A., Bott, C.,
Dombrowski, P., Schauer, P., Pinto, A., Gu, A.Z., 2021. Oligotyping and metagenomics

reveal distinct Candidatus Accumulibacter communities in side-stream versus conventional full-scale enhanced biological phosphorus removal (EBPR) systems. Water Research 206, 117725. https://doi.org/10/gq82kd

Ssekagiri, A., Sloan, W., Zeeshan Ijaz, U., 2017. microbiomeSeq: An R package for analysis of microbial communities in an environmental context. ISCB Africa ASBCB Conference.

Torondel, B., Ensink, J.H.J., Gundogdu, O., Ijaz, U.Z., Parkhill, J., Abdelahi, F., Nguyen, V.-A., Sudgen, S., Gibson, W., Walker, A.W., Quince, C., 2016. Assessment of the influence of intrinsic environmental and geographical factors on the bacterial ecology of pit latrines. Microbial Biotechnology 9, 209–223. https://doi.org/10/f8ttfq

Van Dongen, U., Jetten, M.S.M., Loosdrecht, M.C.M. Van, 2001. Ammonium Rich Wastewater. Growth Lakeland 44, 153–160.

van Kessel, M.A.H.J., Speth, D.R., Albertsen, M., Nielsen, P.H., Op den Camp, H.J.M., Kartal, B., Jetten, M.S.M., Lücker, S., 2015. Complete nitrification by a single microorganism. Nature 528, 555–559. https://doi.org/10/f73qs2

Villanueva, R.A.M., Chen, Z.J., 2019. ggplot2: Elegant Graphics for Data Analysis (2nd ed.). Measurement: Interdisciplinary Research and Perspectives 17, 160–167. https://doi.org/10/ggqrmf

Vollertsen, J., Petersen, G., Borregaard, V.R., 2006. Hydrolysis and fermentation of activated sludge to enhance biological phosphorus removal. Water science and technology 53, 55–64.

Wang, D., Tooker, N.B., Srinivasan, V., Li, G., Fernandez, L.A., Schauer, P., Menniti, A., Maher, C., Bott, C.B., Dombrowski, P., Barnard, J.L., Onnis-Hayden, A., Gu, A.Z., 2019. Side-stream enhanced biological phosphorus removal (S2EBPR) process improves system performance - A full-scale comparative study. Water Research 167, 115109. https://doi.org/10.1016/j.watres.2019.115109

Wang, Q., Duan, H., Wei, W., Ni, B.-J., Laloo, A., Yuan, Z., 2017. Achieving Stable Mainstream Nitrogen Removal via the Nitrite Pathway by Sludge Treatment Using Free Ammonia. Environ. Sci. Technol. 51, 9800–9807. https://doi.org/10/gnn87b

Wang, Q., Ye, L., Jiang, G., Hu, S., Yuan, Z., 2014. Side-stream sludge treatment using free nitrous acid selectively eliminates nitrite oxidizing bacteria and achieves the nitrite pathway. Water Research 55, 245–255. https://doi.org/10/f5z7kk

Wang, Shanyun, Peng, Y., Ma, B., Wang, Shuying, Zhu, G., 2015. Anaerobic ammonium oxidation in traditional municipal wastewater treatment plants with low-strength ammonium loading: Widespread but overlooked. Water Research 84, 66–75. https://doi.org/10/f7s2cs

Wehrfritz, J.-M., Carter, J.P., Spiro, S., Richardson, D.J., 1996. Hydroxylamine oxidation in heterotrophic nitrate-reducing soil bacteria and purification of a hydroxylamine-cytochromec oxidoreductase from aPseudomonas species. Arch. Microbiol. 166, 421–424. https://doi.org/10/bmwckj

Winkler, M.K.H., Bassin, J.P., Kleerebezem, R., Sorokin, D.Y., van Loosdrecht, M.C.M., 2012.
Unravelling the reasons for disproportion in the ratio of AOB and NOB in aerobic
granular sludge. Appl Microbiol Biotechnol 94, 1657–1666. https://doi.org/10/gcbkvb

Xu, G., Xu, X., Yang, F., Liu, S., Gao, Y., 2012. Partial nitrification adjusted by hydroxylamine
in aerobic granules under high DO and ambient temperature and subsequent Anammox
for low C/N wastewater treatment. Chemical Engineering Journal 213, 338–345.
https://doi.org/10.1016/j.cej.2012.10.014

Xu, S., Chai, W., Xiao, R., Smets, B.F., Palomo, A., Lu, H., 2022. Survival strategy of
comammox bacteria in a wastewater nutrient removal system with sludge fermentation
liquid as additional carbon source. Science of The Total Environment 802, 149862.
https://doi.org/10/grv6nq

Yang, Y., Daims, H., Liu, Y., Herbold, C.W., Pjevac, P., Lin, J.-G., Li, M., Gu, J.-D., 2020.
Activity and Metabolic Versatility of Complete Ammonia Oxidizers in Full-Scale
Wastewater Treatment Systems. mBio 11, e03175-19. https://doi.org/10/gn5874

Zhang, T., Wang, B., Li, X., Zhang, Q., Wu, L., He, Y., Peng, Y., 2018. Achieving partial
nitrification in a continuous post-denitrification reactor treating low C/N sewage.
Chemical Engineering Journal 335, 330–337. https://doi.org/10/gdf65t

Zhou, Y., Oehmen, A., Lim, M., Vadivelu, V., Ng, W.J., 2011. The role of nitrite and free
nitrous acid (FNA) in wastewater treatment plants. Water Research 45, 4672–4682.
https://doi.org/10/ff49qr

809