For: eLife

Water-soluble 4-(dimethylaminomethyl)heliomycin exerts greater antitumor effects than parental

heliomycin by targeting the tNOX-SIRT1 axis and apoptosis in oral cancer cells

Atikul Islam [1], Yu-Chun Chang [1], Xiao-Chi Chen [1], Chia-Wei Weng [1,2], Chien-Yu Chen [1], Che-Wei

Wang [3,4], Mu-Kuan Chen [3,4], Alexander S. Tikhomirov [5], Andrey E. Shchekotikhin [5] and Pin Ju

Chueh [1,4,6,7,]*

[1] Institute of Biomedical Sciences, National Chung Hsing University, Taichung, 40227, Taiwan

[2] Institute of Medicine, Chung Shan Medical University, Taichung 40201, Taiwan

[3] Department of Otorhinolaryngology-Head and Neck Surgery, Changhua Christian Hospital,

Changhua 50006, Taiwan

[4] Department of Post-Baccalaureate Medicine, College of Medicine, National Chung Hsing

University, Taichung, 40227, Taiwan

[5] Gause Institute of New Antibiotics, 11 B. Pirogovskaya Street, Moscow 119021, Russia

[6] Department of Medical Research, China Medical University Hospital, Taichung, 40402, Taiwan

[7] Graduate Institute of Basic Medicine, China Medical University, Taichung, 40402, Taiwan

Running Title: Antitumor effects of 4-dimethylaminomethylheliomycin by targeting tNOX-SIRT1

axis

* Correspondence to: Pin Ju Chueh, Ph.D., Institute of Biomedical Sciences, National Chung Hsing

University, Taichung, Taiwan

Tel: +886 4 22840896; Fax: +886 4 22853469

E-mail addresses: islammiu555@gmail.com (AI), gill870715@hotmail.com (YCC),

kiki1232268@gmail.com (XCC), cwweng060304@gmail.com (CWW), z920911419@gmail.com

(CYC), 97302@cch.org.tw (CWW), 53780@cch.org.tw (MKC), tikhomirov.chem@gmail.com

(AST), shchekotikhin@gause-inst.ru (AES), pjchueh@dragon.nchu.edu.tw (PJC)

Abstract

The antibiotic heliomycin (resistomycin), which is generated from *Streptomyces resistomycificus*, has multiple activities, including anticancer effects. Heliomycin was first described in the 1960s, but its clinical applications have been hindered by extremely low solubility. A series of 4-aminomethyl derivatives of heliomycin were synthesized to increase water solubility; studies showed that they had anti-proliferative effects, but the drug targets remained unknown. In this study, we conducted cellular thermal shift assays (CETSA) and molecular docking simulations to identify and validate that heliomycin and its water-soluble derivative, 4-(dimethylaminomethyl)heliomycin (designated compound 4-dmH) engaged and targeted with sirtuin-1 (SIRT1) in p53-functional SAS and p53-mutated HSC-3 oral cancer cells. We further addressed the cellular outcome of SIRT1 inhibition by these compounds and found that, in addition to SIRT1, the water-soluble 4-dmH preferentially targeted a tumor-associated NADH oxidase (tNOX, ENOX2). The direct binding of 4-dmH to tNOX decreased the oxidation of NADH to NAD^+ which diminished NAD^+-dependent SIRT1 deacetylase activity, ultimately inducing apoptosis and significant cytotoxicity in both cell types, as opposed to the parental heliomycin-induced autophagy. We also observed that tNOX and SIRT1 were both upregulated in tumor tissues of oral cancer patients compared to adjacent normal tissues, suggesting their clinical relevance. Finally, the better therapeutic efficacy of 4-dmH was confirmed in tumor-bearing mice, which showed greater tNOX and SIRT1 downregulation and tumor volume reduction when treated with 4-dmH compared to

heliomycin. Taken together, our *in vitro* and *in vivo* findings suggest that the multifaceted properties

of water-soluble 4-dmH enable it to offer superior antitumor value compared to parental heliomycin,

and indicated that it functions through targeting the tNOX-NAD$^+$-SIRT1 axis to induce apoptosis in

oral cancer cells.

Keywords: Apoptosis; Heliomycin; Resistomycin; Sirtuin 1 (SIRT1); Tumor-associated NADH

oxidase (tNOX, ENOX2); Oral cancer cells

Background

Heliomycin (also known as resistomycin) was identified as a secondary metabolite produced by *Streptomyces resistomycificus* and has been reported to have antiviral [1, 2], antifungal [3], antibacterial [4], antimicrobial, RNA synthesis-inhibiting [5], and HIV-1 protease-suppressing [6] activities. It also possesses cytotoxicity against cancer cells from different tissues, such as cervical, gastric, hepatic, and breast cancers [4, 7-9]. Interestingly, heliomycin isolated from marine sponges was reported to inhibit histone deacetylases (HDACs), as assessed by colorimetric assays and *in silico* docking studies [10]. Among the HDACs, sirtuin-1 (SIRT1), a member in the Sir2 family of histone deacetylases, acts in different cellular compartments to remove acetyl groups from histone and non-histone proteins, and thereby modulate various cell functions, including metabolism, healthspan, tumorigenesis, apoptosis, and autophagy [11-19]. Given its position at the intersection between apoptosis and autophagy, it is not surprising that SIRT1 contributes to a complex regulatory network in which conventional signaling molecules undergo multifaceted interactions [20-22], however, no study has investigated the link between SIRT1 and heliomycin.

In an effort to identify cellular targets of heliomycin, we previously conducted a cellular thermal shift assays (CETSA), which is based on the concept that ligand binding augments the heat resistance of a target protein in intact cells [23, 24]. Data from CETSA and molecular docking simulations provided direct and experimental evidence linking heliomycin to SIRT1, which exhibited an enhanced melting temperature when incubated with heliomycin [25]. Our results also

demonstrated that heliomycin treatment downregulated SIRT1 expression and that the binding of

heliomycin to SIRT1 triggered autophagy to repress the growth of bladder cancer cells. Given that

the medical applications of heliomycin have been limited due to its extremely low solubility in

aqueous media, we synthesized a series of chemically modified heliomycin derivatives. We found

that their cytotoxicity was similar to that of the reference anticancer drug doxorubicin in several

cancer cell lines [26, 27]. To continue with this line of investigation, we herein sought to elucidate

the molecular mechanism and identify the cellular targets of the novel water-soluble

4-((dimethylamino)methyl)heliomycin (designated 4-dmH). We report that in addition to targeting

SIRT1, as does its parental heliomycin, 4-dmH engages with the tumor-associated NADH oxidase

(tNOX, ENOX2), as analyzed by CETSAs and molecular docking studies. Upon this binding,

tNOX is ubiquitinylated and degraded, which in turn attenuates signaling by the

tNOX-NAD$^+$-SIRT1 regulatory axis and induces apoptosis in oral cancer cells.

88
89

Results

Both heliomycin and its water-soluble derivative, 4-dmH, target intracellular SIRT1

We previously synthesized a series of heliomycin-derived Mannich bases with enhanced water

solubility and demonstrated that they possessed anticancer activity against different cancer cell lines

[26]; however, the cellular targets and molecular mechanisms of heliomycin and these derivatives

remained unknown. Very recently, SIRT1 was shown to function as a cellular target for heliomycin,

and this targeting was identified as being important for the anticancer activity of heliomycin in

bladder cancer cells [25]. To assess whether the same was true for one of our novel water-soluble

derivatives, 4-dmH, we first generated CETSA-based isothermal dose-response fingerprint curves

(ITDRF$_{CETSA}$). Our results indicated that the heat resistance of SIRT1 was positively and

dose-dependently associated with the water-soluble 4-dmH, with an OC$_{50}$ value of 1.0 μM for SAS

cells and 0.9 μM for HSC-3 cells (Fig. 2a). We next investigated the SIRT1-targeting ability of

4-dmH by determining the melting temperature (T_M) from CETSA melting curves, and found that

the T_M values were elevated from 47.0°C (control) to 56.6°C (4-dmH-exposed) in SAS cells, and

from 47.2 °C (control) to 58.8°C (4-dmH-treated) in HSC-3 cells (Fig. 2b).

ITDRF$_{CETSA}$ also indicated that the parental heliomycin engaged with SIRT1 in oral cancer

cells, as supported by the positive correlation of SIRT1 heat resistance with the heliomycin

concentration (Fig. 3a). Similarly, heliomycin treatment was associated with a greater than a

5-degree increase in T_M values calculated from the CETSA melting curves of two oral cancer cell

lines (Fig. 3b). Our CETSA-based results suggested that the test compounds altered the unfolding

and aggregation properties of SIRT1 in response to heat challenge, indicating that both 4-dmH and

heliomycin biophysically interact with SIRT1 in oral cancer cells.

We next determined whether these compounds affect the *in vitro* deacetylase activity of SIRT1

using a fluorogenic substrate and recombinant protein, and found that both heliomycin and 4-dmH

effectively inhibited SIRT1 deacetylase activity *in vitro* at 20 and 200 μM (Fig. 4a). The degree of

inhibition was comparable to that reported for the specific human SIRT1 inhibitor, sirtinol (IC$_{50}$,

131 μM) [28]. Given that heliomycin was previously reported to target and downregulate SIRT1 in

bladder cancer cells [25], in this present study, we addressed the question of SIRT1 inhibition by

4-dmH and its associated molecular events. We found that 4-dmH markedly attenuated SIRT1

protein expression at 1 and 2 μM for SAS cells and at 2 μM for HSC-3 cells (Fig. 4b), and that

pretreatment with the proteasome inhibitor, MG132, markedly reversed 4-dmH-inhibited SIRT1

expression in both cell lines (Fig. 4c). Our immunoprecipitation results showed that 4-dmH

increased the ubiquitination level of SIRT1 (Fig. 4d), suggesting that 4-dmH may downregulate

SIRT1 via ubiquitin-proteasomal degradation.

Heliomycin provokes autophagy whereas 4-dmH triggers apoptosis

To further address the cellular outcome of 4-dmH-inhibited SIRT1, we explored the cell death

pathway and signaling mechanisms involved in the system. As heliomycin was reported to induce

autophagy in bladder cancer cells [25], we used AO staining (which detects acidic vesicular

organelles) to assess whether heliomycin and/or water-soluble 4-dmH induced autophagy in oral

cancer cells. Our results showed that heliomycin markedly enhanced autophagy in SAS cells at 1

and 2 µM and in HSC-3 cells at 2 µM; on the contrary, water-soluble 4-dmH largely failed to

provoke autophagy in either line at concentrations up to 2 µM (Fig. 5a). The lack of autophagic

activity of 4-dmH prompted us to explore apoptosis, which is a cell death-related pathway

previously reported to be triggered by a different heliomycin derivative in bladder cancer cells [25,

26]. Interestingly, we found that 2 µM of 4-dmH, but not the parental heliomycin, effectively

triggered JC-1 staining; this denotes a loss of mitochondrial membrane potential (MMP), which is

an indicator of apoptosis (Fig. 5b). The heliomycin-induced autophagy was accompanied by

upregulation of autophagy markers, ULK1, Atg5/Atg7, and cleaved LC3-II, but downregulation of

Bcl-2 (Fig. 5c top). The ineffective apoptosis by heliomycin was evidenced by the downregulation

of pro-apoptotic Bak and Puma and a lack of caspases 3-directed PARP cleavage and c-Flip

downregulation (Fig. 5c bottom). Alternatively, the water-soluble 4-dmH upregulated Bax, Noxa,

Puma, and PARP cleavage; this suggested the possible relevance of a p53-dependent apoptotic

pathway in these cells (Fig. 5d). Noteworthily, the 4-dmH-induced downregulation of SIRT1 was

concurrent with an increase in c-Myc acetylation, leading to downregulation of anti-apoptotic c-Flip

in p53-mutated HSC-3 cells, suggesting the possible involvement of a p53-independent pathway

(Fig. 5d right). The absence of upregulation of ULK1, Atg 5, Atg7, and cleaved LC3-II provided

evidence for the inadequate autophagy by 4-dmH (Fig. 5d bottom).

Given that autophagy can either support survival or cause cell death, we analyzed the cellular

outcome of heliomycin-induced autophagy by cell impedance measurements. Our results

demonstrated that heliomycin decreased cell proliferation in both oral cancer cell lines, although the

induced cytotoxic effect was less evident at 1 μM in HSC-3 cells. However, heliomycin did not

exhibit cytotoxicity toward non-cancerous human BEAS-2B cells (Fig. 6a). The water-soluble

4-dmH also effectively diminished cell proliferation in a dose-dependent manner in oral cancer cells,

but not in that of BEAS-2B cells (Fig. 6b). The preferential cytotoxicity of 4-dmH toward cancer

cells was also reported in our previous study, indicating that 4-dmH displayed much higher IC_{50}

values against non-cancerous human dermal microvascular endothelium HMEC-1 cells compared to

those of tumor cells [26]. Additionally, our results from colony-forming assays revealed that both

compounds exhibited high growth-suppressive ability against oral cancer cells (Fig. 6c).

Nevertheless, the lack of changes in cell phase population indicated that the diminution in cell

growth by heliomycin and 4-dmH was least likely to arise from cell cycle arrest (Fig. 6d).

4-dmH, but not heliomycin, targets intracellular tNOX, an upstream regulator of SIRT1

We next addressed the molecular mechanisms underlying SIRT1 inhibition and concurrent cell

death by these two compounds in oral cancer cells. Being an NAD^+-dependent protein deacetylase,

SIRT1 activity is primarily governed by NAD^+/NADH ratio, thus, there exists a positive correlation

between these two [29-37]. We questioned whether these two compounds inhibit SIRT1 by

affecting the intracellular NAD^+/NADH levels, and were surprised to find that 4-dmH, but not

heliomycin, caused a prominent inhibition of intracellular NAD^+/NADH ratio (Fig. 7a). The

discrepancy in their ability to reduce NAD^+ generation led us to explore the role of a

tumor-associated NADH oxidase (tNOX, ENOX2) in 4-dmH-suppressed SIRT1 and apoptosis

induction. We have previously reported that tNOX inhibition reduced the intracellular

NAD^+/NADH ratio and SIRT1 deacetylase activity, increasing p53 acetylation and apoptosis

[38-41]. In the light of this information, we assessed the effect of the compounds on tNOX

expression and found that 4-dmH, but not heliomycin, considerably diminished tNOX protein

expression in a concentration-dependent manner (Fig. 7b). Pretreatment with the proteasome

inhibitor, MG-132, competently recovered the tNOX downregulation triggered by 1 or 2 μM

4-dmH in both cell lines, suggesting that the proteasomal degradation pathway may be associated

with 4-dmH-mediated tNOX downregulation (Fig. 7c). Consistently, 4-dmH increased the

ubiquitination level of tNOX, suggesting the possible involvement of ubiquitination- and

proteasome-driven protein degradation in the 4-dmH-induced downregulation of tNOX (Fig. 7d).

The RNAi-mediated knockdown of tNOX significantly triggered spontaneous autophagy and

apoptosis in SAS cells (Fig. 7e), further validating its role in the regulation of cell death.

Using CETSA, we explored whether tNOX could be a cellular target of 4-dmH. Our results

demonstrated that the heat stability of tNOX was found to be dose-dependently enhanced by 4-dmH

(Fig. 8a), indicating that there was direct binding between the protein and tested compound. The

4-dmH treatment increased the T_M value from 45.7 °C (control) to 50.9 °C (4-dmH-incubated) in

SAS cells, and from 47.7 °C (control) to 56.9°C (4-dmH-incubated) in HSC-3 cells (Fig. 8b),

further corroborating this direct interaction. In contrast, 4-dmH treatments did not seem to increase

the melting temperature of PARP or the nicotinamide adenine dinucleotide phosphate (NADPH)

oxidase-4 (NOX4), excluding those two proteins as potential targets of 4-dmH (Fig. 8c). Similarly,

our results from CETSA indicated that the parental heliomycin did not directly bind to tNOX in

either cell line (Fig. 8d).

Molecular docking simulation predicts better binding of heliomycin with SIRT1, and 4-dmH

with tNOX protein

To study the binding mode of 4-dmH and its parental heliomycin in the binding pockets of the

protein structures of SIRT1 and tNOX, we used molecular docking simulations. The binding poses

of heliomycin and 4-dmH were predicted to have similar orientations for SIRT1 (Fig. 9a). The

docking energy scores indicated that the heliomycin-SIRT1 complex (-10.1 kcal/mol) exhibited

higher affinity than the 4-dmH-SIRT1 complex (-8.0 kcal/mol). The heliomycin-SIRT1 and

4-dmH-SIRT1 complexes were mainly stabilized with hydrophobic interactions, with three

consistent H-bonds predicted in the binding pocket (red circles and ellipses, Fig. 9b). However,

although the 3-hydroxyl, 7-hydroxyl, and 2-carbonyl groups of heliomycin and 4-dmH consistently

formed hydrogen bonds with Ala262, Arg274, and Ile347, respectively, the 4-dmH-SIRT1 complex

exhibited an additional H-bond with Ser441 in the side 4-dimethylaminomethyl group (Fig. 9b).

The additional H-bond was predicted to decrease ligand affinity to SIRT1's binding cavity due to its

interference with the hydrophobic interactions.

Next, we performed a docking study of heliomycin and 4-dmH with tNOX. Previous reports

showed that doxorubicin (adriamycin) could inhibit tNOX activity [42, 43]. Therefore, we

simulated the binding poses of heliomycin, 4-dmH, and doxorubicin (as positive control) using a

blind docking approach. The results of our computational simulations showed that heliomycin,

4-dmH, and doxorubicin could bind in the same pocket of tNOX protein, but with different

orientations of their core moieties (Fig. 9c). The docking energy scores of the 4-dmH-tNOX and

doxorubicin-tNOX complexes were similar to one another (-8.5 and -8.4 kcal/mol, respectively),

while the heliomycin-tNOX complex had a slightly lower affinity (-8.1 kcal/mol). A number of

consistent interaction residues were found in the three docked complexes, including Ile90, Lys98,

Pro111, Pro113, Leu115, Pro117, and Pro118 (Fig. 9d). Both heliomycin and 4-dmH produced

three H-bonds, but the flipped orientation of 4-dmH yielded a higher number of hydrophobic

interactions than the heliomycin-tNOX complex. This suggests that the hydrophobic moieties

surrounding 4-dmH yield a higher affinity for tNOX protein.

Furthermore, for further evaluation of the importance of the consistent interaction residues in

the three docked compound-tNOX complexes, the seven interaction residues on tNOX were

substituted with alanine or glycine amino acids and then simulated the protein structures. The

simulated protein structures appear slightly different from the original tNOX structure. Overall, the

root mean square difference between the original tNOX structure and the structures with residues

substituted by alanine or glycine amino acids was estimated at 3.339 or 4.024 angstroms (Å),

respectively (Fig. S1a). The simulated protein structures were also employed to conduct the docking

analysis for 4-dmH. The results of further docking analysis revealed that 4-dmH could bind within

the same pocket of different types of tNOX structures but with varying orientations (Fig. S1b). This

observation also suggests that the replacement of both key residues with alanine or glycine could

result in a reduction of the binding affinity of 4-dmH to tNOX, with values of -8.2 and -7.6

kcal/mol, respectively. Moreover, the substitution of both key residues with alanine or glycine also

reduces the number of the original interacting residues and interaction forces in core moieties in the

4-dmH-tNOX complexes (Fig. S1c and S1d). Together, our experimental results and molecular

docking simulations are consistent with the notion that 4-dmH possesses a better affinity ability for

tNOX than for SIRT1.

4-dmH exhibits greater therapeutic efficacy in vivo

Given our observations indicating that SIRT1 and tNOX are cellular target(s) for heliomycin

or 4-dmH, we next investigated the therapeutic effects of these compounds *in vivo*. Mice were

inoculated with SAS cells, and tumor-bearing animals (average tumor diameter 0.5-1 cm) were

randomly split into three groups and intratumorally injected with buffer, 200 μg heliomycin in

water, or 200 μg 4-dmH in vehicle buffer. We found that 4-dmH treatment yielded a significant

reduction in tumor sizes at day 6, and continued to do so until animals were terminated at day 10

(Fig. 10a). By comparison, heliomycin treatment appeared to be less effective in diminishing tumor

growth compared to 4-dmH (Fig. 10a and b). Protein analysis of mouse tumor tissues further

showed that 4-dmH exhibited greater inhibition of tNOX expression compared to heliomycin, but

the two compounds were similar in their abilities to downregulate SIRT1 expression (Fig. 10c).

Interestingly, tNOX knockdown was reported to partially attenuate SIRT1 expression and repressed

growth in various cancer cell types, such as lung [44], bladder [45], and stomach [41]. We also

observed that tNOX is acetylated/ubiquitinated under certain stresses and the SIRT1 depletion

affects tNOX expression (data not shown). It is speculated that SIRT1 deacetylates tNOX and

modulates its protein stability. Thus, there is a reciprocal regulation between tNOX and SIRT1.

Although heliomycin does not bind to tNOX, its inhibition of SIRT1 activity/expression might have

an impact on tNOX expression. Consistent with our *in vitro* data, heliomycin enhanced the levels of

autophagy-associated markers, such as Atg 5 and cleaved LC3. 4-dmH treatment was associated *in*

vivo with upregulations of pro-apoptotic Bak and caspase 3-directed PARP cleavage, and the

concurrent downregulation of anti-apoptotic Flip (Fig. 10c), suggesting that the greater inhibition of

tumor volume by this water-soluble derivative of heliomycin was related to apoptosis induction.

Finally, we explored the potential clinical relevance of tNOX and SIRT1 protein in oral cancer.

Protein analysis of tissues from oral cancer patients indicated that both tNOX and SIRT1 were

upregulated in four out of five tumor samples compared to their adjacent normal tissues (Fig. 10d).

Data mining in the Kaplan-Meier plotter (pan-cancer RNA-seq dataset; www.kmplot.com),

revealed that high tNOX expression was associated with a poor prognosis for overall survival

[hazard ratio (HR): 2.81, log-rank $p = 0.008$] among 78 stage-III head-neck cancer patients. The

median overall survival in the low tNOX expression cohort was 213.9 months, compared to 28.7

months in the high tNOX expression cohort (Fig. 10e). However, we did not find any significant

correlation between tNOX and the survival rate in the earlier stages of the disease. The expression

of SIRT1 did not appear to significantly contribute to predicting overall survival at any stage of

head-neck cancer.

In sum, we herein show that in addition to targeting SIRT1, as does its parental heliomycin, the

enhanced water solubility and preferential targeting of tNOX by 4-dmH grants it superior

therapeutic value compared to heliomycin. Upon this binding, tNOX is ubiquitinylated and

degraded, which in turn attenuates signaling by the tNOX-NAD$^+$-SIRT1 regulatory axis and

induces apoptosis in oral cancer cells, as evidenced by the results of *in vitro* and *in vivo* studies.

Discussion

Oral cancers, a disease of epithelial origin, are commonly found in the lining of the lips, mouth, and upper throat. The updated data in GLOBOCAN 2020 estimated that there were nearly 0.38 million new cases and over 0.17 million deaths worldwide from cancers in the lip and oral cavity [46]. Unfortunately, the prognosis and 5-year overall survival of oral cancer patients remain disappointing, despite countless efforts to improve early diagnosis and expand treatment options [47-49]. In conjunction with surgery, radiotherapy, and chemotherapy, targeted therapy has emerged as a promising therapy for oral cancers, such as by targeting proteins that are overexpressed in cancer cells (e.g., EGFR and PD1) [50-52]. In this regard, we explored the possibility of the antibiotic heliomycin and a water-soluble heliomycin derivative as therapeutic strategies and identified their cellular targets. Here, we report that the water-soluble heliomycin derivative, 4-dmH exhibits a higher affinity for tNOX compared to its parental heliomycin, and that tNOX has a stronger affinity than SIRT1 for the 4-dmH. In addition to the action of 4-dmH on SIRT1, its preferential targeting of tNOX resulted in greater therapeutic efficacy in diminishing oral cancer cell proliferation and inhibiting tumor growth by inhibiting tNOX activity, increasing its protein degradation, and ultimately provoking apoptosis in p53-functional and p53-mutated oral cancer systems (Fig. 10f).

Heliomycin (also called resistomycin) was originally isolated from marine sponges and shown to possess a wide range of activities, including HDAC inhibitor activity [10]. HDAC inhibitors are currently under investigation for playing important roles in cancer epigenetic pathways, exhibiting

antitumor activity, and decreasing tumor resistance [53-55]. Our group has explored the anticancer

properties of heliomycin and established its binding with SIRT1 in the native cellular environment

of bladder cancer cells [25, 26]. SIRT1, which belongs to class III of the HDAC super-family, has

the unique feature of being NAD^+-dependent; it contributes to inflammatory oral diseases and oral

cancer by modulating many transcription factors and antioxidant enzymes [56]. In the present study,

we observed upregulation of SIRT1 in tumor tissues of animals and oral cancer patients (Fig. 10).

However, data mining using a pan-cancer RNA-seq dataset (www.kmplot.com) suggests that SIRT1

expression does not have significant importance in predicting overall survival at any stage of

head-neck cancer. A potential correlation between SIRT1 expression and prognosis for head-neck

cancer may warrant further discussion. Using molecular docking simulations and cell-based studies,

we herein demonstrated that both 4-dmH and heliomycin targeted SIRT1 and inhibited its activity

and expression, to attenuate the growth of oral cancer cells. Given that SIRT1 contributes to a

complex regulatory network, it is of primary importance to understand the fundamental regulatory

mechanisms of SIRT1 in cell death regulation [57-62].

Being an NAD^+-dependent protein deacetylase, it is not surprising that SIRT1 activity is

primarily governed by NAD^+/NADH ratio [29-37]. We further investigated the molecular events

underlying the prominent inhibition of intracellular NAD^+/NADH ratio by 4-dmH, but not

heliomycin. tNOX functions as a terminal hydroquinone oxidase, catalyzing not simply the

oxidation of the reduced form of NADH but also the transfer of protons and electrons to molecular

oxygen across the plasma membrane [63]. tNOX exhibits relatively low expression in

non-transformed cells, but is highly correlated with the hallmarks of cancer cells; positively

regulates cell proliferation, and its downregulation is associated with cell death-related pathways

[63-66]. In recent years, efforts to target tNOX with RNA interference or conventional anticancer

drugs have shown promising results in reducing cancer, not just in cell-based functional studies but

also in animals inoculated with cancer cells of colon cancer and melanoma [67, 68]. Specifically,

tNOX-depletion in cancer cells abolishes cancer phenotypes, reducing NAD^+ production,

proliferation, and migration/invasion while increasing apoptosis [44, 45, 65, 69]. Here, we report

consistent findings in oral cancer cells, validating the essential role of tNOX in cell death regulation

(Fig. 7e). Moreover, tNOX-overexpressing in non-cancerous cells stimulates the growth of cells,

decreases doubling time, and enhances cell migration [70-73]. We further substantiated the clinical

relevance of tNOX in oral cancer by showing that the majority of our oral cancer patients exhibit

higher tNOX protein expression in tumor tissues compared to their normal counterparts and there is

a reverse correlation between tNOX expression and overall survival in a Kaplan-Meier analysis (Fig.

10). Given that tNOX contributes to regulating SIRT1 activity, we propose that tNOX targeting is a

better operational strategy for reducing cancer growth in both *in vitro* and *in vivo* studies.

Another significant finding of this study is the cell thermal shift as analysis (CETSA)-based

results showing the direct binding capacities of heliomycin and 4-dmH toward SIRT1 and tNOX.

Moreover, 4-dmH and doxorubicin have very similar predicted affinities for tNOX protein. Based

on our previous report, we postulated that the NADH-binding motif of tNOX is located within the

range from Gly590 to Leu595 [74]. However, our simulation results indicated that the binding site

of the studied compounds is located well away from the NADH-binding motif of the tNOX protein.

The consistent interacting residues involved in the hydrogen bonds and hydrophobic interactions of

the studied compounds were predicted to be Ile90, Lys98, Pro111, Pro113, Leu115, Pro117, and

Pro118 (Fig. 9D), which are not located in the NADH-binding motif. At this point, we are uncertain

of the exact mechanism(s) by which these compounds inhibit tNOX. However, we speculate that

the hydrophobic moieties and extra hydrogen bonds formed between 4-dmH or doxorubicin and the

interacting residues of tNOX protein might affect its substrate binding. To clarify this issue, further

efforts involving the co-crystallization of tNOX/inhibitor complexes are warranted.

Taken together, our present results show that the superior anticancer activity of 4-dmH over its

parental heliomycin resides in its direct targeting of tNOX and SIRT1, and suggest that the

tNOX-NAD$^+$-SIRT1 regulatory axis contributes to inducing apoptosis independent of p53 status, as

evidenced by the results of *in vitro* and *in vivo* studies of oral cancer.

Conclusions

In conclusion, we focused on the identification and validation of the intracellular targets of

heliomycin and its water-soluble derivative, 4-dmH, by CETSA and molecular docking simulations

in p53-functional SAS and p53-mutated HSC-3 oral cancer cells. Our CETSA- and cell-based results

suggested that the targeting and inhibition of SIRT1 by heliomycin provoked autophagy and

attenuated the growth of oral cancer cells. In addition to targeting SIRT1, as does its parental

heliomycin, the enhanced water solubility of 4-dmH can maximize its anticancer effects and inhibit

cooperatively tNOX and SIRT1 to induce apoptosis in oral cancer cells independent of their p53

status. Considering that tNOX not only supports but also exerts functions independent of SIRT1, the

tNOX- and SIRT1-inhibiting function of 4-dmH, thus, results in the different biological outcomes

from the SIRT1-binding heliomycin, as evidenced by the results of *in vitro* and *in vivo* studies of

oral cancer.

Methods

Materials

The anti-SIRT1, anti-Atg5, anti-ULK1, anti-LC3, anti-PARP, anti-Bak, anti-Bax, anti-Puma,

anti-Noxa, anti-Bcl2, anti-c-Flip, anti-c-Myc, and anti-cleaved caspase-3 antibodies were purchased

from Cell Signaling Technology, Inc. (Beverly, MA, USA). The anti-β-actin and anti-acetyl-c-Myc

antibodies were from Millipore Corp. (Temecula, CA, USA). The anti-Atg7 antibody was obtained

from Novus Biologicals (Centennial, CO, USA). The antisera to tNOX used for immunoblotting

were generated as described previously [75]. The commercially available anti-ENOX2 antibody

from Proteintech (Rosemont, IL, USA) was used for immunoprecipitation. The anti-mouse and

anti-rabbit IgG antibodies and other chemicals were purchased from the Sigma Chemical Company

(St. Louis, MO, USA); unless otherwise specified.

Cell culture

SAS (human squamous cell carcinoma of the tongue), HSC-3 (human tongue squamous cell

carcinoma), and BEAS2B (human bronchial epithelial) cells grown in Dulbecco's Modified Eagle

Medium (DMEM), were kindly provided by Dr. Yuen-Chun Li (Department of Biomedical

Sciences, Chung Shan Medical University, Taiwan). Media were supplemented with 10% FBS, 100

units/ml penicillin, and 50 µg/ml streptomycin. Cells were maintained at 37°C in a humidified

atmosphere of 5% CO_2 in the air, and the media were replaced every 2-3 days. Cells were treated

with different concentrations of heliomycin (dissolved in DMSO) or the derivative 4-dmH

(dissolved in water).

ON-TARGETplus tNOX siRNA and negative control siRNA were purchased from Thermo

Scientific, Inc. (Grand Island, NY). Briefly, cells were seeded in 10-cm dishes, allowed to attach

overnight, and then transfected with tNOX siRNA or control siRNA using the Lipofectamine

RNAiMAX Reagent (Gibco/BRL Life Technologies) according to the manufacturer's instructions.

Chemistry

Heliomycin was produced at the Gause Institute of New Antibiotics (Moscow, Russian

Federation)   using   *Actynomyces*   *variabilis*   var.   *heliomycini*   [1].

4-(Dimethylaminomethyl)heliomycin (4-dmH) was prepared from heliomycin by the previously

reported method [26]. The structures of both compounds are illustrated in Fig. 1. The purities of the

tested compounds of heliomycin and the water-soluble derivative were 97 and 99%, respectively, as

examined by HPLC analysis.

Continuous observation of cell proliferation by cell impedance determination*s*

For continuous monitoring of changes in cell proliferation, cells (10^4 cells/well) were plated

onto E-plates, and the plates were incubated for 30 min at room temperature, and placed on an

xCELLigence System (Roche, Mannheim, Germany). The cells were grown overnight and then

exposed to heliomycin (dissolved in DMSO) or 4-dmH (dissolved in water), and impedance was

measured every hour, as previously described [44]. Cell impedance was defined by the cell index

(CI) = $(Z_i - Z_0)$ [Ohm]/15[Ohm], where Z_0 is background resistance and Z_i is the resistance at a

given time point. A normalized cell index was determined as the cell index at a given time point

(CI_{ti}) divided by the cell index at the normalization time point (CI_{nml_time}).

Measurement of mitochondrial membrane potential

Changes in mitochondrial membrane potential (a characteristic of apoptosis) were studied

using JC-1 staining (Theremo Fisher Scientific Inc., Waltham MA, USA). Cells were treated with

different concentrations of heliomycin or 4-dmH for 18 h. The cells were then washed, incubated

with 10 µM JC-1 at 37 °C for 30 min, washed with PBS, pelleted, and resuspended in PBS. The

fluorescence intensity of cells was analyzed using a Beckman Coulter FC500 flow cytometer (Brea,

CA, USA).

Autophagy determination

Autophagosomes, which are acidic intracellular compartments that mediate the degradation of

cytoplasmic materials during autophagy, were visualized by staining with acridine orange (AO;

Sigma Chemical Co.). Cells were washed with PBS, stained with 2 mg/ml AO for 10 min at 37°C,

washed, trypsinized, and analyzed using a Beckman Coulter FC500. The results are expressed as a

percentage of total cells.

Colony formation assay

Onto each 6-cm dish, two hundred cells were cultured in growth medium with various

concentrations of capsaicin at least 10 days. At the end of the experiment, colonies were fixed in

1.25% glutaraldehyde at room temperature for 30 minutes, washed with distilled water, and colored

with a 0.05% methylene blue mixture. The number of colonies was determined and documented.

Cell cycle analysis

In brief, after treatments, 10^6 cells were collected and washed in PBS, slowly fixed in 75%

ethanol, and kept at -20°C for at least 1 hour. The cell pellet was then washed again with PBS, and

centrifuged at 500 × g for 5 minutes. The pellet was resuspended in 200 µl cold PBS and nuclear

DNA was stained in the dark with propidium iodide (PI) solution (20 mM Tris pH 8.0, 1 mM NaCl,

0.1% NP-40, 1.4 mg/mL RNase A, 0.05 mg/mL PI) for 30 minutes on ice. Total cellular DNA

content was analyzed with an FC500 flow cytometer (Beckman Coulter Inc., Brea, CA, USA).

Cellular target identification by cellular thermal shift assay (CETSA)

CETSA was used to identify whether SIRT1 or tNOX are cellular targets of heliomycin and its

water-soluble derivative, 4-dmH. Samples were prepared from control and compound-exposed

cells. For each set, 2×10^7 cells were seeded in a 10-cm culture dish. After 24 h of culture, the cells

were pretreated with 10 µM MG132 for 1 h, washed with PBS, treated with trypsin, collected by

centrifugation at 12,000 rpm for 3 min at room temperature, gently resuspended with 1 ml of PBS,

pelleted by centrifugation at 7,500 rpm for 3 min at room temperature, and resuspended with 1 ml of

lysis buffer (20 mM Tris-HCl pH 7.4, 100 mM NaCl, 5mM EDTA, 2 mM phenylmethylsulfonyl

fluoride (PMSF), 10 ng/ml leupeptin, and 10 µg/ml aprotinin). The samples were transferred to

Eppendorf tubes and subjected to three freeze-thaw cycles; for each cycle, they were exposed to

liquid nitrogen for 3 min, placed in a heating block at 37°C for 3 min, and vortexed briefly. For the

experimental sample set, heliomycin or its derivative was added to a final concentration of 20 µM; the

same volume of vehicle solvent was added for the control sample set. The samples were heated at

37°C for 1 h and dispensed to 100 μl aliquots. Pairs consisting of one control aliquot and one

experimental aliquot were heated at 40°C, 43°C, 46°C, 49°C, 52°C, 55°C, 58°C, 61°C, or 67°C for 3

min. Insoluble proteins were separated by centrifugation at 12,000 rpm for 30 min at 4°C, and the

supernatants (containing soluble proteins) were resolved by SDS-PAGE. Western blot analysis was

performed using commercially available SIRT1 antibodies or tNOX antisera [65, 75]. β-Actin was

detected as a loading control.

An isothermal dose-response fingerprint (ITDRF$_{CETSA}$) was obtained in a manner similar to that

described above for the CETSA melting-curve experiments. Cells were seeded in 60-mm culture

dishes. After 24 h of culture, the cells were pretreated with 10 μM MG132, exposed to the test

compound for 1 h (final concentration, 0.001, 0.01, 0.1, 0.25, 0.5, 0.75, 1, 1.5, or 2 μM), washed with

PBS, treated with trypsin, pelleted at 12,000 rpm for 5 min at room temperature, gently resuspended

with 1 ml of PBS, pelleted at 7,500 rpm for 3 min at room temperature, and resuspended with the

same described above. The samples were subjected to three freeze-thaw cycles, wherein they were

exposed to liquid nitrogen for 3 min, placed in a heating block at 25°C for 3 min, and vortexed

briefly. The samples were then heated at 54°C for 3 min and cooled for 3 min at room temperature.

Insoluble proteins were separated by centrifugation at 12,000 rpm for 30 min at 4°C, and the

supernatants (containing soluble proteins) were subjected to SDS-PAGE and Western blot analysis

using an antibody against SIRT1 or antisera to tNOX. β-Actin was used as a loading control.

Western blot analysis and immunoprecipitation

Cell extracts were prepared in the lysis buffer described above. Volumes of extract containing

equal amounts of proteins (40 µg) were resolved by SDS-PAGE and transferred to PVDF

membranes (Schleicher & Schuell, Keene, NH, USA), and the membranes were blocked, washed,

and probed with the indicated primary antibody overnight. The membranes were washed, incubated

with horseradish peroxidase-conjugated secondary antibody for 1 h, and developed using enhanced

chemiluminescence (ECL) reagents (Amersham Biosciences, Piscataway, NJ, USA) according to

the manufacturer's protocol.

For immunoprecipitation, protein extracted from cells grown in 100-mm dishes were

pre-cleared with 20 µl of Protein G Agarose Beads (for rabbit antibodies) for 1 h at 4°C with

rotation. Ubiquitin antibodies or control IgG were incubated onto beads in 500 µl of lysis buffer,

overnight with rotation at 4°C. Beads were precipitated by centrifugation at 3,000 rpm for 2 min at

4°C. Beads were washed three times with lysis buffer and samples were prepared for Western blotting

analysis.

Determination of SIRT1 deacetylase activity in vitro

SIRT1 deacetylase activity was determined using a SIRT1 Direct Fluorescent Screening Assay

Kit (Cayman Chemical Company, Ann Arbor, MI) according to the manufacturer's protocol.

Briefly, substrate solution was prepared by adding 240 µl of NAD$^+$ solution and 850 µl of diluted

Assay Buffer to 15 µl of the p53 peptide Arg-His-Lys-Lys(εacetyl)-AMC, to yield a final

concentration of 125 µM peptide (substrate) and 3 mM of NAD$^+$. Using a 96-well plate, the

background was determined in wells containing 30 µl of Assay Buffer and 5 µl of DMSO. Maximal

initial activity (defined as 100%) was determined in wells containing 5 µl of diluted human

recombinant SIRT1, 25 µl of Assay Buffer, and 5 µl of DMSO. To measure the ability of the test

compound to modulate SIRT1 activity, 5 µl of 2 µM compound was added to wells containing 25

µl of Assay Buffer and 5 µl of diluted human recombinant SIRT1. Reactions were initiated by

adding 15 µl of Substrate Solution to each well. The plate was then covered and incubated on a

shaker for 45 min at room temperature. Reactions were stopped by adding 50 µl of Stop/Developing

Solution to each well and incubating the plate for 30 min at room temperature. Plates were read in a

fluorimeter using an excitation wavelength of 350-360 nm and an emission wavelength of 450-465

nm.

Measurement of the intracellular NAD$^+$/NADH ratio

The oxidized and reduced forms of intracellular NAD were determined using an NADH/NAD

Quantification Kit (BioVision Inc. Milpitas, CA, USA), as described by the manufacturer. Briefly, 2

$\times 10^5$ cells were washed with cold PBS, pelleted, and extracted by two freeze/thaw cycles with 400

µl of NADH/NAD$^+$ extraction buffer. The samples were vortexed and centrifuged at 14,000 rpm for

5 min. The extracted NADH/NAD$^+$-containing supernatant (200 µl) was transferred to a

microcentrifuge tube, heated to 60°C for 30 min (to decompose NAD$^+$ but not NADH), and then

placed on ice. The samples were then centrifuged and transferred to a multiwell plate. Standards

and an NAD$^+$ cycling mix were prepared according to the manufacturer's protocol. The reaction

mix was distributed at 100 µl/well to wells of a 96-well plate pre-loaded with NADH standards and

samples, and the plate was incubated at room temperature for 5 min to convert NAD$^+$ to NADH.

The provided NADH developer solution was added to each well, and plates were incubated at room

temperature for 15 or 30 min. The reaction was stopped with 10 µl of stop solution per well, and

absorbance was measured at 450 nm.

Molecular docking simulation

A crystal structure of the SIRT1 catalytic domain bound to an EX-527 analog (PDB: 4I5I) [76]

was employed for our molecular docking study. EX-527 is a nanomolar SIRT1 inhibitor with an

IC$_{50}$ value as low as 38 nM [77]; therefore its co-crystal structure of SIRT1 protein was judged to be

suitable for our docking analysis. The water molecule and ligand molecule of the initial crystal

structure were removed using the PyMOL program (https://pymol.org/; accessed on 1 November

2022; PyMOL Molecular Graphics System, Schrodinger, New York, NY) to prepare for the

docking analysis. A predicted full-length structure of the tNOX protein was obtained from the

AlphaFold Protein Structure Database [78, 79] and used in this study. Molecular docking was

performed using the AutoDock Vina package [80] in the PyRx software [81] to assess the probable

binding modes of heliomycin and its derivative in the SIRT1 catalytic domain and the protein

structure of tNOX. The docking site was determined according to the inhibitor binding site of the

EX-527 analog in the crystal structure of SIRT1, which was used as the setting in the grid selection.

Because we lacked information about the ligand-bound pocket, blind docking was performed by

adopting the whole-protein structure of tNOX, as the grid selection. Each compound was optimized

in its molecular geometry, torsional barriers, and intermolecular-interaction geometry using the

MMFF94 forcefield in CHARMM [82]. The best docking conformation of each compound, which

was chosen based on the lowest binding energy, was employed in the interaction analyses. The

interaction diagram was generated using the LigPlot+ software [83] to display the hydrogen bonds

and hydrophobic moieties. The docking pose of each compound was visualized using the PyMOL

program.

Homology modeling of tNOX protein structure

To evaluate the impact of key interacting residues derived from post-docking analysis, the key

residues on the tNOX protein were substituted with alanine or glycine amino acid, and then

conducted the protein structure simulation. The tNOX structure from the AlphaFold database was

used as template during these processes. The 3D structures of the modified tNOX protein sequences

were homology-modeled using the I-TASSER standalone package version 5.2 [84] (https://zhang

group.org/I-TASSER/download/) with default parameters. The simulated structures were also

applied in the molecular docking study to assess possible binding affinity and binding modes

between 4-dmH and the modified tNOX protein

In vivo xenograft studies

Specific pathogen-free (SPF) ASID mice were purchased from the National Laboratory

Animal Center (Taipei, Taiwan). The animal use protocols were approved by the Institutional

Animal Care and Use Committee of National Chung Hsing University (Taichung, Taiwan). Mice

were subcutaneously inoculated with 100 µl containing 2×10^6 live SAS cells in PBS. The

tumor-bearing animals were divided into three groups (n = 4 per group) when the tumor mass

reached an average diameter of 0.5-1 cm. Mice were intratumorally injected with vehicle buffer

(20% DMSO in PBS) as untreated control (Vehicle group), with 200 µg heliomycin in vehicle

buffer (Heliomycin group), with 200 µg 4-dmH in vehicle buffer (4-dmH group). Intratumoral

therapy was performed three times at 1-week intervals. The tumor size was recorded every 2 days

and the tumor volume was calculated using the formula: length $\times$ width^2 $\times$ 0.5. Mice were

euthanized 4 days after the final treatment. The significance of differences in tumor size was

determined by a one-way ANOVA.

Patient specimens

Five pairs of cancer tissues and their corresponding normal counterparts were obtained from

Changhua Christian Hospital in Taiwan (CCH IRB No. 130616). Diagnosis of oral cancer was

based on histological examination of hematoxylin and eosin-stained tissue section. Patient

specimens were immediately frozen in liquid nitrogen after surgery and used for Western blot

analysis

Statistics

All data are expressed as the means ± SEs of three independent experiments. Between-group

comparisons were performed using a one-way analysis of variance (ANOVA) followed by an

appropriate posthoc test. A value of $p<0.05$ was considered to be statistically significant.

Abbreviations

CETSA: Cellular thermal shift assay; ITDRF$_{CETSA}$: CETSA-based isothermal dose-response

fingerprint curves; MMP: Mitochondrial membrane potential; SIRT1: Sirtuin 1; tNOX:

tumor-associated NADH oxidase

Declarations

● Ethics Approval and Consent to Participate: The animal use protocols were approved by the

Institutional Animal Care and Use Committee of National Chung Hsing University. Clinical

specimens were obtained from the Changhua Christian Hospital in Taiwan (CCH IRB No.

130616).

● Consent for publication: Not applicable.

● Availability of data and supporting materials section: The datasets used and analyzed during

the present study are available from the corresponding author upon reasonable request.

● Competing interests: The authors declare that they have no competing interests.

● Funding: Financial support was provided by grants from the Ministry of Sciences and

Technology, Taiwan (MOST 106-2320-B-005-008-MY3 to PJC and MOST

108-2923-B-005-001-MY3 to PJC), the Russian Foundation for Basic Research (Project

19-53-52008 to AES), and the National Chung Hsing University and Changhua Christian

Hospital (NCHU-CCH 11105 to CWW).

● Authors' contributions: Conceived and designed the study: AST, AES, MKC, PJC. Performed

the experiments: AI, YCC, XCC, CYC, CWW, AST. Analyzed the data: AI, YCC, XCC,

CWW, AST, PJC. Wrote the manuscript: AI, YCC, CWW, AST, AES, PJC. All authors

reviewed and approved the final manuscript.

● Acknowledgments: The authors thank Dr. Yuen-Chun Li for providing cell lines.

References

1. Brazhnikova MG, Uspenskaia TA, Sokolova LB, Preobrazhenskaia TP, Gauze GF,
Ukholina RS, Shorin VA, Rossolimo OK, Vertogradova TP: **[New anti-viral antibiotic**
heliomycin]. *Antibiotiki (Mosc)* 1958, **3**(2):29-34.

2. Slesarchuk NA, Khvatov EV, Chistov AA, Proskurin GV, Nikitin TD, Lazarevich AI,
Ulanovskaya AA, Ulashchik EA, Orlov AA, Jegorov AV *et al*: **Simplistic perylene-related**
compounds as inhibitors of tick-borne encephalitis virus reproduction. *Bioorg Med*
Chem Lett 2020, **30**(10).

3. Zhang YL, Li S, Jiang DH, Kong LC, Zhang PH, Xu JD: **Antifungal activities of**
metabolites produced by a termite-associated Streptomyces canus BYB02. *J Agric Food*
Chem 2013, **61**(7):1521-1524.

4. Adinarayana G, Venkateshan MR, Bapiraju VV, Sujatha P, Premkumar J, Ellaiah P, Zeeck
A: **[Cytotoxic compounds from the marine actinobacterium]**. *Bioorg Khim* 2006,
32(3):328-334.

5. Arora SK: **Molecular-Structure of Heliomycin, an Inhibitor of Rna-Synthesis**. *J Antibiot*
1985, **38**(1):113-115.

6. Roggo BE, Petersen F, Delmendo R, Jenny HB, Peter HH, Roesel J:
3-Alkanoyl-5-hydroxymethyl tetronic acid homologues and resistomycin: new
inhibitors of HIV-1 protease. I. Fermentation, isolation and biological activity. *J*
Antibiot (Tokyo) 1994, **47**(2):136-142.

7. Vijayabharathi R, Bruheim P, Andreassen T, Raja DS, Devi PB, Sathyabama S,
Priyadarisini VB: **Assessment of resistomycin, as an anticancer compound isolated and**
characterized from Streptomyces aurantiacus AAA5. *J Microbiol* 2011, **49**(6):920-926.

8. Riaz A, Rasul A, Hussain G, Saadullah M, Rasool B, Sarfraz I, Masood M, Asrar M, Jabeen
F, Sultana T: **Resistomycin, a pentacyclic polyketide, inhibits the growth of triple**
negative breast cancer cells through induction of apoptosis and mitochondrial
dysfunction. *Pak J Pharm Sci* 2020, **33**(3):1233-1238.

9. Liu X, Arai MA, Toume K, Ishibashi M: **Isolation of Resistomycin from a Terrestrial**
Actinomycete with TRAIL Resistance-overcoming Activity. *Nat Prod Commun* 2018,
13(1):65-66.

10. Abdelfattah MS, Elmallah MIY, Faraag AHI, Hebishy AMS, Ali NH: **Heliomycin and**
tetracinomycin D: anthraquinone derivatives with histone deacetylase inhibitory
activity from marine sponge-associated Streptomyces sp SP9. *3 Biotech* 2018, **8**(6).

11. Xu R, Luo X, Ye X, Li H, Liu H, Du Q, Zhai Q: **SIRT1/PGC-1alpha/PPAR-gamma**
Correlate With Hypoxia-Induced Chemoresistance in Non-Small Cell Lung Cancer.
Front Oncol 2021, **11**:682762.

12. Garcia-Peterson LM, Li X: **Trending topics of SIRT1 in tumorigenicity**. *Biochim Biophys*
Acta Gen Subj 2021, **1865**(9):129952.

13. Wang L, Xu C, Johansen T, Berger SL, Dou Z: **SIRT1 - a new mammalian substrate of**
nuclear autophagy. *Autophagy* 2021, **17**(2):593-595.

14. Vaquero A, Sternglanz R, Reinberg D: **NAD(+)-dependent deacetylation of H4 lysine 16**
by class IIIHDACs. *Oncogene* 2007, **26**(37):5505-5520.

15. Houtkooper RH, Pirinen E, Auwerx J: **Sirtuins as regulators of metabolism and**
healthspan. *Nat Rev Mol Cell Bio* 2012, **13**(4):225-238.

16. Donmez G, Guarente L: **Aging and disease: connections to sirtuins**. *Aging cell* 2010,
9(2):285-290.

17. Song NY, Surh YJ: **Janus-faced role of SIRT1 in tumorigenesis**. *Ann N Y Acad Sci* 2012,
1271:10-19.

18. Choupani J, Mansoori Derakhshan S, Bayat S, Alivand MR, Shekari Khaniani M:
Narrower insight to SIRT1 role in cancer: A potential therapeutic target to control
epithelial-mesenchymal transition in cancer cells. *J Cell Physiol* 2018, **233**(6):4443-4457.

19. Lee IH, Cao L, Mostoslavsky R, Lombard DB, Liu J, Bruns NE, Tsokos M, Alt FW, Finkel
T: **A role for the NAD-dependent deacetylase Sirt1 in the regulation of autophagy**. *Proc*
Natl Acad Sci U S A 2008, **105**(9):3374-3379.

20. Sun T, Hu Y, He W, Shang Y, Yang X, Gong L, Zhang X, Gong P, Yang G: **SRT2183**
impairs ovarian cancer by facilitating autophagy. *Aging (Albany NY)* 2020,
12(23):24208-24218.

21. Karbasforooshan H, Roohbakhsh A, Karimi G: **SIRT1 and microRNAs: The role in**
breast, lung and prostate cancers. *Experimental cell research* 2018, **367**(1):1-6.

22. Yousafzai NA, Jin H, Ullah M, Wang X: **Recent advances of SIRT1 and implications in**
chemotherapeutics resistance in cancer. *Am J Cancer Res* 2021, **11**(11):5233-5248.

23. Martinez Molina D, Jafari R, Ignatushchenko M, Seki T, Larsson EA, Dan C, Sreekumar L,
Cao Y, Nordlund P: **Monitoring drug target engagement in cells and tissues using the**
cellular thermal shift assay. *Science* 2013, **341**(6141):84-87.

24. Martinez Molina D, Nordlund P: **The Cellular Thermal Shift Assay: A Novel Biophysical**
Assay for In Situ Drug Target Engagement and Mechanistic Biomarker Studies. *Annu*
Rev Pharmacol Toxicol 2016, **56**:141-161.

25. Lin MH, Islam A, Liu YH, Weng CW, Zhan JH, Liang RH, Tikhomirov AS, Shchekotikhin
AE, Chueh PJ: **Antibiotic heliomycin and its water-soluble 4-aminomethylated**
derivative provoke cell death in T24 bladder cancer cells by targeting sirtuin 1 (SIRT1).
Am J Cancer Res 2022, **12**(3):1042-1055.

26. Nadysev GY, Tikhomirov AS, Lin MH, Yang YT, Dezhenkova LG, Chen HY, Kaluzhny
DN, Schols D, Shtil AA, Shchekotikhin AE *et al*: **Aminomethylation of heliomycin:**
Preparation and anticancer characterization of the first series of semi-synthetic
derivatives. *European journal of medicinal chemistry* 2018, **143**:1553-1562.

27. Tikhomirov AS, Abdelhamid MAS, Nadysev GY, Zatonsky GV, Bykov EE, Chueh PJ,

Waller ZAE, Shchekotikhin AE: **Water-Soluble Heliomycin Derivatives to Target**
i-Motif DNA. *Journal of Natural Products* 2021, **84**(5):1617-1625.

28.　Mai A, Massa S, Lavu S, Pezzi R, Simeoni S, Ragno R, Mariotti FR, Chiani F, Camilloni G,
Sinclair DA: **Design, synthesis, and biological evaluation of sirtinol analogues as class**
III histone/protein deacetylase (sirtuin) inhibitors. *Journal of medicinal chemistry* 2005,
48(24):7789-7795.

29.　Mouchiroud L, Houtkooper RH, Moullan N, Katsyuba E, Ryu D, Canto C, Mottis A, Jo YS,
Viswanathan M, Schoonjans K *et al*: **The NAD(+)/Sirtuin Pathway Modulates Longevity**
through Activation of Mitochondrial UPR and FOXO Signaling. *Cell* 2013,
154(2):430-441.

30.　He S, Gao Q, Wu X, Shi J, Zhang Y, Yang J, Li X, Du S, Zhang Y, Yu J: **NAD(+)**
ameliorates endotoxin-induced acute kidney injury in a sirtuin1-dependent manner via
GSK-3beta/Nrf2 signalling pathway. *J Cell Mol Med* 2022, **26**(7):1979-1993.

31.　Donmez G: **The neurobiology of sirtuins and their role in neurodegeneration**. *Trends*
Pharmacol Sci 2012, **33**(9):494-501.

32.　Teertam SK, Phanithi PB: **Up-regulation of Sirtuin-1/autophagy signaling in human**
cerebral ischemia: possible role in caspase-3 mediated apoptosis. *Heliyon* 2022,
8(12):e12278.

33.　Li BY, Peng WQ, Liu Y, Guo L, Tang QQ: **HIGD1A links SIRT1 activity to adipose**
browning by inhibiting the ROS/DNA damage pathway. *Cell reports* 2023,
42(7):112731.

34.　Bai P, Canto C, Oudart H, Brunyanszki A, Cen Y, Thomas C, Yamamoto H, Huber A, Kiss
B, Houtkooper RH *et al*: **PARP-1 inhibition increases mitochondrial metabolism**
through SIRT1 activation. *Cell Metab* 2011, **13**(4):461-468.

35.　Ma Y, Nie H, Chen H, Li J, Hong Y, Wang B, Wang C, Zhang J, Cao W, Zhang M *et al*:
NAD(+)/NADH metabolism and NAD(+)-dependent enzymes in cell death and
ischemic brain injury: current advances and therapeutic implications. *Curr Med Chem*
2015, **22**(10):1239-1247.

36.　Fulco M, Schiltz RL, Iezzi S, King MT, Zhao P, Kashiwaya Y, Hoffman E, Veech RL,
Sartorelli V: **Sir2 regulates skeletal muscle differentiation as a potential sensor of the**
redox state. *Mol Cell* 2003, **12**(1):51-62.

37.　Yang Y, Liu Y, Wang Y, Chao Y, Zhang J, Jia Y, Tie J, Hu D: **Regulation of SIRT1 and**
Its Roles in Inflammation. *Front Immunol* 2022, **13**:831168.

38.　Tikhomirov AS, Shchekotikhin AE, Lee YH, Chen YA, Yeh CA, Tatarskiy VV, Jr.,
Dezhenkova LG, Glazunova VA, Balzarini J, Shtil AA *et al*: **Synthesis and**
Characterization of 4,11-Diaminoanthra[2,3-b]furan-5,10-diones: Tumor Cell
Apoptosis through tNOX-Modulated NAD(+)/NADH Ratio and SIRT1. *Journal of*
medicinal chemistry 2015, **58**(24):9522-9534.

39.   Chang CF, Islam A, Liu PF, Zhan JH, Chueh PJ: **Capsaicin acts through tNOX (ENOX2)**
to induce autophagic apoptosis in p53-mutated HSC-3 cells but autophagy in
p53-functional SAS oral cancer cells. *Am J Cancer Res* 2020, **10**(10):3230-3247.

40.   Lin CY, Islam A, Su CJ, Tikhomirov AS, Shchekotikhin AE, Chuang SM, Chueh PJ, Chen
YL: **Engagement with tNOX (ENOX2) to Inhibit SIRT1 and Activate p53-Dependent**
and -Independent Apoptotic Pathways by Novel
4,11-Diaminoanthra[2,3-b]furan-5,10-diones in Hepatocellular Carcinoma Cells.
Cancers (Basel) 2019, **11**(3).

41.   Chen HY, Cheng HL, Lee YH, Yuan TM, Chen SW, Lin YY, Chueh PJ: **Tumor-associated**
NADH oxidase (tNOX)-NAD+-sirtuin 1 axis contributes to oxaliplatin-induced
apoptosis of gastric cancer cells. *Oncotarget* 2017, **8**(9):15338-15348.

42.   Morre DJ, Kim C, Paulik M, Morre DM, Faulk WP: **Is the drug-responsive NADH**
oxidase of the cancer cell plasma membrane a molecular target for adriamycin? *J*
Bioenerg Biomembr 1997, **29**(3):269-280.

43.   Hedges KL, Morre DM, Wu LY, Morre DJ: **Adriamycin tolerance in human**
mesothelioma lines and cell surface NADH oxidase. *Life Sci* 2003, **73**(9):1189-1198.

44.   Lee YH, Chen HY, Su LJ, Chueh PJ: **Sirtuin 1 (SIRT1) Deacetylase Activity and**
NAD(+)/NADH Ratio Are Imperative for Capsaicin-Mediated Programmed Cell Death.
J Agric Food Chem 2015, **63**(33):7361-7370.

45.   Lin MH, Lee YH, Cheng HL, Chen HY, Jhuang FH, Chueh PJ: **Capsaicin Inhibits**
Multiple Bladder Cancer Cell Phenotypes by Inhibiting Tumor-Associated NADH
Oxidase (tNOX) and Sirtuin1 (SIRT1). *Molecules* 2016, **21**(7).

46.   Sung H, Ferlay J, Siegel RL, Laversanne M, Soerjomataram I, Jemal A, Bray F: **Global**
cancer statistics 2020: GLOBOCAN estimates of incidence and mortality worldwide
for 36 cancers in 185 countries. *Ca-Cancer J Clin* 2021, **71**(3):209-249.

47.   Sievilainen M, Almahmoudi R, Al-Samadi A, Salo T, Pirinen M, Almangush A: **The**
prognostic value of immune checkpoints in oral squamous cell carcinoma. *Oral Dis*
2019, **25**(6):1435-1445.

48.   Lacas B, Bourhis J, Overgaard J, Zhang Q, Gregoire V, Nankivell M, Zackrisson B,
Szutkowski Z, Suwinski R, Poulsen M *et al*: **Role of radiotherapy fractionation in head**
and neck cancers (MARCH): an updated meta-analysis. *Lancet Oncol* 2017,
18(9):1221-1237.

49.   Carvalho AL, Nishimoto IN, Califano JA, Kowalski LP: **Trends in incidence and**
prognosis for head and neck cancer in the United States: a site-specific analysis of the
SEER database. *Int J Cancer* 2005, **114**(5):806-816.

50.   Li H, Zhang Y, Xu M, Yang D: **Current trends of targeted therapy for oral squamous**
cell carcinoma. *J Cancer Res Clin Oncol* 2022, **148**(9):2169-2186.

51.   Cheng YX, Chen J, Shi YX, Fang XD, Tang ZG: **MAPK Signaling Pathway in Oral**

Squamous Cell Carcinoma: Biological Function and Targeted Therapy. *Cancers* 2022,
14(19).

52. Cao M, Shi E, Wang H, Mao L, Wu Q, Li X, Liang Y, Yang X, Wang Y, Li C:
Personalized Targeted Therapeutic Strategies against Oral Squamous Cell Carcinoma.
An Evidence-Based Review of Literature. *Int J Nanomedicine* 2022, **17**:4293-4306.

53. Basha NJ, Basavarajaiah SM: **An insight into therapeutic efficacy of heterocycles as**
histone-modifying enzyme inhibitors that targets cancer epigenetic pathways. *Chem*
Biol Drug Des 2022, **100**(5):682-698.

54. Sun Y, Hong JH, Ning Z, Pan D, Fu X, Lu X, Tan J: **Therapeutic potential of tucidinostat,**
a subtype-selective HDAC inhibitor, in cancer treatment. *Front Pharmacol* 2022,
13:932914.

55. Contreras-Sanzon E, Prado-Garcia H, Romero-Garcia S, Nunez-Corona D, Ortiz-Quintero B,
Luna-Rivero C, Martinez-Cruz V, Carlos-Reyes A: **Histone deacetylases modulate**
resistance to the therapy in lung cancer. *Front Genet* 2022, **13**:960263.

56. Pan ZJ, Dong H, Huang N, Fang J: **Oxidative stress and inflammation regulation of**
sirtuins: New insights into common oral diseases. *Front Physiol* 2022, **13**.

57. Wang Y, Sui Y, Niu Y, Liu D, Xu Q, Liu F, Zuo K, Liu M, Sun W, Wang Z *et al*:
PBX1-SIRT1 Positive Feedback Loop Attenuates ROS-Mediated HF-MSC Senescence
and Apoptosis. *Stem Cell Rev Rep* 2023, **19**(2):443-454.

58. Wang X, Lu Y, Tuo Z, Zhou H, Zhang Y, Cao Z, Peng L, Yu D, Bi L: **Role of**
SIRT1/AMPK signaling in the proliferation, migration and invasion of renal cell
carcinoma cells. *Oncol Rep* 2021, **45**(6).

59. Lee SC, Kim KH, Kim OH, Lee SK, Hong HE, Choi BJ, Jeong W, Kim SJ: **Everolimus**
Plus Ku0063794 Regimen Promotes Anticancer Effects against Hepatocellular
Carcinoma Cells through the Paradoxical Inhibition of Autophagy. *Cancer Res Treat*
2018, **50**(3):1023-1038.

60. Zhu L, Duan W, Wu GJ, Zhang D, Wang L, Chen D, Chen ZS, Yang B: **Protective effect of**
hydrogen sulfide on endothelial cells through Sirt1-FoxO1-mediated autophagy. *Ann*
Transl Med 2020, **8**(23).

61. Ran D, Zhou D, Liu G, Ma Y, Ali W, Yu R, Wang Q, Zhao H, Zhu J, Zou H *et al*: **Reactive**
Oxygen Species Control Osteoblast Apoptosis through SIRT1/PGC-1alpha/P53(Lys382)
Signaling, Mediating the Onset of Cd-Induced Osteoporosis. *J Agric Food Chem* 2023.

62. Zhang Z, Chen X, Liu S: **Role of Sirtuin-1 in Neonatal Hypoxic-Ischemic**
Encephalopathy and Its Underlying Mechanism. *Med Sci Monit* 2020, **26**:e924544.

63. Hostetler B, Weston N, Kim C, Morré DM, Morré DJ: **Cancer Site-Specific Isoforms of**
ENOX2 (tNOX), A Cancer-Specific Cell Surface Oxidase. *Clinical Proteomics* 2009,
5(1):46-51.

64. Sumiyoshi A, Shibata S, Zhelev Z, Miller T, Lazarova D, Aoki I, Obata T, Higashi T,

Bakalova R: **Targeting Glioblastoma via Selective Alteration of Mitochondrial Redox**
State. *Cancers (Basel)* 2022, **14**(3).

65. Liu SC, Yang JJ, Shao KN, Chueh PJ: **RNA interference targeting tNOX attenuates cell**
migration via a mechanism that involves membrane association of Rac. *Biochem*
Biophys Res Commun 2008, **365**(4):672-677.

66. Ronconi G, Lessiani G, Spinas E, Kritas SK, Caraffa A, Saggini A, Antinolfi P,
Pizzicannella J, Toniato E, Conti P: **ENOX2 (or tNOX): a new and old molecule with**
cancer activity involved in tumor prevention and therapy. *J Biol Regul Homeost Agents*
2016, **30**(3):649-653.

67. Liu NC, Hsieh PF, Hsieh MK, Zeng ZM, Cheng HL, Liao JW, Chueh PJ:
Capsaicin-mediated tNOX (ENOX2) up-regulation enhances cell proliferation and
migration in vitro and in vivo. *J Agric Food Chem* 2012, **60**(10):2758-2765.

68. Islam A, Hsieh PF, Liu PF, Chou JC, Liao JW, Hsieh MK, Chueh PJ: **Capsaicin exerts**
therapeutic effects by targeting tNOX-SIRT1 axis and augmenting ROS-dependent
autophagy in melanoma cancer cells. *Am J Cancer Res* 2021, **11**(9):4199-4219.

69. Cheng HL, Lee YH, Yuan TM, Chen SW, Chueh PJ: **Update on a tumor-associated**
NADH oxidase in gastric cancer cell growth. *World J Gastroenterol* 2016,
22(10):2900-2905.

70. Islam A, Su AJ, Zeng ZM, Chueh PJ, Lin MH: **Capsaicin Targets tNOX (ENOX2) to**
Inhibit G1 Cyclin/CDK Complex, as Assessed by the Cellular Thermal Shift Assay
(CETSA). *Cells* 2019, **8**(10).

71. Su YC, Lin YH, Zeng ZM, Shao KN, Chueh PJ: **Chemotherapeutic agents enhance cell**
migration and epithelial-to-mesenchymal transition through transient up-regulation of
tNOX (ENOX2) protein. *Biochim Biophys Acta* 2012, **1820**(11):1744-1752.

72. Zeng ZM, Chuang SM, Chang TC, Hong CW, Chou JC, Yang JJ, Chueh PJ:
Phosphorylation of serine-504 of tNOX (ENOX2) modulates cell proliferation and
migration in cancer cells. *Experimental cell research* 2012, **318**(14):1759-1766.

73. Chueh PJ, Wu LY, Morre DM, Morre DJ: **tNOX is both necessary and sufficient as a**
cellular target for the anticancer actions of capsaicin and the green tea catechin
(-)-epigallocatechin-3-gallate. *Biofactors* 2004, **20**(4):235-249.

74. Chueh PJ, Morré DM, Morré DJ: **A site-directed mutagenesis analysis of tNOX**
functional domains. *Biochim Biophys Acta* 2002, **1594**(1):74-83.

75. Chen CF, Huang S, Liu SC, Chueh PJ: **Effect of polyclonal antisera to recombinant**
tNOX protein on the growth of transformed cells. *Biofactors* 2006, **28**(2):119-133.

76. Zhao X, Allison D, Condon B, Zhang F, Gheyi T, Zhang A, Ashok S, Russell M, MacEwan
I, Qian Y *et al*: **The 2.5 A crystal structure of the SIRT1 catalytic domain bound to**
nicotinamide adenine dinucleotide (NAD+) and an indole (EX527 analogue) reveals a
novel mechanism of histone deacetylase inhibition. *Journal of medicinal chemistry* 2013,

56(3):963-969.

77.   Solomon JM, Pasupuleti R, Xu L, McDonagh T, Curtis R, DiStefano PS, Huber LJ:
829   **Inhibition of SIRT1 catalytic activity increases p53 acetylation but does not alter cell**
830   **survival following DNA damage**. *Mol Cell Biol* 2006, **26**(1):28-38.

831   78.   Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool K,
832   Bates R, Zidek A, Potapenko A *et al*: **Highly accurate protein structure prediction with**
833   **AlphaFold**. *Nature* 2021, **596**(7873):583-589.

834   79.   Varadi M, Anyango S, Deshpande M, Nair S, Natassia C, Yordanova G, Yuan D, Stroe O,
835   Wood G, Laydon A *et al*: **AlphaFold Protein Structure Database: massively expanding**
836   **the structural coverage of protein-sequence space with high-accuracy models**. *Nucleic*
837   *Acids Res* 2022, **50**(D1):D439-D444.

838   80.   Trott O, Olson AJ: **AutoDock Vina: improving the speed and accuracy of docking with**
839   **a new scoring function, efficient optimization, and multithreading**. *J Comput Chem*
840   2010, **31**(2):455-461.

841   81.   Dallakyan S, Olson AJ: **Small-molecule library screening by docking with PyRx**.
842   *Methods Mol Biol* 2015, **1263**:243-250.

843   82.   Brooks BR, Brooks CL, 3rd, Mackerell AD, Jr., Nilsson L, Petrella RJ, Roux B, Won Y,
844   Archontis G, Bartels C, Boresch S *et al*: **CHARMM: the biomolecular simulation**
845   **program**. *J Comput Chem* 2009, **30**(10):1545-1614.

846   83.   Laskowski RA, Swindells MB: **LigPlot+: multiple ligand-protein interaction diagrams**
847   **for drug discovery**. *J Chem Inf Model* 2011, **51**(10):2778-2786.

848   84.   Yang J, Yan R, Roy A, Xu D, Poisson J, Zhang Y: **The I-TASSER Suite: protein**
849   **structure and function prediction**. *Nat Methods* 2015, **12**(1):7-8.

 Figure Legends

Figure 1

Structures of heliomycin and 4-(dimethylaminomethyl)heliomycin (designated 4-dmH)

Figure 2

CETSA-based determination of binding between 4-dmH and SIRT1 protein. (a) Cells were

incubated with different concentrations of 4-dmH as described in the Methods. Dose-dependent

thermal stabilization of SIRT1 was assessed after heating samples at 54 °C for 3 min in SAS cells and

HSC-3 cells. The band intensities of SIRT1 were normalized with respect to the intensity of actin.

Representative images are shown. (b) CETSA-melting curves of SIRT1 in the presence and absence

of 4-dmH in SAS cells and HSC-3 cells as described in the Methods. The immunoblot intensity was

normalized to the intensity of the 40 °C samples. Representative images are shown. The

denaturation midpoints were determined using a standard process.

Figure 3

CETSA-based determination of binding between heliomycin and SIRT1 protein. (a) Cells were

incubated with different concentrations of heliomycin as described in the Methods. Dose-dependent

thermal stabilization of SIRT1 was assessed after heating samples at 54 °C for 3 min in SAS cells and

HSC-3 cells. The band intensities of SIRT1 were normalized with respect to the intensity of actin.

Representative images are shown. (b) CETSA-melting curves of SIRT1 in the presence and absence

of heliomycin in SAS cells and HSC-3 cells as described in the Methods. The immunoblot intensity

was normalized to the intensity of the 40 °C samples. Representative images are shown. The

denaturation midpoints were determined using a standard process.

Figure 4

SIRT1 inhibition and downregulation by water-soluble 4-dmH. (a) SIRT1 activity was determined

by a SIRT1 Activity Assay Kit (Fluorometric) with control or recombinant protein treated with test

compounds. Values (mean ± S. E.) are from three independent experiments (*p <0.05, ***p

<0.001). (b) 4-dmH markedly attenuated SIRT1 protein expression analyzed by Western blotting. (c)

4-dmH-suppressed SIRT1 expression was reverted by MG132, the proteasome inhibitor, in SAS

cells and HSC-3 cells. Aliquots of cell lysates were resolved by SDS-PAGE and analyzed by

Western blotting. β-Actin was detected as an internal control. Representative images are shown.

Values (means ± SDs) are from at least three independent experiments (*$p < 0.05$, **$p < 0.01$, ***

$p < 0.001$). (d) The lysates of HA-Ub overexpressing cells were immunoprecipitated with

nonimmune IgG or an antibody against SIRT1, and the bound proteins were detected by Western

blotting with anti-HA or anti-SIRT1 antibodies. The total lysates were also immunoblotted with

anti-SIRT1 or anti-β-actin antibodies. Aliquots of cell lysates were resolved by SDS-PAGE and

analyzed by Western blotting. β-actin was detected as an internal control. Representative images are

shown.

Figure 5

Heliomycin and 4-dmH provoked different cell death pathways in oral cancer cells. (a) Cells were

exposed to heliomycin or 4-dmH and the percentage of the autophagic population was determined

by AO staining using flow cytometry. The results are expressed as a percentage of autophagic cells.

Values (mean ± S. E.) are from at least three independent experiments (*p <0.05, **p <0.01). (b)

Cells were exposed to heliomycin or 4-dmH and the percentage of the apoptotic population was

determined by JC-1 staining using flow cytometry. The results are expressed as a percentage of

autophagic cells. Values (mean ± S. E.) are from at least three independent experiments (*p <0.05).

(c, d) Cells were treated with heliomycin for various concentrations, and aliquots of cell lysates

were resolved by SDS-PAGE and analyzed for protein expression by Western blotting. β-actin was

used as an internal loading control to monitor for equal loading. Representative images are provided

from at least three independent experiments.

Figure 6

The effects of heliomycin and 4-dmH on proliferation, colony formation, and cell cycle progression

of oral cancer cells. (a, b) Cell proliferation was dynamically monitored by impedance

measurements in SAS, HSC-3 cells, and BSEA-2B cells as described in the Methods. Shown are the

Figure 7

tNOX inhibition and downregulation by water-soluble 4-dmH. (a) The intracellular NAD^+/NADH

ratio was measured by an NADH/NAD Quantification Kit (BioVision Inc.) with control or lysates

of SAS cells treated with heliomycin or 4-dmH. Values (mean ± S. E.) are from three independent

experiments (*p <0.05). (b) 4-dmH, but not heliomycin, markedly attenuated tNOX protein

expression analyzed by Western blotting. (c) 4-dmH-suppressed tNOX expression was reversed by

MG-132 in SAS cells and HSC-3 cells. Aliquots of cell lysates were resolved by SDS-PAGE and

analyzed by Western blotting. β-Actin was detected as an internal control. Representative images

are shown. Values (means ± SDs) are from at least three independent experiments (*p < 0.05, **p <

0.01, *** p < 0.001). (d) The lysates of HA-Ub overexpressing cells were immunoprecipitated with

nonimmune IgG or an antibody against ENOX2, and the bound proteins were detected by Western

blotting with anti-HA or anti-ENOX2 antibodies. The total lysates were also immunoblotted with

anti-tNOX or anti-β-actin antibodies. (e) The RNA interference-mediated tNOX depletion was

conducted in SAS cells for 48 hours. The percentage of apoptotic/autophagic cells was examined by

flow cytometry. The presented values (mean ± SDs) represent at least three independent

experiments (** p <0.01 for tNOX-depleted cells vs. controls).

Figure 8

CETSA-based determination of binding between 4-dmH and tNOX protein. (a) Cells were

incubated with different concentrations of the 4-dmH as described in the Methods. Dose-dependent

thermal stabilization of tNOX was assessed after heating samples at 54 °C for 3 min in SAS cells and

HSC-3 cells. The band intensities of tNOX were normalized with respect to the intensity of actin.

Representative images are shown. (b) CETSA-melting curves of tNOX in the presence and absence

of 4-dmH in SAS cells and HSC-3 cells as described in the Methods. The immunoblot intensity was

normalized to the intensity of the 40 °C samples. Representative images are shown. The

denaturation midpoints were determined using a standard process. (c) CETSA-melting curves of

PARP or NOX4 in the presence and absence of 4-dmH in SAS cells as described in the Methods.

The immunoblot intensity was normalized to the intensity of the 40 °C samples. Representative

images are shown. The denaturation midpoints were determined using a standard process. (d)

CETSA-melting curves of tNOX in the presence and absence of heliomycin in SAS cells and

HSC-3 cells as described in the Methods. The immunoblot intensity was normalized to the intensity

of the 40 °C samples. Representative images are shown. The denaturation midpoints were

determined using a standard process.

Figure 9

The binding modes of heliomycin and 4-dmH after docking into the pocket of SIRT1 (a, b) and

tNOX (c, d). (a) Superimposition of the docked heliomycin (red) and 4-dmH (blue). (b) Schematic

presentations of possible interactions between test compounds and SIRT1 residues. (c)

Superimposition of the docked heliomycin (red) and 4-dmH (blue), and doxorubicin (purple). (d)

Schematic presentations of possible interactions between test compounds and tNOX residues. The

key residues surrounding the binding pocket of SIRT1 and tNOX were identified via the best

docking pose of each compound. The red circles and ellipses indicate the identical residues that

interacted with different compounds.

Figure 10

Therapeutic potential of heliomycin and 4-dmH in oral cancer. (a-c) In a tumor-bearing mouse

xenograft model, control mice were intratumorally injected with vehicle buffer and treatment group

mice were intratumorally treated with heliomycin or 4-dmH as described in the Methods. The

morphology of the tumor tissues excised from tumor-bearing mice (a) and quantitative analysis of

xenografted tumor volume during the treatment period (b) are shown. (c) Tissues from three sets of

tumor-bearing mice were grounded and prepared for Western blotting analysis. (d) The tumor and

adjacent tissues from oral cancer patients were grounded and prepared for Western blotting analysis.

β-actin was used as an internal loading control to monitor for equal loading. (e) Kaplan-Meier plots

of the association between tNOX (ENOX2) expression and overall survival in 78 stage-III

head-neck cancer patients. (f) Schematic diagram of the mechanism governing 4-dmH-induced

apoptosis and heliomycin-mediated autophagy in oral cancer cells.

Heliomycin

4-dmH

a

b

Tumor volume (cm3)

Vehicle
Heliomycin
4-dmH

Day 0 Day 2 Day 4 Day 6 Day 8 Day 10

c

	Vehicle	Heliomycin	4-dmH
	1 2 3	1 2 3	1 2 3

tNOX
SIRT1
Atg 5
Atg 7
LC3-I
LC3-II
Bak
c-Flip
PARP
Actin

d

Tongue   Buccal

18N 18T 35N 35T   57N 57T 67N 67T 75N 75T

tNOX
SIRT1
Actin

e

ENOX2

HR = 2.81 (1.27 - 6.24)
logrank P = 0.008

Probability

Expression
low
high

Time (months)

Number at risk
low   39   9   2   1   1
high   39   5   1   1   0

f

4-dmH

tNOX → NADH
         NAD+

SIRT1

↑ Bax/Bak   ↑ c-Myc
↑ Noxa   acetylation
↑ Puma   ↓ c-Myc
↓ c-Flip   ↓ c-Flip

Apoptosis   Apoptosis
p53 functional   p53 mutated
SAS cells   HSC-3 cells

Heliomycin

SIRT1

↑ ULK1
↑ Atg5
↑ Atg7
↑ cleaved LC3
↓ Bcl2

Autophagy
SAS and HSC-3 cells