A novel interaction between the 5' untranslated region of the

Chikungunya virus genome and Musashi RNA binding protein

is essential for efficient virus genome replication

Authors: Kaiwen Sun, Francesca Appadoo, Yuqian Liu, Marietta Müller, Catriona

Macfarlane, Mark Harris and Andrew Tuplin*

Correspondence: a.k.tuplin@leeds.ac.uk

Affiliations: Astbury Centre for Structural Molecular Biology and School of

Molecular and Cellular Biology, Faculty of Biological Sciences, University of Leeds,

Leeds LS2 9JT, UK.

Abstract

Chikungunya virus (CHIKV) is a re-emerging, pathogenic alphavirus that is transmitted to humans by *Aedes spp*. mosquitoes—causing fever and debilitating joint pain, with frequent long-term health implications and high morbidity. The CHIKV replication cycle is poorly understood and specific antiviral therapeutics are lacking. In the current study, we identify host cell Musashi RNA binding protein-2 (MSI-2) as a proviral factor. MSI-2 depletion and small molecule inhibition assays demonstrated that MSI-2 is required for efficient CHIKV genome replication. Depletion of both MSI-2 and MSI-1 homologues was found to synergistically inhibit CHIKV replication, suggesting redundancy in their proviral function. Electromobility shift assay (EMSA) competition studies demonstrated that MSI-2 interacts specifically with an RNA binding motif within the 5' untranslated region (5'UTR) of CHIKV and reverse genetic analysis showed that mutation of the binding motif inhibited genome replication and blocked rescue of mutant virus. For the first time, this study identifies the proviral role of MSI RNA binding proteins in the replication of the CHIKV genome, providing important new insight into mechanisms controlling replication of this significant human pathogen and the potential of a novel therapeutic target.

INTRODUCTION

Chikungunya virus (CHIKV) is transmitted by *Aedes spp.* mosquitos and is an alphavirus of the *Togaviridae* family, which includes other medically relevant alphaviruses such as Semliki Forest virus (SFV), Venezuelan equine encephalitis virus (VEEV), and Sindbis virus (SINV) (1,2). CHIKV was first identified in Tanzania in 1952 and symptoms typically include acute febrile symptoms, myalgia, rash and severe arthralgic joint pain, which may persist for months or years (3). CHIKV recently caused epidemic outbreaks across regions of Asia, Africa, the Americas, the Middle East and Southern Europe (3). To date, three phylogenetically distinct lineages of CHIKV have been identified, namely the West African, Asian and the East Central Southern African (ECSA) lineages (4). There remains no clinically approved specific antiviral therapy, due in part to a lack of detailed understanding of the CHIKV replication cycle and its interaction with host cell factors. The first attenuated live CHIKV vaccine (VLA1553) received U.S. FDA approval in November 2023 (5).

CHIKV is an enveloped, positive-sense single-stranded RNA virus with an ~11.8 Kb genome, containing two open reading frames (ORFs) flanked by 5' and 3' untranslated regions (UTRs) (**Fig. 1A**). The 5' UTR is ~ 76 nts in length and is capped by 5' type-0 N-7-methylguanosine. The upstream ORF (ORF-1) encodes the viral non-structural proteins 1-4 (nsP1-4), which are translated directly from the genomic RNA as a single polyprotein that is subsequently proteolytically cleaved into the four mature proteins (6). Through analogy with other alphaviruses, proteolytic cleavage of nsP1-4 in *cis* by nsP2 releases nsP4 which, as the RNA-dependent RNA polymerase

(RdRp), initiates synthesis of the minus-strand intermediate RNA. Subsequent

proteolytic cleavage of the remaining nsP123 polyprotein initiates replication of

genomic and sub-genomic (29S) RNAs from the minus-strand template (7). The

downstream ORF (ORF-2) encodes the structural polyprotein that is processed into

the capsid and envelope proteins, E3, E2, 6K, and E1.

While negative strand replication is initiated at the 3' end of the virus genome, as

with many positive single stranded RNA viruses, regulatory elements and

interactions within the 5' end of the CHIKV genome are required for its initiation and

regulation (8). We and others have previously demonstrated that such regulatory

elements include functional RNA secondary structures and higher order interactions

within the CHIKV 5'UTR and adjacent upstream region of ORF-1 (Fig 1). It has been

speculated that these RNA elements may regulate template specificity and temporal

control of switching, between CHIKV translation and genome replication (9,10).

In a previously published study, we demonstrated by SHAPE mapping and reverse

genetic analysis that the CHIKV 5'UTR and adjacent ORF-1 coding region is highly

structured (10). However, the study also highlighted a single-stranded region (nts 63-

69) exhibiting a high degree of nucleotide conservation that was located

immediately between two RNA structures (Supplementary data 1), which we

demonstrated were essential for initiation of CHIKV genome replication (Fig 1B).

Interestingly, *in silico* analysis noted that nts $_{63}$AUUAAU$_{68}$ were closely homologous

to the Musashi RNA binding protein (MSI) consensus binding sequence ((G/A)U$_{1-}$

$_3$AGU). MSI are highly conserved RNA binding proteins, containing two highly

94 conserved tandem RNA recognition motifs (RNP-1 and RNP-2), that interact with

95 RNA via the same consensus binding motif. Two MSI homologues have been

96 identified, MSI-1 and MSI-2, that share over 90% homology in their RNA-binding

97 domains and a high degree of functional complementarity and redundancy (11,12).

98 MSI have key roles in post-transcriptional regulation of genes involved in

99 development, cell cycle regulation and maintenance of adult neural stem/progenitor

100 cells (13). A recent study demonstrated that MSI-1 promotes Zika Virus (ZIKV)

101 genome replication in neurons, via interaction with the consensus binding site within

102 the viral 3'UTR (14).

104 In the current study, we demonstrate through a range of infectious clone and sub-

105 genomic replicon reporter assays that MSI-2 is required for efficient CHIKV genome

106 replication. A small molecule inhibitor of MSI-1 and MSI-2 reduced infectious virus

107 production by inhibition of CHIKV genome replication. Similarly, MSI-2 silencing by

108 shRNA and siRNA inhibited both infectious virus production and genome replication,

109 while co-silencing of both MSI-2 and MSI-1 exhibited a synergistic effect on inhibition

110 of CHIKV replication. Reverse genetic analysis, in which the putative MSI binding site

111 within the CHIKV 5'UTR was mutated, completely prevented rescue of mutant virus

112 and inhibited CHIKV genome replication. Biochemical MSI-2 binding analysis by

113 competition electromobility shift assay (EMSA) demonstrated that recombinantly

114 expressed MSI-2 specifically interacts with the 5' region of the CHIKV genome and

115 that this interaction is inhibited by mutation of the putative MSI 5'UTR binding site.

116 These findings demonstrate for the first time that interaction between MSI and a

117 binding site within the 5'UTR are required for replication of the CHIKV genome.

Materials and Methods

Cell culture

Human Rhabdomyosarcoma (RD), Human hepatoma (Huh7), Baby Hamster Kidney (BHK) and Human Embryonic Kidney (HEK) 293 were grown in Dulbecco's modified eagle's medium (DMEM; Sigma) containing 10% foetal bovine serum (FBS; PAA), 1x penicillin-streptomycin (Sigma), 25mM HEPES in 0.85% NaCl (Lonza) and 1% non-essential amino acids mixture (NEAA; Lonza). Cells were harvested using trypsin/EDTA (Sigma) and maintained at 37°C in 5% CO_2.

CHIKV cDNA plasmid

The infectious CHIKV (ICRES), CHIKV Dual-luciferase sub-genomic replicon (CHIKV-SGR) and *trans*-complementation (pCHIKV-nsP1234 and pCHIK-Fluc/Gluc) assay were derived from the CHIKV ECSA strain, isolate LR2006 OPY1 (accession number DQ443544)[15]. In CHIKV-SGR the second ORF was replaced by a firefly luciferase gene and a *Renilla* luciferase gene was fused within nsP3 [16]. The *trans*-complementation assay utilised a CMV codon optimised plasmid to supply the CHIKV replicase in *trans* (pCHIKV-nsP1234) and a Pol II expressed reporter plasmid (pCHIKV-FLuc/GLuc), in which the majority of ORF-1 (downstream of nt 320) was replaced by a firefly luciferase gene and all of ORF-2 by a Gaussia Luciferase gene [17]. Plasmid cDNA was purified using GeneJET Plasmid Maxiprep kits (Thermo Fisher Scientific) according to the manufacturer's instructions.

In vitro RNA transcription

To generate CHIKV-SGR and infectious CHIKV RNA, 2 μg of cDNA was linearised with

Not-I HF and used as a template for transcription of 5' [m7G(5')ppp(5')G] capped

(m7G capped) RNA from an upstream SP6 promoter, using an SP6 mMessage

mMachine kit, according to the manufacturer's instructions (Life Technologies).

Uncapped CHIKV 1–337 RNA, used 1 μg of CHIKV 1–337 PCR DNA as a template for _in_

vitro transcription using the SP6-Scribe™ standard RNA IVT kit, according to the

manufacturer's instructions (Lucigen). In all cases, following DNase I treatment, RNA

was purified by LiCl precipitation and analysed by denaturing agarose gel

electrophoresis.

Virus production

1×10^6 BHK cells were trypsinised and resuspended in 400 μl ice-cold DEPC-PBS. Cells

were then electroporated with 1 μg 5'-capped CHIKV ICRES _in vitro_ transcribed RNA

in a 4 mm electro-cuvette, with a single square wave pulse at 260 V for 25 ms using a

Bio-Rad electroporator, before seeding into a T175 flask in 20 ml DMEM. After 24 h,

supernatant was aspirated and virus titre measured by plaque assay on BHK cells.

CHIKV quantification by plaque assay

BHK cells were seeded at 1×10^5 cells per well in 12-well plates and maintained

overnight in 1 ml DMEM. The following day monolayers were washed with PBS,

infected with 10-fold serial dilutions of CHIKV transfection supernatant and

maintained at 37°C. At 1 hour post infection (h.p.i.) monolayers were washed with

PBS and covered with a 0.8% methylcellulose DMEM P/S overlay. Monolayers were

fixed and stained at 48 h.p.i. (5% paraformaldehyde and 0.25% crystal violet

respectively), plaques counted and virus titres expressed in plaque-forming units per

ml (PFU/ml).

Ro 08-2750 cell viability assay

RD cells were seeded in 96-well plate at 8×10^4 cells/well and maintained for 24 h.

Monolayers were then treated with increasing doses (0, 0.5, 1, 3, 5, 7, 10, 20 μM) of

the MSI small molecule inhibitor Ro 08-2750 (TOCRIS) dissolved in DMSO. After 24 h

media/inhibitor was aspirated and replaced with 20 μl of 5 mg/mL 3-(4,5-

dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) (Sigma) and incubated

at 37°C in 5% CO_2 for 3 h. After incubation, MTT solution was replaced with 100 μl

DMSO and the plate shaken at 60 rpm for 5 min. Absorbance at 570 nm was

determined using an Infinite F50 microplate reader (Tecan) and expressed as a

percentage of DMSO control cells. 5 μM Ro 08-2750 was estimated to be the

maximum non-toxic dose in in Huh7 cells and was used for all further assays.

Infectious CHIKV Ro 08-2750 inhibition assays

RD cells were seeded in 12-well plates at 1×10^5 cells/well and incubated overnight in

the presence of Ro 08-2750. Monolayers were then infected with CHIKV at MOI=0.1

and adsorbed to the cells for 1 h at 37°C before aspirating and maintaining in the

presence of Ro 08-2750 for 24 h. Supernatants was then collected and infectious

virus production measured by plaque assay.

siRNA depletion of MSI-1 and MSi-2

RD or Huh7 cells were seeded at 1×10^5 cell/well in 12-well plates in antibiotic-free

medium. After 24 h, cells were washed with PBS and incubated with 1x Opti-MEM +

GlutaMAX (Gibco) for 20 min at 37°C/5% CO_2. For each well, 50 pmol MSI-1 (sc-

106836; Santa Cruz) and/or MSI-2 siRNA (sc-75834; Santa Cruz) were mixed with 100

µl Opti-MEM and incubated at room temperature for 1min. No siRNAs were added

to mock samples and 50 pmol of scrambled siRNA (SI03650318; QIAGEN) was used

as a negative control. In parallel, 3 µl Lipofectamine RNAiMAX (Invitrogen) was

mixed with 100 µl Opti-MEM and incubated at room temperature for 1 min. The

siRNA and Lipofectamine RNAiMAX were mixed and incubated at room temperature

for 5 min before adding to the cells. After 24 h, cells were either lysed to confirm

MSI-1 or MSi-2 depletion by western blot or used for subsequent CHIKV infection

assays.

shRNA depletion of MSI-1 and MSi-2

Human embryonic kidney 293 (HEK 293T) cells were plated in antibiotic-free DMEM

in 6-well plates and maintained until confluency reached ~80%. For each

transfection, in a single tube 300 µL OptiMEM was mixed with 1 µg packaging

plasmid p8.91, 1 µg envelope plasmid pMDG encoding VSV-G protein and 1.5 µg

lentiviral vector plasmid encoding shRNA to MSI-2 (sc-75834-SH; Santa Cruz

Biotechnology). A lentiviral vector plasmid encoding scrambled shRNA sequences

with no cellular targets was used as the negative control (sc-108060; Santa Cruz

Biotechnology). In another tube, 300 µL of OptiMEM was mixed with 5 µL

lipofectamine 2000 (Invitrogen). Both tubes were gently mixed by flicking, incubated

at room temperature for 5 min, mixed together and incubated for 20 min at room

temperature. The antibiotic-free DMEM was aspirated and the monolayers washed

once with PBS. 800 µL Opti-MEM was added to each well before dropwise addition

of the lentiviral plasmids/shRNA mixture. After maintenance at 37°C for 6 h the

media was changed to antibiotic-free DMEM. 48 hpt the lentivirus containing

supernatant was harvested and filtered through a 0.45 µm filter.

RD cells were seeded at 1×10^5 cells/well the day prior to transduction. 1 mL of the

lentivirus supernatant and polybrene (MERCK) were added to each well and

incubated for 6 h before aspirating and replacing with antibiotic free DMEM. After

72 h media was replaced with DMEM containing 2.5 µg/ml puromycin, in which the

cells were then maintained. The efficiency of the shRNA MSI-2 depletion was

confirmed by western blot.

Strand-specific Quantification of CHIKV RNA

Total RNA was extracted from infected cells using TRI Reagent Solution (Applied

Biosystems) according to the manufacturer's instructions. Strand-specific RT-qPCR

(ssRT-qPCR) was performed as previously described (18). Briefly, 500 ng of RNA was

reverse-transcribed with gene specific primers (Supplementary data 2) using the

SCRIPT cDNA Synthesis Kit (Jena Bioscience) according to the manufacturer's

protocol. 100 ng of strand-specific cDNA was used as template for the quantitative

PCR performed with the qPCRBIO SyGreen Blue Mix Lo-ROX (PCR Biosystems), with

gene specific primers amplifying a 94 bp region of the CHIKV nsP1-encoding

sequence using the following PCR program: 95°C for 2 mins, 40 x (95°C for 5 sec,

60°C for 30 sec), dissociation curve 60°C-95°C as pre-defined by the Mx3005P

thermal cycler (Agilent technologies). *In vitro* transcribed CHIKV ICRES RNA was

reverse transcribed and a cDNA dilution series employed as a standard to quantify

copy numbers in the respective samples.

CHIKV Sub-genomic replicon assays

For analysis of Ro 08-2750 inhibition, RD cells were seeded in 24 well plates at

5×10^4 cells per well and maintained overnight in the presence of Ro 08-2750.

Monolayers were washed once in PBS before addition of 400 µl opti-Mem reduced-

serum media and 100 µl transfection media. Transfection media was prepared

according to the manufacturer's instructions, using 1 µl Lipofectamine 2000

(Invitrogen), 250 ng of CHIKV-SGR RNA and appropriate concentrations of Ro 08-

2750, before being made up to 100 µl using opti-Mem. Monolayers were maintained

for 6 hpt before the media was aspirated and replaced with complete DMEM/Ro 08-

2750. At 8 and 24 hpt monolayers were lysed with 100 µl 1 x passive lysis buffer

according to the manufacturer's instructions (Promega), stored at −80°C and

analysed using Dual-luciferase substrate (Promega) in a FLUOstar Optima

luminometer (BMG labTech). For shRNA MSI-2 depleted cell lines the same method

was followed, with the exclusion of Ro 08-2750.

***Trans*-complementation Assay**

The *trans*-complementation assay was performed as previously described (19).

Briefly, for analysis of Ro 08-2750 inhibition, RD cells were seeded in 12 well plates at

5×10^4 cells per well and maintained overnight in the presence of Ro 08-2750.

Monolayers were then co-transfected with 1 µg each of the pCHIKV-nsP1234 and

pCHIKV-FLuc/GLuc using Lipofectamine 2000, as described previously, and assayed

at 8 and 24 hpt. For shRNA MSI-2 depleted cell lines the same method was followed,

with the exclusion of Ro 08-2750.

Expression and purification of recombinant MSI-2 RNA binding domains

The MSI-2 RNA binding domain expressing plasmid pET-22HT-MSI-2 (amino acid 8-

193 and referred to hereafter as MSI-2 8-193) was a kind gift from Prof S. Ryder

(Addgene plasmid # 60356; http://n2t.net/addgene:60356; RRID: Addgene_60356)

(20). The plasmid was transformed into BL21 (DE3) competent cells following the

manufacturer's protocol (NEB), a single colony was inoculated into 10 mL LB

ampicillin (10 mg/ml) and incubated overnight at 37°C before inoculating 1000 ml 1

LB (ampicillin (10 mg/ml) and adding 100 µM Isopropyl β-D-1-thiogalactopyranoside

(IPTG; Thermo Fisher Scientific) when the optical density reached 0.8. The culture

was then incubated overnight at 18°C in an orbital shaker before centrifugation to

pellet the bacteria. The pellet was resuspended in lysis buffer (20U/mL DNase I,

0.6U/mL RNase A, 1mg/mL lysozyme and 1x protease inhibitor) and lysed on ice for

30 min. Following sonication, the suspension was centrifuged twice at 2000 xg for 1

hour at 4°C. The was filtered through a 0.45 µm filter and His-MSI-2 was purified

with HisTrap™ FF column (Cytiva) using the Econo Gradient Pump (BIORAD)

according to the manufacturer's protocol and overnight dialysis. The purified protein

was quantified and the purity and identity assayed by Coomassie SDS-PAGE and

western blot (Supplementary data 3).

After desalting using PD-10 desalting columns (Cytiva), His-MSI-2 protein was further

purified by ion exchange chromatography. Following the manufacturer's

instructions, the column was equilibrated with wash buffer (50 mM MES, 10 mM

NaCl, pH 5.6, degassed) and the protein eluted in the same buffer. Ion exchange

chromatography was performed using HiTrap SP HP cation exchange

chromatography column (Cytiva) following the manufacturer's instructions and

eluted in degassed 50mM MES, 1M NaCl, pH 5.6. The eluted protein fractions were

analysed by Coomassie SDS-PAGE and western blot (Supplementary data 3).

^{32}P RNA end labelling

In vitro transcribed CHIKV RNA (nt's 1-337) was 5' dephosphorylated using Quick CIP

according to the manufacturer's instructions (NEB) before purification using an RNA

Clean & Concentrator column (Zymo Research). 20 pmoles of dephosphorylated RNA

was combined with 2 μl 10 x T4 Polynucleotide Kinase buffer (NEB), 3 μl ATP(γ-^{32}P)

10 mCi/ml, 1 μl T4 Polynucleotide Kinase (NEB) and nuclease free H_2O to a final

volume of 20 μl, incubated at 37°C for 60 min, 65°C for 20 min, purified using an RNA

Clean & Concentrator Kit (Zymo Research) and resuspended in RNase-free H_2O.

Electromobility shift assay (EMSA)

RNA was incubated at 95°C for 2 min and on ice for 2 min before 3.3 x RNA folding

buffer (100 mM HEPES pH 8.0, 100 mM NaCl and 10 mM MgCl2) and RNasin Plus

RNase Inhibitor (Promega) was added to a final volume of 10 µl and incubated for

20 min at 37°C. MSI-2 protein was combined with 3.3 x RNA folding buffer, 0.5 x TE,

100% glycerol, RNasin Plus RNase Inhibitor (Promega) and 5 µg yeast tRNA as non-

specific competitor and incubated at 37°C for 2 min before adding to the RNA

mixture and further incubation at 37°C for 15 min. Finally, samples were analysed by

native PAGE gel electrophoresis at 135 V for ~2 h. The gel was fixed for 30 min and

dried with gel dryer (BIORAD), exposed onto Hyperfilm™ ECL™ (Merck) and

visualised using a Xograph Film Processor. Band shifts in the competition EMSAs

were quantified by measuring the density of the RNA/Protein complex band (as

indicated by square brackets) normalized to the density of the whole lane and

expressed as % of the normalized RNA/Protein complex band in lane 1:0.

Western blotting

Following infection and incubation, monolayers were lysed in IP lysis buffer

(Promega) and incubated at room temperature for 30 min. Protein concentration

was quantified using a Pierce™ BCA Protein Assay Kit (Thermo scientific) according to

the manufacturer's instructions. Equal amounts of protein lysate were analysed by

SDS-PAGE. Protein was transferred onto an Immobilon-FL PVFD transfer membrane

(MERCK) using a TE77X semi-dry transfer (Hoefer) at 15 V for 60 min. Membranes

were blocked using diluted Odyssey® Blocking Buffer in PBS (LI-COR) for 30 min and

probed with primary antibodies against β-actin (1:10,000, mouse monoclonal, Sigma-

Aldrich A1978), MSI-2 (1:1000 rabbit monoclonal, Abcam ab76148) and MSI-1

(1:5000, Abcam ab21628) in diluted Odyssey® Blocking Buffer in PBS (LI-COR)

overnight at 4°C. After overnight incubation, primary antibody was removed and

membranes washed 3 times using PBS. Membranes were stained with secondary

antibodies (IRDye® 800CW Donkey anti-Mouse; IRDye® 680LT Donkey anti-Rabbit; Li-

Cor) for 1 h at room temperature, washed 3 times using 1 x PBS, dried and then

imaged using an Odyssey® Fc Imaging System (Li-Cor).

Statistical analysis

Statistical analysis was carried out using two-tailed Students Test comparisons on

GraphPad Prism version 8.4.0. *P* values of ≤0.05 (*), ≤0.01 (**), ≤0.001 (***) were

used to represent degrees of significance between each drug

treatment/silencing/mutant to wild-type assay.

RESULTS

MSI-2 binds specifically to the predicted $_{63}$AUUAAU$_{68}$ 5'UTR MSI binding site in the

CHIKV 5'UTR

In order to examine the potential for direct interaction between the RNA binding

domains of MSI-2 and nucleotides $_{63}$AUUAAU$_{68}$ within the CHIKV 5'UTR, which had

close sequence homology to the (G/A)U$_{1-3}$AGU MSI consensus nucleotide binding

motif, we used native EMSA (21). $_{63}$AUUAAU$_{68}$ is located in a single-stranded region

of the 5' UTR, 9 nts upstream of the AUG start codon and between two RNA

structures that are essential for CHIKV genome replication (Fig 1B)(10). The first

330nts of the CHIKV genome (RNA-330), representing the 5'UTR and adjacent

upstream region of ORF1, was *in vitro* transcribed, 5' end radiolabelled with ATP-[γ-

32P] and incubated at 37°C - conditions under which correct folding of the functional

RNA-330 structure was previously validated (10). Following incubation with

increasing concentrations of recombinantly expressed and purified MSI-2 8-193 in

the presence of unlabelled tRNA, reaction products were separated by native PAGE

and analysed by autoradiography (Fig 2A). Relative to unbound RNA-330, the

presence of MSI-2 8-193 retarded RNA-330 migration during native PAGE, consistent

with the formation of an RNA-330/MSI-2 8-193 complex, the increasing size of the

retarded band potentially indicating MSI-2 multimerization. In order to investigate

the specificity of the observed interaction, a fixed molar ratio of RNA-330$^{γ-32P}$ and

MSI-2 8-193 was incubated with increasing concentrations of competitor unlabelled

RNA-330 (Fig 2B). Increasing concentrations of unlabelled RNA-330 reduced the

formation of the RNA-330$^{γ-32P}$/MSI-2 8-193 complex, with a corresponding increase

in unbound RNA-330$^{γ-32P}$.

To further investigate the specificity and location of the interaction, the putative

MSI-2 binding site $_{63}$AUUAAU$_{68}$ was mutated to $_{63}$CAACUU$_{68}$ (henceforth termed

$_{63}$CAACUU$_{68}$-mut), which was predicted *by in silico* analysis not to affect local RNA

structure (Supplementary data 4A and B). EMSA competition, with unlabelled

$_{63}$CAACUU$_{68}$-mut RNA-330, was less efficient at competing for MSI-2 8-193 binding

than wild-type CHIKV RNA (Fig 2C and D). These EMSA results are consistent with an

interaction between MSI-2 and the upstream 1-330 nt region of the CHIKV genome.

Furthermore, reduced competition with mutant $_{63}$CAACUU$_{68}$-mut compared with

wild type, was consistent with a specific interaction, in which nts $_{63}$AUUAAU$_{68}$

function as an MSI binding motif.

Replication of CHIKV infectious virus and sub-genomic replicon was restricted by

MSI small molecule inhibitor Ro 08-2750:

In order to investigate the potential role of MSI on CHIKV replication, we first

assessed the effect of a well-characterised MSI small molecule inhibitor, Ro 08-2750,

on CHIKV productive replication in RD cells using both CHIKV infectious virus and a

sub-genomic replicon (SGR) assay (Fig 3A). A number of studies have demonstrated

that Ro interacts with the MSI RNA binding domains and acts as a competitive

inhibitor for its RNA binding activity and subsequent MSI functions within the cell

(22,23) (24). In the current study Ro 08-2750 was used at a maximum non-toxic dose

as determined by MTT assay (Supplementary data 5).

To assay the effect of Ro 08-2750 on early and late stages of CHIKV replication,

productive CHIKV infection was measured at 8 and 24 hpi. Pre-treatment and

incubation in the presence Ro significantly inhibited infectious CHIKV replication by

~10-fold, relative to DMSO treated negative controls (Fig 3B). In order to investigate

the effect of Ro 08-2750 on specific stages of CHIKV replication we used an SGR

construct, in which ORF-2 was replaced by a firefly luciferase (FLuc) gene and a

Renilla luciferase (Rluc) gene was fused in-frame with ORF-1 nsP3 (Fig 3A)(10). The

SGR assay enabled us to measure the effect of Ro 08-2750 on CHIKV genome

replication and translation, in isolation of other stages of virus infection, such as

entry or egress. SGR ORF-1 and ORF-2 expression was significantly inhibited at both 8

and 24 hpt, relative to DMSO treated negative controls, indicating that Ro 08-2750

inhibited CHIKV replication at the level of virus genome replication or translation (Fig

3C).

In order to investigate the possibility that Ro 08-2750 specifically inhibited ORF-1

translation, we used a replication incompetent SGR, incorporating an nsP4 GDD>GAA

substitution in the RdRp active site (CHIKV-SGR (GDD>GAA)). Such an approach

enabled the efficiency of ORF-1 translation to be measured by *Renilla* luciferase

expression, in the presence and absence of Ro 08-2750, under conditions in which

genome replication could not occur. Translation of input SGR 5'capped transcripts

was measured at 6 hpt in RD cells (Fig 3E). No significant difference in level of

translation was observed between the CHIKV-SGR (GDD>GAA) in the presence or

absence of Ro 08-2750. These results indicate that Ro 08-2750 did not influence ORF-

1 translation but inhibited SGR genome replication.

Ro inhibited CHIKV *trans*-complementation assay:

Inhibition of SGR replication demonstrated that Ro was inhibiting CHIKV replication

at the level of virus genome replication or translation. In order to dissect this further,

we utilised a *trans*-complementation assay, that enabled the measurement of CHIKV

genome replication in isolation of virus translation (Fig 4A)(25). The *trans*-

complementation system utilised a CHIKV replicase-expressing plasmid (pCHIKV-

nsP1234). pCHIKV-nsP1234 , expresses the virus nsPs from a CMV promoter and is

codon optimised to disrupt RNA sequence motifs, such as protein binding signals,

and the formation of wild-type CHIKV RNA structure. The expressed nsPs replicate a

CHIKV reporter construct (pCHIKV-FLuc/GLuc), in which the majority of ORF-1 is

replaced by an FLuc gene and ORF-2 by the Gaussia luciferase (GLuc) gene. The effect

of Ro on pCHIKV-FLuc/GLuc expression was measured at 8 and 24 hpt in cells pre-

treated with and maintained in the presence of Ro. Both ORF-1 and ORF-2 expression

was observed to be significantly inhibited relative to the DMSO treated negative

controls, indicating that Ro was specifically inhibiting CHIKV genome replication.

MSI-2 shRNA silencing inhibits infectious CHIKV replication

In order to confirm that Ro induced inhibition of CHIKV genome replication was due

to specific inhibition of MSI, rather than an unrecognised off target effect of Ro, we

next investigated CHIKV replication following shRNA silencing of MSI expression.

Analysis of RD cell total protein extract by western blot, demonstrated that MSI-2

was strongly expressed in RD cells while MSI-1 was expressed at a very low level

(Supplementary data 6). Consequently, we initially investigated the effect of specific

MSI-2 shRNA silencing on CHIKV replication. RD cells were transduced with lentiviral

vectors encoding shRNA against human MSI-2 and successful silencing was

confirmed by western blot (Fig 5A). Negative control cells were treated with either

transfection reagent only (mock) or non-specific scrambled shRNA. Following

infection of MSI-2 depleted and negative control cells, productive CHIKV infection

was measured by plaque assay at 8 and 24 hpi (Fig 5B). Strand specific qRT-PCR was

also used to measure levels of genomic (positive strand) and replication

intermediate (negative strand) CHIKV RNA at both time points (Fig 5C and D). As

observed previously following treatment with Ro, relative to the scrambled shRNA

control productive CHIKV replication was significantly inhibited following shRNA MSI-

2 silencing (Fig 5B). Similarly, strand specific qRT-PCR demonstrated that both

genomic and replication intermediate CHIKV RNA levels were significantly inhibited

(Fig 5C and D). These results are consistent with CHIKV requiring MSI-2 for efficient

genome replication.

MSI-2 is required for efficient CHIKV genome replication

In order to confirm that MSI-2 silencing had the same effect on CHIKV genome

replication as Ro small molecule inhibition, we repeated analysis with the SGR and

trans-complementation systems, following shRNA silencing of MSI-2. As previously

described, MSI-2 shRNA silenced RD cells were transfected with the SGR system and

ORF-1 and ORF-2 expression measured by RLuc and FLuc expression at 8 and 24 hpt

(Fig 6A and B). Expression of both ORF-1 and ORF-2 was significantly inhibited at

both 8 and 24 hpt, consistent with an MSI-2 requirement for CHIKV genome

replication or translation. Analysis at the same time points using the *trans*-

complementation system, following shRNA MSI-2 silencing and measuring ORF1 and

ORF2 expression, confirmed significant inhibition of CHIKV genome replication when

measured in isolation of the effects of other stages of the virus replication cycle (Fig

6C and D)

Both MSI-2 and MSI-1 have a proviral effect on CHIKV genome replication:

While results clearly demonstrated that MSI-2 was required for efficient CHIKV

replication, it remained unclear if MSI homologue MSI-1 was also agonistic for CHIKV

replication. As previously described, western blot analysis showed that MSI-1 was

not highly expressed in RD cells. Consequently, we analysed MSI homologue

redundancy in Huh7 human hepatoma cells. MSI-1 and MSI-2 homologues are

strongly expressed in Huh7 cells (Supplementary data 6) and they are highly

permissive for CHIKV replication (26). Following individual and combined siRNA

silencing of MSI-1 and MSI-2 in Huh7 cells (Supplementary data 7), CHIKV replication

was assayed by plaque assay and ssRT-qPCR at 8 and 24 hpi (Fig 7A, B and C). As was

observed for RD cells, MSI-2 silencing significantly inhibited CHIKV replication,

measured by both plaque assay and strand specific ssRT-qPCR. Interestingly, it was

observed that MSI-1 silencing in Huh7 cells had a similar significant inhibitory effect

and that siRNA co-silencing of both MSI-1 and MSI-2 had a synergistic effect on

inhibition of CHIKV replication. In concordance with previous results, siRNA silencing

of MSI-2 in RD cells significantly inhibited CHIKV replication, by comparable levels

observed following shRNA silencing and inhibition by Ro. Consistent with low MSI-1

expression in RD cells, siRNA co-silencing of both MSI-1 and MSI-2 did not

significantly increase levels of CHIKV inhibition (Fig 7D, E and F).

Reverse genetic analysis of $_{63}$AUUAAU$_{68}$ MSI binding site

In order to further investigate the role of the $_{63}$AUUAAU$_{68}$ MSI binding motif in CHIKV

replication, we took a reverse genetic approach, in which the

$_{63}$AUUAAU$_{68}$>$_{63}$CAACUU$_{68}$ mutations were incorporated into both the *trans*-

complementation assay and infectious CHIKV. While $_{63}$AUUAAU$_{68}$>$_{63}$CAACUU$_{68}$

disrupted the MSI binding site, it was predicted to have no effect on local RNA

structure (Supplementary data 4A and B). Similar to previous results following MSI

inhibition, expression of both ORF-1 and ORF-2 from the *trans*-complementation

system were significantly inhibited in $_{63}CAACUU_{68}$-mut relative to the wild type at 8

and 24 hpt (Fig 8A and B); indicating that mutation of the putative MSI binding site

significantly inhibited CHIKV at the level of virus genome replication. Interestingly,

despite repeated attempts, we were not able to rescue infectious CHIKV $_{63}CAACUU_{68}$-

mut virus, indicating that disruption of the potential MSI binding site completely

inhibited infectious CHIKV replication (Fig 8C). In further reverse genetic analysis, we

analysed the effect of substitution $_{A}67_{G}$ on infectious CHIKV replication. $_{A}67_{G}$

modified the $_{63}AUUAAU_{68}$ to match the published MSI canonical binding site but

disrupted adjacent downstream RNA structure (Supplementary data 4C). Following

infection at MOI 0.1 in RD cells, CHIKV $_{A}67_{G}$ mutant and wild-type control virus was

harvested at 8 and 24 hpi and titred by plaque assay (Fig 8D). Compared to wild-

type, mutant $_{A}67_{G}$ significantly inhibited CHIKV replication at 8 and 24 hpi, indicating

that nucleotides $_{63}AUUAAU_{68}$ are constrained by both a requirement to interact with

MSI-2 and also to maintain essential local RNA structure.

Ro 08-2750 further inhibited CHIKV genome replication in $_{63}CAACUU_{68}$-mut

In previous experiments, we demonstrated that mutations to disrupt the $_{63}AUUAAU_{68}$

5'UTR MSI binding site or MSI-2 depletion/inhibition restricted CHIKV genome

replication. In order to investigate the potential synergistic effects of these

approaches we repeated analysis of $_{63}CAACUU_{68}$-mut *trans*-complementation assay

replication, in the presence of Ro 08-2750 (Fig 9). Interestingly, in the presence of Ro

08-2750, $_{63}CAACUU_{68}$-mut replication was inhibited to a significantly greater level

than $_{63}CAACUU_{68}$-mut in the absence of Ro 08-2750, at both 8 and 24 hpt. These

results are consistent with MSI-2 acting through interaction with $_{63}$AUUAAU$_{68}$ in the

5'UTR and a further region of the CHIKV genome present in the sub-genomic *trans*

complementation reporter (i.e., 5'UTR, intragenic region or 3'UTR).

DISCUSSION

For the first time, in this study we demonstrate that cellular RNA binding protein MSI

is required for efficient CHIKV replication. Our data demonstrates that MSI-2

interacts directly with the 5' end of the CHIKV genome and is consistent with an

interaction at position $_{63}$AUUAAU$_{68}$. A single-stranded region located between two

conserved stem-loops and immediately upstream of the AUG start codon (Fig 1) (10).

Inhibition of MSI-2 expression by siRNA or shRNA, its RNA-binding activity by Ro 08-

2750 and reverse genetic disruption of the CHIKV-5'UTR/MSI-2 interaction, inhibited

CHIKV replication at the level of virus genome replication. In further analysis, we

demonstrated that depletion of both MSI-1 and MSI-2 homologues resulted in a

synergistic increase in the level of CHIKV inhibition, supporting the premise that both

MSI homologues have a redundant pro-viral effect on CHIKV replication.

While depletion of MSI-2 by siRNA and shRNA clearly demonstrated significant

inhibition of CHIKV replication, measured by infectious virus production, CHIKV-SGR

replication or a *trans*-complementation assay, we did not observe complete

inhibition of virus replication. Western blot analysis indicated that we did not

achieve complete ablation of MSI-2 expression by either siRNA or shRNA,

presumably reducing the level of CHIKV inhibition observed. We demonstrated by

western blot that in RD cells MSI-2 is expressed to a high level but the MSI-1

homologue is only expressed to a low level. However, given the synergistic effect

that we observed when silencing expression of both MSI-2 and MSI-1 in Huh7 cells

(in which both are expressed to high levels), the ability of MSI homologues to

complement for each other may also have reduced the level of CHIKV inhibition

observed. This hypothesis is consistent with reverse genetic results, in which

$_{63}AUUAAU_{68}>_{63}CAACUU_{68}$ mutation of the MSI binding site, dramatically inhibited

CHIKV-5'UTR/MSI-2 binding affinity and completely prevented rescue of infectious

mutant virus.

Mutation $_{63}AUUAAU_{68}>_{63}CAACUU_{68}$ also significantly inhibited replication in the

trans-complementation assay. However, genome replication was not completely

ablated. These differences may indicate that this region has a further function during

infectious virus replication, that is not assayed in the *trans*-complementation assay,

such as during virion assembly. Further reverse genetic analysis, in which

substitution $_{A}67_{G}$ engineered a canonical MSI binding site but was predicted to

disrupted downstream RNA structure, significantly inhibited infectious CHIKV

replication, suggesting that $_{63}$AUUAAU$_{68}$ is constrained by both a requirement to bind

MSI and to maintain local RNA structure.

Translation and replication of positive-sense RNA virus genomes, such as those of

CHIKV are mutually exclusive processes – with translation initiating at the 5' end of

the genome and replication at the 3'. Consequently, it is essential that such viruses

have mechanisms for temporal control of both processes. In many such viruses (e.g.

Hepatitis C virus and Polio virus) control involves dynamic interactions between RNA

structures in the virus genome and host/virally expressed protein complexes (8). The

mechanisms and interactions which CHIKV and other alphaviruses use for temporal

control of genome translation and replication remain unclear. However, a recent

study suggested a crucial role for the cellular helicase DHX9. Matkovic et al,

demonstrated that interaction between DHX9 and the 5' end of the virus genome

upregulates CHIKV ORF-1 non-structural protein translation, while simultaneously

inhibiting replication of its genome (27). Build-up of nsP2 caused proteosome

induced degradation of DHX9, although the detailed mechanism for this remains

unclear.

While, genome replication of positive-sense RNA viruses initiates at the 3' end, it is

commonly controlled by promoter elements and interactions with RNA binding

proteins at the 5' end. Results described in this and previous studies are consistent

with a model in which temporal control of CHIKV ORF-1 translation and genome

replication is controlled by a mechanism involving interactions between DHX9, MSI-2

and the 5' region of the virus genome. However, the dynamics and mechanism by

which the opposing roles if MSI-2 and DHX-9 influence CHIKV replication and their

interactions with other regions of the virus genome remains unclear. For example,

EMSA and reverse genetic results presented here clearly demonstrate that MSI-2

binds to nucleotides $_{63}AUUAAU_{68}$ in the 5'UTR and that this interaction is required for

efficient CHIKV genome replication. However, the synergistic effect of Ro 08-2750

and $_{63}CAACUU_{68}$-mut are consistent with MSI-2 also interacting with another region

of the CHIKV genome. Bioinformatic analysis has not predicted further MSI binding

motifs within the CHIKV genome. Consequently, we hypothesise that any such

alternative interactions may be indirect, via a protein complex and are the focus of

ongoing studies.

In summary, using MSI-2 depletion and/or Ro 08-2750 small molecule inhibitor we

demonstrate that MSI-2 RNA binding protein is a critical host factor for efficient

CHIKV replication but is not required for ORF-1 translation. Inhibition of ORF-1 and

ORF-2 signal from the *trans*-complementation assay indicate that MSI-2 is required

at the level of CHIKV genome replication. Furthermore, EMSA and reverse genetic

results are consistent with a direct interaction between MSI-2 and $_{63}AUUAAU_{68}$ in the

positive genomic strand of the virus 5'UTR, suggesting a role in negative strand

synthesis. Results from this study are important both for our understanding of the

fundamental interactions essential to replication of this important human pathogen

and for future studies towards specific antiviral therapies.

Funding

This work was supported by funding from the MRC to AT (MR/NO1054X/1).

Data availability statement

The data underlying this article will be shared on reasonable request to the corresponding author.

REFERENCES

1. Fong, S.W., Kini, R.M. and Ng, L.F.P. (2018) Mosquito Saliva Reshapes Alphavirus Infection and Immunopathogenesis. *Journal of Virology*, **92**.

2. McPherson, R.L., Abraham, R., Sreekumar, E., Ong, S.E., Cheng, S.J., Baxter, V.K., Kistemaker, H.A.V., Filippov, D.V., Griffin, D.E. and Leung, A.K.L. (2017) ADP-ribosylhydrolase activity of Chikungunya virus macrodomain is critical for virus replication and virulence. *P Natl Acad Sci USA*, **114**, 1666-1671.

3. Petitdemange, C., Wauquier, N. and Vieillard, V. (2015) Control of immunopathology during chikungunya virus infection. *Journal of Allergy and Clinical Immunology*, **135**, 846-855.

4. Ng, L.F. (2017) Immunopathology of Chikungunya virus infection: lessons learned from patients and animal models. *Annual review of virology*, **4**, 413-427.

5. Schneider, M., Narciso-Abraham, M., Hadl, S., McMahon, R., Toepfer, S., Fuchs, U., Hochreiter, R., Bitzer, A., Kosulin, K., Larcher-Senn, J. *et al.* (2023) Safety and immunogenicity of a single-shot live-attenuated chikungunya vaccine: a double-blind, multicentre, randomised, placebo-controlled, phase 3 trial. *Lancet*, **401**, 2138-2147.

6. Meshram, C.D., Agback, P., Shiliaev, N., Urakova, N., Mobley, J.A., Agback, T., Frolova, E.I. and Frolov, I. (2018) Multiple Host Factors Interact with the Hypervariable Domain of Chikungunya Virus nsP3 and Determine Viral

Replication in Cell-Specific Mode. *J Virol*, **92**.

7.   Rupp, J.C., Sokoloski, K.J., Gebhart, N.N. and Hardy, R.W. (2015) Alphavirus
RNA synthesis and non-structural protein functions. *Journal of General*
Virology, **96**, 2483-2500.

8.   Tuplin, A. (2015) Diverse roles and interactions of RNA structures during the
replication of positive-stranded RNA viruses of humans and animals. *J Gen*
Virol.

9.   Frolov, I., HARDY, R. and RICE, C.M. (2001) Cis-acting RNA elements at the
5′ end of Sindbis virus genome RNA regulate minus-and plus-strand RNA
synthesis. *Rna*, **7**, 1638-1651.

10.   Kendall, C., Khalid, H., Muller, M., Banda, D.H., Kohl, A., Merits, A.,
Stonehouse, N.J. and Tuplin, A. (2019) Structural and phenotypic analysis of
Chikungunya virus RNA replication elements. *Nucleic Acids Res*, **47**, 9296-
9312.

11.   Troschel, F.M., Minte, A., Ismail, Y.M., Kamal, A., Abdullah, M.S., Ahmed,
S.H., Deffner, M., Kemper, B., Kiesel, L., Eich, H.T. *et al.* (2020) Knockdown
of Musashi RNA Binding Proteins Decreases Radioresistance but Enhances
Cell Motility and Invasion in Triple-Negative Breast Cancer. *Int J Mol Sci*, **21**.

12.   Sutherland, J.M., Siddall, N.A., Hime, G.R. and McLaughlin, E.A. (2015) RNA
binding proteins in spermatogenesis: an in depth focus on the Musashi family.
Asian journal of andrology, **17**, 529-536.

13.   Horisawa, K., Imai, T., Okano, H. and Yanagawa, H. (2010) The Musashi family
RNA-binding proteins in stem cells. *Biomolecular concepts*, **1**, 59-66.

14.   Chavali, P.L., Stojic, L., Meredith, L.W., Joseph, N., Nahorski, M.S., Sanford,
T.J., Sweeney, T.R., Krishna, B.A., Hosmillo, M. and Firth, A.E. (2017)
Neurodevelopmental protein Musashi-1 interacts with the Zika genome and
promotes viral replication. *Science*, **357**, 83-88.

15.   Tsetsarkin, K., Higgs, S., McGee, C.E., De Lamballerie, X., Charrel, R.N. and
Vanlandingham, D.L. (2006) Infectious clones of Chikungunya virus (La
Reunion isolate) for vector competence studies. *Vector Borne Zoonotic Dis*, **6**,
325-337.

16.   Pohjala, L., Utt, A., Varjak, M., Lulla, A., Merits, A., Ahola, T. and Tammela, P.
(2011) Inhibitors of alphavirus entry and replication identified with a stable

Chikungunya replicon cell line and virus-based assays. *PloS one*, **6**, e28923.

17.   Lello, L.S., Utt, A., Bartholomeeusen, K., Wang, S., Rausalu, K., Kendall, C.,
Coppens, S., Fragkoudis, R., Tuplin, A., Alphey, L. *et al.* (2020) Cross-
utilisation of template RNAs by alphavirus replicases. *PLoS pathogens*, **16**,
e1008825.

18.   Muller, M., Slivinski, N., Todd, E., Khalid, H., Li, R., Karwatka, M., Merits, A.,
Mankouri, J. and Tuplin, A. (2019) Chikungunya virus requires cellular chloride
channels for efficient genome replication. *PLoS neglected tropical diseases*, **13**,
e0007703.

19.   Bartholomeeusen, K., Utt, A., Coppens, S., Rausalu, K., Vereecken, K., Arien,
K.K. and Merits, A. (2018) A Chikungunya Virus trans-Replicase System
Reveals the Importance of Delayed Nonstructural Polyprotein Processing for
Efficient Replication Complex Formation in Mosquito Cells. *J Virol*, **92**.

20.   Clingman, C.C., Deveau, L.M., Hay, S.A., Genga, R.M., Shandilya, S.M., Massi,
F. and Ryder, S.P. (2014) Allosteric inhibition of a stem cell RNA-binding
protein by an intermediary metabolite. *eLife*, **3**, e02848.

21.   Zearfoss, N.R., Deveau, L.M., Clingman, C.C., Schmidt, E., Johnson, E.S.,
Massi, F. and Ryder, S.P. (2014) A conserved three-nucleotide core motif defines
Musashi RNA binding specificity. *J Biol Chem*, **289**, 35530-35541.

22.   Arkin, M.R. and Wells, J.A. (2004) Small-molecule inhibitors of protein–
protein interactions: progressing towards the dream. *Nature reviews Drug*
discovery, **3**, 301-317.

23.   Niederhauser, O., Mangold, M., Schubenel, R., Kusznir, E., Schmidt, D. and
Hertel, C. (2000) NGF ligand alters NGF signaling via p75NTR and TrkA.
Journal of neuroscience research, **61**, 263-272.

24.   Minuesa, G., Albanese, S.K., Xie, W., Kazansky, Y., Worroll, D., Chow, A.,
Schurer, A., Park, S.-M., Rotsides, C.Z. and Taggart, J. (2019) Small-molecule
targeting of MUSASHI RNA-binding activity in acute myeloid leukemia.
Nature communications, **10**, 1-15.

25.   Lello, L.S., Utt, A., Bartholomeeusen, K., Wang, S., Rausalu, K., Kendall, C.,
Coppens, S., Fragkoudis, R., Tuplin, A. and Alphey, L. (2020) Cross-utilisation
of template RNAs by alphavirus replicases. *PLoS pathogens*, **16**, e1008825.

26.   Roberts, G.C., Zothner, C., Remenyi, R., Merits, A., Stonehouse, N.J. and Harris,

M. (2017) Evaluation of a range of mammalian and mosquito cell lines for use

in Chikungunya virus research. *Scientific reports*, **7**, 1-13.

27.   Matkovic, R., Bernard, E., Fontanel, S., Eldin, P., Chazal, N., Hersi, D.H.,

Merits, A., Peloponese, J.M. and Briant, L. (2019) The Host DHX9 DExH-Box

Helicase Is Recruited to Chikungunya Virus Replication Complexes for Optimal

Genomic RNA Translation. *Journal of Virology*, **93**.

Figure Legends:

Figure 1. A) Schematic representation of CHIKV genome organisation **B)** Schematic

representation of CHIKV RNA structures within the 5'UTR and adjacent ORF-1 region

of the CHIKV genome (Kendall et al., 2019). RNA replication elements SL3, SL47,

SL88, SL102, SL165, SL194 and SL246 are labelled in black type. The ORF-1 AUG start

codon is labelled by a green arrow and the putative MSI binding site by a red oval.

Figure 2) Native EMSAs between *in vitro* transcribed P^{32} 5' radiolabeled CHIKV RNA

nts 1-330 (*RNA^{WT}) and recombinantly expressed MSI-2 demonstrated an

RNA/protein interaction that was outcompeted by increasing concentrations of

equivalent unlabeled (RNA^{WT}) but less efficiently by the same RNA incorporating

BSM mutation $_{63}AUUAAU_{68} >_{63}CAACUU_{68}$ (RNA^{BSM}). **A)** Increasing concentrations of

MSI-2 intensified the observed band shift to the larger RNA/Protein complex and

decreased the equivalent unbound RNA band. The interaction between a 1:4 ratio of

*RNA^{WT}:MSI-2 was competed with increasing concentrations of unlabelled **B)** RNA^{WT}

or **C)** RNA^{mut} ($_{63}CAACUU_{68}$-mut). **D)** Band shifts in the unlabeled RNA competition

EMSAs were quantified by densitometry and expressed as % change in the density of

the RNA/Protein complex bands, normalized to the equivalent total lane density, for

each competition ratio and compared each time to ratio 1:0. N=3, error bars

represent standard error from the mean and significance was measured by two-

tailed T-test (* = P < 0.05, ** = P < 0.01, *** = P < 0.001). Grey block triangles

indicate increasing concentrations of specific reactants.

Figure 3. Ro 08-2750 signifyingly inhibits replication of infectious CHIKV and the

CHIKV-SGR **A)** Schematic representations of CHIKV infectious clone (top) compared

to the sub-genomic replicon (SGR) (bottom) in which a *Renilla* luciferase (RLuc)

reporter gene is fused within the nsp3 coding sequence and the structural genes of

ORF-2 are replaced by a firefly luciferase (Fluc) reporter gene. Replication is

expressed in Relative Light Units [RLU] **B)** Ro significantly inhibits productive CHIKV

productive replication relative to DMSO treated negative controls at 8 and 24 hpi. **C)**

and **D**) Ro 08-2750 significantly inhibits CHIKV-SGR replication, measured by both

ORF-1 and ORF-2 expression, relative to DMSO treated negative controls at 8 and 24

hpt. **E)** Ro 08-2 750 had no significant (ns) effect on CHIKV-SGR (GDD-GAA)

translation, measured by ORF-1 expression. N=3, error bars represent standard error

from the mean and significance was measured by two-tailed T-test (* = $P < 0.05$, ** =

$P < 0.01$, *** = $P < 0.001$).

Figure 4. Ro signifyingly inhibits CHIKV genome replication. **A)** Schematic

representation of CHIKV *trans*-complementation assay showing codon optimised

pCHIKV-nsP1234 (top) from which the CHIKV nsPs were translated and pCHIK-

Fluc/Gluc (bottom) in which ORF-1 was replaced by an Fluc reporter gene, fused to

the first 77 nts or CHIKV ORF-1 (N77) downstream of the authentic CHIKV 5'UTR.

ORF-2, flanked by the authentic intragenic (SG) and 3' UTRs, was replaced by a Gluc

reporter gene. **B)** and **C)** Ro significantly inhibited CHIKV genome replication of the

trans-complementation assay, measured by both ORF-1 and ORF-2 expression,

relative to DMSO treated negative controls at 8 and 24 hpt. N=3, error bars

represent standard error from the mean and significance was measured by two-

tailed T-test (* = $P < 0.05$, ** = $P < 0.01$, *** = $P < 0.001$).

Figure 5. shRNA suppression of MSI-2 significantly inhibits replication of infectious

CHIKV. **A)** Western blot analysis of total cellular protein extracted from Rd cells and

compared to negative control scrambled shRNA, demonstrated consistent shRNA

knockdown of MSI-2 over 3 serial passages (P1- P3). MSI-2 suppression significantly

inhibited productive CHIKV replication, relative to scrambled shRNA at 8 and 24 hpi

measured by plaque assay **(B)** and strand specific qRT-PCR for the the virus genomic

(C) and negative intermediate **(D)** strands. N=3, error bars represent standard error

from the mean and significance was measured by two-tailed T-test (* = P < 0.05, ** =

P < 0.01, *** = P < 0.001).

Figure 6. shRNA suppression of MSI-2 significantly inhibits CHIKV-SGR replication and

CHIKV genome replication. **A)** and **B)** shRNA suppression of MSI-2 significantly

inhibited CHIKV-SGR replication, measured by both ORF-1 and ORF-2 expression,

relative to scrambled shRNA negative controls at 8 and 24 hpt. **C)** and **D)** shRNA

suppression of MSI-2 significantly inhibited CHIKV genome replication of the *trans-*

complementation assay, measured by both ORF-1 and ORF-2 expression, relative to

DMSO treated negative controls at 8 and 24 hpt. N=3, error bars represent standard

error from the mean and significance was measured by two-tailed T-test (* = P <

0.05, ** = P < 0.01, *** = P < 0.001).

Figure 7: siRNA Depletion of either MSI-1 or MSI-2 significantly inhibited CHIKV

replication in Huh7 cells and co-depletion of both MSI-1 and MSI-2 had a synergistic

effect on CHIKV inhibition in Huh7 cells. siRNA depletion of MSI-2 significantly

inhibited CHIKV replication in RD cells and co-depletion of both MSI-1 and MSI-2 did

not increase the level of CHIKV inhibition. siRNA depletion of MSI-1 and MSI-2

significantly inhibited CHIKV replication in Huh7 cells relative to scrambled siRNA at 8

and 24 hpi measured by plaque assay **(A)** and strand specific qRT-PCR for the the

virus genomic **(B)** and negative intermediate **(C)** strands. siRNA depletion of MSI-2 in

RD cells significantly inhibited CHIKV replication in relative to scrambled siRNA at 8

and 24 hpi measured by plaque assay **(D)** and strand specific qRT-PCR for the the

virus genomic **(E)** and negative intermediate **(F)** strands. N=3, error bars represent

standard error from the mean and significance was measured by two-tailed T-test (*

= P < 0.05, ** = P < 0.01, *** = P < 0.001).

Figure 8. Substitutions disrupting the predicted MSI binding site ($_{63}$CAACUU$_{68}$-mut)

but maintaining local RNA structure prevented virus rescue and significantly

inhibited CHIKV genome replication. Substitution $_A$67$_G$ maintained the predicted MSI-

2 binding motif but disrupted local RNA structure and significantly inhibited CHIKV

replication. **A)** and **B)** Mutation $_{63}$CAACUU$_{68}$ significantly inhibited CHIKV genome

replication of the *trans*-complementation assay, measured by both ORF-1 and ORF-2

expression, relative to wild-type positive controls at 8 and 24 hpt. **C)** Mutation

$_{63}$CAACUU$_{68}$-mut prevented rescue of CHIKV following transfection of capped *in vitro*

transcribed RNA into BHK cells. Released virus was measured by plaque assay of

supernatant 24 hpt and compared to positive control wild-type infectious CHIKV *in*

vitro transcribed RNA, which was transfected and analysed in parallel. **D)** Mutation

^{A}67^G significantly inhibited CHIKV replication at 8 hpi and 24 hpi relative to wild type

CHIKV following infection of RD cells (MOI 0.1). N=3, error bars represent standard

error from the mean and significance was measured by two-tailed T-test (* = P <

0.05, ** = P < 0.01, *** = P < 0.001).

Figure 9. Combining substitutions within the predicted MSI binding site

($_{63}$CAACUU$_{68}$-mut) and Ro 08-2750 had a synergistic affect, significantly increasing

inhibition of CHIKV genome replication. **A)** and **B)** Mutation of the MSI binding site

combined with Ro 08-2750 significantly increased inhibition of the CHIKV *trans*-

complementation assay, measured by both ORF-1 and ORF-2 expression, relative to

$_{63}$CAACUU$_{68}$-mut in the absence of Ro 08-2750 at 8 and 24 hpt. N=3, error bars

represent standard error from the mean and significance was measured by two-

tailed T-test (* = P < 0.05, ** = P < 0.01, *** = P < 0.001).

A:

ORF-1

ORF-2

5'UTR — | nsP1 | nsP2 | nsP3 | nsP4 | | C | E3 | E2 | 6K/TF | E1 | — 3'UTR- Poly[A]-3'

B:

A:

B:

CHIKV Titre [pfu/ml]

8h   24h

*** (8h), *** (24h)

C:

ORF1 [RLU]

8h   24h

** (8h), *** (24h)

D:

ORF2 [RLU]

8h   24h

** (8h), ** (24h)

E:

ORF1 [RLU]

ns

Untreated
DMSO
Ro 08-2750

6h

***

ORF2 [RLU]

Untreated
DMSO
Ro 08-2750

8h   24h

*** (8h), *** (24h)

A:

pCHIKV-nsP1234: IRES — | nsP1 | nsP2 | nsP3 | nsP4 | — Poly[A]-3'

pCHIKV-FLuc/Gluc: 5' UTR — | N77 | Fluc | | Gluc | — 3'UTR- Poly[A]-3'

B: **C:**

Untreated

DMSO

Ro

A:

B:

C:

D:

A: SGR assay — ORF1 [RLU] at 8h and 24h

B: SGR assay — ORF2 [RLU] at 8h and 24h

C: *trans*-complementation assay — ORF1 [RLU] at 8h and 24h

D: *trans*-complementation assay — ORF2 [RLU] at 8h and 24h

Legend:
- Mock
- Scrambled shRNA
- MSI-2 shRNA

Huh7 cells

A:

B:

C:

Mock

Scrambled

MSI1 siRNA

MSI2 siRNA

MS1+2 siRNA

RD cells

D:

E:

F:

MSI2 siRNA

MSI1+2 siRNA

A:

B:

– = Ro 08-2750 Absent
+ = Ro 08-2750 Present

Supplementary data 1: Alignment of CHIKV 5' region (nts 61-120) showing variation within 5'UTR MSI-2 binding site. All complete CHIKV genomes were downloaded from the NCBI, by using the selection for complete genomes and search term for CHIKV. Duplicates were removed, leaving one representative of each, which were aligned using CLUSTAL OMEGA (multiple sequence alignment) on EBI default settings. MSI-2 binding site indicated by orange box.

Accession number   61   120

```
gi|1032899617|gb|KX262992.1|   AGATTAATAAACCCATCATGGATTCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|1168909638|gb|KY703980.1|   AGATTAATAAACCCATCATGGATTCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|2316015176|gb|MZ703085.1|   AGATTAAGAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|1032899611|gb|KX262991.1|   AGATTAATAAACCCATCATGGATTCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|674268615|gb|KF590565.1|    AGATTAATAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|2322557117|gb|OK562602.1|   AGATTAAGAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|1781339200|gb|MK163628.1|   AGATTAATAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|262073953|gb|FJ513632.1|    AGATTAATAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCCGACAGCGCCTTTTT
gi|674268618|gb|KF590566.1|    AGATTAATAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|1050242909|gb|KT324228.1|   AGATTAATAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|655168280|gb|KJ796847.1|    AGATTAATAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|629510202|dbj|AB860301.3|   AGATTAATAAACCCATCATGGATTCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|429324006|emb|FN295483.3|   AGATCAATAAACCCATCATGGATTCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|1655475674|gb|MK690206.1|   AGATTAAGAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|410067116|emb|FR717336.1|   AGATTAATAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|2484235404|gb|OQ605447.1|   AGATTAATAAACCCATCATGGATCCTGTGCACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|1168909306|gb|KY703922.1|   AGATTAATAAACCCATCATGGATTCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|655168286|gb|KJ796850.1|    AGATTAATAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|1240405768|gb|MF499120.1|   AGATTAATAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|2186567288|gb|OL979153.1|   AGATTAATAAACCCATCATGGATCCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|27734686|gb|AF369024.2|     AGATTAAGAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|189409703|gb|EU703760.1|    AGATCAATAAACCCATCATGGATTCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|1189411909|gb|MF001514.1|   AGATTAATAAACCCATCATGGATTCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|2186567368|gb|OL999095.1|   AGATTAATAAACCCATCATGGATCCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|1391207960|gb|MG921596.1|   AGATTAATAAACCCATCATGGATTCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|615794500|gb|KJ451622.1|    AGATTAATAAACCCATCATGGATTCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|1189411921|gb|MF001518.1|   AGATTAATAAACCCATCATGGATTCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|1796551930|gb|MN402890.1|   AGATTAATAAACCCATCATGGATCCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|288572686|gb|FJ445427.2|    AGATTAATAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTCTT
gi|1967465569|gb|MW042254.1|   AGATTAATAAACCCATCATGGATCCTGTGTACGTGGACATAGACACTGACAGCGCCTTTTT
gi|1621066037|gb|MK473621.1|   AGATTAATAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|262410949|gb|FJ807897.1|    AGATTAATAAACCCATCATGGATTCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|2484235419|gb|OQ605452.1|   AGATTAACAAACCCATCATGGATCCTGTGCACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|2484235374|gb|OQ605437.1|   AGATTAACAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|262073950|gb|FJ513629.1|    AGATTAATAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|1677557230|gb|MH329296.1|   AGATTAATAAACCCATCATGGATTCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|745698469|gb|KP003807.1|    AGATTAATAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|564970356|gb|KC614648.1|    AGATTAATAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|2563904640|gb|OR037307.1|   AGATTAATAAACCCATCATGGATCCTGTGTACGTGGATGTAGACGCTGACAGCGCCTTTTT
gi|189409706|gb|EU703761.1|    AGATCAATAAACCCATCATGGATTCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|133779733|gb|EF452494.1|    AGATTAATAAACCCATCATGGATTCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|1021315749|gb|KT308159.1|   AGATTAATAAACCCATCATGGATTCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|1168909620|gb|KY703977.1|   AGATTAATAAACCCATCATGGATTCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|1189411906|gb|MF001513.1|   AGATTAATAAACCCATCATGGATTCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|1189411897|gb|MF001510.1|   AGATTAATAAACCCATCATGGATTCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|1189411903|gb|MF001512.1|   AGATTAATAAACCCATCATGGATTCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|1168909711|gb|KY703992.1|   AGATTAATAAACCCATCATGGATTCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|259014734|gb|FJ959103.1|    AGATTAATAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|1032899593|gb|KX262988.1|   AGATTAATAAACCCATCATGGATTCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|954465478|gb|KR046227.1|    AGATTAATAAACCCATCATGGATTCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|1168909588|gb|KY703972.1|   AGATTAATAAACCCATCATGGATTCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|1333945209|gb|MF773569.1|   AGATTAATAAACCCATCATGGATCTTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|1796551933|gb|MN402891.1|   AGATTAATAAACCCATCATGGATCCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|2329593809|gb|OL840906.1|   AGATTAATAAACCCATCATGGATCCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|1189411900|gb|MF001511.1|   AGATTAATAAACCCATCATGGATTCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|2489916086|gb|OP615961.1|   AGATTAATAAACCCATCATGGATTCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|1189411882|gb|MF001505.1|   AGATTAATAAACCCATCATGGATTCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|2620122017|gb|OR785139.1|   AGATTAATAAACCCATCATGGATTCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|1189411924|gb|MF001519.1|   AGATTAATAAACCCATCATGGATTCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|116047549|gb|DQ443544.2|    AGATTAATAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|2438228262|gb|ON262791.1|   AGATTAATAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|1621066058|gb|MK473628.1|   AGATTAATAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|124295579|gb|EF027135.1|    AGATTAATAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|171918919|gb|EU244823.2|    AGATTAATAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|221229058|gb|FJ000064.1|    AGATTAATAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|256403024|dbj|AB455493.1|   AGATTAATAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTCTT
```

```
gi|262410946|gb|FJ807896.1|    AGATTAATAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|1316030169|dbj|LC259094.1|  AGATTAAGAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|2563904643|gb|OR037308.1|   AGATTAATAAACCCATCATGGATCCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|2675378508|gb|PP319439.1|   AGATTAATAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|1021315758|gb|KT308162.1|   AGATTAATAAACCCATCATGGATTCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|189409700|gb|EU703759.1|    AGATCAATAAACCCATCATGGATTCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|1333945188|gb|MF773562.1|   AGATTAATAAACCCATCATGGATTCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|541987890|gb|KF318729.1|    AGATTAATAAACCCATCATGGATTCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|615794507|gb|KJ451624.1|    AGATTAATAAACCCATCATGGATTCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|1843512200|gb|MN974212.1|   AGATTAATAAACCCATCATGGATCCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|1967465572|gb|MW042255.1|   AGATTAATAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|1843512188|gb|MN974208.1|   AGATTAATAAACCCATCATGGATCCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|1843512170|gb|MN974223.1|   AGATTAATAAACCCATCATGGATCCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|2438228268|gb|ON262793.1|   AGATTAATAAACCCATCATGGATCCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|2563904646|gb|OR037309.1|   AGATTAATAAACCCATCATGGATCCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|1621066055|gb|MK473627.1|   AGATTAATAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|156751972|gb|EU037962.1|    AGATTAATAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|801165883|gb|KP164869.1|    AGATTAATAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|1160525264|gb|KY751908.1|   AGATTAATAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|296124507|gb|HM045794.1|    AGATTAATAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|664818218|gb|KJ941050.1|    AGATTAATAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|429324003|emb|FN295485.3|   AGATTAATAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|2322557111|gb|OK562600.1|   AGATTAAGAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|296124529|gb|HM045802.1|    AGATTAATAAACCCATCATGGATTCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|124295582|gb|EF027136.1|    AGATTAATAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|2186567357|gb|OL999093.1|   AGATTAATAAACCCATCATGGATCCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|2438257538|gb|OQ148650.1|   AGATTAAGAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|2219863227|gb|ON009843.1|   AGATTAAGAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|1240390999|gb|MF774617.1|   AGATTAATAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|296124489|gb|HM045788.1|    AGATTAATAAACCCATCATGGATTCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|2186567214|gb|OL705486.1|   AGATTAATAAACCCATCATGGATCCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|296124564|gb|HM045814.1|    AGATTAATAAACCCATCATGGATTCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|1621066052|gb|MK473626.1|   AGATTAATAAACCCATCATGGATTCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|1992350182|gb|MT666072.1|   AGATTAAGAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|1240390993|gb|MF774615.1|   AGATTAATAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|1240391002|gb|MF774618.1|   AGATTAATAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|1147165987|gb|KY038947.2|   AGATTAAGAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|1677557251|gb|MH329303.1|   AGATTAATAAACCCATCATGGATTCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|1168909471|gb|KY703951.1|   AGATTAATAAACCCATCATGGATTCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|2216793899|gb|MW281311.1|   AGATTAAGAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|1728385259|gb|MK935343.1|   AGATTAATAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|1677557239|gb|MH329299.1|   AGATTAATAAACCCATCATGGATTCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|1168909205|gb|KY703904.1|   AGATTAATAAACCCATCATGGATTCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|1032899626|gb|KX262993.1|   AGATTAATAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|2484235380|gb|OQ605439.1|   AGATTAACAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|1032899599|gb|KX262989.1|   AGATTAATAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|1798045089|gb|MK370032.1|   AGATTAATAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|2186567343|gb|OL999091.1|   AGATTAATAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|1621066043|gb|MK473623.1|   AGATTAATAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|2489916092|gb|OP615963.1|   AGATTAAGAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|2186567294|gb|OL979154.1|   AGATTAATAAACCCATCATGGATCCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|1992350185|gb|MT666073.1|   AGATTAAGAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|1626774529|gb|MH124576.1|   AGATTAAGAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|124295597|gb|EF027141.1|    AGATTAATAAACCCATCATGGATTCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|745698484|gb|KP003812.1|    AGATTAAGAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|1778598293|gb|MK040569.1|   AGATTAATAAACCCATCATGGATCCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|1240391005|gb|MF774619.1|   AGATTAATAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|1798045083|gb|MK370030.1|   AGATTAATAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|1316030139|dbj|LC259084.1|  AGATTAATAAACCCATCATGGATTCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|2118777267|gb|MZ443814.1|   AGATTAATAAACCCATCATGGATCCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|1861718970|gb|MT526804.1|   AGATTAATAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|1032899586|gb|KX262987.1|   AGATTAATAAACCCATCATGGATTCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|1240390988|gb|MF774613.1|   AGATTAATAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|1168909546|gb|KY703965.1|   AGATTAATAAACCCATCATGGATTCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|428670855|emb|HE806461.1|   AGATTAATAAACCCATCATGGATTCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|1685826244|gb|MH359141.1|   AGATTAATAAACCCATCATGGATTCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|2018315610|gb|MW574902.1|   AGATTAATAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|659902278|gb|KJ679578.1|    AGATTAATAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|2438257514|gb|OQ148642.1|   AGATTAAGAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|1279118184|gb|MG280943.1|   AGATTAAGAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|124295576|gb|EF027134.1|    AGATTAATAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|296124486|gb|HM045787.1|    AGATTAATAAACCCATCATGGATTCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|296124549|gb|HM045809.1|    AGATTAAGAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|296124501|gb|HM045792.1|    AGATTAAGAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|124295588|gb|EF027138.1|    AGATTAATAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|2186567333|gb|OL979157.1|   AGATTAATAAACCCATCATGGATCCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|1859767498|gb|MK993755.1|   AGATTAAGAAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
```

61   120

```
gi|1168907703|gb|KY680385.1|   AGATTAATAACCCATCATGGATTCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|1861718958|gb|MT526800.1|   AGATTAATAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|659902275|gb|KJ679577.1|    AGATTAAGAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|296124523|gb|HM045800.1|    AGATTAATAACCCATCATGGATTCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|262073965|gb|FJ513654.1|    AGATTAATAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|262073947|gb|FJ513628.1|    AGATTAATAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|296124495|gb|HM045790.1|    AGATTAATAACCCATCATGGATTCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|262073968|gb|FJ513657.1|    AGATTAATAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|296124541|gb|HM045806.1|    AGATTAAGAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|2484235365|gb|OQ605434.1|   AGATTAACAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|1859767502|gb|MK993757.1|   AGATTAAGAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|2577286133|gb|OR283254.1|   AGATTAAGAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|371985830|gb|JN558834.1|    AGATTAATAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|2484235386|gb|OQ605441.1|   AGATTAACAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|288572683|gb|FJ445426.2|    AGATTAATAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|1992350179|gb|MT666071.1|   AGATTAAGAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|288572704|gb|FJ445443.2|    AGATTAATAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|2674493701|gb|PP193832.1|   AGATTAATAACCCATCATGGATCCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|2322557114|gb|OK562601.1|   AGATTAAGAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|1565065272|gb|MK120202.1|   AGATTAATAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|1565065269|gb|MK120201.1|   AGATTAATAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|1559671006|gb|MH423806.1|   AGATTAATAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|2438257517|gb|OQ148643.1|   AGATTAAGAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|655168290|gb|KJ796852.1|    AGATTAATAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|2118777264|gb|MZ443813.1|   AGATTAATAACCCATCATGGATCCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|296124510|gb|HM045795.1|    AGATTAAGAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|124295594|gb|EF027140.1|    AGATTAATAACCCATCATGGATTCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|1168909139|gb|KY703893.1|   AGATTAATAACCCATCATGGATTCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|1171733075|gb|KX881784.1|   AGATTAATAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|1621066040|gb|MK473622.1|   AGATTAATAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|2484235389|gb|OQ605442.1|   AGATTAACAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|1032899645|gb|KX262996.1|   AGATTAAGAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|1565065260|gb|MK120198.1|   AGATTAATAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGATAGCGCCTTTTT
gi|1032899632|gb|KX262994.1|   AGATTAATAACCCATCATGGATTCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|1621066049|gb|MK473625.1|   AGATTAATAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|1154340972|gb|KY575574.1|   AGATTAAGAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|2438257478|gb|OQ148630.1|   AGATTAAGAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|1168909576|gb|KY703970.1|   AGATTAATAACCCATCATGGATTCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|1168909314|gb|KY703923.1|   AGATTAATAACCCATCATGGATTCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|2438731903|gb|MW588418.1|   AGATTAATAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|1677557221|gb|MH329293.1|   AGATTAATAACCCATCATGGATTCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|1032899652|gb|KX262997.1|   AGATTAATAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|1168909527|gb|KY703961.1|   AGATTAATAACCCATCATGGATTCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|655168284|gb|KJ796849.1|    AGATTAATAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|655168282|gb|KJ796848.1|    AGATTAATAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|1733482778|gb|MK551553.1|   AGATTAATAACCCATCATGGATCCCGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|2438257442|gb|OQ148618.1|   AGATTAATAACCCATCATGGATTCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|1199750720|gb|KX619425.1|   AGATTAATAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|1621066076|gb|MK473634.1|   AGATTAATAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|1733482772|gb|MK518340.1|   AGATTAAGAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|2484235422|gb|OQ605453.1|   AGATTAACAACCCATCATGGATCCTGTGCACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|1626774541|gb|MH124580.1|   AGATTAATAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|2715015475|gb|PP599025.1|   AGATTAATAACCCATCATGGATCCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|1983384050|gb|MT495605.1|   AGATTAATAACCCATCATGGATCCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|2438731973|gb|MW588419.1|   AGATTAATAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|2484235425|gb|OQ605454.1|   AGATTAACAACCCATCATGGATCCTGTGCACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|380467950|gb|JQ067624.1|    AGATTAAGAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|1954890103|gb|MT526903.1|   AGATTAAGAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|1199750723|gb|KX619426.1|   AGATTAATAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|1626774550|gb|MH124583.1|   AGATTAATAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|1626774547|gb|MH124582.1|   AGATTAATAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|2316015179|gb|MZ703086.1|   AGATTAAGAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|1733482775|gb|MK551552.1|   AGATTAATAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|1621066046|gb|MK473624.1|   AGATTAATAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|2186567348|gb|OL999092.1|   AGATTAATAACCCATCATGGATCCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|1168909278|gb|KY703917.1|   AGATCAATAACCCATCATGGATTCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|296124591|gb|HM045823.1|    AGATTAAGAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|296124477|gb|HM045784.1|    AGATTAAGAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|1778598299|gb|MK040571.1|   AGATTAATAACCCATCATGGATCCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|655168288|gb|KJ796851.1|    AGATTAATAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|280983623|gb|GU199350.1|    AGATTAATAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|1168909157|gb|KY703896.1|   AGATTAATAACCCATCATGGATTCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|2287617414|gb|MW557660.1|   AGATTAATAACCCATCATGGATCCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
gi|2154025537|gb|MW473668.1|   AGATTAATAACCCATCATGGATTCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|2484235401|gb|OQ605446.1|   AGATTAACAACCCATCATGGATCCTGTGTACGTGGACATAGACGCTGACAGCGCCTTTTT
gi|1168907836|gb|KY680408.1|   AGATTAACAACCCATCATGGATTCTGTGTACGTGGATATAGACGCTGACAGCGCCTTTTT
```

Supplementary data 2: Primers for the reverse transcription and quantitative PCRs for CHIKV strand-specific detection.

CHIKV (-) strand detection	PCR	Primer sequence (5'-3')
CHIKV FT tag T	reverse transcription	GGC AGT ATC GTG AAT TCG ATG CGA CAC GGA GAC GCC AAC ATT
Tag T	quantitative	GGC AGT ATC GTG AAT TCG ATG C
CHIKV R T	quantitative	AAT AAA TCA TAA GTC TGC TCT CTG TCT ACA TGA
CHIKV (+) strand detection	PCR	Primer sequence (5'-3')
CHIKV RT tag T	reverse transcription	GGC AGT ATC GTG AAT TCG ATG CGT CTG CTC TCT GTC TAC ATG A
CHIKV F T	quantitative	AAT AAA TCA TAA GAC ACG GAG ACG CCA ACA TT
Tag T	quantitative	see above

Supplementary data 3: Expression and purification of MSI-2 Coomassie stained PAGE analysis following **A)** His Tag and **B)** Ion exchange chromatography. **C)** Ion exchange purified MSI-2 (8-193) analyzed by western blot, relative to total protein extracted from RD and Huh7 cells.

Supplementary data 4: CHIKV predicted RNA structure (nts 1-300) in **A)** wild type **B)** mutant [63]CAACUU[68] and **C)** mutant [A]67[G]. RNA structure mapped free energy minimization using Pfold on default settings.

A) Wild type

Predicted MSI binding site

B) Mutant [63]CAACUU[68]

Predicted MSI binding site

Supplementary data 4: CHIKV predicted RNA structure (nts 1-300) in **A)** wild type **B) m**utant [63]CAACUU[68] and **C)** mutant [A]67[G]. RNA structure mapped free energy minimization using Pfold on default settings.

C) Mutant [A]67[G]

Predicted MSI binding site

Supplementary data 5: MTT cytotoxicity assay for Ro 08-2750 in RD cells across a titration of 0, 05, 1, 3, 5 10 and 20uM. N=3, error bars represent standard error from the mean.

Supplementary data 6: A) MSI-1 and **B**) MSI-2 expression in RD and Huh7 cell lysate analyzed by western blot.

Supplementary data 7: Co-inhibition of MSI-2 and MSI-1 by siRNA in **A)** and **B)** Huh7 cells and **C)** RD cells