Deciphering novel TCF4-driven mechanisms underlying a common triplet repeat

expansion-mediated disease

Short title (69/70 characters): Transcriptomic signatures of a common triplet

repeat disease

Authors

Nihar Bhattacharyya[1], Niuzheng Chai[1*], Nathaniel J Hafford-Tear[1*], Amanda N Sadan[1*], Anita

Szabo[1*], Christina Zarouchlioti[1], Jana Jedlickova[2], Szi Kay Leung[3], Tianyi Liao[1], Lubica

Dudakova[2], Pavlina Skalicka[2,4], Mohit Parekh[1], Ismail Moghul[1,5], Aaron R Jeffries[3], Michael E

Cheetham[1], Kirithika Muthusamy[5], Alison J Hardcastle[1,5], Nikolas Pontikos[1,5], Petra Liskova[2,4],

Stephen J Tuft[1,5], Alice E Davidson[1,5†]

Affiliations

[1] University College London Institute of Ophthalmology, London, ECIV 9EL, UK

[2] Department of Paediatrics and Inherited Metabolic Disorders, First Faculty of Medicine,

Charles University and General University Hospital in Prague, Prague, Czech Republic

[3] Faculty of Health and Life Sciences, University of Exeter, Exeter, EX4 4PY, UK

[4] Department of Ophthalmology, First Faculty of Medicine, Charles University and General

University Hospital in Prague, Prague, Czech Republic

[5] Moorfields Eye Hospital, London, EC1V 2PD, UK

[†] To whom correspondence should be addressed:

Dr Alice Davidson, UCL Institute of Ophthalmology, 11-43 Bath Street, London, UK

Email: alice.davidson@ucl.ac.uk

*These authors contributed equally to the manuscript

Keywords: TCF4 dysregulation, RNA toxicity, spliceopathy, triplet-repeat, repeat expansion-

mediated disease, corneal dystrophy, Fuchs endothelial corneal dystrophy, *TCF4,* long-read

sequencing, RNA-Seq, Iso-Seq, rare *TCF4* variants, isoform-specific dysregulation

Abstract: (Max 300 words)

(274/300)

Fuchs endothelial corneal dystrophy (FECD) is an age-related cause of vision loss, and the

most common repeat expansion-mediated disease in humans characterised to date. Up to 80%

of European FECD cases have been attributed to expansion of a non-coding CTG repeat

element (termed CTG18.1) located within the ubiquitously expressed transcription factor

encoding gene, *TCF4*. The non-coding nature of the repeat and the transcriptomic complexity of

TCF4 have made it extremely challenging to experimentally decipher the molecular

mechanisms underlying this disease. Here we comprehensively describe CTG18.1 expansion-

driven molecular components of disease within primary patient-derived corneal endothelial cells

(CECs), generated from a large cohort of individuals with CTG18.1-expanded (Exp+) and CTG

18.1-independent (Exp-) FECD. We employ long-read, short-read, and spatial transcriptomic

techniques to interrogate expansion-specific transcriptomic biomarkers. Interrogation of long-

read sequencing and alternative splicing analysis of short-read transcriptomic data together

reveals the global extent of altered splicing occurring within Exp+ FECD, and unique transcripts

associated with CTG18.1-expansions. Similarly, differential gene expression analysis highlights

the total transcriptomic consequences of Exp+ FECD within CECs. Furthermore, differential

exon usage, pathway enrichment and spatial transcriptomics reveal *TCF4* isoform ratio skewing

solely in Exp+ FECD with potential downstream functional consequences. Lastly, exome data

from 134 Exp- FECD cases identified rare (minor allele frequency <0.005) and potentially

deleterious (CADD>15) *TCF4* variants in 7/134 FECD Exp- cases, suggesting that *TCF4*

variants independent of CTG18.1 may increase FECD risk. In summary, our study supports the

hypothesis that at least two distinct pathogenic mechanisms, RNA toxicity and *TCF4* isoform-

specific dysregulation, both underpin the pathophysiology of FECD. We anticipate these data

will inform and guide the development of translational interventions for this common triplet-

repeat mediated disease.

53 **Author's summary:**

54

55 **200/200**

56 Fuchs endothelial corneal dystrophy (FECD) leads to vision loss and is the most common

57 repeat expansion-mediated disease characterised to date; most individuals with FECD harbour

58 a non-coding CTG repeat expansion within the gene *TCF4*. FECD represents an important

59 paradigm for other rare and devastating neurological repeat-mediated diseases, given its

60 relatively mild and tissue-specific nature. Intriguingly, despite *TCF4* being ubiquitously

61 expressed, individuals with FECD only experience corneal disease, and the biological reason

62 for this tissue-specificity remains elusive. Here, we use tissue from 31 individuals with FECD to

63 perform complementary long-read, short-read and spatial transcriptomic analyses to enhance

64 our understanding of mechanisms underpinning this disease. These data highlight that at least

65 two mechanisms, RNA toxicity and *TCF4* isoform dysregulation, underlie the disease state in

66 affected corneal cells. Furthermore, *TCF4* isoform skewing, with evidence of downregulation,

67 suggests this mechanism in part may explain the unique vulnerability of the cornea. In addition,

68 7/134 FECD expansion negative cases were identified to harbour rare and potentially

69 deleterious *TCF4* variants, further supporting the hypothesis that dysregulation of TCF4 may be

70 key to FECD pathophysiology. Biological insights presented here will guide the development of

71 personalised FECD therapies and may inform the development of repeat-expansion mediated

72 therapies more broadly.

Introduction:

Fuchs endothelial corneal dystrophy (FECD) is a common, age-related, and visually disabling disease that represents a leading indication for corneal transplantation in high-income countries) (1,2). It primarily affects the corneal endothelium, a monolayer of post-mitotic cells on the posterior corneal surface that function to maintain relative corneal dehydration. As the disease progresses, collagenous excrescences (guttae) form on the thickened corneal endothelial basement membrane (Descemet membrane) and corneal endothelial cells (CECs) undergo accelerated cell death, which results in corneal oedema and opacity (3). Affected individuals typically experience symptoms in their 5^{th} to 6^{th} decade, with glare, reduced contrast sensitivity, and blurred vision (4). Transplantation of donor endothelial tissue is the preferred treatment for cases with visually significant symptoms. However, given an increasing societal burden of age-related disease coupled with a global shortage of suitable donor tissue, there is a pressing clinical need for alternative treatments (3,5).

Up to 80% of FECD cases of European descent have one or more expanded copies of a CTG trinucleotide repeat element (termed CTG18.1; MIM #613267) located within an intronic region of the transcription factor gene 4, *TCF4* (4,6). We have previously demonstrated, in our large (n=450) FECD cohort, that an expanded copy of this CTG repeat (defined as ≥50 copies) confers >76-fold increased risk for developing FECD (7). Expansion of non-coding repeat elements within the human genome have been identified to cause more than 30 human diseases to date (8), including fragile X tremor ataxia syndrome (MIM #300623), myotonic dystrophy types 1 and 2 (MIM #160900 and #602668) and *C9orf72*-associated amyotrophic lateral sclerosis (ALS) and frontotemporal dementia (MIM #105550)(8). Expansion-positive (Exp+) FECD represents by far the most common disease within this category. However, to date, it has only been established that it affects the cornea, which is in contrast to the majority of the other much rarer non-coding repeat expansion disorders that typically have devastating

outcomes on neurological and/or neuromuscular systems. Despite these differences, strong

mechanistic molecular parallels exist between CTG18.1-mediated FECD and other non-coding

repeat expansion diseases, whereby multiple pathogenic mechanisms occur concurrently to

drive pathology (8,9).

We, and others, have demonstrated that RNA toxicity comprising splicing dysregulation

attributed to the accumulation of CUG-containing RNA foci and sequestration of splicing factors

MBNL1 and MBNL2 within affected cells, is a key feature of FECD in the presence of one or

more expanded CTG18.1 alleles (6,7,10–12). Alongside this RNA induced toxicity, repeat-

associated non-AUG (RAN) dependent translation of the expanded repeat, and dysregulation of

TCF4 itself have also been hypothesised to contribute to the underlying pathology of CTG18.1

Exp+ mediated FECD(13). However, a comprehensive understanding of consequences and role

of both the spliceopathy and downstream effects of the *TCF4* dysregulation have yet to be

characterised in parallel in the same model system. Furthermore, because *TCF4* comprises >90

alternative spatially and temporally expressed isoforms, deciphering transcript-specific level

dysregulation has proven to be extremely challenging and a consistent pattern of *TCF4*

dysregulation has yet to be associated with CTG18.1 expansions in corneal endothelial cells

(CECs) (14–16). Such diversity is reflected in the varied functional roles TCF4 plays in human

development, alongside the range of human disease associated with *TCF4* variants in addition

to FECD. For example, *TCF4* haploinsufficiency results in Pitt-Hopkins syndrome (MIM

#610954) (17,18), a severe neurodevelopmental disorder, whereas somatic *TCF4* mutations are

commonly identified in Sonic hedgehog medulloblastoma (19) and common germline variants

have been associated with Schizophrenia risk (20,21).

In this study, we used a large cohort of biologically independent primary FECD case-

derived CECs to isolate the effects of the multiple parallel pathogenic processes underlying

CTG18.1-mediated FECD. We present long-read transcriptomic data to enhance our

understanding of the increased isoform diversity observed in Exp+ FECD, and spatial

transcriptomics to demonstrate *TCF4*-isoform dysregulation. For the first time, we also isolate

downstream transcriptomic signatures of dysregulation that we hypothesise to be driven by

TCF4-isoform dysregulation that may, in part, explain the tissue specificity of the disease. We

go on to explore the potential relevance of TCF4 dysregulation to FECD, independent of the

CTG18.1 expansion, by interrogating expansion-negative FECD cases for rare *TCF4* variants

through exome sequencing and subsequently applying a gene-burden style approach.

Results

RNA-seq data from CECs reveal CTG18.1 expansion-mediated FECD to be

transcriptomically distinct from non-expanded FECD

To further increase our understanding of CTG18.1 expansion-associated splicing dysregulation

events, and better understand the relationship between individual events to transcript-specific

isoforms, we generated long-read RNA-seq data from biologically independent unpassaged

primary CEC cultures derived from healthy control (n=4, mean age 68.75 years) and Exp+

FECD (mono-allelic CTG18.1 allele repeat length ≥50) (n=4, mean age 66.8 years) tissues

(**Table 1, S1**)**.** After alignment and stringent quality control, reads that displayed evidence of 5'

degradation, intrapriming and reverse transcription template switching were removed. Exp+ and

control samples were subsequently analysed with SQANTI3 to assess isoform diversity (**Figure**

1A, B and Table 2, S2). We also generated short-read RNA-seq data from unpassaged primary

CEC cultures derived from healthy control (n=4, mean age 58.5 years), Exp+ FECD (mono-

allelic CTG18.1 allele repeat length ≥50) (n=3, mean age 63.3 years) and Exp- FECD (bi-allelic

CTG18.1 allele repeat lengths ≤30) (n=3, mean age 63.3 years) tissues (**Table 1**). Exome data

generated from the Exp- FECD cases utilised in this transcriptomic work were analysed for the

presence of variants in any previously FECD-associated genes, such as *TCF4, COL8A2,*

SLC4A11, ZEB1, and *AGBL1*(4), to identify any potential underlying genetic cause. No rare

(MAF <0.005) or potentially deleterious (CADD >15) variants were identified. Hierarchical

clustering using heatmap visualisation and PCA analysis including sex and ethnicity as

covariates revealed that Exp+ samples grouped separately from not only the controls, but also

the Exp- samples **(Figure 1C, D)**. Driving genes and additional principal components are listed

in Supplemental Table S2. Notably, Exp- samples also clustered together on the PCA plot, but

the heatmap derived from hierarchical clustering highlighted that greater intra-group variability

existed between Exp- samples compared to variability within control and Exp+ groups (**Figure**

1C). We hypothesise that the increased transcriptomic diversity within the Exp- group is due to

potentially distinct, as yet unidentified, genetic causes driving disease for each FECD case-

derived sample. We included all three groups in the downstream comparative pipelines using

the following pairwise comparisons; **(PWC1)** control versus Exp+ FECD, **(PWC2)** Exp+ FECD

versus Exp- FECD, and **(PWC3)** control versus Exp- FECD.

**Table 1: Summary of FECD case and control-derived corneal endothelial cell (CEC)
cultures analysed by short and long-read RNA-seq, and spatial transcriptomics.**

Sample	Sex	Age*	Ethnicity[a]/Ancestry[b]
Long-read			
Control 1	Female	68	White-European[a]
Control 2	Male	64	White-European[a]
Control 3	Female	74	White-European[a]
Control 4	Male	69	White-European[a]
FECD1exp+ (43/90)	Female	74	White-European[a,b]
FECD2exp+ (14/80)	Male	78	White-European[a,b]
FECD3exp+ (18/86)	Female	70	White-European[a,b]
FECD4exp+ (26/78)	Female	45	White-European[b]
Short-read			
Control 1	Female	68	White-European[a]
Control 5 Control 6	Male	55	White-European[a]
Control 7	Female	56	Black-African[a]
FECD5exp+ (16/86)	Female	67	White-European[a,b]
FECD6exp+ (12/106)	Male	69	White-European[a,b]
FECD7exp+ (16/94)	Female	54	White-European[a,b]
FECD1exp- (13/18)	Female	64	European/Black-Carribbean[a] Black-African[b]
FECD2exp- (18/24)	Female	73	White-European[a,b]
FECD3exp- (22/28)	Male	53	Black-African[a,b]
Spatial transcriptomics			
Control 8 (12/16)	Female	45	Black-African[a]
Control 9 (15/33)	Female	46	White-European[a]
Control 10 (25/26)	Male	70	White-European[a]
FECD4exp- (15/15)	Female	48	White-European[a]
FECD5exp- (16/23)	Female	57	Black-African[a]
FECD6exp- (15/16)	Female	58	Black-African[a]
FECD8exp+ (63/88)	Female	75	White-European[a]
FECD9exp+ (12/76)	Male	61	Black-African[a]
FECD10exp+ (18/67)	Male	65	White-European[a]

CTG18.1 expansion status is denoted in brackets for all FECD-case derived CEC lines analysed. *Age at time of corneal tissue excision. [a] self-reported by affected individual. [b] ancestry determined by applying a principal component analysis (PCA) method, FRAPOSA (22). All control samples were derived from donor corneas with very high CEC counts (≥2,600 cells/mm2) and endothelium free of guttae or any other abnormalities.

Figure 1: Exp+ FECD samples are transcriptomically distinct in both short- and long-read datasets when compared to controls

A) Bar chart showing proportion of annotated genes in long read transcriptomic dataset with only one transcript detect vs 2 or more transcripts in Control and Exp+ samples. Exp+ long-read samples have a higher proportion of genes with more than one isoform identified. **B)** Proportion of annotated transcripts with full splice matches. Exp+ FECD and Control samples have similar percentages of full splice matches suggesting additional splicing diversity seen in Exp+ is not predominantly due to aberrant splicing. **C)** Hierarchical clustering heatmap of sample-to-sample distances within short-read RNA-Seq data. Darker colours denote higher similarities between samples, demonstrating that Exp+ and control are more similar within each subgroup than with other samples. Exp- shows a similar result but is less pronounced, likely due to differences in unidentified underlying genetic causes. **D)** Principal component plot (PCA) of all short-read samples with sex, disease state and ethnicity as covariates. All three subsets of data cluster separately in the PCA plot. Control samples appear as circles, FECD Exp+ appear as orange triangles, FECD Exp- appear as green triangles. Male individuals are represented as filled shapes while female individuals are outlined.

Alternative splicing analysis of long-read and short-read transcriptomic data

demonstrate the complexity of increased splicing events underlying CTG18.1 expansion-

mediated disease in CECs

Iso-seq and subsequent SQANTI3 analysis (**Table 2, S3**) revealed a higher proportion of

isoforms per gene in Exp+ samples (1.7 unique isoforms per unique gene in Exp+ vs 1.3 in

control). Additionally, we see a higher proportion of known non-canonical splice junctions in

Exp+ samples. When characterising transcripts based on splice junctions, we see similar

proportions of isoform subtypes (**Table S3**). This is consistent with previous work (6,10–12) and

validated by our short read data, discussed below, (**Figure S1, S2**), whereby Exp+ samples

showed greater isoform diversity compared to control and Exp- samples, but not an overall

skew in the types of alternative splicing observed.

206
207
Table 2: Summary of SQANTI3 analysis of IsoSeq data. In-depth characterisation of isoforms
in control and Exp+ long-read RNA-seq data utilising SQANTI3 pipeline following stringent
quality control. SQANTI3 subdivides splice junctions into known (present in reference
transcriptome) and novel (not in reference transcriptome). Canonical splice junctions refer to the
most common type of splice junction (GTAG, GCAG, ATAC)), while non-canonical refers to
other rarer and tissue specific splice junctions with different donor and acceptor sites. SQANTI3
defines novel as genes or splice junctions that are absent from the reference transcriptome.

	Control (n=4)		**Exp+** FECD (n=3)	
Genes	2,318		4,652	
Isoforms	3,113		8,032	
Gene Classification				
	# of genes		# of genes	
Annotated Genes	2,309		4,642	
Novel Genes	9		10	
Splice Junction Classification				
	# splice junctions	Percent	# splice junctions	Percent
Known canonical	27,552	97.18	52,153	97.35
Known non-canonical	1	0.00001	21	0.04
Novel canonical	798	2.81	1,400	2.61
Novel non-canonical	0	0	0	0

To further characterise and quantify alternative splicing events within our short-read

dataset, we utilised rMATS, a junction-based method that employs a hierarchical framework, to

model unpaired datasets and evaluate and quantify all major alternative splice types (23–25).

Comparing Exp+ to control samples (PWC1) we identified 2,118 significant alternative splicing

events in 1,437 genes (**Figures S1, S2, Table S4)**. Alternative splicing comparisons including

Exp+ (PWC1 and PWC2) yielded the most alternative splice types in every category (**Figure S1,**

S2) further reinforcing the spliceopathy aspects of this genetic subtype of disease.

We then compared the rMATS and Iso-seq results to a subset of published datasets

exploring differential splicing in FECD samples (**Table S5**) and selected 24 published differential

splicing events (from 23 genes) that had previously been identified by bioinformatic or *in vitro*

methods to be strongly associated with CTG18.1 expansion-mediated FECD (4,6,7). Alternative

splicing analysis of our transcriptomic dataset identified that 75.0% (18/24) of these events were

significant in PWC1 (control vs Exp+, **Table S5**). In addition to these 24 selected events, rMATS

identified another 10 novel and significant differential splicing events within Exp+ in these

genes, highlighting the increasing complexity of splicing dysregulation within Exp+ FECD. Long-

read sequencing confirmed the presence of these novel splicing events in Exp+ FECD (**Table**

S6). Also, it was interesting to note that *FN1* was identified to have robust alternative splicing in

Exp+, and further investigation showed that alternatively spliced exons in Exp+ corresponded

with skipping of the Extra Domain A (EDA) and Extra Domain B (EDB) protein domains that play

a critical role in signal transduction and fibrillar formation (26) (**Table S7**).

Skipped exon alternative splicing events, where an additional exon is either included or

excluded, were the most common in all three pairwise comparisons (**Figure S1**). Prior work

regarding alternative splicing in FECD has focused on MBNL1/2 sequestration(6,7,27) as MBNL

proteins are key regulators of exon splicing(28,29). Given their complex role, we attempted to

determine if the flanking intronic sequences of alternatively spliced exons were enriched for

common RNA-binding protein (RBP) binding motifs. We focussed our RBP motif enrichment

analysis (rMAPS2(30)) on the 250bp region upstream of alternatively spliced exons and isolated

those areas where there was enrichment in PWC1 and PWC2, but not PWC3 (i.e. events

enriched within Exp+ CEC lines). By filtering rMAPS results in this manner, a trend whereby

skipped exons are significantly ($p < 0.05$) upregulated by RNA-binding proteins with TT-

containing binding motifs and downregulated by RNA-binding proteins with CG-containing

binding motifs was observed (**Table S8**). As MBNL proteins favour YCGY binding motifs, this

suggests that other RNA-binding proteins with TT-containing motifs could also play a role in

driving the spliceopathy underpinning RNA toxicity, a key mechanism of disease in Exp+ FECD.

Evidence of TCF4 dysregulation appears enriched in CTG18.1-mediated FECD CECs,

highlighting specific candidate biomarkers for this genetic subtype of disease

Differential gene expression was assessed with DESeq2(31) in all three groups including

sex and ethnicity as covariates. For all comparisons an FDR corrected P-value (padj) of <0.05

was used as a threshold for significance (**Table S9**). A total of 4,288 genes were identified to be

differentially expressed between the Exp+ and controls (PWC1; **Figure 2A**). Of these, 2,089

genes had a shrunkLFC value (moderated $\log_2$fold values of expression accounting for high

dispersion and low expression counts) of greater than 1 or less than -1. Overall, PWC2 and

PWC3 groups resulted in lowered differential gene expression, with 1,695 and 1,911

significantly differentially expressed genes, respectively **(Figure 2A; Table S9)**.

Pairwise analysis revealed considerable overlap between dysregulated genes identified

by all three pairwise comparisons **(Figure 2A),** however, highest convergence was observed by

PWC1 (Exp+ versus control) and PWC2 (Exp+ v Exp-), suggesting these events are driven by

CTG18.1 expansions. In total, both pairwise comparisons reveal 1,028 dysregulated genes

uniquely and significantly dysregulated in Exp+ FECD case-derived samples. Conversely, by

comparing the overlap between PWC1 and PWC3 (Exp+ vs control and Exp- vs control) we

identified a further 873 genes that are dysregulated in FECD, irrespective of the underlying

genetic cause, and thus representing more ubiquitous signatures of the disease. Dysregulated

genes in PWC1, including Exp+ unique and common to FECD, did not skew directionally and

were evenly up or down-regulated within our analysis (**Figure 2C**). We isolated 3,292 genes

that are uniquely dysregulated in Exp+ FECD, 2,264 genes dysregulated solely in PWC1 and a

further 1,028 dysregulated genes shared with PWC2; (highlighted in orange **Figure 2A**) and

proceeded with downstream analysis (**Table S9**).

We conducted enrichment analysis using g:profiler (32) (querying gene ontology (GO),

KEGG pathway, and Reactome pathways). GO term enrichment and KEGG pathway analysis of

PWC1 dysregulated genes was consistent with previously published transcriptomes (6,11,12),

with several enriched terms and pathways relevant to RNA toxicity (i.e. GO:0000166 etc) (**Table**

S10). While total *TCF4* gene expression was not significantly downregulated in Exp+ (**Figure**

2B), intriguingly, we observed strong parallels to pathway enrichment analyses highlighted from

a study of *TCF4* neural progenitor knockdown cell model (33) within the 3,292 genes uniquely

dysregulated in Exp+, such as protein processing in endoplasmic reticulum, MAPK signalling,

actin cytoskeletal regulation, endocytosis and apoptosis (**Figure 2D**). Critically, these pathways

are not enriched in PWC3, suggesting that these are unique to Exp+ CECs. Furthermore,

several enriched pathways and GO terms involving Wnt signalling were seen in the 3,292

dysregulated genes unique to Exp+ that are not present in PWC3, again suggesting *TCF4*

function may be modified within Exp+ FECD CECs (**Table S10**). Furthermore, enrichment of

dysregulated genes in pathways involving other repeat expansion-mediated diseases and

misfolded protein aggregate diseases suggest mechanistic parallels which reinforce Exp+ CECs

as a model system with broad relevance to repeatome biology (**Table S10**).

The consistent presence of PI3K-Akt pathway enrichment within KEGG results in Exp+

dysregulated genes (and its consistent absence in PWC3 pathway enrichment) suggests that

Exp+ FECD CECs may interact with growth factors and the extracellular matrix in a unique

manner. Hence, components of the PI3K-Akt pathway may serve as informative biomarkers for

Exp+ FECD in future clinical trial settings, and thus warrant further investigation. Additional

pathways are enriched in this set of genes (and not in PWC3) that involve RNA processing and

binding, other repeat-mediated disease related mechanisms, epithelial to mesenchymal

transition, and multiple signal transduction cascades with extracellular matrix interactions (**Table**

S10).

300   To determine if other RNA-binding proteins may be implicated in the alternative splicing

seen in Exp+ FECD, we cross-referenced our short-read data with a list of ~1500 human RNA-

binding proteins (34), and discovered that there were 141 RBPs uniquely dysregulated in Exp+

(45 downregulated and 106 upregulated), including *MBNL1, CELF1, ESRP2,* and a variety of

hnRNP and DDX encoding genes (**Table S11**). Both the hnRNP and DDX family of RNA-

binding proteins have previously been implicated in other repeat expansion diseases (32,35–

37). It is likely that a combination of these dysregulated RBPs plays a role in driving the

alternative splicing and RNA toxicity seen in Exp+ FECD.

Figure 2: Differential gene expression analysis on Exp+ FECD, Exp- FECD, and control primary corneal endothelial cell (CEC) transcriptomes. **A)** Differential gene expression was assessed via DESeq2. Three pairwise comparisons were conducted: PWC1 (Exp+ vs control), PWC2 (Exp+ vs Exp-), and PWC3 (Exp- vs control). Venn diagrams show overlap of significantly differentially expressed genes (adjusted p-value < 0.05) within these three comparisons. **B)** Total *TCF4* expression showed no significant dysregulation between the different groups. **C)** Volcano plot shows the distribution of gene expression in Exp+ FECD compared to the control group. Horizontal dotted lines denote the boundary between significance (adjusted p-value < 0.05), whereas vertical dotted lines show a shrunkLFC cutoffs for -1 and 1. Dark grey solid markers show significantly expressed genes with a fold change magnitude greater than 1. Solid orange markers delineate uniquely Exp+ differentially

expressed genes. **D)** Pathways enriched in Exp+ vs Control (g:profiler). Shaded grey boxes show direction of pathway enrichment in TCF4 knockdown models (Doostparast Torshizi et al. 2019). Green dots show pathways that are significantly enriched in PWC3 as well. Solid orange dots and open orange circle show that pathway enrichment persists in genes unique to Exp+ (1,028 genes and 3,292 genes, respectively)

Differential TCF4 exon usage suggests isoform ratio disruption of TCF4 in CECs of CTG18.1-mediated FECD individuals

Given signatures of *TCF4* dysregulation were apparent in the pathway enrichment analysis of Exp+ CECs (**Figure 2D**), we then aimed to determine if any tangible patterns of *TCF4* transcript-specific dysregulation was observed in the Exp+ CECs, despite total *TCF4* levels not being significantly different between Exp+, Exp- and control CECs **(Figure 2B).** Due to transcriptional diversity and complexity of TCF4 transcripts, DEXSeq (38) was used to assess differential exon usage utilising our short-read transcriptomic data.

DEXSeq utilises generalised linear models to detect differential exon usage while controlling for biological variation. This analysis allowed us to query gene expression and splicing at the exon level comparing PWC1 (with and without (PWC1_nocov) sex and ethnicity as covariates), PWC2, PWC3, FECD compared to control (FvC) and Exp+ compared to all expansion negative samples (Exp+ v Ctr+Exp-). In concordance with our rMATs and DESeq2 analysis, higher levels of transcriptome-wide differential exon usage events were observed in PWC1 compared to PWC3 (**Figure S3**), suggesting these patterns of differential exon usage are likely attributed to CTG18.1-related pathology.

Our short-read transcriptomic data reveals that multiple *TCF4* exons (n=5) were significantly dysregulated in Exp+ compared to control and Exp- FECD samples (**Figure 3A, Table S12 and S13**), whereas no significant differences were observed between Exp- and control (PWC3, **Figure 3B, Table S12 and S13**), demonstrating for the first time that differential *TCF4* exon usage is driven by CTG18.1 expansions and not a downstream feature of FECD unrelated to the primary genetic driver. A subset of exons contained within a small group of

TCF4 transcripts (4 out of 93) are upregulated in Exp+, while an increased number of more

broadly expressed exons are downregulated (appearing in 56 out of 93 transcripts) (**Figure 3A,**

Table S12 and S13). Transcripts containing downregulated exons are primarily included in

longer isoforms of the gene which contain 3 AD domains, while the upregulated exons

(excluding those surrounding the repeat) appear in shorter *TCF4* isoforms containing only two

AD domains. This suggests that Exp+ CECs may be employing certain compensatory measures

in response to the transcript-specific dysregulation that is induced by the presence of CTG18.1

repeat expansions. Interestingly, our rMATS analysis independently detects this same

phenomenon (**Table S14**), which showed a significant mutually exclusive exon splicing event

whereby the last exon in common with most 3 AD domain-containing isoforms (Exon 6 in

ENST000000354452.8) was favoured in the control group, while the first exon in common with

all isoforms (Exon 7 in ENST000000354452.8) was favoured in the Exp+ group.

We also detect retention of the intron flanking the CTG18.1 repeat, previously described

by Sznajder et al. 2018 (39) in Exp+ FECD **(Table S12 and S13).** The intronic region is

significantly upregulated in Exp+ CECs (E173 and E174) **(Table S12 and S13)**. While this

intron represents the 5' untranslated region (UTR) of several transcripts of *TCF4*, it is also

included within an intronic region in 32 out of 93 currently annotated *TCF4* transcripts.

367
368
Figure 3: CTG18.1 expansion-mediated FECD results in alternative *TCF4* exon usage
within corneal endothelial cells (CECs). A) Differences in exon usage between control (grey
dotted line) and Exp+ (orange dotted line) CECs. Significantly upregulated and downregulated
exons are highlighted with blue and red vertical lines, respectively. The location of CTG18.1 is
depicted with an orange line. Inset boxes magnify dysregulated regions. A genomic schematic
of *TCF4* with transcripts aligned is presented below the axis. Key protein domains highlighted
(green - basic helix loop helix (bHLH) domain, blue - activation domains (ADs), red - main
nuclear localization signal (NLS). **B)** Differences in exon usage between control (grey dotted
line) and Exp- (green dotted line) CECs. No exons were statistically dysregulated.

378
379
380
381

382 ***RNAScope shows robust detection of altered tissue-specific TCF4 isoform expression in***

383 ***CTG18.1-mediated FECD case-derived corneal endothelial cells***

384

385   We performed spatial transcriptomics, via RNAScope, to further validate our hypothesis

386 that a subset of *TCF4* isoforms containing the third AD3 domain are being selectively

387 downregulated due to the presence of an expanded CTG18.1 allele (≥50 repeats). For the first

388 time, we visualised *TCF4* transcripts within FECD and control CECs and fibroblasts using two

389 distinct sets of probes, termed A and B (**Figure 4A**). Probe set A comprises a set of probes that

390 allow the visualisation of all *TCF4* transcripts in the cell (probe A covers all exons in the

391 transcript Isoform B+ which contains 3 AD domains, TCF4-201, ENST00000354452.8,

392 NM_001083962.2). Probe set B targets exons in the canonical longer TCF4-201 *TCF4*

393 transcript which do not overlap with the canonical shorter *TCF4* transcript (Isoform A+/J which

394 contains 2 AD domains, AD1 and AD2, TCF4-204, ENST00000457482.7, NM_001243234.2).

395 Hence, these probe sets enable the detection, visualisation, and quantification of the ratio of

396 longer AD3-containing transcripts, previously identified to be downregulated in Exp+ CECs via

397 our differential exon usage analysis, compared to total *TCF4* transcripts (**Figure 3A**).

398   The *TCF4* transcripts were visualised in control, Exp+ and Exp- FECD case-derived

399 CECs (n=3 for each group) (**Figure 4B, Table 1, S1 and S15)**. Each independent primary CEC

400 culture was also probed with a negative control (targeting bacterial genes) and a $(CAG)_7$-Cy3

401 probe, using a standard FISH protocol, to validate the presence or absence of CUG-specific

402 RNA foci derived from expanded copies of CTG18.1 (**Table S15**). RNA foci were consistently

403 detected in all CTG18.1 Exp+ CECs, and were absent from control and Exp- CECs. Using

404 CellProfiler (version 4), puncta representing *TCF4* transcripts, detected with both probe set A

405 and B, were quantified and puncta/cell values were established for each image and averaged

406 across images with at least 100 nuclei imaged per condition per line. Finally, a ratio was

407 calculated establishing the proportion of longer AD3 containing transcripts compared to total

TCF4 transcripts for each biological replicate. This showed that, on average, Exp+ CECs had a

significantly smaller proportion of the longer AD3-containing transcripts (16.33% ± 4.53)

compared to unaffected control CECs (36% ± 6.9, p=0.0145) and non-expanded FECD CECs

(30.9% ± 5.17, p=0.0213) (**Figure 4C**). To explore potential cell-type specificity of the observed

skewed *TCF4* transcript ratios we also analysed a series of dermal fibroblast lines generated

from unrelated Exp+ patients with FECD (n=6) (7) and control adult dermal fibroblast lines (n=3)

(Table S16). No significant difference (p=0.61) was observed between the Exp+ (44.2% ± 5.68)

and control fibroblast (41.17% ± 12.1) lines, suggesting the dysregulation of *TCF4* is cell-type

specific. This result suggests that the presence of the expanded repeat alone consistently

skews *TCF4* transcript ratios to favour shorter isoforms, specifically in CECs. As this subgroup

of isoforms typically contains only two out of three activation domains, a higher prevalence of

shorter TCF4 isoforms may affect potential dimerization and TCF4 transcriptional function in

Exp+ CECs.

423
Figure 4: RNAScope probes targeting *TCF4* transcripts demonstrate that longer
transcripts containing activation domain 3 (AD3) account for a smaller proportion of total
transcripts specifically in Exp+ FECD corneal endothelial cells (CECs). A) Probe targeting
schematic for *TCF4*. Probe A targets all exons in NM_001083962.2 and visualises total *TCF4*
transcripts, while Probe B targets only the 5' exons, excluding those that are contained within
shorter transcripts (NM_001243234.2). Probe B visualises the longer subset of AD3-containing
transcripts within the cell. CTG18.1 is highlighted in orange. Green bar denotes TCF4 bHLH
region, blue regions highlight TCF4 activation domains, red box shows the bipartite TCF4 NLS
signal and orange shows the genomic region containing the CTG18.1 repeat. Key domains
highlighted (green - basic helix loop helix (bHLH) domain, blue - activation domains (ADs), red -
main nuclear localization signal (NLS)). **B)** Representative confocal images of Exp+, Exp- and
control CECs and Exp+ and control fibroblasts with nuclei (blue, DAPI) and *TCF4* puncta
(visualised with Cy3 but represented here with white). Scale bars represent 20 μm. **C)** Plotting
the proportion of longer AD3-containing *TCF4* transcripts in Exp+ (orange), control (black), and

Exp- (green) shows that Exp+ have significantly lower proportion of AD3-containing transcripts
compared to both control (p = 0.0145) and Exp- (p = 0.0213) CECs, while there is no significant
difference between control and Exp- (p = 0.362). D) Plotting the proportion of longer AD3-
containing *TCF4* transcripts in Exp+ (orange), and control (black) shows that there is no
statistically significant (p=0.61) difference in *TCF4* isoform ratios between the two groups. Open
circle denotes mean and error bars show standard deviation. Diamond datapoint shows bi-allelic
444   expanded FECD CECs. Significance determined with student T-test.

Rare TCF4 variants identified within a genetically refined CTG18.1 expansion-negative

FECD cohort suggest isoform-specific TCF4 dysregulation may be a risk factors for

FECD in the absence of CTG18.1 expansions

We next wanted to explore if rare and potentially deleterious *TCF4* variants could be

contributing to disease in FECD cases without a CTG18.1 expansion, given that Exp+ CECs

display alternative *TCF4* exon usage and skewed ratios of *TCF4* isoforms (**Figure 3 and 4**),

with detectable downstream features of TCF4-driven dysregulation (**Figure 2D**). To test this

hypothesis, we interrogated exome data generated from 134 Exp-FECD cases, acquired from

our ongoing inherited corneal disease research program, representing approximately 22% of

total FECD cases recruited to date. Overall, 8 rare and potentially deleterious (MAF <0.005;

CADD score >15) *TCF4* variants were identified in 7/134 probands (**Table 3**). Sanger

sequencing confirmed the presence of all 8 variants and exome data from these cases were

also interrogated using the same thresholds (MAF <0.005, CADD >15) to determine if any other

rare variants in genes previously associated with FECD could also be contributing to disease (4)

(**Table S17**). The large number of rare and potentially deleterious *TCF4* variants detected in this

cohort seemed intriguing given the high levels of evolutionary constraint on this essential and

ubiquitously expressed transcription factor encoding gene (pLI=1, missense Z=4.1 gnomAD).

We therefore decided to apply CoCoRV, a rare variant analysis framework to determine if the

Exp-FECD cohort was indeed enriched for rare and potentially deleterious *TCF4* variants

compared to the gnomAD dataset. CoCoRV utilises the publicly available genotype summary

counts to prioritise disease-predisposition genes in case cohorts (40). Under a dominant model,

applying the ethnically-stratified (**Table S18**) analysis pipeline, and filtering for rare (MAF

<0.005) and potentially deleterious (CADD score threshold of >15) variants, an inflation factor

estimate (λemp) 1.21 was achieved. At the exome wide level *TCF4* was ranked as 34/17,025

genes with a p-value of 0.001 suggesting the Exp-FECD cohort could be enriched for rare and

potentially deleterious *TCF4* variants. However, after applying CoCoRV's stringent false

discovery rate correction methodology *TCF4* no longer remained significant (corrected p-value =

0.249).

Interestingly rare variants identified in Proband A (Variant 1a/1b), B (Variant 2) and D

(Variant 4) and E (Variant 5) are all predicted to only affect a small fraction of total *TCF4*

transcripts (**Table 3, S19 and Figure 5**). For Proband A, two consecutive *in cis* heterozygous

missense and nonsense variants were identified. These occur within the coding region of only

one (out of 93) *TCF4* transcripts (ENST00000566286.5; c.[57G>T; 58A>T]; p.[(Arg19Ser;

p.Lys20*)]) and in the 5' UTR of a further 5/93 transcripts. The variant identified in Proband B is

only encompassed by the same 6 *TCF4* transcripts (ENST00000566286.5; c.66G>A,

p.(Glu22=)). Despite being synonymous, Splice AI predicts the variant introduces loss of the

splice donor site (SpliceAI Δ score 0.78). Furthermore, SpliceRover also predicts that this could

result in the activation of a cryptic splice donor site downstream (from 0.098 to 0.233) of the

wildtype donor site, which would introduce a short frameshift insertion followed by a premature

termination codon (PTC) c.66_67insGTGCTCGATGAATTTTC, p.(Arg23Valfs*12). The variant

identified in Proband D (Variant 4) is within the protein coding region of three (out of 93) *TCF4*

transcripts and in the 5' UTR of one additional transcript. Variant 5 is only encompassed by the

5' UTR of a single *TCF4* transcript. Collectively, all of these identified variants have the potential

to exert a functional impact on TCF4, but importantly only on a small subset of overall

transcripts.

Table 3: Summary of rare and potentially deleterious *TCF4* variants identified in a FECD
CTG18.1 Exp- cohort (n=134)

Variant ID	FECD Exp-Cases (n=134)	Variant (hg38)	HGVSc, HGVSp	CADD, MAF gnomAD	Included within CoCoRV analysis
1a	Proband A	chr18:55588479-C-A	ENST00000566286.1: c.57G>T, p.(Arg19Ser)	18.3, 0	Yes
1b	Proband A	chr18:55588478-T-A	ENST00000566286.1: c.58A>T, p.(Lys20Ter)	16.45, 0	Yes
2	Proband B	chr18:55588470-C-T	ENST00000566286.1: c.66G>A, p.(Glu22=)	18.39, 0.002122	Yes
3	Proband C	chr18:55464063-T-C	ENST00000354452.3: c.207+13A>G	20.6, 0.0002234	Yes
4	Proband D	chr18:55403689-A-G	ENST00000544241.2: c.26T>C, p.(Ile9Thr)	17.29, 0	Yes
5	Proband E	chr18:55321599-C-T	ENST00000354452.3: c.549+28760G>A	16.74, 0	No*
6	Proband F	chr18:55261560-A-C	ENST00000354452.3: c.923-27T>G	22.3, 0.0007644	Yes
7	Proband G	chr18:55261512-G-A	ENST00000354452.3: c.944C>T, p.(Ala315Val)	24.6, 0.0005734	Yes

Rare (MAF < 0.005 in gnomAD genomes) and potentially deleterious (CADD score threshold of >15) *TCF4* variants identified within a FECD CTG18.1 Exp- cohort (n=134). Abbreviations are as follows: gnomAD, The Genome Aggregation Database; MAF, minor allele frequency, HGVSc, coding DNA sequence based on the Human Genome Variation Society; HGVSp, protein sequence based on the Human Genome Variation Society. *Variant filtered out at Variant Quality Score Recalibration step.

495
496

Figure 5: Schematic map of rare and potentially deleterious variants identified in individuals with molecularly unsolved FECD within *TCF4*, without CTG18.1 expansions. All 8 rare (minor allele frequency (MAF)<0.005) and potentially pathogenic (CADD>15) variants in 7 Exp- Probands are mapped along *TCF4* genomic region and transcriptomic locations contextualised. Proband A variants are found in *cis* within the proband. Conserved intronic transcripts refer to variants located in intronic regions identified as conserved via Genomic Evolutionary Rate Profiling (GERP) conservation scores on Ensembl. Splice modifying refers to transcripts where the variant identified would induce alternative splicing. Transcript numbers where the variant occurred within introns were not included. UTR: untranslated region. NLS: nuclear localization signal. bHLH: basic helix loop helix

Discussion:

FECD is now recognized to be the most common repeat expansion-mediated disease in humans, with up to 80% of European FECD cases harbouring CTG18.1 expansions(4). The non-coding nature of the repeat and the transcriptomic complexity of *TCF4* have made it extremely experimentally challenging to decipher the molecular mechanisms underlying this disease. Thus, many aspects of its pathophysiology remain elusive, despite the strong mechanistic parallels with other repeat-mediated diseases, and the urgent clinical need for new treatments for this common and visually disabling disease. To tackle this challenge, we employed a large, biologically independent set of primary FECD case-derived CEC lines and numerous complementary transcriptomic methodologies to gain novel mechanistic insight into both RNA toxicity driven and *TCF4* dysregulation-related molecular features of this repeat-mediated disease. By including both Exp+ and Exp- FECD data, we have been able to dissect out more generic downstream biomarkers that occur irrespective of CTG18.1 expansion status.

Our extensive long- and short-read RNA-Seq data and spatial transcriptomic analysis of FECD

case-derived CEC lines demonstrates that, in addition to being a spliceopathy, expansion-

positive FECD is concurrently associated with isoform-specific patterns of *TCF4* dysregulation,

hence providing novel mechanistic insights into the pathophysiology of this triplet-repeat

mediated disease. Furthermore, we have identified a subset of Exp- FECD cases with rare and

potentially deleterious *TCF4* variants, supporting the hypothesis that dysregulation of TCF4

itself irrespective of CTG18.1 expansions may also be a risk factor for FECD.

Our robust transcriptomic analysis, presented within a biologically relevant cellular and

genomic context, highlights an array of novel molecular signatures driven by CTG18.1

expansions, as well as validating a range of previously described Exp+-associated

transcriptomic events. We anticipate that many of the novel events identified here could serve

as informative biomarkers to aid the development of CTG18.1 targeted therapeutic approaches.

To date, CUG-containing RNA foci and MBNL1/2 missplicing have been adopted as the

standard for therapeutic outcome measures (7,27,41,42). Data presented here support and

expand the hypothesis that accumulation of repeat expansion-derived RNA structures induce

toxic gain-of-function effects that disrupt cellular homeostasis and induce aberrant RNA

metabolism(7,27). For instance, we have discovered a novel array of RBP encoding transcripts,

including *MBNL3*, *CELF1* and a range of *hnRNP* and *DDX* family members, uniquely

dysregulated in Exp+ CECs that warrant future follow up to elucidate their respective

contributions to aberrant RNA metabolism related mechanisms. However, such biomarkers may

only relate to the spliceopathy aspects of the disease, and our data go on to also illustrate that

translational efforts would also benefit from using more diverse outcome measures. In addition

to RNA metabolism biomarkers, global Exp+-mediated differential gene expression profiles,

alongside *TCF4*-specific dysregulation itself, should also be included as relevant outcome

measures for Exp+ therapies being developed, given we hypothesise these processes are also

pivotal to the pathology. Novel cell surface biomarkers reported here that are specifically

upregulated in Exp+ (e.g. *ITGA5* and *ITGB1*) could have particular utility in a clinical trial setting

due to their accessible locations. Other membrane encoding proteins interfacing with the

extracellular matrix or signal transduction could also be a rich source of potential external

markers, such as cell surface receptors involved in the P13K-Akt pathway (e.g. *EGFR, MET,*

and *LTK*) that we show are specifically upregulated in Exp+ CECs.

Genetic stratification of patient-derived CECs in this study has also highlighted that Exp+

FECD is transcriptionally distinct from Exp- disease. The data show that 76% of all differentially

expressed genes, 81% of skipped exon events, and 67% of all enriched pathways detected in

Exp+ are unique to the repeat expansion disease. For example, in accordance with previous

studies (12,43), we observe that pronounced upregulation of fibronectin (*FN1*) is a robust

transcriptomic feature of FECD, irrespective of genetic underlying cause. However, our

comprehensive transcriptomic analysis has enabled us to determine that the EDA/B+ forms of

fibronectin (26) are significantly favoured in Exp+ FECD. This illustrates that *FN1* expression

and splicing can act as both a general FECD molecular biomarker and an Exp+ specific

biomarker. Similarly, *AQP1* previously documented to be downregulated in FECD (44), has here

been demonstrated to only display marked downregulation in Exp+ CECs, illustrating the

importance of stratifying cases based on CTG18.1 status. Our long-read data suggests that the

splicing diversity seen in Exp+ FECD is not predominantly a result of abnormal missplicing,

given the proportion of annotated full splice match transcripts remains the same between control

and Exp+. This further exemplifies the unique transcriptome of Exp+ FECD endothelial cells.

However, the increased proportion of non-canonical splice junctions seen in Exp+ also suggests

that CTG18.1-specific isoforms may be present within the endothelium and warrants future

investigation. However, transcriptome wide differences in coverage were observed between the

Exp+ FECD versus the control group within the long-read data, thus it will be important for

future long-read approaches to further validate these findings. These examples, alongside the

emerging evidence that the Exp+ FECD displays distinctive pathological features and

epidemiological characteristics(5,7), suggests that Exp+ FECD should be categorised as a

distinct clinical entity. We suggest integrating CTG18.1 genotyping into routine FECD clinical

care pathways is warranted, especially given the numerous international efforts underway to

develop gene-directed therapies for CTG18.1-mediated FECD (27,41,42,45).

Our complementary RNA-seq approaches, in combination with the spatial resolution of *TCF4*

transcripts subgroups, has highlighted a unique and distinctive pattern of TCF4-specific

dysregulation that occurs within Exp+ FECD. Importantly, this signature of dysregulation was

not detected within a patient and control dermal fibroblast cell series, implying it may be unique

to affected cell types. Additionally, our pathway analysis has isolated features of dysregulation

that we hypothesise represent downstream consequences of this isoform shift and an important

pathogenic component of the disease. Notably, we have observed commonality between

signalling transduction pathways dysregulated in our Exp+ CECs (e.g. P13K-Akt and MAPK)

and those dysregulated in neural progenitor cell *TCF4* knockdown transcriptomes(33). We

postulate that this convergence is being driven by TCF4 functional dysregulation, as the shift in

isoform distributions towards shorter (1-2 AD domains) versus longer (2-3 AD domains)

isoforms would result in lowered TCF4 activity within the corneal endothelial cell. TCF4 isoforms

with 3 AD domains have been shown to have an order of magnitude more transcription

activation function than the shorter isoforms containing 1 or 2 activation domains (16).

Furthermore, given that domains in the AD3-containing isoforms can also dictate intracellular

location and dimerization partners of TCF4, we also hypothesise that additional TCF4-driven

functional implications of this shift occur within affected CECs that requires future experimental

investigation. Interestingly, a recent single-cell (sc)RNA-seq study has identified that varying

levels of *TCF4* expression exist within discrete populations of healthy CECs (46). It was not

possible as part of this study to genotype CTG18.1 whilst concurrently analysing transcriptomic

profiles of the CEC populations, thus future development and application of scRNA-seq in

combination with single cell genotyping of CTG18.1 within FECD-case derived CEC samples is

anticipated to shed light on if and how genomic instability of this repeat element within the

corneal endothelium may drive and/or exacerbate the pathophysiology and explain the unique

vulnerability of this cell layer to CTG18.1 expansions. (47,48)

Additionally, we identify rare and predicted deleterious *TCF4* variants in 7/134 Exp-

FECD cases. This was notable to us given the high levels of evolutionary constraint acting on

TCF4 and suggests such rare variants may be acting as risk factors for FECD, independently of

CTG18.1 repeat expansion. This could also in-part explain the established missing heritability of

the disease(4) and does further reinforcing the hypothesis that dysregulation of *TCF4* itself is

pivotal to the pathophysiology of FECD. Despite a rare variant gene-burden style analysis of

these data not reaching significance, we observed an emerging trend whereby a subset of

qualifying variants identified notably affected regions of the gene not included by the vast

majority of *TCF4* isoforms. Hence, they can only be predicted to exert a functional impact on a

small subset of total TCF4 transcripts. This observation is in keeping with the relatively mild and

tissue specific nature of FECD, which is in stark contrast to the severe neurodevelopmental

disorder Pitt-Hopkins associated with total *TCF4* haploinsufficiency(49). Hence, we propose that

TCF4 variants which subtly affect TCF4 functionality may induce CEC-specific disease. Further

experimental investigation of these variants and the potential tissue specific role they may exert

in the corneal endothelium warrants further future investigation.

Parallels can be drawn to other repeat expansion-mediated disorders, such as fragile X

syndrome (*FRM1*)(50), spinocerebellar ataxia type 7 (*ATXN7,* MIM #164500)(51) and *C9orf72*-

mediated ALS(52), whereby rare loss-of-function mutations have been proposed to induce

comparable phenotypic outcomes to non-coding repeat expansions. However, in these

instances loss of total protein isoform function, in combination with toxic gain-of-function

mechanisms, is thought to underlie the respective pathogenic mechanisms, in contrast to this

scenario whereby we hypothesise only a fraction of total *TCF4* isoforms are susceptible to the

effect of either CTG18.1 expansions or *TCF4* variants within the corneal endothelium. In a

limited number of DM1 cases, FECD has been recognised to be part of the phenotypic

spectrum, suggesting that CEC dysfunction can also arise from CTG repeat expansions in

DMPK (53,54). Data presented in this study suggests that *TCF4* variants, independent of the

repeat, may also represent a potential path to FECD. We suggest future FECD genetic

screening approaches should include sequencing of coding and regulatory regions of *TCF4,* to

further elucidate the potential role that rare *TCF4* variants may play in the absence of CTG18.1

expansions.

In conclusion, this study supports the hypothesis that at least two distinct pathogenic

mechanisms, RNA toxicity and *TCF4* isoform-specific dysregulation, underpin the

pathophysiology of the transcriptionally distinct CTG18.1-mediated FECD. In our established

primary corneal endothelial system, we see evidence of *TCF4* isoform dysregulation at the

TCF4 transcript level and can detect the potential downstream effects of the isoform-specific

pattern of TCF4 dysregulation. Our data also suggests that, in rare instances, irrespective of

CTG18.1 expansions, FECD cases can harbour *TCF4* variants that may independently induce

isoform-specific TCF4 dysregulation and act as risk factors for disease. This supports the

hypothesis that *TCF4* dysregulation is a key mechanism underlying this common, age-related

and visually disabling disease. We anticipate that concepts and biomarkers identified by this

study will inform and guide the development of future translational interventions targeting this

triplet-repeat mediated disease, and these in turn may act as exemplars for much rarer and

currently untreatable and severe repeat expansion disorders that typically affect less accessible

neurological/neuromuscular systems.

Materials and Methods:

Subject Recruitment, phenotyping and bio-sample processing

The study adhered to the tenets of the Declaration of Helsinki and was approved by the Research Ethics Committees of University College London (UCL) (22/EE/0090), Moorfields Eye Hospital (MEH) London (13/LO/1084), or the General University Hospital (GUH) Prague (2/19 GACR). All participants provided written informed consent and had either clinical signs of FECD or had a previous corneal transplant for FECD. For participants who had planned endothelial keratoplasty, we collected the diseased corneal endothelium and Descemet membrane (DM) removed as part of the surgical procedure (MEH). All individuals recruited with FECD undergoing DMEK surgery had confluent guttae affecting the central cornea with visual symptoms and were of similar age. Importantly, all individuals at the time of bio-sample recruitment had similarly advanced disease (Krachmer Grade 5).

We applied strict criteria in the selection for control tissues. We obtained control tissue as donor (age >45 years) corneoscleral discs classed as unsuitable for clinical transplantation due to non-CEC related reasons (Miracles in Sight Eye Bank, Texas, USA). We selected the age of the donor tissues to ensure there was no statistically significant differences in the ages between control and case at the time of bio-sample recruitment. All control corneas were examined and were shown to have a high CEC counts ($\geq$2,600 cells/mm^2) with no guttae or other endothelial abnormality. To standardise handling, the control DM and corneal endothelium were excised by a consultant-grade surgical clinician in accordance with standardised clinical practice. All excised case and control tissues were stored in Lebovitz L-15 media (Life Technologies) before transferring to the laboratory for processing within 24 hours of excision from either donor corneas or patients undergoing elective posterior corneal transplantation surgery.

Participant genomic DNA for CTG18.1 genotyping was extracted from whole blood

(QIAgen Gentra Puregene Blood Kit), whereas control DNA was extracted from the residual

corneoscleral disc (QIAgen Blood and Tissue kits).

CTG18.1 genotyping

We determined the CTG18.1 repeat length using a short tandem repeat (STR)

genotyping assay as previously described in Zarouchlioti et al. 2018(7), originally adapted from

Wieben et al. 2012(55). Briefly, STR-PCR was performed using a 6'-FAM-conjugated primer (5'

CAGATGAGTTTGGTGTAAGAT 3') upstream of the repeat and an unlabeled primer (5'

ACAAGCAGAAAGGGGGCTGCAA 3') downstream. Post-PCR, products were combined with

GeneMarker ROX500 ladder and separated on an ABI 3730 Electrophoresis capillary DNA

analyzer (Applied Biosystems). Sizing and data analysis was performed on GeneMarker

software (SoftGenetics). To distinguish between samples with presumed bi-allelic non-

expanded alleles of equal length and samples with a larger repeat length beyond the detection

limit of the STR-PCR, we performed a further triplet-primed PCR assay to confirm the presence

or absence of a CTG18.1 allele above the detection range of the STR assay (approximately 125

repeats), following previously published methods(56).

Primary corneal endothelial cell (CEC) culture

We isolated and cultured FECD and control-derived CECs as described by Zarouchlioti

et al. 2019(7), adapted from Peh et al. 2015(57). Briefly, to dissolve the DM and detach the

CECs, the tissue sample was incubated for 2 to 4 hours in 0.2% collagenase type I. After

incubation, the CECs were resuspended in stabilisation media (containing Human Endothelial-

SFM (Life technologies) supplemented with 5% foetal bovine serum (FBS), 1%

antibiotic/antimycotic, and 0.1% selective ROCK inhibitor Y-27632 (AdooQ BioSciences)), and

then seeded into 6-well plates pre-coated with FNC coating mixture (Stratech). After 24 hours,

we replaced the culture media with expansion media (Ham's F-12 Nutrient Mix with GlutaMAX

Supplement (Life Technologies)/Medium 199 GlutaMAX Supplement (Life Technologies), 20

µg/mL ascorbic acid, 1% insulin-transferrin-selenium (Life Technologies), 5% FBS, 1%

antibiotic/antimycotic, 10 ng/mL bFGF (Life Technologies) and 0.1% selective ROCK inhibitor Y-

27632 (AdooQ BioSciences)). All cells were cultured at 37°C and 5% CO_2 with twice-weekly

media changes.

Fibroblast cell culture

Fibroblast cell lines from dermal skin biopsies were cultured DMEM/F-12, GlutaMAX

(Life Technologies) supplemented with 10% FBS and 1% penicillin/streptomycin. Throughout

culture, cells were kept in an incubator at 37°C, 5% CO_2 and medium was refreshed every 48 hr

until the cells showed appropriate confluence for experimentation or passage.

RNA extraction

All primary CEC monolayer cultures selected for downstream analysis displayed a

consistent morphology characteristic of CECs *in vivo.* Total RNA was extracted from the primary

CEC lines using a NucleoSpin RNA XS kit (Macherey-Nagel), in accordance with the

manufacturer's protocol. The RNA was eluted and stored at -80°C prior to sequencing. To

assess RNA integrity, RNA integrity numbers (RIN) were determined for all RNA samples using

a Bioanalyzer 2100 (Agilent). All samples analysed as part of this study had a RIN value ≥9.6.

Short-read RNA-seq

Stranded RNA-seq libraries were prepared by BGI Genomics using the TruSeq RNA

Library Prep Kit (Illumina) following the manufacturer's protocol. We sequenced these with a

HiSeq 4000 platform (Illumina) with short-read strand-specific library prep with poly A

enrichment (Illumina). Paired-end reads were filtered to remove low-quality reads (Q ≤5) or

those with >10% of N base calls. Adaptor sequences were also removed during data quality

analysis. We checked the quality of the resulting fastq files for the short-read data using FastQC

Quantification(58). Reads were aligned to the human genome (hg38) patch 13 using STAR (59)

(v2.7.10a) and Salmon (V1.4.0) (60). After alignment, hierarchical clustering was performed

using PoiClaClu(61). We also performed principal component analysis (PCA) using

PCAtools(62) (v2.2.0) to determine global sample clustering. Sex and ethnicity have been

factored in as covariates in all applicable downstream pipelines.

We analysed differential gene expression using DESeq2(31) (version 1.8.2), with

independent hypothesis weighting (IHW) testing of the results to generate FDR values(63,64),

and log_2fold change was calculated using the apeglm (v 1.2) package(65). We then determined

significance as FDR corrected-p value (padj) <0.05. In addition, we analysed different exon

usage amongst the subgroups with DEXSeq(38) (v 1.44.0). Significant results were filtered

using an FDR-corrected P-value of <0.05. Pathway analysis was performed by g:Profiler on

significantly dysregulated genes using g:SCS statistics for thresholding (66).

744
We assessed differential splicing at a junction-based level with rMATS-turbo 4.1.2(23)

with the Ensembl hg38 release 105 GTF file as reference. Default rMATS settings were used.

For this study, we focused on splicing events that are fully/partly annotated by Ensembl.

Significant alternative splicing events were identified as an absolute delta percent spliced in

(psi) magnitude ≥ 0.1 and FDR ≤ 0.05. rMATs output was subsequently analysed using

rMAPS2(30) and maser 1.12.1(67). All script utilised to generate data can be found at:

https://github.com/InheritedCornealDisease/TCF4-transcriptomic-paper-2024

Long-read RNA-seq

PacBio Sequel long-read transcriptomic sequencing (Iso-Seq)(68) was applied to

generate full-length cDNA sequences from control and FECD Exp+ primary CEC-derived RNA

samples. Raw reads were processed and de-multiplexed with Iso-Seq (v3) and merged for

transcript collapse using Cupcake(68) (parameters: "-c 0.85 -i 0.95 --dun-merge-5-shorter").

High-quality, full-length transcripts from the merged dataset were then aligned to the human

reference genome (hg38) using Minimap2(69) (v2.17, parameters: "-ax splice -uf --

secondary=no -C5 -O6,24 -B4") and annotated using SQANTI3(70) (v7.4). SQANTI filtering of

artefacts (intra-priming and reverse transcriptase template switching) was then applied. Github

repository for the analysis described here can be found at:

https://github.com/SziKayLeung/UCL_FECD

RNAScope detection of TCF4 transcripts

We seeded primary FECD and control CECs and fibroblasts on chamber slides (Lab-tek)

and confluent cell cultures were stained using standard RNAScope protocols with minor

adjustments (ACD Bio). We used the Hs-TCF4-C2 (cat. 557411) probe set to detect all *TCF4*

isoforms, as well as a custom set of *TCF4* probes to detect all 5' exons in the *TCF4* transcript

NM_001083962.2 excluding all 3' exons contained in the shorter transcript NM_001243234.2.

Akoya TSA Cy3 fluorophores were used to visualise probe staining. Negative control probes

(cat. 320871) were provided by ACD-Bio to verify specificity. All cell lines analysed by

RNAScope were also tested for RNA foci in parallel via fluorescence *in situ* hybridization (FISH)

using a CUG-specific probe (Cy3-$(CAG)_7$)(7). A minimum of 100 cells and four images were

acquired for each cell line and each probe.

Exome Sequencing and Rare Variant Analysis

Exome sequencing libraries were generated from 134 whole blood derived gDNA

samples from individuals with Exp- FECD, using a SureSelect Human All Exome V6 capture kit

(Agilent, USA) or SeqCap EZ MedExome Enrichment Kit (Roche) and sequenced on a HiSeq

4000 or HiSeq 2500 platform (Illumina). All raw FECD case sequencing data was aligned and

annotated following our previously published methods(71). In brief, sequencing reads were

aligned using Novoalign (version 3.02.08) and variants and indels were called according to

Genome Analysis Toolkit (GATK) (version 4.4.0.0) best practices (joint variant calling followed

by variant quality score recalibration) were normalised and quality-controlled, as recommended

by CoCoRV(40). CoCoRV, a rare variant analysis framework, was applied to prioritise disease-

predisposition genes in the Exp-FECD cohort, utilising publicly available genotype summary

counts from gnomAD (genomes version 2.1.1) as a control dataset. For consistency, both case

and control variants were annotated using the Variant Effect Predictor (VEP) (release 110) (72).

The CoCoRV pipeline was then run, applying a dominant model and ethnically stratified in

accordance with ethnicity predictions generated from the case cohort using Fraposa(22).

Fraposa is based on principal component analysis, using 1000 Genomes project as the

reference panel (with known ancestral background, categorised into 5 distinctive

superpopulations: Africans, admixed Americans, East Asians, Europeans and South Asians).

Common variants between the control and exome cohort were identified and filtered (based on

minor allele frequency, linkage equilibrium and missing genotype call rate) by plink and the

principal components were computed with the default OADP approach (Online Augmentation,

Decomposition, and Procrustes Transformation). Variants of interest were defined as having a

CADD score (CADD) >15 (73), and a minor allele frequency (MAF) <0.005 in the Genome

Aggregation Database (version 2.1.1) (74). FECD cases identified to harbour rare *TCF4*

variants were additionally interrogated for rare and potentially disease-associated variants

(MAF<0.005 in gnomAD and Kaviar Genomic Variant Database; CADD>15) in previously

reported FECD-associated genes, including *COL8A2*, *SLC4A11*, *AGBL1* and *ZEB1*(4). Splice

site predictions were subsequently made using SpliceAI(75) and SpliceRover(76) for all

qualifying variants in *TCF4* and any other FECD-associated genes.

Author Contributions: NB and AED conceptualised the study. NB, NJHT, ANS, CZ, JJ,

LD, PS, MP, KM, PL, SJT contributed to the generation of the data. NB, NC, NJHT, AS, SKL,

TL, IM, ARJ, MEC, AJH, NP and AED all contributed to the analysis and interpretation of the

data. AED and NB wrote the manuscript. All authors read and approved the final version of the

document.

Acknowledgments: We would like to thank all affected individuals for participating in

this research. We would also like to acknowledge Miracles for Sight Eye bank for providing

control corneal tissue. We would also like to thank Dr. Christopher Smith from Queen Mary

University for providing a control fibroblast line. This work was funded by a UKRI Future Leader

Fellowship MR/S031820/1 (AED, CZ, AS), Moorfields Eye Charity GR000060, GR001395,

GR001337 (AED, NB, NHT, AS, NP) and Sight Research UK SAC 036 (AED, AJS), the

Rosetrees Trust (AED, AJS), and the The National Institute for Health Research Biomedical

Research Centre at Moorfields Eye Hospital National Health Service Foundation Trust and UCL

Institute of Ophthalmology (AED, NC., SJT, KM, AJH, MEC, NP). JJ, LD., PS, and PL were

supported by GACR 20-19278S, UNCE/MED/007 and SVV 260631. NP was supported by an

NIHR AI Award (AI AWARD02488). This project utilised equipment funded by the UK Medical

Research Council (MRC) Clinical Research Infrastructure Initiative (award number

MR/M008924/1).

825
826
827
828

References:

1. Mathews P, Benbow A, Corcoran K, DeMatteo J, Philippy B, Van Meter W. 2022 Eye Banking Statistical Report—Executive Summary. Eye Bank Corneal Transplant. 2023 Sep;2(3):e0008.

2. NHSBT. NHS Blood and Transplant Annual Activity Report: Cornea Activity [Internet]. Available from: http://nhsbtdbe.blob.core.windows.net/umbraco-assets-corp/27122/section-10-cornea-activity.pdf

3. Matthaei M, Hribek A, Clahsen T, Bachmann B, Cursiefen C, Jun AS. Fuchs Endothelial Corneal Dystrophy: Clinical, Genetic, Pathophysiologic, and Therapeutic Aspects. Annu Rev Vis Sci. 2019;5(1):151–75.

4. Fautsch MP, Wieben ED, Baratz KH, Bhattacharyya N, Sadan AN, Hafford-Tear NJ, et al. TCF4-mediated Fuchs endothelial corneal dystrophy: Insights into a common trinucleotide repeat-associated disease. Prog Retin Eye Res. 2021 Mar 1;81:100883.

5. Thaung C, Davidson AE. Fuchs endothelial corneal dystrophy: current perspectives on diagnostic pathology and genetics—Bowman Club Lecture. BMJ Open Ophthalmol. 2022 Jul 1;7(1):e001103.

6. Wieben ED, Baratz KH, Aleff RA, Kalari KR, Tang X, Maguire LJ, et al. Gene Expression and Missplicing in the Corneal Endothelium of Patients With a TCF4 Trinucleotide Repeat Expansion Without Fuchs' Endothelial Corneal Dystrophy. Invest Ophthalmol Vis Sci. 2019 Aug 30;60(10):3636–43.

7. Zarouchlioti C, Sanchez-Pintado B, Hafford Tear NJ, Klein P, Liskova P, Dulla K, et al. Antisense Therapy for a Common Corneal Dystrophy Ameliorates TCF4 Repeat Expansion-Mediated Toxicity. Am J Hum Genet. 2018 Apr 5;102(4):528–39.

8. Depienne C, Mandel JL. 30 years of repeat expansion disorders: What have we learned and what are the remaining challenges? Am J Hum Genet. 2021 May 6;108(5):764–85.

9. Balendra R, Isaacs AM. C9orf72-mediated ALS and FTD: multiple pathways to disease. Nat Rev Neurol. 2018 Sep;14(9):544–58.

10. Wieben ED, Aleff RA, Tang X, Butz ML, Kalari KR, Highsmith EW, et al. Trinucleotide Repeat Expansion in the Transcription Factor 4 (TCF4) Gene Leads to Widespread mRNA Splicing Changes in Fuchs' Endothelial Corneal Dystrophy. Invest Ophthalmol Vis Sci. 2017 Jan 24;58(1):343–52.

11. Wieben ED, Aleff RA, Tang X, Kalari KR, Maguire LJ, Patel SV, et al. Gene expression in the corneal endothelium of Fuchs endothelial corneal dystrophy patients with and without expansion of a trinucleotide repeat in TCF4. PLOS ONE. 2018 Jul 2;13(7):e0200005.

12. Chu Y, Hu J, Liang H, Kanchwala M, Xing C, Beebe W, et al. Analyzing pre-symptomatic tissue to gain insights into the molecular and mechanistic origins of late-onset degenerative trinucleotide repeat disease. Nucleic Acids Res. 2020 Jul 9;48(12):6740–58.

13. Soragni E, Petrosyan L, Rinkoski TA, Wieben ED, Baratz KH, Fautsch MP, et al. Repeat-
Associated Non-ATG (RAN) Translation in Fuchs' Endothelial Corneal Dystrophy. Invest
Ophthalmol Vis Sci. 2018 Apr 1;59(5):1888–96.

14. Westin IM, Viberg A, Byström B, Golovleva I. Lower Fractions of TCF4 Transcripts
Spanning over the CTG18.1 Trinucleotide Repeat in Human Corneal Endothelium. Genes.
2021 Dec 17;12(12):2006.

15. Sirp A, Leite K, Tuvikene J, Nurm K, Sepp M, Timmusk T. The Fuchs corneal dystrophy-
associated CTG repeat expansion in the TCF4 gene affects transcription from its alternative
promoters. Sci Rep. 2020 Oct 28;10(1):18424.

16. Sepp M, Kannike K, Eesmaa A, Urb M, Timmusk T. Functional Diversity of Human Basic
Helix-Loop-Helix Transcription Factor TCF4 Isoforms Generated by Alternative 5′ Exon
Usage and Splicing. Kashanchi F, editor. PLoS ONE. 2011 Jul 15;6(7):e22138.

17. Zweier C, Peippo MM, Hoyer J, Sousa S, Bottani A, Clayton-Smith J, et al.
Haploinsufficiency of TCF4 causes syndromal mental retardation with intermittent
hyperventilation (Pitt-Hopkins syndrome). Am J Hum Genet. 2007 May;80(5):994–1001.

18. Amiel J, Rio M, de Pontual L, Redon R, Malan V, Boddaert N, et al. Mutations in TCF4,
encoding a class I basic helix-loop-helix transcription factor, are responsible for Pitt-Hopkins
syndrome, a severe epileptic encephalopathy associated with autonomic dysfunction. Am J
Hum Genet. 2007 May;80(5):988–93.

19. Hellwig M, Lauffer MC, Bockmayr M, Spohn M, Merk DJ, Harrison L, et al. TCF4 (E2-2)
harbors tumor suppressive functions in SHH medulloblastoma. Acta Neuropathol (Berl).
2019 Apr 1;137(4):657–73.

20. Stefansson H, Ophoff RA, Steinberg S, Andreassen OA, Cichon S, Rujescu D, et al.
Common variants conferring risk of schizophrenia. Nature. 2009 Aug 6;460(7256):744–7.

21. Schizophrenia Psychiatric Genome-Wide Association Study (GWAS) Consortium. Genome-
wide association study identifies five new schizophrenia loci. Nat Genet. 2011 Sep
18;43(10):969–76.

22. Zhang D, Dey R, Lee S. Fast and robust ancestry prediction using principal component
analysis. Bioinforma Oxf Engl. 2020 Jun 1;36(11):3439–46.

23. Shen S, Park JW, Lu Z xiang, Lin L, Henry MD, Wu YN, et al. rMATS: Robust and flexible
detection of differential alternative splicing from replicate RNA-Seq data. Proc Natl Acad Sci.
2014 Dec 23;111(51):E5593–601.

24. Shen S, Park JW, Huang J, Dittmar KA, Lu Z xiang, Zhou Q, et al. MATS: a Bayesian
framework for flexible detection of differential alternative splicing from RNA-Seq data.
Nucleic Acids Res. 2012 Apr 1;40(8):e61.

25. Park JW, Tokheim C, Shen S, Xing Y. Identifying Differential Alternative Splicing Events
from RNA Sequencing Data Using RNASeq-MATS. In: Shomron N, editor. Deep
Sequencing Data Analysis [Internet]. Totowa, NJ: Humana Press; 2013 [cited 2022 Oct 31].

p. 171–9. (Methods in Molecular Biology). Available from: https://doi.org/10.1007/978-1-
62703-514-9_10

26. Dalton CJ, Lemmon CA. Fibronectin: Molecular Structure, Fibrillar Structure and
Mechanochemical Signaling. Cells. 2021 Sep;10(9):2443.

27. Hu J, Rong Z, Gong X, Zhou Z, Sharma VK, Xing C, et al. Oligonucleotides targeting TCF4
triplet repeat expansion inhibit RNA foci and mis-splicing in Fuchs' dystrophy. Hum Mol
Genet. 2018 Mar 15;27(6):1015–26.

28. Konieczny P, Stepniak-Konieczna E, Sobczak K. MBNL proteins and their target RNAs,
interaction and splicing regulation. Nucleic Acids Res. 2014;42(17):10873–87.

29. Konieczny P, Stepniak-Konieczna E, Sobczak K. MBNL expression in autoregulatory
feedback loops. RNA Biol. 2018 Jan 2;15(1):1–8.

30. Hwang JY, Jung S, Kook TL, Rouchka EC, Bok J, Park JW. rMAPS2: an update of the RNA
map analysis and plotting server for alternative splicing regulation. Nucleic Acids Res. 2020
Jul 2;48(W1):W300–6.

31. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-
seq data with DESeq2. Genome Biol. 2014 Dec 5;15(12):550.

32. Mori K, Lammich S, Mackenzie IRA, Forné I, Zilow S, Kretzschmar H, et al. hnRNP A3
binds to GGGGCC repeats and is a constituent of p62-positive/TDP43-negative inclusions
in the hippocampus of patients with C9orf72 mutations. Acta Neuropathol (Berl). 2013
Mar;125(3):413–23.

33. Doostparast Torshizi A, Armoskus C, Zhang H, Forrest MP, Zhang S, Souaiaia T, et al.
Deconvolution of transcriptional networks identifies TCF4 as a master regulator in
schizophrenia. Sci Adv. 2019 Sep 11;5(9):eaau4139.

34. Gerstberger S, Hafner M, Tuschl T. A census of human RNA-binding proteins. Nat Rev
Genet. 2014 Dec;15(12):829–45.

35. Li M, Zhuang Y, Batra R, Thomas JD, Li M, Nutter CA, et al. HNRNPA1-induced
spliceopathy in a transgenic mouse model of myotonic dystrophy. Proc Natl Acad Sci. 2020
Mar 10;117(10):5472–7.

36. Laurent FX, Sureau A, Klein AF, Trouslard F, Gasnier E, Furling D, et al. New function for
the RNA helicase p68/DDX5 as a modifier of MBNL1 activity on expanded CUG repeats.
Nucleic Acids Res. 2012 Apr;40(7):3159–71.

37. Pettersson OJ, Aagaard L, Andrejeva D, Thomsen R, Jensen TG, Damgaard CK. DDX6
regulates sequestered nuclear CUG-expanded DMPK-mRNA in dystrophia myotonica type
1. Nucleic Acids Res. 2014 Jun;42(11):7186–200.

38. Anders S, Reyes A, Huber W. Detecting differential usage of exons from RNA-seq data.
Genome Res. 2012 Oct;22(10):2008–17.

39. Sznajder ŁJ, Thomas JD, Carrell EM, Reid T, McFarland KN, Cleary JD, et al. Intron
retention induced by microsatellite expansions as a disease biomarker. Proc Natl Acad Sci.
2018 Apr 17;115(16):4234–9.

40. Chen W, Wang S, Tithi SS, Ellison DW, Schaid DJ, Wu G. A rare variant analysis
framework using public genotype summary counts to prioritize disease-predisposition
genes. Nat Commun. 2022 May 11;13(1):2592.

41. ProQR Therapeutics. Open-Label, Single-Dose, Exploratory Study With QR-504a to
Evaluate Safety, Tolerability, and Corneal Endothelium Molecular Biomarker(s) in Subjects
With Fuchs Endothelial Corneal Dystrophy With Trinucleotide Repeat Expansion in the
TCF4 Gene (FECD3) [Internet]. clinicaltrials.gov; 2022 May [cited 2023 Mar 6]. Report No.:
NCT05052554. Available from: https://clinicaltrials.gov/ct2/show/NCT05052554

42. Powers A, Rinkoski TA, Cheung K, Schehr H, Osgood N, Livelo C, et al. GeneTAC small
molecules reduce toxic nuclear foci and restore normal splicing in corneal endothelial cells
derived from patients with Fuchs endothelial corneal dystrophy (FECD) harboring repeat
expansions in transcription factor 4 (TCF4). Invest Ophthalmol Vis Sci. 2022 Jun
1;63(7):2753-A0242.

43. Goyer B, Thériault M, Gendron SP, Brunette I, Rochette PJ, Proulx S. Extracellular Matrix
and Integrin Expression Profiles in Fuchs Endothelial Corneal Dystrophy Cells and Tissue
Model. Tissue Eng Part A. 2018 Apr;24(7–8):607–15.

44. Thériault M, Gendron SP, Brunette I, Rochette PJ, Proulx S. Function-Related Protein
Expression in Fuchs Endothelial Corneal Dystrophy Cells and Tissue Models. Am J Pathol.
2018 Jul 1;188(7):1703–12.

45. Angelbello AJ, Benhamou RI, Rzuczek SG, Choudhary S, Tang Z, Chen JL, et al. A Small
Molecule that Binds an RNA Repeat Expansion Stimulates Its Decay via the Exosome
Complex. Cell Chem Biol. 2021 Jan 21;28(1):34-45.e6.

46. Wang Q, Dou S, Zhang B, Jiang H, Qi X, Duan H, et al. Heterogeneity of human corneal
endothelium implicates lncRNA NEAT1 in Fuchs endothelial corneal dystrophy. Mol Ther -
Nucleic Acids. 2022 Mar 8;27:880–93.

47. Handsaker R, Kashin S, Reed N, Lee WS, McDonald TM, Tan S, et al. Somatic DNA repeat
expansion underlies Huntington's Disease neuropathology (Plenary Abstract). In
Washington DC; 2023.

48. Gall-Duncan T, Sato N, Yuen RKC, Pearson CE. Advancing genomic technologies and
clinical awareness accelerates discovery of disease-associated tandem repeat sequences.
Genome Res. 2022 Jan;32(1):1–27.

49. Sirp A, Roots K, Nurm K, Tuvikene J, Sepp M, Timmusk T. Functional consequences of
TCF4 missense substitutions associated with Pitt-Hopkins syndrome, mild intellectual
disability, and schizophrenia. J Biol Chem [Internet]. 2021 Dec 1 [cited 2022 Nov 10];297(6).
Available from: https://www.jbc.org/article/S0021-9258(21)01187-X/abstract

50. Tekendo-Ngongang C, Grochowsky A, Solomon BD, Yano ST. Beyond Trinucleotide
Repeat Expansion in Fragile X Syndrome: Rare Coding and Noncoding Variants in FMR1
and Associated Phenotypes. Genes. 2021 Oct 22;12(11):1669.

51. Nath S, Caron NS, May L, Gluscencova OB, Kolesar J, Brady L, et al. Functional
characterization of variants of unknown significance in a spinocerebellar ataxia patient using
an unsupervised machine learning pipeline. Hum Genome Var. 2022 Apr 14;9(1):1–12.

52. Liu F, Liu Q, Lu CX, Cui B, Guo XN, Wang RR, et al. Identification of a novel loss-of-function
C9orf72 splice site mutation in a patient with amyotrophic lateral sclerosis. Neurobiol Aging.
2016 Nov 1;47:219.e1-219.e5.

53. Winkler NS, Milone M, Martinez-Thompson JM, Raja H, Aleff RA, Patel SV, et al. Fuchs'
Endothelial Corneal Dystrophy in Patients With Myotonic Dystrophy, Type 1. Invest
Ophthalmol Vis Sci. 2018 Jun 1;59(7):3053–7.

54. Mootha VV, Hansen B, Rong Z, Mammen PP, Zhou Z, Xing C, et al. Fuchs' Endothelial
Corneal Dystrophy and RNA Foci in Patients With Myotonic Dystrophy. Invest Ophthalmol
Vis Sci. 2017 Sep 1;58(11):4579–85.

55. Wieben ED, Aleff RA, Tosakulwong N, Butz ML, Highsmith WE, Edwards AO, et al. A
Common Trinucleotide Repeat Expansion within the Transcription Factor 4 (TCF4, E2-2)
Gene Predicts Fuchs Corneal Dystrophy. PLOS ONE. 2012 Nov 21;7(11):e49083.

56. Vasanth S, Eghrari AO, Gapsis BC, Wang J, Haller NF, Stark WJ, et al. Expansion of
CTG18.1 Trinucleotide Repeat in TCF4 Is a Potent Driver of Fuchs' Corneal Dystrophy.
Invest Ophthalmol Vis Sci. 2015 Jul;56(8):4531–6.

57. Peh GSL, Chng Z, Ang HP, Cheng TYD, Adnan K, Seah XY, et al. Propagation of human
corneal endothelial cells: A novel dual media approach. Cell Transplant. 2015;24(2):287–
304.

58. Babraham Bioinformatics - FastQC A Quality Control tool for High Throughput Sequence
Data [Internet]. [cited 2022 Oct 31]. Available from:
https://www.bioinformatics.babraham.ac.uk/projects/fastqc/

59. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast
universal RNA-seq aligner. Bioinforma Oxf Engl. 2013 Jan 1;29(1):15–21.

60. Patro R, Duggal G, Love MI, Irizarry RA, Kingsford C. Salmon provides fast and bias-aware
quantification of transcript expression. Nat Methods. 2017 Apr;14(4):417–9.

61. Witten DM. Classification and clustering of sequencing data using a Poisson model. Ann
Appl Stat. 2011 Dec;5(4):2493–518.

62. Blighe K. PCAtools: everything Principal Component Analysis [Internet]. 2023 [cited 2023
Feb 20]. Available from: https://github.com/kevinblighe/PCAtools

63. Ignatiadis N, Huber W. Covariate powered cross-weighted multiple testing. J R Stat Soc Ser
B Stat Methodol. 2021;83(4):720–51.

64. Ignatiadis N, Klaus B, Zaugg JB, Huber W. Data-driven hypothesis weighting increases
detection power in genome-scale multiple testing. Nat Methods. 2016 Jul;13(7):577–80.

65. Zhu A, Ibrahim JG, Love MI. Heavy-tailed prior distributions for sequence count data:
removing the noise and preserving large differences. Bioinformatics. 2019 Jun;35(12):2084–
92.

66. Raudvere U, Kolberg L, Kuzmin I, Arak T, Adler P, Peterson H, et al. g:Profiler: a web server
for functional enrichment analysis and conversions of gene lists (2019 update). Nucleic
Acids Res. 2019 Jul 2;47(W1):W191–8.

67. Veiga DT. maseR: Mapping Alternative Splicing Events to pRoteins [Internet]. 2022 [cited
2023 Feb 20]. Available from: https://github.com/DiogoVeiga/maser

68. Gordon SP, Tseng E, Salamov A, Zhang J, Meng X, Zhao Z, et al. Widespread Polycistronic
Transcripts in Fungi Revealed by Single-Molecule mRNA Sequencing. PLOS ONE. 2015 Jul
15;10(7):e0132628.

69. Li H. Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics. 2018 Sep
15;34(18):3094–100.

70. Tardaguila M, Fuente L de la, Marti C, Pereira C, Pardo-Palacios FJ, Risco H del, et al.
SQANTI: extensive characterization of long-read transcript sequences for quality control in
full-length transcriptome identification and quantification. Genome Res. 2018 Jan
3;28(3):396–411.

71. Pontikos N, Yu J, Moghul I, Withington L, Blanco-Kelly F, Vulliamy T, et al. Phenopolis: an
open platform for harmonization and analysis of genetic and phenotypic data.
Bioinformatics. 2017 Aug 1;33(15):2421–3.

72. McLaren W, Gil L, Hunt SE, Riat HS, Ritchie GRS, Thormann A, et al. The Ensembl Variant
Effect Predictor. Genome Biol. 2016 Jun 6;17(1):122.

73. Kircher M, Witten DM, Jain P, O'Roak BJ, Cooper GM, Shendure J. A general framework for
estimating the relative pathogenicity of human genetic variants. Nat Genet. 2014
Mar;46(3):310–5.

74. Glusman G, Caballero J, Mauldin DE, Hood L, Roach JC. Kaviar: an accessible system for
testing SNV novelty. Bioinformatics. 2011 Nov 15;27(22):3216–7.

75. Jaganathan K, Kyriazopoulou Panagiotopoulou S, McRae JF, Darbandi SF, Knowles D, Li
YI, et al. Predicting Splicing from Primary Sequence with Deep Learning. Cell. 2019 Jan
24;176(3):535-548.e24.

76. Zuallaert J, Godin F, Kim M, Soete A, Saeys Y, De Neve W. SpliceRover: interpretable
convolutional neural networks for improved splice site prediction. Bioinformatics. 2018 Dec
15;34(24):4180–8.

1051
1052
1053
1054

Supplemental Material Figure and Table legends

Figure S1: Pairwise comparison of rMATS alternative splicing analysis demonstrates increased levels of alternative splicing in Exp+ FECD compared to both Exp- FECD and control primary corneal endothelial cells. Short-read RNA-seq data generated from primary corneal endothelial cells were analysed by rMATS. **A)** Table of rMATS events for each pairwise comparison (PWC). Significance denoted by FDR ≤ 0.05 and deltapsi magnitude larger than 0.1. Values in brackets show percentage of total splice events **B)** Venn diagram of skipped exon event coordinates between all PWCs. The largest overlap is observed between PWC1 and PWC2 highlighting an enrichment of skipped exon events in Exp+ FECD compared to Exp-FECD and controls.

Figure S2: Alternative splicing analyses of short-read CEC RNA-seq data demonstrate increased levels of alternative splicing in Exp+ FECD. A) Summary of global differences in splicing events categorised by rMATS**.** N=4 for controls, N=3 each for FECD Exp+ and FECD Exp-. The most common splice category of alternative splicing observed between all pairwise comparisons was skipped exon, representing ~70% of all significant alternative splicing events detected. Alt 3' SS: alternative 3' splice site, Alt 5' SS: alternative 5' splice site, Ret. Intron: retained intron, MXE: mutually exclusive exon **B)** Volcano plot of statistically significant skipped exon events in PWC1 (Control vs Exp+). The dpsi value denotes the magnitude of change for each dysregulated skipped exon event identified. A positive dspi denotes decreased levels of exon inclusion in Exp+, whereas a negative dspi denotes increased levels of exon inclusion in Exp+.

Figure S3: Pairwise comparison of differentially expressed exons identified by DEXSeq2 demonstrates increased levels of alternative splicing in Exp+ FECD compared to both

1081 **Exp- FECD and control primary corneal endothelial cells.   A)** Table of DEXSeq2 events

1082 and genes for each pairwise comparison (PWC). Significance denoted by p-adj ≤ 0.05. **B)** Venn

1083 diagram of dysregulated exons coordinates between all three PWCs. The largest overlap is

1084 observed between PWC1 and PWC2, highlighting an enrichment of skipped exon events in

1085 Exp+ FECD compared to Exp- FECD and controls.

1086

1087 **Table S1:** Clinical details of individuals with Fuchs endothelial corneal dystrophy used to

1088 established primary corneal endothelial cell cultures used for downstream analysis.

1089

1090 **Table S2:** Driving genes and additional principal components for PCA analysis of short-read

1091 transcriptomic data

1092

1093 **Table S3:** Summary of SQANTI3 analysis of IsoSeq data. In depth characterization of isoforms

1094 in control and Exp+ long-read RNA-seq utilising SQANTI3 pipeline following stringent quality

1095 control.

1096

1097 **Table S4:** Summary of rMATs results. The top five splice events detected for the following

1098 pairwise comparisons (PWC) are presented; (PWC1) control versus Exp+ FECD, (PWC2) Exp+

1099 FECD versus Exp- FECD, and (PWC3) control versus Exp- FECD.

1100

1101 **Table S5:** rMATS identified significant differentially spliced events in Exp+ matching published

1102 events with strong association to CTG18.1-expansion mediated FECD.

1103 **Table S6:** Details on the pairwise comparison(s) where rMATS identified new significant

1104 differentially spliced events matching published differentially spliced genes with strong

1105 association to CTG18.1-expansion mediated FECD.

1106

1107 **Table S7:** Differential gene expression and alternative splicing of fibronectin (*FN1*) in all three

1108 pairwise comparisons.

1109

1110 **Table S8:** rMAPS-identified RNAbinding motif enrichment in PWC1 and PWC2, which was also

1111 absent in PWC3.

1112

1113 **Table S9:** Differentially expressed genes for all three pairwise comparisons including list of

1114 dysregulated genes that are common to FECD in general.

1115

1116 **Table S10:** Pathway enrichment (GO, KEGG, and Reactome) for PWC1, PWC3, and

1117 dysregulated genes unique to Exp+.

1118

1119 **Table S11:** Dysregulated RNA binding proteins (RBPs) uniquely in PWC1.

1120

1121 **Table S12:** Summary of all *TCF4* DEXSeq runs

1122

1123 **Table S13:** DEXSeq results for *TCF4* in all three pairwise comparisons showing dysregulation

1124 of exons within the gene.

1125

1126 **Table S14:** rMATS *TCF4* mutually exclusive exon events demonstrating a shift between longer

1127 isoforms containing 2+ AD domains and shorter isoforms containing 1-2 AD domains detected

1128 via alternative splicing pipelines.

1129

1130 **Table S15**: *TCF4* isoform RNAScope sample summary with results of FISH with probe targeting

1131 repeat and negative RNAScope control experiments (with probes targeting bacterial genes).

1132

1133 **Table S16:** *TCF4* isoform RNAScope sample summary with results of FISH with probe targeting

1134 repeat and negative RNAScope experiments (with probe targeting bacterial genes) in adult

1135 human dermal fibroblasts.

1136

1137 **Table S17:** Summary of rare, potentially deleterious, variants in FECD-associated genes

1138 identified in Proband B.

1139

1140 **Table S18:** A summary of FRAPOSA-derived ancestry information generated for all CTG18.1

1141 Exp- using genome-wide SNP data extracted from exome sequencing data.

1142

1143 **Table S19:** Clinical data of probands with Fuchs endothelial corneal dystrophy identified to have

1144 rare and potentially deleterious heterozygous *TCF4* variants.