1
12 **Deterministic Genetic Barcoding for Multiplexed Behavioral and Single-Cell**
13 **Transcriptomic Studies**
14
15

16 Jorge Blanco Mendana[1], Margaret Donovan[1], Lindsey Gengelbach[1], Benjamin Auch[1,†], John
17 Garbe[1], Daryl M. Gohl[1,2,3]
18
19 [1] University of Minnesota Genomics Center, Minneapolis, MN, 55455, USA
20 [2] Department of Genetics, Cell Biology, and Development, University of Minnesota, Minneapolis,
21 MN 55455, USA
22
23
24 [3] Corresponding author: dmgohl@umn.edu
25 [†] Present address: Phase Genomics, Seattle, WA, 98109, USA

26
27

Abstract

Advances in single-cell sequencing technologies have provided novel insights into the dynamics of gene expression throughout development, been used to characterize somatic variation and heterogeneity within tissues, and are currently enabling the construction of transcriptomic cell atlases. However, despite these remarkable advances, linking anatomical information to transcriptomic data and positively identifying the cell types that correspond to gene expression clusters in single-cell sequencing data sets remains a challenge. We describe a straightforward genetic barcoding approach that takes advantage of the powerful genetic tools available in Drosophila to allow *in vivo* tagging of defined cell populations. This method, called Targeted Genetically-Encoded Multiplexing (TaG-EM), involves inserting a DNA barcode just upstream of the polyadenylation site in a Gal4-inducible *UAS-GFP* construct so that the barcode sequence can be read out during single-cell sequencing, labeling a cell population of interest. By creating many such independently barcoded fly strains, TaG-EM will enable a number of potential applications that will improve the quality and information content of single-cell transcriptomic data including positive identification of cell types in cell atlas projects, identification of multiplet droplets, and barcoding of experimental timepoints, conditions, and replicates. Furthermore, we demonstrate that the barcodes from TaG-EM fly lines can be read out using next-generation sequencing to facilitate population-scale behavioral measurements. Thus, TaG-EM has the potential to enable large-scale behavioral screens in addition to improving the ability to reliably annotate cell atlas data, expanding the scope, and improving the robustness of single-cell transcriptomic experiments.

Introduction

Spatially and temporally regulated gene expression patterns are a hallmark of multicellular life and function to orchestrate patterning, growth, and differentiation throughout development (Ingham, 1988; Reeves et al., 2006). In mature organisms, spatial expression patterns both in tissues and within cells define functionally distinct compartments and determine many aspects of cellular and organismal physiology (Martin and Ephrussi, 2009). In addition, such expression patterns differentiate healthy and diseased tissue and impact disease etiology (Marusyk et al., 2012). Spatial and temporal expression patterns, which can be used to distinguish between cell types and provide insight into cellular function, also provide a means to understand the organization and physiology of complex tissues such as the brain (Thompson et al., 2014). Thus, robust and scalable tools for measuring spatial and temporal gene expression patterns at a genome-wide scale and at high resolution would be transformative research tools across many biological disciplines.

Single-cell sequencing technologies have provided insights into the dynamics of gene expression throughout development, been used to characterize somatic variation and heterogeneity within tissues, and are currently enabling the construction of transcriptomic cell atlases (Klein et al., 2015; Macosko et al., 2015; Zheng et al., 2017). However, linking anatomical information to transcriptomic data and positively identifying the cell types that correspond to gene expression clusters in single-cell sequencing data sets remains a challenge. The cellular identities of gene expression clusters identified in cell atlas data sets are typically inferred from the expression of distinctive gene sets (Hung et al., 2020; Li et al., 2022; Ma et al., 2021), and the lack of positive identification of gene expression clusters introduces an element of uncertainty in this analysis. Moreover, this process of manual annotation is labor-intensive and often requires additional experiments to determine or confirm the expression patterns of marker genes. Emerging spatial genomics technologies hold promise in linking anatomical and transcriptomic information (Lee et al., 2015; Lein et al., 2017). Several of the emerging commercial spatial genomics technologies rely on *in situ* sequencing of marker genes allowing droplet-based single-cell transcriptomic data to be mapped onto a tissue. However, these technologies currently suffer from constraints related to cost, content, or applicability to specific model systems.

In addition to descriptive cell atlas projects, studies involving multiple experimental timepoints throughout development and aging, or to assess the effects of experimental exposures or genetic manipulations would benefit from increased ability to multiplex samples. Given the fixed costs of droplet-based single-cell sequencing, generating data for single-cell transcriptomic time courses or experimental manipulations can be costly. These costs are also a barrier to including

replicates to assess biological variability; consequently, a lack of replicates is a common shortcoming of single-cell sequencing experiments. Antibody-based cell hashing or feature barcoding approaches have been developed to allow multiplexing of samples in droplet-based single-cell sequencing reactions (Stoeckius et al., 2018, 2017). In addition, other multiplexing strategies for single-cell sequencing based on alternative methods for tagging cells (Cheng et al., 2021) or making use of natural genetic variation have been used (Kurmangaliyev et al., 2020). While such approaches can reduce per-sample costs, typically samples are barcoded at a population level and thus do not enable labeling of cell subpopulations within a sample.

We have developed a straightforward genetic barcoding approach that takes advantage of the powerful genetic tools available in Drosophila to allow deterministic *in vivo* tagging of defined cell populations. This method, called <u>Ta</u>rgeted <u>G</u>enetically-<u>E</u>ncoded <u>M</u>ultiplexing (TaG-EM), involves inserting a DNA barcode just upstream of the poly-adenylation site in a Gal4-inducible *UAS-GFP* construct so that the barcode sequence can be read out during droplet-based single-cell sequencing, labeling a cell population of interest.

Genetic barcoding approaches have been employed in many unicellular systems, cell culture, and viral transfection, to facilitate high throughput screening using sequencing-based readouts (Bhang et al., 2015; Smith et al., 2009; van Opijnen et al., 2009). In multicellular animals, techniques such as GESTALT have enabled lineage tracing by using CRISPR to create unique barcodes in differentiating tissue (McKenna et al., 2016), and barcode sequencing has also been employed to map connectivity in the brain (Chen et al., 2019). In addition to improving single-cell sequencing-based measurements, genetically barcoded fly lines can enable highly multiplexed behavioral assays read out using high throughput sequencing. Flies carrying TaG-EM barcodes can be exposed to different experimental perturbations and then run through assays where flies, larvae, or embryos are fractionated based on behavioral outcomes or other phenotypes. Thus, TaG-EM has the potential to enable large-scale next-generation sequencing (NGS)-based behavioral or other fractionation screens analogous to BAR-Seq or Tn-Seq approaches employed in microbial organisms.

Results

TaG-EM: A novel genetic barcoding strategy for multiplexed behavioral and single-cell transcriptomics

We cloned a fragment containing a PCR handle sequence and a diverse 14 bp barcode sequence into the SV40 3' untranslated region (UTR) sequence just upstream of the polyadenylation sites in the 10xUAS-myr::GFP (pJFRC12, (Pfeiffer et al., 2010)) backbone (Figure 1A). A pool containing 29 unique barcode-containing plasmids was injected into Drosophila embryos for PhiC31-mediated integration into the attP2 landing site (Groth et al., 2004) and transgenic lines were isolated and confirmed by Sanger sequencing (Supplemental Figure 1). We recovered 20 distinctly barcoded Drosophila lines, with some barcodes recovered from multiple crosses (Supplemental Figure 1). Such barcoded fly lines have the potential to enable population behavioral measurements, where different exposures, experimental timepoints, and genetic or neural perturbations can be multiplexed and analyzed by measuring barcode abundance in sequencing data (Figure 1B). In addition, the barcodes, which reside on a Gal4-inducible *UAS-GFP* construct, can be expressed tissue-specifically and read out during droplet-based single-cell sequencing, labeling a cell population and/or an experimental condition of interest (Figure 1C).

Testing the accuracy and reproducibility of TaG-EM behavioral measurements using structured pools

We conducted initial experiments to optimize amplification of the genetic barcodes using primers targeting the PCR handle inserted just upstream of the 14 bp barcode sequence and PCR primers downstream of the TaG-EM barcode in the SV40 3' UTR sequence (Supplemental Figure 2). To test the accuracy and reproducibility of sequencing-based measurements of TaG-EM barcodes, we constructed structured pools containing defined numbers of flies pooled either evenly with each of the 20 barcode constructs comprising 5% of the pool, or in a staggered manner with sets of barcodes differing in abundance in 2-fold increments (Figure 2A). To examine the impact of technical steps such as DNA extraction and PCR amplification on TaG-EM barcode measurements, even pools were made and extracted in triplicate and amplicon sequencing libraries were made in triplicate for each independently extracted DNA sample for both the even and staggered pools. The resulting data indicated that TaG-EM measurements are highly accurate and reproducible. Technical replicates (indicated by error bars in Figure 2B-E) showed minimal variability. Likewise, the three independently extracted replicates of the even

pools produced consistent data with all 20 barcodes detected at levels close to the expected 5%

abundance (Figure 2B-C). Barcode abundance values for the staggered structured pools was

generally consistent with the input values and in most cases, the 2-fold differences between the

different groups of barcodes could be distinguished (Figure 2D-E). For the staggered pools,

abundances correlated well with the expected values, particularly when multiple barcodes for an

input level were averaged, in which case R^2 values were >0.99 (Figure 2D-E, inset plots). This

indicates that a high level of quantitative accuracy can be attained using sequencing-based

analysis of TaG-EM barcode abundance, particularly when averaging data for three to four

independent barcodes for an experimental condition.

TaG-EM measurement of phototaxis behavior correlate well with video-based
measurements

Next, we tested whether TaG-EM could be used to measure a phototaxis behavior. A

mixture of barcoded wild type or blind *norpA* mutant flies were run together through a phototaxis

assay. At the end of a period of light exposure, test tubes facing toward or away from the light

were capped, DNA was isolated, and barcodes were amplified and sequenced for each tube.

Raw read counts were scaled in proportion to the number of flies per tube and a preference

index was calculated for each barcode (Figure 3A). In parallel, individual preference indices

were calculated based on manual scoring of videos recorded for each line (Figure 3B).

Preference indices calculated for the pooled, NGS-based TaG-EM measurements were nearly

identical to conventional behavioral measurements for both wild type and *norpA* mutants (Figure

3A-B).

TaG-EM measurement of oviposition behavior and age-dependent fecundity

We next tested whether NGS-based pooled measurements of egg laying could be made.

Fertilized females from each of the 20 barcode lines were placed together in egg laying cups,

embryos were collected, aged for 12 hours to enable cell numbers to stabilize in the developing

eggs, and then DNA was extracted from both the pooled adult flies and the embryos. In general,

TaG-EM measurements of oviposition correlated with fly numbers, with the exception of

barcode 14 which had reduced barcode abundance across multiple trials (Supplemental Figure

3). This suggests that despite the fact that the genetic barcode constructs are inserted in a

common landing site, differences with respect to specific behaviors may exist among the lines

and thus one should test to make sure given lines are appropriate to use in specific behavioral

assays.

To determine whether TaG-EM could be used to measure age-dependent fecundity, we

collected flies from twelve different TaG-EM barcode lines at four time points separated by one

week (three barcode lines per timepoint). We collected eggs from these fly lines individually and

scored the number of viable eggs per female. Next, we pooled the barcoded flies from all

timepoints and collected eggs from the pooled flies. These eggs were aged, DNA was

extracted, and the TaG-EM barcodes were amplified and sequenced. While measurements from

individual barcode lines were noisy, both for manual counts and sequencing based

measurements, there was a general trend toward declining fecundity over time (Supplemental

Figure 4), consistent with published reports (David et al., 1975). Manually scored viable egg

numbers and TaG-EM barcode abundances were well correlated across two independent

experimental trials (R^2 values of 0.52-0.61 for Trial 1 and 0.74-0.84 for Trial 2). When barcodes

from each individual timepoint were averaged, R^2 values for the correlation between manual and

sequencing-based measurements were 0.95 for Trial 1 and 0.99 for Trial 2 (Figure 3C,

Supplemental Figure 4).

Tissue-specific expression of TaG-EM GFP constructs

To facilitate representation of the TaG-EM barcodes in single-cell sequencing data,

genetic barcodes were placed just upstream of the polyadenylation signal sequences and polyA

cleavage sites (Figure 4A). To verify that the inserted sequences did not interfere with Gal4-

driven GFP expression, we crossed each of the barcoded TaG-EM lines to *decapentaplegic-*

Gal4 (*dpp-Gal4*). We observed GFP expression in the expected characteristic central stripe

(Teleman and Cohen, 2000) in the wing imaginal disc for 19/20 lines at similar expression levels

to the base pJFRC12 *UAS-myr::GFP* construct inserted in the same landing site (Figure 4B,

Supplemental Figure 6). No GFP expression was visible for TaG-EM barcode number 8. Gal4-

driven expression levels of TaG-EM barcoded GFP constructs were also similar to that of the

pJFRC12 base construct for multiple driver lines (Supplemental Figure 7) indicating that the

presence of the barcode does not generally impair expression of GFP.

Boosting the GFP signal of TaG-EM constructs to enable robust cell sorting

While with some driver lines, expression of the myr::GFP from the TaG-EM construct

may be too weak to allow robust enrichment of the tagged cells, adding an additional hexameric

GFP construct (Shearin et al., 2014) could boost expression of weak driver lines to levels that

are sufficient for robust detection of labeled flies or larvae (Figure 4C) and for labeling of

dissociated cells for flow cytometry (Supplemental Figure 8). Stocks with an additional UAS

hexameric GFP construct recombined onto the same chromosome as the TaG-EM construct

have been established for all 20 TaG-EM barcode lines.

Correlation between expression of TaG-EM barcodes and intestinal cell marker genes in

single-cell sequencing data

To test whether we could detect TaG-EM barcodes in single-cell sequencing data, we

crossed three TaG-EM barcode lines to two different gut Gal4 driver lines (Ariyapala et al.,

2020), one expressing in the enterocytes (EC-Gal4) and the other in intestinal precursor cells

(PC-Gal4), including stem cells and enteroblasts (EBs). Due to weak GFP expression with the

EC-Gal4 driver, we did not see visible GFP positive cells for this driver line. The PC-Gal4 driver

line contained an additional UAS-Stinger (2xGFP) construct and expressed GFP at a level

sufficient for flow sorting when crossed to the TaG-EM line (Supplemental Figure 7). Larval guts

were dissected, dissociated, stained with propidium iodide (PI) to label dead cells, and flow

sorted to recover PI negative and GFP positive cells. Approximately 10,000 cells were loaded

into a 10x Genomics droplet generator and a single-cell library was prepared and sequenced.

Two clusters were observed in the resulting sequencing data, one of which had high read

counts from mitochondrial genes suggesting that this cluster consisted of mitochondria, debris,

or dead and dying cells, after filtering the cells with high mitochondrial reads, a single cluster

remained (Supplemental Figure 9). This cluster expressed known intestinal precursor cell

markers such as *escargot* (*esg*), *klumpfuss* (*klu*), and *Notch* pathway genes like *E(spl)mbeta-*

HLH (Supplemental Figure 9). Expression of all three PC-Gal4 driven TaG-EM barcodes was

observed in this cluster (Supplemental Figure 9) indicating that TaG-EM barcodes can be

detected in single-cell sequencing data. Interestingly, TaG-EM barcode 8, for which no GFP

expression was observed was represented in the single-cell sequencing data indicating that the

insertion of this specific barcode sequence may affect GFP translation, but not mRNA

expression for this line.

A previous study used droplet-based single-cell sequencing to characterize the cell

types that make up the adult midgut (Hung et al., 2020). This study took advantage of two

fluorescent protein markers, an *escargot* (*esg*)-GFP fusion protein and a *prospero* (*pros*)-Gal4

driven RFP to label the intestinal stem cells (ISCs) and enteroendocrine cells (EECs),

respectively (Hung et al., 2020). The authors compared the resulting clusters to a list of known

marker genes in the literature, including antibody staining, GFP, LacZ, and Gal4 reporter

expression patterns to classify the cells in individual clusters, and also found that the *esg*-GFP

expression was present in a broader subset of cells than anticipated. Thus, most of these cell

classifications relied upon inference as opposed to direct positive labeling. Recently, a large

collection of split-*Gal4* lines were screened for expression in the adult and larval gut (Ariyapala

et al., 2020). These include pan-midgut driver lines, split-*Gal4* lines specific for the EBs, ECs,

EECs, and ISC/EBs, as well as driver lines with regionalized gene expression. We crossed four

different TaG-EM barcode lines with the pan-midgut driver (PMG-Gal4), and one barcode line to

each of the precursor cell (PC-Gal4), enterocyte (EC-Gal4), enteroblast (EB-Gal4) and

enteroendocrine (EE-Gal4) drivers (Ariyapala et al., 2020). Larval guts were dissected from

these lines and cells were dissociated, flow sorted as described above to select live, GFP

positive cells, and approximately 30,000 cells were loaded into a 10x Genomics droplet

generator for single-cell sequencing. Using the additional hexameric GFP construct to boost

GFP expression resulted in visible fluorescent signal for all eight barcode Gal4 line

combinations.

An advantage of cell barcoding both for cell hashing (Stoeckius et al., 2018) and for

TaG-EM *in vivo* barcoding is that such labeling facilitates the identification and removal of

multiplets, which are an artifact of droplet-based single-cell sequencing approaches. After

filtering and removing cells with a high percentage of mitochondrial or ribosomal reads, we

searched for cells that co-expressed multiple TaG-EM barcodes. Out of 8,847 cells, we

identified and removed 854 cells (9.65%) that expressed two different TaG-EM barcodes

(Figure 5A, Supplemental Figure 10). We additionally attempted amplifying the TaG-EM

barcodes in parallel to preparation of the gene expression library, following a modified cell

hashing enrichment protocol. However, this enrichment was unsuccessful (resulting sequencing

reads were predominantly off-target) and thus amplification of TaG-EM barcodes will require

further troubleshooting.

After doublet removal, the remaining cells were clustered (Figure 5B) and analyzed

using ScanPy (Wolf et al., 2018). Analysis of differentially expressed genes identified clusters

expressing marker genes previously reported for adult gut cell types (Hung et al., 2020). These

included genes associated with precursor cells (*Notch* pathway genes), enterocytes (trypsins,

serine proteases, amalyse, mannosidases), and enteroendocrine cells (neuropeptides and

neuropeptide receptors) (Supplemental Figure 11, Supplemental Figure 12).

As expected, the barcodes expressed by the pan-midgut driver were broadly distributed

across the cell clusters (Supplemental Figure 13). In contrast, expression of the cell type-

specific barcodes showed more restricted patterns of expression among the cell clusters and

were co-localized with known marker genes for these cell types (Figure 5C-N). For instance,

TaG-EM barcode 4 expression (Figure 5F), which was driven by the EC-Gal4 line, was seen in

clusters that expressed enterocyte markers such as the serine protease *Jon99Ciii* (Figure 5C)

and other enterocyte marker genes such as the amylase, *Amy-d* (not shown). TaG-EM barcode

4 did not label other classes of enterocyte cells such as *beta-Trypsin* or *LManVI* positive

enterocytes (Figures 5D-E). TaG-EM barcode 6 (Figure 5J), driven by the EB-Gal4 line, was

strongly expressed in cells that also expressed precursor cell markers such as *esg* (Figure 5G),

klu (Figure 5H), and Notch pathway genes such as *e(spl)mbeta-HLH* (Figure 5I) and *e(spl)m3-*

HLH (not shown). Finally, expression of TaG-EM barcode 9 (Figure 5N), which was expressed

using the EE-Gal4 driver line, was observed in clusters of cells that also expressed

enteroendocrine cell derived neuropeptide genes such as *Dh31* (Figure 5K) and other

enteroendocrine markers such as *IA-2*, a tyrosine phosphatase involved in the secretion of

insulin-like peptide (Figure 5L). As with the enterocyte labeling, the EE-Gal4 expressed TaG-EM

barcode 9 did not label all classes of enteroendocrine cells and other clusters of presumptive

enteroendocrine cells expressing other neuropeptides such as *Orcokinin* (Figure 5M), *AstA* (not

shown), and *AstC* (not shown), or neuropeptide receptors such as *CCHa2* (not shown) were

also observed. The EE-Gal4 driver uses *Dh31* regulatory elements, so it is not surprising that

the TaG-EM barcodes specifically labeled *Dh31*-positive enteroendocrine cells and this result

further highlights the ability to target specific genetically defined cell types using TaG-EM based

on *in vivo* cell labeling. Taken together, these results demonstrate that TaG-EM can be used to

label specific cell populations *in vivo* for subsequent identification in single-cell sequencing data.

Discussion

Advances in next-generation sequencing, as well as single-cell and spatial genomics are

enabling new types of detailed analyses to study important biological processes such as

development and nervous system function. Here we describe TaG-EM, a genetic barcoding

strategy that enables novel capabilities in several different experimental contexts (Figure 1).

We demonstrate that the genetic barcodes can be quantified from mixtures of barcoded

fly lines using next-generation sequencing. Analysis of structured pools of flies with defined

inputs suggests that TaG-EM barcode measurements are highly accurate and reproducible,

particularly in cases where multiple barcodes are used to label an experimental condition and

averaged (Figure 2). Sequencing-based TaG-EM measurements recapitulated more laborious,

one-at-a-time measurements in both a phototaxis assay and an age-dependent fecundity assay,

demonstrating that TaG-EM can be used to measure behavior or other phenotypes in

multiplexed, pooled populations (Figure 3). We did note that one line (TaG-EM barcode 14)

exhibited poor performance in oviposition assays, suggesting that barcode performance should

be verified for a specific assay, though other strategies such as averaging across multiple

barcode lines or permutation of barcode assignment across replicates could also mitigate such

deficiencies. Currently, up to twenty conditions can be multiplexed in a single pooled experiment

with existing TaG-EM lines, but because sequencing indices can be added after amplification in

a separate indexing PCR step, hundreds or even thousands of such experiments can be

multiplexed in a single sequencing run.

In addition, we show that TaG-EM barcodes can be expressed by tissue-specific Gal4

drivers and used to tag specific cell populations upstream of single-cell sequencing (Figures 4-

5). This capability will allow for positive identification of cell clusters in cell atlas projects and will

facilitate multiplexing of single-cell sequencing experiments. Recently, a conceptually similar

approach called RABID-Seq (Clark et al., 2021) has been described, which allows trans-

synaptic labeling of neural circuits using barcoded viral transcripts. However, one distinction

between the two approaches is that RABID-Seq relies on stochastic viral infection of

mammalian cells while TaG-EM allows reproducible targeting of defined cell populations

allowing unambiguous cell identification and potentially allowing the same cell populations to be

assessed at different timepoints or in the context of different experimental manipulations. One

current limitation is that TaG-EM barcodes are not observed in all cells in single-cell gene

expression data and low expressing driver lines may result in particularly sparse labelling. In

antibody-conjugated oligo cell hashing approaches, this sparsity of barcode representation is

overcome by spiking in an additional primer at the cDNA amplification step and amplifying the

hashtag oligo by PCR. A similar enrichment approach could be employed to generate dense

labeling of the TaG-EM barcodes, improving upon the current mapping of barcode-containing

transcripts observed in the gene expression data, and facilitating comparison between

replicates and robust detection of multiplets.

In the future, generation of additional TaG-EM lines will enable higher levels of

multiplexing. In addition, while the original TaG-EM lines were made using a membrane-

localized myr::GFP construct, variants that express GFP in other cell compartments such as the

cytoplasm or nucleus could be constructed to enable increased expression levels or purification

of nuclei. Nuclear labeling could also be achieved by co-expressing a nuclear GFP construct

with existing TaG-EM lines in analogy to the use of hexameric GFP described above.

In summary, combined with the large collections of Gal4 and split-Gal4 lines that have

been established in Drosophila that enable precise targeting of a high proportion of cell types

(Ariyapala et al., 2020; Aso et al., 2014; Davis et al., 2020; Gohl et al., 2011; Namiki et al., 2018;

Pfeiffer et al., 2010, 2008; Venken et al., 2011), TaG-EM provides a means to target and label

cells *in vivo* for subsequent detection in single-cell sequencing. Moreover, these genetic

barcodes can be used to multiplex behavioral or other phenotypic measurements. Thus, TaG-

EM provides a flexible system for barcoding cells and organisms.

Methods

Drosophila stocks and maintenance

Drosophila stocks were grown at 22ºC on cornmeal agar unless otherwise indicated. The

following stocks were used in this study:

Stock name   **Source**

IsoD1   Clandinin Lab, Stanford University

w-; +; + (IsoD1)   Clandinin Lab, Stanford University

attP2 line   BDSC #25710

norpA   BDSC# 9048

UAS-myr::GFP (pJFRC12)   BDSC #32197

Hexameric GFP lines   BDSC #91402, #91403

dpp-Gal4 driver   BDSC #1553

Act-Gal4 driver   BDSC #4414

Tub-Gal4 driver   BDSC #5138

Mhc-Gal4 driver   BDSC #55132

PC-Gal4 driver   BDSC #91400 + #91401

EC-Gal4 driver   BDSC #91405

EB-Gal4 driver   BDSC #91398 + #91404

EE-Gal4 driver   BDSC #91402 + #91403

PMG-Gal4 driver   BDSC #91398 + #91399

TaG-EM lines (x20)   This study

TaG-EM lines + 6xGFP (x20)   This study

Design and cloning of TaG-EM constructs

A gBlock with the following sequence containing a part of the SV40 3' UTR with a PCR handle

(uppercase, below) and a 14 bp randomer sequence just upstream of the SV40 polyadenylation

site (bold and underlined, below) was synthesized (Integrated DNA Technologies, IDT):

caaaggaaaaagctgcactgctatacaagaaaattatggaaaaatatttgatgtatagtgccttgactagagatcataatcagccata

ccacatttgtagaggtttttacttgctttaaaaaacctcccacacctcccctgaacctgaaacataaaatgaatgcaattgttgttgttaact

tgtttattgcagcttataaCTTCCAACAACCGGAAGTGANNNNNNNNNNNNNNtggttaca<u>aataaa</u>gcaatag

catcacaaatttcaca<u>aataaa</u>gcattttttttcactg**ca**ttctagttgtggtttgtccaaactcatcaatgtatcttatcatgtctggatcgatct

ggccggccgtttaaacgaattcttgaagacgaaagggcctcgtgatacgcctattttttataggttaatgtcatgataataatg

The gBlock was resuspended in 20 µl EB, incubated at 50ºC for 20 minutes and then cut with

*Psi*I and *EcoR*I (New England Biolabs, NEB) using the following reaction conditions: 4 µl gBlock

DNA (35 ng), 2 µl 10x CustSmart buffer (NEB), 1 µl *Eco*RI enzyme (NEB), 1 µl *Psi*I enzyme

(NEB), and 12 µl nuclease-free water were mixed and incubated at 37ºC for 1 hour followed by

65ºC for 20 minutes to heat inactivate the restriction enzymes. pJFRC12-10XUAS-IVS-

myr::GFP plasmid (Addgene, Plasmid #26222) (Pfeiffer et al., 2010) was digested with the

following reaction conditions: 5 µl pJFRC12-10XUAS-IVS-myr::GFP plasmid DNA (~3 µg), 5 µl

10x CutSmart buffer (NEB), 1 µl *Psi*I enzyme (NEB), 1 µl *Eco*RI enzyme (NEB), and 38 µl

nuclease-free water, were mixed and incubated at 37ºC for 1 hour, followed by addition of 1 µl

of CIP and incubation for an additional 30 minutes. The digested vector backbone was gel

purified using the QiaQuick Gel Purification Kit (Qiagen). The digested gBlock was ligated into

the digested pJFRC12-10XUAS-IVS-myr::GFP backbone using the following reactions

conditions: 4 µl T4 ligase buffer (10x) (NEB), 20 µl plasmid backbone DNA (0.005 pmol), 5 µl

gBlock digest DNA (0.03 pmol), 2 µl of T4 DNA ligase (NEB), and 9 µl nuclease-free water were

mixed and incubated at 22ºC for 2 hours. 2 µl of the ligation reaction was transformed into 50 µl

of TOP10 competent cells (Invitrogen), and the cells were incubated on ice for 30 minutes, then

heat shocked at 42ºC for 30 seconds, and incubated on ice for 5 minutes. 250 µl SOC was

added and the cells were plated on LB+ampicillin plates and incubated overnight at 37ºC. DNA

was isolated from 36 pJFRC12-gBlock colonies using a QIAprep Spin MiniPrep kit (Qiagen).

Expected construct size was verified by diagnostic digest with *Eco*RI and *Apa*LI. DNA

concentration was determined using a Quant-iT PicoGreen dsDNA assay (Thermo Fisher

Scientific) and the randomer barcode for each of the constructs was determined by Sanger

sequencing using the following primers:

SV40_post_R: GCCAGATCGATCCAGACATGA

SV40_5F: CTCCCCCTGAACCTGAAACA

Generation of TaG-EM transgenic lines

29 sequence verified constructs were normalized, pooled evenly, and injected as a pool into

embryos (Rubin and Spradling, 1982) expressing PhiC31 integrase and the carrying the *attP2*

landing site (BDSC #25710). Injected flies were outcrossed to w- flies, and up to three *white*+

progeny per cross were identified, and the transgenic lines were homozygosed. DNA was

extracted (GeneJET genomic DNA purification Kit, Thermo Scientific) and the region containing

the DNA barcode was amplified with the following PCR reaction: 2.5 µl 1:10 diluted template

DNA, 2 µl 10x Reaction Buffer (Qiagen), 0.2 µl dNTP mix (10 µM), 1 µl 10 µM SV40_5F primer

(10 µM), 1 µl SV40_post_R primer (10 µM), 0.8 µl MgCl$_2$ (3 mM), 0.1 µl Taq polymerase

(Qiagen), 12.4 µl nuclease-free water. Reactions were amplified using the following cycling

conditions: 95ºC for 5 minutes, followed by 30 cycles of 94ºC for 30 seconds, 55ºC for 30

seconds, 72ºC for 30 seconds, followed by 72ºC for 5 minutes. PCR products were treated with

Exo-CIP using the following reaction conditions: 5 µl PCR product, 1 µl Exo-CIP Tube A (NEB),

1 µl Exo-CIP Tube B (NEB) were mixed and incubated at 37ºC for 4 minutes, followed by 80ºC

for 1 minute. The barcode sequence for each of the independent transgenic lines was

determined by Sanger sequencing using the SV40_5F and SV40_PostR primers. Transgenic

lines containing 20 distinct DNA barcodes were recovered (Supplemental Figure 1).

Optimizing amplification of TaG-EM barcodes for next-generation sequencing

The following primers were evaluated to amplify the TaG-EM barcodes upstream of NGS:

Forward primer pool; four primers with frameshifting bases to increase library sequence

diversity in initial sequencing cycles were normalized to 10 µM and pooled evenly to make a

B2_3'F1_Nextera_0-6 primer pool:

B2_3'F1_Nextera:

TCGTCGGCAGCGTCAGATGTGTATAAGAGACAGCTTCCAACAACCGGAAG*TGA

B2_3'F1_Nextera_2

TCGTCGGCAGCGTCAGATGTGTATAAGAGACAGAGCTTCCAACAACCGGAAG*TGA

B2_3'F1_Nextera_4

TCGTCGGCAGCGTCAGATGTGTATAAGAGACAGTCGACTTCCAACAACCGGAAG*TGA

B2_3'F1_Nextera_6

TCGTCGGCAGCGTCAGATGTGTATAAGAGACAGGAAGAGCTTCCAACAACCGGAAG*TGA

The following two reverse primers were tested:

SV40_pre_R_Nextera:

GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAGATTTGTGAAATTTGTGATGCTATTGC*T

TT

SV40_post_R_Nextera:

GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAGGCCAGATCGATCCAGACA*TGA

SV40_pre_R_nextera is designed to produce a shorter amplicon (200 bp with Illumina adapters

and indices added) and SV40-post_R_Nextera is designed to produce a longer amplicon (290

bp with Illumina adapters and indices added).

An initial test was performed with three different polymerases (Qiagen Q5, KAPA HiFi, and

Qiagen Taq) at two different annealing temperatures and with both the B2_3'F1_Nextera/

SV40_pre_R_Nextera and B2_3'F1_Nextera/ SV40_post_R_Nextera primer sets to determine

whether the primers amplify as expected (Supplemental Figure 1). Two different samples were

tested:

1. Pool of 8 putative transformant samples (pooled 5 ul each of 1:10 diluted sample)

2. OreR (wild type - diluted 1:10)

Set up the following PCR reactions:

Q5 polymerase: 2.5 µl template DNA, 1 µl 10 µM Forward primer (10 µM), 1 µl Reverse primer

(10 µM), 10 µl 2x Q5 Master Mix (Qiagen), 5.5 µl nuclease-free water. Reactions were amplified

using the following cycling conditions: 98ºC for 30 seconds, followed by 30 cycles of 98ºC for 20

seconds, 55ºC or 60ºC for 15 seconds, 72ºC for 30 seconds, followed by 72ºC for 5 minutes.

KAPA HiFi polymerase: 2.5 µl template DNA, 1 µl 10 µM Forward primer (10 µM), 1 µl Reverse

primer (10 µM), 10 µl 2x KAPA HiFi ReadyMix (Roche), 5.5 µl nuclease-free water. Reactions

were amplified using the following cycling conditions: 95ºC for 5 minutes, followed by 30 cycles

of 98ºC for 20 seconds, 55ºC or 60ºC for 15 seconds, 72ºC for 30 seconds, followed by 72ºC for

5 minutes.

Taq polymerase: 2.5 µl template DNA, 2 µl 10x Reaction Buffer (Qiagen), 0.2 µl dNTP mix (10

µM), 1 µl 10 µM Forward primer (10 µM), 1 µl Reverse primer (10 µM), 0.8 µl $MgCl_2$ (3 mM), 0.1

µl Taq polymerase (Qiagen), 12.4 µl nuclease-free water. Reactions were amplified using the

following cycling conditions: 95ºC for 5 minutes, followed by 30 cycles of 94ºC for 30 seconds,

55ºC or 60ºC for 30 seconds, 72ºC for 30 seconds, followed by 72ºC for 5 minutes.

Samples were run on a 2% agarose gel to verify amplification products (Supplemental Figure 1).

Next, the TaG-EM barcode lines were pooled in either an even or staggered manner. To

optimize reaction conditions for the barcode measurements, either 5 ng or 50 ng of DNA was

amplified in triplicate for each pool for either 20, 25, or 30 cycles with either KAPA HiFi using the

B2_Nextera_F 0-6 forward primer pool together with either the SV40_pre_R_Nextera or the

SV40_post_R_Nextera reverse primer. Next, PCR reactions were diluted 1:100 in nuclease-free

water and amplified in the following indexing reactions: 3 µl PCR 1 (1:100 dilution), 1 µl indexing

primer 1 (5 µM), 1 µl indexing primer 2 (5 µM), and 5 µl 2x Q5 master mix. The following

indexing primers were used (X indicates the positions of the 8 bp indices):

Forward indexing primer:

AATGATACGGCGACCACCGAGATCTACACXXXXXXXXTCGTCGGCAGCGTC

Reverse indexing primer:

CAAGCAGAAGACGGCATACGAGATXXXXXXXXGTCTCGTGGGCTCGG

Reactions were amplified using the following cycling conditions: 98°C for 30 seconds, followed

by 10 cycles of 98°C for 20 seconds, 55°C for 15 seconds, 72°C for 1 minute, followed by 72°C

for 5 minutes. Amplicons were then purified and normalized using a SequalPrep normalization

plate (Thermo Fisher Scientific), followed by elution in 20 µl of elution buffer. An even volume of

the normalized libraries was pooled and concentrated using 1.8x AmpureXP beads (Beckman

Coulter). Pooled libraries were quantified using a Qubit dsDNA high sensitivity assay (Thermo

Fisher Scientific) and libraries were normalized to 2 nM for sequencing on the Illumina MiSeq

(see below).

Structured fly pool experiments

Male or female flies from TaG-EM barcode lines were pooled in either an even or staggered

manner (Figure 2A). For the even pools, three independently pooled samples were constructed

in order to assess sample-to-sample variability. DNA was extracted from these structured pools

using a protocol adapted from Huang *et al.*, 2009 (Huang et al., 2009), using homemade SPRI

beads (DeAngelis et al., 1995) in the last purification step and amplified in triplicate using 2.5 µl

template DNA (50 ng), 1 µl 10 µM B2_Nextera_F 0-6 primers (10 µM), 1 µl

SV40_post_R_Nextera (10 µM), 10 µl 2x KAPA HiFi ReadyMix (Roche), 5.5 µl nuclease-free

water. Reactions were amplified using the following cycling conditions: 98ºC for 5 minutes,

followed by 30 cycles of 98ºC for 20 seconds, 60ºC for 15 seconds, 72ºC for 30 seconds,

followed by 72ºC for 5 minutes. Amplicons were indexed, normalized, quantified, and prepared

for sequencing as described above.

Phototaxis experiments

<u>Video-based measurements</u>

A pair of white LED strip lights with Muzata LED Strip Light Diffusers (U1SW WW 1M, LU1)

were mounted withing a light-tight box and controlled using an Vilros Uno Rev 3 microcontroller.

Test tubes containing flies were held in place with Acoustic Foam Panels (1"x12"x12",

ALPOWL). Videos and images were acquired using an Arducam 1080P Day & Night Vision

USB Camera with an IR filter and using Photo Booth software (Apple). Wild type and *norpA* flies

carrying one of 4 different TaG-EM barcodes were tested in three independent experimental

replicates. 20 male flies of each genotype were transferred into 25 mm x 150 mm glass test

tubes, incubated at 34ºC for 10 minutes and then run in the phototaxis assay, where a light at

one end of the chamber was turned on for 30 seconds. Videos of all tests were recorded

through the end of the 30 second light pulse. Videos were independently scored by two

observers to determine the number of flies in the light-facing or dark-facing tubes and the results

were averaged. A preference index (P.I.) was calculated using the following formula: [(number

of flies in light tube) - (number of flies in dark tube)]/(total number of flies).

<u>TaG-EM measurements</u>

For TaG-EM barcode-based phototaxis measurements, the following genotypes were

consolidated into a single test tube:

<u>Pool A</u>

norpA/Y;TaG-EM BC4/+

norpA/Y;TaG-EM BC3/+

w-/Y;TaG-EM BC2/+

w-/Y;TaG-EM BC1/+

<u>Pool B</u>

norpA/Y;TaG-EM BC2/+

norpA/Y;TaG-EM BC1/+

w-/Y;TaG-EM BC4/+

w-/Y;TaG-EM BC3/+

These pools were individually incubated at 34ºC for 10 minutes and then run in the phototaxis

assay. Videos of all tests were recorded and at the end of a 30 second light pulse the two test

tubes were quickly separated and capped. Flies in each of these tubes were counted, then DNA

was extracted from the flies from the light-facing or dark-facing tubes and amplified using 2.5 µl

template DNA (50 ng), 1 µl 10 µM B2_Nextera_F 0-6 primers (10 µM), 1 µl

SV40_post_R_Nextera (10 µM), 10 µl 2x KAPA HiFi ReadyMix (Roche), 5.5 µl nuclease-free

water. Reactions were amplified using the following cycling conditions: 95ºC for 5 minutes,

followed by 30 cycles of 98ºC for 20 seconds, 60ºC for 15 seconds, 72ºC for 30 seconds,

followed by 72ºC for 5 minutes. Amplicons were indexed, normalized, quantified, and prepared

for sequencing as described above.

Oviposition experiments

Newly hatched flies (males and females) from three barcode lines were collected at one-week

intervals during 4 consecutive weeks (12 barcode lines in total). Fresh fly food was provided

every 3-4 days. Ten days after the last collection, 10 females from each barcode line were taken

and pooled together in a collection cage (10 females x 12 barcode lines= 120 females). The

remaining females from each barcode line were separated from the males and put in individual

collection cages. Two days later, the experiment started and run for 3 consecutive days. Each

day a 1-1.5 hour pre-collection was followed by a 6 hour collection, both at 25ºC. 100 embryos

from each individual collection plate were transferred to new plates and incubated for 2 days at

18ºC. The number of hatched larvae were counted and used to calculate the egg survival rate.

The pooled collection plate was also incubated at 18ºC and the next day the embryos were

dechorionated and frozen. The 12 individual collection plates were kept at 4ºC and the number

of embryos counted in the following days. For the barcode measurements, DNA was extracted

from the embryos, and amplified using 2.5 µl template DNA (50 ng), 1 µl 10 µM B2_Nextera_F

0-6 primers (10 µM), 1 µl SV40_post_R_Nextera (10 µM), 10 µl 2x KAPA HiFi ReadyMix

(Roche), 5.5 µl nuclease-free water. Reactions were amplified using the following cycling

conditions: 95ºC for 5 minutes, followed by 30 cycles of 98ºC for 20 seconds, 60ºC for 15

seconds, 72ºC for 30 seconds, followed by 72ºC for 5 minutes. Amplicons were indexed,

normalized, quantified, and prepared for sequencing as described above.

Cell dissociation and isolation

Midguts from 3rd instar larvae were dissected in phosphate-buffered saline (PBS) and

transferred to microcentrifuge tubes on ice containing PBS + 30% normal goat serum (NGS).

After dissection, 150 µL of 2.7 mg/mL elastase was added to each sample tube. The tubes were

then incubated at 27°C for one hour. During incubation, samples were mixed by pipetting ~30

times every 15 minutes to improve elastase dissociation of the cells. Samples were then filtered

through a 40 µM FlowMi tip filter (Bel-Art) to reduce debris. Afterwards, the samples were

quantified on the LUNA-FL Dual Fluorescence Cell Counter (Logos Biosystems) using 9 µL of

sample to 1µL AO/PI dye to ensure there were enough viable cells for flow sorting.

Once quantified, the samples were brought up to a volume of ~1.1mL with the PBS + 30% NGS

solution to facilitate flow sorting. The samples were then fluorescently sorted on a FACSAria II

Cell Sorter (BD Biosciences) to isolate GFP+ cells. Following sorting, samples were centrifuged

at 300x g for 10 minutes to concentrate the cells. The supernatant was aspirated off until 50 µL

cell concentrate remained in each sample. Then, the samples were carefully resuspended using

wide bore pipette tips before being combined into one sample tube. This sample was quantified

on the LUNA-FL Dual Fluorescence Cell Counter (Logos Biosystems) as described above. If

necessary, cells were centrifuged, concentrated, and re-counted.

Preparation of single-cell sequencing libraries

The resulting pool was prepared for sequencing following the 10X Genomics Single Cell 3'

protocol (version CG000315 Rev C). Custom nested primers were used to attempt enrichment

of TaG-EM barcodes after cDNA creation using PCR. The remaining cDNA followed the 10X 3'

protocol without modification while the enriched TaG-EM portion followed BioLegends' "Total-

Seq-A Antibodies and Cell Hashing with 10X Single Cell 3' Reagents Kit v3 or v3.1" Protocol

starting at step III. The resulting 3' gene expression library and TaG-EM enrichment library were

sequenced together following Scenario 1 of the BioLegends protocol. Later, the TaG-EM library

was sequenced alone following Scenario 2 from the same protocol.

Sequencing

Libraries for TaG-EM barcode analysis from structured pools or from phototaxis or oviposition

experiments were denatured with NaOH and prepared for sequencing according to the

protocols described in the Illumina MiSeq Denature and Dilute Libraries Guides. Single-cell

libraries were sequenced on the Illumina NextSeq 2000 or Illumina NovaSeq 6000.

Data analysis

Behavioral experiments: Demultiplexed fastq files were generated using bcl2fastq or bcl-

convert. TaG-EM barcode data was analyzed using custom R and Python scripts and BioPython

(Cock et al., 2009). Leading primer sequences were trimmed using cutadapt (Martin, 2011) and

the first 14 bp of the remaining trimmed read were compared to a barcode reference file, with a

maximum of 2 mismatches allowed, using a custom script which is available via Github:

https://github.com/darylgohl/TaG-EM.

Single-cell experiments:

Data was initially mapped and analyzed using the Cellranger software package (version 7.0.1,

10X Genomics). A custom reference Drosophila genome was created by starting with the

BDGP6.28 reference genome assembly and Ensembl gene annotations. Custom gene

definitions for each of the TaG-EM barcodes were added to the fasta genome file and gtf gene

annotation file and a Cellranger reference package was generated with the cellranger mkref

command. Subsequent analysis was performed using Scanpy (version 1.9.3) (Wolf et al., 2018).

Cells expressing less than 200 genes and genes expressed in fewer than three cells were

filtered from the expression matrix. Next, cells expressing more than 4,000 genes, cells

expressing more than 15% mitochondrial transcripts, and cells expressing more than 30%

ribosomal (RpL) genes were filtered out. Expression counts were log normalized and highly

variable genes were filtered out (min_mean=0.0125, max_mean=3, min_disp=0.5). Cells

expressing multiple TaG-EM barcodes were identified and removed and cells were clustered

using the Leiden algorithm (Traag et al., 2019) and projected with UMAP. Scanpy was used to

call and visualize differentially expressed genes.

Declarations

Availability of data, code, and materials

Sequencing data for this project is available through the National Center for Biotechnology Information (NCBI) Sequence Read Archive BioProject PRJNA912199. Fly stocks containing each of the 20 TaG-EM barcodes together with an additional UAS hexameric GFP expression construct will be available from the Bloomington Drosophila Stock Center. The TaG-EM barcode analysis script and barcode reference fasta file are available via Github: https://github.com/darylgohl/TaG-EM.

Competing interests

None to declare.

Funding

This study was supported by a grant from the Winston and Maxine Wallin Neuroscience Discovery Fund.

Author contributions

J.B.M. conceived and designed experiments, conducted experiments, analyzed data, revised manuscript, M.D. conducted experiments, revised manuscript, L.G. conducted experiments, revised manuscript, B.A. conducted experiments, revised manuscript, J.G. analyzed data, revised manuscript, D.M.G. conceived and designed experiments, conducted experiments, analyzed data, wrote and revised the manuscript.

Acknowledgements

We thank our colleagues in the University of Minnesota Genomics Center (RRID:SCR_012413), in particular Aaron Becker, Dylan Cole, and Logan Silber for help with DNA sequencing, Emma Stanley, Fernanda Rodriguez, and Patrick Grady for assistance and advice on single-cell sequencing, and Kenneth Beckman, Andrew Alegria, Troy Louwagie, and Aaron Barnes for helpful feedback and discussions. This work was supported by the resources and staff at the University of Minnesota University Imaging Centers (RRID:SCR_020997). The authors acknowledge the Minnesota Supercomputing Institute (MSI) at the University of Minnesota for providing resources that contributed to the research results reported within this paper. URL:

http://www.msi.umn.edu. Stocks obtained from the Bloomington Drosophila Stock Center (NIH

P40OD018537) were used in this study.

References

Ariyapala IS, Holsopple JM, Popodi EM, Hartwick DG, Kahsai L, Cook KR, Sokol NS. 2020.
Identification of split-GAL4 drivers and enhancers that allow regional cell type
manipulations of the drosophila melanogaster intestine. *Genetics* **216**:891–903.
doi:10.1534/genetics.120.303625
Aso Y, Hattori D, Yu Y, Johnston RM, Iyer NA, Ngo TTB, Dionne H, Abbott LF, Axel R,
Tanimoto H, Rubin GM. 2014. The neuronal architecture of the mushroom body provides
a logic for associative learning. *eLife* **3**:e04577. doi:10.7554/ELIFE.04577
Bhang HC, Ruddy DA, Krishnamurthy Radhakrishna V, Caushi JX, Zhao R, Hims MM, Singh
AP, Kao I, Rakiec D, Shaw P, Balak M, Raza A, Ackley E, Keen N, Schlabach MR,
Palmer M, Leary RJ, Chiang DY, Sellers WR, Michor F, Cooke VG, Korn JM, Stegmeier
F. 2015. Studying clonal dynamics in response to cancer therapy using high-complexity
barcoding. *Nature Medicine* **21**:440–448. doi:10.1038/nm.3841
Chen X, Sun Y-C, Zhan H, Kebschull JM, Fischer S, Matho K, Huang ZJ, Gillis J, Zador AM.
2019. High-Throughput Mapping of Long-Range Neuronal Projection Using In Situ
Sequencing. *Cell* **179**:772-786.e19. doi:10.1016/j.cell.2019.09.023
Cheng J, Liao J, Shao X, Lu X, Fan X. 2021. Multiplexing Methods for Simultaneous Large-
Scale Transcriptomic Profiling of Samples at Single-Cell Resolution. *Advanced Science*
8:2101229. doi:10.1002/advs.202101229
Clark IC, Gutiérrez-Vázquez C, Wheeler MA, Li Z, Rothhammer V, Linnerbauer M, Sanmarco
LM, Guo L, Blain M, Zandee SEJ, Chao CC, Batterman KV, Schwabenland M, Lotfy P,
Tejeda-Velarde A, Hewson P, Polonio CM, Shultis MW, Salem Y, Tjon EC, Fonseca-
Castro PH, Borucki DM, de Lima KA, Plasencia A, Abate AR, Rosene DL, Hodgetts KJ,
Prinz M, Antel JP, Prat A, Quintana FJ. 2021. Barcoded viral tracing of single-cell
interactions in central nervous system inflammation. *Science* **372**.
Cock PJA, Antao T, Chang JT, Chapman BA, Cox CJ, Dalke A, Friedberg I, Hamelryck T, Kauff
F, Wilczynski B, de Hoon MJL. 2009. Biopython: freely available Python tools for
computational molecular biology and bioinformatics. *Bioinformatics (Oxford, England)*
25:1422–3. doi:10.1093/bioinformatics/btp163
David J, Cohet Y, Fouillet P. 1975. The variability between individuals as a measure of
senescence: A study of the number of eggs laid and the percentage of hatched eggs in
the case of Drosophila melanogaster. *Experimental Gerontology* **10**:17–25.
doi:10.1016/0531-5565(75)90011-X
Davis FP, Nern A, Picard S, Reiser MB, Rubin GM, Eddy SR, Henry GL. 2020. A genetic,
genomic, and computational resource for exploring neural circuit function. *eLife* **9**.
DeAngelis MM, Wang DG, Hawkins TL. 1995. Solid-phase reversible immobilization for the
isolation of PCR products. *Nucleic Acids Res* **23**:4742–4743.
Gohl DM, Silies MA, Gao XJ, Bhalerao S, Luongo FJ, Lin C-C, Potter CJ, Clandinin TR. 2011. A
versatile in vivo system for directed dissection of gene expression patterns. *Nature*
Methods **8**:231–237. doi:10.1038/nmeth.1561
Groth AC, Fish M, Nusse R, Calos MP. 2004. Construction of transgenic Drosophila by using
the site-specific integrase from phage phiC31. *Genetics* **166**:1775–82.
doi:10.1534/genetics.166.4.1775
Huang AM, Rehm EJ, Rubin GM. 2009. Quick preparation of genomic DNA from Drosophila.
Cold Spring Harb Protoc **2009**:pdb.prot5198. doi:10.1101/pdb.prot5198
Hung RJ, Hu Y, Kirchner R, Liu Y, Xu C, Comjean A, Tattikota SG, Li F, Song W, Sui SH,
Perrimon N. 2020. A cell atlas of the adult Drosophila midgut. *Proceedings of the*
National Academy of Sciences of the United States of America **117**:1514–1523.
doi:10.1073/pnas.1916820117

Ingham PW. 1988. The molecular genetics of embryonic pattern formation in Drosophila. *Nature*
720   **335**:25–34. doi:10.1038/335025a0
721   Klein AM, Mazutis L, Akartuna I, Tallapragada N, Veres A, Li V, Peshkin L, Weitz DA, Kirschner
722   MW. 2015. Droplet barcoding for single-cell transcriptomics applied to embryonic stem
723   cells. *Cell* **161**:1187–1201. doi:10.1016/j.cell.2015.04.044
724   Kurmangaliyev YZ, Yoo J, Valdes-Aleman J, Sanfilippo P, Zipursky SL. 2020. Transcriptional
725   Programs of Circuit Assembly in the Drosophila Visual System. *Neuron* **108**:1045-
726   1057.e6. doi:10.1016/J.NEURON.2020.10.006
727   Lee JH, Daugharthy ER, Scheiman J, Kalhor R, Ferrante TC, Terry R, Turczyk BM, Yang JL,
728   Lee HS, Aach J, Zhang K, Church GM. 2015. Fluorescent in situ sequencing (FISSEQ)
729   of RNA for gene expression profiling in intact cells and tissues. *Nature Protocols*
730   **10**:442–458. doi:10.1038/nprot.2014.191
731   Lein E, Borm LE, Linnarsson S. 2017. The promise of spatial transcriptomics for neuroscience
732   in the era of molecular cell typing. *Science* **358**:64–69. doi:10.1126/science.aan6827
733   Li H, Janssens J, De Waegeneer M, Kolluru SS, Davie K, Gardeux V, Saelens W, David FPA,
734   Brbić M, Spanier K, Leskovec J, McLaughlin CN, Xie Q, Jones RC, Brueckner K, Shim J,
735   Tattikota SG, Schnorrer F, Rust K, Nystul TG, Carvalho-Santos Z, Ribeiro C, Pal S,
736   Mahadevaraju S, Przytycka TM, Allen AM, Goodwin SF, Berry CW, Fuller MT, White-
737   Cooper H, Matunis EL, DiNardo S, Galenza A, O'Brien LE, Dow JAT, FCA Consortium,
738   Jasper H, Oliver B, Perrimon N, Deplancke B, Quake SR, Luo L, Aerts S. 2022. Fly Cell
739   Atlas: A single-nucleus transcriptomic atlas of the adult fruit fly. *Science* **375**:eabk2432.
740   doi:10.1126/science.abk2432
741   Ma D, Przybylski D, Abruzzi KC, Schlichting M, Li Q, Long X, Rosbash M. 2021. A
742   transcriptomic taxonomy of drosophila circadian neurons around the clock. *eLife* **10**:1–
743   19. doi:10.7554/eLife.63056
744   Macosko EZ, Basu A, Satija R, Nemesh J, Shekhar K, Goldman M, Tirosh I, Bialas AR,
745   Kamitaki N, Martersteck EM, Trombetta JJ, Weitz DA, Sanes JR, Shalek AK, Regev A,
746   McCarroll SA. 2015. Highly Parallel Genome-wide Expression Profiling of Individual
747   Cells Using Nanoliter Droplets. *Cell* **161**:1202–1214. doi:10.1016/j.cell.2015.05.002
748   Martin KC, Ephrussi A. 2009. mRNA localization: gene expression in the spatial dimension. *Cell*
749   **136**:719–30. doi:10.1016/j.cell.2009.01.044
750   Martin M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads.
751   *EMBnet.journal* **17**:10–12.
752   Marusyk A, Almendro V, Polyak K. 2012. Intra-tumour heterogeneity: a looking glass for
753   cancer? *Nature Reviews Cancer* **12**:323–334. doi:10.1038/nrc3261
754   McKenna A, Findlay GM, Gagnon JA, Horwitz MS, Schier AF, Shendure J. 2016. Whole-
755   organism lineage tracing by combinatorial and cumulative genome editing. *Science*
756   **353**:aaf7907. doi:10.1126/science.aaf7907
757   Namiki S, Dickinson MH, Wong AM, Korff W, Card GM. 2018. The functional organization of
758   descending sensory-motor pathways in drosophila. *eLife* **7**.
759   Pfeiffer BD, Jenett A, Hammonds AS, Ngo T-TB, Misra S, Murphy C, Scully A, Carlson JW,
760   Wan KH, Laverty TR, Mungall C, Svirskas R, Kadonaga JT, Doe CQ, Eisen MB,
761   Celniker SE, Rubin GM. 2008. Tools for neuroanatomy and neurogenetics in Drosophila.
762   *Proceedings of the National Academy of Sciences of the United States of America*
763   **105**:9715–20. doi:10.1073/pnas.0803697105
764   Pfeiffer BD, Ngo T-TB, Hibbard KL, Murphy C, Jenett A, Truman JW, Rubin GM. 2010.
765   Refinement of tools for targeted gene expression in Drosophila. *Genetics* **186**:735–55.
766   doi:10.1534/genetics.110.119917
767   Reeves GT, Muratov CB, Schüpbach T, Shvartsman SY. 2006. Quantitative Models of
768   Developmental Pattern Formation. *Developmental Cell* **11**:289–300.
769   doi:10.1016/J.DEVCEL.2006.08.006

Rubin GM, Spradling AC. 1982. Genetic transformation of Drosophila with transposable element
vectors. *Science (New York, NY)* **218**:348–53. doi:10.1126/science.6289436
Shearin HK, MacDonald IS, Spector LP, Steven Stowers R. 2014. Hexameric GFP and mCherry
reporters for the Drosophila GAL4, Q, and LexA transcription systems. *Genetics*
196:951–960. doi:10.1534/GENETICS.113.161141/-/DC1/GENETICS.113.161141-
5.PDF
Smith AM, Heisler LE, Mellor J, Kaper F, Thompson MJ, Chee M, Roth FP, Giaever G, Nislow
C. 2009. Quantitative phenotyping via deep barcode sequencing. *Genome research*
19:1836–42. doi:10.1101/gr.093955.109
Stoeckius M, Hafemeister C, Stephenson W, Houck-Loomis B, Chattopadhyay PK, Swerdlow H,
Satija R, Smibert P. 2017. Simultaneous epitope and transcriptome measurement in
single cells. *Nature Methods* **14**:865–868. doi:10.1038/nmeth.4380
Stoeckius M, Zheng S, Houck-Loomis B, Hao S, Yeung BZ, Mauck WM, Smibert P, Satija R.
2018. Cell Hashing with barcoded antibodies enables multiplexing and doublet detection
for single cell genomics. *Genome Biology* **19**:1–12. doi:10.1186/s13059-018-1603-1
Teleman AA, Cohen SM. 2000. Dpp gradient formation in the drosophila wing imaginal disc.
Cell **103**:971–980. doi:10.1016/S0092-8674(00)00199-9
Thompson CL, Ng L, Menon V, Martinez S, Lee C-K, Glattfelder K, Sunkin SM, Henry A, Lau C,
Dang C, Garcia-Lopez R, Martinez-Ferre A, Pombero A, Rubenstein JLR, Wakeman
WB, Hohmann J, Dee N, Sodt AJ, Young R, Smith K, Nguyen T-N, Kidney J, Kuan L,
Jeromin A, Kaykas A, Miller J, Page D, Orta G, Bernard A, Riley Z, Smith S, Wohnoutka
P, Hawrylycz MJ, Puelles L, Jones AR. 2014. A High-Resolution Spatiotemporal Atlas of
Gene Expression of the Developing Mouse Brain. *Neuron* **83**:309–323.
doi:10.1016/J.NEURON.2014.05.033
Traag VA, Waltman L, van Eck NJ. 2019. From Louvain to Leiden: guaranteeing well-connected
communities. *Sci Rep* **9**:5233. doi:10.1038/s41598-019-41695-z
van Opijnen T, Bodi KL, Camilli A. 2009. Tn-seq: high-throughput parallel sequencing for fitness
and genetic interaction studies in microorganisms. *Nature methods* **6**:767–72.
doi:10.1038/nmeth.1377
Venken KJT, Schulze KL, Haelterman NA, Pan H, He Y, Evans-Holm M, Carlson JW, Levis RW,
Spradling AC, Hoskins RA, Bellen HJ. 2011. MiMIC: a highly versatile transposon
insertion resource for engineering Drosophila melanogaster genes. *Nature methods*
8:737–43.
Wolf FA, Angerer P, Theis FJ. 2018. SCANPY: large-scale single-cell gene expression data
analysis. *Genome Biology* **19**:15. doi:10.1186/s13059-017-1382-0
Zheng GXY, Terry JM, Belgrader P, Ryvkin P, Bent ZW, Wilson R, Ziraldo SB, Wheeler TD,
McDermott GP, Zhu J, Gregory MT, Shuga J, Montesclaros L, Underwood JG,
Masquelier DA, Nishimura SY, Schnall-Levin M, Wyatt PW, Hindson CM, Bharadwaj R,
Wong A, Ness KD, Beppu LW, Deeg HJ, McFarland C, Loeb KR, Valente WJ, Ericson
NG, Stevens EA, Radich JP, Mikkelsen TS, Hindson BJ, Bielas JH. 2017. Massively
parallel digital transcriptional profiling of single cells. *Nature Communications* **8**:1–12.
doi:10.1038/ncomms14049
812

Figures

Figure 1. Overview of TaG-EM system. A) Schematic illustrating the design of the TaG-EM constructs, where a barcode sequence is inserted in the 3' UTR of a UAS-GFP construct and inserted in a specific genomic locus using PhiC31 integrase. B) Use of TaG-EM barcodes for sequencing-based population behavioral assays. C) Use of TaG-EM barcodes expressed with tissue-specific Gal4 drivers to label cell populations *in vivo* upstream of cell isolation and single-cell sequencing.

Figure 2. Structured pool tests. A) Overview of the construction of the structured pools for assessing the quantitative accuracy of TaG-EM barcode measurements. Male and female even pools were constructed and extracted in triplicate. The table shows the number of flies that were pooled for each experimental condition. B) Barcode abundance data for three independent replicates of the female even pool. C) Barcode abundance data for three independent replicates

of the male even pool. D) Barcode abundance data for the female staggered pool. Inset plot

shows the average observed barcode abundance among lines pooled at each level compared

to the expected abundance. E) Barcode abundance data for the male staggered pool. Inset plot

shows the average observed barcode abundance among lines pooled at each level compared

to the expected abundance. For all plots, bars indicate the mean barcode abundance for three

technical replicates of each pool, error bars are +/- S.E.M.

Preference Index (P.I.) = $\dfrac{(\text{Light} - \text{Dark})}{(\text{Light} + \text{Dark})}$

Figure 3. TaG-EM barcode-based behavioral measurements. A) TaG-EM barcode lines in

either a wild-type or *norpA* background were pooled and tested in a phototaxis assay. After 30

seconds of light exposure, flies in tubes facing the light or dark side of the chamber were

collected, DNA was extracted, and TaG-EM barcodes were amplified and sequenced. Barcode

abundance values were scaled to the number of flies in each tube and used to calculate a

preference index (P.I.). Average P.I. values for four different TaG-EM barcode lines in both the

wild-type and *norpA* backgrounds are shown (n=3 biological replicates, error bars are +/-

S.E.M.). B) The same eight lines used for the sequencing-based TaG-EM barcode

measurements were independently tested in the phototaxis assay and manually scored videos

were used to calculate a P.I. for each genotype. Average P.I. values for each line are shown

(n=3 biological replicates, error bars are +/- S.E.M.). C) Flies carrying different TaG-EM

barcodes were collected and aged for one to four weeks and then eggs were collected and egg

number and viability was manually scored for each line. In parallel the barcoded flies from each

timepoint were pooled, and eggs were collected, aged, and DNA was extracted, followed by

TaG-EM barcode amplification and sequencing. Average number of viable eggs per female

(manual counts) and average barcode abundance are shown both as a bar plot and scatter plot

(n=3 biological replicates for 3 barcodes per condition, error bars are +/- S.E.M.).

Figure 4. Gal4 driven expression of GFP from TaG-EM lines. A) Detailed view of the 3' UTR

of the TaG-EM constructs showing the position of the 14 bp barcode sequence (green highlight)

relative to the polyadenylation signal sequences (underlined) and polyA cleavage sites (red

highlights). pJFRC12 backbone schematic is from (Pfeiffer et al., 2010). B) Comparison of

endogenous GFP expression and GFP antibody staining in the wing imaginal disc for the

original pJFRC12 construct inserted in the attP2 landing site or for a TaG-EM barcode line

driven by *dpp-Gal4*. Wing discs are counterstained with DAPI. C) Expression of GFP from either

a TaG-EM barcode construct (left column), a hexameric GFP construct (middle column), or both

a line carrying both a TaG-EM barcode construct and a hexameric GFP construct (right column)

driven by the indicated gut driver line (PMG-Gal4: Pan-midgut driver; EC-Gal4: Enterocyte

driver; EE-Gal4: Enteroendocrine driver; EB-Gal4: Enteroblast driver).

Figure 5. Expression of TaG-EM genetic barcodes in larval intestinal cell types. A)

Number of doublets removed due to co-expression of each of the pairwise combinations of

TaG-EM barcodes. B) UMAP plot of Drosophila larval gut cell clusters after TaG-EM barcode-

based doublet removal. UMAP plots showing gene expression levels of enterocyte marker

genes C) *Jon99Ciii*, D) *betaTry*, E) *LManVI*, and F) the TaG-EM barcode (BC4) driven by the

EC-Gal4 line. UMAP plots showing gene expression levels of enteroblast/ISC marker genes G)

esg, H) *klu*, I) *E(spl)mbeta-HLH*, and J) the TaG-EM barcode (BC6) driven by the EB-Gal4 line.

UMAP plots showing gene expression levels of enteroendocrine cell marker genes K) *Dh31*, L)

IA-2, M) *Orcokinin*, and N) the TaG-EM barcode (BC9) driven by the EE-Gal4 line.

Supplemental Figures

A

Barcode sequence	Transgenic line
ATTTTGCGCCTTAG	1-2
CCCTAAAGTACCGT	4-3
TAGTTAGGCGGAAA	5-4
TTGCTCTTTCGTTT	6-4
TTGCTGCGCACGGC	7-1
CGTTTATATCCTGT	10-1
AGCAGGTGATGTAC	1-1, 1-3, 10-2
TATGCCTCATTTCT	13-2
GTTTGATCAGTACC	13-4
GGTTTACCACAAAG	14-2
AGACACATCCTGTG	15-1
GAATTTATCAGTTA	15-2
TCTTCATGACCAGC	2-1, 2-2, 2-3, 11-1, 11-2, 11-4
TATGTCAATATGGA	3-1, 3-2, 3-3, 9-2
ACTCTCTCTTAAGG	4-1, 4-4
CTTTAATTGCAGTC	5-3, 5-5, 9-1, 9-3
TACTGTACATGCTC	6-2, 10-4, 14-3, 14-4
TACCTTTGAATTCG	6-3, 12-2, 12-4
CATCGAGTTCACCT	8-1, 12-3
GTCCTTCTACCCCG	8-2, 8-4

B

Supplemental Figure 1. Sanger sequencing identification of TaG-EM barcode lines. A)
Summary of barcode pool injections. Barcode sequence and transgenic vial identifier in which
the barcode was identified are shown. B) Sanger sequencing-based confirmation of the barcode
sequence and PCR handle in TaG-EM transgenic lines.

Supplemental Figure 2. Optimization of TaG-EM barcode amplification. A) Gels showing

bands produced when amplifying TaG-EM flies or a wild type control with the indicated

polymerase, annealing temperature, and primer pair (short = B2_3'F1_Nextera/

SV40_pre_R_Nextera; long = B2_3'F1_Nextera/ SV40_post_R_Nextera). B-E) Mean error

886   (R.M.S.D. root mean squared deviation from expected value) for even pool amplified with the

887   indicated primer set, input amount, and cycle number using KAPA HiFi polymerase (n=3, error

888   bars are +/- S.E.M.). F-G) Mean error (R.M.S.D. root mean squared deviation from expected

889   value) for staggered pool amplified with the indicated primer set, input amount, and cycle

890   number using KAPA HiFi polymerase (n=3, error bars are +/- S.E.M.).

Supplemental Figure 3. Oviposition tests with TaG-EM barcode lines. Plots showing mean

TaG-EM barcode abundance for adult females used in oviposition experiments (top) and eggs

collected from these females (bottom). Data from two independent trials is shown (n=3 for each

trial, error bars are +/- S.E.M.).

Supplemental Figure 4. Fecundity data for individual TaG-EM lines. Manually collected data for mean number of viable eggs per female, barcode abundance data, and barcode abundance data normalized to adult fly barcode data for each of the TaG-EM barcode lines used in the age-dependent fecundity experiment. Scatterplots show correlations between manually collected data and barcode sequencing results. Data from two independent trials is shown (n=3 for each trial, error bars are +/- S.E.M.).

906 **Supplemental Figure 5. Average age-dependent fecundity data for Trial 1.** Average number

907 of viable eggs per female (manual counts) and average barcode abundance are shown both as

908 a bar plot and scatter plot (n=3 biological replicates for 3 barcodes per condition, error bars are

909 +/- S.E.M.). Data from Trial 2 is shown in Figure 3C.

Supplemental Figure 6. Expression driven by *dpp-Gal4* for 20 TaG-EM lines. GFP antibody
staining in the wing imaginal disc for the indicated TaG-EM barcode line driven by *dpp-Gal4*.
Wing discs are counterstained with DAPI.

A

B

Supplemental Figure 7. TaG-EM line GFP expression driven by different Gal4 drivers. A) Comparison of GFP expression in larvae for the original pJFRC12 construct inserted in the attP2 landing site (left) or for a TaG-EM barcode line (right) expressed under the control of the indicated driver line. B) GFP expression of the PC-Gal (Precursor-Gal4) driver line together with either UAS-2xGFP or a combination of UAS-2xGFP and a TaG-EM barcode line.

922 **Supplemental Figure 8. Dissociated intestinal cell viability.** A) GFP expression visualized in

923 dissociated cells from gut driver lines crossed to hexameric GFP and TaG-EM line. B)

924 Proportion of live (left) and dead (right) cells post-isolation and flow sorting as assessed by GFP

925 expression and propidium iodide staining.

Supplemental Figure 9. Expression of TaG-EM genetic barcodes in larval intestinal precursor cells. UMAP plots showing gene expression levels of A) enteroblast/ISC marker genes *esg*, *klu*, and *E(spl)mbeta-HLH*; and B) the TaG-EM barcodes 7, 8, and 9 driven by the PC-Gal4 line.

Supplemental Figure 10. Identification of doublets based on co-expression of TaG-EM barcodes. UMAP plots of all 28 pairwise barcode combinations showing cells co-expressing both of the indicated barcodes in blue (True).

937 **Supplemental Figure 11. Differentially expressed genes** Dotplot showing the top three

938 differentially expressed genes by log-fold change for each cluster.

Supplemental Figure 12. Top 25 marker genes for each cell cluster. Plots showing t-test-

based rankings of the top 25 differentially expressed genes for each cluster.

Supplemental Figure 13. Expression of the PMG-Gal4 driven TaG-EM barcodes. UMAP plots showing expression of the four PMG-Gal4 driven TaG-EM barcodes (BC1, BC1, BC3, and BC7).