1
12 **Deterministic Genetic Barcoding for Multiplexed Behavioral and Single-Cell**
13 **Transcriptomic Studies**
14
15

16 Jorge Blanco Mendana[1], Margaret Donovan[1], Lindsey G. O'Brien[1], Benjamin Auch[1,†], John
17 Garbe[1], Daryl M. Gohl[1,2,3]
18
19 [1] University of Minnesota Genomics Center, Minneapolis, MN, 55455, USA
20 [2] Department of Genetics, Cell Biology, and Development, University of Minnesota, Minneapolis,
21 MN 55455, USA
22
23
24 [3] Corresponding author: dmgohl@umn.edu
25 [†] Present address: Phase Genomics, Seattle, WA, 98109, USA
26
27

Abstract

Advances in single-cell sequencing technologies have provided novel insights into the dynamics of gene expression throughout development, been used to characterize somatic variation and heterogeneity within tissues, and are currently enabling the construction of transcriptomic cell atlases. However, despite these remarkable advances, linking anatomical information to transcriptomic data and positively identifying the cell types that correspond to gene expression clusters in single-cell sequencing data sets remains a challenge. We describe a straightforward genetic barcoding approach that takes advantage of the powerful genetic tools available in Drosophila to allow *in vivo* tagging of defined cell populations. This method, called Targeted Genetically-Encoded Multiplexing (TaG-EM), involves inserting a DNA barcode just upstream of the polyadenylation site in a Gal4-inducible *UAS-GFP* construct so that the barcode sequence can be read out during single-cell sequencing, labeling a cell population of interest. By creating many such independently barcoded fly strains, TaG-EM will enable a number of potential applications that will improve the quality and information content of single-cell transcriptomic data including positive identification of cell types in cell atlas projects, identification of multiplet droplets, and barcoding of experimental timepoints, conditions, and replicates. Furthermore, we demonstrate that the barcodes from TaG-EM fly lines can be read out using next-generation sequencing to facilitate population-scale behavioral measurements. Thus, TaG-EM has the potential to enable large-scale behavioral screens in addition to improving the ability to reliably annotate cell atlas data, expanding the scope, and improving the robustness of single-cell transcriptomic experiments.

Introduction

Spatially and temporally regulated gene expression patterns are a hallmark of multicellular life and function to orchestrate patterning, growth, and differentiation throughout development (Ingham, 1988; Reeves et al., 2006). In mature organisms, spatial expression patterns both in tissues and within cells define functionally distinct compartments and determine many aspects of cellular and organismal physiology (Martin and Ephrussi, 2009). In addition, such expression patterns differentiate healthy and diseased tissue and impact disease etiology (Marusyk et al., 2012). Spatial and temporal expression patterns, which can be used to distinguish between cell types and provide insight into cellular function, also provide a means to understand the organization and physiology of complex tissues such as the brain (Thompson et al., 2014). Thus, robust and scalable tools for measuring spatial and temporal gene expression patterns at a genome-wide scale and at high resolution would be transformative research tools across many biological disciplines.

Single-cell sequencing technologies have provided insights into the dynamics of gene expression throughout development, been used to characterize somatic variation and heterogeneity within tissues, and are currently enabling the construction of transcriptomic cell atlases (Klein et al., 2015; Macosko et al., 2015; Zheng et al., 2017). However, linking anatomical information to transcriptomic data and positively identifying the cell types that correspond to gene expression clusters in single-cell sequencing data sets remains a challenge. The cellular identities of gene expression clusters identified in cell atlas data sets are typically inferred from the expression of distinctive gene sets (Hung et al., 2020; Li et al., 2022; Ma et al., 2021), and the lack of positive identification of gene expression clusters introduces an element of uncertainty in this analysis. Moreover, this process of manual annotation is labor-intensive and often requires additional experiments to determine or confirm the expression patterns of marker genes. Emerging spatial genomics technologies hold promise in linking anatomical and transcriptomic information (Lee et al., 2015; Lein et al., 2017). Several of the emerging commercial spatial genomics technologies rely on *in situ* sequencing of marker genes allowing droplet-based single-cell transcriptomic data to be mapped onto a tissue. However, these technologies currently suffer from constraints related to cost, content, or applicability to specific model systems.

In addition to descriptive cell atlas projects, studies involving multiple experimental timepoints throughout development and aging, or studies assessing the effects of experimental exposures or genetic manipulations would benefit from increased ability to multiplex samples. Given the fixed costs of droplet-based single-cell sequencing, generating data for single-cell transcriptomic time courses or experimental manipulations can be costly. Outside of descriptive

studies, these costs are also a barrier to including replicates to assess biological variability;
consequently, a lack of biological replicates derived from independent samples is a common
shortcoming of single-cell sequencing experiments. Antibody-based cell hashing or feature
barcoding approaches have been developed to allow multiplexing of samples in droplet-based
single-cell sequencing reactions (Stoeckius et al., 2018, 2017). In addition, other multiplexing
strategies for single-cell sequencing based on alternative methods for tagging cells (Cheng et al.,
2021) or making use of natural genetic variation have been used (Kurmangaliyev et al., 2020).
While such approaches can reduce per-sample costs, typically samples are barcoded at a
population level and thus do not enable labeling of cell subpopulations within a sample.
We have developed a straightforward genetic barcoding approach that takes advantage
of the powerful genetic tools available in Drosophila to allow deterministic *in vivo* tagging of
defined cell populations. This method, called Targeted Genetically-Encoded Multiplexing (TaG-
EM), involves inserting a DNA barcode just upstream of the poly-adenylation site in a Gal4-
inducible *UAS-GFP* construct so that the barcode sequence can be read out during droplet-
based single-cell sequencing, labeling a cell population of interest.
Genetic barcoding approaches have been employed in many unicellular systems, cell
culture, and viral transfection to facilitate high throughput screening using sequencing-based
readouts (Bhang et al., 2015; Smith et al., 2009; van Opijnen et al., 2009). In multicellular
animals, techniques such as GESTALT have enabled lineage tracing by using CRISPR to
create unique barcodes in differentiating tissue (McKenna et al., 2016), and barcode
sequencing has also been employed to map connectivity in the brain (Chen et al., 2019).
Genetically barcoded fly lines can also be used to enable highly multiplexed behavioral
assays which can be read out using high throughput sequencing. Flies carrying TaG-EM
barcodes can be exposed to different experimental perturbations and then tested in assays
where flies, larvae, or embryos are fractionated based on behavioral outcomes or other
phenotypes. Thus, TaG-EM has the potential to enable large-scale next-generation sequencing
(NGS)-based behavioral or other fractionation screens analogous to BAR-Seq or Tn-Seq
approaches employed in microbial organisms.
112

Results

TaG-EM: A novel genetic barcoding strategy for multiplexed behavioral and single-cell transcriptomics

We cloned a fragment containing a PCR handle sequence and a diverse 14 bp barcode sequence into the SV40 3' untranslated region (UTR) sequence just upstream of the polyadenylation sites in the 10xUAS-myr::GFP (pJFRC12, (Pfeiffer et al., 2010)) backbone (Figure 1A). A pool containing 29 unique barcode-containing plasmids was injected into Drosophila embryos for PhiC31-mediated integration into the attP2 landing site (Groth et al., 2004) and transgenic lines were isolated and confirmed by Sanger sequencing (Figure 1B, Supplemental Figure 1). We recovered 20 distinctly barcoded Drosophila lines, with some barcodes recovered from multiple crosses (Supplemental Figure 1). Such barcoded fly lines have the potential to enable population behavioral measurements, where different exposures, experimental timepoints, and genetic or neural perturbations can be multiplexed and analyzed by measuring barcode abundance in sequencing data (Figure 1C). In addition, the barcodes, which reside on a Gal4-inducible *UAS-GFP* construct, can be expressed tissue-specifically and read out during droplet-based single-cell sequencing, labeling a cell population and/or an experimental condition of interest (Figure 1D).

Testing the accuracy and reproducibility of TaG-EM behavioral measurements using structured pools

We conducted initial experiments to optimize amplification of the genetic barcodes using primers targeting the PCR handle inserted just upstream of the 14 bp barcode sequence and PCR primers downstream of the TaG-EM barcode in the SV40 3' UTR sequence (Supplemental Figure 2). To test the accuracy and reproducibility of sequencing-based measurements of TaG-EM barcodes, we constructed structured pools containing defined numbers of flies pooled either evenly with each of the 20 barcode constructs comprising 5% of the pool, or in a staggered manner with sets of barcodes differing in abundance in 2-fold increments (Figure 2A). To examine the impact of technical steps such as DNA extraction and PCR amplification on TaG-EM barcode measurements, even pools were made and extracted in triplicate and amplicon sequencing libraries were made in triplicate for each independently extracted DNA sample for both the even and staggered pools. The resulting data indicated that TaG-EM measurements are highly accurate and reproducible. Technical replicates (indicated by error bars in Figure 2B-E) showed minimal variability. Likewise, the three independently extracted replicates of the even

pools produced consistent data with all 20 barcodes detected at levels close to the expected 5%

abundance (Figure 2B-C). Barcode abundance values for the staggered structured pools was

generally consistent with the input values and in most cases, the 2-fold differences between the

different groups of barcodes could be distinguished (Figure 2D-E). The coefficients of variation

were largely consistent for groups of TaG-EM barcodes pooled evenly or at different levels

within the staggered pools (Supplemental Figure 3). For the staggered pools, abundances

correlated well with the expected values, particularly when multiple barcodes for an input level

were averaged, in which case R^2 values were >0.99 (Figure 2D-E, inset plots). This indicates

that a high level of quantitative accuracy can be attained using sequencing-based analysis of

TaG-EM barcode abundance, particularly when averaging data for three to four independent

barcodes for an experimental condition.

TaG-EM measurement of phototaxis behavior correlate well with video-based

measurements

Next, we tested whether TaG-EM could be used to measure a phototaxis behavior. A

mixture of barcoded wild type or blind *norpA* mutant flies were run together through a phototaxis

assay. At the end of a period of light exposure, test tubes facing toward or away from the light

were capped, DNA was isolated, and barcodes were amplified and sequenced for each tube.

Raw read counts were scaled in proportion to the number of flies per tube and a preference

index was calculated for each barcode (Figure 3A). In parallel, individual preference indices

were calculated based on manual scoring of videos recorded for each line (Figure 3B).

Preference indices calculated for the pooled, NGS-based TaG-EM measurements were nearly

identical to conventional behavioral measurements for both wild type and *norpA* mutants (Figure

3A-B).

TaG-EM measurement of oviposition behavior and age-dependent fecundity

We next tested whether NGS-based pooled measurements of egg laying could be made.

Fertilized females from each of the 20 barcode lines were placed together in egg laying cups,

embryos were collected, aged for 12 hours to enable cell numbers to stabilize in the developing

eggs, and then DNA was extracted from both the pooled adult flies and the embryos. In general,

TaG-EM measurements of oviposition correlated with fly numbers, with the exception of

barcode 14 which had reduced barcode abundance across multiple trials (Supplemental Figure

4). This suggests that despite the fact that the genetic barcode constructs are inserted in a

common landing site, differences with respect to specific behaviors may exist among the lines,

and thus one should test to make sure given lines are appropriate to use in specific behavioral

assays.

To determine whether TaG-EM could be used to measure age-dependent fecundity, we

collected flies from twelve different TaG-EM barcode lines at four time points separated by one

week (three barcode lines per timepoint). We collected eggs from these fly lines individually and

scored the number of viable eggs per female. Next, we pooled the barcoded flies from all

timepoints and collected eggs from the pooled flies. These eggs were aged, DNA was

extracted, and the TaG-EM barcodes were amplified and sequenced. While measurements from

individual barcode lines were noisy, both for manual counts and sequencing based

measurements, there was a general trend toward declining fecundity over time (Supplemental

Figure 5), consistent with published reports (David et al., 1975). Manually scored viable egg

numbers and TaG-EM barcode abundances were well correlated across two independent

experimental trials (R^2 values of 0.52-0.61 for Trial 1 and 0.74-0.84 for Trial 2). When barcodes

from each individual timepoint were averaged, R^2 values for the correlation between manual and

sequencing-based measurements were 0.95 for Trial 1 and 0.99 for Trial 2 (Figure 3C,

Supplemental Figure 6).

Quantifying food transit time in the larval gut using TaG-EM

Gut motility defects underlie a number of functional gastrointestinal disorders in humans

(Keller et al., 2018). To study gut motility in Drosophila, we have developed an assay based on

the time it takes a food bolus to transit the larval gut (Figure 4A), similar to approaches that

have been employed for studying the role of the microbiome in human gut motility (Asnicar et

al., 2021). Third instar larvae were starved for 90 minutes and then fed food containing a blue

dye. After 60 minutes, larvae in which a blue bolus of food was visible were transferred to plates

containing non-dyed food, and food transit (indicated by loss of the blue food bolus) was scored

every 30 minutes for five hours (Supplemental Figure 7).

Because this assay is highly labor-intensive and requires hands-on effort for the entire

five-hour observation period, there is a limit on how many conditions or replicates can be scored

in one session (~8 plates maximum). Thus, we decided to test whether food transit could be

quantified in a more streamlined and scalable fashion by using TaG-EM (Figure 4B). Using the

manual assay, we observed that while caffeine-containing food is aversive to larvae, the

presence of caffeine reduces transit time through the gut (Figure 4C, Supplemental Figure 7).

This is consistent with previous observations in adult flies that bitter compounds (including

caffeine) activate enteric neurons via serotonin-mediated signaling and promote gut motility

215 (Yao and Scott, 2022). We tested whether TaG-EM could be used to measure the effect of

216 caffeine on food transit time in larvae. As with prior behavioral tests, the TaG-EM data

217 recapitulated the results seen in the manual assay (Figure 4D). Conducting the transit assay via

218 TaG-EM enables several labor-saving steps. First, rather than counting the number of larvae

219 with and without a food bolus at each time point, one simply needs to transfer non-bolus-

220 containing larvae to a collection tube. Second, because the TaG-EM lines are genetically

221 barcoded, all the conditions can be tested at once on a single plate, removing the need to

222 separately count each replicate of each experimental condition. This reduces the hands-on time

223 for the assay to just a few minutes per hour. A summary of the anticipated cost and labor

224 savings for the TaG-EM-based food transit assay is shown in Supplemental Figure 8.

226 **Tissue-specific expression of TaG-EM GFP constructs**

227  To facilitate representation of the TaG-EM barcodes in single-cell sequencing data,

228 genetic barcodes were placed just upstream of the polyadenylation signal sequences and polyA

229 cleavage sites (Figure 1A). To verify that the inserted sequences did not interfere with Gal4-

230 driven GFP expression, we crossed each of the barcoded TaG-EM lines to *decapentaplegic-*

231 *Gal4* (*dpp-Gal4*). We observed GFP expression in the expected characteristic central stripe

232 (Teleman and Cohen, 2000) in the wing imaginal disc for 19/20 lines at similar expression levels

233 to the base pJFRC12 *UAS-myr::GFP* construct inserted in the same landing site (Figure 5A,

234 Supplemental Figure 9). No GFP expression was visible for TaG-EM barcode number 8, which

235 upon molecular characterization had an 853 bp deletion within the GFP coding region (data not

236 shown). We generated and tested GFP expression of an additional 156 TaG-EM barcode lines

237 (Alegria et al., 2024), by crossing them to Mhc-Gal4 and observing expression in the adult

238 thorax. All 156 additional TaG-EM lines had robust GFP expression (data not shown). Gal4-

239 driven expression levels of TaG-EM barcoded GFP constructs were also similar to that of the

240 pJFRC12 base construct for multiple driver lines (Supplemental Figure 10) indicating that the

241 presence of the barcode does not generally impair expression of GFP.

243 **Boosting the GFP signal of TaG-EM constructs to enable robust cell sorting**

244  While with some driver lines, expression of the myr::GFP from the TaG-EM construct

245 may be too weak to allow robust enrichment of the tagged cells, adding an additional hexameric

246 GFP construct (Shearin et al., 2014) could boost expression of weak driver lines to levels that

247 are sufficient for robust detection of labeled flies or larvae (Figure 5B) and for labeling of

248 dissociated cells for flow cytometry (Supplemental Figure 11). Stocks with an additional UAS

249   hexameric GFP construct recombined onto the same chromosome as the TaG-EM construct
250   have been established for 20 TaG-EM barcode lines.
251
252   **Correlation between expression of TaG-EM barcodes and intestinal cell marker genes in**
253   **single-cell sequencing data**
254   To test whether we could detect TaG-EM barcodes in single-cell sequencing data, we
crossed three TaG-EM barcode lines to two different gut Gal4 driver lines (Ariyapala et al.,
2020), one expressing in the enterocytes (EC-Gal4: TaG-EM barcodes 1, 2, and 3) and the
other in intestinal precursor cells (PC-Gal4: TaG-EM barcodes 7, 8, and 9), which includes stem
cells and enteroblasts (EBs). Due to weak GFP expression with the EC-Gal4 driver, we did not
see visible GFP positive cells for this driver line. The PC-Gal4 driver line contained an additional
UAS-Stinger (2xGFP) construct and expressed GFP at a level sufficient for flow sorting when
crossed to the TaG-EM line (Supplemental Figure 12). Larval guts were dissected, dissociated,
stained with propidium iodide (PI) to label dead cells, and flow sorted to recover PI negative and
GFP positive cells. Approximately 10,000 cells were loaded into a 10x Genomics droplet
generator and a single-cell library was prepared and sequenced. Two clusters were observed in
the resulting sequencing data, one of which had high read counts from mitochondrial genes
suggesting that this cluster consisted of mitochondria, debris, or dead and dying cells. After
filtering the cells with high mitochondrial reads, a single cluster remained (Supplemental Figure
13). This cluster expressed known intestinal precursor cell markers such as *escargot* (*esg*),
klumpfuss (*klu*), and *Notch* pathway genes like *E(spl)mbeta-HLH* (Supplemental Figure 13).
Expression of all three PC-Gal4 driven TaG-EM barcodes was observed in this cluster
(Supplemental Figure 13) indicating that TaG-EM barcodes can be detected in single-cell
sequencing data. Interestingly, TaG-EM barcode 8, for which no GFP expression was observed,
was represented in the single-cell sequencing data indicating that the lesion in the GFP coding
region does not prevent mRNA expression for this line.
275   A previous study used droplet-based single-cell sequencing to characterize the cell
types that make up the adult midgut (Hung et al., 2020). This study took advantage of two
fluorescent protein markers, an *escargot* (*esg*)-GFP fusion protein and a *prospero* (*pros*)-Gal4
driven RFP to label the intestinal stem cells (ISCs) and enteroendocrine cells (EEs),
respectively (Hung et al., 2020). The authors compared the resulting clusters to a list of known
marker genes in the literature, including antibody staining, GFP, LacZ, and Gal4 reporter
expression patterns to classify the cells in individual clusters, and also found that the *esg*-GFP
expression was present in a broader subset of cells than anticipated. Thus, most of these cell

classifications relied upon inference as opposed to direct positive labeling. Recently, a large
collection of split-*Gal4* lines were screened for expression in the adult and larval gut (Ariyapala
et al., 2020). These include pan-midgut driver lines, split-*Gal4* lines specific for the EBs, ECs,
EEs, and ISC/EBs, as well as driver lines with regionalized gene expression. We crossed four
different TaG-EM barcode lines with the pan-midgut driver (PMG-Gal4: TaG-EM barcodes 1, 2,
3, and 7), and one barcode line to each of the precursor cell (PC-Gal4: TaG-EM barcode 5),
enterocyte (EC-Gal4: TaG-EM barcode 4), enteroblast (EB-Gal4: TaG-EM barcode 6) and
enteroendocrine (EE-Gal4: TaG-EM barcode 9) drivers (Ariyapala et al., 2020). Larval guts were
dissected from these lines and cells were dissociated, flow sorted as described above to select
live, GFP positive cells, and approximately 30,000 cells were loaded into a 10x Genomics
droplet generator for single-cell sequencing (Supplemental Figure 14). Using the additional
hexameric GFP construct to boost GFP expression resulted in visible fluorescent signal for all
eight barcode Gal4 line combinations.
An advantage of cell barcoding both for cell hashing (Stoeckius et al., 2018) and for
TaG-EM *in vivo* barcoding is that such labeling facilitates the identification and removal of
multiplets, which are an artifact of droplet-based single-cell sequencing approaches. After
filtering and removing cells with a high percentage of mitochondrial or ribosomal reads, we used
DoubletFinder (McGinnis et al., 2019) to computationally identify multiplet droplets. In parallel,
we searched for cells that co-expressed multiple TaG-EM barcodes. DoubletFinder identified
2,019 multiplet droplets, while TaG-EM barcodes identified 298 such droplets, 198 of which
(66.4%) overlapped with those identified by DoubletFinder (Supplemental Figure 15). Thus,
TaG-EM help identify an additional 100 doublets that would have otherwise been overlooked
using computational doublet identification methods.
After doublet removal, the remaining cells were clustered (Figure 6A, Supplemental
Figure 16) and analyzed using Seurat (Satija et al., 2015). Analysis of differentially expressed
genes identified clusters expressing marker genes previously reported for adult gut cell types
(Hung et al., 2020). These included genes associated with precursor cells (*Notch* pathway
genes), enterocytes (trypsins, serine proteases, amalyse, mannosidases), and enteroendocrine
cells (neuropeptides and neuropeptide receptors) (Supplemental Figure 17, data not shown).
TaG-EM barcodes derived from the eight multiplexed genotypes were observed in
approximately one-quarter of the cells (Supplemental Figure 17).
In antibody-conjugated oligo cell hashing approaches, sparsity of barcode representation
is overcome by spiking in an additional primer at the cDNA amplification step and amplifying the
hashtag oligo by PCR. We employed a similar approach to attempt to enrich for TaG-EM

barcodes in an additional library sequenced separately from the 10x Genomics gene expression

library. Our initial attempts at barcode enrichment using spike-in and enrichment primers

corresponding to the TaG-EM PCR handle were unsuccessful (Supplemental Figure 18).

However, we subsequently optimized the TaG-EM barcode enrichment by 1) using a longer

spike-in primer that more closely matches the annealing temperature used during the 10x

Genomics cDNA creation step, and 2) using a nested PCR approach to amplify the cell-barcode

and unique molecular identifier (UMI)-labeled TaG-EM barcodes (Supplemental Figure 18).

Using the enriched library, TaG-EM barcodes were detected in nearly 100% of the cells

at high sequencing depths (Supplemental Figure 19). However, although we used a polymerase

that has been engineered to have high processivity and that has been shown to reduce the

formation of chimeric reads in other contexts (Gohl et al., 2016), it is possible that PCR

chimeras could lead to unreliable detection events for some cells. Indeed, many cells had a

mixture of barcodes detected with low counts and single or low numbers of associated UMIS.

To assess the reliability of detection, we analyzed the correlation between barcodes detected in

the gene expression library and the enriched TaG-EM barcode library as a function of the purity

of TaG-EM barcode detection for each cell (the percentage of the most abundant detected TaG-

EM barcode, Supplemental Figure 19). For TaG-EM barcode detections where the most

abundance barcode was a high percentage of the total barcode reads detected (~75%-99.99%),

there was a high correlation between the barcode detected in the gene expression library and

the enriched TaG-EM barcode library. Below this threshold, the correlation was substantially

reduced.

In the enriched library, we identified 26.8% of cells with a TaG-EM barcode reliably

detected, a very modest improvement over the gene expression library alone (23.96%),

indicating that at least for this experiment, the main constraint is sufficient expression of the

TaG-EM barcode and not detection. To identify TaG-EM barcodes in the combined data set, we

counted a positive detection as any barcode either identified in the gene expression library or

any barcode identified in the enriched library with a purity of >75%. In the case of conflicting

barcode calls, we assigned the barcode that was detected directly in the gene expression

library. This increased the total fraction of cells where a barcode was identified to approximately

37% (Figure 6B).

As expected, the barcodes expressed by the pan-midgut driver were broadly distributed

across the cell clusters (Supplemental Figure 20). However, the number of cells recovered

varied significantly among the four pan-midgut driver associated barcodes. Expression of the

cell type-specific barcodes showed more restricted patterns of expression among the cell

clusters and were co-localized with known marker genes for these cell types (Figure 6C-N). For

instance, TaG-EM barcode 6, driven by the EB-Gal4 line, was expressed primarily in cells that

were annotated as enteroblasts (Figure 6C-D) and that expressed precursor cell markers such

as *esg* (Figure 6E), *klu* (Figure 6F), and Notch pathway genes such as *e(spl)mbeta-HLH* and

e(spl)m3-HLH (not shown).

TaG-EM barcode 4 expression, which was driven by the EC-Gal4 line, was seen

primarily in a cluster that was annotated as enterocytes (Figure 6G-H) and that expressed

enterocyte markers such as the serine protease *Jon99Ciii* and other enterocyte marker genes

such as the amylase, *Amy-d* (not shown), but not the *beta-Trypsin (betaTry)* gene (Figure 6I-J).

Detailed characterization of the EC-Gal4 line indicated that although this line labeled a high

percentage of enterocytes, expression was restricted to an area at the anterior and middle of

the midgut, with gaps between these regions and at the posterior (Supplemental Figure 21).

This could explain the absence of subsets of enterocytes, such as those labeled by *betaTry*,

which exhibits regional expression in R2 of the adult midgut (Buchon et al., 2013).

Finally, expression of TaG-EM barcode 9, which was expressed using the EE-Gal4

driver line, was observed in clusters annotated as enteroendocrine cells (Figure 6K-L) and that

also expressed enteroendocrine cell derived neuropeptide genes such as *Dh31* (Figure 6M) and

other enteroendocrine markers such as *IA-2*, a tyrosine phosphatase involved in the secretion of

insulin-like peptide (Figure 6N). Detailed characterization of the EE-Gal4 driver line indicated

that ~80-85% of Prospero-positive enteroendocrine cells are labeled in the anterior and middle

of the larval midgut, with a lower percentage (~65%) of Prospero-positive cells labeled in the

posterior midgut (Supplemental Figure 21). As with the enterocyte labeling, and consistent with

the Gal4 driver expression pattern, the EE-Gal4 expressed TaG-EM barcode 9 did not label all

classes of enteroendocrine cells and other clusters of presumptive enteroendocrine cells

expressing other neuropeptides such as *Orcokinin*, *AstA*, and *AstC*, or neuropeptide receptors

such as *CCHa2* (not shown) were also observed. The EE-Gal4 driver uses *Dh31* regulatory

elements, so it is not surprising that the TaG-EM barcodes specifically labeled *Dh31*-positive

enteroendocrine cells and this result further highlights the ability to target specific genetically

defined cell types using TaG-EM based on *in vivo* cell labeling. Taken together, these results

demonstrate that TaG-EM can be used to label specific cell populations that correlate with Gal4-

driven expression patterns *in vivo* for subsequent identification in single-cell sequencing data.

Discussion

383
Advances in next-generation sequencing, as well as single-cell and spatial genomics are
enabling new types of detailed analyses to study important biological processes such as
development and nervous system function. Here we describe TaG-EM, a genetic barcoding
strategy that enables novel capabilities in several different experimental contexts (Figure 1).
We demonstrate that the genetic barcodes can be quantified from mixtures of barcoded
fly lines using next-generation sequencing. Analysis of structured pools of flies with defined
inputs suggests that TaG-EM barcode measurements are highly accurate and reproducible,
particularly in cases where multiple barcodes are used to label an experimental condition and
averaged (Figure 2). Sequencing-based TaG-EM measurements recapitulated more laborious,
one-at-a-time measurements in a phototaxis assay, an age-dependent fecundity assay, and a
gut motility assay, demonstrating that TaG-EM can be used to measure behavior or other
phenotypes in multiplexed, pooled populations (Figures 3-4). We did note that one line (TaG-EM
barcode 14) exhibited poor performance in oviposition assays, suggesting that barcode
performance should be verified for a specific assay. We excluded this poor performing barcode
line from the fecundity tests, however, backcrossing is often used to bring reagents into a
consistent genetic background for behavioral experiments and could also potentially be used to
address behavior-specific issues with specific TaG-EM lines. In addition, other strategies such
as averaging across multiple barcode lines (3-4 per condition, which yielded R^2 values >0.99 in
tests with structured pools) or permutation of barcode assignment across replicates could also
mitigate such deficiencies. Currently, up to 176 conditions can be multiplexed in a single pooled
experiment with existing TaG-EM lines, but because sequencing indices can be added after
amplification in a separate indexing PCR step, many hundreds or even thousands of such
experiments can be multiplexed in a single sequencing run. While the utility of TaG-EM
barcode-based quantification will vary based on the number of conditions being analyzed and
the ease of quantifying the behavior or phenotype by other means, we demonstrate that TaG-
EM can be employed to cost-effectively streamline labor-intensive assays and to quantify
phenotypes with small effect sizes (Figure 4, Supplemental Figure 8). An additional benefit of
multiplexed TaG-EM behavioral measurements is that the experimental conditions are
effectively blinded as the multiplexed conditions are intermingled in a single assay.
In addition, we show that TaG-EM barcodes can be expressed by tissue-specific Gal4
drivers and used to tag specific cell populations upstream of single-cell sequencing (Figures 5-
6). This capability will allow for positive identification of cell clusters in cell atlas projects and will
facilitate multiplexing of single-cell sequencing experiments. Recently, a conceptually similar

approach called RABID-Seq (Clark et al., 2021) has been described, which allows trans-

synaptic labeling of neural circuits using barcoded viral transcripts. However, one distinction

between the two approaches is that RABID-Seq relies on stochastic viral infection of

mammalian cells while TaG-EM allows reproducible targeting of defined cell populations

allowing unambiguous cell identification and potentially allowing the same cell populations to be

assessed at different timepoints or in the context of different experimental manipulations. One

current limitation is that TaG-EM barcodes are not observed in all cells in single-cell gene

expression data. It is likely that the strength of the Gal4 driver contributes to the labeling density.

However, we also observed variable recovery of TaG-EM barcodes that were all driven by the

same pan-midgut Gal4 driver (Supplemental Figure 20). For single cell RNA-Seq experiments,

the cost savings of multiplexing is roughly the cost of a run divided by the number of

independent lines multiplexed, plus labor savings by also being able to multiplex upstream flow

cytometry, minus loss of unbarcoded cells. Our experiments indicated that for the specific

drivers we tested TaG-EM barcodes are detected in around one quarter of the cells if relying on

endogenous expression in the gene expression library, though this fraction was higher (~37%) if

sequencing an enriched TaG-EM barcode library in parallel (Figure 6, Supplemental Figures 18-

19).

In the future, generation of additional TaG-EM lines will enable higher levels of

multiplexing. In addition, while the original TaG-EM lines were made using a membrane-

localized myr::GFP construct, variants that express GFP in other cell compartments such as the

cytoplasm or nucleus could be constructed to enable increased expression levels or purification

of nuclei. Nuclear labeling could also be achieved by co-expressing a nuclear GFP construct

with existing TaG-EM lines in analogy to the use of hexameric GFP described above.

In summary, combined with the large collections of Gal4 and split-Gal4 lines that have

been established in Drosophila that enable precise targeting of a high proportion of cell types

(Ariyapala et al., 2020; Aso et al., 2014; Davis et al., 2020; Gohl et al., 2011; Kanca et al., 2022;

Namiki et al., 2018; Pfeiffer et al., 2010, 2008; Venken et al., 2011; Zirin et al., 2024), TaG-EM

provides a means to target and label cells *in vivo* for subsequent detection in single-cell

sequencing. Moreover, these genetic barcodes can be used to multiplex behavioral or other

phenotypic measurements. Thus, TaG-EM provides a flexible system for barcoding cells and

organisms.

Methods

Drosophila stocks and maintenance

Drosophila stocks were grown at 22ºC on cornmeal agar unless otherwise indicated. The

following stocks were used in this study:

Stock name	Source
IsoD1	Clandinin Lab, Stanford University
w-; +; + (IsoD1)	Clandinin Lab, Stanford University
attP2 line	BDSC #25710
norpA	BDSC# 9048
UAS-myr::GFP (pJFRC12)	BDSC #32197
Hexameric GFP lines	BDSC #91402, #91403
UAS-6XmCherry-HA	BDSC #52268
UAS-GFP.nls	BDSC #4776
esg-GFP.FPTB	BDSC #83386
dpp-Gal4 driver	BDSC #1553
Act-Gal4 driver	BDSC #4414
Tub-Gal4 driver	BDSC #5138
Mhc-Gal4 driver	BDSC #55132
PC-Gal4 driver	BDSC #73356 + #75528
PC-Gal4 driver (with UAS-Stinger)	BDSC #91400 + #91401
EC-Gal4 driver	BDSC #91406
EB-Gal4 driver	BDSC #91398 + #91404
EE-Gal4 driver	BDSC #91402 + #91403
PMG-Gal4 driver	BDSC #91398 + #91399
TaG-EM lines (x176)	This study, (Alegria et al., 2024)
TaG-EM lines + 6xGFP (x20)	This study

Design and cloning of TaG-EM constructs

A gBlock with the following sequence containing a part of the SV40 3' UTR with a PCR handle

(uppercase, below) and a 14 bp randomer sequence just upstream of the SV40 polyadenylation

site (bold and underlined, below) was synthesized (Integrated DNA Technologies, IDT):

caaaggaaaaagctgcactgctatacaagaaaattatggaaaaatatttgatgtatagtgccttgactagagatcataatcagccata

ccacatttgtagaggttttacttgctttaaaaaacctcccacacctccccctgaacctgaaacataaaatgaatgcaattgttgttgttaact

tgtttattgcagcttataaCTTCCAACAACCGGAAGTGANNNNNNNNNNNNNNNtggttaca<u>aataaag</u>caatag

catcacaaatttcaca<u>aataaag</u>cattttttttcactg**ca**ttctagttgtggtttgtccaaactcatcaatgtatcttatcatgtctggatcgatct

ggccggccgtttaaacgaattcttgaagacgaaagggcctcgtgatacgcctattttataggttaatgtcatgataataatg

The gBlock was resuspended in 20 µl EB, incubated at 50ºC for 20 minutes and then cut with

*Psi*I and *EcoR*I (New England Biolabs, NEB) using the following reaction conditions: 4 µl gBlock

DNA (35 ng), 2 µl 10x CustSmart buffer (NEB), 1 µl *Eco*RI enzyme (NEB), 1 µl *Psi*I enzyme

(NEB), and 12 µl nuclease-free water were mixed and incubated at 37ºC for 1 hour followed by

65ºC for 20 minutes to heat inactivate the restriction enzymes. pJFRC12-10XUAS-IVS-

myr::GFP plasmid (Addgene, Plasmid #26222) (Pfeiffer et al., 2010) was digested with the

following reaction conditions: 5 µl pJFRC12-10XUAS-IVS-myr::GFP plasmid DNA (~3 µg), 5 µl

10x CutSmart buffer (NEB), 1 µl *Psi*I enzyme (NEB), 1 µl *Eco*RI enzyme (NEB), and 38 µl

nuclease-free water, were mixed and incubated at 37ºC for 1 hour, followed by addition of 1 µl

of CIP and incubation for an additional 30 minutes. The digested vector backbone was gel

purified using the QiaQuick Gel Purification Kit (Qiagen). The digested gBlock was ligated into

the digested pJFRC12-10XUAS-IVS-myr::GFP backbone using the following reactions

conditions: 4 µl T4 ligase buffer (10x) (NEB), 20 µl plasmid backbone DNA (0.005 pmol), 5 µl

gBlock digest DNA (0.03 pmol), 2 µl of T4 DNA ligase (NEB), and 9 µl nuclease-free water were

mixed and incubated at 22ºC for 2 hours. 2 µl of the ligation reaction was transformed into 50 µl

of TOP10 competent cells (Invitrogen), and the cells were incubated on ice for 30 minutes, then

heat shocked at 42ºC for 30 seconds, and incubated on ice for 5 minutes. 250 µl SOC was

added and the cells were plated on LB+ampicillin plates and incubated overnight at 37ºC. DNA

was isolated from 36 pJFRC12-gBlock colonies using a QIAprep Spin MiniPrep kit (Qiagen).

Expected construct size was verified by diagnostic digest with *Eco*RI and *Apa*LI. DNA

concentration was determined using a Quant-iT PicoGreen dsDNA assay (Thermo Fisher

Scientific) and the randomer barcode for each of the constructs was determined by Sanger

sequencing using the following primers:

SV40_post_R: GCCAGATCGATCCAGACATGA

SV40_5F: CTCCCCCTGAACCTGAAACA

Generation of TaG-EM transgenic lines

29 sequence verified constructs were normalized, pooled evenly, and injected as a pool into

embryos (Rubin and Spradling, 1982) expressing PhiC31 integrase and the carrying the *attP2*

landing site (BDSC #25710). Injected flies were outcrossed to w- flies, and up to three *white+*

progeny per cross were identified, and the transgenic lines were homozygosed. DNA was

extracted (GeneJET genomic DNA purification Kit, Thermo Scientific) and the region containing

the DNA barcode was amplified with the following PCR reaction: 2.5 µl 1:10 diluted template

DNA, 2 µl 10x Reaction Buffer (Qiagen), 0.2 µl dNTP mix (10 µM), 1 µl 10 µM SV40_5F primer

(10 µM), 1 µl SV40_post_R primer (10 µM), 0.8 µl $MgCl_2$ (3 mM), 0.1 µl Taq polymerase

(Qiagen), 12.4 µl nuclease-free water. Reactions were amplified using the following cycling

conditions: 95ºC for 5 minutes, followed by 30 cycles of 94ºC for 30 seconds, 55ºC for 30

seconds, 72ºC for 30 seconds, followed by 72ºC for 5 minutes. PCR products were treated with

Exo-CIP using the following reaction conditions: 5 µl PCR product, 1 µl Exo-CIP Tube A (NEB),

1 µl Exo-CIP Tube B (NEB) were mixed and incubated at 37ºC for 4 minutes, followed by 80ºC

for 1 minute. The barcode sequence for each of the independent transgenic lines was

determined by Sanger sequencing using the SV40_5F and SV40_PostR primers. Transgenic

lines containing 20 distinct DNA barcodes were recovered (Supplemental Figure 1). An

additional 156 TaG-EM barcode lines were isolated and sequence verified as described in a

separate publication (Alegria et al., 2024).

Optimizing amplification of TaG-EM barcodes for next-generation sequencing

The following primers were evaluated to amplify the TaG-EM barcodes upstream of NGS:

Forward primer pool; four primers with frameshifting bases to increase library sequence

diversity in initial sequencing cycles were normalized to 10 µM and pooled evenly to make a

B2_3'F1_Nextera_0-6 primer pool:

B2_3'F1_Nextera:

TCGTCGGCAGCGTCAGATGTGTATAAGAGACAGCTTCCAACAACCGGAAG*TGA

B2_3'F1_Nextera_2

TCGTCGGCAGCGTCAGATGTGTATAAGAGACAGAGCTTCCAACAACCGGAAG*TGA

B2_3'F1_Nextera_4

TCGTCGGCAGCGTCAGATGTGTATAAGAGACAGTCGACTTCCAACAACCGGAAG*TGA

B2_3'F1_Nextera_6

TCGTCGGCAGCGTCAGATGTGTATAAGAGACAGGAAGAGCTTCCAACAACCGGAAG*TGA

The following two reverse primers were tested:

SV40_pre_R_Nextera:

GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAGATTTGTGAAATTTGTGATGCTATTGC*T

TT

SV40_post_R_Nextera:

GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAGGCCAGATCGATCCAGACA*TGA

SV40_pre_R_nextera is designed to produce a shorter amplicon (200 bp with Illumina adapters

and indices added) and SV40-post_R_Nextera is designed to produce a longer amplicon (290

bp with Illumina adapters and indices added).

An initial test was performed with three different polymerases (Qiagen Q5, KAPA HiFi, and

Qiagen Taq) at two different annealing temperatures and with both the B2_3'F1_Nextera/

SV40_pre_R_Nextera and B2_3'F1_Nextera/ SV40_post_R_Nextera primer sets to determine

whether the primers amplify as expected (Supplemental Figure 1). Two different samples were

tested:

1. Pool of 8 putative transformant samples (pooled 5 ul each of 1:10 diluted sample)

2. OreR (wild type - diluted 1:10)

Set up the following PCR reactions:

Q5 polymerase: 2.5 µl template DNA, 1 µl 10 µM Forward primer (10 µM), 1 µl Reverse primer

(10 µM), 10 µl 2x Q5 Master Mix (Qiagen), 5.5 µl nuclease-free water. Reactions were amplified

using the following cycling conditions: 98ºC for 30 seconds, followed by 30 cycles of 98ºC for 20

seconds, 55ºC or 60ºC for 15 seconds, 72ºC for 30 seconds, followed by 72ºC for 5 minutes.

KAPA HiFi polymerase: 2.5 µl template DNA, 1 µl 10 µM Forward primer (10 µM), 1 µl Reverse

primer (10 µM), 10 µl 2x KAPA HiFi ReadyMix (Roche), 5.5 µl nuclease-free water. Reactions

were amplified using the following cycling conditions: 95ºC for 5 minutes, followed by 30 cycles

of 98ºC for 20 seconds, 55ºC or 60ºC for 15 seconds, 72ºC for 30 seconds, followed by 72ºC for

5 minutes.

Taq polymerase: 2.5 µl template DNA, 2 µl 10x Reaction Buffer (Qiagen), 0.2 µl dNTP mix (10

µM), 1 µl 10 µM Forward primer (10 µM), 1 µl Reverse primer (10 µM), 0.8 µl $MgCl_2$ (3 mM), 0.1

µl Taq polymerase (Qiagen), 12.4 µl nuclease-free water. Reactions were amplified using the

following cycling conditions: 95ºC for 5 minutes, followed by 30 cycles of 94ºC for 30 seconds,

55ºC or 60ºC for 30 seconds, 72ºC for 30 seconds, followed by 72ºC for 5 minutes.

Samples were run on a 2% agarose gel to verify amplification products (Supplemental Figure 1).

Next, the TaG-EM barcode lines were pooled in either an even or staggered manner. To

optimize reaction conditions for the barcode measurements, either 5 ng or 50 ng of DNA was

amplified in triplicate for each pool for either 20, 25, or 30 cycles with either KAPA HiFi using the

B2_Nextera_F 0-6 forward primer pool together with either the SV40_pre_R_Nextera or the

SV40_post_R_Nextera reverse primer. Next, PCR reactions were diluted 1:100 in nuclease-free

water and amplified in the following indexing reactions: 3 µl PCR 1 (1:100 dilution), 1 µl indexing

primer 1 (5 µM), 1 µl indexing primer 2 (5 µM), and 5 µl 2x Q5 master mix. The following

indexing primers were used (X indicates the positions of the 8 bp indices):

Forward indexing primer:

AATGATACGGCGACCACCGAGATCTACACXXXXXXXXXTCGTCGGCAGCGTC

Reverse indexing primer:

CAAGCAGAAGACGGCATACGAGATXXXXXXXXGTCTCGTGGGCTCGG

Reactions were amplified using the following cycling conditions: 98°C for 30 seconds, followed

by 10 cycles of 98°C for 20 seconds, 55°C for 15 seconds, 72°C for 1 minute, followed by 72°C

for 5 minutes. Amplicons were then purified and normalized using a SequalPrep normalization

plate (Thermo Fisher Scientific), followed by elution in 20 µl of elution buffer. An even volume of

the normalized libraries was pooled and concentrated using 1.8x AmpureXP beads (Beckman

Coulter). Pooled libraries were quantified using a Qubit dsDNA high sensitivity assay (Thermo

Fisher Scientific) and libraries were normalized to 2 nM for sequencing on the Illumina MiSeq

(see below).

Structured fly pool experiments

Male or female flies from TaG-EM barcode lines were pooled in either an even or staggered

manner (Figure 2A). For the even pools, three independently pooled samples were constructed

in order to assess sample-to-sample variability. DNA was extracted from these structured pools

using a protocol adapted from Huang *et al*., 2009 (Huang et al., 2009), using homemade SPRI

beads (DeAngelis et al., 1995) in the last purification step and amplified in triplicate using 2.5 µl

template DNA (50 ng), 1 µl 10 µM B2_Nextera_F 0-6 primers (10 µM), 1 µl

SV40_pre_R_Nextera (10 µM), 10 µl 2x KAPA HiFi ReadyMix (Roche), 5.5 µl nuclease-free

water. Reactions were amplified using the following cycling conditions: 98ºC for 5 minutes,

followed by 30 cycles of 98ºC for 20 seconds, 60ºC for 15 seconds, 72ºC for 30 seconds,

followed by 72ºC for 5 minutes. Amplicons were indexed, normalized, quantified, and prepared

for sequencing as described above.

Phototaxis experiments

<u>Video-based measurements</u>

A pair of white LED strip lights with Muzata LED Strip Light Diffusers (U1SW WW 1M, LU1)

were mounted withing a light-tight box and controlled using an Vilros Uno Rev 3 microcontroller.

Test tubes containing flies were held in place with Acoustic Foam Panels (1"x12"x12",

ALPOWL). Videos and images were acquired using an Arducam 1080P Day & Night Vision

USB Camera with an IR filter and using Photo Booth software (Apple). Wild type and *norpA* flies

carrying one of 4 different TaG-EM barcodes were tested in three independent experimental

replicates. 20 male flies of each genotype were transferred into 25 mm x 150 mm glass test

tubes, incubated at 34ºC for 10 minutes and then run in the phototaxis assay, where a light at

one end of the chamber was turned on for 30 seconds. Videos of all tests were recorded

through the end of the 30 second light pulse. Videos were independently scored by two

observers to determine the number of flies in the light-facing or dark-facing tubes and the results

were averaged. A preference index (P.I.) was calculated using the following formula: [(number

of flies in light tube) - (number of flies in dark tube)]/(total number of flies).

<u>TaG-EM measurements</u>

For TaG-EM barcode-based phototaxis measurements, the following genotypes were

consolidated into a single test tube:

<u>Pool A</u>

norpA/Y;TaG-EM BC4/+

norpA/Y;TaG-EM BC3/+

w-/Y;TaG-EM BC2/+

w-/Y;TaG-EM BC1/+

<u>Pool B</u>

norpA/Y;TaG-EM BC2/+

norpA/Y;TaG-EM BC1/+

w-/Y;TaG-EM BC4/+

w-/Y;TaG-EM BC3/+

These pools were individually incubated at 34ºC for 10 minutes and then run in the phototaxis

assay. Videos of all tests were recorded and at the end of a 30 second light pulse the two test

tubes were quickly separated and capped. Flies in each of these tubes were counted, then DNA

was extracted from the flies from the light-facing or dark-facing tubes and amplified using 2.5 µl

template DNA (50 ng), 1 µl 10 µM B2_Nextera_F 0-6 primers (10 µM), 1 µl

SV40_pre_R_Nextera (10 µM), 10 µl 2x KAPA HiFi ReadyMix (Roche), 5.5 µl nuclease-free

water. Reactions were amplified using the following cycling conditions: 95ºC for 5 minutes,

followed by 30 cycles of 98ºC for 20 seconds, 60ºC for 15 seconds, 72ºC for 30 seconds,

followed by 72ºC for 5 minutes. Amplicons were indexed, normalized, quantified, and prepared

for sequencing as described above.

Oviposition experiments

Newly hatched flies (males and females) from three barcode lines were collected at one-week

intervals during 4 consecutive weeks (12 barcode lines in total). Fresh fly food was provided

every 3-4 days. Ten days after the last collection, 10 females from each barcode line were taken

and pooled together in a collection cage (10 females x 12 barcode lines= 120 females). The

remaining females from each barcode line were separated from the males and put in individual

collection cages. Two days later, the experiment started and run for 3 consecutive days. Each

day a 1-1.5 hour pre-collection was followed by a 6 hour collection, both at 25ºC. 100 embryos

from each individual collection plate were transferred to new plates and incubated for 2 days at

18ºC. The number of hatched larvae were counted and used to calculate the egg survival rate.

The pooled collection plate was also incubated at 18ºC and the next day the embryos were

dechorionated and frozen. The 12 individual collection plates were kept at 4ºC and the number

of embryos counted in the following days. For the barcode measurements, DNA was extracted

from the embryos, and amplified using 2.5 µl template DNA (50 ng), 1 µl 10 µM B2_Nextera_F

0-6 primers (10 µM), 1 µl SV40_pre_R_Nextera (10 µM), 10 µl 2x KAPA HiFi ReadyMix

(Roche), 5.5 µl nuclease-free water. Reactions were amplified using the following cycling

conditions: 95ºC for 5 minutes, followed by 30 cycles of 98ºC for 20 seconds, 60ºC for 15

seconds, 72ºC for 30 seconds, followed by 72ºC for 5 minutes. Amplicons were indexed,

normalized, quantified, and prepared for sequencing as described above.

Larval gut motility experiments

<u>Preparing Yeast Food Plates</u>

Yeast agar plates were prepared by making a solution containing 20% Red Star Active Dry

Yeast 32oz (Red Star Yeast) and 2.4% Agar Powder/Flakes (Fisher) and a separate solution

containing 20% Glucose (Sigma-Aldrich). Both mixtures were autoclaved with a 45-minute liquid

cycle and then transferred to a water bath at 55ºC. After cooling to 55ºC, the solutions were

combined and mixed, and approximately 5 mL of the combined solution was transferred into

100 x 15 mm petri dishes (VWR) in a PCR hood or contamination-free area. For blue-dyed

yeast food plates, 0.4% Blue Food Color (McCormick) was added to the yeast solution. For the

caffeine assays, 300 µL of a solution of 100 mM 99% pure caffeine (Sigma-Aldrich) was

pipetted onto the blue-dyed yeast plate and allowed to absorb into the food during the 90-minute

starvation period.

<u>Manual Gut Motility Assay</u>

Third instar Drosophila larvae were transferred to empty conical tubes that had been misted with

water to prevent the larvae from drying out. After a 90-minute starvation period the larvae were

moved from the conical to a blue-dyed yeast plate with or without caffeine and allowed to feed

for 60 minutes. Following the feeding period, the larvae were transferred to an undyed yeast

plate. Larvae were scored for the presence or absence of a food bolus every 30 minutes over a

5-hour period. Up to 8 experimental replicates/conditions were scored simultaneously.

<u>TaG-EM Gut Motility Assay</u>

Third instar larvae were starved and fed blue dye-containing food with or without caffeine as

described above. An equal number of larvae from each experimental condition/replicate were

transferred to an undyed yeast plate. During the 5-hour observation period, larvae were

examined every 30 minutes and larvae lacking a food bolus were transferred to a

microcentrifuge tube labeled for the timepoint. Any larvae that died during the experiment were

placed in a separate microcentrifuge tube and any larvae that failed to pass the food bolus were

transferred to a microcentrifuge tube at the end of the experiment. DNA was extracted from the

larvae in each tube and TaG-EM barcode libraries were prepared and sequenced as described

above.

Dissection and immunostaining

Midguts from third instar larvae of driver lines crossed to UAS-GFP.nls or UAS-mCherry were

dissected in 1xPBS and fixed with 4% paraformaldehyde (PFA) overnight at 4ºC. Fixed samples

were washed with 0.1% PBTx (1xPBS + 0.1% Triton X-100) three times for 10 minutes each

and blocked in PBTxGS (0.1% PBTx + 3% Normal Goat Serum) for 2–4 hours at RT. After

blocking, midguts were incubated in primary antibody solution overnight at 4ºC. The next day

samples were washed with 0.1% PBTx three times for 20 minutes each and were incubated in

secondary antibody solution for 2–3 hours at RT (protected from light) followed by three washes

with 0.1% PBTx for 20 minutes each. One µg/ml DAPI solution prepared in 0.1% PBTx was added to the sample and incubated for 10 minutes followed by washing with 0.1% PBTx three times for 10 minutes each. Finally, samples were mounted on a slide glass with 70% glycerol and imaged using a Nikon AX R confocal microscope. Confocal images were processed using Fiji software.

The primary antibodies used were rabbit anti-GFP (A6455,1:1000 Invitrogen), mouse anti-mCherry (3A11, 1:20 DSHB), mouse anti-Prospero (MR1A, 1:50 DSHB) and mouse anti-Pdm1 (Nub 2D4, 1:30 DSHB). The secondary antibodies used were goat anti-mouse and goat anti-rabbit IgG conjugated to Alexa 647 and Alexa 488 (1:200) (Invitrogen), respectively. Five larval gut specimens per Gal4 line were dissected and examined.

Cell dissociation and isolation

Midguts from 3rd instar larvae were dissected in phosphate-buffered saline (PBS) and transferred to microcentrifuge tubes on ice containing PBS + 30% normal goat serum (NGS). After dissection, 150 µL of 2.7 mg/mL elastase was added to each sample tube. The tubes were then incubated at 27°C for one hour. During incubation, samples were mixed by pipetting ~30 times every 15 minutes to improve elastase dissociation of the cells. Samples were then filtered through a 40 µM FlowMi tip filter (Bel-Art) to reduce debris. Afterwards, the samples were quantified on the LUNA-FL Dual Fluorescence Cell Counter (Logos Biosystems) using 9 µL of sample to 1µL AO/PI dye to ensure there were enough viable cells for flow sorting.

Once quantified, the samples were brought up to a volume of ~1.1mL with the PBS + 30% NGS solution to facilitate flow sorting. The samples were then fluorescently sorted on a FACSAria II Cell Sorter (BD Biosciences) to isolate GFP+ cells. Following sorting, samples were centrifuged at 300x g for 10 minutes to concentrate the cells. The supernatant was aspirated off until 50 µL cell concentrate remained in each sample. Then, the samples were carefully resuspended using wide bore pipette tips before being combined into one sample tube. This sample was quantified on the LUNA-FL Dual Fluorescence Cell Counter (Logos Biosystems) as described above. If necessary, cells were centrifuged, concentrated, and re-counted.

Preparation of single-cell sequencing libraries

The resulting pool was prepared for sequencing following the 10x Genomics Single Cell 3' protocol (version CG000315 Rev C), At step 2.2 of the protocol, cDNA amplification, 1 µl of

TaG-EM spike-in primer (10 μM) was added to the reaction to amplify cDNA with the TaG-EM

barcode. Gene expression cDNA and TaG-EM cDNA were separated using a double-sided

SPRIselect (Beckman Coulter) bead clean up following 10x Genomics Single Cell 3' Feature

Barcode protocol, step 2.3 (version CG000317 Rev E). The gene expression cDNA was created

into a library following the CG000315 Rev C protocol starting at section 3. Custom nested

primers were used for enrichment of TaG-EM barcodes after cDNA creation using PCR.

The following primers were tested (see Supplemental Figure 18):

UMGC_IL_TaGEM_SpikeIn_v1:

GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTCTTCCAACAACCGGAAGT*G*A

UMGC_IL_TaGEM_SpikeIn_v2:

GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTGCAGCTTATAACTTCCAACAACCGGAA

GT*G*A

UMGC_IL_TaGEM_SpikeIn_v3:

TGTGCTCTTCCGATCTGCAGCTTATAACTTCCAACAACCGGAAGT*G*A

D701_TaGEM:

CAAGCAGAAGACGGCATACGAGATCGAGTAATGTGACTGGAGTTCAGACGTGTGCTCTTC

CGATCTGCAGC*T*T

SI PCR Primer: AATGATACGGCGACCACCGAGATCTACACTCTTTCCCTACACGACGC*T*C

UMGC_IL_DoubleNest:

GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTGCAGCTTATAACTTCCAACAACCGG*A*

A

P5: AATGATACGGCGACCACCGA

D701:

GATCGGAAGAGCACACGTCTGAACTCCAGTCACATTACTCGATCTCGTATGCCGTCTTCTG

CTTG

D702:

GATCGGAAGAGCACACGTCTGAACTCCAGTCACTCCGGAGAATCTCGTATGCCGTCTTCT

GCTTG

After multiple optimization trials, the following steps yielded ~96% on-target reads for the TaG-

EM library (Supplemental Figure 18, note that for the enriched barcode data shown in Figure 6

and Supplemental Figure 19, a similar amplification protocol was used TaG-EM barcodes were

amplified from the gene expression library cDNA and not the SPRI-selected barcode pool).

TaG-EM cDNA was amplified with the following PCR reaction: 5 μl purified TaG-EM cDNA, 50

μl 2x KAPA HiFi ReadyMix (Roche), 2.5 μl UMGC_IL_DoubleNest primer (10 μM), 2.5 μl

SI_PCR primer (10 µM), and 40 µl nuclease-free water. The reaction was amplified using the

following cycling conditions: 98ºC for 2 minutes, followed by 15 cycles of 98ºC for 20 seconds,

63ºC for 30 seconds, 72ºC for 20 seconds, followed by 72ºC for 5 minutes. After the first PCR,

the amplified cDNA was purified with a 1.2x SPRIselect (Beckman Coulter) bead cleanup with

80% ethanol washes and eluted into 40 µL of nuclease-water. A second round of PCR was run

with following reaction: 5 µl purified TaG-EM cDNA, 50 µl 2x KAPA HiFi ReadyMix (Roche), 2.5

µl D702 primer (10 µM), 2.5 µl p5 Primer (10 µM), and 40 µl nuclease-free water. The reaction

was amplified using the following cycling conditions: 98ºC for 2 minutes, followed by 10 cycles

of 98ºC for 20 seconds, 63ºC for 30 seconds, 72ºC for 20 seconds, followed by 72ºC for 5

minutes. After the second PCR, the amplified cDNA was purified with a 1.2x SPRIselect

(Beckman Coulter) bead cleanup with 80% ethanol washes and eluted into 40uL of nuclease-

water. The resulting 3' gene expression library and TaG-EM enrichment library were sequenced

together following Scenario 1 of the BioLegend "Total-Seq-A Antibodies and Cell Hashing with

10x Single Cell 3' Reagents Kit v3 or v3.1" protocol. Additional sequencing of the enriched TaG-

EM library also done following Scenario 2 from the same protocol.

Sequencing

Libraries for TaG-EM barcode analysis from structured pools or from phototaxis or oviposition

experiments were denatured with NaOH and prepared for sequencing according to the

protocols described in the Illumina MiSeq Denature and Dilute Libraries Guides. Single-cell

libraries were sequenced on the Illumina NextSeq 2000 or Illumina NovaSeq 6000. One of the

single cell enriched TaG-EM barcode libraries was sequenced on an Element Aviti sequencer

following the manufacturers loading instructions.

Data analysis

Behavioral experiments: Demultiplexed fastq files were generated using bcl2fastq or bcl-

convert. TaG-EM barcode data was analyzed using custom R and Python scripts and BioPython

(Cock et al., 2009). Leading primer sequences were trimmed using cutadapt (Martin, 2011) and

the first 14 bp of the remaining trimmed read were compared to a barcode reference file, with a

maximum of 2 mismatches allowed, using a custom script (TaG-EM_barcode_analysis.py)

which is available via Github: https://github.com/darylgohl/TaG-EM.

Single-cell experiments:

Datasets were first mapped and analyzed using the Cell Ranger analysis pipeline (10x

Genomics). A custom Drosophila genome reference was made by combining the BDGP.28

reference genome assembly and Ensembl gene annotations. Custom gene definitions for each

of the TaG-EM barcodes were added to the fasta genome file and .gtf gene annotation file. A

Cell Ranger reference package was generated with the Cell Ranger *mkref*

command. Subsequent single-cell data analysis was performed using the R package Seurat

(Satija et al., 2015). Cells expressing less than 200 genes and genes expressed in fewer than

three cells were filtered from the expression matrix. Next, percent mitochondrial reads, percent

ribosomal reads cells counts, and cell features were graphed to determine optimal filtering

parameters. DecontX (Yang et al., 2020) was used to identify empty droplets, to evaluate

ambient RNA contamination, and to remove empty cells and cells with high ambient RNA

expression. DoubletFinder (McGinnis et al., 2019) to identify droplet multiplets and remove cells

classified as multiplets. Clustree (Zappia and Oshlack, 2018) was used to visualize different

clustering resolutions and to determine the optimal clustering resolution for downstream

analysis. Finally, SingleR (Aran et al., 2019) was used for automated cell annotation with a gut

single-cell reference from the Fly Cell Atlas (Li et al., 2022). The dataset was manually

annotated using the expression patterns of marker genes known to be associated with cell types

of interest. To correlate TaG-EM barcodes with cell IDs in the enriched TaG-EM barcode library,

a custom Python script was used (TaG-EM_barcode_Cell_barcode_correlation.py), which is

available via Github: https://github.com/darylgohl/TaG-EM.

Declarations

Availability of data, code, and materials

Sequencing data for this project is available through the National Center for Biotechnology Information (NCBI) Sequence Read Archive BioProject PRJNA912199. Fly stocks containing 20 of the TaG-EM barcodes together with an additional UAS hexameric GFP expression construct will be available from the Bloomington Drosophila Stock Center. Additional TaG-EM barcode stocks are available upon request. Single cell analysis code and the TaG-EM barcode analysis script and barcode reference fasta files are available via Github: https://github.com/darylgohl/TaG-EM.

Competing interests

None to declare.

Funding

This study was supported by a grant from the Winston and Maxine Wallin Neuroscience Discovery Fund.

Author contributions

J.B.M. conceived and designed experiments, conducted experiments, analyzed data, revised manuscript, M.D. conducted experiments, analyzed data, revised manuscript, L.G. conducted experiments, revised manuscript, B.A. conducted experiments, revised manuscript, J.G. analyzed data, revised manuscript, D.M.G. conceived and designed experiments, conducted experiments, analyzed data, wrote and revised the manuscript.

Acknowledgements

We thank our colleagues in the University of Minnesota Genomics Center (RRID:SCR_012413), in particular Aaron Becker, Dylan Cole, and Logan Silber for help with DNA sequencing, Emma Stanley, Fernanda Rodriguez, and Patrick Grady for assistance and advice on single-cell sequencing, and Thomas Clandinin, Nam Chul Kim, Kenneth Beckman, Andrew Alegria, Troy Louwagie, and Aaron Barnes for helpful feedback and discussions. This work was supported by the resources and staff at the University of Minnesota University Imaging Centers (RRID:SCR_020997). The authors acknowledge the Minnesota Supercomputing Institute (MSI) at the University of Minnesota for providing resources that contributed to the research results

873   reported within this paper. URL: http://www.msi.umn.edu. Stocks obtained from the Bloomington

874   Drosophila Stock Center (NIH P40OD018537) were used in this study. Antibodies were

875   obtained from the Developmental Studies Hybridoma Bank (DSHB). Prospero (MR1A) was

876   deposited to the DSHB by C.Q. Doe and Nub 2D4 was deposited to the DSHB by Michalis

877   Averof.

References

Alegria AD, Joshi AS, Mendana JB, Khosla K, Smith KT, Auch B, Donovan M, Bischof J, Gohl DM, Kodandaramaiah SB. 2024. High-throughput genetic manipulation of multicellular organisms using a machine-vision guided embryonic microinjection robot. *Genetics* **226**:iyae025. doi:10.1093/genetics/iyae025

Aran D, Looney AP, Liu L, Wu E, Fong V, Hsu A, Chak S, Naikawadi RP, Wolters PJ, Abate AR, Butte AJ, Bhattacharya M. 2019. Reference-based analysis of lung single-cell sequencing reveals a transitional profibrotic macrophage. *Nat Immunol* **20**:163–172. doi:10.1038/s41590-018-0276-y

Ariyapala IS, Holsopple JM, Popodi EM, Hartwick DG, Kahsai L, Cook KR, Sokol NS. 2020. Identification of split-GAL4 drivers and enhancers that allow regional cell type manipulations of the drosophila melanogaster intestine. *Genetics* **216**:891–903. doi:10.1534/genetics.120.303625

Asnicar F, Leeming ER, Dimidi E, Mazidi M, Franks PW, Al Khatib H, Valdes AM, Davies R, Bakker E, Francis L, Chan A, Gibson R, Hadjigeorgiou G, Wolf J, Spector TD, Segata N, Berry SE. 2021. Blue poo: impact of gut transit time on the gut microbiome using a novel marker. *Gut* **70**:1665–1674. doi:10.1136/gutjnl-2020-323877

Aso Y, Hattori D, Yu Y, Johnston RM, Iyer NA, Ngo TTB, Dionne H, Abbott LF, Axel R, Tanimoto H, Rubin GM. 2014. The neuronal architecture of the mushroom body provides a logic for associative learning. *eLife* **3**:e04577. doi:10.7554/ELIFE.04577

Bhang HC, Ruddy DA, Krishnamurthy Radhakrishna V, Caushi JX, Zhao R, Hims MM, Singh AP, Kao I, Rakiec D, Shaw P, Balak M, Raza A, Ackley E, Keen N, Schlabach MR, Palmer M, Leary RJ, Chiang DY, Sellers WR, Michor F, Cooke VG, Korn JM, Stegmeier F. 2015. Studying clonal dynamics in response to cancer therapy using high-complexity barcoding. *Nature Medicine* **21**:440–448. doi:10.1038/nm.3841

Buchon N, Osman D, David FPA, Yu Fang H, Boquete J-P, Deplancke B, Lemaitre B. 2013. Morphological and Molecular Characterization of Adult Midgut Compartmentalization in *Drosophila*. *Cell Reports* **3**:1725–1738. doi:10.1016/j.celrep.2013.04.001

Chen X, Sun Y-C, Zhan H, Kebschull JM, Fischer S, Matho K, Huang ZJ, Gillis J, Zador AM. 2019. High-Throughput Mapping of Long-Range Neuronal Projection Using In Situ Sequencing. *Cell* **179**:772-786.e19. doi:10.1016/j.cell.2019.09.023

Cheng J, Liao J, Shao X, Lu X, Fan X. 2021. Multiplexing Methods for Simultaneous Large-Scale Transcriptomic Profiling of Samples at Single-Cell Resolution. *Advanced Science* **8**:2101229. doi:10.1002/advs.202101229

Clark IC, Gutiérrez-Vázquez C, Wheeler MA, Li Z, Rothhammer V, Linnerbauer M, Sanmarco LM, Guo L, Blain M, Zandee SEJ, Chao CC, Batterman KV, Schwabenland M, Lotfy P, Tejeda-Velarde A, Hewson P, Polonio CM, Shultis MW, Salem Y, Tjon EC, Fonseca-Castro PH, Borucki DM, de Lima KA, Plasencia A, Abate AR, Rosene DL, Hodgetts KJ, Prinz M, Antel JP, Prat A, Quintana FJ. 2021. Barcoded viral tracing of single-cell interactions in central nervous system inflammation. *Science* **372**.

Cock PJA, Antao T, Chang JT, Chapman BA, Cox CJ, Dalke A, Friedberg I, Hamelryck T, Kauff F, Wilczynski B, de Hoon MJL. 2009. Biopython: freely available Python tools for computational molecular biology and bioinformatics. *Bioinformatics (Oxford, England)* **25**:1422–3. doi:10.1093/bioinformatics/btp163

David J, Cohet Y, Fouillet P. 1975. The variability between individuals as a measure of senescence: A study of the number of eggs laid and the percentage of hatched eggs in the case of Drosophila melanogaster. *Experimental Gerontology* **10**:17–25. doi:10.1016/0531-5565(75)90011-X

Davis FP, Nern A, Picard S, Reiser MB, Rubin GM, Eddy SR, Henry GL. 2020. A genetic, genomic, and computational resource for exploring neural circuit function. *eLife* **9**.

DeAngelis MM, Wang DG, Hawkins TL. 1995. Solid-phase reversible immobilization for the
isolation of PCR products. *Nucleic Acids Res* **23**:4742–4743.
Gohl DM, Silies MA, Gao XJ, Bhalerao S, Luongo FJ, Lin C-C, Potter CJ, Clandinin TR. 2011. A
versatile in vivo system for directed dissection of gene expression patterns. *Nature*
Methods **8**:231–237. doi:10.1038/nmeth.1561
Gohl DM, Vangay P, Garbe J, MacLean A, Hauge A, Becker A, Gould TJ, Clayton JB, Johnson
TJ, Hunter R, Knights D, Beckman KB. 2016. Systematic improvement of amplicon
marker gene methods for increased accuracy in microbiome studies. *Nature*
Biotechnology **34**:942–949. doi:10.1038/nbt.3601
Groth AC, Fish M, Nusse R, Calos MP. 2004. Construction of transgenic Drosophila by using
the site-specific integrase from phage phiC31. *Genetics* **166**:1775–82.
doi:10.1534/genetics.166.4.1775
Hristova K, Wimley WC. 2023. Determining the statistical significance of the difference between
arbitrary curves: A spreadsheet method. *PLoS One* **18**:e0289619.
doi:10.1371/journal.pone.0289619
Huang AM, Rehm EJ, Rubin GM. 2009. Quick preparation of genomic DNA from Drosophila.
Cold Spring Harb Protoc **2009**:pdb.prot5198. doi:10.1101/pdb.prot5198
Hung RJ, Hu Y, Kirchner R, Liu Y, Xu C, Comjean A, Tattikota SG, Li F, Song W, Sui SH,
Perrimon N. 2020. A cell atlas of the adult Drosophila midgut. *Proceedings of the*
National Academy of Sciences of the United States of America **117**:1514–1523.
doi:10.1073/pnas.1916820117
Ingham PW. 1988. The molecular genetics of embryonic pattern formation in Drosophila. *Nature*
335:25–34. doi:10.1038/335025a0
Kanca O, Zirin J, Hu Y, Tepe B, Dutta D, Lin W-W, Ma L, Ge M, Zuo Z, Liu L-P, Levis RW,
Perrimon N, Bellen HJ. 2022. An expanded toolkit for Drosophila gene tagging using
synthesized homology donor constructs for CRISPR-mediated homologous
recombination. *eLife* **11**:e76077. doi:10.7554/eLife.76077
Keller J, Bassotti G, Clarke J, Dinning P, Fox M, Grover M, Hellström PM, Ke M, Layer P,
Malagelada C, Parkman HP, Scott SM, Tack J, Simren M, Törnblom H, Camilleri M,
International Working Group for Disorders of Gastrointestinal Motility and Function.
2018. Expert consensus document: Advances in the diagnosis and classification of
gastric and intestinal motility disorders. *Nat Rev Gastroenterol Hepatol* **15**:291–308.
doi:10.1038/nrgastro.2018.7
Klein AM, Mazutis L, Akartuna I, Tallapragada N, Veres A, Li V, Peshkin L, Weitz DA, Kirschner
MW. 2015. Droplet barcoding for single-cell transcriptomics applied to embryonic stem
cells. *Cell* **161**:1187–1201. doi:10.1016/j.cell.2015.04.044
Kurmangaliyev YZ, Yoo J, Valdes-Aleman J, Sanfilippo P, Zipursky SL. 2020. Transcriptional
Programs of Circuit Assembly in the Drosophila Visual System. *Neuron* **108**:1045-
1057.e6. doi:10.1016/J.NEURON.2020.10.006
Lee JH, Daugharthy ER, Scheiman J, Kalhor R, Ferrante TC, Terry R, Turczyk BM, Yang JL,
Lee HS, Aach J, Zhang K, Church GM. 2015. Fluorescent in situ sequencing (FISSEQ)
of RNA for gene expression profiling in intact cells and tissues. *Nature Protocols*
10:442–458. doi:10.1038/nprot.2014.191
Lein E, Borm LE, Linnarsson S. 2017. The promise of spatial transcriptomics for neuroscience
in the era of molecular cell typing. *Science* **358**:64–69. doi:10.1126/science.aan6827
Li H, Janssens J, De Waegeneer M, Kolluru SS, Davie K, Gardeux V, Saelens W, David FPA,
Brbić M, Spanier K, Leskovec J, McLaughlin CN, Xie Q, Jones RC, Brueckner K, Shim J,
Tattikota SG, Schnorrer F, Rust K, Nystul TG, Carvalho-Santos Z, Ribeiro C, Pal S,
Mahadevaraju S, Przytycka TM, Allen AM, Goodwin SF, Berry CW, Fuller MT, White-
Cooper H, Matunis EL, DiNardo S, Galenza A, O'Brien LE, Dow JAT, FCA Consortium,
Jasper H, Oliver B, Perrimon N, Deplancke B, Quake SR, Luo L, Aerts S. 2022. Fly Cell

Atlas: A single-nucleus transcriptomic atlas of the adult fruit fly. *Science* **375**:eabk2432.
doi:10.1126/science.abk2432
Ma D, Przybylski D, Abruzzi KC, Schlichting M, Li Q, Long X, Rosbash M. 2021. A
transcriptomic taxonomy of drosophila circadian neurons around the clock. *eLife* **10**:1–
19. doi:10.7554/eLife.63056
Macosko EZ, Basu A, Satija R, Nemesh J, Shekhar K, Goldman M, Tirosh I, Bialas AR,
Kamitaki N, Martersteck EM, Trombetta JJ, Weitz DA, Sanes JR, Shalek AK, Regev A,
McCarroll SA. 2015. Highly Parallel Genome-wide Expression Profiling of Individual
Cells Using Nanoliter Droplets. *Cell* **161**:1202–1214. doi:10.1016/j.cell.2015.05.002
Martin KC, Ephrussi A. 2009. mRNA localization: gene expression in the spatial dimension. *Cell*
136:719–30. doi:10.1016/j.cell.2009.01.044
Martin M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnet.journal **17**:10–12.
Marusyk A, Almendro V, Polyak K. 2012. Intra-tumour heterogeneity: a looking glass for
cancer? *Nature Reviews Cancer* **12**:323–334. doi:10.1038/nrc3261
McGinnis CS, Murrow LM, Gartner ZJ. 2019. DoubletFinder: Doublet Detection in Single-Cell
RNA Sequencing Data Using Artificial Nearest Neighbors. *Cell Systems* **8**:329-337.e4.
doi:10.1016/j.cels.2019.03.003
McKenna A, Findlay GM, Gagnon JA, Horwitz MS, Schier AF, Shendure J. 2016. Whole-
organism lineage tracing by combinatorial and cumulative genome editing. *Science*
353:aaf7907. doi:10.1126/science.aaf7907
Namiki S, Dickinson MH, Wong AM, Korff W, Card GM. 2018. The functional organization of
descending sensory-motor pathways in drosophila. *eLife* **7**.
Pfeiffer BD, Jenett A, Hammonds AS, Ngo T-TB, Misra S, Murphy C, Scully A, Carlson JW,
Wan KH, Laverty TR, Mungall C, Svirskas R, Kadonaga JT, Doe CQ, Eisen MB,
Celniker SE, Rubin GM. 2008. Tools for neuroanatomy and neurogenetics in Drosophila.
Proceedings of the National Academy of Sciences of the United States of America
105:9715–20. doi:10.1073/pnas.0803697105
Pfeiffer BD, Ngo T-TB, Hibbard KL, Murphy C, Jenett A, Truman JW, Rubin GM. 2010.
Refinement of tools for targeted gene expression in Drosophila. *Genetics* **186**:735–55.
doi:10.1534/genetics.110.119917
Reeves GT, Muratov CB, Schüpbach T, Shvartsman SY. 2006. Quantitative Models of
Developmental Pattern Formation. *Developmental Cell* **11**:289–300.
doi:10.1016/J.DEVCEL.2006.08.006
Rubin GM, Spradling AC. 1982. Genetic transformation of Drosophila with transposable element
vectors. *Science (New York, NY)* **218**:348–53. doi:10.1126/science.6289436
Satija R, Farrell JA, Gennert D, Schier AF, Regev A. 2015. Spatial reconstruction of single-cell
gene expression data. *Nature Biotechnology* **33**:495–502. doi:10.1038/nbt.3192
Shearin HK, MacDonald IS, Spector LP, Steven Stowers R. 2014. Hexameric GFP and mCherry
reporters for the Drosophila GAL4, Q, and LexA transcription systems. *Genetics*
196:951–960. doi:10.1534/GENETICS.113.161141/-/DC1/GENETICS.113.161141-
5.PDF
Smith AM, Heisler LE, Mellor J, Kaper F, Thompson MJ, Chee M, Roth FP, Giaever G, Nislow
C. 2009. Quantitative phenotyping via deep barcode sequencing. *Genome research*
19:1836–42. doi:10.1101/gr.093955.109
Stoeckius M, Hafemeister C, Stephenson W, Houck-Loomis B, Chattopadhyay PK, Swerdlow H,
Satija R, Smibert P. 2017. Simultaneous epitope and transcriptome measurement in
single cells. *Nature Methods* **14**:865–868. doi:10.1038/nmeth.4380
Stoeckius M, Zheng S, Houck-Loomis B, Hao S, Yeung BZ, Mauck WM, Smibert P, Satija R.
2018. Cell Hashing with barcoded antibodies enables multiplexing and doublet detection
for single cell genomics. *Genome Biology* **19**:1–12. doi:10.1186/s13059-018-1603-1

Teleman AA, Cohen SM. 2000. Dpp gradient formation in the drosophila wing imaginal disc.
Cell **103**:971–980. doi:10.1016/S0092-8674(00)00199-9
Thompson CL, Ng L, Menon V, Martinez S, Lee C-K, Glattfelder K, Sunkin SM, Henry A, Lau C,
Dang C, Garcia-Lopez R, Martinez-Ferre A, Pombero A, Rubenstein JLR, Wakeman
WB, Hohmann J, Dee N, Sodt AJ, Young R, Smith K, Nguyen T-N, Kidney J, Kuan L,
Jeromin A, Kaykas A, Miller J, Page D, Orta G, Bernard A, Riley Z, Smith S, Wohnoutka
P, Hawrylycz MJ, Puelles L, Jones AR. 2014. A High-Resolution Spatiotemporal Atlas of
Gene Expression of the Developing Mouse Brain. *Neuron* **83**:309–323.
doi:10.1016/J.NEURON.2014.05.033
van Opijnen T, Bodi KL, Camilli A. 2009. Tn-seq: high-throughput parallel sequencing for fitness
and genetic interaction studies in microorganisms. *Nature methods* **6**:767–72.
doi:10.1038/nmeth.1377
Venken KJT, Schulze KL, Haelterman NA, Pan H, He Y, Evans-Holm M, Carlson JW, Levis RW,
Spradling AC, Hoskins RA, Bellen HJ. 2011. MiMIC: a highly versatile transposon
insertion resource for engineering Drosophila melanogaster genes. *Nature methods*
8:737–43.
Yang S, Corbett SE, Koga Y, Wang Z, Johnson WE, Yajima M, Campbell JD. 2020.
Decontamination of ambient RNA in single-cell RNA-seq with DecontX. *Genome Biology*
21:57. doi:10.1186/s13059-020-1950-6
Yao Z, Scott K. 2022. Serotonergic neurons translate taste detection into internal nutrient
regulation. *Neuron* **110**:1036-1050.e7. doi:10.1016/j.neuron.2021.12.028
Zappia L, Oshlack A. 2018. Clustering trees: a visualization for evaluating clusterings at multiple
resolutions. *GigaScience* **7**:giy083. doi:10.1093/gigascience/giy083
Zheng GXY, Terry JM, Belgrader P, Ryvkin P, Bent ZW, Wilson R, Ziraldo SB, Wheeler TD,
McDermott GP, Zhu J, Gregory MT, Shuga J, Montesclaros L, Underwood JG,
Masquelier DA, Nishimura SY, Schnall-Levin M, Wyatt PW, Hindson CM, Bharadwaj R,
Wong A, Ness KD, Beppu LW, Deeg HJ, McFarland C, Loeb KR, Valente WJ, Ericson
NG, Stevens EA, Radich JP, Mikkelsen TS, Hindson BJ, Bielas JH. 2017. Massively
parallel digital transcriptional profiling of single cells. *Nature Communications* **8**:1–12.
doi:10.1038/ncomms14049
Zirin J, Jusiak B, Lopes R, Ewen-Campen B, Bosch JA, Risbeck A, Forman C, Villalta C, Hu Y,
Perrimon N. 2024. Expanding the Drosophila toolkit for dual control of gene expression.
eLife **12**:RP94073. doi:10.7554/eLife.94073
1064
1065

 Figures

A

NNN... - TaG-EM barcode
ca - polyA cleavage sites

B

TaG-EM constructs → PhiC31 integration → Individually barcoded Drosophila stocks

C Population behavioral barcoding

Experimental manipulation
• Exposure
• Time course
• Genetic perturbation
• Neural inactivation/activation

Behavioral assay → Extract DNA → Amplify and sequence barcodes

D *In vivo* deterministic tagging for single cell sequencing

Experimental manipulation
• Exposure
• Time course
• Genetic perturbation
• Replicate

Gal4-driven tissue-specific expression

Dissociate tissue → Enrich labeled cells (FACS) → Droplet cell capture → Sequence

Figure 1. Overview of TaG-EM system. A) Detailed view of the 3' UTR of the TaG-EM

constructs showing the position of the 14 bp barcode sequence (green highlight) relative to the

polyadenylation signal sequences (underlined) and polyA cleavage sites (purple highlights).

pJFRC12 backbone schematic is modified from Pfeiffer *et al.*, 2010. B) Schematic illustrating

the design of the TaG-EM constructs, where a barcode sequence is inserted in the 3' UTR of a

UAS-GFP construct and inserted in a specific genomic locus using PhiC31 integrase. C) Use of

TaG-EM barcodes for sequencing-based population behavioral assays. D) Use of TaG-EM

barcodes expressed with tissue-specific Gal4 drivers to label cell populations *in vivo* upstream

of cell isolation and single-cell sequencing.

1077

1078

A **Structured pool tests**

	Barcode																			
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
Female even	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
Male even	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
Female staggered	64	64	64	32	32	32	16	16	16	16	8	8	8	8	4	4	4	2	2	2
Male staggered	2	2	2	4	4	4	8	8	8	8	16	16	16	16	32	32	32	64	64	64

1079

Figure 2. Structured pool tests. A) Overview of the construction of the structured pools for

assessing the quantitative accuracy of TaG-EM barcode measurements. Male and female even

pools were constructed and extracted in triplicate. The table shows the number of flies that were

pooled for each experimental condition. B) Barcode abundance data for three independent

replicates of the female even pool. C) Barcode abundance data for three independent replicates

of the male even pool. D) Barcode abundance data for the female staggered pool. Inset plot

shows the average observed barcode abundance among lines pooled at each level compared

to the expected abundance. E) Barcode abundance data for the male staggered pool. Inset plot

shows the average observed barcode abundance among lines pooled at each level compared

to the expected abundance. For all plots, bars indicate the mean barcode abundance for three

technical replicates of each pool, error bars are +/- S.E.M.

Preference Index (P.I.) = $\dfrac{(\text{Light} - \text{Dark})}{(\text{Light} + \text{Dark})}$

Figure 3. TaG-EM barcode-based behavioral measurements. A) TaG-EM barcode lines in either a wild-type or *norpA* background were pooled and tested in a phototaxis assay. After 30

seconds of light exposure, flies in tubes facing the light or dark side of the chamber were

collected, DNA was extracted, and TaG-EM barcodes were amplified and sequenced. Barcode

abundance values were scaled to the number of flies in each tube and used to calculate a

preference index (P.I.). Average P.I. values for four different TaG-EM barcode lines in both the

wild-type and *norpA* backgrounds are shown (n=3 biological replicates, error bars are +/-

S.E.M.). B) The same eight lines used for the sequencing-based TaG-EM barcode

measurements were independently tested in the phototaxis assay and manually scored videos

were used to calculate a P.I. for each genotype. Average P.I. values for each line are shown

(n=3 biological replicates, error bars are +/- S.E.M.) for TaG-EM-based quantification (top) and

manual video-based quantification (bottom). C) Flies carrying different TaG-EM barcodes were

collected and aged for one to four weeks and then eggs were collected and egg number and

viability was manually scored for each line. In parallel the barcoded flies from each timepoint

were pooled, and eggs were collected, aged, and DNA was extracted, followed by TaG-EM

barcode amplification and sequencing. Average number of viable eggs per female (manual

counts) and average barcode abundance are shown both as a bar plot and scatter plot (n=3

biological replicates for 3 barcodes per condition, error bars are +/- S.E.M.).

1110

1111

Figure 4. TaG-EM barcode-based quantification of larval gut motility. Schematics depicting

A) manual and B) TaG-EM-based assays for quantifying food transit time in Drosophila larvae.

1114   C) Transit time of a food bolus in the presence and absence of caffeine measured using the

1115   manual assay (p = 0.0340). D) Transit time of a food bolus in the presence and absence of

1116   caffeine measured using the TaG-EM assay (p = 0.0488). A modified Chi-squared method was

1117   used for statistical testing (Hristova and Wimley, 2023).

1118

Figure 5. Gal4 driven expression of GFP from TaG-EM lines. A) Comparison of endogenous
GFP expression and GFP antibody staining in the wing imaginal disc for the original pJFRC12
construct inserted in the attP2 landing site or for a TaG-EM barcode line driven by *dpp-Gal4*.
Wing discs are counterstained with DAPI. B) Endogenous expression of GFP from either a TaG-
EM barcode construct (left column), a hexameric GFP construct (middle column), or a line
carrying both a TaG-EM barcode construct and a hexameric GFP construct (right column)
driven by the indicated gut driver line (PMG-Gal4: Pan-midgut driver; EC-Gal4: Enterocyte
driver; EE-Gal4: Enteroendocrine driver; EB-Gal4: Enteroblast driver).

1129

Figure 6. Expression of TaG-EM genetic barcodes in larval intestinal cell types. A) UMAP

plot of Drosophila larval gut cell types. B) Annotation of cells associated with a TaG-EM barcode

across all 8 multiplexed experimental conditions using data from the gene expression library

and an enriched TaG-EM barcode library. C) Annotated enteroblast cells. D) Presence of TaG-

EM barcode (BC6) driven by the EB-Gal4 line using data from the gene expression library and

an enriched TaG-EM barcode library. Gene expression levels of enteroblast marker genes E)

esg, F) *klu*. G) Annotated enterocyte cells. H) Presence of TaG-EM barcode (BC4) driven by the

EC-Gal4 line using data from the gene expression library and an enriched TaG-EM barcode

library. Gene expression levels of enterocyte marker genes I) *betaTry*, J) *Jon99Ciii*. K)

Annotated enteroendocrine cells. L) Presence of TaG-EM barcode (BC9) driven by the EE-Gal4

line using data from the gene expression library and an enriched TaG-EM barcode library. Gene

expression levels of enteroendocrine cell marker genes M) *Dh31*, N) *IA-2*.

Supplemental Figures

A

Barcode sequence	Transgenic line
ATTTTGCGCCTTAG	1-2
CCCTAAAGTACCGT	4-3
TAGTTAGGCGGAAA	5-4
TTGCTCTTTCGTTT	6-4
TTGCTGCGCACGGC	7-1
CGTTTATATCCTGT	10-1
AGCAGGTGATGTAC	1-1, 1-3, 10-2
TATGCCTCATTTCT	13-2
GTTTGATCAGTACC	13-4
GGTTTACCACAAAG	14-2
AGACACATCCTGTG	15-1
GAATTTATCAGTTA	15-2
TCTTCATGACCAGC	2-1, 2-2, 2-3, 11-1, 11-2, 11-4
TATGTCAATATGGA	3-1, 3-2, 3-3, 9-2
ACTCTCTCTTAAGG	4-1, 4-4
CTTTAATTGCAGTC	5-3, 5-5, 9-1, 9-3
TACTGTACATGCTC	6-2, 10-4, 14-3, 14-4
TACCTTTGAATTCG	6-3, 12-2, 12-4
CATCGAGTTCACCT	8-1, 12-3
GTCCTTCTACCCCG	8-2, 8-4

B

Supplemental Figure 1. Sanger sequencing identification of TaG-EM barcode lines. A) Summary of barcode pool injections. Barcode sequence and transgenic vial identifier in which the barcode was identified are shown. B) Sanger sequencing-based confirmation of the barcode sequence and PCR handle in TaG-EM transgenic lines. Because the TaG-EM barcode constructs were injected as a pool of 29 purified plasmids, some of the transgenic lines had

1149   inserts of the same construct. In total 20 unique lines were recovered from this round of

1150   injection.

1151

1152

1153

Supplemental Figure 2. Optimization of TaG-EM barcode amplification. A) Gels showing

bands produced when amplifying TaG-EM flies or a wild type control with the indicated

polymerase, annealing temperature, and primer pair (short = B2_3'F1_Nextera/

SV40_pre_R_Nextera; long = B2_3'F1_Nextera/ SV40_post_R_Nextera). B-E) Mean error

(R.M.S.D. root mean squared deviation from expected value) for even pool amplified with the

indicated primer set, input amount, and cycle number using KAPA HiFi polymerase (n=3, error

bars are +/- S.E.M.). F-G) Mean error (R.M.S.D. root mean squared deviation from expected

value) for staggered pool amplified with the indicated primer set, input amount, and cycle

number using KAPA HiFi polymerase (n=3, error bars are +/- S.E.M.).

1163

1164

1165 **Supplemental Figure 3. Coeffient of variation for TaG-EM structured pools.** Plot showing

1166 coefficient of variation for different groups of TaG-EM barcodes in the structured pools. Dashed

1167 line indicates the mean coefficient of variation across all conditions.

1168

1169

1170

1171 **Supplemental Figure 4. Oviposition tests with TaG-EM barcode lines.** Plots showing mean

1172 TaG-EM barcode abundance for adult females used in oviposition experiments (top) and eggs

1173 collected from these females (bottom). Data from two independent trials is shown (n=3 for each

1174 trial, error bars are +/- S.E.M.). Dashed lines indicate the expected abundance values.

1175

1176

Supplemental Figure 5. Fecundity data for individual TaG-EM lines. Manually collected data

for mean number of viable eggs per female, barcode abundance data, and barcode abundance

data normalized to adult fly barcode data for each of the TaG-EM barcode lines used in the age-

dependent fecundity experiment. Scatterplots show correlations between manually collected

data and barcode sequencing results. Data from two independent trials is shown (n=3 for each

trial, error bars are +/- S.E.M.).

1183

1184 **Supplemental Figure 6. Average age-dependent fecundity data for Trial 1.** Average number

1185 of viable eggs per female (manual counts) and average barcode abundance are shown both as

1186 a bar plot and scatter plot (n=3 biological replicates for 3 barcodes per condition, error bars are

1187 +/- S.E.M.). Data from Trial 2 is shown in Figure 3C.

1188

1189

Supplemental Figure 7. Larval gut motility assay parameters. A) Images of larvae fed with

blue-dyed yeast agar. B) Effect of dye concentration on food transit time. C) Effect of starvation

1192   time on feeding and uptake of the dyed food bolus. D) Effect of liquid versus solid diet on food

1193   transit time. E) Aversive effect of caffeine on food bolus uptake.

A Cost per Datapoint Analysis

B Monetary Cost Analysis ($USD)

C Labor Time Cost Analysis (Hours)

1194

1195 **Supplemental Figure 8. Cost comparisons for manual and TaG-EM gut motility assays.** A)

1196 Cost per data point as a function of the number of data points generated and the number of

1197   experimental conditions screened. B) Overall experiment cost and C) labor effort as a function

1198   of the number of data points generated and the number of experimental conditions screened.

1199

Supplemental Figure 9. Expression driven by *dpp-Gal4* for 20 TaG-EM lines. GFP antibody staining in the wing imaginal disc for the indicated TaG-EM barcode line driven by *dpp-Gal4*. Wing discs are counterstained with DAPI.

A

B

1203

1204 **Supplemental Figure 10. TaG-EM line GFP expression driven by different Gal4 drivers.** A)

1205 Comparison of endogenous GFP expression in larvae for the original pJFRC12 construct

1206 inserted in the attP2 landing site (left) or for a TaG-EM barcode line (right) expressed under the

1207 control of the indicated driver line. B) GFP expression of the PC-Gal (Precursor-Gal4) driver line

1208 together with either UAS-2xGFP or a combination of UAS-2xGFP and a TaG-EM barcode line.

1209

1210

1211 **Supplemental Figure 11. Dissociated intestinal cell viability.** A) GFP expression visualized

1212 in dissociated cells from gut driver lines crossed to hexameric GFP and TaG-EM line. B)

1213 Proportion of live (left) and dead (right) cells post-isolation and flow sorting as assessed by GFP

1214 expression and propidium iodide staining.

1215

1216

1217 **Supplemental Figure 12. BD FACSDiva 8.0.1 gating for sorted cells.** A) GFP gating created

1218 by analyzing a pool of GFP positive and negative cells. B) Flow gating for Drosophila gut cells

1219 with TaG-EM GFP expression driven in intestinal precursor cells (PC-Gal4) and enterocytes

1220 (EC-Gal4).

1221

1222

1223

Supplemental Figure 13. Expression of TaG-EM genetic barcodes in larval intestinal

precursor cells. UMAP plots showing gene expression levels of A) enteroblast/ISC marker

genes *esg*, *klu*, and *E(spl)mbeta-HLH*; and B) the TaG-EM barcodes 7, 8, and 9 driven by the

PC-Gal4 line.

1228

1229

1230

1231 **Supplemental Figure 14. BD FACSDiva 8.0.1 gating for sorted cells.** A) Dead cell gating

1232 created by staining sample with propidium iodine (PI). B) Flow gating for *Drosophila* gut cells

1233 with TaG-EM and hexameric GFP expression driven by the pan-midgut, enteroblast, enterocyte,

1234 enteroendocrine, and precursor cell drivers.

1235

1236

1237

Supplemental Figure 15. TaG-EM-based doublet identification. UMAP plots pre-doublet

removal showing A) doublets uniquely identified by DoubletFinder, B) all doublets identified by

1240   DoubletFinder, C) doublets uniquely identified by TaG-EM barcodes, D) all doublets identified

1241   by TaG-EM barcodes, E) doublets mutually found by TaG-EM and DoubletFinder, F) Venn

1242   diagram of overlap between doublets identified by TaG-EM and DoubletFinder.

1243

1244

1245

Supplemental Figure 16. Clustering and automated annotation. A) UMAP plots clustered at

different resolutions. B) Clustree analysis of the effect of clustering resolution. C) Automated cell

type annotation using data from the Fly Cell Atlas.

1249

1250

Supplemental Figure 17. Expression of TaG-EM genetic barcodes in larval intestinal cell

types. A) UMAP plot of Drosophila larval gut cell types. B) Annotation of cells associated with a

TaG-EM barcode across all 8 multiplexed experimental conditions using data from the gene

expression library only. C) Annotated enteroblast cells. D) Expression level of TaG-EM barcode

(BC6) driven by the EB-Gal4 line using data from the gene expression library only. Gene

expression levels of enteroblast marker genes E) *esg*, F) *klu*. G) Annotated enterocyte cells. H)

Expression level of TaG-EM barcode (BC4) driven by the EC-Gal4 line using data from the gene

expression library only. Gene expression levels of enterocyte marker genes I) *betaTry*, J)

Jon99Ciii. K) Annotated enteroendocrine cells. L) Expression level of TaG-EM barcode (BC9)

driven by the EE-Gal4 line using data from the gene expression library only. Gene expression

levels of enteroendocrine cell marker genes M) *Dh31*, N) *IA-2*.

1262

1263

Supplemental Figure 18. Optimizing amplification of the TaG-EM barcode library. A)

Workflow for single-cell capture; cDNA amplification with added spike-in primer for TaG-EM

library followed by a SPRI size-selection clean-up, then PCR(s) to create library for sequencing.

B) Spike-in primers and amplification primers used to enrich TaG-EM barcodes. Table

1268   summarizes different protocols tested to amplify the TaG-EM barcodes and create an enriched

1269   sequencing library. C) Percent of on-target reads for each enriched TaG-EM barcode library.

1270

1271

1272

Supplemental Figure 19. Performance of the enriched TaG-EM barcode library. A)
Proportion of cells with at least one barcode read assigned as a function of read depth for the
enriched TaG-EM barcode library. Dashed line indicated percentage of cells with TaG-EM
barcodes detected in the gene expression library B) Number of unique UMIs observed as a
function of read depth. C) Correlation between barcodes detected in the gene expression (GEX)
library and the enriched TaG-EM barcode library as a function of the purity of TaG-EM barcode
assignment to the corresponding cell barcode. Dashed line indicates the threshold used for
TaG-EM barcode calling in the enriched TaG-EM barcode library.

A

Pan-midgut-associated TaG-EM barcode expression
(BC1, BC2, BC3, BC7)

B

1281

1282   **Supplemental Figure 20. Expression of the PMG-Gal4 driven TaG-EM barcodes.** UMAP

1283   plots showing expression of the four PMG-Gal4 driven TaG-EM barcodes (BC1, BC2, BC3, and

1284   BC7) either A) in aggregate or B) individually.

1285

Supplemental Figure 21. Characterization of Gal4 line expression in the larval gut. A) Confocal images of third instar midguts showing Gal4-driven fluorophore expression (GFP or mCherry) and comparison with immunostainings of the gut cell markers Prospero (enteroendocrine), Pdm1 (enterocyte) and Esg-GFP (progenitor cell). For each image, Z projections of the stacks recorded along the length of the midgut were manually stitched together. B) Representative single frames confocal images of a small region of the midgut showing immunostainings of the different gut cell markers and the Gal4-driven fluorophores. Quantification of overlapping and non-overlapping expression between the Gal4-driver fluorophore expression

1295   and the cell type marker in the anterior (A), middle (M), and posterior (P) regions for C)

1296   enteroendocrine cells (EC-Gal4), D) enterocytes (EC-Gal4), E) precursor cells (PC-Gal4). Five

1297   specimens for each Gal4 line were examined. In the case of the enterocyte-specific driver, only

1298   anterior and middle regions were analyzed since the driver is largely inactive in the posterior part

1299   of the midgut.

1300