Estimating receptive fields of simple and complex cells in early visual cortex: A convolutional neural network model with parameterized rectification

running head: Convolutional neural network model of A17/18 cells

Philippe Nguyen[1a#], Jinani Sooriyaarachchi[2#], Qianyu Huang[3b], & Curtis L. Baker Jr.[4*]

[1]Department of Biomedical Engineering, McGill University, Montreal, Quebec, Canada

[2]Department of Physiology, McGill University, Montreal, Quebec, Canada

[3]Department of Biology, McGill University, Montreal, Quebec, Canada

[4]Department of Ophthalmology and Visual Sciences, McGill University, Montreal, Quebec, Canada

[a]Current Address: Resson Aerospace | AgTech, 46 Dineen Dr, Frederickton, New Brunswick, Canada H3B 9W4

[b]Current Address: SSW Consulting, 406 Xintiandi St, Hangzhou, Zhejiang Province, China

[#] Joint first authors

[*] Corresponding Author:

Email: curtis.baker@mcgill.ca

Abstract

Neurons in the primary visual cortex respond selectively to simple features of visual stimuli, such as orientation and spatial frequency. Simple cells, which have phase-sensitive responses, can be modeled by a single receptive field filter in a linear-nonlinear model. However, it is challenging to analyze phase-invariant complex cells, which require more elaborate models having a combination of nonlinear subunits. Estimating parameters of these models is made more difficult by cortical neurons' trial-to-trial response variability.

We develop a simple convolutional neural network method to estimate receptive field models for both simple and complex visual cortex cells from their responses to natural images. The model consists of a spatiotemporal filter, a parameterized rectifier unit (PReLU), and a two-dimensional Gaussian "map" of the receptive field envelope. A single model parameter determines the simple vs. complex nature of the receptive field, capturing complex cell responses as a summation of homogeneous subunits, and collapsing to a linear-nonlinear model for simple type cells.

The convolutional method predicts simple and complex cell responses to natural image stimuli as well as grating tuning curves. The model estimates yield a continuum of values for the PReLU parameter across the sampled neurons, showing that the simple/complex nature of cells can vary in a continuous manner. We demonstrate that complex cells respond less reliably than simple cells - compensation for this unreliability reveals good predictive performance on novel sets of natural images, with predictive performance for complex cells proportionately closer to that for simple cells. Most spatial receptive field structures are well fit by Gabor functions, whose parameters confirm well-known properties of cat A17/18 receptive fields.

New and Noteworthy

Methods for recording increasingly many visual cortex neurons are advancing rapidly, demanding new approaches to characterize diverse receptive fields. We present a compact convolutional neural network model of early cortical neurons, which is uniformly applicable to simple and complex cells, and whose parameters are simple to interpret and readily estimated from responses to natural image stimuli. This novel approach introduces a single estimated model parameter to capture the simple/complex nature of a neuron's receptive field, revealing a continuum of simple vs complex-like behaviour. We show that almost all complex-like cells exhibit a lower response reliability to repeated presentations of the same stimuli, compared to more reliable responses from simple-like cells. Accounting for this "noise ceiling" brings predictive performance for complex cells proportionately closer to that for simple cells. This model estimation approach reveals a diversity of spatial receptive field shapes, which are nevertheless generally well fit by Gabor functions. The results extend earlier findings with laboratory stimuli (white noise, gratings) to responses to natural images, for both simple and complex cells in cat A17 and A18.

Keywords

visual cortex, receptive field, simple cell, complex cell, convolutional neural networks

Introduction

Neurons in the striate cortex are the first stage of visual processing to provide a rich representation of the elementary features needed to efficiently encode natural images (Olshausen, 1996). Since the work of (Hubel & Wiesel, 1962) these neurons have been extensively studied with simple stimuli such as bars, sinewave gratings, or white noise. However, predicting responses to arbitrarily complex stimuli such as natural images require quantitative models that are selective to both location in visual space and stimulus features within those locations, including estimation of the model parameters through system identification (Wu et al., 2006). Simple cells, a subset of striate cortex neurons, have distinct excitatory and inhibitory receptive field regions with linear spatial summation, and are phase-sensitive, i.e., give modulated responses to drifting gratings. Thus they are well modeled by a linear-nonlinear (LN) model (Simoncelli et al., 2004), which is relatively easy to experimentally estimate. Complex cells, however, are phase-insensitive, and exhibit mixed excitatory/inhibitory responses throughout the receptive field - they are generally thought to be integrating responses from several nearby linear-nonlinear subunits, which are more difficult to estimate. The task is further complicated by significant variability in responses of cortical neurons (Carandini, 2004; Tolhurst et al., 1983; Vogels et al., 1989).

The development of improved methods to quantitatively characterize the diverse receptive fields of visual cortex neurons is becoming more important, in order to effectively exploit newer recording technologies that simultaneously record large numbers of single neurons, such as two-photon imaging and high-density electrophysiology (Blanche et al., 2005; Jun et al., 2017). In such situations the neurons may differ in multiple ways, which could render conventional tuning curve approaches problematic.

Receptive fields of simple type cells have been traditionally estimated by spike-triggered averaging (STA) of responses to white noise (e.g. DeAngelis et al., 1993; Marmarelis & Marmarelis, 1978; Schwartz et al., 2006). The nonlinear subunit structure of complex cell receptive fields, however, has generally required more elaborate methods such as spike-triggered covariance (Schwartz et al., 2006). However, these methods produce more complicated models (multiple eigenfunctions), requiring many more parameters than the

linear-nonlinear model, and are therefore harder to estimate and may give poorer predictive performance. The use of different methods to analyze simple and complex cells is at odds with evidence indicating that simple and complex cells may not be wholly physiologically different (Mechler & Ringach, 2002). A combination of STA and STC approaches can be used on both simple and complex cells (Fournier et al., 2014; Schwartz et al., 2006), but the resultant models are even more heavily parameterized, as well as being difficult to interpret. It would be highly desirable to have a method that works uniformly on simple or complex cells, or intermediate types, providing estimated models without over-parameterization.

Receptive fields of visual neurons vary in important ways not only in their spatial properties, but also in their temporal dynamics - for example, sustained vs transient or lagged vs non-lagged behaviour (Saul & Humphrey, 1992), spatiotemporal non-separability (McLean et al., 1994), or temporal frequency response (Movshon et al, 1978). Many previous system identification studies of visual cortex neurons have only considered spatially 2D models, without time dynamics, or have included temporal properties at the expense of employing spatially 1-D stimuli (e.g., random bar stimuli) to estimate a "space-time" receptive field (e.g., Baker, 1988; DeAngelis et al, 1993). A full 3D analysis (space-space-time) has been challenging due to the much larger number of model parameters to be estimated. However, this would be most desirable for revealing the full spatiotemporal properties of receptive fields.

Most previous system identification methods for early visual neurons have necessitated the use of spatiotemporally uncorrelated ("white") stimuli. However, it is desirable to have methods that work for more complex stimuli such as natural images, which potentially provide results more relevant to normal operating ranges of image statistics (David et al., 2004; Touryan et al., 2005). Furthermore, neurons beyond the very earliest stages of visual processing may give poor responses to white noise and require complex stimuli to give responses more suitable for system identification (Talebi & Baker, 2012; Zhang et al., 2019). It is possible to modify STA and STC-style approaches for responses to natural images (Lochmann et al., 2013; Oliver, 2014; Theunissen et al., 2001; Touryan et al., 2005; Willmore & Smyth, 2003). Another approach involves estimating a general linear model that sums responses of a wavelet basis

designed to mimic early-stage neurons (Ikezoe et al., 2018; Oliver, 2014; Willmore et al., 2010), though this requires strong prior assumptions about the nature of the earlier-stage filters.

Recently, convolutional neural networks (CNNs) have become widely used in engineering/computer science applications due to their state of the art performance, particularly on image-related problems (Krizhevsky et al., 2012). These models typically use a succession of layers, each having multiple spatially homogeneous convolutional filters, separated by half-wave rectifiers. Such multi-layer "deep" neural networks (DNNs) have been used to estimate responses of neurons in striate cortex (Cadena et al, 2017; Ecker et al, 2018; Kindel et al, 2019; Zhang & Young, 2020). However, it can be unclear how to interpret intermediate stages of such complicated model architectures. In addition, such DNNs have previously been estimated only for 2-D spatial receptive field models (i.e., without time dynamics). To extend such approaches to full 3-D space-space-time models, it would seem problematic to estimate the multitude of parameters in such complex models using limited sets of data from the noisy responses of cortical neurons.

Here we present a compact convolutional neural network model, with parameters that are simple to interpret: a single initial 3-D spatiotemporal filter captures the feature selectivity, a single parameter explicitly models the simple/complex behaviour of the neuron (phase sensitivity), and a final layer localizes the filter activity (overall envelope of receptive field). The parameters of the model can be solved directly by gradient descent, using responses to natural images. The convolutional model prediction of neuron response performs well for most single neurons from cat A17 and A18, both for natural images and for tuning curves from sinewave gratings. The phase sensitivity estimates of the convolutional model trained on natural images is highly correlated with neurons' simple vs. complex type phase responses to sinewave gratings. Our analysis also shows that complex cell responses are inherently less reliable than those for simple cells. Accounting for this "noise ceiling" brings predictive performance for complex cells proportionately closer to that for simple cells.

We also demonstrate how the approach can be used to measure cortical receptive field properties such as size and optimal spatial frequency, and how the distributions of these properties are related to simple vs. complex type responses and to the cortical area from which

they are recorded. The results reveal receptive field shapes that may be Gabor-like, with diverse aspect ratios, as well as numerous instances of non-oriented receptive fields (e.g., centre-surround organization).

Materials and Methods

Animal Preparation

The procedures for animal surgery, recording, and maintenance were conventional and have been described in detail (St-Amand & Baker, 2022), so here they will be only briefly summarized. Anesthesia in adult cats of either sex was induced with isofluorane/oxygen (3–5%), followed by intravenous (i.v.) cannulation. Surgical anesthesia was then provided with i.v. propofol - first a bolus i.v. injection (5 mg·kg^{-1}), followed by a continuous infusion (5.3 mg·kg^{-1}·h^{-1}), and supplemental doses of propofol as necessary. Intravenous doses of glycopyrrolate (30 µg) and dexamethasone (1.8 mg) were provided, and a tracheal cannula or intubation tube was positioned for a secure airway. Body temperature was maintained with a thermostatically controlled heating pad, and heart rate was monitored (Vet/Ox Plus 4700).

The animal was then secured in a stereotaxic apparatus, and a craniotomy and small duratomy were performed over the cortical region of interest (area 17, P3/L1, or area 18, A3/L4), which was then protected by 2% agarose and petroleum jelly. Body temperature was kept constant throughout, using a thermister-controlled heating pad. All surgical sites were infused with local injections of bupivacaine (0.50%).

After completion of all surgical procedures, the animal was connected to a ventilator (Ugo Basile 6025) supplying oxygen/nitrous oxide (70:30). Additional anaesthesia was provided with i.v. remifentanil (bolus injection, 1.25 µg·kg^{-1}, followed by infusion, 3.7 µg·kg^{-1}·h^{-1}), with a reduced propofol infusion rate (5.3 mg·kg^{-1}·h^{-1}). Paralysis was then induced and maintained with i.v. gallamine triethiodide (bolus: to effect, followed by infusion: 10mg·kg^{-1}·h^{-1}).

Additional doses of glycopyrrolate (16 µg) and dexamethasone (1.8 mg), intramuscular, were also provided daily. Recording experiments were conducted over a period of 3-4 days,

during which vital signs (temperature, heart rate, expired CO_2 and EEG activity) were monitored and kept at normal levels.

Topical ophthalmic carboxymethylcellulose (1%) was used for protection of the corneas during initial surgery. Neutral contact lenses were later inserted for long-term corneal protection. Spectacle lenses were used to bring the eyes into focus at 57 cm, the distance at which visual stimuli would be presented. The optical quality was further improved with artificial pupils (2.5 mm). Topical phenylephrine hydrochloride (2.5%), to retract nictitating membranes, a mydiatic (atropine sulfate, 1%, or cyclopentolate, 1.0 %, in later experiments) were applied daily.

All animal procedures were approved by the Animal Care Committee of the Research Institute, McGill University Health Centre, and are in accordance with the guidelines of the Canadian Council on Animal Care.

Visual Stimuli and Electrophysiology

Visual stimuli were presented on a gamma-corrected CRT monitor (NEC FP1350, 20 inches, 640x480 pixels, 150 Hz, 36 cd/m^2) at a viewing distance of 57 cm. Stimuli were produced by an Apple Macintosh computer (MacPro, 2.66 GHz, 6 GB, MacOSX ver. 10.6.8, NVIDIA GeForce GT 120) using custom software written in MATLAB (ver. 2012b) with the Psychophysics Toolbox (ver. 3.0.10; Brainard, 1997; Pelli, 1997). A photometer (United Detector Technology) was used to measure the monitor's gamma nonlinearity, which was corrected using inverse lookup tables. Signals from a photocell (TAOS, TSL12S) in one corner of the monitor were used for temporal registration of stimulus onset/offset timing and spike recordings, and to verify the absence of dropped frames.

Extracellular recordings were obtained via 32-channel multielectrode probes (NeuroNexus, linear array: A1x32-6mm-100-177, or polytrode: A1x32-Poly2-5mm-50s-177), and a Plexon Recorder data acquisition system. A single-channel window discriminator was used to isolate spiking responses from a single channel (300-5kHz) of interest during the experiment ("on-line"). However all 32 broadband (3Hz-5kHz) data channels (as well as CRT photocell signals, TAOS TSL12S) were streamed to hard disk for later off-line spike waveform classification

using SpikeSorter software (Swindale & Spacek, 2015), and detailed analyses of each neuron's responses.

Electrodes were inserted in approximately vertical, near-columnar penetrations, so that neurons generally had similar receptive field location, preferred spatial frequency, and preferred orientation (DeAngelis et al., 1999; Hubel & Wiesel, 1962; Hubel & Wiesel, 1968; Tootell et al., 1981). An interactively controlled bar stimulus was used to get an initial estimate of the neurons' receptive field location and dominant eye, from a representative channel. The non-dominant eye was occluded, and the CRT centered approximately on the receptive field.

For subsequent quantitative experiments, visual stimuli (gratings, natural images) were presented on the CRT display. The responses of one neuron on one channel were chosen to guide the optimization of grating stimuli. Drifting sinewave gratings were circularly cosine-tapered against a blank background at mean luminance. Tuning curve measurements were obtained from gratings presented in random order in 1 sec trials, with at least 10 repetitions of each condition. Only the background was presented for "blank" trials which were used to estimate spontaneous activity. Each neuron's "AC/DC ratio" (first harmonic divided by mean response) was calculated from the response to its optimal grating (Skottun et al., 1991).

Natural image stimuli were constructed from monochrome 480x480 pixel subsets of photographs from the McGill Calibrated Color Image Database (Olmos & Kingdom, 2004). Images having low pixel standard deviations were discarded, since they were nearly blank. The remaining images had their mean luminance subtracted and were RMS-normalized and clipped to 8 bits. These images were the same as used previously (Talebi & Baker, 2012), but with a higher RMS contrast. Ensembles of 375 such images were presented in 5 sec trials, preceded by 100 msec of mean luminance blank screen in order to estimate spontaneous activity. Though the CRT refresh rate was 150Hz, each stimulus frame was presented twice for an effective rate of 75Hz. Stimulus image ensembles were divided into 3 sets, designed to be used for training, regularization, and testing (see below). There were 20 estimation ensembles, each presented 5 times. Both regularization and test sets consisted of 5 ensembles and were presented 20 times each. The number of ensembles and repetitions were chosen as a trade-off between maximizing the distinct stimuli presented to the neuron while reducing the effect of trial-by-

trial variability. Image ensemble sets were presented in pseudorandom order over a period of ca. 45 minutes.

Model estimation

Convolutional neural network (CNN) model

To handle both complex and simple type cells, we introduce a "convolutional" model (Figure 1):

$$\hat{R}_t = N(\sum_{x,y} w[x,y]G(\sum_{j,k,\tau} c[j,k,\tau]s[x-j,y-k,t-\tau]) \tag{1}$$

In this model, the stimulus **s** is first spatiotemporally convolved with the filter **c**. The output of this convolution, at a given time *t*, is a two-dimensional image, which is then passed through a static nonlinearity *G*. It can be seen that at a fixed point (x, y), the convolution $(c^T s_t(x,y))$ is similar to the linear component of the linear-nonlinear model for a small region of the full stimulus. Similarly, $G(c^T s_t(x,y))$ can be thought of as similar to the linear-nonlinear model operating on a small region of the full stimulus. Note that the full convolution operation, $G(c * s_t)$, applies the linear-nonlinear model to every possible *nxn* sub-window of the stimulus, where *n* is the size of the filter **c**. Thus, the operation $G(c * s_t)$ can be considered as the activity of a homogeneous array of linear-nonlinear subunits in response to the stimulus s_t. Each subunit will have the same filter **c**, where the subunits are tiled across the stimulus space. The final set of weights, **w**, parameterizes the regions of visual space where the subunit activity is predictive of the neuron's response **R**. The final output nonlinearity, N, is parameterized as a rectified power law exponent.

Figure 1. Convolutional model for simple and complex type visual cortex neurons. Stimulus frames are spatiotemporally convolved with the filter layer, then passed through a Parametric Rectified Linear Unit (PReLU) non-linearity. The dot product between the PReLU layer output and the spatial map layer generates the neuron response. A final rectified power law nonlinearity is added to account for the spiking threshold and possible nonlinear spike frequency response. Cyan elements represent estimatable parameters.

Instead of fixing $G(x)$ as a half-wave rectifier, we make a slight adjustment to allow for estimation of the linear-nonlinear model as well as the subunit model. We parameterize $G(x)$ as a "rectified linear unit" (PReLU), a nonlinearity that has been shown to improve the performance of artificial neural networks trained to classify images (He et al., 2015). The PReLU is a simple static piecewise-linear rectifier:

$$G(x) = \begin{cases} x, & if\ x > 0 \\ \alpha x & if\ x \le 0 \end{cases} \tag{2}$$

The PReLU has a single parameter α which denotes the slope of the negative component of the rectifier. If $\alpha = 0$, then G acts as a half-wave rectifier; this would correspond to a simple cell instance of the subunit model as mentioned above. If $\alpha = 1$, then G is the identity function $(G(x) = x)$ and the linear weights of both **w** and **c** will be in a linear cascade, which is equivalent to a single linear layer - thus the convolutional model then collapses to the linear-nonlinear model.

The nonlinearity G could be modeled more generally, for example as a set of piecewise linear B-spline ("tent") basis functions (De Boor, 1978; Vintch et al., 2015), which provide a local approximation of high order polynomials. However, when using a tent basis, G cannot be estimated concurrently (i.e., within the same gradient descent) as the convolutional filter c and map weights w. This problem can be solved by using an EM-like algorithm such as alternating least squares (Takane et al., 1977; Vintch et al., 2015), that alternately keeps one part fixed while optimizing the other. However, such methods take longer to calculate, might incur a greater risk of convergence on local optima, and may be less robust with choice of initial conditions. Modelling G as a PReLU allows us to estimate α jointly with the other parameters of the model, since it can be incorporated into the same gradient descent backpropagation that estimates the filter and map functions.

We refer to w as the "map" layer, because it indicates the regions in visual space where the subunit filter c is operative. Prior knowledge of the receptive field shape can be used to define model parameters (DiMattina & Zhang, 2011; St-Yves & Naselaris, 2018). The map layer embodies the spatial envelope of the neuron's receptive field, which for early visual cortex neurons can be parameterized as a two-dimensional Gaussian (DeAngelis et al., 1991). Thus, the map layer w can be defined with only six parameters - two for the means (μ_X, μ_Y), three for the variances (covariance matrix Σ), and a scale factor β:

$$w[x,y] = \beta \cdot \eta_{\mu,\Sigma}(x,y) = \beta \cdot \frac{\exp\left(-\frac{1}{2}(x-\mu_X; y-\mu_Y)\Sigma^{-1}(x-\mu_X; y-\mu_Y)^T\right)}{2\pi\sqrt{|\Sigma|}} \tag{3}$$

We also estimate the N power law exponent in the final layer of the model. Simultaneously estimating N with other model parameters using backpropagation is problematic due to "gradient explosion". Therefore, the final output nonlinearity N is estimated in two stages. In the first stage, N is set to be a simple ReLU activation function and an initial estimate of the receptive field and other model parameters are obtained. In the second stage, the pre-trained receptive field weights are loaded, and the model is re-trained with N set to be an adaptive exponent output nonlinearity defined in a custom Keras layer with two trainable parameters gain (g) and power law exponent (exp) as:

$$N(x) = \begin{cases} g.x^{exp}, & if \ x > 0 \\ 0 & if \ x \leq 0 \end{cases}$$

(4)

Estimation of model parameters

Data files were converted from the Plexon file format (.plx) to MATLAB data files (.mat) using in-house code. Spikes were sorted (Swindale & Spacek, 2015) independently for each stimulus set (i.e. natural image ensembles, tuning curves for different grating parameters). Sorted waveforms were then matched (Euclidean similarity) to establish corresponding neurons across the different stimulus sets. Only matches with a high degree of confidence were accepted for use in those analyses requiring comparisons of responses to different stimuli. The neuron response was averaged over the repetitions of each natural movie stimulus ensemble to be used in receptive field estimation.

In methods for estimating receptive field models from responses to complex stimuli, in which the stimuli could be much larger than the receptive field, it is important to first "crop" the stimulus images, utilizing only the parts of the stimulus that encompass the receptive field (David et al., 2004), prior to further analysis. This procedure is necessary because visual stimulus pixels are expensive in terms of model estimation: using too many pixels can result in overparameterization (Bishop, 2006), i.e. estimating unnecessary parameters, leading to a loss in accuracy of the estimated model. Here we employ a three-pass cropping procedure. In the first pass, we estimate the model parameters using the full 480x480 stimulus images downsampled to 30x30. We then manually identify a square-shaped cropping window that encloses an area slightly larger than the apparent receptive field, and downsample the image within it to 30x30. This cropped image is then used to re-train the model in the second pass. In the third pass, we adjust the cropping window based on the model estimate obtained in the second pass to identify accurate boundaries of the receptive field. To handle very small receptive fields in which this size was a pixel or less, we imposed a minimum cropping window size of 120 pixels. However, the majority of A18 neurons and a subset of A17 neurons having large receptive fields required only the first or two passes.

The amount of downsampling employed after cropping was then determined by the choice of filter size - if the filter size is too low, the estimated filters are too small, noisy, and may appear clipped off at the edges. On the other hand, an overly large filter size can result in overparameterization with resultant loss of predictive performance. Preliminary analyzes indicated best results with the filter size between 10 to 15. Here we evaluated for each neuron, filter sizes of 11 and 15 pixels and chose the one giving filters which are minimally clipped the edges.

Bias terms ("dc offsets") are included where appropriate and early stopping with the regularization dataset was used to prevent overfitting (Bishop, 2006; Finnoff et al., 1993). We solve for all parameters in the model with gradient descent, using a loss function of the mean square error between the neuron's measured response and the model's estimated response for the training dataset. The gradient of the error is backpropagated in order to estimate each parameter in each layer (Rumelhart et al., 1986). Optimization of the convolutional model was implemented in Python using tools from the TensorFlow optimization package, version 2.0.0 (Abadi et al., 2016) and Keras, version 1.1.2 (Chollet et al, 2018) - see https://www.tensorflow.org/ and https://keras.io/. TensorFlow is used for its ability to automatically compute the gradient with respect to each parameter, including the α parameter of the PReLU and the Gaussian parameters of the map layer, as well as its built-in support for GPU processing. The optimization was performed using the Adam optimizer (Kingma & Ba, 2014), with a "patience" parameter of 50. The SciPy/NumPy Python packages are also extensively used (Jones et al., 2014). The weights for the filter are initialized randomly (Glorot & Bengio, 2010), with a taper applied at the edges to encourage non-zero weights towards the center of the filter array. The PReLU α is initialized at 0.5, and the Gaussian map layer is initialized to the center of the image, with standard deviation equal to the length of the image (after cropping and downsampling, i.e., 30 pixels). Code for a working example of the model estimation method can be found at our GitHub (https://github.com/JinaniSooriyaarachchi/CNNwithPReLU_RFestimateV1).

Model validation

The parameters of the model are optimized using the training dataset, with early stopping based on predictive ability for the regularization set. The test set is held out until all training procedures are completed, to give an unbiased estimate of the models' predictive accuracy. The model accuracy is measured by the variance accounted for (VAF), which is computed as the square of the correlation between the neuron's measured response and the response of the estimated model. This "raw" VAF (averaged over all 20 repetitions of the test set, and expressed as a percentage), however, does not take account of degradation by noisy responses of the neuron.

To measure the trial-by-trial variability of a neuron's response, we estimate a "noise ceiling" as follows. For each of the 20 repetitions of the validation set, the neuron's response on that repetition is used to predict the average response for the 19 other repetitions of the same stimulus, and a VAF (R^2) is calculated for this prediction. The average of these 20 VAFs provides an estimate of the noise ceiling, i.e., an upper bound on the explainable variance in the neuron's response, R^2_{neuron}. This value is similar in principle to the "reliability ratio" (Borst & Theunissen, 1999; Lesica et al., 2007).

The model's performance is computed similarly to R^2_{neuron}. For each of the 20 repetitions of the validation set, the neuron's response on that repetition is compared with the model's prediction of the neuron response, and a VAF (R^2) is calculated for this prediction. The average of all 20 of these VAF values is denoted as R^2_{model}. The ratio between R^2_{model} and R^2_{neuron} is an estimate of the proportion of "explainable" variance achieved by the model (Freeman et al., 2015). This value, again expressed as a percentage, is used to quantify a "noise-corrected" estimate of the model performance.

Neuron Inclusion Criteria

The estimated reliability value, R^2_{neuron}, was also used as a criterion for the inclusion of neurons for further analysis. R^2_{neuron} values were obtained for the natural image stimulus regularization set (20 repetitions), natural image stimulus test set (20 repetitions), and the grating stimuli (10 repetitions). These datasets were included in further analysis only if the R^2_{neuron} was greater

than 1%. In addition, we also require that the neurons' average firing rates in response to visual stimuli were at least 0.5 spikes/second greater than to the blank controls.

Simple-Complex Measure

For purposes of comparison with the estimated PReLU α values, the ratio of first harmonic to mean response to an optimized grating, or "AC/DC" value, has been widely used to classify a cortical neuron as a simple or complex type cell (Skottun et al., 1991). Neurons with an AC/DC less than unity were classified as complex, otherwise they were classified as simple. We calculated the AC/DC value for neurons on which sinewave grating responses were collected.

Analysis of spatial receptive field properties

In the convolutional analysis, both the filter and map layers contribute to the receptive field properties of a neuron. For example, when the filter has oriented, alternating excitatory and inhibitory regions and the map function is an isotropic Gaussian, the optimal orientation will be determined by the filter, but the overall size of the receptive field will depend jointly on both the filter and map functions. For archetypal simple type cells (alpha = 1.0) which can be modeled as a linear filter followed by a pointwise nonlinearity ("LN" model), the spatiotemporal receptive field can be "reconstructed" by taking the convolution of the filter function **c** with the map function **w**. The receptive fields of complex type cells cannot, in general, be characterized in this way, since their phase-invariance precludes an LN model. However, we note that the PReLU can be considered as a weighted sum of a linear function and a full-wave rectifier, which give linear and nonlinear responses, respectively, analogously to first and second Volterra kernels (Marmarelis & Marmarelis, 1978). For the linear part, the PReLU contributes only a scale factor, and again we can take the convolution of the filter function with the map function ("reconstruction") to provide a first-order indication of the receptive field properties, analogous to a first Volterra kernel. Here we will examine the reconstruction functions, which provide a consistent and uniform approach to characterizing properties of both simple and complex type cells, as well as those having intermediate properties, without having to treat them as categorically distinct.

We analyze spatial receptive field properties, by considering the reconstruction at the time lag having maximal variance (DeAngelis et al., 1993; Ringach, 2002). To characterize some of the conventional parameters of visual receptive fields, we find a best-fitting Gabor function (Eq 5, below) for each RF reconstruction. These functions fit most receptive fields well (e.g., Anzai et al., 1999; Jones & Palmer, 1987; Kulikowski & Bishop, 1981; Marĉelja, 1980), provide parameters of interest directly in the fitted function, and also have a rich theoretical underpinning (Marĉelja, 1980). We parameterize the fitted Gabor function as a sinusoid multiplied by a Gaussian envelope, with a change of variable to handle orientation:

$$\mathrm{h}(x', y') = A \exp\left(-\left(\frac{x'}{\sqrt{2}\sigma_{x'}}\right)^2 - \left(\frac{y'}{\sqrt{2}\sigma_{y'}}\right)^2\right) cos(2\pi f x' + \varphi) + d \qquad (5)$$

where

$$x' = (x - x_0)\, cos\theta + (y - y_0)\, sin\theta$$

$$y' = -(x - x_0)\, sin\theta + (y - y_0)\, cos\theta$$

and there are 9 free parameters, whose values are fit to a given spatial filter:

A is the overall amplitude or gain

f is the spatial frequency of the sinusoid

θ is the orientation of the sinusoid

φ is the phase (in radians) of the sinusoid

$\sigma_{x'}$ and $\sigma_{y'}$ are the standard deviations of the Gaussian along orthogonal axes, i.e.,
width and length respectively

x_0 and y_0 are the center location

d is an offset

To find values of the parameters that minimize the summed square error between a measured spatial filter and fitted Gabor, we employed a modified version of Kendrick Kay's *fitGabor2d* and its associated functions *(https://github.com/kendrickkay/knkutils*). This code uses Matlab's *lsqcurvefit* function which finds best fitted parameters in the least-square sense. We modified this code to avoid falling into local minima during fitting, by evaluating results from multiple

different initial values for the phase parameter, φ, to obtain the best fit.

Goodness of fit is indicated by the fraction of variance unexplained, FVU (Ringach, 2002):

$$FVU = 1 - R^2 \qquad\qquad (6)$$

where R^2 is the coefficient of determination. FVU ranges from 0 to 1.0, with decreasing values indicating better fits. As will be seen below, in majority of the cases the fitting algorithm converged, and gave good fitting functions that corresponded well with the spatial receptive fields.

Results

A total of 271 neurons were isolated from multielectrode recordings in 12 cats (some of which also contributed to recordings that were part of other concurrent studies). Of these, 206 neurons (127 from area 17 and 79 from area 18) exhibited sufficient response reliability (R^2_{neuron} > 1.0% for both the regularization and validation datasets) to be included. Neurons with an explainable VAF less than 1.0% were also excluded, to help ensure reliability of the estimated model parameters. The accepted 168 neurons (104 from area 17 and 64 from area 18) were used in further analysis as explained in the following sections.

Convolutional model predictive performance

The three-pass cropping procedure played an important role in characterizing a neuron's receptive field with high resolution and substantially increased predictive performance. Detailed results from an example neuron are shown in Figure 2. In the first pass, the estimated receptive field filter has poor resolution, the gaussian map layer is confined within a single pixel, and the linear reconstruction appears - such results are unsatisfactory for characterizing receptive field properties, because the lack of cropping leads to inappropriate downsampling. The estimated filter and map layer in the second pass are somewhat better with an acceptable linear reconstruction for the receptive field. However, estimated weights of a third pass yield a much better resolution of the structure of both the filter and map layers, and

a more satisfactory linear reconstruction (Figure 2A, 2B). The VAF accuracy on a hold-back test dataset for this neuron improved from 46.2% in the first pass, 58.1% in the second pass to 61.3% in the third pass. Figure 2C illustrates the overall improvement of VAF accuracy across three passes for a subset of area 17 neurons which had sufficiently small receptive fields to be mapped through all the three passes (n=83; mean VAFs: 15.1% in pass 1, 26.8% in pass 2 and 33.9% in pass 3). Consequently, the remainder of the paper will present results from the third pass.

Figure 2. Three pass procedure to estimate a better cropping window for an example cell from area 17. **A.** Estimated receptive field filters and linear reconstructions for three passes with cropping windows of 480x480 in first pass (VAF=46.2%), 240x240 in second pass (VAF=58.1%) and 120x120 in third pass (VAF=61.3%), all downsampled to 30x30. **B.** Map layer for three passes, estimated as a parameterized 2D Gaussian. **C.** Model VAF improvement for a subset of area 17 neurons with sufficiently small receptive fields (n = 83) across three passes. Distributions indicated as box and whisker plots, in which black triangles indicate mean values, whiskers indicate standard deviations, box bounds indicate 25th and 75th percentiles, and horizontal lines indicate medians.

An example of detailed results from a simple-type cell are shown in Figure 3. The estimated convolutional model performed well, with a VAF of 61.3%. Its estimated receptive field filter (Figure 3A) contained a central inhibitory region and surrounding excitation. The map layer (Figure 3C), parameterized as a Gaussian, localizes the activity of the filter. Note that the size of the filter layer (15x15) to produce the map layer (16x16 pixels, after the input image of 30x30 convolved is with a filter of 15x15) was hyperparameters whose values were optimized for this neuron - in general the size of the map layer depends on the size of the filter due to the convolution. The PReLU (Figure 3B) had an α of 0.8, corresponding to an approximately linear relationship. Since α is closer to 1.0 ($G(x) = x$) for this neuron we can simply neglect the PReLU and reconstruct the receptive field by convolving the filter with the map layer. Figure 3E shows the resulting linear receptive field reconstruction for this neuron. This reconstruction appears less noisy and has a more clearly delineated excitatory surround. In the convolutional model, the model architecture implicitly embodies an important constraint, i.e., that signal filtering properties of the neuron are determined by a relatively small filter, which is operative on only a limited subset (map layer, receptive field) of the input image. The Gaussian map layer applies a constraint on the locality of the filter activity, which reduces the noise outside of the neuron's receptive field. These factors together result in the linear reconstruction (Figure 3E) having smoother excitatory and inhibitory regions, on a background that is less noisy. The convolutional output from the map layer passes through the final output nonlinearity (exponent 1.2; Figure 3D) to generate the model's predicted neural response.

Figure 3. Visualization of models estimated for an example simple cell from area 17. **A.** Filter layer, 15x15 pixels. **B.** Intermediate nonlinearity (parameterized rectifier unit, PReLU) between filter and map layers, having a fixed positive unity gain for positive inputs (red) and an estimated gain parameter, α, for negative inputs (cyan); here, the estimated α = 0.8. **C.** Map layer, estimated as a parameterized 2D Gaussian; note different spatial scale. **D.** Final output nonlinearity (exponent 1.2). **E.** Linear filter reconstruction for the convolutional model (30x30 pixels), from convolution of filter and map layers. The convolutional method gave a raw VAF of 61.3%.

Results of the convolutional model parameters estimated for an example complex type cell are shown in Figure 4 with a raw VAF of 15.1%. The filter layer (Figure 4A) is noisy compared to the above example simple cell filter, but nevertheless shows a vertically oriented inhibitory zone in the third time lag. The PReLU (Figure 4B) has an α = -0.12, indicating a substantial nonlinearity and thus a large deviation from the linear-nonlinear model conventionally used for simple-type cells. The notions often used for complex cells are more appropriate here: the filter function can be considered a subunit, with multiple such (rectified) subunits contributing to the response of the neuron, as proposed by Hubel and Wiesel (1962). Considering the PReLU as a weighted sum of a linear proportionality and a full wave rectifier

corresponding to first and second order Volterra kernels (see Methods), the linear reconstruction of the complex cell (Figure 4E) provides a first order indication of the receptive field properties. The final output nonlinearity (exponent of 1.5) for this complex cell example is shown in Figure 4D.

Figure 4. Visualization of model estimated for an example complex cell from area 17. **A.** Filter layer, 15x15 pixels. **B.** PReLU nonlinearity, illustrated as for Figure 2B; here, estimated α = -0.12. **C.** Map layer **D.** final output nonlinearity of (exponent of 1.5) **E.** Linear filter reconstruction (30x30 pixels), from convolution of filter and map layers. The convolutional model gave a raw VAF of 15.1%.

Across the sample of neurons, there was a very wide variation in predictive performance (Figure 5A), with VAF values ranging from near zero to ca 70% in both area 17 and area 18 neurons. Previous system identification (Fournier et al., 2014) generally found much lower VAFs for complex than for simple type cells. Since our sample includes a mixture of both kinds of cells, it then seems likely that many or most of the neurons with lower VAFs in Figure 5A might be complex cells. Figure 5B compares the raw VAF of the convolutional method with a subset of neurons whose AC/DC values were obtained for optimal gratings. The raw VAFs for simple cells

(AC/DC >= 1.0, n = 21) were almost always better than those for complex cells (AC/DC < 1.0, n = 8). The simple cells had an average raw VAF of 44.3% whereas the complex cells had an average raw VAF of 21.9% (p = 0.0047, significant by two-sample t-test).

Figure 5. Predictive accuracy for a holdback test dataset. **A**, Percentage of variance accounted for (VAF) in neurons' responses for Area 17 (mean VAF=30.2%, n=104) and area 18 neurons (mean VAF=36.5%, n=64), respectively. Distributions indicated as box and whisker plots, in which black triangles indicate mean values, whiskers indicate standard deviations, box bounds indicate 25th and 75th percentiles, and horizontal lines indicate medians. **B**, Comparison between the model prediction accuracy (raw VAF) and the neuron's phase sensitivity (AC/DC) to sinewave gratings (n=29).

Prediction of grating responses

We additionally assessed the convolutional model by measuring how well it predicts neurons' responses to grating stimuli, for those neurons on which both kinds of responses were available. These models were estimated from the responses to the natural image stimuli, then simulations of the models were used to predict their responses to grating stimuli. The quality of prediction is assessed by the VAF between a predicted spatial frequency tuning curve and the neuron's measured tuning curve.

An example of such a comparison for spatial frequency tuning is shown in Figure 6A for an example area 17 simple cell. Here the neuron's measured response (black) and the convolutional model prediction (red) are quite similar (VAF = 91%), with clear bandpass tuning

to about 1.0 cpd. Results for spatial frequency tuning prediction in 39 neurons (21 from area 17, 18 from area 18) are shown in Figure 6B. Considering the neurons in both brain areas together, the convolutional model performed with an average VAF of 58.2%. Considering only the area 18 neurons, the average VAF the convolutional model is 51.3%, and for the area 17 neurons, the average VAF for the convolutional model is 64.1%. In summary, the convolutional model is reliable in predicting spatial frequency tuning curves for both area 17 and area 18 neurons.

Figure 6. Prediction of grating tuning curves by estimated models. **A**, Spatial frequency tuning curve (average firing rate, normalized to maximum) for an example A17 neuron. Temporal frequency, 2 Hz; average of 10 repetitions of 1 sec each. Solid black line for neuron's response to the grating stimuli, red for tuning curve predicted by convolutional model (VAF = 91%). **B**, Model performance (VAFs) for spatial frequency tuning predictions (n = 18 for area 18, n = 21 for area 17). Distributions indicated as box and whisker plots, in which black triangles indicate mean values, whiskers indicate standard deviations, box bounds indicate 25th and 75th percentiles, and horizontal lines indicate medians.

Noise ceiling analysis and simple vs. complex cells

Lower VAFs for complex cells could be expected since inadequacies of the associated model architecture might be greater than for simple cells. However, another possibility is that responses of simple cells might be less noisy, i.e. exhibit less trial to trial variability, than for complex cells. We examined this idea using neurons in our sample for which we had responses to gratings as well as natural images. Figure 7A shows raster plots of neuronal firing across repeated trials in response to a subset of the natural image ensembles, for example neurons identified as simple or complex type cells based on their AC/DC ratios for optimized gratings. It can be seen that the raster points for the example simple cell (top) are relatively well aligned

vertically, indicating fairly reliable responses, while the complex cell (bottom) gives much more variability in spike firing times across repetitions of the same stimulus ensemble. We quantified the trial-wise reliability by measuring how well the response on a given trial was correlated with the average response across all the repeated trials, using the R^2_{neuron} index (see Methods) - higher values of R^2_{neuron} indicate greater reliability. For the example neurons in Figure 7A, R^2_{neuron} was 59.4% for the simple cell and 3.1% for the complex cell.

Figure 7. Trial-by-trial reliability of neuronal responses for simple and complex type cells. *A*, Spike raster plots for responses of example simple and complex cells (same neurons as in Figures 2 and 3, respectively) to 20 repetitions of a natural image ensemble (only first 2 sec shown). Top traces are post stimulus time histograms (bins = 6.67 ms) for the neurons' spiking responses (black) and red

(convolutional). ***B***, Neurons' reliability index, R^2_{neuron}, for natural image responses as a function of grating phase sensitivity, AC/DC (n = 29). Larger values of R^2_{neuron} correspond to greater reliability. ***C***, R^2_{neuron} calculated from responses to each neuron's preferred grating (n = 100). ***D***, Population averages of response reliability, R^2_{neuron}, of simple and complex type cells to gratings and to natural images. Conventions for box and whisker plots are same as in Figure *5A.*

We similarly measured R^2_{neuron} values from natural image responses for all of the sampled neurons for which responses to gratings were also available. Neurons with R^2_{neuron} less than 1.0% on either the natural stimuli datasets (regularization and validation), or less than 1.0% on the preferred grating stimuli (see below), were excluded. Results for 29 neurons (20 from area 17, 9 from area 18), shown in Figure 7B, reveal generally greater reliability in the simple cells (AC/DC >= 1.0) than in the complex cells (AC/DC < 1.0). For neurons classified as simple, the average R^2_{neuron} (34.59%, n = 21) was significantly greater (t-test, p = 0.0105) than for neurons classified as complex (average R^2_{neuron} = 14.56%, n = 8) - this difference is illustrated in the bar and whisker plots on the left side of Figure 7D.

Conceivably this difference in reliability between simple and complex cells might be peculiar to their responses to natural images. To evaluate this idea, we performed the same analysis on responses to drifting sinewave gratings across repeated trials. Since this analysis required only grating responses, we were able to utilize a much larger sample of 100 neurons (73 from area 17, 27 from area 18). Cells with R^2_{neuron} less than 1.0%, were removed to ensure valid AC/DC measurements. Figure 7C shows these neurons' R^2_{neuron} values for responses to their optimized gratings, as a functions of each neuron's AC/DC ratio. Again, it is clear that many of the simple cells show higher R^2_{neuron} values than complex cells. For neurons classified as simple, the average R^2_{neuron} was 7.6% (n = 54). For neurons classified as complex, the average R^2_{neuron} was 3.3% (n = 46). The distribution for simple cells was significantly greater (t-test, p = 1.77078E-10), again showing that complex cells were noisier and less reliable.

Thus the poorer predictive performance for complex cells (Figure 4) might be, at least in part, due to their greater trial-wise variability, instead of (or in addition to) an inadequacy of model architecture.

Relationship between PReLU α and simple/complex type responses

In many situations, suitable grating data for obtaining AC/DC values for individual neurons may not be available. As explained earlier, the PReLU α parameter of the convolutional model distinguishes between a linear-nonlinear model and a hierarchical (subunit summation) model, which are generally thought to reasonably represent simple and complex cells, respectively. Thus, the PReLU α value of the convolutional model itself, as estimated from responses to broadband stimuli such as natural images, could provide a surrogate for the AC/DC value.

In order to examine the interpretability of the PReLU α, Figure 8A shows a comparison between the convolutional model's PReLU α obtained from natural image responses, and AC/DC values measured with grating stimuli for 29 neurons (20 from area 17, 9 from area 18). Neurons with an explainable VAF less than 1.0% were excluded, to help ensure reliability of the estimated model parameters. As expected, simple type cells (AC/DC > 1.0) indeed had higher PReLU α values, typically near unity, while complex type cells usually had values less than unity, sometimes less than zero. While the relationship was not strictly linear, the PReLU α and AC/DC were highly correlated, with a Pearson's correlation coefficient of r = 0.84 (p = 1.12e-08, n =29). In terms of ranked values, these measures were also highly related, with a Spearman's correlation coefficient of r = 0.82 (p = 6.0816e-08, n =29). Thus, the PReLU α estimated from natural image responses could be used to distinguish simple vs. complex type cells with reasonable (if not perfect) reliability.

Since complex cells have a higher trial-to-trial response variability compared to simple cells, we can expect lower raw VAFs for complex cells with lower PReLU α values compared to simple cells with higher PReLU α values. Figure 8B shows this for the population of neurons, with lower predictive performance in terms of raw VAF for complex cells and higher predictive performance for simple cells. Across the population of neurons, we observed a significant correlation coefficient of r=0.58 between raw VAF and the estimated PReLU α values (p= 3.49e-16, n=168). In terms of ranked values, these measures were also highly related, with a Spearman's correlation coefficient of r = 0.59 (p = 0.0, n =168).

Figure 8. Relationship between model prediction accuracy, simple-vs-complex type responses, and model PReLU. **A**, Comparison between the convolutional model's PReLU α parameter and neurons' phase sensitivity (AC/DC) (n=29). **B**, Comparison between the model prediction accuracy and the convolutional model's PReLU α parameter (n=168). **C**, Histogram of alpha values for A17 neurons (n=104) **D**, Histogram of alpha values for A18 neurons (n=64).

The distributions of estimated alpha values for the full sample of neurons are shown as histograms in Figure 8C, D. The A17 sample (Figure 8C) appears to have a range of alpha values, with many neurons having alpha values close to unity, indicative of simple-type response properties, and others having values below zero, corresponding to complex-type responses. The distribution for the A18 cells (Figure 8D) shows a similar range of alpha values. However, the distributions of both A17 and A18 cells do not indicate statistically significant bimodality by Hartigan's dip test (A17: p=0.45, n=104, dip at 0.03; A18: p=0.83, n=64, dip at 0.03).

Spatial receptive field properties from the convolutional model

So far, we have emphasized the ability of the fitted CNN models to predict neurons' responses to novel visual stimuli which were not used for training - this is important for establishing the predictive power of the fitted models for a wide variety of visual stimuli. However, an important utility of system identification is also to gain insight into each neuron's selectivity for particular properties of visual stimuli, and how these preferences vary from one neuron to another, and from one brain area to another.

As an illustration of how the CNN results can be used to characterize receptive fields, we fit Gabor functions to the reconstructed filter functions at the optimal temporal lag for each neuron (see Methods). To help avoid spurious function fits to noisy results, we included in this analysis only those neurons whose fitted CNN models predicted the test datasets with VAF values greater than 10% (94 out of 104 A17 cells, 57 out of 64 A18 cells). Two examples of such Gabor-fits are seen in Figure 9A, in each case showing the CNN reconstructed filter (left), best-fitting Gabor (middle), and residual (difference between them, right). A typical A17 cell with alternating, elongated excitatory and inhibitory regions is shown in the top row of Figure 9A - this was a complex-type cell (alpha = -0.12), giving a VAF = 15.1% and a very good Gabor function fit (fraction of unexplained variance, FVU = 0.04). The bottom row illustrates a simple-type cell (alpha = 0.8) with a relatively non-oriented, center-surround spatial receptive field, giving a VAF = 61.3% and a very good fit by the Gabor function (FVU = 0.09).

Figure 9. Fitting of 2-D Gabor functions to receptive field reconstructions. **A**, 2-D Gabor fits to estimated spatial receptive field (convolution of filter with map layer) at optimal time lags, for two example neurons. Images in left column show the estimated receptive fields, with best-fitting Gabors in the middle column, and the residuals in the right column. **B**, Histogram of FVU (fraction of variance unaccounted) values, for A17 sample. **C**, scatterplot of FVU vs alpha values for A17 sample, with symbol for each neuron colored to indicate VAF for model's prediction of test dataset. **D**, as in **B**, but for A18 sample. **E**, as in **C**, but for A18 sample.

In agreement with previous studies (e.g, (Jones & Palmer, 1987; Ringach, 2002), most A17 receptive fields were fairly well fit by a Gabor function, as seen in the histogram of FVU values (Figure 9B) - the FVU values were small, ranging from 0.0148 to 0.445 (median = 0.0732). Such good fits were in spite of some neurons having receptive fields with very little orientation selectivity (Ringach, 2002) or having "crescent" shaped flanking regions (Conway & Livingstone, 2006). These relatively good fits were found across the full range of PReLU α values (Figure 9C), showing that Gabor functions are also appropriate descriptions for the selectivity of complex as well as simple type cells. FVU was not significantly correlated with alpha in the A17 cells (r = -0.245, p = 0.018, n=94). For A18 neurons, the FVU histogram in Figure 9D shows that Gabor functions gave similarly good fits in most cases, with FVU values from 0.0238 to 0.4641 (median=0.0906). We excluded from further analysis only two A18 neurons for giving a particularly poor fit (FVU > 0.5). For the remaining A18 neurons, the FVU was not significantly correlated with alpha, as seen in Figure 9E (r = -0.374, p = 0.048, n=55).

The simplest receptive field parameter to examine is its size. Here we estimate RF size from the envelope of the best-fitting Gabor, as $(\sigma_{x'}^2 + \sigma_{y'}^2)^{0.5}$, where $\sigma_{x'}$ and $\sigma_{y'}$ are the envelope sigma parameters along the length and width, respectively. The estimated RF sizes for A17 neurons, illustrated in the histogram of Figure 10A, ranged from 0.5381 to 10.0469 deg (median 2.3785 deg) - these values largely agree with previous results from cat A17 (e.g., (Baker, 1988; Sasaki & Ohzawa, 2007). A scatterplot of these RF sizes against the alpha values for the A17 sample is shown in Figure 10B, showing that simple and complex type cells had highly overlapping ranges of RF sizes, but with substantially larger values in a minority of complex cells. As a result, resultant correlation of RF size with alpha was statistically significant (r = -0.3576, p = 4e-04, n=94). Area 18 neurons had a wide range of RF sizes (1.4037 to 13.0555 deg, median = 5.1752 deg), as seen in the histogram of Figure 10C. The ranges of sizes for lower vs. higher values of PReLU α were highly overlapping (Figure 10D), resulting in a non-significant correlation (r =-0.334, p = 0.012, n=55). Comparison of the histograms in Figure 10A and 10C makes it clear that RF sizes tended to be larger in A18 than A17, confirmed statistically with an unequal variance (Welch) t test (p= 9.4e-8).

Figure 10. Receptive field sizes estimated for simple and complex-type A17/18 cells, from envelopes of fitted Gabors. **A**. Histogram of estimated receptive field sizes for A17 cells (N=94). **B**, Relationship of A17 receptive field sizes to PReLU α values. Each plotted point represents one neuron, with colour indicating VAF. **C, D**, same as *A, B* for A18 cells (N=55).

A characteristic spatial property of early visual cortex neurons is their selectivity for spatial frequency. In our analysis a neuron's optimal spatial frequency would be captured explicitly by the frequency parameter (*f*) of the best-fitting Gabor. Most neurons in A17 had optimal values ranging from 0.0049 to 0.689 cpd (Figure 11A), with a median of 0.1871cpd (n=94). This range is consistent with previous reported measurements (e.g., Ferster & Jagadeesh, 1991; Sasaki & Ohzawa, 2007; Zhou & Baker, 1996). Optimal spatial frequencies for lower vs. higher values of PReLU α (Figure 11B) overlapped greatly, with more complex-like cells apparently extending to lower frequencies - this was confirmed by a non-significant correlation with alpha (r=0.2003, p= 0.0530, n=94). Most neurons in A18 had estimated optimal

spatial frequencies between 0.0088 and 0.2896 cpd (median 0.0646 cpd), as seen in the histogram of Figure 11C. Neurons with smaller, more complex-like PReLU α values and neurons with larger alpha values (simple-like cells) spanned a larger but overlapping range of frequencies (Figure 11D). This relationship did not show a significant correlation (r = 0.1735, p = 0.2051, n=55). Comparison of the distributions for A17 vs A18 (Figure 11A, C) indicates a lower range of optimal spatial frequencies in A18 than those in A17, consistent with previous findings for single units (Ferster & Jagadeesh, 1991; Movshon et al., 1978; Zhou & C. L. Baker, 1996) as well as optical imaging (Issa et al., 2000; Zhan & Baker, 2008) in the cat.

Figure 11. Optimal spatial frequencies estimated for simple and complex-type A17/18 cells, from envelopes of fitted Gabors. **A**. Histogram of estimated optimal spatial frequencies for A17 cells (N=87). **B**, Relationship of A17 optimal spatial frequencies to alpha values (Pearson's r=0.200, p= 0.0530). Each plotted point represents one neuron, with colour indicating VAF. **C, D**, same as *A, B* for A18 cells (N=49). (r = 0.173, p = 0.205).

In general, the above results broadly agree with previous published reports, serving as a further validation of the methodology. In addition, these results extend earlier findings with

laboratory stimuli (white noise, gratings) to responses to natural images, for both simple and complex cells in cat A17 and A18.

Discussion

We have shown how a simple convolutional model can estimate receptive fields of both simple and complex type visual cortex neurons from responses to natural images and predict responses to holdback natural image ensembles as well as responses to sinewave gratings. We also demonstrate that the generally poorer predictive performance for complex cells is due in large part to a lower trial-wise reliability of responses from these neurons.

The model's architectural simplicity (Figure 1) allows for a straightforward interpretation of its parameters. The filter layer can be thought of as a spatially delocalized simple cell, which primarily determines the neuron's selectivity for stimulus properties such as orientation and spatial frequency. The map layer indicates the region of space where the filter is active, i.e., the overall location of the receptive field. Most importantly, we find that a single parameter, the PReLU, can capture a neuron's phase sensitivity. Not only does the PReLU α correlate well with the neuron's AC/DC value measured from grating responses, this parameter is related to well-known models for early visual cortical neurons. The PReLU α can be considered a rough representation of the balance between the the filter and it's opposite-polarity counterpart. When the two are balanced (α near 1.0), the PReLU approximates a linear relationship, so simple cell-like responses are generated.

As the PReLU α deviates from unity, the neurons' responses will begin to deviate from linearity. In this case, the model is more similar to the idea (Adelson & Bergen, 1985; Hubel & Wiesel, 1962) that complex cell responses arise from the integration of multiple simple cells that act as nonlinear subunits. With complex cells, the interposed nonlinearity enables the possibility for system identification to retrieve the subunit filters. However, with simple cells, due to the linearity, subunit filters cannot be explicitly recovered, since two linear filters in a cascade are equivalent to a single composite linear filter. The case where the PReLU α is near

zero corresponds to Hubel and Wiesel's complex cell model, i.e., a linear summation of simple cells, all having the same filter properties but different positional offsets. Consider the filter layer, w, and a window of the stimulus, s_1. If the PReLU is a half-wave rectifier, $N(x)$, then $N(w \cdot s_1)$ acts like a linear-nonlinear model of a simple cell. The map layer indicates the region of visual space where the simple cell-like filter is active. If the PReLU α is negative, the model becomes somewhat more similar to an energy model (Adelson & Bergen, 1985) in which an opponent filter also contributes to the neuron's response.

However, this does not necessarily imply that the convolutional model corresponds to the underlying neural circuitry of visual cortex neurons - there are many different connectivity models that can generate simple and complex cell responses. Rather, this convolutional model is a simple and reliable method of estimating the properties of early visual cortex cells. While the PReLU may show a simple relationship between simple and complex cells, its utility in the model does not provide concrete evidence for a particular underlying neural connectivity.

Although the convolutional model can perform well in predicting both simple and complex cell response properties, it does not provide a complete description of all early visual neurons. The intermediate nonlinearity is likely to be more complex than the PReLU, e.g. a polynomial function such as a square law, as in the (Adelson & Bergen, 1985) energy model. However, such more elaborate functions generally increase the difficulty of the model parameter estimation. For example, the tent basis as used by Vintch et al. (2015) could approximate a square law, however the required estimation procedure becomes more complicated. The simplicity of the PReLU allows for a smooth gradient calculation, and only adds one parameter to the model, which can be trained jointly with the other parameters. Another advantage of the PReLU is that its single parameter is simply related to the simple vs. complex cell distinction. Our method also does not require any additional initialization procedure; the weights can be initialized randomly (filter) or at a fixed value (PReLU/Gaussian).

A slight complication in estimating the convolutional model is the need for correct hyperparameters (filter size and amount of downsampling). Fortunately, nearby neurons in the visual cortex typically have fairly similar sizes and/or preferred spatial frequencies (DeAngelis et

al., 1999; Issa et al., 2000; Tootell et al., 1981), so the model filters of neighboring neurons are likely to be similar in size. Thus, a hyperparameter search may not be necessary for every neuron. However in some cases, a limited hyperparameter search may still be valuable, though the model can still perform fairly well with sub-optimal hyperparameters.

It should be noted that despite the advantageous aspects of the convolutional model, the raw VAFs obtained for complex cells were lower than those for simple cells. However, these raw VAFs are in a similar range as those reported for other recently proposed estimation procedures (Fournier et al., 2014; Vintch et al., 2015). We have shown that the complex cells are more noisy than simple cells in their trial-by-trial reliability. Using VAFs adjusted for this unreliability, we found that the relative difference between complex and simple cells was lowered, though not eliminated. This suggests that there is scope for improvement of our model architecture, to more fully explain complex cell responses.

Vintch et al. (2015) proposed a convolutional model for V1 receptive fields, having two filters capable of characterizing both simple and complex cells' responses to sparse white noise. However, estimation of the model parameters apparently required carefully chosen initial conditions and iterative back-and-forth estimation of different parts of the model; in addition, the model parameters are not straightforwardly related to the simple/complex distinction. Such a model would probably provide somewhat better predictive performance than the one described here, but it is more difficult to estimate. A future possibility might be to use the approach described here to obtain good initial conditions for a more complicated model such as that of Vintch et al. (2015).

An important advantage of the system identification approach used here is that it does not make an *apriori* assumption of a functional filter form, e.g., 2-D Gabor function, but rather it estimates all points in the filter from the neuron's responses. Consequently, this allows the method to reveal other kinds of receptive field shapes, for example those that are non-oriented, or even "crescent" shapes (Conway & Livingstone, 2006). Then subsequently, we can examine how well candidate functions can serve as descriptions of the receptive field (e.g., Figure 9).

Since our convolutional model has only a single filter, it cannot capture properties such as cross-orientation inhibition (Morrone et al., 1982), which requires at least two filters. Other properties such as surround suppression (Jones et al., 2001; Nurminen & Angelucci, 2014; Sengpiel et al., 1997), contrast gain control (Ohzawa et al., 1982), or second-order responses (Li et al., 2014), also would require more complex model architectures. However with improvements such as greater amounts of data, or additional methods of regularization (Sietsma & Dow, 1991; Srivastava et al., 2014), it may become possible to resolve greater model structure. Because the model is constructed within the framework of deep neural networks, with widely used tools for machine learning, further improvements to it can benefit from on-going rapid advances in software algorithms and GPU parallel processing support for this kind of learning algorithm.

Neurons in higher-tier visual areas require models with more filters and multiple layers (Oliver & Gallant, 2016; Yamins et al., 2014). For example, neurons in higher stages of visual processing become more selective for particular types of stimuli, but less positionally selective within larger receptive fields (Chelazzi et al., 1993; Desimone & Schein, 1987). A well-known example are neurons selective for faces, regardless of position or rotation (Desimone, 1991). Models for these higher regions generally involve a hierarchy of processing stages (layers) beyond early-stage simple and complex cells, to build more complex and object-specific responses (Cadieu et al., 2007; Oliver & Gallant, 2016). In a convolutional neural network model, this is achieved by adding more convolutional layers. Interestingly, it has been found that the activity of intermediate-layer neurons in large multilayered convolutional neural networks have similarities to ventral stream cell responses (Yamins et al., 2014).

Our convolutional model architecture has relatively few additional parameters beyond those of the linear-nonlinear model, making it straightforward to estimate. The PReLU imposes negligible computational cost - it has a single additional parameter that can be estimated within the gradient descent framework. Another important advantage of this simplicity is that the estimation is relatively robust with choice of initial conditions - therefore pre-training methods are unnecessary. The estimation of the convolutional model is sufficiently fast that it could conceivably be employed on-line during recording experiments, to provide a rapid estimate of

neuronal filtering and receptive field locations. This could be particularly valuable in the context of multi-neuron recording with multielectrode or 2-photon imaging, when different neurons may have quite different stimulus selectivities and receptive field locations.

References:

Abadi, M., Agarwal, A., Barham, P., Brevdo, E., Chen, Z., Citro, C., Corrado, G. S., Davis, A., Dean, J., Devin, M., Ghemawat, S., Goodfellow, I., Harp, A., Irving, G., Isard, M., Jia, Y., Jozefowicz, R., Kaiser, L., Kudlur, M., . . . Zheng, X. (2016). TensorFlow: Large-Scale Machine Learning on Heterogeneous Distributed Systems. arXiv:1603.04467. Retrieved March 01, 2016, from https://ui.adsabs.harvard.edu/abs/2016arXiv160304467A

Adelson, E. H., & Bergen, J. R. (1985). Spatiotemporal energy models for the perception of motion. *Journal of the Optical Society of America A*, *2*(2), 284-299. https://doi.org/10.1364/JOSAA.2.000284

Anzai, A., Ohzawa, I., & Freeman, R. D. (1999). Neural mechanisms for processing binocular information I. Simple cells. *J Neurophysiol*, *82*(2), 891-908. https://doi.org/10.1152/jn.1999.82.2.891

Baker, C. L., Jr. (1988). Spatial and temporal determinants of directionally selective velocity preference in cat striate cortex neurons. *J Neurophysiol*, *59*(5), 1557-1574. https://doi.org/10.1152/jn.1988.59.5.1557

Bishop, C. M. (2006). *Pattern Recognition and Machine Learning*. Springer. https://books.google.ca/books?id=kTNoQgAACAAJ

Blanche, T. J., Spacek, M. A., Hetke, J. F., & Swindale, N. V. (2005). Polytrodes: high-density silicon electrode arrays for large-scale multiunit recording. *J Neurophysiol*, *93*(5), 2987-3000. https://doi.org/10.1152/jn.01023.2004

Borst, A., & Theunissen, F. E. (1999). Information theory and neural coding. *Nature Neuroscience*, *2*(11), 947-957.

Brainard, D. H. (1997). The Psychophysics Toolbox. *Spatial Vision*, *10*, 433-436. https://doi.org/10.1163/156856897X00357

Cadena, S.A., Denfield, G.H., Walker, E.Y., Gatys, L.A., Tolias, A.S., Bethge, M., Echer, A.S. (2017) Deep convolutional models improve predictions of macaque V1 responses to natural images. PLOS Comp Biol 15(4):e1006897.

Cadieu, C., Kouh, M., Pasupathy, A., Connor, C. E., Riesenhuber, M., & Poggio, T. (2007). A model of V4 shape selectivity and invariance. *Journal of Neurophysiology*, *98*, 1733-1750. https://doi.org/10.1152/jn.01265.2006

Carandini, M. (2004). Amplification of Trial-to-Trial Response Variability by Neurons in Visual Cortex. *PLOS Biology*, *2*(9), e264. https://doi.org/10.1371/journal.pbio.0020264

Chelazzi, L., Miller, E. K., Duncan, J., & Desimone, R. (1993). A neural basis for visual search in inferior temporal cortex. *Nature*, *363*(6427), 345-347. https://doi.org/10.1038/363345a0

Chollet, F., & others. (2018). *Keras: The Python Deep Learning library*. https://ui.adsabs.harvard.edu/abs/2018ascl.soft06022C

Conway, B. R., & Livingstone, M. S. (2006). Spatial and Temporal Properties of Cone Signals in Alert Macaque Primary Visual Cortex. *The Journal of Neuroscience*, *26*(42), 10826-10846. https://doi.org/10.1523/jneurosci.2091-06.2006

David, S. V., Vinje, W. E., & Gallant, J. L. (2004). Natural Stimulus Statistics Alter the Receptive Field Structure of V1 Neurons. *The Journal of Neuroscience*, *24*(31), 6991-7006. https://doi.org/10.1523/jneurosci.1422-04.2004

De Boor, C. (1978). *A practical guide to splines* (Vol. 27). Springer-Verlag New York.

DeAngelis, G. C., Ghose, G. M., Ohzawa, I., & Freeman, R. D. (1999). Functional micro-organization of primary visual cortex: receptive field analysis of nearby neurons. *Journal of Neuroscience*, *19*(10), 4046-4064.

DeAngelis, G. C., Ohzawa, I., & Freeman, R. (1993). Spatiotemporal organization of simple-cell receptive fields in the cat's striate cortex. I. General characteristics and postnatal development. *Journal of Neurophysiology*, *69*(4), 1091-1117.

DeAngelis, G. C., Ohzawa, I., & Freeman, R. D. (1991). Depth is encoded in the visual cortex by a specialized receptive field structure. *Nature*, *352*(6331), 156-159.

Desimone, R. (1991). Face-selective cells in the temporal cortex of monkeys. *J Cogn Neurosci*, *3*(1), 1-8. https://doi.org/10.1162/jocn.1991.3.1.1

Desimone, R., & Schein, S. J. (1987). Visual properties of neurons in area V4 of the macaque: sensitivity to stimulus form. *J Neurophysiol*, *57*(3), 835-868. https://doi.org/10.1152/jn.1987.57.3.835

DiMattina, C., & Zhang, K. (2011). Active Data Collection for Efficient Estimation and Comparison of Nonlinear Neural Models. *Neural Computation*, *23*(9), 2242-2288. https://doi.org/10.1162/NECO_a_00167

Ecker, A.S., Sinz, F.H., Froudarakis, E., Fahey, P.G., Cadena, S.A., Walker, E.Y., Cobos, E., Reimer, J., Tolias, A.S., Bethge, M. (2018) A rotation-equivalent convolutional neural network model of primary visual cortex. arXiv: 1809.10504

Ferster, D., & Jagadeesh, B. (1991). Nonlinearity of spatial summation in simple cells of areas 17 and 18 of cat visual cortex. *J Neurophysiol*, *66*(5), 1667-1679. https://doi.org/10.1152/jn.1991.66.5.1667

Finnoff, W., Hergert, F., & Zimmermann, H. G. (1993). Improving model selection by nonconvergent methods. *Neural Networks*, *6*(6), 771-783. https://doi.org/https://doi.org/10.1016/S0893-6080(05)80122-4

Fournier, J., Monier, C., Levy, M., Marre, O., Sári, K., Kisvárday, Z. F., & Frégnac, Y. (2014). Hidden complexity of synaptic receptive fields in cat v1. *The Journal of Neuroscience*, *34*(16), 5515-5528.

Freeman, J., Field, G. D., Li, P. H., Greschner, M., Gunning, D. E., Mathieson, K., Sher, A., Litke, A. M., Paninski, L., & Simoncelli, E. P. (2015). Mapping nonlinear receptive field structure in primate retina at single cone resolution. *Elife*, *4*, e05241.

Glorot, X., & Bengio, Y. (2010). Understanding the difficulty of training deep feedforward neural networks. Aistats,

He, K., Zhang, X., Ren, S., & Sun, J. (2015). Delving deep into rectifiers: Surpassing human-level performance on imagenet classification. Proceedings of the IEEE International Conference on Computer Vision,

Hubel, D. H., & Wiesel, T. N. (1962). Receptive fields, binocular interaction and functional architecture in the cat's visual cortex. *The Journal of Physiology*, *160*(1), 106-154.

Hubel, D. H., & Wiesel, T. N. (1968). Receptive fields and functional architecture of monkey striate cortex. *The Journal of Physiology*, *195*(1), 215-243.

Ikezoe, K., Amano, M., Nishimoto, S., & Fujita, I. (2018). Mapping stimulus feature selectivity in macaque V1 by two-photon Ca2+ imaging: Encoding-model analysis of fluorescence

responses to natural movies. *NeuroImage*, *180*, 312-323.
https://doi.org/https://doi.org/10.1016/j.neuroimage.2018.01.009

Issa, N. P., Trepel, C., & Stryker, M. P. (2000). Spatial frequency maps in cat visual cortex. *The Journal of Neuroscience*, *20*, 8504-8514.

Jones, E., Oliphant, T., & Peterson, P. (2014). {SciPy}: open source scientific tools for {Python}.

Jones, H. E., Grieve, K. L., Wang, W., & Sillito, A. M. (2001). Surround suppression in primate V1. *J Neurophysiol*, *86*(4), 2011-2028. https://doi.org/10.1152/jn.2001.86.4.2011

Jones, J. P., & Palmer, L. A. (1987). An evaluation of the two-dimensional Gabor filter model of simple receptive fields in cat striate cortex. *J Neurophysiol*, *58*(6), 1233-1258. https://doi.org/10.1152/jn.1987.58.6.1233

Jun, J. J., Steinmetz, N. A., Siegle, J. H., Denman, D. J., Bauza, M., Barbarits, B., Lee, A. K., Anastassiou, C. A., Andrei, A., Aydın, Ç., Barbic, M., Blanche, T. J., Bonin, V., Couto, J., Dutta, B., Gratiy, S. L., Gutnisky, D. A., Häusser, M., Karsh, B., . . . Harris, T. D. (2017). Fully integrated silicon probes for high-density recording of neural activity. *Nature*, *551*(7679), 232-236. https://doi.org/10.1038/nature24636

Kindel, W.F., Christensen, E.D., Zylberberg, J. (2019) Using deep learning to probe the neural code for images in primary visual cortex. J. Vision 19(4):29,1-12.

Kingma, D. P., & Ba, J. (2014). Adam: A method for stochastic optimization. *arXiv preprint arXiv:1412.6980*.

Krizhevsky, A., Sutskever, I., & Hinton, G. E. (2012). Imagenet classification with deep convolutional neural networks. Advances in neural information processing systems,

Kulikowski, J. J., & Bishop, P. O. (1981). Linear analysis of the responses of simple cells in the cat visual cortex. *Experimental Brain Research*, *44*(4), 386-400. https://doi.org/10.1007/BF00238831

Lesica, N. A., Jin, J., Weng, C., Yeh, C.-I., Butts, D. A., Stanley, G. B., & Alonso, J.-M. (2007). Adaptation to stimulus contrast and correlations during natural visual stimulation. *Neuron*, *55*(3), 479-491.

Li, G., Yao, Z., Wang, Z., Yuan, N., Talebi, V., Tan, J., Wang, Y., Zhou, Y., & Baker, C. L. (2014). Form-Cue Invariant Second-Order Neuronal Responses to Contrast Modulation in Primate Area V2. *The Journal of Neuroscience*, *34*(36), 12081-12092. https://doi.org/10.1523/jneurosci.0211-14.2014

Lochmann, T., Blanche, T. J., & Butts, D. A. (2013). Construction of Direction Selectivity through Local Energy Computations in Primary Visual Cortex. *PLOS ONE*, *8*(3), e58666. https://doi.org/10.1371/journal.pone.0058666

Marĉelja, S. (1980). Mathematical description of the responses of simple cortical cells*. *Journal of the Optical Society of America*, *70*(11), 1297-1300. https://doi.org/10.1364/JOSA.70.001297

Marmarelis, P. Z., & Marmarelis, V. Z. (1978). Analysis of Physiological Signals. In *Analysis of Physiological Systems: The White-Noise Approach* (pp. 11-69). Springer US. https://doi.org/10.1007/978-1-4613-3970-0_2

McLean, J., Raab, S., & Palmer, L. A. (1994). Contribution of linear mechanisms to the specification of local motion by simple cells in areas 17 and 18 of the cat. *Vis Neurosci*, *11*(2), 271-294. https://doi.org/10.1017/s0952523800001632

Mechler, F., & Ringach, D. L. (2002). On the classification of simple and complex cells. *Vision Research*, *42*(8), 1017-1033.

Morrone, M. C., Burr, D. C., & Maffei, L. (1982). Functional implications of cross-orientation inhibition of cortical visual cells. I. Neurophysiological evidence. *Proc R Soc Lond B Biol Sci*, *216*(1204), 335-354. https://doi.org/10.1098/rspb.1982.0078

Movshon, J. A., Thompson, I. D., & Tolhurst, D. J. (1978). Spatial and temporal contrast sensitivity of neurones in areas 17 and 18 of the cat's visual cortex. *J Physiol*, *283*, 101-120. https://doi.org/10.1113/jphysiol.1978.sp012490

Nurminen, L., & Angelucci, A. (2014). Multiple components of surround modulation in primary visual cortex: Multiple neural circuits with multiple functions? *Vision Research*, *104*, 47-56. https://doi.org/https://doi.org/10.1016/j.visres.2014.08.018

Ohzawa, I., Sclar, G., & Freeman, R. D. (1982). Contrast gain control in the cat visual cortex. *Nature*, *298*(5871), 266-268. https://doi.org/10.1038/298266a0

Oliver, M. (2014). Nonlinear Modeling of the Early and Mid-Level Visual System. *Vision Science. California, Berkeley: University of California.*

Oliver, M., & Gallant, J. (2016). A deep convolutional energy model of V4 responses to natural movies. *Journal of Vision*, *16*(12), 876-876. https://doi.org/10.1167/16.12.876

Olmos, A., & Kingdom, F. (2004). *McGill calibrated colour image database*. http://tabby.vision.mcgill.ca

Olshausen, B. A. (1996). Emergence of simple-cell receptive field properties by learning a sparse code for natural images. *Nature*, *381*(6583), 607-609.

Pelli, D. G. (1997). The VideoToolbox software for visual psychophysics: Transforming numbers into movies. *Spatial Vision*, *10*, 437-442. https://doi.org/10.1163/156856897X00366

Ringach, D. L. (2002). Spatial Structure and Symmetry of Simple-Cell Receptive Fields in Macaque Primary Visual Cortex. *Journal of Neurophysiology*, *88*(1), 455-463. https://doi.org/10.1152/jn.2002.88.1.455

Rumelhart, D. E., Hinton, G. E., & Williams, R. J. (1986). Learning representations by back-propagating errors. *Nature*, *323*(6088), 533-536.

Sasaki, K. S., & Ohzawa, I. (2007). Internal spatial organization of receptive fields of complex cells in the early visual cortex. *J Neurophysiol*, *98*(3), 1194-1212. https://doi.org/10.1152/jn.00429.2007

Saul, A. B., & Humphrey, A. L. (1992). Evidence of input from lagged cells in the lateral geniculate nucleus to simple cells in cortical area 17 of the cat. *Journal of Neurophysiology*, *68*(4), 1190-1208. https://doi.org/10.1152/jn.1992.68.4.1190

Schwartz, O., Pillow, J. W., Rust, N. C., & Simoncelli, E. P. (2006). Spike-triggered neural characterization. *Journal of Vision*, *6*(4), 13-13.

Sengpiel, F., Sen, A., & Blakemore, C. (1997). Characteristics of surround inhibition in cat area 17. *Experimental Brain Research*, *116*(2), 216-228. https://doi.org/10.1007/PL00005751

Sietsma, J., & Dow, R. J. F. (1991). Creating artificial neural networks that generalize. *Neural Networks*, *4*(1), 67-79. https://doi.org/https://doi.org/10.1016/0893-6080(91)90033-2

Simoncelli, E. P., Paninski, L., Pillow, J., & Schwartz, O. (2004). Characterization of neural responses with stochastic stimuli. *The cognitive neurosciences*, *3*, 327-338.

Skottun, B. C., De Valois, R. L., Grosof, D. H., Movshon, J. A., Albrecht, D. G., & Bonds, A. (1991). Classifying simple and complex cells on the basis of response modulation. *Vision Research*, *31*(7), 1078-1086.

Srivastava, N., Hinton, G., Krizhevsky, A., Sutskever, I., & Salakhutdinov, R. (2014). Dropout: a simple way to prevent neural networks from overfitting. *J. Mach. Learn. Res.*, *15*(1), 1929–1958.

St-Amand, D., & Baker, C. L. (2022). Transient inhibition to light explains stronger V1 responses to dark stimuli. *bioRxiv*, 2021.2011.2021.469446. https://doi.org/10.1101/2021.11.21.469446

St-Yves, G., & Naselaris, T. (2018). The feature-weighted receptive field: an interpretable encoding model for complex feature spaces. *Neuroimage*, *180*(Pt A), 188-202. https://doi.org/10.1016/j.neuroimage.2017.06.035

Swindale, N. V., & Spacek, M. A. (2015). Spike detection methods for polytrodes and high density microelectrode arrays. *Journal of Computational Neuroscience*, *38*(2), 249-261.

Takane, Y., Young, F. W., & De Leeuw, J. (1977). Nonmetric individual differences multidimensional scaling: An alternating least squares method with optimal scaling features. *Psychometrika*, *42*(1), 7-67.

Talebi, V., & Baker, C. L., Jr. (2012). Natural versus synthetic stimuli for estimating receptive field models: a comparison of predictive robustness. *J Neurosci*, *32*(5), 1560-1576. https://doi.org/10.1523/jneurosci.4661-12.2012

Theunissen, F. E., David, S. V., Singh, N. C., Hsu, A., Vinje, W. E., & Gallant, J. L. (2001). Estimating spatio-temporal receptive fields of auditory and visual neurons from their responses to natural stimuli. *Network*, *12*(3), 289-316.

Tolhurst, D. J., Movshon, J. A., & Dean, A. F. (1983). The statistical reliability of signals in single neurons in cat and monkey visual cortex. *Vision Res*, *23*(8), 775-785. https://doi.org/10.1016/0042-6989(83)90200-6

Tootell, R. B., Silverman, M. S., & De Valois, R. L. (1981). Spatial frequency columns in primary visual cortex. *Science*, *214*(4522), 813-815.

Touryan, J., Felsen, G., & Dan, Y. (2005). Spatial Structure of Complex Cell Receptive Fields Measured with Natural Images. *Neuron*, *45*(5), 781-791. https://doi.org/https://doi.org/10.1016/j.neuron.2005.01.029

Vintch, B., Movshon, J. A., & Simoncelli, E. P. (2015). A convolutional subunit model for neuronal responses in macaque V1. *The Journal of Neuroscience*, *35*(44), 14829-14841.

Vogels, R., Spileers, W., & Orban, G. A. (1989). The response variability of striate cortical neurons in the behaving monkey. *Experimental Brain Research*, *77*(2), 432-436. https://doi.org/10.1007/BF00275002

Willmore, B., & Smyth, D. (2003). Methods for first-order kernel estimation: simple-cell receptive fields from responses to natural scenes. *Network*, *14*(3), 553-577.

Willmore, B. D. B., Prenger, R. J., & Gallant, J. L. (2010). Neural representation of natural images in visual area V2. *The Journal of neuroscience : the official journal of the Society for Neuroscience*, *30*(6), 2102-2114. https://doi.org/10.1523/JNEUROSCI.4099-09.2010

Wu, M. C., David, S. V., & Gallant, J. L. (2006). Complete functional characterization of sensory neurons by system identification. *Annu Rev Neurosci*, *29*, 477-505. https://doi.org/10.1146/annurev.neuro.29.051605.113024

Yamins, D. L., Hong, H., Cadieu, C. F., Solomon, E. A., Seibert, D., & DiCarlo, J. J. (2014). Performance-optimized hierarchical models predict neural responses in higher visual cortex. *Proc Natl Acad Sci U S A*, *111*(23), 8619-8624. https://doi.org/10.1073/pnas.1403112111

Zhan, C. A., & Baker, C. L., Jr. (2008). Critical spatial frequencies for illusory contour processing in early visual cortex. *Cereb Cortex*, *18*(5), 1029-1041. https://doi.org/10.1093/cercor/bhm139

Zhang, Y., Lee, T. S., Li, M., Liu, F., & Tang, S. (2019). Convolutional neural network models of V1 responses to complex patterns. *Journal of Computational Neuroscience*, *46*(1), 33-54. https://doi.org/10.1007/s10827-018-0687-7

Zhang, Y., & Young, L.-S. (2020). DNN-assisted statistical analysis of a model of local cortical circuits. *Scientific Reports*, *10*(1), 20139. https://doi.org/10.1038/s41598-020-76770-3

Zhou, Y. X., & C. L. Baker, J. (1996). Spatial properties of envelope-responsive cells in area 17 and 18 neurons of the cat. *Journal of Neurophysiology*, *75*(3), 1038-1050. https://doi.org/10.1152/jn.1996.75.3.1038