Microbiome diversity: A barrier to the environmental spread of antimicrobial resistance?

Uli Klümper[1,*], Giulia Gionchetta[2,*], Elisa C. P. Catao[3,4,*], Xavier Bellanger[3], Irina Dielacher[5], Peiju Fang[1], Sonia Galazka[6], Agata Goryluk-Salmonowicz[7], David Kneis[1], Uchechi Okoroafor[8], Elena Radu[5,9], Mateusz Szadziul[7], Edina Szekeres[10], Adela Teban-Man[10], Cristian Coman[10], Norbert Kreuzinger[5], Magdalena Popowska[7], Julia Vierheilig[5,11], Fiona Walsh[8], Markus Woegerbauer[6], Helmut Bürgmann[2], Christophe Merlin[3], Thomas U. Berendonk[1,#]

[1] Technische Universität Dresden, Institute for Hydrobiology, Zellescher Weg 40, 01217 Dresden, Germany

[2] Eawag, Swiss Federal Institute of Aquatic Science and Technology, Department of Surface Waters – Research and Management, 6047 Kastanienbaum, Switzerland

[3] Université de Lorraine, CNRS, LCPME, UMR 7564, Villers-lès-Nancy, France

[4] Université de Toulon, MAPIEM, CS 60584, Toulon, France

[5] Institute of Water Quality and Resource Management, TU Wien, Karlsplatz 13/2261, 1040 Vienna, Austria

[6] Department for Integrative Risk Assessment, Division for Risk Assessment, Data and Statistics, AGES – Austrian Agency for Health and Food Safety, Spargelfeldstraße 191, 1220 Vienna, Austria

[7] University of Warsaw, Faculty of Biology, Institute of Microbiology, Department of Bacterial Physiology, Miecznikowa 1, 02-096 Warsaw, Poland

[8] Department of Biology, Kathleen Lonsdale Institute for Human Health, Maynooth University, Maynooth, Co. Kildare, Ireland

[9] Institute of Virology Stefan S. Nicolau, Romanian Academy of Science, 285 Mihai Bravu Avenue, 030304, Bucharest, Romania

[10] NIRDBS, Institute of Biological Research Cluj-Napoca, 400015, Cluj-Napoca, Romania

[11] Interuniversity Cooperation Centre Water & Health, Austria

* Contributed equally to this work

[#] Corresponding author

Corresponding author address:

Prof. Thomas Berendonk (ORCID: 0000-0002-9301-1803)

Technische Universität Dresden

Institute of Hydrobiology

Zellescher Weg 40

01217 Dresden

Germany

Tel.   +49 351 463-34956

Fax   +49 351 463-37108

Mail: thomas.berendonk@tu-dresden.de

Abstract

Background: In the environment, microbial communities are constantly exposed to invasion by antimicrobial resistant bacteria (ARB) and their associated antimicrobial resistance genes (ARGs) that were enriched in the anthroposphere. A successful invader has to overcome the biotic resilience of the habitat, which is more difficult with increasing biodiversity. The capacity to exploit resources in a given habitat is enhanced when communities exhibit greater diversity, reducing opportunities for invaders, leading to a lower persistence. In the context of antimicrobial resistance (AMR) dissemination, exogenous ARB reaching a natural community may persist longer if the biodiversity of the autochthonous community is low, increasing the chance of ARGs to transfer to community members. Reciprocally, high microbial diversity could serve as a natural long-term barrier towards invasion by ARB and ARGs.

Results: To test this hypothesis, a sampling campaign across seven European countries was carried out to obtain 172 environmental samples from sites with low anthropogenic impact. Samples were collected from contrasting environments: stationary structured forest soils, or dynamic river biofilms and sediments. Microbial diversity and relative abundance of 27 ARGs and 5 mobile genetic element marker genes were determined. In soils, higher diversity, evenness and richness were all significantly negatively correlated with the relative abundance of the majority (>85%) of ARGs. Furthermore, the number of detected ARGs per sample was inversely correlated with diversity. However, no such effects were found for the more dynamic, regularly mixed rivers.

Conclusions: In conclusion, we demonstrate that diversity can serve as barrier towards AMR dissemination in the environment. This effect is mainly observed in stationary, structured environments, where long-term, diversity-based resilience against invasion can evolve. Such barrier effects can in the future be exploited to limit the environmental proliferation of AMR.

Keywords

Biodiversity; Microbial barrier; Microbial Invasion, Riverbed microbiome; Soil microbiome; ARG

Background

The spread of antibiotic resistance genes (ARGs) represents one of the biggest challenges to future human and animal health [1,2]. It has become clear that tackling the issues brought on by the expansion of antimicrobial resistance (AMR) requires a global effort that spans the fields of human, veterinary, and environmental health [3,4]. This perspective, also known as the "One Health" approach, is frequently used as a framework for worldwide efforts to contain the spread of AMR. However, integrating the environmental sphere has, due to its high complexity, proven particularly challenging. Understanding the existing ecological dispersal barriers is necessary to restrict the propagation of AMR in the environment [5,6].

ARGs are ancient, naturally occurring in environmental bacteria and evolved over billions of years [7]. However, during the last century, environmental microbial communities have been constantly subjected to invasion events by antibiotic resistant bacteria (ARB), and their associated ARGs, which have been enriched or released through anthropogenic activities. For example, release and reuse of wastewater effluents or the application of manure to soils are known to be primary conduits of AMR spread in the aquatic and terrestrial environments [8–11]. But even habitats with no direct anthropogenic impact are regularly exposed to lower frequencies of such invasion events, for example, through wildlife and aerial depositions. These natural diffuse sources of AMR-pollution promote the widespread dispersal of AMR over time at low levels [12–16].

Invasion has been defined, both on the micro- and the macro-biological scale, as a process consisting of a sequence of successive steps, namely, (i) introduction, (ii) establishment, (iii) growth and spread, and (iv) impact [17]. For it to be successful, the invader has to overcome the "biotic resistance" towards invasion of the habitat [18]. In theory, this process becomes more difficult with increasing biodiversity [19]. The capacity to exploit the resources provided by a habitat is enhanced when communities exhibit a greater diversity, which in turn reduces

opportunities for invaders, hence lowering their persistence [20]. If an invader is phylogenetically,

thus presumably also physiologically, close to established community members occupying its

specific niche, the invasion process has been shown to become merely stochastic [21]. In

contrast, even small-scale disturbance events can considerably increase invasion success by

affecting the niche occupancy of resident species [22]. However, such effects strongly depend on

the niche partitioning within the given environment rather than diversity alone [23], and might

hence be far more pronounced in structured environments with a long-term established niche

occupation.

In the context of AMR dissemination, it is reasonable to assume that - as any invader -

ARB reaching a natural community may persist longer when the biodiversity of the autochthonous

community is low, as this usually coincides with a lower rate of niche occupation. This could result

in a longer-term establishment of the ARB invaders, and thus increase the relative abundance of

ARGs. Even when the invasion process is not successful in the long term, a slightly prolonged

residence time of the invader could have a significant impact on the likelihood of ARGs being

horizontally transferred to the endemic microbiota [24]. Alternatively, future invasion events could

be favored by reducing the community resilience [25]. Reciprocally, high microbial diversity could

impede the spread of ARB and ARGs, thus serving as a potential natural barrier.

Diversity as a limiting factor for AMR invasion was demonstrated in the short-term for

laboratory soil microcosms inoculated at different diversities using an artificial dilution-to-

extinction approach. Less diverse soil microcosms displayed a far higher likelihood of being

invaded by ARBs [26]. Regarding the aquatic dimension, increased invasion success of a

resistant *E. coli* strain into river biofilm communities was shown to coincide with a loss in diversity

of these communities under stress conditions [27]. Further, diverse microbial communities with a

high degree of functional niche coverage, such as activated sludge, have been suggested to

provide natural barriers for the proliferation of AMR [28]. However, data suggests that activated

sludge communities have also incorporated a particularly high diversity of ARGs encoded on

mobile genetic elements (MGEs) [29].

Based on this theoretical and experimental knowledge, we hypothesize that the invasion

success and, in consequence, the pervasiveness of AMR into environmental microbiomes is

inversely correlated to the diversity of the communities in question. To our knowledge, no prior

field-based study has explored whether AMR dissemination is in the long-term indeed related to

microbial diversity in terrestrial or aquatic environmental microbiomes. To this end, low impacted

environmental samples with no direct anthropogenic depositions were collected. Their ARG

levels, beyond natural background levels, are hypothesized to result from the accumulation of

invasion success over time from invaders introduced through the previously mentioned low level

dispersion routes. Therefore, ARG levels would rise, if after a successful introduction, these

invaders were able to either establish themselves long-term in the indigenous microbial

communities or transferred their mobile ARG load. Consequently, 172 of such low anthropogenic

impact environmental samples were collected during fall/winter 2020/21 across seven European

countries. Half of these were taken from forest soils, representing a stationary, structured

environment, while the other half was obtained from river sediments and biofilms representing a

more dynamic environment. For each sample, the microbial diversity was assessed through

bacterial 16S rRNA gene-based amplicon sequencing, while the abundance of 27 AMR markers

and 5 MGEs was determined through high-throughput chip-based qPCR to ultimately determine

how microbial diversity as a potential barrier determines ARG abundance in low-impact

environments.

Methods

Soil sampling and processing

The terrestrial campaign consisted of collecting 74 forest soil samples from the seven

countries during fall 2020 (Fig. S1). The aim was to obtain sample sets that included samples

across a gradient of high and low microbial diversity that are of relatively low anthropogenic impact

(Table S1).

From each forest location five single core samples (Pürckhauer drill, Buerkle™, Germany)

were extracted from a depth of 0-25 cm along two 10 m virtual diagonals laid across the sampling

location in the form of an X-pattern. 200 g of each of these five subsamples were combined in an

aseptic plastic bag, thoroughly homogenized and transferred to the laboratory at 10 °C. From the

composite sample aliquots of 20 g were sieved (2 mm mesh size) and stored at -20 °C. DNA

extraction was performed using the DNeasy PowerSoil Pro Kit (Qiagen, Germany) according to

the manufacturer's instructions. To obtain DNA from a total of 1 g of each sieved soil sample four

replicates of 0.25 g each were extracted in parallel and combined thereafter. The quality and

quantity of the extracted DNA was assessed spectrophotometrically.

Riverbed material sampling and processing

The aquatic campaign included the collection of 98 river samples from seven countries

(Austria, France, Germany, Ireland, Poland, Romania, and Switzerland) during the fall/winter

2020/21 (Fig. S1). The locations were selected to obtain samples across a gradient of high and

low microbial diversity that are of relatively low anthropogenic impact (e.g., not exposed to

wastewater treatment plant effluents) (Table S2). At each site, the substrate best representing

sessile, non-phototrophic, oxygenated microbial communities in the chosen riverbeds, was

sampled. Either epilithic biofilms from the undersides of rocks to avoid phototrophic communities

for those streams dominated by rock/gravel, or oxygenated sediment for streams dominated by

fine sediment were collected. Specifically, for epilithic biofilm samples, five individual rocks,

collected from a shadowed sample area from a riverbed length of approximately 10 meters, were

gently scraped from the bottom surface using a sterile toothbrush, and combined to create a

composite river biofilm sample. Repeated rinsing with sterile water in a 50 mL falcon tube was

performed to collect the biomass. If no rocks or rock biofilms were available, fine surface sediment

from shaded areas was sampled. In this case, the upper layer (~ 5 cm) of sediment was collected

using a 50 mL falcon tube. Five sediment cores were combined at equal weight to obtain one

composite sample.

All collected samples were gently homogenized and transported to the laboratory on ice.

Then, samples were centrifuged (4,000 rpm for 5 min at 4 °C), the supernatant removed, pellets

weighted and stored at -20 °C. DNA extraction was performed using the DNeasy PowerSoil Pro

Kit (Qiagen, Germany) according to the manufacturer's instructions. Extraction blanks were used

to confirm the absence of DNA contamination. The quality and quantity of extracted DNA was

assessed spectrophotometrically.

Amplicon sequencing and analyses of sequence datasets

To analyze the microbial diversity and taxonomic composition of the samples, DNA

extracts were sent to the IKMB sequencing facility (minimum 10,000 reads per sample; Kiel

University, Germany). Illumina MiSeq amplicon sequencing of the bacterial 16S rRNA gene was

performed using primers targeting the V3-V4 region (V3F: 5′-CCTACGGGAGGCAGCAG-3′ V4R:

5′-GGACTACHVGGGTWTCTAAT-3) [30]. Sequences were collectively analyzed using DADA2

[31] in QIIME2 [32]. Forward and reverse reads were merged into amplicon-sequence variants

(ASV), at 99% sequence similarity. Prior to downstream analyses, the *filter-features* and *filter-*

table scripts were applied in QIIME2 to clean ASV and taxa tables by removing unclassified and

rare ASVs with a frequency of less than 0.1% of the mean sample depth. The corresponding river

and soil ASV tables consisted of 15,473 ASVs (river dataset) and 14,464 ASVs (soil dataset). All

sequencing data was submitted to the NCBI sequencing read archive under project accession

number PRJNA948643.

High-throughput qPCR of ARGs and genetic markers for MGEs

To determine the relative abundance of target genes in each sample, DNA extracts were

sent to Resistomap Oy (Helsinki, Finland) for HT-qPCR analysis using a SmartChip Real-time

PCR system (TaKaRa Bio, Japan). The target genes included 27 ARGs and 5 MGEs (Table S3)

[33]. In addition, the 16S rRNA gene and the anthropogenic fecal pollution indicator crAssphage

[34,35] were quantified. The protocol was as follows: PCR reaction mixture (100 nL) was prepared

using SmartChip TB Green Gene Expression Master Mix (TaKara Bio, Japan), nuclease-free

PCR-grade water, 300 nM of each primer, and 2 ng/µL DNA template. After initial denaturation at

95 °C for 10 min, PCR comprised 40 cycles of 95 °C for 30 s and 60 °C for 30 s, followed by

melting curve analysis for each primer set. A cycle threshold (CT) of 31 was selected as the

detection limit [36,37]. Amplicons with non-specific melting curves or multiple peaks were

excluded. The relative abundances of the detected gene to 16S rRNA gene were estimated using

the ΔCT method based on mean CTs of three technical replicates [38].

Data analysis and visualization

To characterize the alpha diversity of the terrestrial and aquatic microbial communities,

the diversity indices Chao1 richness, Shannon diversity and Pielou evenness were calculated

using the *core-metrics-phylogenetic* script in QIIME2 [32] on rarefied data, with samples with an

insufficient sampling depth to reliably assess diversity removed from the datasets. The taxonomy

of these microbial communities was classified based on the SILVA classifier (version 138, [39]).

Beta-diversity dissimilarities among microbial communities were assessed using a principal

coordinate analysis (PCoA) based on Bray-Curtis distance matrices [40].

Differences in the relative ARG abundance for aquatic and terrestrial samples were

transformed to a $\log_{10}$ scale, and displayed using the package *ComplexHeatmap* [41]. To explore

the degree of correlation present among the ARGs in both environments, pairwise correlations

analysis based on Spearman rank were assessed. Only correlations showing coefficients |ρ| >

0.75 were considered significant at $p < 0.05$ after Benjamini-Hochberg correction for multiple

testing. The connectivity in relative abundance among ARGs in aquatic and terrestrial samples

was displayed through network analysis, performed using the *picante* package [42] based on

significant Spearman correlations. The network visualization was generated using the open-

source software Gephi v8.2. To test the correlation between resistome and bacterial diversity,

correlation analysis between relative ARG abundance and the calculated alpha diversity metrics

was performed based on Spearman rank correlation or Pearson correlation followed by Bonferroni

correction for multiple testing. Individual groups of data were compared using One-Way ANOVA

with post-hoc Tukey HSD tests. Significant differences of correlations from 0 were tested using a

t-test. All statistics and plots were produced using R version 4.2.0 [43].

Results

Assessing diversity in the river and soil dataset

Two complementary sets of samples were obtained from a total of 94 river as well as 70

soil samples. When assessing the beta diversity of the microbial communities, soil samples

differed significantly and with a large effect size from those obtained from river biofilms

(PERMANOVA, pseudo-F = 14.73, p < 0.001, n = 148) and river sediments (pseudo-F = 5.53, p

< 0.001, n = 78) (Fig. 1A). No clear distinction was observed comparing sediments and epilithic

biofilms (pseudo-F = 2.77, n = 94). Consequently, these samples were subsequently grouped to

create the combined river dataset. No clear trends with regards to the countries of origin were

observed for either dataset, but the spread of the entire dataset exceeded the spread of any

sample set from a single country and some river communities from Romania formed a distinct

cluster (Fig. 1A).

The aquatic samples contained 19 phyla with an average relative abundance above 1%

and were, throughout, dominated by bacteria belonging to the phyla *Proteobacteria*, *Bacteroidota*

and *Actinobacteriota* (0.26 ± 0.09, 0.17 ± 0.13, 0.12 ± 0.09) (Fig. S2). In the soil dataset

Acidobacteria, *Actinobacteriota* and *Proteobacteria* (0.18 ± 0.04, 0.15 ± 0.02, 0.13 ± 0.03)

dominated (Fig. S3). No significant differences in dominant phyla between samples based on

country of origin were observed. CrAssphage, a common indicator for recent anthropogenic fecal

pollution, was undetected in the entirety of soil samples and in 78% of the river samples. For the

latter it remained at low relative abundance below 10^{-5} copies per copy of the bacterial 16S rRNA

gene, confirming that samples were indeed of low anthropogenic impact origin. For the three main

alpha-diversity metrics - Chao1 richness, Shannon diversity and Pielou evenness - high and low

biodiversity samples for each of the two datasets, rivers and soils, were obtained (Fig. 1B). The

main distinction between the two datasets was the significantly higher level of Pielou evenness in

the dataset from the structurally stable soil (0.95 ± 0.02) compared to the dynamic river

environment (0.89 ± 0.08) ($p < 0.0001$, $f = 48.78$, One-Way ANOVA). The differences obtained

subsequently allowed these diversity metrics to be used as test variables for correlation with ARG

abundance.

Figure 1. Diversity of the river and soil datasets. Symbols depict sample type, colors code for the country of origin. A) PCoA of the beta diversity based on Bray Curtis distance of ASV relative abundance data from riverbed materials (sediments and biofilms) and soil. B) Alpha-diversity indices (Chao1 richness, Shannon diversity and Pielou evenness) from riverbed materials (top) and soil (bottom) collected from the seven countries.

Resistome diversity and abundances

In both resistome datasets, the *aac(3)-VI* gene conferring aminoglycoside-resistance was

the most abundant (Fig. 2). In the river dataset, genetic determinants for sulfonamide (*sul1*),

vancomycin (*vanA*), colistin (*mcr1*) and phenicol (*floR*) resistance clustered together as dominant,

followed by other ARGs that promote resistance to macrolides, lincosamides and streptogramins

B - (*mphA*), β-Lactam (*bla$_{CTX-M2}$, bla$_{CTX-M1}$, bla$_{CMY2}$*), phenicol (*cmlA*) and aminoglycoside (*aac(6')-*

lb3, a*ph(3')-lb*) antibiotic classes. ARGs conferring resistance to quinolone, trimethoprim and

tetracycline were less abundant, although in certain soils, particularly from Poland and Romania,

the corresponding abundance of individual genes (e.g., *tet(W), qnrS, drfA*) was higher (Fig. 2A).

In general, considerable clustering of samples was observed. The Irish samples clustered

together with a few Romanian and one German sample separately from the rest, and displayed

an overall lower relative ARG abundance with the exception of the *blaTEM* gene, which displayed

a particularly high abundance (Fig. 2A).

In the soil sample set, the *acc(3)-VI* (aminoglycoside), *qnrS1* (quinolone), *mphA* (MLS$_B$)

and *dfrA1* (trimethoprim) genes clustered together as the most abundant determinants in most

countries (Fig. 2B). ARGs conferring resistance to colistin (*mcr-1*), Phenicol (*cmlA2*), β-Lactams

(*bla$_{CMY-2}$, bla$_{CTX-M}$*) and aminoglycoside (a*ph(3')-lb, aac(6')-lb3*) were detected in most countries

at intermediate abundances, but with highest values in Switzerland and France. The colistin ARG

mcr-3 was the sole ARG below the limit of detection for all samples, therefore it was not used for

any further analysis. Accordingly, the Swiss and French soil resistomes clustered together as

most of the ARGs were found at higher and similar abundances compared to the other countries

(Fig. 2B). Irish soils displayed again the lowest number of ARGs detected and similar to the river

dataset the abundance of *blaTEM* was significantly elevated in Ireland compared to other

countries (Fig. 2B).

Figure 2. Heatmap of relative ARG abundances in the river (A) and soil (B) dataset. Values are displayed after transformation to log10 scale. The list of ARGs is presented based on similarity in abundance patterns and displayed from high abundance (red) to below the detection limit (blue). Color coding on the right displays the class of antibiotics they confer resistance to. Samples are ordered according to similarity in ARG profiles and color coded based on country of origin displayed on top.

Higher degree of correlation between relative ARG abundances in soils compared to rivers

In both datasets, a high number of significant correlations were found between the relative abundance of different ARGs. Out of all potential comparisons between individual relative ARG abundances 140 out of 351 (27.40%) for rivers and 144 out of 325 (44.31%) for soils were significantly positively correlated with each other ($p < 0.05$, Spearman) (Fig. 3 A,B). Only in the case of aminoglycosides resistance gene *aph(3')-Ib* and trimethoprim resistance gene *dfrA1* in the river dataset, a single negative correlation in relative abundance could be detected. While in the river environment the distribution of correlations did not follow a clear pattern, in soil a distinct cluster of 17 ARGs that are highly correlated among each other was observed. This cluster included ARGs conferring resistance to antibiotics belonging to different classes, namely vancomycin, tetracyclines, quinolones, polymyxins, phenicols, MLS_B, β-Lactams and aminoglycosides. Further the average correlation coefficient of the significant comparisons (Rs) among all soil ARGs was significantly higher ($R_s = 0.40 \pm 0.33$) than among the river dataset ($Rs = 0.30 \pm 0.29$; $p < 0.0001$, ANOVA) (Fig. 3). This increased level and higher degree of consistency of correlations between ARGs in soil was further confirmed using network analysis, where the average degree of connections of each ARG with the remaining ones was significantly higher in soil (10.533) compared to the river dataset (5.692) (Fig. S4).

No significant correlation of the observed ARG number or the relative abundance of any ARGs with the relative abundance of crAssphage was obtained for the river dataset (all $p > 0.05$,

Spearman) while crAssphage was absent in the soil dataset. Thus, it again demonstrates that

results are not directly impacted by recent anthropogenic fecal pollution.

Figure 3. Correlation analysis between relative ARG abundances: Pairwise correlations in

river (A) and soil (B) microbial communities based on Spearman rank correlation. Only significant

comparisons (p < 0.05 after Benjamini-Hochberg correction for multiple testing) are shown.

Intensity of colors displays strength of correlation with orange depicting positive and blue depicting

negative correlation.

Diversity as a barrier to ARG spread

To determine whether higher diversity could lower the long-term invasion success of

ARGs by the communities, we examined correlations between the diversity metrics and the

number of detected ARGs as well as the relative abundance of each individual gene in both

datasets. On average, 18.44 ± 5.61 of the 27 ARGs tested were successfully detected in samples

from the river dataset. No clear trend could be observed in correlation between the number of

detected ARGs and any of the three diversity metrics. While all three correlations were negative,

none were statistically significant (r_{Pielou} = -0.02, $r_{Shannon}$ = -0.04, r_{Chao1} = -0.11, all p > 0.05; Pearson

correlation; Fig. 4 A-C).

Figure 4. Correlation analysis of the number of ARGs detected per sample with diversity metrics

based on Pearson correlation with Bonferroni correction for multiple testing. Linear correlations

from river environmental samples with Pielou Evenness (A), Shannon Diversity (B) and Chao1

Richness (C). Linear correlations from soil environmental samples with Pielou Evenness (D),

Shannon Diversity (E) and Chao1 Richness (F). Colors depict the country of sample origin and the

symbols depict the sample type.

Slightly, but significantly, less ARGs per sample (15.95 ± 6.05; p = 0.014, f = 6.11, One-

Way ANOVA) were successfully detected in the soil dataset. Contrary to the river dataset, higher

diversity in soils correlated with a lower number of detected ARGs. This negative correlation was

significant based on Spearman rank correlation analysis for Pielou evenness (r = -0.35, p =

0.0042) and Shannon diversity (r = -0.37, p = 0.0027) (Fig. 4 D,E). Similarly, for Chao1 Richness

an inverse correlation with the number of ARGs detected was observed, however, barely not

significant ($r = -0.25$, $p = 0.051$) (Fig. 4 F). These results provided a first indication that diversity-

based barrier effects might indeed exist, at least in the more structured soil environment. Although

diversity had a significant impact, the effect sizes of 25-37% suggest that, as expected for complex

environmental datasets, diversity is only one of multiple interacting drivers of the observed trends.

To further test this hypothesis the relative abundance of each individual ARG was

correlated with the obtained diversity metrics. To account for zero-inflation during correlation, only

those ARGs that were found in at least 25% of the samples of the respective dataset were tested.

In the river dataset, similar to the number of ARGs no clear trends were observed for the relative

abundance of any of the tested ARGs (Fig. 5 A-C). The only correlation considered significant

based on Spearman rank correlation (with Bonferroni correction for multiple testing) was a positive

one between the *blaTEM* gene and the Chao1 richness ($R_S = 0.42$, $p = 0.0003$). For the remaining

combinations of ARGs and diversity metrics only slight negative or slight positive correlation

trends ($R_S = -0.24 - 0.33$, all $p > 0.05$) were observed. Among those non-significant trends, no

obvious patterns emerged. In fact, the average Spearman's *rho* of the tested ARGs for each of

the three diversity metrics was near 0 ($p > 0.05$; t-test with Bonferroni correction for multiple

testing).

In contrast, a high number of significant negative correlations of relative ARG abundance

with the different diversity indices were observed based on Spearman Rank correlation analysis

in the soil dataset (Fig. 5 D-F). Pielou evenness and Shannon diversity displayed the most

significant correlations with 13 of the 18 tested ARG relative abundances being negatively

correlated, while six ARGs were negatively correlated with Chao1 richness. ARGs negatively

correlated with diversity were widely distributed across antibiotic classes. Similar to the river

environment, the *blaTEM* gene was the main exception from the observed trend and positively

correlated to either diversity metric ($R_S = 0.47-0.51$, all $p < 0.05$, Fig. 5 D-F). Despite this outlier, a

general trend for the correlation between relative ARG abundance and diversity was observed for

the soil dataset, with the average correlation coefficients of all tested ARGs being both negative

and significantly different from 0 for Chao1 richness (R_S = -0.257 ± 0.239, p = 0.0004), Pielou

evenness (R_S = -0.234 ± 0.228, p = 0.0001) and Shannon diversity (R_S = -0.267 ± 0.223, p =

0.0003, Fig. 5 D-F). Finally, contrary to ARGs, none of the five indicator genes for MGEs tested

(the class1 integron integrase gene *intI1*, the *IncP* plasmid *oriT*, the *IncW* plasmid *trwAB* gene,

the *ORF37* gene of *IS26* and the Tn*5* transposase gene) displayed any correlation with either of

the diversity indices in any of the datasets (all p>0.05).

Figure 5. Correlation analysis of relative ARG abundance with observed diversity metrics based on Spearman rank correlation with Bonferroni correction for multiple testing. Correlations from river environmental samples with Pielou Evenness (A), Shannon Diversity (B) and Chao1 Richness (C). Correlations from soil environmental samples with Pielou Evenness (D), Shannon Diversity (E) and Chao1 Richness (F). Filled bars represent significant, while hatched bars represent non-significant correlations. Colors depict the class of antibiotic the ARG confers resistance to. Only ARGs that were detected in at least 25% of samples of a dataset were tested.

Discussion

Here we demonstrate based on analysis of a pan-European sampling campaign that communities of high bacterial diversity are more resistant to ARG pervasiveness, and that diversity might serve as a barrier to the long-term invasion and establishment of ARGs into environmental endemic microbiomes. Both, the number of detectable ARGs as well as a majority of the individual relative ARG abundances were negatively correlated with the diversity indices observed in soil. Among these indices, Pielou evenness was most significantly negatively correlated to the number of detected ARGs and their relative abundances. While this possible effect of community diversity on long-term invasion and establishment of resistant bacteria was highly visible and frequently statistically significant in the structured soil environment, it was barely observed in the more dynamic river environment.

Within the context of these results, the effects of diversity on invasion of AMR needs to be assessed individually for the successive steps that make up a successful invasion event, namely 1) introduction of the invader, 2) its establishment, 3) its growth and spread along with 4) its impact in the new microbial community [17]. The initial introduction of each invader is primarily of stochastic nature and does not rely on biological interactions with the indigenous community [17,20]. Consequently, the success of these initial introduction events depends on the quantity of invaders present, also known as the propagule pressure, together with the level of physical interaction of these invaders [21,44]. In this study, the samples originated from low impacted soils and rivers across Europe. Here, we define low impact as being not in direct proximity to the release of bacteria enriched in ARGs through anthropic action such as treated wastewater effluents [45,46] or manure [37,47]. The propagule pressure - the number of invaders harboring ARGs that were introduced into these environments - can be assumed low at the time of sampling and was likely low in the past. However, there is a high probability that bacteria with ARGs acquired in the antibiotic era occurred, nevertheless, at some rate (e.g. through human presence

or transport by wild and domestic animals, including defecation, wet and dry atmospheric deposition). Consequently, it can be assumed that any increase in the resistomes in our samples are unlikely to stem from recent pollution events, but rather from past invasion events that manifest on top of the more or less universal background levels of resistance recently determined for a number of environments [48,49]. Increases in ARG occurrence and relative abundance would hence result from the accumulation of invasion success of previous repetitive introductions of invaders over time that have been able to establish themselves in the autochthonous microbiome or left their mobile ARG load behind, if we consider that bacteria from the human or animal spheres are regularly not fully fit to be long-term maintained in environmental microbiomes.

Contrary to the original introduction step of the invader, where biological interactions play only a minor role, the interactions with the local community are highly relevant during the subsequent establishment and growth phases. The internal resistance of the indigenous community towards invasion, e.g., its biotic barriers, have to be overcome to lead to the successful establishment of the invader [17,20] and can lead to the maintenance of ARGs in the community, when transient invasion success is long enough to allow for gene transfer [24]. Here we suggest that in the long term, microbial diversity might provide a biotic barrier, hindering ARG success of invasion in the low impacted soil microbial communities. However, no such effect could be observed for river communities. In the context of ARB such diversity effects have earlier been mainly demonstrated in short-term laboratory experiments for both types of environments using soil microcosms [26] and laboratory river flume experiments [27]. Still, in these experiments, diversity was artificially lowered to non-natural levels, which made it difficult to evaluate if such effects would equally be observable in the environment across natural biodiversity gradients in the long-term. In our analysis we make the implicit assumption that the present-day diversity is indicative also of past diversity, or at least of differences in diversity between sites and that both soil and riverbed microbiomes act as records of the long-term impacts endured by those microbial communities. This assumption appears likely to be correct in the case of forest soils, which are

supposedly an environment typically stable over decades or more [50]. However, the assumption

could be challenged regarding riverbeds, which are more likely exposed to considerably different

conditions in the past (considering e.g., droughts, changes in water quality of European rivers

over recent decades, etc.) [51]. This aspect may have contributed to the contrasting results in

these environments.

The observed differences between the two datasets regarding the correlations between

diversity and ARGs abundances likely originate from the different nature of the two environments.

The resistance of the community to the invasion processes is directly related to the number of

available niches for invader establishment, with more diverse communities providing less

available niche spaces, referred to in macro-ecology as the "diversity-invasion effect" [20]. In the

stationary and structured soil environment, the amount of available niches rarely changes and is,

once occupied by a diverse community, able to reach a steady-state [52]. Under these

circumstances it is unlikely that niches open up for invaders as no major loss of community

members is to be expected. Within this context it is also unsurprising that community evenness

is more strongly correlated with lower ARG abundance in soil than community richness. In highly

even communities, a higher number of bacterial species are abundant enough to fully occupy

their specific available niche space [53]. Rich communities have an intrinsically increased

potential for their populations to occupy a higher number of different niches [54]. However, once

a certain level of richness is reached, niche occupation moves towards saturation with additional

species playing only a minor role [55]. Increased richness might then lead to increased

competition and potential shifts within the metabolic networks, with novel niches becoming

available to be exploited by the invading strains. With these two mechanisms at play, the negative

correlation of richness with relative ARG abundances remains weaker than that observed for

evenness or Shannon diversity.

In the river microbiomes, microbial diversity and available niches are potentially rather

transient than long-term established due to the constant currents, biofilm adhesion and microbial

dispersion events and alterations in nutrient availability [56]. Therefore, the effect of diversity on

the long-term establishment of ARGs, which was observed for soil, may in the river microbiomes

be masked by the dynamic nature of this environment. This difference between less (soil) and

more (riverbed) dynamic environments is also displayed by the high number of ARGs being

correlated in abundance in soil compared to the river dataset. While in soil the majority of

abundant ARGs are present and affected by the described processes, in rivers ARGs rather

appear to be more independent of each other based on dynamic processes.

466   Unlike for biotic invaders, such as alien bacteria, the invasion success of genetic invaders,

such as ARGs, goes far beyond the successful establishment and growth of their hosts in the

novel environment. A prolonged residence time of the host can favor the spread of mobile ARGs

via horizontal gene transfer from the invading to the indigenous bacteria, thus leading to an even

higher persistence of ARGs [24]. This can even be the case if the host's invasion is only transiently

successful and invaders are lost after some time due to high community resilience in the observed

environment. Mobile ARGs encoded on plasmids are able to spread from an invading donor strain

to highly diverse proportions of soil and water derived microbial communities, with bacteria

belonging to over 25 different phyla able to receive individual resistance encoding plasmids [57–

61]. However, effects of community diversity on the efficiency of horizontal gene transfer and the

maintenance of plasmids in the community consist of a complex interplay of different mechanisms

and remain difficult to predict. On the one hand, at higher diversity an increased number of

potential plasmid hosts and conjugation partners are available that can lead to increased plasmid

maintenance and transferability in the community [62,63] increasing the chance of transfer to a

highly competitive host. On the other hand, in more diverse communities it can be harder to

encounter a permissive conjugation partner, which reduces transferability due to this dilution

effect [64]. Further, competition with other community members might increase the costs of

resistance [65] and could ultimately drive the loss of ARG hosting plasmids from the community

[66]. This loss process would be expected to be elevated in more diverse communities with better

competitors. The complex interplay of mechanisms leading to a poor predictability of effects is

equally represented in our soil dataset, where unlike for ARGs no clear correlation for MGEs (e.g.,

intI1, IncP plasmid *oriT*) with diversity could be established. However, a good indication that

community diversity might also limit the horizontal acquisition of mobile ARGs from invading

bacteria is that diversity is also negatively correlated with the number of detected ARGs in the soil

dataset. This is according to ecological theory, where species diversity is not always immediately

implying a higher degree of genetic diversity [67,68]. Still, assuming that, in the long-term,

invaders harboring the tested ARGs reach each of the tested communities it becomes apparent

that an increasing number of ARGs are not successfully retained in those communities of higher

diversity. If this is due to a shorter residence time of the invader, the above discussed increased

competition or dilution effects needs future research.

Conclusions

In summary, we display that the microbial diversity within a given environment could affect

the proliferation of AMR within and through this environment. Considering sites of low

anthropogenic impact, the observed negative correlation of diversity with detection and

abundance of ARGs in soils compared to river ecosystems can be directly connected to the

intrinsic characteristics of the specific environments within the framework of invasion theory as

well as horizontal gene transfer dynamics.

Natural environments, such as rivers and soils, may play a key role in AMR development

and proliferation. The characteristics of the individual environment, its texture, its dynamism as

well as the diversity of the resident microbial community could define its role as a source or a

barrier to AMR dissemination. We present support that in the structured soil environment, a high

bacterial diversity might indeed serve as a barrier to the long-term invasion and establishment of

ARGs in the autochthonous microbiome. Our results point to a previously overlooked benefit of

healthy environments, with diverse microbial communities, providing natural barrier effects to the

proliferation of AMR, thus clearly displaying how environmental and human health are

immediately interconnected through the One Health concept. Furthermore, such barrier effects

can be exploited within soil ecosystem management, for example, in defining optimal locations

for aquifer recharge through wastewater reuse. Here choosing locations with a high intrinsic

diversity could be beneficial in limiting the spread of wastewater born ARGs. To achieve this, the

role of microbial diversity in the dissemination and mobilization of AMR markers requires a closer

look through targeted experiments aimed at elucidating the exact mechanisms that limit the

proliferation of resistance determinants and how exploiting such natural barrier effects could have

cascading effects on the ecosystem biodiversity.

Declarations

<u>Ethics approval and consent to participate</u>

Not applicable

<u>Consent for publication</u>

Not applicable

<u>Availability of data and material</u>

The datasets supporting the conclusions of this article are included within the article and its

additional files. Original sequencing data is available in the NCBI sequencing read archive under

project   accession   number   PRJNA948643

(https://www.ncbi.nlm.nih.gov/bioproject/PRJNA948643), with individual sample identifiers given

in tables S1 & S2.

<u>Competing interests</u>

The authors declare that they have no competing interests.

Funding

This work was supported by the ANTIVERSA project (BiodivERsa2018-A-452) here funded by

the Bundesministerium für Bildung, und Forschung of Germany [01LC1904A], the French Agence

Nationale de la Recherche [ANR-19-EBI3-0005-04], the Swiss National Science Foundation

[186531], the Austrian Science Fund (FWF) [I 4374-B], the Irish Environmental Protection Agency

[2019-NC-MS-9], the National Science Centre (NCN) of Poland [UMO-2019/32/Z/NZ8/00011], the

Romanian National Authority for Scientific Research and Innovation (CCCDI – UEFISCDI)

[117/2020]. UK & TUB were supported through the Explore-AMR project funded by the

Bundesministerium für Bildung, und Forschung under grant number 01DO2200. AT-M and ES

were supported by the Ministry of Research, Innovation and Digitization through the Core Project

BIORESGREEN, subproject BioClimpact no. 7/30.12.2022, code 23020401. PF was supported

through the China Scholarship Council (CSC) under grant number 202004910327. DK was

supported through the Urban Resistome project funded by the Deutsche

Forschungsgemeinschaft (DFG) under project number 460816351. Responsibility for the

information and views expressed in the manuscript lies entirely with the authors.

Author contributions

Conceptualization of the study and sampling strategy: UK, GG, EC, XB, SG, AGS, ES, CC, NK,

MP, FW, MW, HB, CM, TUB; Identification of national sampling locations, sampling, metadata

collection & sample processing: UK, GG, EC, XB, ID, SG, AGS, UO, ER, MS, ES, AT-M, CC, NK,

MP, JV, FW, MW, HB, CM, TUB; Sampling coordination, sequencing and HT-qPCR data

acquisition: UK, TUB; 16S sequence analysis: GG, EC; HT-qPCR analysis: UK; Data curation

and validation: UK, GG, EC; Correlation and network analysis: UK, GG, EC, DK, PF; Data

interpretation: UK, GG, EC, HB, CM, TUB; Visualization of data: UK, GG, EC, DK, PF; Funding

acquisition: CC, NK, MP, FW, MW, HB, CM, TUB; Supervision: UK, CC, NK, MP, JV, FW, MW,

HB, CM, TUB; Writing - original draft: UK, GG, EC; Writing - review and editing: all authors. All

authors have read and approved the final version of the manuscript.

Acknowledgements

We give a special thanks to Rosi Siber for creating the GIS Map (Figure S1). We thank all the

local experts on soils and rivers in the different countries for advice regarding sample location

identification.

References

1. Laxminarayan R, Duse A, Wattal C, Zaidi AKM, Wertheim HFL, Sumpradit N, et al. Antibiotic

resistance—the need for global solutions. Lancet Infect Dis. Elsevier; 2013;13:1057–98.

2. WHO. Antimicrobial Resistance Global Report on Surveillance. 2014.

3. Hernando-Amado S, Coque TM, Baquero F, Martínez JL. Defining and combating antibiotic

resistance from One Health and Global Health perspectives. Nat Microbiol. Nat Microbiol;

2019;4:1432–42.

4. WHO. Global Action Plan on Antimicrobial Resistance. Microbe Mag. Switzerland;

2015;10:354–5.

5. Berendonk TU, Manaia CM, Merlin C, Fatta-Kassinos D, Cytryn E, Walsh F, et al. Tackling

antibiotic resistance: the environmental framework. Nat Rev Microbiol. Nature Publishing Group;

2015;13:310–7.

6. Smalla K, Cook K, Djordjevic SP, Klümper U, Gillings M. Environmental dimensions of

antibiotic resistance: assessment of basic science gaps. FEMS Microbiol Ecol. 2018;94.

7. D'Costa VM, King CE, Kalan L, Morar M, Sung WWL, Schwarz C, et al. Antibiotic resistance

is ancient. Nature. Nature Publishing Group, a division of Macmillan Publishers Limited. All

Rights Reserved.; 2011;477:457–61.

8. Rizzo L, Manaia C, Merlin C, Schwartz T, Dagot C, Ploy MCC, et al. Urban wastewater

treatment plants as hotspots for antibiotic resistant bacteria and genes spread into the

environment: A review. Sci Total Environ. Elsevier; 2013;447:345–60.

9. Kampouris ID, Klümper U, Agrawal S, Orschler L, Cacace D, Kunze S, et al. Treated

wastewater irrigation promotes the spread of antibiotic resistance into subsoil pore-water.

Environ Int. Pergamon; 2021;146:106190.

10. Heuer H, Schmitt H, Smalla K. Antibiotic resistance gene spread due to manure application

on agricultural fields. Curr Opin Microbiol. 2011;14:236–43.

11. Kampouris IDID, Agrawal S, Orschler L, Cacace D, Kunze S, Berendonk TUTU, et al.

Antibiotic resistance gene load and irrigation intensity determine the impact of wastewater

irrigation on antimicrobial resistance in the soil microbiome. Water Res. Pergamon;

2021;193:116818.

12. Kormos D, Lin K, Pruden A, Marr LC. Critical review of antibiotic resistance genes in the

atmosphere. Environ Sci Process Impacts. The Royal Society of Chemistry; 2022;24:870–83.

13. Gao M, Zhang X, Yue Y, Qiu T, Wang J, Wang X. Air path of antimicrobial resistance related

genes from layer farms: Emission inventory, atmospheric transport, and human exposure. J

Hazard Mater. Elsevier; 2022;430:128417.

14. Plaza-Rodríguez C, Alt K, Grobbel M, Hammerl JA, Irrgang A, Szabo I, et al. Wildlife as

Sentinels of Antimicrobial Resistance in Germany? Front Vet Sci. Frontiers Media S.A.;

2021;7:1251.

15. Vittecoq M, Godreuil S, Prugnolle F, Durand P, Brazier L, Renaud N, et al. Antimicrobial

resistance in wildlife. J Appl Ecol. John Wiley & Sons, Ltd; 2016;53:519–29.

16. Jarma D, Sánchez MI, Green AJ, Peralta-Sánchez JM, Hortas F, Sánchez-Melsió A, et al.

Faecal microbiota and antibiotic resistance genes in migratory waterbirds with contrasting

habitat use. Sci Total Environ. Elsevier; 2021;783:146872.

17. Mallon CA, Elsas JD van, Salles JF. Microbial invasions: the process, patterns, and

mechanisms. Trends Microbiol. 2015;23:719–29.

18. Kinnunen M, Dechesne A, Proctor C, Hammes F, Johnson D, Quintela-Baluja M, et al. A

conceptual framework for invasion in microbial communities. ISME J 2016 1012. Nature

Publishing Group; 2016;10:2773–9.

19. Hector A, Dobson K, Minns A, Bazeley-White E, Lawton JH. Community diversity and

invasion resistance: An experimental test in a grassland ecosystem and a review of comparable

studies. Ecol Res 2001 165. Springer; 2001;16:819–31.

20. Mallon CA, Le Roux X, Van Doorn GS, Dini-Andreote F, Poly F, Salles JF. The impact of

failure: Unsuccessful bacterial invasions steer the soil microbial community away from the

invader's niche. ISME J. Nature Publishing Group; 2018;12:728–41.

21. Kinnunen M, Dechesne A, Albrechtsen HJ, Smets BF. Stochastic processes govern

invasion success in microbial communities when the invader is phylogenetically close to

resident bacteria. ISME J 2018 1211. Nature Publishing Group; 2018;12:2748–56.

22. Altman S, Whitlatch RB. Effects of small-scale disturbance on invasion success in marine

communities. J Exp Mar Bio Ecol. Elsevier; 2007;342:15–29.

23. Finke DL, Snyder WE. Niche partitioning increases resource exploitation by diverse

communities. Science (80-).   American Association for the Advancement of Science   ;

2008;321:1488–90.

24. Bellanger X, Guilloteau H, Bonot S, Merlin C. Demonstrating plasmid-based horizontal gene

transfer in complex environmental matrices: A practical approach for a critical review. Sci Total

Environ. Elsevier; 2014;493:872–82.

25. Coyte KZ, Schluter J, Foster KR. The ecology of the microbiome: Networks, competition,

and stability. Science. Science; 2015;350:663–6.

26. Chen QL, An XL, Zheng BX, Gillings M, Peñuelas J, Cui L, et al. Loss of soil microbial

diversity exacerbates spread of antibiotic resistance. Soil Ecol Lett 2019 11. Springer; 2019;1:3–

13.

27. Bagra K, Bellanger X, Merlin C, Singh G, Berendonk TU, Klümper U. Exposure to

environmental stress decreases the resistance of river microbial communities towards invasion

by antimicrobial resistant bacteria. bioRxiv. Cold Spring Harbor Laboratory;

2022;2022.11.19.517188.

28. Dai D, Brown C, Bürgmann H, Larsson DGJ, Nambi I, Zhang T, et al. Long-read

metagenomic sequencing reveals shifts in associations of antibiotic resistance genes with

mobile genetic elements from sewage to activated sludge. Microbiome. BioMed Central Ltd;

2022;10:1–16.

29. Liu Z, Klümper U, Liu Y, Yang Y, Wei Q, Lin JG, et al. Metagenomic and metatranscriptomic

analyses reveal activity and hosts of antibiotic resistance genes in activated sludge. Environ Int.

Pergamon; 2019;129:208–20.

30. Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Lozupone CA, Turnbaugh PJ, et al.

Global patterns of 16S rRNA diversity at a depth of millions of sequences per sample. Proc Natl

Acad Sci U S A. National Academy of Sciences; 2011;108:4516–22.

31. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. DADA2: High-

resolution sample inference from Illumina amplicon data. Nat Methods. Nature Publishing

Group; 2016;13:581–3.

32. Bolyen E, Rideout JR, Dillon MR, Bokulich NA, Abnet CC, Al-Ghalith GA, et al.

Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. Nat

Biotechnol 2019 378. Nature Publishing Group; 2019;37:852–7.

33. Stedtfeld RD, Guo X, Stedtfeld TM, Sheng H, Williams MR, Hauschild K, et al. Primer set

2.0 for highly parallel qPCR array targeting antibiotic resistance genes and mobile genetic

elements. FEMS Microbiol Ecol. Oxford Academic; 2018;94.

34. Stachler E, Bibby K. Metagenomic Evaluation of the Highly Abundant Human Gut

Bacteriophage CrAssphage for Source Tracking of Human Fecal Pollution. Environ Sci Technol

Lett. American Chemical Society; 2014;1:405–9.

35. Karkman A, Pärnänen K, Larsson DGJ. Fecal pollution can explain antibiotic resistance

gene abundances in anthropogenically impacted environments. Nat Commun. Nature

Publishing Group; 2019;10:80.

36. Fang P, Xiao P, Tan F, Mo Y, Chen H, Klümper U, et al. Biogeographical Patterns of

Bacterial Communities and Their Antibiotic Resistomes in the Inland Waters of Southeast

China. Microbiol Spectr. American Society for Microbiology; 2022;10.

37. Zhu YG, Johnson TA, Su J-Q, Qiao M, Guo G-X, Stedtfeld RD, et al. Diverse and abundant

antibiotic resistance genes in Chinese swine farms. PNAS. National Academy of Sciences;

2013;110:3435–40.

38. Schmittgen TD, Livak KJ. Analyzing real-time PCR data by the comparative CT method. Nat

Protoc 2008 36. Nature Publishing Group; 2008;3:1101–8.

39. Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, et al. The SILVA ribosomal

RNA gene database project: Improved data processing and web-based tools. Nucleic Acids

Res. Oxford University Press; 2013;41:D590-6.

40. Bray JR, Curtis JT. An ordination of the upland forest communities of Southern Wisconsin.

Ecol Monogr. Wiley-Blackwell; 1957;27:325–49.

41. Gu Z. Complex heatmap visualization. iMeta. John Wiley & Sons, Ltd; 2022;1:e43.

42. Kembel SW, Cowan PD, Helmus MR, Cornwell WK, Morlon H, Ackerly DD, et al. Picante: R

tools for integrating phylogenies and ecology. Bioinformatics. Oxford Academic; 2010;26:1463–

4.

43. Team RC. R: A language and environment for statistical computing. R Found Stat Comput

Vienna, Austria. 2013;

44. Acosta F, Zamor RM, Najar FZ, Roe BA, Hambright KD. Dynamics of an experimental

microbial invasion. Proc Natl Acad Sci U S A. National Academy of Sciences; 2015;112:11594–

9.

45. Cacace D, Fatta-Kassinos D, Manaia CM, Cytryn E, Kreuzinger N, Rizzo L, et al. Antibiotic

resistance genes in treated wastewater and in the receiving water bodies: A pan-European

survey of urban settings. Water Res. Elsevier Ltd; 2019;162:320–30.

46. Munk P, Brinch C, Møller FD, Petersen TN, Hendriksen RS, Seyfarth AM, et al. Genomic

analysis of sewage from 101 countries reveals global landscape of antimicrobial resistance. Nat

Commun. Nature Publishing Group; 2022;13:7251.

47. Wolters B, Widyasari-Mehta A, Kreuzig R, Smalla K. Contaminations of organic fertilizers

with antibiotic residues, resistance genes, and mobile genetic elements mirroring antibiotic use

in livestock? Appl Microbiol Biotechnol. Springer Berlin Heidelberg; 2016;100:9343–53.

48. Keenum I, Liguori K, Calarco J, Davis BC, Milligan E, Harwood VJ, et al. A framework for

standardized qPCR-targets and protocols for quantifying antibiotic resistance in surface water,

recycled water and wastewater. https://doi.org/101080/1064338920212024739. Taylor &

Francis; 2022;52:4395–419.

49. Abramova A, Berendonk TU, Bengtsson-Palme J. Meta-analysis reveals the global picture

of antibiotic resistance gene prevalence across environments. bioRxiv. Cold Spring Harbor

Laboratory; 2022;2022.01.29.478248.

50. Uroz S, Buée M, Deveau A, Mieszkin S, Martin F. Ecology of the forest microbiome:

Highlights of temperate and boreal ecosystems. Soil Biol Biochem. Pergamon; 2016;103:471–

88.

51. Worischka S, Schöll F, Winkelmann C, Petzoldt T. Twenty-eight years of ecosystem

recovery and destabilisation: Impacts of biological invasions and climate change on a temperate

river. Sci Total Environ. Elsevier; 2023;875:162678.

52. Baquero F, Coque TM, Galán JC, Martinez JL. The Origin of Niches and Species in the

Bacterial World. Front Microbiol. Frontiers Media S.A.; 2021;12:566.

53. Mattingly WB, Hewlate R, Reynolds HL, Mattingly WB, Hewlate R, Reynolds HL. Species

evenness and invasion resistance of experimental grassland communities. Oikos. John Wiley &

Sons, Ltd; 2007;116:1164–70.

54. Fox BJ. Niche Parameters and Species Richness. Ecology. John Wiley & Sons, Ltd;

1981;62:1415–25.

55. Furness EN, Garwood RJ, Mannion PD, Sutton MD. Productivity, niche availability, species

richness, and extinction risk: Untangling relationships using individual-based simulations. Ecol

Evol. John Wiley & Sons, Ltd; 2021;11:8923–40.

56. Battin TJ, Besemer K, Bengtsson MM, Romani AM, Packmann AI. The ecology and

biogeochemistry of stream biofilms. Nat Rev Microbiol 2016 144. Nature Publishing Group;

2016;14:251–63.

57. Klümper U, Riber L, Dechesne A, Sannazzarro A, Hansen LH, Sørensen SJ, et al. Broad

host range plasmids can invade an unexpectedly diverse fraction of a soil bacterial community.

ISME J. International Society for Microbial Ecology; 2015;9:934–45.

58. Klümper U, Dechesne A, Riber L, Brandt KK, Gülay A, Sørensen SJ, et al. Metal stressors

consistently modulate bacterial conjugal plasmid uptake potential in a phylogenetically

conserved manner. ISME J. International Society for Microbial Ecology; 2017;11:152–65.

59. Li L, Dechesne A, He Z, Madsen JS, Nesme J, Sørensen SJ, et al. Estimating the transfer

range of plasmids encoding antimicrobial resistance in a wastewater treatment plant microbial

community. Environ Sci Technol Lett. American Chemical Society; 2018;5:260–5.

60. Song J, Klümper U, Riber L, Dechesne A, Smets BF, Sørensen SJ, et al. A converging

subset of soil bacterial taxa is permissive to the IncP-1 plasmid pKJK5 across a range of soil

copper contamination. FEMS Microbiol Ecol. 2020;96.

61. Wang Y, Yu Z, Ding P, Lu J, Klümper U, Murray AK, et al. Non-antibiotic pharmaceuticals

promote conjugative plasmid transfer at a community-wide level. Microbiome. BioMed Central

Ltd; 2022;10:1–15.

62. Hall JPJ, Wood AJ, Harrison E, Brockhurst MA. Source-sink plasmid transfer dynamics

maintain gene mobility in soil bacterial communities. Proc Natl Acad Sci U S A. National

Academy of Sciences; 2016;113:8260–5.

63. Kottara A, Hall JPJ, Brockhurst MA. The proficiency of the original host species determines

community-level plasmid dynamics. FEMS Microbiol Ecol. Oxford Academic; 2021;97:26.

64. Kottara A, Carrilero L, Harrison E, Hall JPJ, Brockhurst MA. The dilution effect limits plasmid

horizontal transmission in multispecies bacterial communities. Microbiol (United Kingdom).

Microbiology Society; 2021;167:1086.

65. Klümper U, Recker M, Zhang L, Yin X, Zhang T, Buckling A, et al. Selection for antimicrobial

resistance is reduced when embedded in a natural microbial community. ISME J. Nature

Publishing Group; 2019;13:2927–37.

66. Walker-Sünderhauf D, Klümper U, Gaze WH, Westra ER, Houte S van. Interspecific

competition can drive the loss of conjugative plasmids from a focal species in a microbial

community. bioRxiv. Cold Spring Harbor Laboratory; 2022;2022.06.06.494957.

67. Taberlet P, Zimmermann NE, Englisch T, Tribsch A, Holderegger R, Alvarez N, et al.

Genetic diversity in widespread species is not congruent with species richness in alpine plant

communities. Ecol Lett. John Wiley & Sons, Ltd; 2012;15:1439–48.

68. Lawrence ER, Fraser DJ. Latitudinal biodiversity gradients at three levels: Linking species

richness, population richness and genetic diversity. Glob Ecol Biogeogr. John Wiley & Sons,

Ltd; 2020;29:770–88.