**Delineating the functional role of the *PPE50 (Rv3135) - PPE51 (Rv3136)* gene cluster in the pathophysiology of *Mycobacterium tuberculosis***

Ravi Prasad Mukku[1], Kokavalla Poornima[1], Sangya Yadav[1a], and Tirumalai R. Raghunand[1,2*]

[1]CSIR - Centre for Cellular and Molecular Biology, Uppal Road, Hyderabad 500007, India

[2]Academy of Scientific and Innovative Research (AcSIR), Ghaziabad 201002, India

**Email information:**

Ravi Prasad Mukku - raviprasad.mukku@ccmb.res.in

Kokavalla Poornima - poornima.kchowdary@gmail.com

Sangya Yadav - sangyayadav20@gmail.com

Tirumalai R. Raghunand - raghu@ccmb.res.in. *Correspondence and reprints

[a]Current Address: University of Colorado, Anschutz Medical Campus, 13001 E 17th Pl, Aurora, CO 80045, USA

**Keywords**: *Mycobacterium tuberculosis*; PE_PPE family; Macrophages; TLR1; Immunomodulation, immune surveillance

**ABSTRACT**

The extraordinary success of *Mycobacterium tuberculosis* (*M. tb*) has been attributed to its ability to modulate host immune responses. The genome of *M. tb* encodes multiple immunomodulatory factors, including several proteins of the multigenic PE_PPE family, which comprise about 10% of its coding potential. The presence of these proteins in pathogenic mycobacteria strongly suggests that they play a role in disease pathogenesis. To understand its role in *M. tb* physiology we have begun to characterise the *PPE50 (Rv3135)-PPE51 (Rv3136)* gene cluster, one of nine *PPE-PPE* clusters in the *M. tb* genome. We demonstrate that this cluster encodes a co-transcriptional unit and that PPE50 and PPE51 interact both *in vitro* and *in vivo*, the first demonstration of PPE-PPE interaction. THP-1 macrophages infected with recombinant *M. smegmatis* strains expressing *PPE50* and *PPE51* showed less intracellular viability than the control strain containing the vector alone, the decline in viable counts correlating with an increase in transcript levels of inducible nitric oxide synthase (*iNOS2*). Macrophages infected with the recombinant strains exhibited an upregulation in levels of the anti-inflammatory cytokine *IL-10*, indicating an immunomodulatory role for these proteins. Using pull-down assays, we discovered TLR1 to be the cognate receptor for PPE50, with signalling through the receptor being indicated by an increase in IRAK1 phosphorylation. All the phenotypes observed on infection of THP-1 macrophages including the decrease in CFUs, the increase in *iNOS2* and *IL-10* levels, as well as signalling through the receptor, were reversed on treatment of macrophages with an anti-TLR1 antibody prior to infection, validating the functional outcome of PPE50-TLR1 interaction. Our data points to a TLR1 dependent role for the *PPE50-PPE51* cluster in promoting bacillary persistence, *via* CFU reduction and a concomitant upregulation of the anti-inflammatory response - a two-pronged strategy to circumvent host immune surveillance.

**INTRODUCTION**

*M. tb*, the causative agent of human tuberculosis, has achieved unprecedented success due to its ability to modify host immunological responses, thereby promoting its long-term survival (1). Understanding the pathogenic processes of *M. tb* depends on identifying and characterizing bacillary elements involved in immune evasion (2), and their interaction with host defense components during infection. In this context, the multigenic PE/PPE proteins, which account for 10% of the coding potential of the *M. tb* genome, have emerged as key players because several family members have been linked to evasion of host immune processes. These proteins are named after the conserved Proline-Glutamate and Proline-Proline-Glutamate residues at their N-termini (3, 4), and the *M. tb* genome is reported to encode 169 PE/PPE proteins. A wide variety of *PE/PPE* genes have been discovered within pathogenic Mycobacteria. Most of them are arranged sporadically in the genome, and a few are arranged in clusters. *M. tb* has total of 40 such *PE/PPE* gene clusters, with loci including *PE-PPE* [15], *PE-PE* [5] and *PPE-PPE* [9] genes arrayed in tandem; among them are uncommon clusters like a *PPE* gene followed by another *PPE*, and a *PE* gene sandwiched between two *PPE* genes (3). Putative operonic clusters containing only *PE/PPE* genes are found for 22 of the 69 PPE proteins (3, 5). The 14 gene pairs with a 5' *PE-PPE* 3' orientation are separated by 100 bases or less, and are likely to be co-transcribed and be functionally linked (6). Evidence from comparative genomics suggests that the expansion of the *PE/PPE* family is linked to the five *Esx* clusters spread out over the *M. tb* genome, which encode components of the Type VII secretion system. In contrast to the extensive studies on the PE_PGRS sub-group in the literature, no studies have been conducted on the role of *PPE-PPE* clusters in *M. tb* pathophysiology. We have selected the *Rv3135* (*PPE50*) - *Rv3136* (*PPE51*) cluster for its functional characterisation and to understand the involvement of the two genes in *M. tb* physiology, owing to their peculiar cluster arrangement. In a microarray study, these genes were observed to be upregulated on exposure to palmitic acid. Repression in expression of both genes was seen on exposure of *M. tb* to DETA/NO, $H_2O_2$, KCN, and under stationary phase growth conditions (7). More recent reports have suggested a role for PPE51 in drug tolerance (8), nutrient transport (9), and in the inhibition of autophagy (10). No functional studies have yet been described for the individual genes, or for the gene cluster as a whole.

## MATERIALS AND METHODS

### Bacterial strains, media and growth conditions

*M. smegmatis* mc$^2$6 and *M. tb* H37Rv were cultivated in Middlebrook, 7H9 broth and 7H10 agar (Difco) containing albumin dextrose complex (5 g BSA, 2 g glucose, and 0.85 g NaCl/L), 0.5% (v/v) glycerol, and 0.5% (v/v) Tween 80. Luria Bertani medium was used to cultivate *E. coli* strains - both *E. coli* and mycobacteria were cultured by shaking at 37° C. When required, 200 µg/ml of ampicillin and 50 µg/ml of kanamycin (for *E. coli*) and 15 µg/ml of kanamycin (for mycobacteria) were added to the culture medium.

### DNA techniques

All restriction enzymes, T4 DNA ligase, and Taq polymerase were purchased from New England Biolabs (NEB) and Invitrogen, respectively. DNA manipulation protocols including plasmid DNA preparation, restriction endonuclease digestion, agarose gel electrophoresis, ligation of DNA fragments and *E. coli* transformation were performed according to (11). The PCR amplifications were performed according to the manufacturer's instructions. During each cycle, 95° C was held for 30 s, 60° C for 30 s, and 72° C for 1 min, followed by a 72° C extension cycle for 10 min. The Qiagen gel extraction kit was used to purify DNA fragments for cloning reactions according to the manufacturer's instructions. *M. smegmatis* was transformed by electroporation.

### *In silico* analyses

Multiple Sequence Alignments were performed using the ClustalW2 algorithm(12). All *M. tb* sequences were obtained from the Mycobrowser database (https://mycobrowser.epfl.ch). The TM-pred (13) and Kyte& Doolittle (14) algorithms were used to identify trans-membrane domains and regions of hydrophobicity respectively.

### Sub-cellular localisation and Proteinase K Sensitivity Assay

In order to determine the subcellular location of these proteins, the *M. tb PPE50*, *PPE51* ORFs were cloned between *BamHI* and *EcoRI* sites of pJEX55 (15) and the recombinant plasmids were transformed into *M. smegmatis*. The recombinant *M.*

*smegmatis* strains expressing c-myc tagged PPE50, PPE51 were harvested at the

logarithmic phase of growth, washed and resuspended in PBS. The samples were

divided into two identical aliquots and incubated at 37°C for 30 min with or without

Proteinase K (Sigma). With the addition of 2 mM EGTA, the reaction was stopped,

and the sub-cellular fractions of these samples were isolated as described in (16). By

denaturing SDS PAGE, the individual fractions were separated, and fusion proteins

were detected by western blotting with anti-c-myc antibodies (sc40, Santa Cruz).

**Expression of PPE50, PPE51 in *M. smegmatis***

*M. tb PPE50, PPE51* ORFs were amplified using gene-specific primers (Table S1)

and cloned between the *BamHI* and *EcoRI* sites of pMV261(17) These constructs

were transformed into *M. smegmatis* for overexpression.

**Expression and purification of PPE50 and PPE51**

The full-length *M. tb PPE50* ORF was cloned as C-terminal 6xHis-tagged fusion in

pET22b, and *PPE51* was cloned as N-terminal Glutathione-S-Transferase

(GST) tagged fusion in pGEX6p1. Following sequence confirmation, the constructs

were transformed into *E. coli* BL21(DE3). For induction, cultures were allowed to

grow to an OD of 0.5, induced with 0.5 mM IPTGm following which PPE50 was

purified using Ni-NTA affinity chromatography, and PPE51 was purified using

glutathione agarose beads. Purified proteins were used for pull-down experiments.

***In vitro* growth kinetics**

To functionally characterise *M. tb PPE50, PPE51* their ORFs were amplified from *M.*

*tb* H37Rv genomic DNA using gene specific primers (Table S1), cloned between the

*BamHI* and *EcoRI* sites of pMV261 (18) and transformed into *M.smegmatis*. To

examine their growth patterns, recombinant *M. smegmatis* strains were grown until

late exponential phase, diluted to an $A_{600}$ (OD) of 0.2 and cultured in Middlebrook

7H9 broth containing 15 µg/ml kanamycin. Growth curves were generated by CFU

measurements and plotted against time. At each designated time point, cultures

were harvested for RNA extraction and gene expression analyses performed. All

CFU enumeration assays were performed in the presence of 15 µg/ml kanamycin.

**Macrophage infection**

THP-1 monocytes were cultured at 37° C in 5% $CO_2$ in RPMI 1640 medium
supplemented with 10% (v/v) Fetal bovine serum and antibiotics (60 µg/ml penicillin
G sodium, 50 µg/ml streptomycin sulphate, and 30 µg/ml gentamycin sulphate). For
the infection experiments, monocytes were seeded at a density of $2x10^6$/well for 24
h, differentiated with 5 ng/ml PMA, and then infected 72 h later. Cultures of
exponentially growing bacteria were pelleted, washed, resuspended in RPMI
medium (without antibiotics) to an OD of 1.0. Single cell suspensions of
recombinant *M. smegmatis* strains were prepared by passing culture suspensions
through needles of 26 1/2 gauge 5-6 times. At each step, CFU counts were
performed to determine bacterial viability. Based on pilot infections with multiple
MOIs we performed with the cell lines used, equal numbers of each strain were used
for infections (Input counts) at a MOI of 1:100. Post-infection CFU counts were
determined by lysis of infected cells after 2 h of incubation with bacteria (T0
counts). In order to kill extracellular bacteria, complete RPMI containing gentamycin
was added. Infected cells were lysed with 0.1% Triton X-100 and plated on
Middlebrook 7H10 agar at the designated time points to determine CFU counts. For
each experiment, macrophages infected with *M. smegmatis* expressing the empty
vector were used as controls.

**Real-time RT-PCR analysis**

The expression profiles of *PPE50*, *PPE51* in *M. tb* and recombinant strains of *M.*
*smegmatis* expressing pMV261-*PPE50*, pMV261-*PPE51* as a function of growth,
were determined using real-time transcript measurements. For this, cells were
harvested at the OD 0.8, 1.5, 2, 3, 3.5 and at 4, 8, 12, 24, 48 and 72 h time points
respectively, and total RNA isolated from each culture using TRIzol reagent
(Invitrogen) as per the manufacturer's protocol. Following treatment with RNAse free
DNAse I, cDNA synthesis was performed using the iScript cDNA synthesis kit (Bio-
Rad) and subsequently used as a template for SYBR green based PCR amplification
using *PPE50*, *PPE51* specific primers (Table S1) to generate 200 bp amplicons.
Gene specific transcript levels in *M. smegmatis* were normalised to the *sigA*
transcript in each sample and in *M. tb* to the 16s rRNA transcript. In *M. smegmatis*

the relative fold change in transcript levels at each time point was calculated with
respect to the levels at 4 h. In the *M. tb* expression profile, the relative fold change in
transcript at each OD point was calculated with respect to the levels at OD 0.8. To
quantify host cytokines and *iNOS2* transcript levels, total RNA was isolated from
infected macrophages using TRIzol reagent. Following treatment with RNAse free
DNAse I, cDNA synthesis was performed using the iScript cDNA synthesis kit (Bio-
Rad) and subsequently used as a template for SYBR green based PCR amplification
using primers specific to *iNOS*, *IL-10*, *IL-12* (Table S1) designed to generate 200 bp
amplicons. The levels of each mRNA were normalised to the transcript levels of
GAPDH and β-actin. Relative fold change was calculated with reference to
macrophages infected with *M. smegmatis* expressing the empty vector.
202

**Receptor interaction assay**
To extract the membrane fraction for receptor identification, PMA-differentiated THP-
1 cells were resuspended in cell lysis buffer (50 mM HEPES buffer pH 7.4, 150 mM
NaCl, 5 mM EDTA, PIC), sonicated at 20 percent amplitude with 10 s on and off for 5
minutes, centrifuged at 13000 rpm for 1 h, and the pellet was resuspended in lysis
buffer containing 1% CHAPS. For co-immunoprecipitation assays, 50 μg of PPE50-
6xHis was combined with 500 μg of cell membrane fractions in lysis buffer to a final
volume of 500 μl. The reactions were incubated at $37^o$ C for 6 h, after which anti-
His antibodies (CST) were used to conduct pull-downs. Before adding antibodies, 60
μl of each sample was aliquoted for use as loading controls. Each sample was then
mixed with Protein A agarose beads and incubated overnight at 4° C. Following this,
after three washes with PBS, the beads were boiled in 50 μl of sample buffer,
resolved on SDS-PAGE. Following transfer onto PVDF membranes, the co-
immunoprecipitated proteins were identified using antibodies to TLR1/2/ 4/ 6. Equal
loading of the cell membrane fraction was evaluated using western blotting using
antibodies to CD81. In reverse pulldown assays we incubated PPE50 with the
membrane fraction along with a TLR1 Ab (Abcam) at $4^0$ C overnight, bound the mix
with Protein A beads, washed and resolved the bead bound proteins on SDS-PAGE
and transferred them onto PVDF membranes. The blot was then probed with an anti-
His antibody (CST).
223

**TLR1 Receptor inhibition**

THP-1 monocytes were grown in RPMI 1640 media supplemented with 10% (v/v)
Fetal bovine serum and antibiotics (60 µg/ml penicillin G sodium, 50 µg/ml
streptomycin sulphate, and 30 µg/ml gentamycin sulphate) at 37°C and 5% $CO_2$.
These monocytes were seeded at a density of $2x10^6$ per well for 24 h, differentiated
with 5 ng/ml PMA for 72 h, and then infected. Prior to infection, we treated these
differentiated cells for 6 with 0.5µg/ml of a polyclonal anti-TLR1 antibody (Abcam).
231

**Western blotting**

The levels of total and phosphorylated IRAK-1 in infected THP-1 macrophages were
determined by western blot analysis. Each sample was lysed in RIPA lysis buffer,
following which, equal protein amounts from each cell lysate subjected to SDS-
PAGE and transferred onto PVDF membranes. Each blot was sequentially probed
with antibodies specific to phosphorylated or total IRAK1 (Cell Signalling Technology
& Abcam) and β-actin. Proteins were detected using Enhanced Chemiluminescence
(Santa Cruz). For the inhibition studies, THP-1 cells were treated with 0.5 µg/ml anti-
TLR1 Ab (Abcam) 6 h prior to infection. Recombinant *M. smegmatis* containing
pMV261 was used as an empty vector/ mock control. For quantitation, the intensity
of all bands was densitometrically measured using ImageJ and normalised to levels
of β-actin. The normalised levels of phosphorylated IRAK1 in control samples
(pMV261) were assigned a value of 1, and the fold change in the levels of these
proteins in the test samples (PPE50, PPE51) were derived with respect to the
control.
247

**Co-transcriptional analysis**

249 Total RNA extracted from exponentially growing *M. tb* H37Rv was used to assess
250 the transcriptional state of the *PPE50-PPE51* cluster. RNA treated with DNAse I was
251 used as a template for cDNA synthesis with the primers *PPE51* R1 (Table S1). The
252 cDNA was utilised for PCR amplification with gene-specific primer combinations. The
253 analysis includes appropriate negative controls (-RT).
254

**Co-immunoprecipitation**

256 To perform *in vivo* pull down assays, 150 ml of independent *M. smegmatis* cultures
257 co-transformed with pJEX55+pSCW54, pJEX55-*PPE51*+pSCW54,
258 pJEX55+pSCW54-*PPE50*, pJEX55-*PPE51*+pSCW54-*PPE50* were grown in

Middlebrook 7H9 broth containing kanamycin and hygromycin to log phase and induced with 0.2% acetamide for 6 h. Induced cells were pelleted, resuspended in 1.2 ml PBS containing protease inhibitors and lysed by sonication. Half of this lysate was used to prepare a cell envelope fraction by ultracentrifugation (16) and the pellets were resuspended in 150 µl of PBS containing protease inhibitors and 0.5% CHAPS. 400 µl of cell lysates were mixed with 50 µl of the solubilised cell envelope fraction and the protein amounts in the mix were quantitated using a Bradford assay. For the co-IP assay 500 µg of protein in a total volume of 500 µl PBS was incubated with an anti c-myc antibody; prior to antibody addition, 30 µl of the reaction mix was removed and saved as a loading control. Protein A agarose beads were then added to each sample and incubated at 4°C for 6 h. Following three washes with PBS, the beads were boiled in 50 µl of sample buffer and subjected to SDS-PAGE. Separated proteins were transferred to PVDF and probed with an anti-His antibody.

**Mycobacterial Protein Fragment Complementation (MPFC) assay**

To assess the interaction of PPE50-PPE51 at *in vivo PPE50*, *PPE51* were cloned into the *E. coli - Mycobacterium* shuttle vectors pUAB400 and pUAB300 respectively. Both constructs were co-transformed in *M. smegmatis* mc$^2$155 and the interaction assay was performed as described (19). Appropriate positive (pUAB100+pUAB200) and negative (pUAB100+pUAB400, pUAB400+ pUAB300) controls were included in the assay.

**RESULTS**

***PPE50* (*Rv3135*) and *PPE51* (*Rv3136*) are co-operonic and interact with each other**

Among the 40 PE/PPE clusters in the *M. tb* genome, *PPE50-PPE51* is one of 9 such PPE-PPE clusters. The two proteins share 62% identity (71/114) and 77% homology (88/114) in their PPE domains with PPE51 (380 aa) possessing a C-terminal extension that PPE50 (132 aa) lacks (Fig. 1). Since *PPE50-PPE51* was among the nine *PPE-PPE* gene pairs expected to be co-operonic based on their co-repression in drug tolerance research, it was hypothesised that they would be cooperative (8). To confirm this hypothesis, RT-PCR analyses were performed with RNA isolated from exponentially growing *M. tb* H37Rv using primer pairs (Table S1) designed to amplify ORF and junction-specific areas of the gene pair (Fig. 2A) (*PPE50*-200 bp,

*PPE51*-200 bp, and *PPE50-PPE51* junction-200 bp). We observed amplification
products in the test lanes but not in the -RT control lanes, leading us to infer that this
gene pair was transcribed as a mono-cistronic message (Fig. 2B). Since co-operonic
protein pairs are known to often physically interact (20-22) the potential interaction of
the PPE50-PPE51 protein pair was investigated by performing co-
immunoprecipitation of C-terminally His-tagged PPE50 with N-terminally GST-tagged
PPE50 (Fig. S1)  along with the appropriate controls. Co-precipitation of PPE51 and
PPE50 suggested these two proteins directly interact *in vitro* (Fig. 2C). To verify this
finding, *in vivo* pull-down experiments were performed utilising cell wall enriched cell
lysates of *M. smegmatis* expressing C-terminal c-myc fused PPE51 and C-terminal
6xHis-tagged PPE50. Co-precipitation of PPE50 and PPE51 suggested that the two
proteins interact *in vivo* as well (Fig. 2D). We ensured that in both the *in vitro* and *in*
*vivo* pull-down analyses, equal amounts of each fraction were loaded (Fig. S2). In
addition, the mycobacterial protein fragment complementation (M-PFC) assay was
used to examine the interaction between these proteins (19). Co-transformants of *M.*
*smegmatis* mc$^2$155 expressing two protein pairs fused to the murine dihydrofolate
reductase domains F [1,2] and F [3] were resistant to 40 g/ml trimethoprim (Figure 2
E). Negative control samples exhibited no growth, validating the *in vivo* interaction
between PPE50 and PPE51.

**M. smegmatis strains expressing *PPE50* and *PPE51* show reduced survival in**
**macrophages**
To investigate the possible involvement of PPE50 and PPE51 in mycobacterial
pathophysiology, we used the saprophyte *M. smegmatis,* which has been
extensively used as a surrogate model to study the roles of this family of proteins,
including by our own research group (20, 21, 23-27). Since it lacks the homologues
of most PE-PPE proteins, including *PPE50* and PPE51, we expressed these proteins
in *M. smegmatis* and examined their localisation. To do so, recombinant *M.*
*smegmatis* strains expressing c-myc tagged PPE50 and PPE51 were treated with
Proteinase K, and sub-cellular fractionation was carried out. Prior to western blotting
with a c-myc antibody, we ensured equal loading for each fraction (Fig. S3A, B). The
blots revealed that both proteins were localized to the cell wall and, that their signals
were greatly diminished in the cell wall-associated fractions treated with Proteinase
K (Fig. S34, B). This indicated that PPE50 and PPE51 are exposed to the cell

surface, making this a suitable system to evaluate their functions as modulators of
pathogenicity *via* contact with the host. This finding was consistent with the predicted
hydrophobicity and transmembrane domain prediction of these proteins generated *in*
*silico* (Fig. S5A, B). *M. smegmatis* strains expressing untagged PPE50 and PPE51
under the control of the constitutive *hsp60* promoter were then used to determine
whether these proteins mediate host-pathogen interactions (17, 18). CFU
enumeration and growth profiles (Fig. S5A, B) of recombinant *M. smegmatis* strains
expressing *PPE50*, *PPE51*, *PPE50-PPE51*, and the empty vector revealed that over-
expression of these genes did not result in growth compromise *in vitro*. Both genes
were expressed at comparable levels in these strains as quantified by real-time RT-
PCR using gene-specific primers (Table S1), suggesting that their expression did not
vary as a function of growth (Fig. S6C, D, E). Using the THP-1 macrophage infection
model, we assessed the intracellular survival of the recombinant *M. smegmatis*
strains, and the ability of the proteins to modulate host functions. Curiously, bacillary
counts of strains expressing *PPE50*, *PPE51*, and *PPE50-PPE51* were lower than
those of the empty vector control at 24, 48, and 72 hours post-infection in THP-1
macrophages (Fig. 3A), Though unexpected, this observation correlated with the
elevated levels of inducible nitric oxide synthase-2 (*iNOS2*) expression in these cells
(Fig. 3B). This is significant since iNOS2 is an important factor in the control of
intracellular bacillary survival (28-30).
347

**PPE50 and PPE51 are immunomodulators that promote IL-10 production**
**through TLR1 signalling**
Our observation that PPE50 and PPE51 are engaged in reducing intra-macrophage
bacterial survival, led us to explore if they also affect immunological responses, as
has been observed for other members of this protein family (5, 20, 21, 25, 31). To
test their potential immune modulatory function, we infected THP-1 macrophage with
recombinant *M. smegmatis* strains expressing *PPE50*, *PPE51* and *PPE50-PPE51*,
and examined the levels of pro- and anti-inflammatory cytokines in infected cells. As
shown in Fig. 4A, we observed an upregulation of the anti-inflammatory cytokine IL-
10 in macrophages infected with recombinant *M. smegmatis* expressing *PPE50* at
24 h,48 h and 72 h post-infection, and the strain expressing *PPE50-PPE51* at the 24
h time point, in comparison to controls. Working on the hypothesis that this induction
must happen *via* ligand-receptor interaction, we used purified PPE50 and PPE51

(Fig. S1) to identify their cognate receptors. In pull-down and reverse pull-down

experiments with macrophage membrane fractions, we identified TLR1 to be the

binding partner for PPE50 (Fig. 4B, C); in the same experiments, no TLR interaction

could be identified for PPE51 (data not shown). Equal loading was ensured for all

fractions in these experiments, by quantifying levels of CD81 (Fig. S7). To test the

functionality of PPE50:TLR1 interaction,  we monitored IRAK1 phosphorylation as a

marker for  induction of TLR1 signalling (32), in macrophages infected with the

above-listed recombinant *M. smegmatis* strains. We observed an increase in

phosphorylation of IRAK-1 in THP-1 macrophages infected with the recombinant

PPE50-expressing *M. smegmatis* strain, implying a functional consequence for

PPE50:TLR1 interaction; no induction of IRAK-1 phosphorylation was seen in the

other two infection samples (Fig. 4D, E). To test if the induction of *iNOS2* and *IL-10*

we observed in earlier experiments was mediated by PPE50:TLR1 interaction, we

performed TLR1 blocking experiments by incubating macrophages with an anti-TLR1

antibody, prior to infection with the *PPE50* expressing recombinant *M. smegmatis*

strain. As depicted in figure 5, all the phenotypes associated with infection of

macrophages with recombinant *M. smegmatis* expressing *PPE50*, were reversed on

TLR1 blocking compared to the controls. These included the reduction in viability

(Figure 5A, B), the increase in *iNOS2* transcripts levels (Figure 5C, D), as well as the

elevation in the *IL-10* transcript (Figure 5E, F). Moreover, the increase in IRAK1

phosphorylation was lost on antibody blocking as well (Figure 9G, H), clearly

indicating that PPE50 activates the IRAK1 pathway through TLR1 to upregulate the

expression of *iNOS2* and *IL-10*.

**DISCUSSION**

While there is experimental evidence for the individual PE and PPE proteins, and PE

proteins encoded in operons, in the modulation of host immunity and pathogenesis

(20, 21, 33), there are no reports examining the involvement of the unusual

PPE/PPE clusters in *M. tb* pathogenesis. Despite long-standing evidence for *PPE50*

upregulation under conditions of hypoxia (7), and regulation of *PPE51* expression by

the dormancy and hypoxia-induced DevRS TCS  (2), the potential role of the *PPE50-*

*PPE51* cluster in the pathophysiology of *M. tb* had not been investigated thus far.

Our RT-PCR analysis revealed that *PPE50* and *PPE51* are expressed as one
transcriptional unit. Although this could have been predicted from the genomic
coordinates with only 61 bp separating *PPE50* and *PPE51* (genomic coordinates
3501794 -3502936), our data confirmed their operonic status. This observation was
further validated by performing growth dependent expression analyses of these
genes in *M.tb* H37Rv, where the two genes were seen to be coordinately expressed
in all growth phases from log to stationary (Fig S8). Based on our experience of
finding interaction between co-operonic members of this protein family (20, 21), we
tested this possibility, and established using both genetic and biochemical
approaches, that PPE50 and PPE51 indeed interact, the first record of a PPE-PPE
heterodimeric pair. This opened up the possibility that the two proteins might play a
role in *M. tb* physiology both independently as well as in a complex. To examine their
possible role in mediating interactions at the mycobacterium-macrophage interface,
we used the *M. smegmatis* model of macrophage infection that our laboratory has
used extensively to interrogate the role of this family of proteins in host-pathogen
interactions (20, 21, 34, 35). On infecting THP-1 macrophages with recombinant *M.*
*smegmatis* expressing *PPE50*, *PPE51*, and the entire cluster, we observed that all
these strains showed reduced survival compared to the control strain expressing the
empty vector. We were surprised with this observation since the predominant
phenotypes of recombinant *M. smegmatis* expressing the PE-PPE class of proteins
in macrophages is that of enhanced survival compared to controls. Since this
phenotype was reproducible and the reduced CFU counts correlated with an
enhancement in *iNOS2* levels, we were convinced that this finding was genuine. We
searched to see if any other such observation existed in the literature, and
discovered that *M. smegmatis* expressing the stand alone genes *PPE10, PPE24*,
showed low CFUs than their controls in RAW macrophages (31). However, in this
report, the authors do not provide a corresponding measure of *iNOS2* levels for this
data. Although unusual, we believe that this phenotype signifies that *M. tb* possesses
the ability to reduce its infection load as a possible strategy for immune evasion.

In addition to reduction in bacillary viability, we also found that the anti-inflammatory
cytokine IL-10 was consistently upregulated in macrophages infected with
recombinant *M. smegmatis* expressing *PPE50* as a function of time, whereas strains
expressing *PPE51* and the entire cluster showed none or inconsistent elevation in

expression of *IL-10*. Since IL-10 inhibits the production of host-protective pro-
inflammatory cytokines (31, 36, 37), the ability of PPE50 to induce an elevation in its
levels makes it an important component of the anti-inflammatory arsenal of *M. tb*.
Operating on the hypothesis that PPE50 mediates *IL-10* induction through ligand-
receptor interaction, we performed receptor pull-down experiments and discovered
TLR1 to be the cognate receptor for PPE50, with signal activation through the
receptor being indicated by an increase in IRAK1 phosphorylation. In the signaling
assay we also noted that IRAK1 phosphorylation was not induced in macrophages
infected with recombinant *M. smegmatis* strains expressing either *PPE51*, or the
entire operon. This observation, coupled with the lack of IL-10 upregulation in these
very macrophages, led us to infer a negative regulatory role for PPE51 on PPE50
function when the two are in a complex (Fig. 4A). The reversal of all the phenotypes
we observed on infection of THP-1 macrophages, on pre-treatment of macrophages
with an anti-TLR1 antibody prior to infection, unequivocally demonstrated the
functional outcome of PPE50-TLR1 interaction. This establishes PPE50 as a novel
TLR1 ligand, and to the best of our knowledge only the second member of the PE-
PPE family after PPE17 (38) that can bind and signal through this receptor.

Our findings suggest that the *PPE50-PPE51* cluster could play a role in facilitating
bacillary persistence through decreasing of bacillary load and upregulation of the
anti-inflammatory response, a two-pronged strategy to evade host immune
surveillance. These observations strongly suggest that PPE50 and PPE51 are
immunomodulatory proteins and are likely to participate in either the establishment
or persistence of infection by hampering the mounting of immune response to *M.*
*tb* infection *via* an altered TLR1 signaling pathway (Fig. 6). Performing macrophage
and mouse infections with knock-out strains of *M. tb* carrying deletions in *PPE50*,
*PPE51*, and the entire operon, would help solidify this hypothesis - these
experiments are proposed in the next phase of this study.

Hypoxia, non-replicative persistence, thiol (diamide) stress, oxidative stress, iron
deficiency, and starvation are all factors important to *M. tb* physiology, and have all
been shown to affect the expression of different PE/PPE genes (2, 7). This suggests
that the functions of these proteins are not redundant, but rather they serve distinct
purposes during different stages of disease progression, depending on the

pathogen's niche. The pathogenic process is probably enhanced when these proteins and other PPE proteins operate together. It has been hypothesized that both cognate and non-cognate ESX-secreted PE/PPE complexes and their constituent proteins have virulence-effector roles (39). Recent reports have shown the YxxxD/E motif to be a conserved secretion signal, which has been found in all known substrates or substrate complexes of the mycobacterial type VII secretion system (40). Since this signature is conserved in PPE51 as well, the localization/ export of these proteins could be tracked in *M. tb* strains with mutations in the *esx* genes to test this theory.

In summary, this study represents the first demonstration of an independent immunomodulatory function of a PPE-PPE protein cluster and strongly suggests that the *PPE50-PPE51* cluster could potentially play a pivotal role in the establishment and persistence of *M. tb* infection in the host.

**Conflict of interest**

The authors declare that there are no conflicts of interest.

**Funding information**

This work was supported by grant from the Council of Scientific and Industrial Research (CSIR) (BSC104 -SpLEnDID), Government of India and Institut Merieux (GAP 371) (to T. R. R.). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. K. P. was supported by a fellowship from CSIR (MLP 0144), S. Y. was supported by fellowships from Institut Merieux (GAP371) and CSIR (BSC104), R.P.M. was supported by a Senior Research Fellowship from the DBT.

**Author contributions**

T. R. R. designed the study. R. P. M., K. P. and S. Y. performed the experiments. T. R. R. and R.P.M. analysed the data. T.R.R. and R. P. M wrote the manuscript.

**FIGURE LEGENDS**

**Figure 1: Sequence homology of *M. tb* PPE50/PPE51.** Boxshade representation of PPE50, PPE51 protein sequence alignment.

**Figure 2: *PPE50* and *PPE51* are co-transcribed, and their gene products interact**. (A) Schematic depiction of the genomic structure of the *PPE50-PPE51* region in the *M. tb* genome showing the sites of the primers used in the investigation. (B) RT-PCR amplification products of the *PPE50-PPE51* gene pair : Lane 1- (50F+50R), Lane 2- (50JF+51JR), Lane 3 - (51F+51R), Lanes 5, 6, and 7 correspond to -RT controls for these respective primer pairs, Lane 4-100 bp DNA ladder. (C) Western blot of *in vitro* co-immunoprecipitation reactions of PPE50-6xHis and GST-PPE51 pulled with an anti-GST antibody and probed with anti-His antibody. (D) Western blot of PPE50-PPE51 co-IP from *M. smegmatis* expressing PPE51-myc and PPE50-6xHis. The complex was pulled down using an anti-c-myc antibody, and the blot was probed with an anti-His antibody. (E) M-PFC analysis of the interaction of PPE50 and PPE51. The image depicts the growth pattern of *M. smegmatis* co-transformants on Middlebrook 7H11 agar plates with (+) or without (-) 40µg/ml trimethoprim (TRIM). The findings are based on at least two biological replicates.

**Figure 3** : **Intra macrophage growth of recombinant *M. smegmatis* expressing *PPE50* and *PPE51*** CFU counts of *M. smegmatis* expressing pMV261, *PPE50*, *PPE51* and *PPE50-PPE51*, 24 h, 48 h and 72 h post infection in THP-1 macrophages (A) Input and $T_0$ (post-infection) CFU counts of infecting bacilli (± SEM) (B). *iNOS2* transcript levels in THP-1 macrophages infected with *M. smegmatis* expressing pMV261, *PPE50*, *PPE51* and *PPE50-PPE51*, 24 h, 48 h and72 h post infection (C). The data is representative of two biological replicates. *$p \leq 0.05$, **$p \leq 0.005$, ***$p \leq 0.001$   Error bars represent mean ± SEM of three biological replicates.

**Figure 4: Immunomodulatory effects of the PPE50- PPE51 cluster.** (A) *IL-10* transcript levels following THP-1 infection with recombinant *M. smegmatis* strains expressing *PPE50, PPE51, PPE50PPE51,*and the empty vector pMV261. (B) Receptor recognition by PPE50 Western blot depicting co-IP reactions of PPE50-His and solubilised cell-membrane fractions of THP-1 macrophages- an anti-His

antibody was used pull-down the complex and the blot was probed with an anti-TLR1

Ab. (C) Reverse pull-down for receptor recognition – an anti-TLR1 antibody was

used to pull down the complex and the blot was probed with an anti-His antibody. (D)

Western blots showing levels of phospho-IRAK1, Total IRAK1 and β-actin levels in

lysates from Macrophages infected with recombinant *M. smegmatis* strains

expressing *PPE50* (50), *PPE51* (51), *PPE50-PPE51* (50/51) and the empty vector

pMV261 (EV). The densitometric quantitation of this data is represented in (E). Error

bars represent ± SEM from two biological replicates. ***$p<0.001$, * $p<0.05$.

**Figure 5: Effect of TLR1 blocking on CFU counts, iNOS*2* & *IL-10* transcript**

**levels, and IRAK-1 phosphorylation.**  Growth index (A, B), *iNOS2* transcript levels

(C, D), transcript levels of *IL-10* in THP-1 macrophages (E, F) infected with a

recombinant *M. smegmatis* strain expressing the empty vector pMV261 (Ev) and

*PPE50*, with and without antibody blocking. (G) Western blots showing levels of

phospho-IRAK1, Total IRAK1 and β-actin levels in lysates from TLR1 blocked

infected macrophages (H) Densitometric quantitation of the blots in (G). The data

represents two biological replicates. Error bars represent ± SEM from two biological

replicates. ***$p<0.001$, **$p<0.005$, * $p<0.05$.

**Figure 6: Model depicting the immunomodulatory/ immune evasion role of the**

***PPE50-PPE51* cluster in *M. tb* pathophysiology.** MΦ - Macrophage**.**

**BIBLIOGRAPHY**

1. Lin PL, Flynn JL. 2010. Understanding latent tuberculosis: a moving target. J Immunol 185:15-22.

2. Majumdar SD, Vashist A, Dhingra S, Gupta R, Singh A, Challu VK, Ramanathan VD, Kumar P, Tyagi JS. 2012. Appropriate DevR (DosR)-mediated signaling determines transcriptional response, hypoxic viability and virulence of Mycobacterium tuberculosis. PLoS One 7:e35847.

3. Akhter Y, Ehebauer MT, Mukhopadhyay S, and Hasnain SE. 2012. The PE/PPE multigene family codes for virulence factors and is a possible source of mycobacterial antigenic variation: perhaps more? Biochimie 94:110-6.

4. Medha, Sharma S, Sharma M. 2021. Proline-Glutamate/Proline-Proline-Glutamate (PE/PPE) proteins of Mycobacterium tuberculosis: The multifaceted immune-modulators. Acta Trop 222:106035.

5. Tundup S, Pathak N, Ramanadham M, Mukhopadhyay S, Murthy KJ, Ehtesham NZ, Hasnain SE. 2008. The co-operonic PE25/PPE41 protein complex of *Mycobacterium tuberculosis* elicits increased humoral and cell mediated immune response. PLoS One 3:e3586.

6. Riley R, Pellegrini M, Eisenberg D. 2008. Identifying cognate binding pairs among a large set of paralogs: the case of PE/PPE proteins of Mycobacterium tuberculosis. PLoS Comput Biol 4:e1000174.

7. Voskuil MI, Schnappinger D, Rutherford R, Liu Y, Schoolnik GK. 2004. Regulation of the *Mycobacterium tuberculosis* PE/PPE genes. Tuberculosis (Edinb) 84:256-62.

8. Martini MC, Hicks ND, Xiao J, Alonso MN, Barbier T, Sixsmith J, Fortune SM, Shell SS. 2022. Loss of RNase J leads to multi-drug tolerance and accumulation of highly structured mRNA fragments in Mycobacterium tuberculosis. PLoS Pathog 18:e1010705.

9. Wang Q, Boshoff HIM, Harrison JR, Ray PC, Green SR, Wyatt PG, Barry CE, 3rd. 2020. PE/PPE proteins mediate nutrient transport across the outer membrane of Mycobacterium tuberculosis. Science 367:1147-1151.

10. Strong EJ, Wang J, Ng TW, Porcelli SA, Lee S. 2022. Mycobacterium tuberculosis PPE51 Inhibits Autophagy by Suppressing Toll-Like Receptor 2-Dependent Signaling. mBio 13:e0297421.

583    11.    Sambrook J, Fritsch EF, Maniatis T. 1989. Molecular Cloning: A Laboratory

584            Manual, 2nd edition. Cold Spring Harbor Laboratory Press, Cold Spring

585            Harbor, New York.

586    12.    Larkin MA, Blackshields G, Brown NP, Chenna R, McGettigan PA, McWilliam

587            H, Valentin F, Wallace IM, Wilm A, Lopez R, Thompson JD, Gibson TJ,

588            Higgins DG. 2007. Clustal W and Clustal X version 2.0. Bioinformatics

589            23:2947-8.

590    13.    Hofmann K, and W. Stoffel 1993. TMbasc—a database of membrane

591            spanning protein segments. Biol Chem Hoppe-Seyler 374:166.

592    14.    Kyte J, Doolittle RF. 1982. A simple method for displaying the hydropathic

593            character of a protein. J Mol Biol 157:105-32.

594    15.    Spratt JM, Ryan AA, Britton WJ, Triccas JA. 2005. Epitope-tagging vectors for

595            the expression and detection of recombinant proteins in mycobacteria.

596            Plasmid 53:269-73.

597    16.    Cascioferro A, Delogu G, Colone M, Sali M, Stringaro A, Arancia G, Fadda G,

598            Palu G, Manganelli R. 2007. PE is a functional domain responsible for protein

599            translocation and localization on mycobacterial cell wall. Mol Microbiol

600            66:1536-47.

601    17.    Stover CK, Bansal GP, Hanson MS, Burlein JE, Palaszynski SR, Young JF,

602            Koenig S, Young DB, Sadziene A, Barbour AG. 1993. Protective immunity

603            elicited by recombinant bacille Calmette-Guerin (BCG) expressing outer

604            surface protein A (OspA) lipoprotein: a candidate Lyme disease vaccine. J

605            Exp Med 178:197-209.

606    18.    Stover CK, de la Cruz VF, Fuerst TR, Burlein JE, Benson LA, Bennett LT,

607            Bansal GP, Young JF, Lee MH, Hatfull GF, et al. 1991. New use of BCG for

608            recombinant vaccines. Nature 351:456-60.

609    19.    Singh A, Mai D, Kumar A, Steyn AJ. 2006. Dissecting virulence pathways of

610            *Mycobacterium tuberculosis* through protein-protein association. Proc Natl

611            Acad Sci U S A 103:11346-51.

612    20.    Tiwari B, Ramakrishnan UM, Raghunand TR. 2015. The Mycobacterium

613            tuberculosis protein pair PE9 (Rv1088)-PE10 (Rv1089) forms heterodimers

614            and induces macrophage apoptosis through Toll-like receptor 4. Cell Microbiol

615            17:1653-69.

21.  Tiwari B, Soory A, Raghunand TR. 2014. An immunomodulatory role for the
Mycobacterium tuberculosis region of difference 1 locus proteins PE35
(Rv3872) and PPE68 (Rv3873). FEBS J 281:1556-70.

22.  Tundup S, Akhter Y, Thiagarajan D, Hasnain SE. 2006. Clusters of PE and
PPE genes of *Mycobacterium tuberculosis* are organized in operons:
evidence that PE Rv2431c is co-transcribed with PPE Rv2430c and their
gene products interact with each other. FEBS Lett 580:1285-93.

23.  Chaturvedi R, Bansal K, Narayana Y, Kapoor N, Sukumar N,
Togarsimalemath SK, Chandra N, Mishra S, Ajitkumar P, Joshi B, Katoch VM,
Patil SA, Balaji KN. 2010. The multifunctional PE_PGRS11 protein from
*Mycobacterium tuberculosis* plays a role in regulating resistance to oxidative
stress. J Biol Chem 285:30389-403.

24.  Daim S, Kawamura I, Tsuchiya K, Hara H, Kurenuma T, Shen Y, Dewamitta
SR, Sakai S, Nomura T, Qu H, Mitsuyama M. 2011. Expression of the
*Mycobacterium tuberculosis* PPE37 protein in *Mycobacterium smegmatis*
induces low tumour necrosis factor alpha and interleukin 6 production in
murine macrophages. J Med Microbiol 60:582-91.

25.  Nair S, Ramaswamy PA, Ghosh S, Joshi DC, Pathak N, Siddiqui I, Sharma P,
Hasnain SE, Mande SC, Mukhopadhyay S. 2009. The PPE18 of
Mycobacterium tuberculosis interacts with TLR2 and activates IL-10 induction
in macrophage. J Immunol 183:6269-81.

26.  North RJ. 1998. Mice incapable of making IL-4 or IL-10 display normal
resistance to infection with Mycobacterium tuberculosis. Clin Exp Immunol
113:55-8.

27.  Singh PP, Parra M, Cadieux N, Brennan MJ. 2008. A comparative study of
host response to three Mycobacterium tuberculosis PE_PGRS proteins.
Microbiology 154:3469-79.

28.  Anes E, Peyron P, Staali L, Jordao L, Gutierrez MG, Kress H, Hagedorn M,
Maridonneau-Parini I, Skinner MA, Wildeman AG, Kalamidas SA, Kuehnel M,
Griffiths G. 2006. Dynamic life and death interactions between *Mycobacterium
smegmatis* and J774 macrophages. Cell Microbiol 8:939-60.

29.  Gupta D, Sharma S, Singhal J, Satsangi AT, Antony C, Natarajan K. 2010.
Suppression of TLR2-induced IL-12, reactive oxygen species, and inducible
nitric oxide synthase expression by *Mycobacterium tuberculosis* antigens

expressed inside macrophages during the course of infection. J Immunol
184:5444-55.

30.  Voskuil MI, Schnappinger D, Visconti KC, Harrell MI, Dolganov GM, Sherman
DR, Schoolnik GK. 2003. Inhibition of respiration by nitric oxide induces a
*Mycobacterium tuberculosis* dormancy program. J Exp Med 198:705-13.

31.  Meng L, Tong J, Wang Q, Niu C, Gao Q. 2017. Diverse effects of
mycobacterial proline-proline-glutamic acid proteins upon interaction with host
macrophages. FEMS Microbiol Lett 364.

32.  Hayes M, Salvato M. 2012. Arenavirus evasion of host anti-viral responses.
Viruses 4:2182-96.

33.  Mukhopadhyay S, Balaji KN. 2011. The PE and PPE proteins of
Mycobacterium tuberculosis. Tuberculosis (Edinb) 91:441-7.

34.  Tiwari BM, Kannan N, Vemu L, Raghunand TR. 2012. The Mycobacterium
tuberculosis PE proteins Rv0285 and Rv1386 modulate innate immunity and
mediate bacillary survival in macrophages. PLoS One 7:e51686.

35.  Yeruva VC, Kulkarni A, Khandelwal R, Sharma Y, Raghunand TR. 2016. The
PE_PGRS Proteins of Mycobacterium tuberculosis Are Ca(2+) Binding
Mediators of Host-Pathogen Interaction. Biochemistry 55:4675-87.

36.  Murray PJ, Young RA. 1999. Increased antimycobacterial immunity in
interleukin-10-deficient mice. Infect Immun 67:3087-95.

37.  Redpath S, Ghazal P, Gascoigne NR. 2001. Hijacking and exploitation of IL-
10 by intracellular pathogens. Trends Microbiol 9:86-92.

38.  Udgata A, Qureshi R, Mukhopadhyay S. 2016. Transduction of Functionally
Contrasting Signals by Two Mycobacterial PPE Proteins Downstream of TLR2
Receptors. J Immunol 197:1776-87.

39.  Sampson SL. 2011. Mycobacterial PE/PPE proteins at the host-pathogen
interface. Clin Dev Immunol 2011:497203.

40.  Daleke MH, Ummels R, Bawono P, Heringa J, Vandenbroucke-Grauls CM,
Luirink J, Bitter W. 2012. General secretion signal for the mycobacterial type
VII secretion pathway. Proc Natl Acad Sci U S A 109:11342-7.

**Figure 1**

**Figure 2**

**B**

| | pMV261 | pMV261- *ppe50* | pMV261- *ppe51* | pMV261- *ppe50-ppe51* |
|---|---|---|---|---|
| **Input** | 2.09 ± 0.2 X10⁸ | 1.98 ± 0.5 X10⁸ | 2.13 ± 0.5 X10⁸ | 2.60 ± .3 X10⁸ |
| T0 | 29 ± 3X10⁵ | 20.9 ± 9.5X10⁵ | 19.5 ± 6.7X10⁵ | 31.1 ± 1.6X10⁵ |

**Figure 3**

**Figure 4**

**Figure 5**

**Figure 6**