Dual targeting of MAPK and PI3K pathways unlocks redifferentiation of *Braf-*

mutated thyroid cancer organoids

Hélène Lasolle [1], Andrea Schiavo [1], Adrien Tourneur[1], Pierre Gillotay [1], Bárbara de Faria

da Fonseca [1], Lucieli Ceolin [1,2], Olivier Monestier [1], Benilda Aganahi [1], Laura Chomette

[1], Marina Malta Letro Kizys [1], Lieven Haenebalcke [3], Tim Pieters [3,4], Steven Goossens

[3,5], Jody Haigh [6], Vincent Detours [1], Ana Luiza Silva Maia [2], Sabine Costagliola [1] and

Mírian Romitti [1*]

[1] Institut de Recherche Interdisciplinaire en Biologie Humaine et Moléculaire (IRIBHM),

Université Libre de Bruxelles (ULB), Brussels, Belgium

[2] Thyroid Section, Endocrine Division, Hospital de Clínicas de Porto Alegre (HCPA),

Universidade Federal do Rio Grande do Sul (UFRGS), Porto Alegre, Brazil

[3] VIB, Flanders Institute for Biotechnology, Ghent University, Ghent, Belgium

[4] Department of Biomolecular Medicine, Faculty of Medicine and Health Sciences, Ghent

University, Ghent, Belgium

[5] Department of Diagnostic Sciences, Faculty of Medicine and Health Sciences, Ghent

University, Ghent, Belgium

[6] CancerCare Manitoba Research Institute, Department of Pharmacology and

Therapeutics, Rady Faculty of Health Sciences, University of Manitoba, Winnipeg,

Manitoba, Canada

* Corresponding author: mirian.romitti@ulb.be

The authors declare no potential conflicts of interest

Thyroid cancer is the most common endocrine malignancy and several genetic events

have been described to promote the development of thyroid carcinogenesis. Besides the

effects of specific mutations on thyroid cancer development, the molecular mechanisms

controlling tumorigenesis, tumor behavior, and drug resistance are still largely unknown.

Cancer organoids have been proposed as a powerful tool to study aspects related to tumor

development and progression and appear promising to test individual responses to

therapies. Here, using mESC-derived thyroid organoids, we developed a $Braf^{V637E}$-

inducible model able to recapitulate the features of papillary thyroid cancer *in vitro*.

Overexpression of the murine $Braf^{V637E}$ mutation, equivalent to $Braf^{V600E}$ in humans,

rapidly triggers to MAPK activation, cell dedifferentiation, and disruption of follicular

organization. $Braf^{V637E}$-expressing organoids show a transcriptomic signature for p53,

focal adhesion, ECM-receptor interactions, EMT, and inflammatory signaling pathways.

Finally, PTC-like thyroid organoids were used for drug screening assays. The

combination of MAPK and PI3K inhibitors reversed *$Braf^{V637E}$* oncogene-promoted cell

dedifferentiation while restoring thyroid follicle organization and function *in vitro*. Our

results demonstrate that pluripotent stem cells-derived thyroid cancer organoids can

mimic tumor development and features while providing an efficient tool for testing novel

targeted therapies.

INTRODUCTION

Pluripotent Stem Cells (PSCs) emerged as a model to dissect and recapitulate the

molecular events and gene networks that regulate cell fate determination, cell

differentiation and organogenesis. Guided differentiation of PSCs into 3D organized

tissue by inducible overexpression of specific transcription or growth factors and

inhibitors allows the derivation of *in vitro* organoids and the understanding of

mechanisms regulating cell differentiation and tissue formation[1]. Due to their remarkable

self-organizing structures and functional properties, organoid technology became a

powerful tool to model organ development and disease 'in a dish' [2,3].

Our group has demonstrated the generation of functional thyroid organoids

derived from mouse and human embryonic stem cells (mESC and hESC)[4,5]. The

differentiation protocols rely on transient induction of *Nkx2.1* and *Pax8* thyroid

transcription factors. This approach enables an efficient generation of thyroid follicular

cells that organize into three-dimensional follicular structures capable of thyroid hormone

production *in vitro* and *in vivo*. These models, along with the further development of

thyroid organoids derived from healthy murine and human pluripotent or adult stem cells,

(PSCs/aSC)[6–16] open new perspectives to explore thyroid developmental and pathological

processes, including thyroid cancer.

In the last decade, the use of organoids for cancer research emerged opening new

possibilities to better understand tumor behavior. Initially, colon, prostate, pancreatic,

ovarian, lung and thyroid cancer-derived organoids have been efficiently generated,

which resemble phenotypically and genetically the tumor of origin[17–26]. A large set of

healthy and tumour-matching organoids has been generated and is available through

biobanks. These patient-derived organoids are particularly interesting in testing

individual responses to therapies[21,26,27].

Recently, studies have reported the generation of cancer models arising from

healthy (aSCs and PSCs) cells by controlling oncogene expression[28–30]. By using shRNA

and CRISPR-Cas9, stomach cancer (*Cdh1*[-/-]; *Tp53*[-/-])[31], colon cancer (*APC, TP53, KRAS*

and *Smad4* mutations)[28,29,32], pancreatic cancer (*KRAS* and/or *TP53* mutations)[17,33] and

lung adenocarcinoma organoids (*HER2* overexpression) could be efficiently generated[30].

Compared to patient-derived organoids, healthy stem cells-derived cancer organoids are

suitable to address additional questions such as precise effects of oncogenes and early

events driving tumorigenesis; the role of cancer stem cells on tumor induction, genomic

stability, the effect of treatments at different stages of carcinogenesis and screening for

new therapeutical tools[34]. Besides, they allow multiple reproducible experiments free

from inter-individual variability.

Thyroid cancer is the most frequent endocrine malignancy, and several genetic

events have been described as driving thyroid carcinogenesis. Papillary thyroid

carcinoma (PTC) is the most common malignant thyroid tumor[35], and aberrant activation

of the MAPK pathway is a hallmark[36]. The $BRAF^{V600E}$ mutation is the most frequent

genetic event, accounting for around 50% of PTCs[37–39] and results in an oncogenic protein

with markedly elevated kinase activity that constitutively induces MEK/ERK signaling

transduction[40–42].

BRAF-mutated thyroid tumors are often less differentiated, mainly due to the

lower expression levels of thyroid functional genes *NIS, TSHR, TG* or *TPO,* leading to a

low Thyroid Differentiation Score (TDS)[43]. This dedifferentiated state is associated with

a worse clinical condition and with a higher rate of radioactive iodine (RAI) refractory

tumors[44,45]. However, its impact on prognosis is still uncertain, as $BRAF^{V600E}$ alone has

not been proven to be an independent prognostic factor[46,47]. On the other hand, *BRAF* and

TERT promoter mutated tumors are associated with higher specific mortality[48]. Also,

studies have shown a small proportion (around 20%) of $BRAF^{V600E}$ mutated tumors

presenting a higher level of differentiation associated with less aggressive behavior and

a preserved RAI uptake ability[43,49]. This suggests a possible cell heterogeneity among the

BRAF^{V600E}-mutated tumors.

So far, efficient, fast, and cost-effective tools are lacking to functionally

characterize a large set of candidate genes in the context of thyroid carcinogenesis.

Thyroid cancer organoids/spheroids models have been recently described based on

patient tumor-derived models[50–54]. Tumor-derived organoids can be used to explore

patient-specific tumor behavior and the treatment response, but present high

heterogeneity and low reproducibility. Saito et al. described a mouse model of poorly

differentiated thyroid carcinoma obtained after transplantation of thyroid organoids

derived from $Tp53^{-/-}$ mice with *NRAS* activating mutation[25]. Recently, Veschi *et al.* [55]

generated PTC and FTC organoids by inducing *BRAF*, *NRAS* and *TP53* mutations in

thyroid progenitors derived from hESC. After transplantation, the tumors resembled

Papillary and Follicular tumors and, transcriptomics analysis suggested a cooperative

effect of Kisspeptin receptor (KISS1R) and Tissue Inhibitor of Metalloproteinase 1

(TIMP1)/Matrix metallopeptidase 9 (MMP9)/Cluster of differentiation 44 (CD44), on

tumor development and progression[55].

In this study, we have established an *in vitro* organoid model for PTC to streamline

the process of screening potential targets and pharmaceutical agents. We achieved this by

leveraging our established model of thyroid generation, which involves the inducible

expression of the $Braf^{V637E}$ mutation in functional thyroid follicles derived from mouse

ESCs.

RESULTS

mESC_BRAF^{V637E} cell line generation and characterization

We combined the previously described mESC transient induction thyroid differentiation

strategy[4] with a $Braf^{V637E}$-ERT^2 inducible model[56] to generate a double-inducible

recombinant mESC_Braf^{V637E} cell line. The murine Braf^{V637E} mutation corresponds to the

human Braf^{V600E} mutation. The resulting mESC_Nkx2-1/Pax8_bTg_Braf^{V637E}_ERT^2

clones were initially selected and characterized by the response to doxycycline (Dox)

treatment, confirmed by the overexpression of Nkx2-1 and Pax8; and ability to

spontaneously differentiate into cells from the three germ-layers (Supplementary Fig. 1A-

C). In addition, a control mESC line was generated in which Braf^{V637E}_ERT^2 system was

replaced by the eGFP sequence (mESC_Nkx2-1/Pax8_bTg_eGFP). Similarly, the control

line responded properly to Dox induction and maintained the pluripotency capacity

(Supplementary Fig. 1D-F).

Thyroid differentiation protocol

The capability of the new cell line to generate thyroid follicles and express the Braf^{V637E}

oncogene was demonstrated following the mESC thyroid differentiation protocol[4,57,58].

Since we used a newly modified mESC line, the steps of the differentiation protocol were

tested and adapted accordingly (Fig. 1A). Here, using the mESC_Nkx2-

1/Pax8_bTg_Braf^{V637E} line, we observed that Dox stimulation for 5 days leads to higher

exogenous and endogenous *Nkx2-1*, *Pax8* and *Tg* mRNA compared to 3 or 4 days of

treatment (Supplementary Fig. 2A-D). Similar results were obtained using the control

mESC line (data not shown). Next, two weeks of treatment with cAMP (from day 9 to

23) markedly induced the expression levels of *Nkx2-1, Pax8, Tg, TshR, Nis, Tpo* and

exogenous *Braf^{V637E}* when compared to the control (-Dox -cAMP) (Fig. 1B).

Immunofluorescence staining also showed Nkx2-1 and Tg-expressing cells (Fig. 1C)

organized into follicular structures with intraluminal iodinated-Tg (Tg-I) accumulation

(Fig. 1D). Since our protocol also generates other cell types than thyroid[58], a follicle

enrichment (FE) step was added to the protocol to enrich the follicular structures and

induce the Braf^{V637E} oncogene in a purer population. This protocol led to an increase in

thyroid markers levels (Fig. 1E) when compared to the non-enriched control condition

(No FE), while the follicular structures and functionality were preserved as demonstrated

by Nkx2-1, Nis and Tg-I stainings (Fig. 1F-G). Notably, after re-embedding in Matrigel

(MTG), follicles could be maintained in culture for at least 28 additional days without

major histological changes (data not shown).

Braf^{V637E} induction on thyroid organoids leads to PTC-like phenotype

To ensure that Braf^{V637E} oncogene is expressed specifically among follicular thyroid cells,

we used the bovine Tg (bTg) as a promoter controlling Braf^{V637E}_ERT^2 expression.

Effectively, exogenous *Braf^{V637E}* mRNA expression was rapidly induced after Dox

treatment (Day 9; Supplementary Fig. 2E), maintaining the expression levels stable over

time (Fig.1B and Supplementary Fig. 3A), while *Tg* expression increased following the

differentiation program (Supplementary Fig. 3A). In the absence of tamoxifen (4OHT)

Braf^{V637E} protein is not active/phosphorylated since it is maintained in the nucleus in an

inactive complex with HSP90. The addition of 4OHT induces nuclear translocation of the

Braf^{V637E} mutant to the cytoplasm (NES signal) [56], then driving MAPK activation[56] (Fig.

2A). Indeed, 48 hours after 4OHT addition, we observed an increase in ERK

phosphorylation (pERK) compared to the control condition (cAMP) (Fig. 2B).

One of the hallmarks of thyroid cancer is the downregulation of thyroid

differentiation markers[41,59]. In our organoid model, the activation of Braf^{V637E} rapidly led

to a progressive time-dependent decrease of *Tpo*, *TSHR* and *Tg* mRNA expression (Fig.

2C-E and Supplementary Fig. 3B), detected as early as 6 hours after 4OHT addition (Fig.

2C). Notably, more significant downregulation compared to the cAMP control was

observed after 48 hours and 7 days while maintained for at least 21 days (Fig. 2D-F,

respectively). As for PTCs, in our model, the expression of thyroid transcription factors

Pax8 and *Nkx2-1* was also globally maintained. However, a partial decrease of *Nkx2.1*

levels was observed at 7 days, which corroborates recent findings suggesting that thyroid

cells transiently downregulated NKX2.1 in early tumor stages[60]. Conversely, Galectin 3

(Lgals3) expression, which has been suggested as a marker for thyroid malignancies and,

more specifically, for PTC[61], was significantly increased at 48 h and 7 days of 4OHT

treatment (Fig. 2D-E). Also, an increase in Fibronectin 1 (*Fn1*) expression, associated

with aggressive thyroid cancer[62,63], was observed after 7 days of Braf[V637E] induction,

suggesting a more advanced dedifferentiated cell state in our organoid model (Fig. 2E).

The generation of thyroid organoids was equally successful when differentiating

mESC_Nkx2-1/Pax8_bTg_Braf[V637E]_ERT2 cells using hrTSH instead of cAMP, as

demonstrated in Supplementary Figure 4A and C. Furthermore, the addition of 4OHT to

the conditioned media (for a duration of 7 days, as illustrated in Supplementary Figure

4B-C) led to dedifferentiation and histological changes.

Cell proliferation assessment demonstrated an increased proportion of Nkx2-

1/BrdU+ cells within the Braf[V637E]-expressing organoids (4OHT) as compared to the

controls (cAMP) after 7 days of treatment (Fig. 2G). Interestingly, there were no

observable variations in proliferation even after 21 days of 4OHT treatment (Fig. 2G).

Furthermore, considering that prior studies have indicated that the induction of oncogenes

in organoids can lead to cell death[64], our analysis of apoptosis showed no significant

difference in the proportions of Nkx2.1/Caspase3+ between cAMP and 4OHT-treated

organoids both at 7 and 21 days of 4OHT treatment (Fig. 2H). However, a notable

increase in the proportion of Nkx2.1/Caspase3+ cells was evident when cAMP-treated

organoids were exposed for 24h to staurosporine, a highly potent inducer of apoptosis.

(Fig. 2H). Given that tumor growth arrest and antiapoptotic phenotype are linked to

oncogene-induced senescence (OIS)[65,66], we examined p21 expression at both 7 and 21

days of 4OHT treatment. Notably, we observed no co-expression of p21 with Nkx2-1 in

either the control or 4OHT-treated thyroid organoids. However, a small proportion of

non-thyroid cells (Nkx2.1 negative) were positive for p21 and used as positive controls

(Supplementary Fig. 5A-C).

Histological characterization performed after 48 hours of continuous activation of

Braf^{V637E} by 4OHT demonstrated disruption of the follicular organization in non-enriched

thyroid organoids compared to the control condition (cAMP) (Fig. 3A). Nkx2-1 and Tg

staining revealed that Braf^{V637E} oncogene strongly disturbs thyroid follicles resulting in

elongated and unorganized structures. In addition, heterogeneous expression of Tg was

observed among the Nkx2-1 cells (Fig. 3A). Further analysis, performed in follicle

enriched (FE) population, confirmed the effect of Braf^{V637E} oncogene in promoting cell

dedifferentiation and loss of follicle organization. Tg levels were overall reduced and

heterogeneous among the unorganized Nkx2-1+ cells (Fig. 3B). Interestingly, the cell

disorganization caused by Braf^{V637E} oncogene might pass by an initial expansion of the

follicular size with infiltration of Nkx2-1 cells inside of the lumen compartment, which

also presents a heterogeneous distribution of Tg (Supplementary Fig. 3C) since this

feature was observed among some "follicular-like" remaining structures after 48 h of

4OHT treatment. Reassuringly, staining for E-Cadherin (epithelial) and Zo-1 (intra-

luminal) adhesion markers showed that Braf^{V637E}-expressing cells are not able to preserve

polarization and, consequently lumen space is not well defined in most of the structures

(Fig. 3C and D, respectively). Therefore, the functionality of those cells was impaired

and, Nis expression and Tg-I accumulation were significantly reduced compared to the

control condition (Fig. 3E-F). Of note, despite the strong effect of Braf^{V637E} oncogene on

thyroid follicle disruption, we observed that few follicles still preserved a certain degree

of organization (Supplementary Fig. 3D) after 48 h of 4OHT treatment. As expected, after

7 days of Braf^{V637E} continuous stimulation, a higher level of cell dedifferentiation and

lack of organization could be detected, with a lower proportion of the Nkx2-1 cells

expressing Tg compared to control (cAMP) (Fig. 3G) and to the previous time point (48

h; Fig. 3A). However, despite the absence of proper follicular organization, there was an

increased proportion of Tg-expressing cells within the Braf^{V637E}-expressing organoids

after 21 days of 4OHT treatment compared to earlier time points (48 h and 7 days; Fig.

3H). This observation, in conjunction with the gene expression and the proliferation data

(Fig. 2F-G) implies a partial arrest in the tumorigenic processes at a later stage.

The above-described experiments were also performed using the control TRE-

Nkx2-1-Pax8_bTg-eGFP cell line. Adding 4OHT to mature follicles for 7 days did not

impair the levels of the thyroid differentiation markers and the follicles' organization

(Supplementary Fig. 6A-B). Furthermore, there was a slight increase in the proportions

of Nkx2.1 and GFP-expressing cells when WT organoids were treated with 4OHT (7

days; Supplementary Fig. 6C). It's worth noting, as previously demonstrated[58], that

bTg_GFP-expressing cells do not encompass the entire Tg-expressing cells population

(Supplementary Fig. 6D). This could be attributed to the lack of regulatory regions not

included in our construct.

Effect of transient Braf^{V637E}-induction on cell differentiation

Since our system for Braf^{V637E} activation is dependent on continuous treatment with

4OHT, we transiently treated the cells with 4OHT for 2 days, followed by 5 days of

culture in a 4OHT-free medium (cAMP) in order to explore if the oncogenic effect is

maintained when the protein is not active. Surprisingly, we did not observe recovery in

Nkx2-1, TSHR, Tg or *Tpo* gene expression levels (Supplementary Fig. 7A, C-E) after

4OHT removal. However, *Pax8* and *Slc5a5/Nis* expression was partially recovered

(Supplementary Fig. 7B and F), confirming the specific inhibitory effect of Braf^{V637E}

oncogene on *Slc5a5/Nis* expression. These findings indicate that *Nis* regulatory

mechanisms are maintained in our system and suggest it as a tool for Nis-reactivation

studies.

Screening for Nis re-expression using signaling pathways inhibitors

Studies in mouse models and humans have demonstrated that tumors carrying *BRAF*

mutations show reduced expression of *Nis* and, consequently a higher rate of radioiodine

(RAI) refractoriness[41,44,67]. Here, we used our organoids model to explore new strategies

to reactivate *Nis* expression in Braf[V637E]-expressing cells by screening distinct categories

of inhibitors previously described to be involved in *Nis* regulation [68,69]. PD0325901

(MEK inhibitor), LY204002 (PI3K inhibitor), VPA (HDAC inhibitor), NAC (N-acetyl

cysteine, antioxidant compound), 5-AZA-2' deoxycytidine and RG108 (DMNT

inhibitors) were tested isolated and/or in combination for 3 days (in addition to

cAMP+4OHT) using Braf[V637E]-expressing organoids (previously treated for 4 days with

4OHT; Fig. 4A). Initially, we observed that VPA alone was the only treatment able to

completely restore *Nis* expression levels (Fig. 4B, C and Supplementary Fig. 8A).

However, when combined, PD0325901 and LY204002 inhibitors; PD0325901,

LY204002 and VPA; and PD0325901 and NAC resulted in a great increase of *Nis* mRNA

(Fig. 4B, C and Supplementary Fig. 8A). IF staining shows Nis protein re-expression

among Braf[V637E]-induced cells treated with VPA alone, but localization was not restricted

to the basolaterall membrane. In contrast, when cells were treated with MEK (PD) of

PI3K (LY) inhibitors, associated or not with VPA, we observed Nis protein correctly

localized at the basolateral membrane and surprisingly, it restored the follicular structure

(Fig. 4C).

MAPK and PI3K signaling inhibition cause Braf^{V637E}-induced cell redifferentiation and function recovery

Since we observed that HDAC, MEK, and PI3K inhibition seem to favor follicle reorganization, we evaluated if such treatments also induce cell redifferentiation and function recovery. Initially, we analyzed the gene expression levels of thyroid differentiation markers strongly affected by Braf^{V637E} induction. The effects of isolated and combined drugs on *Tg, TSHR, Tpo, Nkx2-1* and *Duox* genes appear to be variable and pathway-dependent (Supplementary Fig. 8B-F). As for *Slc5a5/Nis*, MEK and PI3K inhibition in combination with or without VPA resulted in complete recovery of *Tg*, *TSHR* and *Tpo* mRNA levels (Fig. 4D-F), indicating cell redifferentiation. While *TSHR* levels can be restored by treatment with isolated inhibitors (Fig. 4E), *Tpo* expression appears to be dependent on PI3K signaling (Fig. 4F). On the other hand, *Tg* mRNA expression was recovered only when the inhibitors were used in combination (Fig. 4D). Indeed, immunostaining showed that isolated inhibitors could not restore Tg levels. Nevertheless, a higher proportion of Tg-expressing cells was observed with the MEK inhibitor PD0325901 (Supplementary Fig. 8G). Conversely, combined MEK and PI3K inhibitors restored Tg protein levels and led to a significant reorganization of the cells into follicles comparable to the control (cAMP) condition (Fig. 4G-H). Considering that inhibition of MEK and PI3K pathways (+/-VPA) recovered thyroid differentiation and follicular organization, we tested the functionality of reorganized follicles. Remarkably, we observed an accumulation of Tg-I in the lumen of follicles treated with the combinations of inhibitors (Fig. 4I). The organification assay confirmed that the co-treatment restored iodide uptake and ^{125}I binding to proteins resulting in organification levels comparable to the control (Fig. 4J-L).

Dabrafenib and trametinib effect on redifferentiation is potentialized by PI3K inhibition

The treatment of advanced thyroid carcinomas as PDTC and ATC with BRAF inhibitor dabrafenib and MEK inhibitor trametinib have shown significant redifferentiation and response rates in *BRAF*-mutated tumors[70–72]. In this study, we examined the effect of isolated dabrafenib and trametinib inhibitors, as well as their combination with PI3K inhibition, on thyroid cancer organoids. Using the same experimental approach (Fig. 4A), we noted that both dabrafenib and trametinib treatments successfully restored *Nis* expression to levels comparable to the control group (treated with cAMP). Notably, the combination of trametinib with a PI3K inhibitor yielded an even more significant recovery, resulting in a threefold increase compared to the control group (Fig. 5A). Furthermore, isolated treatments with dabrafenib and trametinib elicited a moderate restoration of the main thyroid differentiation markers' expression. However, the most substantial recovery of *Tg, Tpo*, and *TSHR* levels was observed when organoids were subjected to a combination of dabrafenib and trametinib along with a PI3K inhibitor, with the latter showing the most pronounced effect (Fig. 5B-D). Interestingly, both Tg and Nis proteins were detectable in all experimental conditions, with some cells organized into functional follicles able to produce Tg-I (Fig. 5E). However, due to the complexity of the cell composition and organization of our 3D organoids, accurately quantifying the proportion of reorganized follicles proved to be very challenging and could not be efficiently assessed in the present study. Furthermore, our proliferation and apoptosis assays revealed that co-treatment with Trametinib and LY led to a decrease in proliferation (Nkx2.1/BrdU+ cells) and a marginal increase in apoptosis (Nkx2.1/Caspase 3 + cells). Comparable outcomes were achieved through incubation with MEK and PI3K inhibitors (PD+LY). Nevertheless, it's worth noting that apoptosis induction was notably

more pronounced when compared to the effects observed with Trametinib+LY treatment

(Fig. 5F-G). Western blot analysis distinctly revealed an increase in pERK levels after a

7-day 4OHT treatment. However, when Trametinib and LY were combined in the

treatment, it led to a mild inhibition of pERK compared to the conditions involving 4OHT

alone or in combination with PD+LY (Fig. 5H).

Transcriptomic characterization of the PTC model and drug screening

Transcriptomic analysis of control (cAMP), Braf^{V637E}-induced (4OHT) and inhibitors-

treated (PD+LY) organoids (Fig. 6A) confirmed downregulation of thyroid genes under

Braf^{V637E} stimulation (4OHT condition) whereas, global recovery of expression was

observed under inhibitors treatment (Fig. 6A). Accordingly, thyroid differentiation scores

(TDS and enhanced (e)TDS) were decreased under Braf^{V637E} induction while they

recovered in the presence of inhibitors (Fig. 6B). Conversely, the ERK activation score

was higher in the Braf^{V637E} samples and decreased sharply under inhibitor conditions (Fig.

6C). Since bulk RNAseq was performed using the follicle-enriched population, the

presence of non-thyroid cells is reduced compared with the whole original organoids but

still present. It may explain the modest increase in ERK score among Braf^{V637E}-activated

cells while the apparent decrease under the condition with inhibitors reflects the effect of

the treatment on each cell type.

Differential expression analysis identified 321 Differential Expressed Genes

(DEGs; 156 upregulated and 165 downregulated genes) in the 4OHT condition compared

to cAMP; and 853 DEGs in inhibitors compared to 4OHT (287 upregulated and 566

downregulated genes). Gene enrichment classification analysis of DEGs results is

presented in Fig. 6D-G. Briefly, among the upregulated genes in Braf^{V637E}-induced

compared to control, we observed gene signature for the hyperactivation of PI3K-AKT-

mTOR, TNF, and cytokines signaling and promotion of Epithelial-Mesenchymal

transition (EMT) (Fig. 6D). While genes associated with thyroid hormone production,

TGF-beta, Wnt/Beta-Catenin pathway, and regulation of angiogenesis were down-

regulated (Fig. 6E). Conversely, the inhibitor's treatment, compared to 4OHT condition,

evidenced upregulation of genes associated with thyroid hormone production, cell-cell

contact, extra-cellular matrix organization, and angiogenesis processes (Fig. 6F), while

PI3K, MAPK, TNF, cytokines signaling, ECM-receptor interactions and EMT-related

genes were downregulated (Fig. 6G). The list of the DEGs for each gene enrichment

classification is provided in Supplementary Table 4.

Given our transcriptomics analysis, which indicated elevated expression of

inflammation-related genes and downregulation of the Wnt/β-catenin pathway in

$Braf^{V637E}$-expressing cells, we tested the effect of anti-inflammatory drugs, specifically

dexamethasone and CC-5013 (Lenalidomide; TNF-alpha inhibitor) as well as CHIR-

99021 (Wnt/β-catenin activator) on our cancer organoids. Following the same

experimental strategy as with the other inhibitors (Fig. 4A), we observed that co-treatment

with dexamethasone and CC-5013 led to approximately a 3.5 fold increase in *Nis*

expression compared to the control condition (cAMP) (Supplementary Fig. 9A),

confirming the previously described inhibitory effect of inflammation on *Nis* expression.

On the other hand, for the other main thyroid differentiation genes, namely *Tg, TSHR* and

Tpo, treatment with anti-inflammatory drugs and the Wnt/β-catenin activator did not

result in their re-expression (Supplementary Fig. 9B-D). This suggests that these

alterations are likely a consequence of the oncogenic process rather than being driving

factors.

DISCUSSION

Here, we described a Braf[V637E] oncogene-derived thyroid cancer organoid model recapitulating patient tumor features. *In vitro* thyroid cancer models that recapitulate tumor development and behavior can facilitate the identification of early tumor drivers and enable the screening of several new drugs for the treatment of thyroid cancer without the need to use too many animal models. Combining our previous thyroid organoid model derived from mESCs with an inducible system, the Braf[V637E] oncogene could be induced explicitly in mature Tg-expressing cells to obtain a 3D *in vitro* cancer model. Overexpression of Braf[V637E] rapidly led to MAPK activation with increased pERK, cell dedifferentiation, and disruption of follicular organization. Similar effects have been described in genetically modified mouse models[41,59,73]. Furthermore, the gene expression signature of our cancer organoids confirmed the findings found in the PTC samples, which also showed enrichment of genes associated to p53, focal adhesion, ECM-receptor interactions, EMT and inflammatory pathways[74–76].

Activating mutations in the *BRAF* gene are found in approximately 7% of all solid human tumors, particularly common in PTCs, ATCs, and melanomas. In addition, they have been reported less frequently in colorectal cancers (CRCs), lung cancers, pediatric low-grade gliomas (PLGGs), glioblastomas, breast cancers, and serous ovarian cancers[77–83]. In PTCs and melanomas, BRAF[V600E] mutation seems to be associated with a higher degree of dedifferentiation and more aggressive histological patterns. However, its prognostic role is still debated as it was not found independently of histological features[46,47]. Nevertheless, studies suggest that BRAF[V600E] might predict response to tyrosine kinase inhibitors (TKIs) in melanoma and lung cancer[84,85].

Surgery remains the first choice for thyroid cancer therapy. Recommended post-operative treatment includes TSH suppression and RAI ablation, particularly as adjuvant

treatment for patients at high risk for tumor recurrence and to treat patients with

persistent/recurrent or metastatic disease[86]. The benefit of RAI has been demonstrated in

patients by reducing the risk of recurrence and disease-related mortality[87]. However,

approximately two-thirds of metastatic DTC become radioiodine refractory (RR-DTC),

defined by the absence of iodine uptake or tumor progression despite uptake[88,89]. There

is no curative treatment for radioiodine refractory DTCs. The recommended first-line

systemic treatment when local therapies are not possible is targeted therapies using

mainly multitarget TKIs, such as Lenvatinib showed a progression-free survival of 18

months [67,90–92].

NIS, a basal membrane iodide transporter, plays a critical role in radioiodine

accumulation in DTC cells and its level is closely related to response to RAI (I^{131}) therapy.

Studies in mouse models and humans have shown that the presence of *BRAF* mutations

results in lower *Nis* levels and, consequently a higher rate of RAI-refractory tumours[41,44].

Here, we confirmed that this regulation is preserved in our cancer organoids, as disruption

of Braf^{V637E} activation led to in recovery of *Nis* levels. As TKIs, retinoic acids (RA),

histone deacetylase inhibitors (HDAC), peroxisome proliferator-activated receptor-

gamma (PPARG) have been tested to promote redifferentiation and *NIS* re-expression of

RR-DTCs, to suggest RAI treatment after a short, targeted treatment[90,93–95]. This strategy

could lead to tumor response while limiting adverse effects and, several clinical trials are

ongoing. However, the re-expression of NIS is insufficient to explain the response to

redifferentiation therapeutics and RAI treatment. Membrane trafficking and the cellular

machinery that concentrate and retain iodine must be preserved[96] in a follicular

organization. Our model is proving to be a potential tool for redifferentiation studies

because of its ability to test large sets and combinations of treatments and to assess

follicular reorganization and iodide organification capacity using a functional assay that

is likely to be more strongly associated with response to RAI treatment than *NIS* mRNA

levels. Braf[V637E]-expressing cells treated with VPA, MAPK, PI3K, Dabrafenib,

Trametinib, and inflammation inhibitors were shown to restore the expression of *Nis*.

Interestingly, the combination of MAPK (BRAF and MEK) and PI3K inhibitors also

promoted the restoration of thyroid markers, follicular organization, and iodide

organification ability. Interestingly, an ongoing clinical trial tests the effect of BKM120,

a PI3K inhibitor, in patients with progressive, metastatic, refractory, follicular or poorly

differentiated thyroid cancers (NCT01830504). However, the association with MAPK

inhibitors is still not evaluated in patients with thyroid cancer.

In *BRAF*-mutated tumors, studies have shown that the combination of TKIs

therapies appears more effective and circumvents primary and acquired resistance to TKI

therapy. Often, resistance is due to reactivation of the MAPK/ERK pathway or activation

of other signaling pathways such as PTEN, NF-1 or RAS. It may also result from the

hyperactivation of tyrosine kinase receptors, such as PDFRβ, IGF-1R and HGF, which

lead to activation of the AKT/PI3K pathway[97]. Studies in patients with advanced

melanoma carrying a BRAF[V600E] mutation have shown that combining BRAF and MEK

inhibitors resulted in a higher rate of complete/partial responses and median progression-

free survival compared with monotherapy groups[98–100]. Such significant results lead to

the use of drug combinations as standard treatment for patients[101,102]. Likewise, phase II

clinical trials have demonstrated noteworthy response rates in advanced *BRAF*-mutated

thyroid carcinomas when treated with a combination of the BRAF inhibitor dabrafenib

and MEK inhibitor trametinib[70]. Furthermore, recent findings have shown that the

administration of dabrafenib-trametinib treatment followed by surgery can yield 24-

month overall survival (OS) rates as high as 80% in ATCs[103.]

In summary, we have developed a Braf^{V637E} oncogene-expressing thyroid cancer

organoid *in vitro* model from mESCs that recapitulates transcriptomic and histological

features of PTCs at early and advanced stages. Moreover, we demonstrated that the

combination of MEK and PI3K inhibitors promotes *Nis* re-expression and cell re-

differentiation leading to the restoration of follicular structures and thyroid functionality.

Considering the robustness of this *in vitro* model, which allows controlled induction of

the major thyroid oncogene in a three-dimensional system, its efficiency and simplicity,

our model can be used to study mechanisms associated with thyroid cancer development

and progression, thyroid redifferentiation, and drug screening.

MATERIAL AND METHODS

Cell culture and mESC_Braf^{V637E} line generation

The previously genetically modified G4 RosaLuc mouse embryonic stem cell line

(mESC)[104] was initially cultured on gamma-irradiated mouse embryonic fibroblasts

(MEFs) feeder using mouse stem cell medium[4,58] and incubated at 37 °C, 5 % CO^2 and

>95 % humidity. Cells were routinely tested for mycoplasma. To insert the target

sequences into the modified Rosa26 locus of G4 RosaLuc mESCs, around $1x10^5$ cells

were transfected using the Flpe expressing vector[104] and the target vector containing the

rtTA-TRE induction system, thyroid transcription factors, Nkx2-1 and Pax8 and bTg-

NES-Braf^{V637E}-ERT^2 sequences (Supplementary Fig. 1A) following the lipofectamine

3000 protocol (Thermo Scientific). The Braf^{V637E} mutation here used for mouse cells is

equivalent to the human BRAF^{V600E} mutation[56]. Briefly, $1x10^6$ mESCs were transfected

(in suspension) with 20 μg of each vector and cultured on Neomycin-resistant MEFs.

G418 (Neomycin; 300 μg/ml) was applied 48 h after transfection, for 10 days, and

individual clones were isolated (colonies were separated from the MEFs using insulin

needles) and further expanded. Clones were screened by immunofluorescence for Nkx2-

1 and Pax8 expression after three days of incubation with 1 µg/ml Doxycyclin (Dox)

(Supplementary Fig. 1B). Positive clones were then characterized according to

pluripotency maintenance (Supplementary Fig. 1C), *Nkx2-1* and *Pax8* expression, and

efficiency of thyroid differentiation. In addition, we generated an mESC control line

where the sequences of NES-BRAF^{V637E}-ERT^2 were replaced by the eGFP sequence,

resulting in the TRE-Nkx2-1-Pax8_bTg-eGFP line (Supplementary Fig. 1D). The mESC

control line generation, selection and characterization were performed as described above

(Supplementary Fig. 1E, F).

Thyroid differentiation protocol

G4   RosaLuc   TRE-Nkx2-1-Pax8_bTg-NES-Braf^{V637E}-ERT^2 and   TRE-Nkx2-1-

Pax8_bTg-eGFP cells were cultured and differentiated as described previously by

Antonica *et al.* (2012)[4] with few modifications. Briefly, modified mESCs cultured in

mESC media on top of MEFs were split using Trypsin EDTA (TE). Then, cells were

resuspended in mESC media[57] and seeded into a 10-cm petri dish for 30-45 min allowing

the attachment of most of the MEFs. mESC-enriched supernatant was collected,

centrifuged (500g for 5min), and resuspended in differentiation media[57], cells were

counted and finally diluted to 40,000 cells/ml. Embryoid bodies (EBs), were then

generated by hanging drops (25µl drops containing 1,000 cells), collected after four days

and embedded in growth factor reduced Matrigel (GFR MTG; BD Biosciences); 50µl

MTG drops (containing around 20 EBs) were plated into 12-wells plates. EBs were

differentiated using a differentiation medium (1 ml/well) initially supplemented with

1µg/ml of Doxycycline (Sigma) for five days, followed by two weeks of maturation

induction by using 0.3 µmol of 8-Br-cAMP (BioLog Life Science Institute) or 1mU ml^{-1}

of thyrotropin (hrTSH; Genzyme). The culture medium was changed every 2 days.

Thyroid differentiation and functionality were evaluated by transcriptomics (RT-PCR and

bulk RNA sequencing), immunofluorescence, and iodide organification.

Braf^{V637E} induction and drug screening

After full thyroid maturation (Day 23), MTG drops were washed twice with Hanks's

balanced salt solution (HBSS, containing calcium and magnesium; Invitrogen) and

incubated in a HBSS solution (1 ml per well) containing 10 U/ml of dispase II (Roche)

and 125 U/ml of collagenase type IV (Sigma) for 30 min at 37°C. Then the enzymes were

inactivated by adding 10% FBS and cells were centrifuged at 500g for 3 min. Cells were

rinsed twice with HBSS, and the follicle population was enriched by filtering using 30

μm (single cell removal) and 100 μm reverse strainer (Pluriselect). Finally, enriched

structures were re-embedded in GFR MTG (50μl) and plated into 12-well plates. Twenty-

four hours later, cells were incubated with 1 μM of 4-Hydroxytamoxifen (4OHT) (Sigma)

and 8-Br-cAMP or hrTSH for 24h, 48h,7 days or 3 weeks to promote Braf^{V637E}-induced

phosphorylation of MAPK pathway[56]. The control condition, treated with cAMP and/or

hrTSH, was also treated with vehicle ethanol for 4OHT treatment comparisons.

Braf^{V637E}-induced thyroid organoids (previously treated for four days with 4OHT) were

also cultured in addition to distinct compounds, previously suggested to regulate Nis re-

expression and inhibit Braf^{V637E} oncogenic effects[68,69]. Among the screened drugs,

several cell processes were targeted by the following compounds: PD0325901 (MEK

inhibitor; 250nM; Stem Cell), LY204002 (PI3K inhibitor; 5μM; Selleckchem), VPA

(HDAC inhibitor; 250μM; Sigma), NAC (N-acetyl cysteine, antioxidant compound;

2mM; Sigma), 5-AZA-2' deoxycytidine (DNA methyltransferase (DMNT) inhibitor;

1μM; Sigma) and RG108 (DMNT inhibitor; 10μM; Sigma). Furthermore, experiments

were also conducted using inhibitors that are already clinically approved for the treatment

of thyroid cancer, namely dabrafenib (BRAF inhibitor; 100nM; Selleckchem) and/or

trametinib (MEK inhibitor; 20nM; Selleckchem).

Disrupted pathways identified by transcriptomics analysis in 4OHT condition compared

to cAMP control were inhibited/activated by treatment with dexamethasone (anti-

inflammatory drug; 50nM; Sigma), CC-5013 (Lenalidomide; TNF-alpha inhibitor;

10μM; Selleckchem) and CHIR- 99021(Wnt/β-catenin activator; 3μM; Selleckchem). All

inhibitor treatments involved organoids that had previously been exposed to

cAMP+4OHT for four days, and they were subsequently subjected to continuous

incubation with cAMP+4OHT+inhibitors for an additional 3 days. Culture media was

replenished every 2 days. All experiments were performed following the same time points

of incubation described above.

Flow cytometry

Proliferation assay: Organoids subjected to various treatments, including controls

(cAMP), 4OHT, and 4OHT+inhibitors, at both 7 days (day 30) and 21 days (day 54),

were exposed to BrdU (1mM) for a duration of 3 hours. Subsequently, a proliferation

assay was conducted in accordance with the BrdU Flow Kit staining protocol (BD)

instructions. Briefly, Matrigel drops (pool of at least four replicates per condition) were

first digested with HBSS solution containing 10 U/ml dispase II and 125 U/ml

collagenase type IV for 30-60 min at 37°C; then a single cell suspension was obtained by

dissociation with TripLE Express (10-15 min incubation at 37°C), the enzymes were

inactivated by addition of differentiation medium. After centrifugation (500 g for 3 min),

samples were rinsed with PBS, and then the BrdU Flow Kit staining protocol (BD

Biosciences) was used according to the manufacturer's instructions. In order to identify

the BrdU incorporated cells among the thyroid population, cells were also stained using

primary Nkx2-1 antibody (1:100; Abcam) and donkey anti-rabbit IgG Cy3-conjugated

(1:300; Jackson Immunoresearch). As controls, we used cells untreated with BrdU and

stained with the APC anti-BrdU antibody for BrdU gating while for Nkx2.1 gating, cells

were incubated only with the secondary antibody. Data are presented as a percentage of

double-positive (BrdU/Nkx2.1) cells. For each experiment, at least four wells from each

condition were used.

Apoptosis assay: Organoids subjected to various treatments, including controls (cAMP),

4OHT, and 4OHT+inhibitors, at both 7 days (day 30) and 21 days (day 54) were collected

and dissociated into single cells following the protocol described above (proliferation

assay   section).   Cells   were   then   fixed   and   permeabilized   using   the   BD

Cytofix/Cytoperm™ Fixation/Permeabilization Kit (BD Biosciences) and stained using

Nkx2-1 antibody (1:100; Abcam) and donkey anti-rabbit IgG Cy3-conjugated secondary

antibody (1:300; Jackson Immunoresearch) combined with FITC Rabbit Anti-Active

Caspase-3 conjugated antibody (1:5; BD Biosciences). As a positive control, cAMP-

treated organoids were incubated with Staurosporine (10nM; Sigma) for 16 hours and

collected at the respective time points following the procedures described above.

Untreated cells and cells incubated only with the secondary antibody (isotype) were used

as negative controls for the gating strategy. For each experiment, at least four wells from

each condition were used.

Proportion of bTg-GFP cells: The proportions of Nkx2.1 and GFP cells in the control

mESC line (TRE-Nkx2-1-Pax8_bTg-eGFP) were evaluated at day 30. Controls (cAMP)

and 4OHT-treated (7 days) organoids were dissociated and stained for Nkx2.1 following

the steps described above (apoptosis assay session). GFP detection was based on the

endogenous expression (bTg-eGFP). Undifferentiated cells (mESCs) and cells incubated

only with the secondary antibody (isotype) were used as negative controls for the gating

strategy. For each experiment, at least four wells from each condition were used.

All analyses were performed using the LSR-Fortessa X-20 flow cytometer and BD

FACSDiva software was used for the quantification analysis. Data are presented as a

percentage of positive cells.

Immunofluorescence staining

Organoids embedded in MTG were fixed in 4% paraformaldehyde (PFA; Sigma) for 1 h

at room temperature (RT) and washed three times in PBS. Samples were either stained as

whole mount (MTG drop) or embedded in 4% low-melting agarose and cut using

Vibratome (80-100µm; Sigma). Blocking was performed using a solution of PBS

containing 3% bovine serum albumin (BSA; Sigma), 5% horse serum (Invitrogen) and

0.3% Triton X-100 (Sigma) for 30 min at RT. The primary and secondary antibodies were

diluted in a solution of PBS containing 3% BSA, 1% horse serum and 0.1% Triton X-

100. Primary antibodies were incubated overnight at 4°C followed by incubation with

secondary antibodies and Hoechst for 2 h at RT. Slides were mounted with Glycergel

(Dako).

For paraffin embedding, organoids were fixed in 4% PFA for 1 h at 4°C and kept in 70%

ethanol at 4°C before processing. Samples were then embedded in paraffin, sectioned (5

µm), mounted on glass slides, deparaffinized, and rehydrated following standard

protocols. For immunostaining, antigen retrieval was performed by incubating the

sections for 10 min in the microwave (850 W) in Sodium Citrate Buffer (10 mM Sodium

Citrate, 0.05% Tween 20, pH 6.0). After cooling, the sections were rinsed with PBS and

blocked with 1% BSA and 10% horse serum PBS solution for 1 h at RT. Primary

antibodies (Supplementary Table 1) were diluted in the blocking solution and incubated

overnight at 4°C. The sections were rinsed three times in PBS and incubated with Hoechst

and secondary antibodies diluted in blocking solution for 1 h at RT. Slides were mounted

with Glycergel (Dako). Imaging was performed using a Zeiss LSM510 META confocal

microscope and a Leica DMI6000 microscope with DFC365FX camera. The antibodies

specifications and dilutions are listed in the supplementary Table 1.

Western blot

Follicle enriched-organoids were initially isolated (collagenase IV/dispase II solution,

described above) from at least 6 distinct wells and pooled together for protein extraction

using RIPA buffer. The protein concentration was determined according to Pierce 660

nm protein assay reagent protocol (Thermo Scientific). Briefly, for each sample, 30–50

μg of protein was fractionated by 10% SDS–PAGE and blotted onto an Immobilon PVDF

membrane (Millipore). Non-specific binding sites were blocked by incubation with 5%

BSA in Tris-buffered saline (TBS) 0.1% Tween-20 (TBS-T) for 1 h. Thereafter, the

membrane was incubated overnight at 4 °C with primary antibodies against phospho-

ERK1/2 (1:1,000; Cell Signaling), ERK1/2 (1:400; Santa Cruz Biotechnology), and β-

Actin (1:1,1000; Cell Signaling) in a blocking solution. Next, the membrane was

incubated with horseradish peroxidase-conjugated anti-rabbit (1:1,000; Cell Signaling) or

anti-mouse antibody (1:1,000; Cell Signaling) in a blocking solution, for 60 min at RT.

The antigen-antibody complexes were visualized using an enhanced chemiluminescence

system (GE Healthcare) and captured by Azure 500 system (Azure Biosystems).

Iodide organification assay

Thyroid organoids treated with cAMP, cAMP+4OHT, cAMP+4OHT+PD+LY and

cAMP+4OHT+PD+LY+VPA were tested for the ability of iodide uptake and

organification as previously described[4,58]. Briefly, cells were washed with HBSS and

incubated with 1 ml per well of an organification medium containing 1,000,000 c.p.m.

per ml ^{125}I (PerkinElmer) and 100 nM sodium iodide (NaI, Sigma) in HBSS for 2 h at 37

°C. The reaction was stopped by the addition of 1 ml of methimazole (MMI; 4 mM,

Sigma). After two washes with cold PBS, organoids were dissociated by incubation with

trypsin/EDTA (Invitrogen) for 10 min at 37 °C. For iodide uptake quantification, cells

were collected, and radioactivity was measured using a gamma-counter. Subsequently,

proteins were precipitated by adding 100 μl of gamma-globulins (10 mg/ml; Sigma) and

2 ml of 20% TCA followed by centrifugation at 2,000 r.p.m. for 10 min, at 4°C and the

radioactivity of protein-bound ^{125}I (PBI) was measured. Iodide organification was

calculated as an iodide uptake/PBI ratio and, the values were expressed as a percentage.

Cells were also treated with 1 mM sodium perchlorate (Nis inhibitor; NaClO4, Sigma-

Aldrich) and 2 mM MMI (Tpo inhibitor; Sigma-Aldrich) as iodide uptake and protein-

binding controls, respectively. The experiments were performed in triplicates for each

condition.

Gene expression analysis

Real-time PCR (RT-qPCR) was performed on cDNA from thyroid organoids from at least

three independent experiments. Total RNA was extracted from thyroid organoids by the

addition of lysis RLT Lysis buffer (Qiagen) + 1% 2-mercaptoethanol directly on the MTG

drop containing the organoids. For longer cultures (21 days), organoids were initially

incubated with a collagenase IV/Dispase II solution (described above) for 30-45 min.

Enzymes were inactivated by the addition of differentiation medium, cells were

centrifuged (500 g for 3 min), washed with PBS, and resuspended in RLT buffer. RNA

was isolated using the RNeasy micro kit (Qiagen) according to the manufacturer's

instructions. cDNA was generated by reverse transcription using the Superscript II kit

(Invitrogen). qPCR was performed in triplicates on cDNA (1:10 dilution) using Takyon™

No ROX SYBR 2X MasterMix blue dTTP (Eurogentec) and CFX Connect Real-Time

System (Biorad). Results are presented as linearized values normalized to the

housekeeping gene, β2-microglobulin and the indicated reference value (2-DDCt). cAMP

condition was used as the control for all the comparisons. Primer sequences are described

in Supplementary Table 2.

Bulk RNA sequencing and analysis

Bulk RNA-seq was performed on day 31, using the cAMP, cAMP+4OHT (5 days) and

cAMP+4OHT+PD+LY (3 days) conditions (see protocol, Fig. 5A). RNA extraction was

performed as previously described (section "*Gene expression analysis*"), and

experiments were performed in duplicates. At least three distinct wells were pooled

together for each condition. RNA quality and concentration were evaluated using

Bioanalyser 2100 (Agilent) and RNA 6000 Nano Kit (Agilent). RNA integrity was

preserved, and no genomic DNA contamination was detected. Indexed cDNA libraries

were obtained using the TruSeq Stranded mRNA Sample Prep kit (Illumina) using an S2

flow cell, and sequences were produced using a 200 cycles reagent kit. The resulting high-

quality indexed cDNA libraries were quantified with the Quant-iT PicoGreen kit (Life

Sciences) and Infinite F200 Proplate reader (Tecan); DNA fragment size distribution was

examined with the 2100 Bioanalyzer (Agilent) using the DNA 1000 kit (Agilent).

Multiplexed libraries (10ρM) were loaded onto flow cells and sequenced on the HiSeq

1500 system (Illumina) in high-output mode using the HiSeq Cluster Kit v4 (Illumina).

The sequenced data were uploaded on the galaxy web platform version 22.05.1, and the

public server, Data independent acquisition proteomics workbench (RRID:SCR_021862,

https://usegalaxy.eu), was used for mapping and counting analysis. Approximately 10

million paired-end reads were obtained per sample. After the removal of low-quality

bases and Illumina adapter sequences using Trimmomatic software[105]

(RRID:SCR_011848), paired-end reads were mapped against the mouse reference

genome mm10 (GRCm38) using HISAT2 sofware[106], allowing soft clipping. Raw read

counts were obtained using HTseq count software with unstranded option[107]. Low-

expressed genes were filtered using the EdgeR package in R Project for Statistical

Computing (https://www.r-project.org). Then, differential expression analysis was

performed using DESeq2 package[108]. Fold change and adjusted p-value thresholds of 2

and 0.05, respectively were used to select differentially expressed genes (DEG). Gene

ontology and pathways enrichment analysis in up and down differentially expressed gene

lists was realized using EnrichR (https://maayanlab.cloud/Enrichr/). Transformed counts

corresponding to log2 (normalized counts +4) using the DESeq2 package were used for

further analysis and heatmap visualization. Thyroid differentiation score (TDS),

Enhanced Thyroid differentiation score (eTDS) and ERK activation score were calculated

as previously described[67]. The list of the genes used for the score's calculation is provided

in Supplementary Table 3.

Data availability

Bulk RNA-seq data have been deposited in the NCBI Gene Expression Omnibus (GEO;

RRID:SCR_005012) under accession number GSE228281.

Statistical analysis

All statistical analysis were performed using GraphPad Prism 9. Comparison between

two groups and its statistical significance was tested using the nonparametric Mann-

Whitney U test. In contrast, comparisons between multiple groups was performed using

the nonparametric Kruskal-Wallis test. Data are displayed as median (IQR). Differences

were considered significant at p<0.05 and presented as follows: *p < 0.05, **p < 0.01,

***p < 0.001, ****p < 0.0001. All the data presented are from at least three independent

experiments.

Acknowledgements and Funding

The authors acknowledge the ULB flow cytometry platform (C. Dubois), the ULB

genomic core facility (F. Libert and A. Lefort), LiMIF platform for confocal microscopy

(J.-M. Vanderwinden) and Eduardo Andrés Rios Morris for the contribution with the

imaging. We acknowledge the funding agencies that supported this study. The Belgian

National Fund for Scientific Research (FNRS) (PDR T.0140.14; PDR T.0230.18, CDR

J.0068.22, Televie 7.4633.17/7.4526.19) and the Fonds d'Encouragement à la Recherche

de l'Université Libre de Bruxelles (FER-ULB). The Belgian Fondation contre le cancer

(F/2020/1402; V.D.). FNRS (Chargé de Recherche, No.825745; M.R.). The cooperation

program CAPES-WBI (S.C. and A.L.M). The Brazilian National Council for Scientific

and Technological Development (CNPq; M.R. and A.L.M), the Coordination for the

Improvement of Higher Education Personnel (CAPES; M.R. and A.L.M), the Brazilian

Society of Endocrinology and Metabolism (SBEM; M.R. and A.L.M) and the Fundação

de Amparo à Pesquisa do Estado do Rio Grande do Sul (FAPERGS; A.L.M). S.C is

Research Director at FNRS. M.R is researcher at ULB.

Author contributions

H.L., S.C. and M.R. developed the project, designed the experiments and analyzed the

data. L.H., T.P., S.G. and J.H. generated the plasmids and the G4 Rosaluc mESC line.

M.R. and A.S. generated the G4 RosaLuc TRE-Nkx2-1-Pax8_bTg-NES-Braf^{V637E}-ERT^2

and TRE-Nkx2-1-Pax8_bTg-eGFP cell lines. H.L., M.R., B.F.F., A.S., O.M., L.C.,

M.K.P. B.A., L.C., performed the *in vitro* experiments and protocol set up. M.R. and P.G.

obtained confocal images. H.L., M.R. and A.T. performed bulk RNA-Sequencing and

analyzed the results. H.L. and A.T. performed the bioinformatics analysis. H.L. and M.R.

wrote the first draft and, A.L.M. and S.C. edited the manuscript. S.C., M.R. and A.L.M.

acquired funding for the project. All authors contributed to the article and approved the

submitted version.

Competing interests

All authors declare no competing interests.

Correspondence and requests for materials should be addressed to Mírian Romitti.

REFERENCES

1   Schutgens F, Clevers H. Human Organoids: Tools for Understanding Biology
and Treating Diseases. *Annual Review of Pathology: Mechanisms of Disease*
2020; **15**: 211–234.
2   Clevers H. Modeling Development and Disease with Organoids. *Cell* 2016; **165**:
1586–1597.
3   Kim J, Koo BK, Knoblich JA. Human organoids: model systems for human
biology and medicine. Nat Rev Mol Cell Biol. 2020; **21**: 571–584.
4   Antonica F, Kasprzyk DF, Opitz R, Iacovino M, Liao XH, Dumitrescu AM *et al.*
Generation of functional thyroid from embryonic stem cells. *Nature* 2012; **491**:
66–71.
5   Romitti M, Tourneur A, de Faria da Fonseca B, Doumont G, Gillotay P, Liao X-
H *et al.* Transplantable human thyroid organoids generated from embryonic stem
cells to rescue hypothyroidism. *Nat Commun* 2022; **13**: 7057.
6   Longmire TA, Ikonomou L, Hawkins F, Christodoulou C, Cao Y, Jean JC *et al.*
Efficient derivation of purified lung and thyroid progenitors from embryonic
stem cells. *Cell Stem Cell* 2012; **10**: 398–411.
7   Ma R, Latif R, Davies TF. Human embryonic stem cells form functional thyroid
follicles. *Thyroid* 2015; **25**: 455–461.
8   Ma R, Latif R, Davies TF. Thyroid follicle formation and thyroglobulin
expression in multipotent endodermal stem cells. *Thyroid* 2013; **23**: 385–391.
9   Ma R, Morshed SA, Latif R, Davies TF. TAZ Induction Directs Differentiation
of Thyroid Follicular Cells from Human Embryonic Stem Cells. *Thyroid* 2017;
27: 292–299.

10   Ma R, Shi R, Morshed SA, Latif R, Davies TF. Derivation and 97% Purification
of Human Thyroid Cells From Dermal Fibroblasts. *Front Endocrinol (Lausanne)*
2020; **11**: 446.

11   Kurmann AA, Serra M, Hawkins F, Rankin SA, Mori M, Astapova I *et al.*
Regeneration of Thyroid Function by Transplantation of Differentiated
Pluripotent Stem Cells. *Cell Stem Cell* 2015; **17**: 527–542.

12   Serra M, Alysandratos KD, Hawkins F, McCauley KB, Jacob A, Choi J *et al.*
Pluripotent stem cell differentiation reveals distinct developmental pathways
regulating Lung-Versus Thyroid-Lineage specification. *Development*
(Cambridge) 2017; **144**: 3879–3893.

13   Dame K, Cincotta S, Lang AH, Sanghrajka RM, Zhang L, Choi J *et al.* Thyroid
Progenitors Are Robustly Derived from Embryonic Stem Cells through
Transient, Developmental Stage-Specific Overexpression of Nkx2-1. *Stem Cell*
Reports 2017; **8**: 216–225.

14   Ogundipe VML, Groen AH, Hosper N, Nagle PWK, Hess J, Faber H *et al.*
Generation and Differentiation of Adult Tissue-Derived Human Thyroid
Organoids. *Stem Cell Reports* 2021; **16**: 913–925.

15   Liang J, Qian J, Yang L, Chen X, Wang X, Lin X *et al.* Modeling Human
Thyroid Development by Fetal Tissue-Derived Organoid Culture. *Advanced*
Science 2022; **9**. doi:10.1002/advs.202105568.

16   van der Vaart J, Bosmans L, Sijbesma SF, Knoops K, van de Wetering WJ, Otten
HG *et al.* Adult mouse and human organoids derived from thyroid follicular cells
and modeling of Graves' hyperthyroidism. *Proceedings of the National Academy*
of Sciences 2021; **118**. doi:10.1073/pnas.2117017118.

17   Boj SF, Hwang C-I, Baker LA, Chio IIC, Engle DD, Corbo V *et al.* Organoid
Models of Human and Mouse Ductal Pancreatic Cancer. *Cell* 2015; **160**: 324–
338.

18   Gao D, Vela I, Sboner A, Iaquinta PJ, Karthaus WR, Gopalan A *et al.* Organoid
Cultures Derived from Patients with Advanced Prostate Cancer. *Cell* 2014; **159**:
176–187.

19   Sato T, Stange DE, Ferrante M, Vries RGJ, van Es JH, van den Brink S *et al.*
Long-term Expansion of Epithelial Organoids From Human Colon, Adenoma,
Adenocarcinoma, and Barrett's Epithelium. *Gastroenterology* 2011; **141**: 1762–
1772.

20   van de Wetering M, Francies HE, Francis JM, Bounova G, Iorio F, Pronk A *et al.*
Prospective Derivation of a Living Organoid Biobank of Colorectal Cancer
Patients. *Cell* 2015; **161**: 933–945.

21   Drost J, Clevers H. Organoids in cancer research. *Nat Rev Cancer* 2018; **18**: 407–
418.

22   Gleave AM, Ci X, Lin D, Wang Y. A synopsis of prostate organoid
methodologies, applications, and limitations. *Prostate* 2020; **80**: 518–526.

23   Maenhoudt N, Vankelecom H. Protocol for establishing organoids from human
ovarian cancer biopsies. *STAR Protoc* 2021; **2**: 100429.

24   Gunti S, Hoke ATK, Vu KP, London NR. Organoid and Spheroid Tumor
Models: Techniques and Applications. *Cancers (Basel)* 2021; **13**: 874.

25   Saito Y, Onishi N, Takami H, Seishima R, Inoue H, Hirata Y *et al.* Development
of a functional thyroid model based on an organoid culture system. *Biochem*
Biophys Res Commun 2018; **497**: 783–789.

808	26	Chen D, Su X, Zhu L, Jia H, Han B, Chen H *et al.* Papillary thyroid cancer
809		organoids harboring BRAFV600E mutation reveal potentially beneficial effects
810		of BRAF inhibitor-based combination therapies. *J Transl Med* 2023; **21**: 9.
811	27	Vlachogiannis G, Hedayat S, Vatsiou A, Jamin Y, Fernández-Mateos J, Khan K
812		*et al.* Patient-derived organoids model treatment response of metastatic
813		gastrointestinal cancers. *Science (1979)* 2018; **359**: 920–926.
814	28	Drost J, van Jaarsveld RH, Ponsioen B, Zimberlin C, van Boxtel R, Buijs A *et al.*
815		Sequential cancer mutations in cultured human intestinal stem cells. *Nature*
816		2015; **521**: 43–47.
817	29	Matano M, Date S, Shimokawa M, Takano A, Fujii M, Ohta Y *et al.* Modeling
818		colorectal cancer using CRISPR-Cas9–mediated engineering of human intestinal
819		organoids. *Nat Med* 2015; **21**: 256–262.
820	30	Miura A, Yamada D, Nakamura M, Tomida S, Shimizu D, Jiang Y *et al.*
821		Oncogenic potential of human pluripotent stem cell-derived lung organoids with
822		<scp>HER2</scp> overexpression. *Int J Cancer* 2021; **149**: 1593–1604.
823	31	Nadauld LD, Garcia S, Natsoulis G, Bell JM, Miotke L, Hopmans ES *et al.*
824		Metastatic tumor evolution and organoid modeling implicate TGFBR2as a cancer
825		driver in diffuse gastric cancer. *Genome Biol* 2014; **15**: 428.
826	32	Li X, Nadauld L, Ootani A, Corney DC, Pai RK, Gevaert O *et al.* Oncogenic
827		transformation of diverse gastrointestinal tissues in primary organoid culture. *Nat*
828		*Med* 2014; **20**: 769–777.
829	33	Huang L, Holtzinger A, Jagan I, BeGora M, Lohse I, Ngai N *et al.* Ductal
830		pancreatic cancer modeling and drug screening using human pluripotent stem
831		cell– and patient-derived tumor organoids. *Nat Med* 2015; **21**: 1364–1371.
832	34	Lo Y-H, Karlsson K, Kuo CJ. Applications of organoids for cancer biology and
833		precision medicine. *Nat Cancer* 2020; **1**: 761–773.
834	35	Lim H, Devesa SS, Sosa JA, Check D, Kitahara CM. Trends in Thyroid Cancer
835		Incidence and Mortality in the United States, 1974-2013. *JAMA* 2017; **317**: 1338.
836	36	ROMITTI M, CEOLIN L, SIQUEIRA DR, FERREIRA CV, WAJNER SM,
837		MAIA AL. Signaling pathways in follicular cell-derived thyroid carcinomas. *Int*
838		*J Oncol* 2013; **42**: 19–28.
839	37	Kimura ET, Nikiforova MN, Zhu Z, Knauf JA, Nikiforov YE, Fagin JA. High
840		prevalence of BRAF mutations in thyroid cancer: genetic evidence for
841		constitutive activation of the RET/PTC-RAS-BRAF signaling pathway in
842		papillary thyroid carcinoma. *Cancer Res* 2003; **63**: 1454–7.
843	38	Fagin JA, Mitsiades N. Molecular pathology of thyroid cancer: diagnostic and
844		clinical implications. *Best Pract Res Clin Endocrinol Metab* 2008; **22**: 955–969.
845	39	Scheffel RS, Cristo AP de, Romitti M, Vargas CVF, Ceolin L, Zanella AB *et al.*
846		The BRAFV600E mutation analysis and risk stratification in papillary thyroid
847		carcinoma. *Arch Endocrinol Metab* 2020. doi:10.20945/2359-3997000000285.
848	40	Xing M. The T1799A BRAF mutation is not a germline mutation in familial
849		nonmedullary thyroid cancer. *Clin Endocrinol (Oxf)* 2005; **63**: 263–266.
850	41	Chakravarty D, Santos E, Ryder M, Knauf JA, Liao X-H, West BL *et al.* Small-
851		molecule MAPK inhibitors restore radioiodine incorporation in mouse thyroid
852		cancers with conditional BRAF activation. *Journal of Clinical Investigation*
853		2011; **121**: 4700–4711.
854	42	Zaman, Wu, Bivona. Targeting Oncogenic BRAF: Past, Present, and Future.
855		*Cancers (Basel)* 2019; **11**: 1197.

43   Agrawal N, Akbani R, Aksoy BA, Ally A, Arachchi H, Asa SL *et al.* Integrated
Genomic Characterization of Papillary Thyroid Carcinoma. *Cell* 2014; **159**: 676–
690.

44   Romei C, Fugazzola L, Puxeddu E, Frasca F, Viola D, Muzza M *et al.*
Modifications in the Papillary Thyroid Cancer Gene Profile Over the Last 15
Years. *J Clin Endocrinol Metab* 2012; **97**: E1758–E1765.

45   Sabra MM, Dominguez JM, Grewal RK, Larson SM, Ghossein RA, Tuttle RM *et*
al. Clinical Outcomes and Molecular Profile of Differentiated Thyroid Cancers
With Radioiodine-Avid Distant Metastases. *J Clin Endocrinol Metab* 2013; **98**:
E829–E836.

46   Xing M. BRAFV600E Mutation and Papillary Thyroid Cancer. *JAMA* 2013; **310**:
535.

47   Scheffel RS, Cristo AP de, Romitti M, Vargas CVF, Ceolin L, Zanella AB *et al.*
The BRAFV600E mutation analysis and risk stratification in papillary thyroid
carcinoma. *Arch Endocrinol Metab* 2021. doi:10.20945/2359-3997000000285.

48   Liu R, Bishop J, Zhu G, Zhang T, Ladenson PW, Xing M. Mortality Risk
Stratification by Combining *BRAF* V600E and *TERT* Promoter Mutations in
Papillary Thyroid Cancer. *JAMA Oncol* 2017; **3**: 202.

49   Boucai L, Seshan V, Williams M, Knauf JA, Saqcena M, Ghossein RA *et al.*
Characterization of Subtypes of BRAF-Mutant Papillary Thyroid Cancer Defined
by Their Thyroid Differentiation Score. *J Clin Endocrinol Metab* 2022; **107**:
1030–1039.

50   Chen D, Tan Y, Li Z, Li W, Yu L, Chen W *et al.* Organoid Cultures Derived
From Patients With Papillary Thyroid Cancer. *J Clin Endocrinol Metab* 2021;
106: 1410–1426.

51   Sondorp LHJ, Ogundipe VML, Groen AH, Kelder W, Kemper A, Links TP *et al.*
Patient-Derived Papillary Thyroid Cancer Organoids for Radioactive Iodine
Refractory Screening. *Cancers (Basel)* 2020; **12**: 3212.

52   Pecce V, Sponziello M, Bini S, Grani G, Durante C, Verrienti A. Establishment
and maintenance of thyroid organoids from human cancer cells. *STAR Protoc*
2022; **3**: 101393.

53   Chen D, Su X, Zhu L, Jia H, Han B, Chen H *et al.* Papillary thyroid cancer
organoids harboring BRAFV600E mutation reveal potentially beneficial effects
of BRAF inhibitor-based combination therapies. *J Transl Med* 2023; **21**: 9.

54   Yang H, Liang Q, Zhang J, Liu J, Wei H, Chen H *et al.* Establishment of
papillary thyroid cancer organoid lines from clinical specimens. *Front*
Endocrinol (Lausanne) 2023; **14**. doi:10.3389/fendo.2023.1140888.

55   Veschi V, Turdo A, Modica C, Verona F, Di Franco S, Gaggianesi M *et al.*
Recapitulating thyroid cancer histotypes through engineering embryonic stem
cells. *Nat Commun* 2023; **14**: 1351.

56   Ortiz O, Wurst W, Kühn R. Reversible and tissue-specific activation of MAP
kinase signaling by tamoxifen in braf [V637] ER [T2] mice. *genesis* 2013; **51**: 448–
455.

57   Antonica F, Kasprzyk DF, Schiavo AA, Romitti M, Costagliola S. Generation of
Functional Thyroid Tissue Using 3D-Based Culture of Embryonic Stem Cells.
2017, pp 85–95.

58   Romitti M, Eski SE, Fonseca BF, Gillotay P, Singh SP, Costagliola S. Single-
Cell Trajectory Inference Guided Enhancement of Thyroid Maturation In Vitro
Using TGF-Beta Inhibition. *Front Endocrinol (Lausanne)* 2021; **12**.
doi:10.3389/fendo.2021.657195.

59 Knauf JA, Ma X, Smith EP, Zhang L, Mitsutake N, Liao X-H *et al.* Targeted
Expression of BRAFV600E in Thyroid Cells of Transgenic Mice Results in
Papillary Thyroid Cancers that Undergo Dedifferentiation. *Cancer Res* 2005; **65**:
4238–4245.

60 Schoultz E, Liang S, Carlsson T, Filges S, Ståhlberg A, Fagman H *et al.* Tissue
specificity of oncogenic BRAF targeted to lung and thyroid through a shared
lineage factor. *iScience* 2023; **26**: 107071.

61 Tang W-W, Huang C, Tang C, Xu J, Wang H. Galectin-3 may serve as a
potential marker for diagnosis and prognosis in papillary thyroid carcinoma: a
meta-analysis. *Onco Targets Ther* 2016; : 455.

62 Xia S, Wang C, Postma EL, Yang Y, Ni X, Zhan W. Fibronectin 1 promotes
migration and invasion of papillary thyroid cancer and predicts papillary thyroid
cancer lymph node metastasis. *Onco Targets Ther* 2017; **Volume 10**: 1743–1755.

63 Sponziello M, Rosignolo F, Celano M, Maggisano V, Pecce V, De Rose RF *et al.*
Fibronectin-1 expression is increased in aggressive thyroid cancer and favors the
migration and invasion of cancer cells. *Mol Cell Endocrinol* 2016; **431**: 123–132.

64 Reischmann N, Andrieux G, Griffin R, Reinheckel T, Boerries M, Brummer T.
BRAFV600E drives dedifferentiation in small intestinal and colonic organoids
and cooperates with mutant p53 and Apc loss in transformation. *Oncogene* 2020;
39: 6053–6070.

65 Chandeck C, Mooi WJ. Oncogene-induced Cellular Senescence. *Adv Anat Pathol*
2010; **17**: 42–48.

66 Bellelli R, Vitagliano D, Federico G, Marotta P, Tamburrino A, Salerno P *et al.*
Oncogene-induced senescence and its evasion in a mouse model of thyroid
neoplasia. *Mol Cell Endocrinol* 2018; **460**: 24–35.

67 Dunn LA, Sherman EJ, Baxi SS, Tchekmedyian V, Grewal RK, Larson SM *et al.*
Vemurafenib Redifferentiation of *BRAF* Mutant, RAI-Refractory Thyroid
Cancers. *J Clin Endocrinol Metab* 2019; **104**: 1417–1428.

68 Kogai T, Brent GA. The sodium iodide symporter (NIS): Regulation and
approaches to targeting for cancer therapeutics. *Pharmacol Ther* 2012; **135**: 355–
370.

69 Azouzi N, Cailloux J, Cazarin JM, Knauf JA, Cracchiolo J, Al Ghuzlan A *et al.*
NADPH Oxidase NOX4 Is a Critical Mediator of BRAF [V600E]-Induced
Downregulation of the Sodium/Iodide Symporter in Papillary Thyroid
Carcinomas. *Antioxid Redox Signal* 2017; **26**: 864–877.

70 Subbiah V, Kreitman RJ, Wainberg ZA, Cho JY, Schellens JHM, Soria JC *et al.*
Dabrafenib plus trametinib in patients with BRAF V600E-mutant anaplastic
thyroid cancer: updated analysis from the phase II ROAR basket study. *Annals of*
Oncology 2022; **33**: 406–415.

71 Leboulleux S, Do Cao C, Zerdoud S, Attard M, Bournaud C, Lacroix L *et al.* A
Phase II Redifferentiation Trial with Dabrafenib-Trametinib and 131I in
Metastatic Radioactive Iodine Refractory *BRAF* p.V600E-Mutated Differentiated
Thyroid Cancer. *Clinical Cancer Research* 2023; **29**: 2401–2409.

72 Califano I, Smulever A, Jerkovich F, Pitoia F. Advances in the management of
anaplastic thyroid carcinoma: transforming a life-threatening condition into a
potentially treatable disease. *Rev Endocr Metab Disord* 2023.
doi:10.1007/s11154-023-09833-1.

73 Saqcena M, Leandro-Garcia LJ, Maag JLV, Tchekmedyian V, Krishnamoorthy
GP, Tamarapu PP *et al.* SWI/SNF Complex Mutations Promote Thyroid Tumor

955		Progression and Insensitivity to Redifferentiation Therapies. *Cancer Discov*
956		2021; **11**: 1158–1175.
957	74	Qiu W, Yang Z, Fan Y, Zheng Q. ZNRF3 is downregulated in papillary thyroid
958		carcinoma and suppresses the proliferation and invasion of papillary thyroid
959		cancer cells. *Tumor Biology* 2016; **37**: 12665–12672.
960	75	Han J, Chen M, Wang Y, Gong B, Zhuang T, Liang L *et al.* Identification of
961		Biomarkers Based on Differentially Expressed Genes in Papillary Thyroid
962		Carcinoma. *Sci Rep* 2018; **8**: 9912.
963	76	Yoo S-K, Song YS, Lee EK, Hwang J, Kim HH, Jung G *et al.* Integrative
964		analysis of genomic and transcriptomic characteristics associated with
965		progression of aggressive thyroid cancer. *Nat Commun* 2019; **10**: 2764.
966	77	Davies H, Bignell GR, Cox C, Stephens P, Edkins S, Clegg S *et al.* Mutations of
967		the BRAF gene in human cancer. *Nature* 2002; **417**: 949–954.
968	78	Dhomen N, Marais R. New insight into BRAF mutations in cancer. *Curr Opin*
969		*Genet Dev* 2007; **17**: 31–39.
970	79	Nikiforov YE, Nikiforova MN. Molecular genetics and diagnosis of thyroid
971		cancer. *Nat Rev Endocrinol* 2011; **7**: 569–580.
972	80	Cheng L, Lopez-Beltran A, Massari F, MacLennan GT, Montironi R. Molecular
973		testing for BRAF mutations to inform melanoma treatment decisions: a move
974		toward precision medicine. *Modern Pathology* 2018; **31**: 24–38.
975	81	Schirripa M, Biason P, Lonardi S, Pella N, Pino MS, Urbano F *et al.* Class 1, 2,
976		and 3 BRAF-Mutated Metastatic Colorectal Cancer: A Detailed Clinical,
977		Pathologic, and Molecular Characterization. *Clinical Cancer Research* 2019; **25**:
978		3954–3961.
979	82	Pisapia P, Pepe F, Malapelle U, Troncone G. Mutations in Lung Cancer. *Acta*
980		*Cytol* 2019; **63**: 247–250.
981	83	Pisapia P, Pepe F, Iaccarino A, Sgariglia R, Nacchio M, Russo G *et al.* BRAF: A
982		Two-Faced Janus. *Cells* 2020; **9**: 2549.
983	84	Pavlick AC, Fecher L, Ascierto PA, Sullivan RJ. Frontline Therapy for BRAF-
984		Mutated Metastatic Melanoma: How Do You Choose, and Is There One Correct
985		Answer? *American Society of Clinical Oncology Educational Book* 2019; : 564–
986		571.
987	85	Anguera G, Majem M. BRAF inhibitors in metastatic non-small cell lung cancer.
988		*J Thorac Dis* 2018; **10**: 589–592.
989	86	Haugen BR, Alexander EK, Bible KC, Doherty GM, Mandel SJ, Nikiforov YE *et*
990		*al.* 2015 American Thyroid Association Management Guidelines for Adult
991		Patients with Thyroid Nodules and Differentiated Thyroid Cancer: The American
992		Thyroid Association Guidelines Task Force on Thyroid Nodules and
993		Differentiated Thyroid Cancer. *Thyroid* 2016; **26**: 1–133.
994	87	Schmidbauer B, Menhart K, Hellwig D, Grosse J. Differentiated Thyroid
995		Cancer—Treatment: State of the Art. *Int J Mol Sci* 2017; **18**: 1292.
996	88	Kim TY, Kim WG, Kim WB, Shong YK. Current Status and Future Perspectives
997		in Differentiated Thyroid Cancer. *Endocrinology and Metabolism* 2014; **29**: 217.
998	89	Ahn B-C. Personalized Medicine Based on Theranostic Radioiodine Molecular
999		Imaging for Differentiated Thyroid Cancer. *Biomed Res Int* 2016; **2016**: 1–9.
1000	90	Paeng JC, Kang KW, Park DJ, Oh SW, Chung J-K. Alternative Medical
1001		Treatment for Radioiodine-Refractory Thyroid Cancers. *Nucl Med Mol Imaging*
1002		2011; **45**: 241–247.
1003	91	Brose MS, Nutting CM, Jarzab B, Elisei R, Siena S, Bastholt L *et al.* Sorafenib in
1004		radioactive iodine-refractory, locally advanced or metastatic differentiated

thyroid cancer: a randomised, double-blind, phase 3 trial. *The Lancet* 2014; **384**:
319–328.

92 Schlumberger M, Tahara M, Wirth LJ, Robinson B, Brose MS, Elisei R *et al.*
Lenvatinib versus Placebo in Radioiodine-Refractory Thyroid Cancer. *New*
England Journal of Medicine 2015; **372**: 621–630.

93 Ho AL, Grewal RK, Leboeuf R, Sherman EJ, Pfister DG, Deandreis D *et al.*
Selumetinib-Enhanced Radioiodine Uptake in Advanced Thyroid Cancer. *New*
England Journal of Medicine 2013; **368**: 623–632.

94 Rothenberg SM, Daniels GH, Wirth LJ. Redifferentiation of Iodine-Refractory
BRAF V600E-Mutant Metastatic Papillary Thyroid Cancer with Dabrafenib—
Response. *Clinical Cancer Research* 2015; **21**: 5640–5641.

95 Buffet C, Wassermann J, Hecht F, Leenhardt L, Dupuy C, Groussin L *et al.*
Redifferentiation of radioiodine-refractory thyroid cancers. *Endocr Relat Cancer*
2020; **27**: R113–R132.

96 Riesco-Eizaguirre G, Santisteban P. A perspective view of sodium iodide
symporter research and its clinical implications. *Eur J Endocrinol* 2006; **155**:
495–512.

97 Czarnecka AM, Bartnik E, Fiedorowicz M, Rutkowski P. Targeted Therapy in
Melanoma and Mechanisms of Resistance. *Int J Mol Sci* 2020; **21**: 4576.

98 Flaherty KT, Infante JR, Daud A, Gonzalez R, Kefford RF, Sosman J *et al.*
Combined BRAF and MEK Inhibition in Melanoma with BRAF V600
Mutations. *New England Journal of Medicine* 2012; **367**: 1694–1703.

99 Larkin J, Ascierto PA, Dréno B, Atkinson V, Liszkay G, Maio M *et al.*
Combined Vemurafenib and Cobimetinib in BRAF-Mutated Melanoma. *New*
England Journal of Medicine 2014; **371**: 1867–1876.

100 Dummer R, Ascierto PA, Gogas HJ, Arance A, Mandala M, Liszkay G *et al.*
Encorafenib plus binimetinib versus vemurafenib or encorafenib in patients with
BRAF -mutant melanoma (COLUMBUS): a multicentre, open-label, randomised
phase 3 trial. *Lancet Oncol* 2018; **19**: 603–615.

101 Smalley KSM. Increased immunity and BRAF inhibition: Yet another argument
for combination therapy? *Pharmacol Res* 2016; **113**: 719–720.

102 Roskoski R. Targeting oncogenic Raf protein-serine/threonine kinases in human
cancers. *Pharmacol Res* 2018; **135**: 239–258.

103 Zhao X, Wang JR, Dadu R, Busaidy NL, Xu L, Learned KO *et al.* Surgery After
BRAF-Directed Therapy Is Associated with Improved Survival in BRAF [V600E]
Mutant Anaplastic Thyroid Cancer: A Single-Center Retrospective Cohort Study.
Thyroid 2023; **33**: 484–491.

104 Haenebalcke L, Goossens S, Naessens M, Kruse N, Ghahremani MF, Bartunkova
S *et al.* Efficient ROSA26-Based Conditional and/or Inducible Transgenesis
Using RMCE-Compatible F1 Hybrid Mouse Embryonic Stem Cells. *Stem Cell*
Rev Rep 2013; **9**: 774–785.

105 Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics* 2014; **30**. doi:10.1093/bioinformatics/btu170.

106 Kim D, Langmead B, Salzberg SL. HISAT: a fast spliced aligner with low
memory requirements. *Nat Methods* 2015; **12**. doi:10.1038/nmeth.3317.

107 Hao Y, Hao S, Andersen-Nissen E, Mauck WM, Zheng S, Butler A *et al.*
Integrated analysis of multimodal single-cell data. *Cell* 2021; **184**: 3573-
3587.e29.

108 Love MI, Huber W, Anders S. Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biol* 2014; **15**: 550.

1055

1056

1057

1058

1059 **FIGURE LEGENDS**

1060

Fig. 1 mESC_Nkx2-1/Pax8_bTg_Braf^{V637E} cell line differentiation into thyroid

organoids. Scheme of the thyroid differentiation protocol and Braf^{V637E} induction in

mature thyroid follicles (A). Differentiation of mESC_Nkx2-1/Pax8_bTg_Braf^{V637E} cell

line promotes expression of the main thyroid genes and Braf^{V637E} exogenous (B). Control

corresponds to the - Dox condition. Immunofluorescence staining showing Nkx2-1 and

Tg co-expressing cells organized in follicular structures (C), which are accumulating the

thyroid hormone precursor, Tg-I, inside the lumen compartment (D). The follicular

enrichment (FE) procedure significantly increased the expression levels of thyroid genes

(E) while keeping the structural organization of the follicles evidenced by Nis basolateral

localization (F) and its functionality, with Tg-I accumulation (G). Values represent the

median (IQR) of 3 independent experiments with individual values shown (*$p<0.05$;

**$p<0.01$; ***$p<0.001$; Mann-Whitney U test). Scale bars, 50 μm and 10 μm for high

magnification follicles.

1074

Fig. 2 Effect of the Braf^{V637E} oncogene induction on mature thyroid follicles.

Schematic representation of thyroid differentiation, Braf^{V637E} oncogene expression on

thyroid cells, and its activation under tamoxifen (4OHT) treatment (A). Western blot

demonstrates an increase of phospho-ERK (pERK) 48 hours after the addition of 4OHT

to the organoids culture (B). Beta-actin was used as a loading control for the immunoblot

experiments. The image represents one experiment from 3 experimental replicates. Gene

expression analysis showing the inhibitory effect of Braf^{V637E}-oncogene activation on

thyroid genes after 6 hours (C), 48 hours (D), 7 days (E) and 21 days (F) of 4OHT

treatment. For each time point gene expression levels of the cAMP+4OHT treated cells

were compared to the control (cAMP) levels. Bar graphs represent the median (IQR) of

at least 3 independent experiments with individual values shown. ($p<0.05$; $p<0.01$;

***p<0.001; ***p<0.0001; Mann-Whitney U test). Proportions of proliferating

(Nkx2.1/BrDU+) (G) and apoptotic (Nkx2.1/Caspase3+) (H) cells among control

(cAMP) and Braf^{V637E}-induced (4OHT) organoids after 7 and 21 days of 4OHT treatment.

For the proliferation assay, isotype and -BrdU conditions were used as negative controls

for flow cytometry gating. As a positive control for apoptosis, staurosporine treatment

(24 h) was used to determine Caspase 3 expression. Bar graphs represent the median

(IQR) of at least 3 independent experiments with individual values shown. cAMP was

used as control for comparisons. (**p<0.01; Mann-Whitney U test).

1094

Fig. 3 Morphological changes on thyroid follicles caused by Braf^{V637E} activation.

cAMP-treated thyroid cells show follicular organization with Nkx2-1 nuclear expression

and Tg accumulation in the luminal compartment. In contrast, after 48 h of Braf^{V637E}

oncogene induction by 4OHT, most of the cells are not organized into follicular structures

and a great proportion is expressing very low levels of Tg (A). Higher magnification

images showing the follicular organization of the thyroid cells in the control condition

(48 h), with proper expression of Nkx2-1 and/or Tg (B), E-cadherin (C), Zo-1 (D), Nis

(E) and Tg-I (F) accumulation in the lumen. While in the 4OHT condition the follicular

organization is disrupted as well as its function. Nkx2-1 and Tg co-staining in control

(cAMP) and Braf^{V637E}-induced (4OHT) cells for 7 days (G) and 21 days (H) shows clear

changes in thyroid morphology evidenced by the heterogeneity of Nkx2-1 cells which

mostly do not express Tg or at low levels at day 7, while at day 21 a higher proportion of

Nkx2-1 cells are Tg positive. Hoescht (shown in blue) was used for nuclei staining. Scale

bars, 20 μm (A), 10 μm (B-F) and 50 μm (G-H).

1109

Fig. 4 Drug screening reveals that VPA and MAPK/PI3K inhibition can restore Nis

expression in Braf^{V637E}-induced cells. Schematic representation of the protocol used for

drug screening experiments (A). qPCR data show that 4OHT-treated organoids (4 days)

treated with MEK (PD0325901; PD) and PI3K (LY294002; LY) inhibitors isolated

increase but do not restore *Nis* expression to control (cAMP) levels. However, when

combined, *Nis* expression reached cAMP levels. On the other hand, the HDAC inhibitor

(VPA) can recover *Nis* expression (B) by itself. Bar graphs represent the median (IQR)

of at least 4 independent experiments with individual values shown. Comparisons were

performed against the cAMP condition. (**$p < 0.05$; ***$p < 0.001$; Mann-Whitney U test).

Confocal images show downregulation of the Nis transporter in 4OHT condition, which

is restored by VPA treatment. However, the expression pattern differs from the control

(cAMP) condition. Co-inhibition of MAPK and PI3K pathways associated or not to VPA

treatment restores Nis protein expression at the basolateral membrane of the properly

organized follicles (C). Scale bars, 10 μm. Gene expression analysis shows that

PD0325901 and LY294002 co-treatment recover the *Tg* (D), *TSHR* (E), and *Tpo* (F)

mRNA to cAMP levels. Bar graphs represent the median (IQR) of at least 3 independent

experiments with individual values shown. Comparisons were performed against the

cAMP condition. (*$p < 0.05$; **$p < 0.01$; ***$p < 0.001$; ***$p < 0.0001$; Mann-Whitney U

test). Immunostaining for Nkx2-1 and Tg shows that proteins levels are similar to controls

(cAMP) while follicular organization (G, H) and luminal Tg-I accumulation were

restored under PD+LY and PD+LY+VPA conditions (I). Scale bars, 20 μm (G), 50 μm

(H), and 10 μm (I). Organification assay shows recovery of Iodine uptake (J), Protein-

bound to 125I (K), and % of iodine Organification (L) in PD+LY and PD+LY+VPA

conditions. Bar graphs represent the median (IQR) of at least 3 independent experiments

with individual values shown. Comparisons were performed against the cAMP condition.

(**p<0.01 ; ***p<0.0001; Mann-Whitney U test).

1136

Fig. 5 Dabrafenib and trametinib effect on redifferentiation of Braf^V637E^-expressing

cells. Braf^V637E^-expressing organoids treated with Dabrafenib and Trametinib restore the

expression of *Nis* to cAMP levels. However, a greater increase is observed under

Trametinib+LY co-treatment (A). Recovery of *Tg* (B)*, TSHR* (C) and *Tpo* (D) mRNA to

control levels was observed under co-treatment of Dabrafenib and Trametinib with the

PI3K inhibitor (LY). Bar graphs represent the median (IQR) of at least 4 independent

experiments with individual values shown. Comparisons were performed against the

cAMP condition. (*p<0.05 ; **p<0.01; Mann-Whitney U test). Confocal images show

that Dabrafenib and Trametinib isolated or combined to PI3K inhibitor (LY) also induce

Tg and Nis protein levels while restoring the follicular structure (in a proportion of cells)

and Tg-I accumulation in the lumen (E). Hoescht (shown in blue) was used for nuclei

staining. Scale bars, 20 μm (Tg and Tg-I) and 10 μm (Nis). Proportions of proliferating

(Nkx2.1/BrDU+) (F) and apoptotic (Nkx2.1/Caspase3+) (G) cells among control

(cAMP), Braf^V637E^-induced (4OHT) and inhibitors (4OHT+inhibitors)-treated organoids

after. For the proliferation assay, isotype and -BrdU conditions were used as negative

controls for flow cytometry gating. As a positive control for apoptosis, staurosporine

treatment (24 h) was used to determine Caspase 3 expression. Bar graphs represent the

median (IQR) of at least 3 independent experiments with individual values shown.

Comparisons were performed against the 4OHT condition. (*p<0.05 ; **p<0.01 ; Mann-

Whitney U test). Western blot shows an increase of phospho-ERK (pERK) 7 days after

the addition of 4OHT to the organoids culture when compared to cAMP control.

Conversely, the treatment with Trametinib+LY resulted in pERK reduction compared to

the 4OHT condition. (B). Beta-actin was used as a loading control for the immunoblot

experiments. The image represents one experiment from 3 experimental replicates.

1161

Fig. 6 Transcriptomics analysis confirms thyroid redifferentiation of Braf^{V637E}-

expressing cells treated with MAPK and PI3K inhibitors and suggests by which

mechanisms. Heatmap of normalized bulk RNA-Seq expression of normal thyroid cells

(cAMP), Braf^{V637E} expressing cells (4OHT), and Braf^{V637E} expressing cells treated with

PD0325901+LY204002 inhibitors. Rows represent markers and columns represent

specific conditions. Color values in the heatmap represent mean expression levels (A).

Thyroid differentiation (eTDS and TDS) (B) and ERK (C) scores were calculated among

the different conditions. Classification of upregulated and downregulated genes

comparing 4OHT *vs.* cAMP (D, E) and PD+LY *vs.* 4OHT (F, G). Colors represent the

classification method and scale the -log10 (adj p-value).