Electrophysiological indices of hierarchical speech processing differentially reflect the comprehension of speech in noise

Abbreviated title: EEG speech indices differentially reflect comprehension

Shyanthony R. Synigal[1,2], Andrew J. Anderson[1,2,3], Edmund C. Lalor[1,2,4,5]

[1]Department of Neuroscience, University of Rochester, Rochester, NY 14642, USA.

[2]Del Monte Institute for Neuroscience, University of Rochester, Rochester, NY 14642, USA.

[3]Department of Neurology, Medical College of Wisconsin, Milwaukee, WI 53226, USA.

[4]Department of Biomedical Engineering, University of Rochester, Rochester, NY 14627, USA.

[5]Center for Visual Science, University of Rochester, Rochester, NY 14627, USA.

Author Contributions: ECL and SRS designed research; SRS performed research; AJA contributed analytic tools, SRS analyzed data; SRS, AJA, ECL wrote the paper

Corresponding Author: elalor@ur.rochester.edu (ECL)

Number of figures: 6 in main text, 1 supplementary
Number of tables: 2 in main text, 3 supplementary
Number of multimedia: 0
Number of words for abstract: 244
Number of words for significance statement: 116
Number of words for introduction: 678
Number of words for discussion: 1,955

Acknowledgements: The authors thank Ms. Xueying Wang and Dr. Aaron Nidiffer for some assistance with data collection, and Dr. Aaron Nidiffer and Dr. Madeline Cappelloni for helpful comments on the manuscript.

Authors report no conflict of interest.

Funding sources: University of Rochester Department of Biomedical Engineering, University of Rochester Del Monte Institute for Neuroscience, The Del Monte Institute Pilot Grant OP346161

ABSTRACT

The past few years have seen an increase in the use of encoding models to explain neural responses to natural speech. The goal of these models is to characterize how the human brain converts acoustic energy into distinct linguistic representations that enable everyday speech comprehension. For example, researchers have shown that electroencephalography (EEG) data can be modeled in terms of acoustic features of speech, such as its amplitude envelope or spectrogram, linguistic features such as phonemes and phoneme probability, and higher-level linguistic features like context-based word predictability. However, it is unclear how reliably EEG indices of these speech feature representations reflect comprehension in different listening conditions. To address this, we recorded EEG from neurotypical adults who listened to segments of an audiobook in various levels of background noise. We modeled how their EEG responses reflected a range of acoustic and linguistic speech features and how this tracking varied with behavior across noise levels. EEG tracking of nearly all examined features showed SNR-dependent changes in unique variance explained, with the largest changes occurring for linguistic features. We hypothesized that only higher-level feature tracking would predict behavior but instead found that both high and low-level features were associated with behavioral scores depending on the noise level. EEG markers of the influence of top-down, context-based prediction on bottom-up acoustic processing also correlated with behavior. These findings help characterize the relationship between brain and behavior by comprehensively linking hierarchical indices of neural speech processing to language comprehension metrics.

SIGNIFICANCE STATEMENT

Acoustic and linguistic features of speech have been shown to be consistently tracked by neural activity even in noisy conditions. However, it is unclear how signatures of low- and high-level features covary with one another and relate to behavior across these listening conditions. Here, we find that linguistic (phonetic feature and word probability-based feature) processing is affected by noise more than low-level acoustic feature processing. We also find that behavioral performance is associated with acoustic, phonetic, and lexical surprisal tracking, and that these associations depend on background noise levels. These results extend our understanding of how various speech features are comparatively reflected in electrical brain activity and how they relate to perception in challenging listening conditions.

INTRODUCTION

Given the importance of speech communication in human life, tremendous amounts of research have focused on characterizing the neurophysiology of language comprehension (Hickok, 2015). This research has revealed a network of brain areas that are functionally specialized for processing different hierarchical levels of speech and language (Hickok & Poeppel, 2007). For example, work has shown that low level acoustic and spectrotemporal features of speech are chiefly processed in early auditory cortex (de Heer et al., 2017), with various phonological features being processed in secondary areas such as the superior temporal gyrus (Hamilton et al., 2018; Mesgarani et al., 2014) and some prefrontal areas (Burton, 2009; de Heer et al., 2017), and meaning being represented across large areas of cortex (Anderson et al., 2017; Huth et al., 2016; Pereira et al., 2018).

While much of the above knowledge has been obtained from functional neuroimaging and invasive recordings in neurosurgical patients, parallel efforts have been made to obtain noninvasive magneto- and electrophysiological (MEG/EEG) markers reflecting the hierarchical processing of speech. This includes modeling how EEG and MEG track the amplitude envelope of natural speech (Lalor & Foxe, 2010) and how neural responses reflect the spectrotemporal (Daube et al., 2019; Di Liberto et al., 2015), phonetic (Di Liberto et al., 2015), phonotactic (Brodbeck, Hong, et al., 2018; Di Liberto et al., 2019; Gwilliams et al., 2022), lexical (Dou et al., 2025; Heilbron et al., 2022), prosodic (Teoh et al., 2019), and semantic (Broderick et al., 2018; Heilbron et al., 2022) features of natural speech.

Some advantages of EEG are that it is significantly cheaper and easier to use in applied research in different cohorts (Peck et al., 2021; Salisbury et al., 2002). Consequently, there has been considerable interest in exploring how different EEG markers of speech processing reflect speech intelligibility (Verschueren et al., 2021) and language comprehension (Ahissar et al., 2001;

Broderick et al., 2022). Many of these studies altered the intelligibility or comprehensibility of

speech by adding background noise (Iotzov & Parra, 2019; Schmitt et al., 2022; Zhang et al., 2023)

or by degrading the speech signal itself (Karunathilake et al., 2023; Viswanathan et al., 2021). For

example, some work has shown that cortical tracking of the speech envelope decreases as noise

levels increase (Etard & Reichenbach, 2019; Lesenfants et al., 2019; Vanthornhout et al., 2018;

Zou et al., 2019), although others have suggested such tracking remains robust until the

background noise is more than twice as loud as the speech it masks (Ding & Simon, 2013).

Meanwhile, experiments that have explored EEG indices of semantic processing appear to show a

strong correlation with speech intelligibility and/or understanding (Broderick et al., 2018). Very

few studies, however, have systematically explored how EEG indices of both low- and high-level

speech processing covary across different levels of speech comprehension (Mesik et al., 2021;

Sohoglu & Davis, 2020; Strauß et al., 2022; Yasmin et al., 2023). This is the goal of the present

study.

Our study explores how EEG markers of speech envelope, spectrogram, acoustic onset,

phonetic feature, and lexical surprisal vary with subjective measures of the percentage of words

heard (PWH, as a proxy for intelligibility) and objective measures of comprehension across

different background noise conditions. We hypothesize that EEG measures of higher-level

processing (e.g., lexical surprisal and phonetic features) will more strongly correlate with behavior

than lower-level measures (e.g., spectrogram). We also test how listeners exploit linguistic context

to process noisy speech and how that effect might manifest in EEG. Here, we leverage a previously

introduced measure of predictive speech perception that quantifies how the tracking of low-level

speech features varies as a function of the context-based semantic content of that speech (Broderick

et al., 2019), but this time using lexical surprisal. We hypothesize that this measure strengthens for

speech in moderate levels of noise (when speech is still intelligible) relative to speech in quiet,

before declining at high levels of background noise (when speech is no longer intelligible). With

this study we seek to extend our understanding of the hierarchical processing of continuous speech

under a range of realistic listening conditions.

MATERIALS AND METHODS

Participants

28 healthy adults (9 males, 18-35 years old [mean = 22.08, SD = 4.63]) participated in this

study. One participant was excluded due to an insufficient amount of data and two were excluded

due to technical issues, resulting in a dataset of 25 participants. Each participant provided written

informed consent and reported having normal hearing, normal or corrected-to-normal vision,

English as their first and main language, and no history of neurological disorders. Participants were

also compensated for their participation. All procedures were approved by the University of

Rochester Research Subjects Review Board.

Stimuli and experimental procedure

Participants listened to 70 minutes of *A Wrinkle in Time* by Madeleine L'Engle which was

read by an American female speaker. The stimulus was presented across 70 trials, each being one

minute in duration. Each trial was presented at one of five noise levels: quiet (no noise) and +3

dB, -3 dB, -6 dB, and -9 dB signal-to-noise ratios (SNRs). The background noise was spectrally

matched stationary noise, which was estimated from the clean speech using a 46[th] order forward

linear predictive model (Crosse, Di Liberto, & Lalor, 2016; Ding & Simon, 2013). Prediction order

was defined based on the audio sampling rate as $p_{order} = (Fs/1000)+2$, to capture vocal tract and

non-vocal tract related contributions to speech signals (Atal & Hanauer, 1971). There were 14

minutes' worth of audio for each of the five noise conditions. The storyline was preserved from

trial to trial, but the conditions were pseudorandomized such that no noise level occurred

consecutively. Participants estimated the percentage of words they heard (on a scale of 0-100%)

and answered two multiple choice comprehension questions after each trial. The comprehension

questions used here were the same questions used from a previous study (Maddox & Lee, 2018),

except we presented only two out of the four original questions created for each trial. The stimuli

were presented through Sennheiser HD650 headphones at a sampling rate of 44.1 kHz using

Psychtoolbox (Kleiner et al., 2007) and custom MATLAB scripts (MathWorks, 2019).

Data acquisition and preprocessing

EEG data were recorded from 128 scalp electrodes (plus two mastoid channels that were

not analyzed in this work). The data were acquired at a 1024 Hz sampling rate with the BioSemi

Active Two system. The data were preprocessed using the PREP pipeline and its default

parameters (Bigdely-Shamlo et al., 2015). This pipeline first detrended the data at 1 Hz, then

removed the 60 Hz line noise. Afterwards, robust re-referencing was applied which allows the data

to be referenced to an average of all channels except those contaminated with noise. This function

identifies and interpolates noisy channels in an iterative manner such that the re-referencing itself

is not affected by the noise. The cleaned data was then low pass filtered at 8 Hz, using a Chebyshev

Type II filter with an 8.5 Hz cutoff frequency and an 80 dB stopband attenuation. Next, the data

were epoched and independent component analysis (ICA) was applied using EEGLAB's *picard*

function (Delorme & Makeig, 2004; Pion-Tonachini et al., 2019) to remove muscle and eye

artifacts. Lastly, the data were downsampled to 128 Hz.

Speech stimulus characterization

Speech is organized in a hierarchical manner where sounds can form syllables, syllables

form words, words form sentences, and so on. To assess how our brains might concurrently process

speech across levels of this hierarchy, we chose to model EEG responses to speech based on

several different representations of that speech, all of which were computed using the clean (no

noise) versions of each trial.

Envelope. We first calculated the speech envelope, a well-established feature shown to be

robustly tracked by cortical activity (Aiken & Picton, 2008; Destoky et al., 2019; Di Liberto et al.,

2015; Ding & Simon, 2013; Etard & Reichenbach, 2019; Lalor & Foxe, 2010; Nourski et al., 2009;

Pasley et al., 2012) and to be important for speech recognition and intelligibility (Ahissar et al.,

2001; Drullman et al., 1994; Shannon et al., 1995). The speech signal was first lowpass filtered at

20 kHz (22.05 kHz cutoff frequency, 1 dB passband attenuation, 60 dB stopband attenuation). The

broadband speech envelope was calculated using a gammachirp auditory filterbank to mimic the

filtering properties of the cochlea (Irino & Patterson, 2006). This filterbank was used to filter the

speech into 16 bands from 250 Hz to 8 kHz with an equal loudness contour (essentially creating a

spectrogram). Lastly, we calculated the absolute value of the Hilbert transform for each frequency

band and averaged this result across all 16 bands.

Acoustic onsets and spectrogram. We chose to model two additional acoustic features,

acoustic onsets and spectrogram, which were shown to be reflected in cortical activity above and

beyond the speech envelope (Brodbeck et al., 2020; Di Liberto et al., 2015; Sohoglu & Davis,

2020). Acoustic onsets were approximated by computing the first derivative of the speech

envelope and then half-wave rectifying the result (Hertrich et al., 2012). A 16-band spectrogram

was computed using the same filterbank and parameters as the speech envelope, just without the

final averaging step.

Phonetic features. To calculate phonetic features, the Montreal Forced Aligner (McAuliffe

et al., 2017) was first used to partition and time align each word in the story into phonemes

according to the International Phonetic Alphabet for American English. Then, each phoneme was

linearly mapped onto a set of 19 binary phonetic features based on the University of Iowa's

phonetics project (http://www.uiowa.edu/~acadtech/phonetics/english/english.html/).

Lexical surprisal. Lastly, we calculated the surprisal of each word based on its preceding

context using the Transformer-XL model (Dai et al., 2019). This model contains a recurrence

mechanism that allows it to build and reuse memory from previous segments and learn longer-

term dependencies, while preserving the temporal information of previous word embeddings. This

model was chosen because it can predict the probability of an upcoming word using the context

from all preceding words. The softmax of the values from the output layer of the model were taken

to estimate the probability of each word, and the negative log of a word's probability was computed

to estimate its lexical surprisal (Dai et al., 2019).

Modeling the relationship between speech features and EEG responses

One goal of the present study was to determine how neural representations of individual

speech features change with SNR and how those changes relate to comprehension and what

participants reported hearing. To index the fidelity of individual speech feature representations,

we used a forward encoding model (in the form of a linear filter or kernel, sometimes known as a

temporal response function) which describes the transformation from the speech features to the

EEG responses recorded at each electrode. Here, acoustic onsets, the spectrogram, phonetic

features, and lexical surprisal were modeled. An additional vector with impulses placed at each

word's onset was included to capture any acoustic related onset responses at word boundaries that

the acoustic onset predictor may have missed. The forward encoding models can be expressed as

follows:

$$r(t,n) = \sum_{\tau} w(\tau, n) \, s(t - \tau) + \varepsilon(t, n) \qquad (1)$$

where $s(t - \tau)$ is the speech feature of interest, $r(t, n)$ is the neural response on each channel, n,

$w(\tau, n)$ is the temporal response function to be estimated, and $\varepsilon(t, n)$ is the residual recorded EEG

not explained by the model, which is assumed to be Gaussian.

Because our five different speech features of interest (spectrogram, acoustic onsets,

phonetic features, lexical surprisal, and word onsets)—like almost all speech features—are

correlated with each other in time, we wanted to use an approach where we could identify the

unique contribution that each speech feature made to the resulting EEG responses. To do this, we

used a partial correlation approach. Specifically, our strategy involved first modeling the data from

each of our experimental (SNR) conditions *separately* using each of our five speech features. This

resulted in five TRF models for each condition, calculated from training data. We then used the

five speech features from left-out trials, and the corresponding TRF models, to generate five

separate predictions of the EEG responses for those trials.

Then, to calculate how much each speech feature uniquely contributed to the recorded brain

activity, we computed a partial correlation between the real EEG and the predicted EEG from each

speech feature, while accounting for the predicted EEG from all other speech features. For

example, to assess how much the acoustic onsets [ons_A] uniquely contributed to the EEG

responses, we concatenated the predicted EEG from the other four feature models (spectrogram

[sgram], phonetic features [fea], surprisal [surp], word onsets [ons_W]) into a matrix, Z. We then

used MATLAB's partialcorr (X,Y,Z) function (Fisher, 1924) to determine the partial (Pearson's)

correlation between the actual EEG (X) and the predicted EEG from the model trained on the

acoustic onsets (Y), while controlling for the activity that can be explained by all other features

(Z). We repeated this process to identify the unique contribution of each speech feature, while

controlling for all other speech features.

In terms of training and testing the models, nested leave-one-out cross-validation and ridge

regression were used. Each speech feature was normalized between 0 and 1 by dividing by its

maximum value, and the EEG data were z-scored. The features and EEG were partitioned into

train and test sets. The stimuli were lagged from -100-700ms to capture both short and long latency

responses to acoustic and linguistic features. Given the amount of data, models were validated on

the clean condition and tested on the remaining noise conditions. Specifically, cross-validation

was conducted on the *clean* condition to select the optimal regularization (ridge) parameter, λ,

which ranged from 10^{-1}–10^{8}. We identified the regularization parameter that resulted in the highest

prediction accuracy for each individual test fold. We then selected the parameter that produced the

highest prediction accuracy most often (across all test folds) so that we could use one parameter

to train the models for each condition. Using the same parameter for all folds and conditions

(within a participant) allowed for a fairer comparison of model performance since it results in the

predictions across trials being on the same scale for each participant. This method was particularly

important for the low SNR conditions, as the models would be more prone to overfitting, especially

when training on multidimensional feature spaces.

Our primary analysis aimed to assess how different hierarchical speech features related to

EEG activity across listening conditions. However, given its prevalence in the literature and

relative ease of modeling, we also examined how speech envelope tracking varied with SNR. To

do this, we used backward modeling to reconstruct an estimate of the speech envelope from the

recorded EEG for each trial. The backward model or decoder, $g(\tau, n)$, described the

transformation from lagged EEG responses at all electrodes, $r(t + \tau, n)$, to an estimate of the

speech envelope, $\hat{s}(t)$. As detailed elsewhere (Crosse, Di Liberto, Bednar, et al., 2016; Crosse et

al., 2021), the modeling procedure can be expressed as:

$$\hat{s}(t) = \sum_{n}\sum_{\tau} r(t + \tau, n)g(\tau, n)$$ (2)

Similar to the forward models, this analysis was conducted separately for each

experimental (SNR) condition using 14-fold leave-one-out cross-validation. First, the stimuli and

responses were normalized and then partitioned into train and test sets. Here the EEG data were

lagged, τ, from -100–300ms to focus on short latency responses to the speech envelope, s(t).

Cross-validation and regularization parameter (10^{-1}–10^8) selection followed the same procedure as

described above for forward modeling. A decoder was trained using the training set and selected

regularization parameter to reconstruct an estimate of the speech envelope. After testing the

decoder on held-out data, model performance was assessed by computing the Pearson correlation

between the actual speech envelope and the reconstructed speech envelope. This model

performance, also known as reconstruction accuracy, was averaged across folds and participants

(Figure 1).

Figure 1. Methods. EEG data were recorded while participants listened to an audiobook in different levels

of noise. They also reported the percentage of words they heard (%) and answered comprehension questions

(Q&A). Forward modeling (with a temporal response function, TRF) was used to estimate EEG responses

(R) from the clean representations of the speech features (S), which were acoustic onsets (ons$_A$), word

onsets (ons$_W$), spectrogram (sgram), phonetic features (fea), and word surprisal (surp). Model performance

(r) was assessed by calculating the partial correlation between the actual EEG and predicted EEG from one

feature's model, while controlling for the predictions generated by all other features. These partial

correlations were averaged across the selected electrode channels marked in the head plots. Backward

modeling (with a decoder, g) was also used to reconstruct an estimate of the clean speech envelope (env,

S). Model performance (reconstruction accuracy, ρ_w) was assessed by calculating the correlation between

the original speech envelope and the reconstructed speech envelope. A linear mixed-effects model (LME)

was then used to determine the influence of word surprisal on envelope reconstruction accuracy.

Assessing the role of lexical context on acoustic encoding in different levels of background

noise

Another major goal of the present study was to test how listeners might rely more on lexical

context to—perhaps predictively—encode speech that is masked by moderate levels of

background noise. To do this, we used a variant of a recently introduced approach that involved

quantifying how the tracking of low-level speech features varies as a function of the context-based

semantic content of that speech (Broderick et al., 2019). In our variation, we used a linear mixed-

effects (LME) model to explore the extent to which word predictability in the form of lexical

surprisal as opposed to semantic similarity influenced how the envelope of that word was reflected

in EEG and how that influence changed across SNRs. By using an LME model in this way, one

can measure the relationship between the main variable(s) of interest while controlling for

variability caused by random factors. We used the *lmerTest* package in R (Kuznetsova et al., 2017)

to model the following equation:

$$\rho_w \sim 1 + snr + surp + envStd + f_{rel} + res + f_{rel} * envStd +$$
$$f_{rel} * res + surp * envStd + surp * res + surp * snr + \quad\quad\quad (3)$$
$$(1|participant) + (1|word)$$

The dependent variable is word reconstruction accuracy which was calculated as the

Spearman's correlation between the actual word envelope and the predicted word envelope for the

first 100 ms of each word. As detailed below, the independent variables are SNR, lexical surprisal

(*surp*), envelope variability (*envStd*), relative pitch (f_{rel}), resolvability (*res*), and various

interactions such as the interaction between surprisal and SNR.

Envelope variability, relative pitch, and resolvability were selected as nuisance regressors.

This was because these measures can correlate with surprisal, with one another, and with envelope

tracking; so, following Broderick and colleagues, they are included here and as interactions to

ensure that they aren't inherently contaminating the lexical surprisal effects (Broderick et al.,

2019). Envelope variability is represented as the standard deviation of the speech envelope.

Relative pitch is pitch normalized to the vocal range of the speaker (Tang et al., 2017) and

resolvability measures whether a sound's harmonics are processed between distinct (resolved) or

within the same (unresolved) filters of the cochlea (Shackleton & Carlyon, 1994). Prosodic cues

such as relative pitch and resolvability can uniquely predict EEG activity even after accounting for

other acoustic and phonetic features (Teoh et al., 2019) which was yet another reason to include

them in this LME model. Relative pitch was extracted using Praat (Boersma & Weenink, 2013).

Once the software estimated absolute pitch, this result was then z-scored, resulting in relative pitch.

Resolvability was extracted using custom scripts based on a model of the human auditory

periphery (McDermott & Simoncelli, 2011; Teoh et al., 2019).

The LME model also included by-word and by-participant random intercepts as some

words may be easier to reconstruct than others and some participants may, on average, have higher

reconstruction accuracies than other participants (based on cortical folding and other biophysical

factors). No random slopes were included, as they caused the model to not converge (calculate a

solution) even with the addition of an optimizer (additional functions the model can use to reach a

solution (Brown, 2021)). Like Broderick and colleagues, the speech envelope and the nuisance

regressors were calculated for the duration of the trials. Afterward, word reconstruction accuracy

was calculated over the first 100 ms following each word's onset and the nuisance regressors were

segmented over the same interval (Broderick et al., 2019). There were 11,748 words total with an

average length of 292.3 ms and a standard deviation of 182.3 ms.

Statistical analysis

All statistical analyses were performed in R (version 4.5.0) and in MATLAB R2024b

(MathWorks, 2024). Due to the skewed distribution of the behavioral scores, comparisons were

calculated using a nonparametric Friedman test, followed by a Wilcoxon Rank Sum test with Holm

correction. Pairwise tests were also used to determine differences in envelope reconstruction

accuracy and EEG partial correlations between conditions. Permutation testing was performed to

test if the partial correlation coefficients were greater than chance. The predicted EEG trials were

shuffled, and the partial correlation was computed between the actual EEG (in its correct order)

and the predicted EEG (which was out of order). This was repeated 1,000 times for each

participant. The null partial correlations were averaged across folds, electrodes (12 parietal

electrodes for the surprisal model and 12 frontotemporal electrodes for all other feature models,

all selected *a priori*, Figure 1), and participants, and we assessed the proportion of permuted

coefficients that were greater than the actual group-averaged coefficients. Lexical surprisal is

known to be reflected on more parietal areas of the scalp and displays more N400-like responses,

and thus the surprisal partial correlations were averaged across the selected parietal electrodes

(Broderick et al., 2022). All other features were averaged across the frontotemporal electrodes as

those regions have been shown to reflect cortical auditory responses (Di Liberto et al., 2015).

Permutation testing was also performed to find if the speech envelope reconstruction

accuracies were greater than chance. A null model was created for each SNR by shuffling the

speech envelope between trials and calculating a new model with each shuffle. This procedure,

including regularization, was repeated 1,000 times. The null reconstruction accuracies were

averaged across folds and participants and then we assessed the proportion of reconstruction

accuracies that were greater than the actual reconstruction accuracies.

One question of interest was how the neural representation of each speech feature varied

with noise. To test this, the partial correlations for each trial were used in an LME model. This

model (Equation 4) included SNR, speech feature type, and the interaction between the two as

fixed effects, and a by-participant random intercept.

$$r \sim snr + feature + snr{:}feature + (1|participant) \qquad\qquad (4)$$

To compare the changes in partial correlations across SNRs between each feature's model

(essentially comparing the slopes of the SNR by feature interaction), we computed their estimated

marginal means and performed pairwise comparisons with Tukey Honestly Significant Difference

(HSD) correction using the *emmeans* package in R (Lenth et al., 2019).

Generalized linear mixed models (GLMM) were also used to model the relationships

between behavior and partial correlations/reconstruction accuracies (Equation 5) and the

surprisal*SNR coefficients from Equation 3 (Equation 6).

$$behavior \sim (r_{feature} + \cdots + r_{feature\,N}) * snr + (1|participant) \tag{5}$$

$$behavior \sim \beta_{surp:snr} + (1|participant) \tag{6}$$

Equations 5 and 6 utilized ordered beta distributions with a logistic link function (which

can support the skewed and bounded nature of our behavioral data (Smithson & Verkuilen, 2006))

and the template model builder (glmmTMB) function. The *glmmTMB* package allowed us to

model the ordered beta distribution with the addition of random effects (Brooks et al., 2017).

Each mixed-effects model in this study was calculated using z-scored predictors, the

default parameters which included fitting the models with restricted maximum likelihood (REML),

and Satterthwaite's method for the t-tests. Marginal and conditional R^2 values are provided via the

performance package (Lüdecke et al., 2021) to indicate the variance explained by fixed effects

alone (R^2_m) and by both fixed and random effects (R^2_c) (Nakagawa et al., 2017). The remaining

packages used to perform statistical analysis and plotting in R were completed using *ggeffects*

(Lüdecke, 2018), *ggplot2* (Wickham, 2011), and *patchwork* (Pedersen, 2019).

RESULTS

Speech perception decreases as SNR decreases

Behavioral scores were collected to show how participants' perception of the story changed

as listening conditions became more challenging. After each one-minute-long trial, participants

estimated the PWH on a scale from 0-100% and answered two multiple-choice comprehension

questions, each of which had four possible answers. As expected, there was a significant reduction

in the PWH ($X^2(4) = 98.894$, $p < 2.20 \times 10^{-16}$) and comprehension scores ($X^2(4) = 83.44$, $p < 2.20$

x 10⁻¹⁶, Friedman test followed by Wilcoxon rank sum test with Holm correction, Figure 2) with

the addition of noise. After the experiment, participants reported hearing very few words in the -9

dB SNR condition but attempted to use context clues from previous trials to answer the questions

in the -9 dB SNR trials. To control for this potential confound in our measure of comprehension,

we used a procedure similar to Orf and colleagues, where nine additional volunteers answered the

comprehension questions for each trial without listening to the story (Orf et al., 2022). Based on

the performance of those new participants, we determined a new empirical chance level of 28%

rather than 25%. All participants performed above chance in the quiet, +3 dB SNR, and -3 dB SNR

conditions. Comprehension scores were not above chance for five participants in the -6 dB SNR

condition (p ≥ 0.066) and 13 participants in the -9 dB SNR condition (p ≥ 0.066). Although

participants reported hearing fewer words in the +3 dB SNR condition compared to speech in

quiet, they performed similarly in terms of comprehension (p = 1.000, S1 Table).

Figure 2. Behavioral results. (A) Average percentage of words participants reported hearing in each

condition. (B) Average percentage of correctly answered comprehension questions for each condition. The

dashed gray line is the chance level at 28% which was calculated using a separate set of participants who

answered the comprehension questions without listening to the audiobook. For both plots, significance is

indicated by * if p < 0.05, ** if p < 0.01, and *** if p < 0.001 using pairwise Wilcoxon rank sum tests. The

black dots in each plot correspond to individual participants and the box and whiskers depict the median

across participants, in addition to the 75th and 25th percentiles.

Hierarchical speech feature encoding declines across SNRs

Many EEG/MEG speech studies have focused on modeling brain responses to isolated

speech features, perhaps most frequently the speech envelope. Recent work, however, is

increasingly using various methods to model acoustic and linguistic features simultaneously

(Brodbeck, Hong, et al., 2018; Brodbeck, Presacco, et al., 2018; de Heer et al., 2017; Gillis et al.,

2021; Heilbron et al., 2022; Verschueren et al., 2022) to find how the brain uniquely encodes a

feature of interest while accounting for others, as some features may be correlated and explain

similar neural activity (Daube et al., 2019). Since few studies have modeled the encoding of

simultaneous features in challenging listening conditions (Brodbeck et al., 2020; Karunathilake et

al., 2023; Sohoglu & Davis, 2020), we were interested in how a range of acoustic and linguistic

features were encoded in noise.

To test this, we quantified the unique contribution that each speech feature makes to the

EEG responses using partial correlation calculations. Specifically, separate models were fit on the

acoustic onset, speech spectrogram, phonetic feature, word surprisal, and word onset features.

Those models were tested on left out data to predict unseen EEG. The partial correlation was then

calculated between the actual EEG and the predicted EEG for one feature while controlling for the

effects of all others. The acoustic onset, spectrogram, phonetic feature, and word onset model

partial correlations were averaged across 12 frontotemporal electrodes (Di Liberto et al., 2015)

and the surprisal partial correlations were averaged across 8 parietal electrodes (Broderick et al.,

2022) plus 4 additional electrodes in the same area so that both analyses consisted of 12 electrodes

(electrodes shown in Figure 1). Model partial correlation coefficients were above chance for each

SNR ($p \leq 0.021$, permutation tests) except for the phonetic feature model in the -9 dB SNR ($p =$

0.157) and the word onset model in the -6 dB and -9 dB SNRs (p = 0.146, p = 0.333). Unlike the

other features, there were no significant differences in the average partial correlation coefficients

for the spectrogram model between SNRs (Figure 3A, within feature paired tests with Tukey HSD

correction).

Figure 3. Forward modeling results. (A) Partial correlations were calculated based on actual EEG and

predicted EEG from models based on acoustic onsets, spectrogram, phonetic features, surprisal, and word

onset using MATLAB's partialcorr function. Significance is indicated by * if $p < 0.05$, ** if $p < 0.01$, and

*** if p < 0.001 using within-feature pairwise t-tests with Tukey's HSD correction. (B top) Spectrogram

and (B bottom) phonetic feature TRF model weights across the experimental conditions. The model weights

were averaged across the 12 frontotemporal electrodes depicted in Figure 1.

A subsequent LME model analysis (Equation 4) was conducted to examine how the neural

representation of different speech features decreases with increasing levels of noise. This helps

account for the fact that EEG partial correlation coefficients can vary greatly between participants

(based on, for example, cortical folding or skull/scalp geometry). Since we were interested in

trends across successive noise levels, SNR was represented using a linear polynomial contrast

which tests for a monotonic trend across these ordered noise levels. In this case, the order was set

from quiet to -9 dB so that the results would be easily comparable to the partial correlation results

shown in Figure 3A. Due to the large number of variables included in the model, a reference level

was also needed. The acoustic onset category of the *feature* variable was set as the reference level

using a dummy coding scheme. Since the reference level also serves as the model intercept when

dummy coding, every other category of the *feature* variable represents the adjustment to that

intercept. Similarly, the interaction coefficients can be interpreted as an adjustment to the *SNR*

variable because, due to the dummy coding scheme, it represents how partial correlations for the

acoustic onset model varied across SNRs (Brown, 2021).

The LME model (R^2_m = 0.086, R^2_c = 0.111) results are shown in Table 1. The raw fixed

effect outputs are listed in Section 1. Section 2 reports the estimated marginal slopes for each

feature by SNR interaction, along with statistics testing whether each slope differs from zero.

Finally, Section 3 reports Tukey HSD corrected pairwise comparisons between the interaction

effects from Section 2. With acoustic onsets as the reference level and SNR coded using a linear

polynomial contrast, the main effect coefficients quantify overall differences between acoustic

onsets and every other feature evaluated at the mean SNR. As such, each feature's main effects

differed significantly from the acoustic onset reference, indicating feature-specific differences in

partial correlation strength at the mean SNR ($p = 0.010$ for spectrogram and $p < 2.22 \times 10^{-16}$ for

phonetic features, surprisal, and word onset).

We were particularly interested in the interaction effects, since we wanted to know how

the partial correlations for each model varied across SNRs. EEG predictions based on the speech

spectrogram had the flattest slope across noise conditions ($\beta = -3.533 \times 10^{-4}$, $p = 0.561$, Table 1,

Section 2) suggesting that after controlling for all other features, the speech spectrogram exhibited

the least change in how it was reflected in the EEG as noise levels increased. Another notable

finding was that the changes in partial correlation coefficients for both the phonetic feature ($\beta = -$

6.511×10^{-3}) and surprisal ($\beta = -5.355 \times 10^{-3}$, Table 1, Section 2) models were similar to one

another ($p = 0.663$), yet different from all other models ($p < 0.05$, Table 1, Section 3). The partial

correlation coefficient trends for the phonetic feature and surprisal models decreased at a faster

rate compared to the other features which suggests that phonetic feature and surprisal

representations were more sensitive to noise. Altogether, the LME model analysis and post-hoc

comparisons showed that neural representations of the speech spectrogram were least affected by

noise and that the higher-level speech feature representations were more sensitive to increasing

levels of noise than the lower-level acoustic feature representations.

Table 1. LME model results which show how the partial correlation coefficients changed across SNRs in
addition to post-hoc contrasts between each feature.

Section 1: LME model outputs					
Fixed Effect	Estimate	Std. Error	t-value	p-value	
Intercept (Ons$_A$)	2.038×10^{-2}	1.046×10^{-3}	19.486	$< 2.22 \times 10^{-16}$	***
Sgram	2.219×10^{-3}	8.590×10^{-4}	2.583	9.798×10^{-3}	**
Fea	-1.106×10^{-2}	8.590×10^{-4}	-12.872	$< 2.22 \times 10^{-16}$	***
Surp	-7.084×10^{-3}	8.590×10^{-4}	-8.246	$< 2.22 \times 10^{-16}$	***
Ons$_W$	-1.559×10^{-2}	8.590×10^{-4}	-18.150	$< 2.22 \times 10^{-16}$	***
SNR (Ons$_A$)	-2.981×10^{-3}	6.074×10^{-4}	-4.907	9.406×10^{-7}	***
SNR:Sgram	2.628×10^{-3}	8.591×10^{-4}	3.059	2.230×10^{-3}	**
SNR:Fea	-3.530×10^{-3}	8.591×10^{-4}	-4.109	4.007×10^{-5}	***
SNR:Surp	-2.374×10^{-3}	8.591×10^{-4}	-2.764	5.721×10^{-3}	**
SNR:Ons$_W$	-8.956×10^{-6}	8.591×10^{-4}	-0.010	0.992	
Section 2: Interaction variable statistics					
Feature	SNR:Feature Slope	Std. Error	z-ratio	p-value	
Ons$_A$	-2.981×10^{-3}	6.074×10^{-4}	-4.907	< 0.001	***
Sgram	-3.533×10^{-4}	6.074×10^{-4}	-0.582	0.561	
Fea	-6.511×10^{-3}	6.074×10^{-4}	-10.718	< 0.001	***
Surp	-5.355×10^{-3}	6.074×10^{-4}	-8.816	< 0.001	***
Ons$_W$	-2.990×10^{-3}	6.074×10^{-4}	-4.922	< 0.001	***
Section 3: Comparisons between interaction variables					
Contrast	Estimate	Std. Error	t-ratio	p-value	
Ons$_A$ – Sgram	-2.628×10^{-3}	8.591×10^{-4}	-3.059	0.019	*
Ons$_A$ – Fea	3.530×10^{-3}	8.591×10^{-4}	4.109	4×10^{-4}	***
Ons$_A$ – Surp	2.374×10^{-3}	8.591×10^{-4}	2.764	0.045	*
Ons$_A$ – Ons$_W$	8.956×10^{-6}	8.591×10^{-4}	0.010	1.000	
Sgram – Fea	6.158×10^{-3}	8.591×10^{-4}	7.168	$< 1 \times 10^{-4}$	***
Sgram – Surp	5.002×10^{-3}	8.591×10^{-4}	5.823	$< 1 \times 10^{-4}$	***
Sgram – Ons$_W$	2.637×10^{-3}	8.591×10^{-4}	3.069	0.018	*
Fea – Surp	-1.156×10^{-3}	8.591×10^{-4}	-1.345	0.663	
Fea – Ons$_W$	-3.521×10^{-3}	8.591×10^{-4}	-4.099	4×10^{-4}	***
Surp – Ons$_W$	-2.365×10^{-3}	8.591×10^{-4}	-2.754	0.047	*

Another way to examine the effect of SNR on speech feature encoding is to visualize the
TRF models themselves. In fact, recent work has shown that the amplitude and latency of acoustic
onset and semantic TRFs are influenced by speech SNR (Yasmin et al., 2023). Given that the

encoding of specific frequency bands or phonetic properties may decrease differentially with noise,

we examined the change in the spectrogram and phonetic feature TRF weights at each SNR. Since

the main positive peaks are seen within the first 200 ms, the model weights were averaged from

0-200ms to better examine how the TRF weights changed across that time range (S1 Figure).

In quiet, we saw the strongest TRF weights in the lowest frequency bands, suggesting the

importance of low frequency spectral tracking when no noise is present. Notably, model weights

from 250-1,800 Hz seem to diminish in the -6 dB and -9 dB SNR conditions compared to quiet (p

< 0.05 and $p < 0.001$, pairwise tests with Tukey correction). Those same model weights were

different in -9 dB compared to the +3 dB condition ($p < 0.01$, paired tests with Tukey correction,

Figure 3B top, S1 Figure). These results support the importance of lower frequencies in speech

encoding which is reasonable given evidence that this range is useful under marginally intelligible

conditions (Bröhl & Kayser, 2021) and when supplementing listening at higher noise levels

(Chang et al., 2006; Turner et al., 2004). However, it is important to note that although there was

a reduction in spectrogram TRF weights across SNRs, the spectrogram partial correlation

coefficients did not show a similar pattern. This is likely due to the TRF being fit on the individual

feature, whereas the partial correlation analysis controlled for the contribution of all other features.

Therefore, the TRF weight results should be interpreted with caution.

In the phonetic feature case, TRF amplitudes decreased for the majority of features as noise

levels increased (Figure 3B bottom). Voicing and vowel backness features (central, back, and

diphthong) on average remained largely intact in the +3 dB condition compared to quiet ($p > 0.05$

for each, paired t-test with Tukey correction, S1 Figure) but decreased in all other conditions

(including the front feature, $p < 0.05$ for each, paired t-test with Tukey correction, S1 Figure 1,

Figure 3B bottom). In the -6 dB SNR condition, only some fricative and vowel backness weights

remained whereas all other features here and in the -9 dB SNR condition diminished (Figure 3B

bottom). Although TRF weights for some individual features decreased in the +3 dB SNR, we

witnessed a larger reduction in TRF amplitudes in the -6 dB and -9 dB SNRs similar to

Swaminathan and Heinz who found their greatest lapse in phonetic feature reception around their

-5 dB and -10 dB SNRs (Swaminathan & Heinz, 2012).

Feature-specific brain–behavior relationships change with SNR

One of the primary hypotheses of this study was that EEG signatures of linguistic

processing would be more closely related to behavior than EEG measures of low-level acoustic

processing, particularly as noise levels increased. To test this, we modeled the relationship between

behavioral scores and each speech feature's partial correlations across SNRs using GLMMs

(Equation 5) with logistic link functions. There is one GLMM for each behavioral metric and all

partial correlations are included in each model to control for the effects of the other speech features.

In this framework, the main effect of each feature reflects the association between partial

correlations and behavior averaged across noise levels, whereas the feature by SNR interaction

tests whether the strength of this association depends on SNR. SNR was again represented using

a linear polynomial contrast to test for a monotonic trend across noise levels. Here, the order of

this variable was set from -9 dB to quiet to represent an increase in SNR. Both behavioral metrics

were scaled between 0 and 1 and the predictors were z-scored prior to modeling.

S2 Table displays information about each GLMM such as the intercept, fixed effect

estimates, standard error for the fixed effect, z-value, and p-value. Figure 4 depicts the main effect

of each feature, with significant interactions shown in inset plots. The predicted lines in each

subplot were generated from the GLMMs using *ggpredict* (from the *ggeffects* package) and

represent the expected behavioral scores for each combination of predictors, averaged over

participant-level random effects. Because logistic link functions were used, the fixed effect

estimates represent the amount that the log odds of the behavioral scores change given a unit

increase in the partial correlation and partial correlation by SNR coefficients.

The PWH GLMM ($R^2_m = 0.944$, $R^2_c = 0.997$) showed robust overall effects of SNR, such

that PWH scores increased as SNR increased ($\beta = 2.132$, $p < 2.22 \times 10^{-16}$). Among the neural

tracking predictors, a significant main effect was observed only for surprisal-related neural

tracking, indicating that increased neural tracking of lexical surprisal was associated with higher

PWH scores ($\beta = 0.398$, $p = 3.652 \times 10^{-5}$, Figure 4, top). Critically, however, this relationship

between behavior and neural tracking varied across noise levels as reflected by a significant

surprisal by SNR interaction, demonstrating that the relationship between lexical surprisal tracking

and PWH strengthened as SNR increased ($\beta = 0.265$, $p = 7.067 \times 10^{-3}$). A significant interaction

was also observed for acoustic onsets, such that the relationship between acoustic onset tracking

and PWH weakened as SNR increased ($\beta = -0.258$, $p = 4.820 \times 10^{-3}$). Intuitively, this pattern

suggests that as listening conditions improve, word report scores become less linked to low-level

acoustic onset tracking and increasingly linked to high-level lexical surprisal tracking.

For the comprehension score GLMM ($R^2_m = 0.771$, $R^2_c = 0.986$), behavioral performance

again increased as SNR increased ($\beta = 0.830$, $p < 0.001$). While positive effects of spectrogram (β

$= 0.128$, $p = 0.042$) and surprisal ($\beta = 0.122$, $p = 0.046$) tracking indicated overall associations

with comprehension scores across SNRs, only phonetic-feature tracking exhibited a significant

interaction with SNR ($\beta = -0.124$, $p = 0.033$). Phonetic-feature tracking appears to play a more

prominent role in supporting comprehension when speech is acoustically degraded, but it becomes

less informative as SNR improves. Overall, these results suggest a noise-dependent division of

labor across neural feature representations, with PWH score reliance shifting from acoustic to

lexical tracking as listening conditions improve, and comprehension score reliance on phonetic-

level tracking decreasing as listening conditions improve.

Figure 4. The relationship between the partial correlation coefficients and behavior. GLMMs were used to

relate the neural tracking of each speech feature to participants' PWH (top) and comprehension scores

(bottom). Both models were fit using an ordered beta distribution and a logistic link function. The black

lines represent GLMM-predicted behavioral scores (via *ggpredict*) that show the expected effect of each

speech feature's tracking while controlling for SNR and tracking of all other features. The shaded grey

regions are 95% confidence intervals, and the colored dots represent each participant's average partial

correlation coefficient in each SNR. The inset plots show predicted behavioral scores for features whose

relationship with behavior significantly varied across SNRs. Each SNR is represented by a fitted line, with

lightly shaded areas indicating the 95% confidence intervals.

 Envelope reconstruction accuracy maps well to measures of behavior

Numerous studies have shown that cortical activity tracks the temporal modulations of the

speech envelope (Ahissar et al., 2001; Aiken & Picton, 2008; Destoky et al., 2019; Di Liberto et

al., 2015; Ding & Simon, 2013; Etard & Reichenbach, 2019; Lalor & Foxe, 2010; Nourski et al.,

2009; Pasley et al., 2012). Speech envelope reconstruction is a powerful method of indexing

speech tracking, given its advantage of using all scalp data to provide a better signal to noise ratio.

Yet, it is unknown exactly what information the speech envelope carries because it has been shown

to relate to syllabic boundaries (Hertrich et al., 2012; Oganian & Chang, 2019), phonetic feature

information (Rosen, 1992), and prosodic cues (Myers et al., 2019). Nevertheless, due to its

common usage in speech research and its improved SNR, we were interested in how speech

envelope tracking would change across our experimental conditions and how it would relate to

behavior apart from the other speech features.

Figure 5. Envelope modeling results. (A) Colored boxplots show envelope reconstruction accuracies for

each condition, and the black points represent individual participants. Significance is indicated by * if p <

0.05, ** if p < 0.01, and *** if p < 0.001 using pairwise t-tests with Holm correction. (B, C) The black lines

indicate GLMM-predicted PWH (B) and comprehension scores (C) via *ggpredict* as a function of envelope

tracking while controlling for SNR. The shaded regions represent the 95% confidence intervals, and each

colored dot represents each participant's average reconstruction accuracy in each SNR.

The first step in this analysis was to use a backward modeling approach to determine how

speech envelope tracking was affected by different levels of background noise. Speech in quiet

and in the +3 dB SNR condition were reconstructed to a similar degree ($p = 0.305$, paired t-test

with Holm correction) and were reconstructed more reliably than all other SNRs ($p < 0.05$). The -

3 dB SNR and -6 dB SNR accuracies were not significantly different from each other ($p = 0.305$)

but were both different from the -9 dB SNR condition ($p = 1.748 \times 10^{-7}$ and $p = 2.347 \times 10^{-6}$,

respectively, Figure 5A).

Next, we tested if the fidelity of envelope tracking was indicative of how well participants

could hear and understand the story. This was tested using a GLMM model (Equation 5) that

contained envelope reconstruction accuracies, SNR (using a linear polynomial contrast), and the

interaction between the two as fixed effects and participants as the random effect. Envelope

tracking showed a positive relationship with both PWH ($\beta = 0.356$, $p = 1.991 \times 10^{-3}$, $R^2_m = 0.926$,

$R^2_c = 0.996$) and comprehension scores ($\beta = 0.178$, $p = 0.040$, $R^2_m = 0.804$, $R^2_c = 0.983$). Increases

in SNR were also associated with increases in behavioral scores (PWH: $\beta = 2.019$, $p = 2.268 \times 10^{-15}$

; comprehension: $\beta = 0.840$, $p = 2.190 \times 10^{-6}$). The interaction between reconstruction accuracy

and SNR, however, was not significant in the PWH ($\beta = 0.096$, $p = 0.247$) GLMM nor the

comprehension ($\beta = 0.043$, $p = 0.416$) GLMM. This indicates that the relationship between

envelope tracking and behavioral scores does not vary across SNRs. In sum, neural tracking of the

speech envelope provides a consistent predictor of both PWH and comprehension scores,

independent of listening condition.

Lexical surprisal's influence on early auditory encoding decreases at high noise levels

Previous work by Broderick and colleagues has shown that the cortical tracking of an

individual word's acoustic representations was enhanced the more semantically similar it was to

its preceding context (Broderick et al., 2019). Since higher-level representations can bias

perception when incoming stimuli are noisy (de Lange et al., 2018), perhaps participants in the

present study relied more on a higher-level feature (i.e., next word probability/surprisal) when

noise levels slightly increased—thereby strengthening the influence of word surprisal on lower-

level feature encoding. This influence of lexical surprisal should then decrease the noisier the

speech becomes, i.e., as people begin to fail to understand the speech. We aimed to test these

hypotheses using a two-stage regression analysis that consisted of stimulus envelope

reconstruction followed by linear mixed effects modeling.

The backward modeling technique helps improve overall signal to noise ratio by

simultaneously training on all 128 electrodes to reconstruct the (single channel) speech envelope.

Because of this benefit, backward models were used in the two-stage regression rather than

forward models. The decoders were used to reconstruct an estimate of the speech envelopes.

Spearman's correlation was then computed between the first 100 ms of the reconstructed and actual

envelopes for each individual word. All words in the story were scored on their lexical surprisal,

which was calculated as the negative logarithm of a word's probability given the words that came

before it. Envelope variability, relative pitch, and resolvability (the nuisance regressors) were also

calculated to control for acoustic related properties of the stimuli, many of which could correlate

with word surprisal and word reconstruction accuracy. Surprisal, SNR, and the nuisance regressors

were included in an LME model to predict word reconstruction accuracy for the first 100 ms of

each word.

There was a significant positive relationship between lexical surprisal and word

reconstruction accuracy in quiet ($\beta = 2.850$ x 10^{-2}, p < 2.22 x 10^{-16}, $R^2_m = 0.017$, $R^2_c = 0.030$, Table

2). In other words, the more surprising a word (or the less probable that word was given its

preceding context), the better that word's envelope was decoded from the EEG. The proportion of

variance explained by the present LME model, $R^2_m = 0.017$, is consistent with what was found by

Broderick and colleagues, $R^2 = 0.0171$ (Broderick et al., 2019). The interaction between surprisal

and SNR shows how the *surp* coefficient changes (or is adjusted) across noise levels (Figure 6A,

Table 2, S3 Table). For example, whereas surprisal in quiet has a coefficient of 2.850 x 10^{-2},

surprisal in the +3 dB condition has a coefficient (or slope) that is 2.172x 10^{-4} units lower (S3

Table, top). The influence of lexical surprisal on word reconstruction accuracy in quiet was similar

to the +3 dB ($\beta = -2.172$x 10^{-4}, p $= 0.951$) and -3 dB ($\beta = -4.767$x 10^{-3}, p $= 0.173$) SNR conditions

and greater than the -6 dB ($\beta = -9.900$ x 10^{-3}, p $= 0.005$) and -9 dB ($\beta = -1.600$ x 10^{-2}, p $= 4.890$ x

10^{-6}, Table 2) SNR conditions. This shows that lexical surprisal impacts envelope tracking in quiet

and in low levels of background noise (+3 dB and -3 dB SNRs), but that this influence significantly

decreases with moderate to high levels of noise. It is also important to note that the nuisance

regressors and additional interactions included in the LME model also significantly influenced

word reconstruction accuracy, reinforcing the value of controlling for those variables and ensuring

our results were not confounded by other acoustic related measures.

Table 2. LME model showing the relationship between lexical surprisal and envelope tracking in each

SNR.

Fixed Effect	Estimate	Std. Error	t-value	p-value	
Intercept (Quiet)	8.992×10^{-2}	5.451×10^{-3}	16.496	$< 2 \times 10^{-16}$	***
+3 dB	-4.703×10^{-3}	3.514×10^{-3}	-1.338	0.18077	
-3 dB	-1.671×10^{-2}	3.524×10^{-3}	-4.724	2.12×10^{-6}	***
-6 dB	-3.196×10^{-2}	3.496×10^{-3}	-9.141	$< 2 \times 10^{-16}$	***
-9 dB	-5.822×10^{-2}	3.511×10^{-3}	-16.586	$< 2 \times 10^{-16}$	***
Surp	2.850×10^{-2}	2.709×10^{-3}	10.519	$< 2 \times 10^{-16}$	***
Surp:+3 dB	-2.172×10^{-4}	3.528×10^{-3}	-0.062	0.95091	
Surp:-3 dB	-4.767×10^{-3}	3.497×10^{-3}	-1.363	0.17287	
Surp:-6 dB	-9.900×10^{-3}	3.514×10^{-3}	-2.817	0.00484	**
Surp:-9 dB	-1.600×10^{-2}	3.501×10^{-3}	-4.570	4.89×10^{-6}	***
EnvStd	6.042×10^{-2}	1.430×10^{-3}	42.251	$< 2 \times 10^{-16}$	***
F_{rel}	8.697×10^{-3}	1.576×10^{-3}	5.519	3.42×10^{-8}	***
Res	-1.605×10^{-2}	1.453×10^{-3}	-11.051	$< 2 \times 10^{-16}$	***
EnvStd:f_{rel}	6.692×10^{-3}	1.148×10^{-3}	5.832	5.49×10^{-9}	***
F_{rel}:res	-6.327×10^{-3}	1.450×10^{-3}	-4.363	1.28×10^{-5}	***
Surp:envStd	1.624×10^{-2}	1.305×10^{-3}	12.448	$< 2 \times 10^{-16}$	***
Surp:res	-6.872×10^{-3}	1.325×10^{-3}	-5.186	2.17×10^{-7}	***

Figure 6. The relationship between lexical surprisal and envelope tracking at different levels of noise. (A)

The surprisal coefficient in quiet and the interaction between surprisal and SNR using a 100 ms word

window. The error bars are the standard errors of each measure, calculated using the LME model.

Significance is indicated by * if $p < 0.05$, ** if $p < 0.01$, and *** if $p < 0.001$ using R's *emtrends* function

(*emmeans* package) to compare estimated marginal means of linear trends. (B, C) The relationship between

the surprisal coefficients across SNRs and behavior. The percentage of words heard (PWH) scores are in B

and comprehension scores are in C.

Despite studies showing that low SNRs benefit from the utilization of context (Mayo et al.,

1997), our finding that lexical surprisal significantly influences envelope tracking for the -6 dB (z-

ratio = 6.814, p < 0.001) and -9 dB conditions (z-ratio = 4.615, p < 0.001, Holm corrected test that

each SNR by surprisal trend is greater than 0, S3 Table) is unexpected. Participants reported

hearing either no words or very few words in, for instance, the -9 dB SNR condition, yet we still

see a significant influence of lexical surprisal on envelope tracking. We did not control for any

other features (including surprisal) in our envelope reconstruction model, so these unaccounted-

for features may have contributed to the significantly positive interaction coefficients in the -6 dB

and -9 dB SNR conditions. Or the fact that some participants were able to hear some words in both

conditions may have been enough to cause this significant effect. Lastly, we found that single

subject surprisal coefficients also predicted their self-reported percentage of words heard ($\beta =$

0.358, $p = 0.001$, $R^2_m = 0.384$, Figure 6B) and comprehension scores ($\beta = 3.558$, $p = 1.906$ x 10^{-5},

$R^2_m = 0.533$, $R^2_c = 686$) across SNRs (Equation 6, Figure 6C). That is to say, the stronger the

influence of surprisal on envelope tracking, the better the participants were able to hear and

comprehend the story.

DISCUSSION

This study sought to establish how well indices of hierarchical neural speech processing

reflect language comprehension—advancing on prior work that has typically tested specific

hierarchical levels without controlling for the others. First, the encoding of a range of hierarchical

speech features was characterized to assess how this encoding changed with noise and if those

changes related to behavior. Linguistic feature representations were more affected by noise than

the acoustic representations. In addition, associations between behavioral performance and speech

feature tracking were task- and condition-dependent: PWH scores became decreasingly linked to

acoustic onset tracking and increasingly linked to lexical surprisal tracking as SNR improved,

whereas comprehension scores became increasingly linked to phonetic feature as SNR worsened.

Speech envelope tracking fidelity was significantly associated with behavioral scores, and this

relationship remained consistent across SNRs. The final set of analyses investigated the influence

of lexical surprisal on neural tracking of the speech envelope. Results showed that the envelopes

of more unexpected words were strongly reflected in the EEG in quiet and in low levels of noise,

but this relationship weakened as noise levels increased.

Neural representations of acoustic and phonetic features show dissociable sensitivity to noise

Acoustic feature representations were hypothesized to be more invariant to background

noise than linguistic feature representations; this pattern was supported by the present results. The

degree to which spectrogram information was reflected in EEG remained unchanged across SNRs.

This result contrasts with studies such as Lesenfants and colleagues whose spectrogram model

performance increased monotonically (as noise levels *decreased*) when training on delta band EEG

and increased in a more S-shaped pattern when training on theta band EEG (Lesenfants et al.,

2019). These differences may be partly explained by the efforts in the present study to partial out

the contribution of other speech features when assessing model performance.

Interestingly, phonetic features were uniquely represented in neural activity even when

accounting for the speech spectrogram and acoustic onsets. This is consistent with previous studies

showing that the addition of phonetic features to spectrotemporal representations improves EEG

prediction (Di Liberto et al., 2015; Sohoglu & Davis, 2020; Tezcan et al., 2023) and correlations

with speech intelligibility (Lesenfants et al., 2019), and that phonetic features can uniquely predict

EEG responses even when attending to a specific talker (Teoh et al., 2022). However, the present

phonetic feature results contrast with previous work which suggested that responses to

articulations could be explained by simpler acoustic features (Daube et al., 2019). Moreover, the

differential decline of phonetic feature and spectrogram encoding, along with their distinct

relationships to behavior, provides further evidence that the two features are dissociable.

When assessing speech envelope decoding on the other hand, the reconstruction accuracies

first began to decrease in the -3 dB SNR and decreased yet again in the -9 dB SNR. This pattern

was surprising given a prior report that the synchronization between neural activity and the speech

envelope remained largely unaffected until the speech signal reached an SNR of -9 dB (Ding &

Simon, 2013). These contrasting results may be attributed to a combination of factors: neural

recording modality, data preprocessing, model training and testing procedures between conditions,

or the regularization method used (e.g., boosting versus ridge regression). Instead, the present

results show a gradual decrease in envelope tracking across SNRs similar to findings from

Vanthornhout and colleagues (Vanthornhout et al., 2018). However, the present results indicate

that envelope reconstruction accuracies were significantly associated with participants' behavioral

scores which is consistent with many studies that have shown that the speech envelope (using

stimulus reconstruction or cross-correlation) contributes and relates to speech intelligibility and

comprehension (Ahissar et al., 2001; Decruy et al., 2020; Iotzov & Parra, 2019; Lesenfants et al.,

2019; Muncke et al., 2022; Vanthornhout et al., 2018).

Behavioral relevance of acoustic and linguistic features varies with noise level

When further examining the relationship between feature-specific neural tracking and

behavioral performance, the present results revealed a more nuanced pattern than originally

hypothesized. Rather than observing a uniform dominance of higher-level linguistic feature

tracking across listening conditions, the behavioral relevance of neural representations depended

on both listening condition and the behavioral metric being assessed. The association between

what participants reported hearing and lexical surprisal tracking strengthened as speech became

cleaner (less noisy). This finding is consistent with prior work which demonstrated that lexical

surprisal tracking is prominent when speech is comprehensible (Gillis et al., 2023; Mesik et al.,

2021) and that this tracking weakens as speech becomes harder to understand (Verschueren et al.,

2022). In contrast, the relationship between PWH and acoustic onset tracking weakened as speech

became cleaner. This suggests that this low-level cue was more behaviorally relevant when the

speech signal was degraded (Davis & Johnsrude, 2007; Mattys et al., 2005).

A different pattern was observed for comprehension scores. Both spectrogram and lexical

surprisal based representations were positively associated with comprehension scores across

listening conditions, suggesting stable contributions of acoustic and linguistic processing to speech

understanding. Notably, however, only phonetic feature tracking showed a significant interaction

with SNR such that its association with comprehension decreased as noise levels decreased. This

pattern indicates that the ability to convert acoustic speech sounds into phonetic-level

representations in noisy conditions supports comprehension of that speech under those conditions.

Across behavioral measures, these findings suggest that the behavioral relevance of neural

representations shifts across hierarchical levels as listening conditions change. Specifically, lower-

level acoustic and phonetic information contributes more strongly under noisy conditions, whereas

higher-level linguistic predictions become most informative when speech is in quiet. Consistent

with prior work showing that speech in noise perception is best predicted by models that combine

neural measures across multiple levels (Lesenfants et al., 2019), the present study complements

this finding by showing that speech perception and comprehension rely in parallel on acoustic and

phonetic representations. Our approach reveals how the behavioral relevance of hierarchical

speech feature tracking is differentially weighted across listening conditions, an insight that may

be less apparent in models combining multiple speech features.

Lexical context can predict the fidelity of word-level acoustic tracking, but this effect decreases

as noise levels increase

Another key hypothesis of our study was that participants would use lexical context to

predict and encode the acoustic features of each word. This hypothesis was supported: the LME

analysis (stage two of the two-stage regression) revealed that the more unexpected a word, the

better that word's envelope was reconstructed. However, we also hypothesized that participants

would rely more on these predictions for speech in moderate levels of noise (when speech is still

intelligible) relative to speech in quiet, before declining at high levels of background noise (when

speech is no longer intelligible). This result was only partially borne out. Specifically, the use of

lexical context in processing speech acoustics did decrease as the speech became noisier, but there

was no evidence to support a stronger reliance on context in moderate levels of noise. In particular,

while there was no difference in comprehension scores between the quiet and +3 dB conditions,

there was no increase in the influence of surprisal on envelope tracking for the latter condition

compared to speech in quiet. However, a larger spread of the percentage of words heard scores

was observed across participants in the +3 dB condition. This raised the possibility that participants

who were starting to struggle might put forth more effort to understand and process the story by

relying more on context (and thus might have a higher surprisal weight in Figure 6A) than those

who remained at ceiling. Nevertheless, no significant difference was observed to support this

possibility. This was a little surprising given that context has been known to affect behavior

(Golestani et al., 2013) and neural activity (Boulenger et al., 2011; Koskinen et al., 2020; Strauß

et al., 2022) in challenging listening conditions. Future work with larger participant numbers and

perhaps even lower levels of background noise (e.g., + 6 dB SNR) might reveal such an effect.

The present second-stage regression analysis based on word surprisal seems to be at odds

with Broderick and colleagues who found that the envelopes of words that were more semantically

similar to their context were better reflected in the EEG. That is to say, envelope tracking is

enhanced for words that share a similar meaning with their context (Broderick et al., 2019).

However, semantic similarity and lexical surprisal tend to share a moderate, and sometimes weak

negative correlation (Frank & Willems, 2017). Indeed, a re-analysis of Broderick et al.'s original

EEG data has revealed that both semantic similarity and lexical surprisal play complementary

(positive) roles in estimating when envelope tracking is enhanced (Broderick & Lalor, 2020).

Nevertheless, the nature of this duality remains mysterious, and we hope it will provide the grounds

for an exciting body of future work. We anticipate it will take a substantial battery of future

experiments to shape a unifying explanation, with stimuli that can disentangle correlations between

semantic similarity, lexical surprisal, and other linguistic factors that could come into play (e.g.,

semantic content, next-word entropy, phonetic surprisal, next-phoneme entropy). In any case, what

seems clear in the present results is that lexical context influences the neural tracking of speech

acoustics on a word-by-word basis, and this influence drops as speech becomes unintelligible.

Methodological limitations

The present study aimed to characterize how hierarchical speech feature encoding changes

in increasing levels of background noise. One inherent challenge was the selection of speech

features, particularly given the relatively low signal to noise ratio of EEG and the limited amount

of EEG data per participant. Numerous candidate features have been used in prior work such as

the spectrogram derivative (Teoh et al., 2022), log-mel spectrogram (Daube et al., 2019), phoneme

onset, cohort entropy (Brodbeck, Hong, et al., 2018), and semantic dissimilarity (Broderick et al.,

2018). As such, we opted to focus on a set of commonly used features spanning acoustic, phonetic,

and lexical levels to balance interpretability with model complexity.

A related challenge is that speech features are likely to correlate with one another and

explain similar activity in the EEG. This motivated our use of the partial correlation analyses to

estimate the unique contribution of each feature to the neural responses while controlling for the

effect of others. While this approach provides a principled way to disentangle correlated

representations, results will partly depend on the specific feature set included.

Another potential limitation of this study was the mode of regularization. We optimized

the regularization parameter on the clean speech and used this same parameter on all other

conditions. This choice was motivated by the desire to reduce overfitting, as this would be

particularly prudent at lower SNRs (i.e., -6 dB and -9 dB) where multidimensional feature models

would be more susceptible to noise-driven fits. Although this approach may reduce sensitivity to

condition-specific changes in optimal regularization, it provides a conservative and unbiased basis

for comparing model performance across listening conditions.

Conclusion

In summary, the current results showed that neural representations of lexical surprisal and

phonetic features were more susceptible to noise than acoustic speech feature representations.

Behavioral performance reflected contributions from multiple hierarchical levels of speech

processing, with the relative predictive value of acoustic, phonetic, and lexical features varying

with noise level and task. We also found that the encoding of certain phonetic features decreased

even in low levels of noise, and that the encoding of frequencies below 1.3k essentially disappeared

in high levels of noise. Lastly, we showed that context—in the form of a word predictability—

influenced how the acoustics of those words were encoded. This influence lessens in high

background noise levels. Future work will aim to further characterize how people might rely more

or less on top-down context to process bottom-up speech input as a function of stimulus type, task,

and listening conditions.

REFERENCES

Ahissar, E., Nagarajan, S., Ahissar, M., Protopapas, A., Mahncke, H., & Merzenich, M. M. (2001).
Speech comprehension is correlated with temporal response patterns recorded from
auditory cortex. *Proceedings of the National Academy of Sciences*, *98*(23), 13367–13372.

Aiken, S. J., & Picton, T. W. (2008). Human cortical responses to the speech envelope. *Ear Hear*,
29(2), 139–157. https://doi.org/10.1097/aud.0b013e31816453dc

Anderson, A. J., Binder, J. R., Fernandino, L., Humphries, C. J., Conant, L. L., Aguilar, M., Wang,
X., Doko, D., & Raizada, R. D. S. (2017). Predicting Neural Activity Patterns Associated
with Sentences Using a Neurobiologically Motivated Model of Semantic Representation.
Cereb Cortex, *27*(9), 4379–4395. https://doi.org/10.1093/cercor/bhw240

Atal, B. S., & Hanauer, S. L. (1971). Speech analysis and synthesis by linear prediction of the
speech wave. *The journal of the acoustical society of America*, *50*(2B), 637–655.

Bigdely-Shamlo, N., Mullen, T., Kothe, C., Su, K. M., & Robbins, K. A. (2015). The PREP pipeline:
standardized preprocessing for large-scale EEG analysis. *Front Neuroinform*, *9*, 16.
https://doi.org/10.3389/fninf.2015.00016

Boersma, P., & Weenink, D. (2013). Praat: doing phonetics by computer [Computer program].
Version 5.3. 51. *Online: http://www. praat. org/retrieved, last viewed on*, *12*.

Boulenger, V., Hoen, M., Jacquier, C., & Meunier, F. (2011). Interplay between acoustic/phonetic
and semantic processes during spoken sentence comprehension: An ERP study. *Brain
and language*, *116*(2), 51–63.

Brodbeck, C., Hong, L. E., & Simon, J. Z. (2018). Rapid Transformation from Auditory to Linguistic
Representations of Continuous Speech. *Curr Biol*, *28*(24), 3976–3983 e3975.
https://doi.org/10.1016/j.cub.2018.10.042

Brodbeck, C., Jiao, A., Hong, L. E., & Simon, J. Z. (2020). Neural speech restoration at the cocktail
party: Auditory cortex recovers masked speech of both attended and ignored speakers.
PLoS Biol, *18*(10), e3000883. https://doi.org/10.1371/journal.pbio.3000883

Brodbeck, C., Presacco, A., & Simon, J. Z. (2018). Neural source dynamics of brain responses to
continuous stimuli: Speech processing from acoustics to comprehension. *Neuroimage*,
172, 162–174. https://doi.org/10.1016/j.neuroimage.2018.01.042

Broderick, M. P., Anderson, A. J., Di Liberto, G. M., Crosse, M. J., & Lalor, E. C. (2018).
Electrophysiological Correlates of Semantic Dissimilarity Reflect the Comprehension of
Natural, Narrative Speech. *Curr Biol*, *28*(5), 803–809 e803.
https://doi.org/10.1016/j.cub.2018.01.080

Broderick, M. P., Anderson, A. J., & Lalor, E. C. (2019). Semantic Context Enhances the Early
Auditory Encoding of Natural Speech. *J Neurosci*, *39*(38), 7564–7575.
https://doi.org/10.1523/JNEUROSCI.0584-19.2019

Broderick, M. P., & Lalor, E. C. (2020). Co-existence of prediction and error signals in
electrophysiological responses to natural speech. *bioRxiv*, 2020.2011. 2020.391227.

Broderick, M. P., Zuk, N. J., Anderson, A. J., & Lalor, E. C. (2022). More than words:
Neurophysiological correlates of semantic dissimilarity depend on comprehension of the
speech narrative. *European journal of neuroscience*, *56*(8), 5201–5214.

Bröhl, F., & Kayser, C. (2021). Delta/theta band EEG differentially tracks low and high frequency
speech-derived envelopes. *Neuroimage*, *233*, 117958.

Brooks, M. E., Kristensen, K., Van Benthem, K. J., Magnusson, A., Berg, C. W., Nielsen, A.,
Skaug, H. J., Mächler, M., & Bolker, B. M. (2017). glmmTMB balances speed and flexibility
among packages for zero-inflated generalized linear mixed modeling.

Brown, V. A. (2021). An introduction to linear mixed-effects modeling in R. *Advances in Methods
and Practices in Psychological Science*, *4*(1), 2515245920960351.

Burton, M. W. (2009). Understanding the role of the prefrontal cortex in phonological processing.
Clin Linguist Phon, *23*(3), 180–195. https://doi.org/10.1080/02699200802394963

Chang, J. E., Bai, J. Y., & Zeng, F. G. (2006). Unintelligible low-frequency sound enhances
simulated cochlear-implant speech recognition in noise. *IEEE Trans Biomed Eng*, *53*(12
Pt 2), 2598–2601. https://doi.org/10.1109/TBME.2006.883793

Crosse, M. J., Di Liberto, G. M., Bednar, A., & Lalor, E. C. (2016). The Multivariate Temporal
Response Function (mTRF) Toolbox: A MATLAB Toolbox for Relating Neural Signals to
Continuous Stimuli. *Front Hum Neurosci*, *10*, 604.
https://doi.org/10.3389/fnhum.2016.00604

Crosse, M. J., Di Liberto, G. M., & Lalor, E. C. (2016). Eye Can Hear Clearly Now: Inverse
Effectiveness in Natural Audiovisual Speech Processing Relies on Long-Term
Crossmodal Temporal Integration. *J Neurosci*, *36*(38), 9888–9895.
https://doi.org/10.1523/JNEUROSCI.1396-16.2016

Crosse, M. J., Zuk, N. J., Di Liberto, G. M., Nidiffer, A. R., Molholm, S., & Lalor, E. C. (2021).
Linear modeling of neurophysiological responses to speech and other continuous stimuli:
methodological considerations for applied research. *Frontiers in Neuroscience*, 1350.

Dai, Z., Yang, Z., Yang, Y., Carbonell, J. G., Le, Q., & Salakhutdinov, R. (2019). Transformer-xl:
Attentive language models beyond a fixed-length context. Proceedings of the 57th annual
meeting of the association for computational linguistics,

Daube, C., Ince, R. A. A., & Gross, J. (2019). Simple acoustic features can explain phoneme-
based predictions of cortical responses to speech. *Current Biology*, *29*(12), 1924–1937.
e1929.

Davis, M. H., & Johnsrude, I. S. (2007). Hearing speech sounds: top-down influences on the
interface between audition and speech perception. *Hearing research*, *229*(1-2), 132–147.

de Heer, W. A., Huth, A. G., Griffiths, T. L., Gallant, J. L., & Theunissen, F. E. (2017). The
Hierarchical Cortical Organization of Human Speech Processing. *J Neurosci*, *37*(27),
6539–6557. https://doi.org/10.1523/JNEUROSCI.3267-16.2017

de Lange, F. P., Heilbron, M., & Kok, P. (2018). How Do Expectations Shape Perception? *Trends
Cogn Sci*, *22*(9), 764–779. https://doi.org/10.1016/j.tics.2018.06.002

Decruy, L., Lesenfants, D., Vanthornhout, J., & Francart, T. (2020). Top-down modulation of
neural envelope tracking: The interplay with behavioral, self-report and neural measures
of listening effort. *Eur J Neurosci*, *52*(5), 3375–3393. https://doi.org/10.1111/ejn.14753

Delorme, A., & Makeig, S. (2004). EEGLAB: an open source toolbox for analysis of single-trial
EEG dynamics including independent component analysis. *J Neurosci Methods*, *134*(1),
9–21. https://doi.org/10.1016/j.jneumeth.2003.10.009

Destoky, F., Philippe, M., Bertels, J., Verhasselt, M., Coquelet, N., Vander Ghinst, M., Wens, V.,
De Tiege, X., & Bourguignon, M. (2019). Comparing the potential of MEG and EEG to

uncover brain tracking of speech temporal envelope. *Neuroimage*, *184*, 201–213.
https://doi.org/10.1016/j.neuroimage.2018.09.006

Di Liberto, G. M., O'sullivan, J. A., & Lalor, E. C. (2015). Low-frequency cortical entrainment to
speech reflects phoneme-level processing. *Current Biology*, *25*(19), 2457–2465.

Di Liberto, G. M., Wong, D., Melnik, G. A., & de Cheveigné, A. (2019). Low-frequency cortical
responses to natural speech reflect probabilistic phonotactics. *Neuroimage*, *196*, 237–
247.

Ding, N., & Simon, J. Z. (2013). Adaptive temporal encoding leads to a background-insensitive
cortical   representation   of   speech.   *J   Neurosci*,   *33*(13),   5728–5735.
https://doi.org/10.1523/JNEUROSCI.5297-12.2013

Dou, J., Anderson, A. J., White, A. S., Norman-Haignere, S. V., & Lalor, E. C. (2025). Dynamic
modeling of EEG responses to natural speech reveals earlier processing of predictable
words. *PLOS Computational Biology*, *21*(4), e1013006.

Drullman, R., Festen, J. M., & Plomp, R. (1994). Effect of reducing slow temporal modulations on
speech   reception.   *J   Acoust   Soc   Am*,   *95*(5   Pt   1),   2670–2680.
https://doi.org/10.1121/1.409836

Etard, O., & Reichenbach, T. (2019). Neural Speech Tracking in the Theta and in the Delta
Frequency Band Differentially Encode Clarity and Comprehension of Speech in Noise. *J
Neurosci*, *39*(29), 5750–5759. https://doi.org/10.1523/JNEUROSCI.1828-18.2019

Fisher, R. A. (1924). The distribution of the partial correlation coefficient. *Metron*, *3*, 329–332.

Frank, S. L., & Willems, R. M. (2017). Word predictability and semantic similarity show distinct
patterns of brain activity during language comprehension. *Language, Cognition and
Neuroscience*, *32*(9), 1192–1203.

Gillis, M., Vanthornhout, J., & Francart, T. (2023). Heard or understood? Neural tracking of
language features in a comprehensible story, an incomprehensible story and a word list.
eNeuro, *10*(7).

Gillis, M., Vanthornhout, J., Simon, J. Z., Francart, T., & Brodbeck, C. (2021). Neural Markers of
Speech Comprehension: Measuring EEG Tracking of Linguistic Speech Representations,
Controlling   the   Speech   Acoustics.   *J   Neurosci*,   *41*(50),   10316–10329.
https://doi.org/10.1523/JNEUROSCI.0812-21.2021

Golestani, N., Hervais-Adelman, A., Obleser, J., & Scott, S. K. (2013). Semantic versus
perceptual interactions in neural processing of speech-in-noise. *Neuroimage*, *79*, 52–61.
https://doi.org/10.1016/j.neuroimage.2013.04.049

Gwilliams, L., King, J. R., Marantz, A., & Poeppel, D. (2022). Neural dynamics of phoneme
sequences reveal position-invariant code for content and order. *Nat Commun*, *13*(1),
6606. https://doi.org/10.1038/s41467-022-34326-1

Hamilton, L. S., Edwards, E., & Chang, E. F. (2018). A Spatial Map of Onset and Sustained
Responses to Speech in the Human Superior Temporal Gyrus. *Curr Biol*, *28*(12), 1860–
1871 e1864. https://doi.org/10.1016/j.cub.2018.04.033

Heilbron, M., Armeni, K., Schoffelen, J.-M., Hagoort, P., & De Lange, F. P. (2022). A hierarchy of
linguistic predictions during natural language comprehension. *Proceedings of the National*
Academy of Sciences, *119*(32), e2201968119.

Hertrich, I., Dietrich, S., Trouvain, J., Moos, A., & Ackermann, H. (2012). Magnetic brain activity
phase-locked to the envelope, the syllable onsets, and the fundamental frequency of a
perceived   speech   signal.   *Psychophysiology*,   *49*(3),   322–334.
https://doi.org/10.1111/j.1469-8986.2011.01314.x

Hickok, G. (2015). *Neurobiology of language*. Elsevier.

Hickok, G., & Poeppel, D. (2007). The cortical organization of speech processing. *Nature Reviews*
Neuroscience, *8*(5), 393–402.

Huth, A. G., De Heer, W. A., Griffiths, T. L., Theunissen, F. E., & Gallant, J. L. (2016). Natural
speech reveals the semantic maps that tile human cerebral cortex. *Nature*, *532*(7600),
453–458.

Iotzov, I., & Parra, L. C. (2019). EEG can predict speech intelligibility. *J Neural Eng*, *16*(3), 036008.
https://doi.org/10.1088/1741-2552/ab07fe

Irino, T., & Patterson, R. D. (2006). A Dynamic Compressive Gammachirp Auditory Filterbank.
IEEE   Trans   Audio   Speech   Lang   Process,   *14*(6),   2222–2232.
https://doi.org/10.1109/TASL.2006.874669

Karunathilake, I. D., Kulasingham, J. P., & Simon, J. Z. (2023). Neural tracking measures of
speech intelligibility: Manipulating intelligibility while keeping acoustics unchanged.
Proceedings of the National Academy of Sciences, *120*(49), e2309166120.

Kleiner, M., Brainard, D., & Pelli, D. (2007). What's new in Psychtoolbox-3?

Koskinen, M., Kurimo, M., Gross, J., Hyvarinen, A., & Hari, R. (2020). Brain activity reflects the
912   predictability of word sequences in listened continuous speech. *Neuroimage*, *219*,
913   116936. https://doi.org/10.1016/j.neuroimage.2020.116936

914   Kuznetsova, A., Brockhoff, P. B., & Christensen, R. H. (2017). lmerTest package: tests in linear
915   mixed effects models. *Journal of statistical software*, *82*, 1–26.

916   Lalor, E. C., & Foxe, J. J. (2010). Neural responses to uninterrupted natural speech can be
917   extracted with precise temporal resolution. *European Journal of Neuroscience*, *31*(1),
918   189–193. https://doi.org/10.1111/j.1460-9568.2009.07055.x

919   Lenth, R., Singmann, H., Love, J., Buerkner, P., & Herve, M. (2019). Package 'emmeans'. *R
920   package version*, *1*(3.2).

921   Lesenfants, D., Vanthornhout, J., Verschueren, E., Decruy, L., & Francart, T. (2019). Predicting
922   individual speech intelligibility from the cortical tracking of acoustic- and phonetic-level
923   speech   representations.   *Hear   Res*,   *380*,   1–9.
924   https://doi.org/10.1016/j.heares.2019.05.006

925   Lüdecke, D. (2018). ggeffects: Tidy data frames of marginal effects from regression models.
926   *Journal of open source software*, *3*(26), 772.

927   Lüdecke, D., Ben-Shachar, M. S., Patil, I., Waggoner, P., & Makowski, D. (2021). performance:
928   An R package for assessment, comparison and testing of statistical models. *Journal of
929   open source software*, *6*(60).

930   Maddox, R. K., & Lee, A. K. C. (2018). Auditory Brainstem Responses to Continuous Natural
931   Speech in Human Listeners. *eNeuro*, *5*(1). https://doi.org/10.1523/ENEURO.0441-
932   17.2018

933   MathWorks. (2019). *MATLAB*. In (Version 9.6.0 (R2019a)) The MathWorks, Inc.
934   https://www.mathworks.com

935   MathWorks. (2024). *MATLAB*. In (Version 24.2.0 (R2024b)) The MathWorks, Inc.
936   https://www.mathworks.com

937   Mattys, S. L., White, L., & Melhorn, J. F. (2005). Integration of multiple speech segmentation cues:
938   a hierarchical framework. *Journal of Experimental Psychology: General*, *134*(4), 477.

Mayo, L. H., Florentine, M., & Buus, S. (1997). Age of second-language acquisition and
perception of speech in noise. *J Speech Lang Hear Res*, *40*(3), 686–693.
https://doi.org/10.1044/jslhr.4003.686

McAuliffe, M., Socolof, M., Mihuc, S., Wagner, M., & Sonderegger, M. (2017). Montreal Forced
Aligner: Trainable Text-Speech Alignment Using Kaldi. Interspeech,

McDermott, J. H., & Simoncelli, E. P. (2011). Sound texture perception via statistics of the auditory
periphery: evidence from sound synthesis. *Neuron*, *71*(5), 926–940.
https://doi.org/10.1016/j.neuron.2011.06.032

Mesgarani, N., Cheung, C., Johnson, K., & Chang, E. F. (2014). Phonetic feature encoding in
human superior temporal gyrus. *Science*, *343*(6174), 1006–1010.
https://doi.org/10.1126/science.1245994

Mesik, J., Ray, L., & Wojtczak, M. (2021). Effects of Age on Cortical Tracking of Word-Level
Features of Continuous Competing Speech. *Front Neurosci*, *15*, 635126.
https://doi.org/10.3389/fnins.2021.635126

Muncke, J., Kuruvila, I., & Hoppe, U. (2022). Prediction of Speech Intelligibility by Means of EEG
Responses to Sentences in Noise. *Front Neurosci*, *16*, 876421.
https://doi.org/10.3389/fnins.2022.876421

Myers, B. R., Lense, M. D., & Gordon, R. L. (2019). Pushing the Envelope: Developments in
Neural Entrainment to Speech and the Biological Underpinnings of Prosody Perception.
Brain Sci, *9*(3). https://doi.org/10.3390/brainsci9030070

Nakagawa, S., Johnson, P. C. D., & Schielzeth, H. (2017). The coefficient of determination R(2)
and intra-class correlation coefficient from generalized linear mixed-effects models
revisited and expanded. *J R Soc Interface*, *14*(134). https://doi.org/10.1098/rsif.2017.0213

Nourski, K. V., Reale, R. A., Oya, H., Kawasaki, H., Kovach, C. K., Chen, H., Howard, M. A., &
Brugge, J. F. (2009). Temporal envelope of time-compressed speech represented in the
human auditory cortex. *J Neurosci*, *29*(49), 15564–15574.
https://doi.org/10.1523/JNEUROSCI.3065-09.2009

Oganian, Y., & Chang, E. F. (2019). A speech envelope landmark for syllable encoding in human
superior temporal gyrus. *Sci Adv*, *5*(11), eaay6279.
https://doi.org/10.1126/sciadv.aay6279

Orf, M., Wöstmann, M., Hannemann, R., & Obleser, J. (2022). Auditory neural tracking reflects
target enhancement but not distractor suppression in a psychophysically augmented

continuous-speech   paradigm.   *bioRxiv*,   2022.2006.2018.496558.
https://doi.org/10.1101/2022.06.18.496558

Pasley, B. N., David, S. V., Mesgarani, N., Flinker, A., Shamma, S. A., Crone, N. E., Knight, R.
T., & Chang, E. F. (2012). Reconstructing speech from human auditory cortex. *PLoS Biol*,
10(1), e1001251. https://doi.org/10.1371/journal.pbio.1001251

Peck, F. C., Gabard-Durnam, L. J., Wilkinson, C. L., Bosl, W., Tager-Flusberg, H., & Nelson, C.
A. (2021). Prediction of autism spectrum disorder diagnosis using nonlinear measures of
language-related EEG at 6 and 12 months. *Journal of neurodevelopmental disorders*,
13(1), 1–13.

Pedersen, T. L. (2019). Patchwork: The composer of plots. *CRAN: Contributed Packages*.

Pereira, F., Lou, B., Pritchett, B., Ritter, S., Gershman, S. J., Kanwisher, N., Botvinick, M., &
Fedorenko, E. (2018). Toward a universal decoder of linguistic meaning from brain
activation. *Nat Commun*, *9*(1), 963. https://doi.org/10.1038/s41467-018-03068-4

Pion-Tonachini, L., Kreutz-Delgado, K., & Makeig, S. (2019). ICLabel: An automated
electroencephalographic independent component classifier, dataset, and website.
Neuroimage, *198*, 181–197. https://doi.org/10.1016/j.neuroimage.2019.05.026

Rosen, S. (1992). Temporal information in speech: acoustic, auditory and linguistic aspects.
Philos   Trans   R   Soc   Lond   B   Biol   Sci,   *336*(1278),   367–373.
https://doi.org/10.1098/rstb.1992.0070

Salisbury, D. F., Shenton, M. E., Griggs, C. B., Bonner-Jackson, A., & McCarley, R. W. (2002).
Mismatch negativity in chronic schizophrenia and first-episode schizophrenia. *Arch Gen
Psychiatry*, *59*(8), 686–694. https://doi.org/10.1001/archpsyc.59.8.686

Schmitt, R., Meyer, M., & Giroud, N. (2022). Better speech-in-noise comprehension is associated
with enhanced neural speech tracking in older adults with hearing impairment. *Cortex*,
151, 133–146.

Shackleton, T. M., & Carlyon, R. P. (1994). The role of resolved and unresolved harmonics in
pitch perception and frequency modulation discrimination. *J Acoust Soc Am*, *95*(6), 3529–
3540. https://doi.org/10.1121/1.409970

Shannon, R. V., Zeng, F. G., Kamath, V., Wygonski, J., & Ekelid, M. (1995). Speech recognition
with   primarily   temporal   cues.   *Science*,   *270*(5234),   303–304.
https://doi.org/10.1126/science.270.5234.303

Smithson, M., & Verkuilen, J. (2006). A better lemon squeezer? Maximum-likelihood regression with beta-distributed dependent variables. *Psychol Methods*, *11*(1), 54–71. https://doi.org/10.1037/1082-989X.11.1.54

Sohoglu, E., & Davis, M. H. (2020). Rapid computations of spectrotemporal prediction error support perception of degraded speech. *Elife*, *9*. https://doi.org/10.7554/eLife.58077

Strauß, A., Wu, T., McQueen, J. M., Scharenborg, O., & Hintz, F. (2022). The differential roles of lexical and sublexical processing during spoken-word recognition in clear and in noise. *Cortex*, *151*, 70–88. https://doi.org/10.1016/j.cortex.2022.02.011

Swaminathan, J., & Heinz, M. G. (2012). Psychophysiological analyses demonstrate the importance of neural envelope coding for speech perception in noise. *J Neurosci*, *32*(5), 1747–1756. https://doi.org/10.1523/JNEUROSCI.4493-11.2012

Tang, C., Hamilton, L. S., & Chang, E. F. (2017). Intonational speech prosody encoding in the human auditory cortex. *Science*, *357*(6353), 797–801. https://doi.org/10.1126/science.aam8577

Teoh, E. S., Ahmed, F., & Lalor, E. C. (2022). Attention Differentially Affects Acoustic and Phonetic Feature Encoding in a Multispeaker Environment. *J Neurosci*, *42*(4), 682–691. https://doi.org/10.1523/JNEUROSCI.1455-20.2021

Teoh, E. S., Cappelloni, M. S., & Lalor, E. C. (2019). Prosodic pitch processing is represented in delta-band EEG and is dissociable from the cortical tracking of other acoustic and phonetic features. *Eur J Neurosci*, *50*(11), 3831–3842. https://doi.org/10.1111/ejn.14510

Tezcan, F., Weissbart, H., & Martin, A. E. (2023). A tradeoff between acoustic and linguistic feature encoding in spoken language comprehension. *Elife*, *12*, e82386.

Turner, C. W., Gantz, B. J., Vidal, C., Behrens, A., & Henry, B. A. (2004). Speech recognition in noise for cochlear implant listeners: benefits of residual acoustic hearing. *J Acoust Soc Am*, *115*(4), 1729–1735. https://doi.org/10.1121/1.1687425

Vanthornhout, J., Decruy, L., Wouters, J., Simon, J. Z., & Francart, T. (2018). Speech Intelligibility Predicted from Neural Entrainment of the Speech Envelope. *J Assoc Res Otolaryngol*, *19*(2), 181–191. https://doi.org/10.1007/s10162-018-0654-z

Verschueren, E., Gillis, M., Decruy, L., Vanthornhout, J., & Francart, T. (2022). Speech understanding oppositely affects acoustic and linguistic neural tracking in a speech rate manipulation paradigm. *Journal of Neuroscience*, *42*(39), 7442–7453.

Verschueren, E., Vanthornhout, J., & Francart, T. (2021). The effect of stimulus intensity on neural
envelope tracking. *Hear Res*, *403*, 108175. https://doi.org/10.1016/j.heares.2021.108175

Viswanathan, V., Bharadwaj, H. M., Shinn-Cunningham, B. G., & Heinz, M. G. (2021). Modulation
masking and fine structure shape neural envelope coding to predict speech intelligibility
across diverse listening conditions. *J Acoust Soc Am*, *150*(3), 2230.
https://doi.org/10.1121/10.0006385

Wickham, H. (2011). ggplot2. *Wiley interdisciplinary reviews: computational statistics*, *3*(2), 180–
185.

Yasmin, S., Irsik, V. C., Johnsrude, I. S., & Herrmann, B. (2023). The effects of speech masking
on neural tracking of acoustic and semantic features of natural speech. *Neuropsychologia*,
186, 108584.

Zhang, X., Li, J., Li, Z., Hong, B., Diao, T., Ma, X., Nolte, G., Engel, A. K., & Zhang, D. (2023).
Leading and following: Noise differently affects semantic and acoustic processing during
naturalistic speech comprehension. *Neuroimage*, *282*, 120404.

Zou, J., Feng, J., Xu, T., Jin, P., Luo, C., Zhang, J., Pan, X., Chen, F., Zheng, J., & Ding, N.
(2019). Auditory and language contributions to neural encoding of speech features in noisy
environments. *Neuroimage*, *192*, 66–75.
https://doi.org/10.1016/j.neuroimage.2019.02.047

1051

1052   **SUPPLEMENTARY INFORMATION**

1053   **S1 Table**. Wilcoxon rank sum p-values from behavioral score comparisons between conditions.

% Words Heard				
	Quiet	+3 dB	-3 dB	-6 dB
+3 dB	3.180×10^{-4}	-	-	-
-3 dB	1.294×10^{-4}	1.294×10^{-4}	-	-
-6 dB	1.294×10^{-4}	1.294×10^{-4}	1.294×10^{-4}	-
-9 dB	1.294×10^{-4}	1.294×10^{-4}	1.294×10^{-4}	1.294×10^{-4}
Comprehension Scores				
	Quiet	+3 dB	-3 dB	-6 dB
+3 dB	5.991×10^{-1}	-	-	-
-3 dB	1.899×10^{-4}	4.904×10^{-4}	-	-
-6 dB	1.259×10^{-4}	1.259×10^{-4}	1.411×10^{-4}	-
-9 dB	1.259×10^{-4}	1.259×10^{-4}	1.259×10^{-4}	2.244×10^{-4}

1054

1055

1056

1057   **S1 Figure.** Spectrogram (Sgram, left) and Phonetic Feature (Fea, right) TRF model weights at each signal

1058   to noise ratio. The model weights were averaged across time (0 to 200 ms) and the 12 frontotemporal

1059   electrodes depicted in Figure 1. Frequency bands and feature spaces are plotted on the x-axes and model

1060   weights (in arbitrary units) are on the y-axes.

1061

1062

1063 **S2 Table.** Results from GLMMs that related partial correlation coefficients to behavioral scores

Percent words heard score GLMM					
Fixed Effect	Estimate	Std. Error	t-value	p-value	
Intercept	1. 609	0.138	11.635	$< 2 \times 10^{-16}$	***
Ons_A	-0.129	0.086	-1.497	0.134	
Sgram	0.117	0.088	1.322	0.186	
Fea	0.104	0.102	1.022	0.307	
Surp	0.398	0.097	4.128	3.652×10^{-5}	***
Ons_W	0.055	0.070	0.779	0.436	
SNR	-2.133	0.104	-20.454	$< 2 \times 10^{-16}$	***
SNR:Ons_A	0.258	0.092	2.819	0.005	**
SNR:Sgram	-0.038	0.073	-0.528	0.597	
SNR:Fea	-0.098	0.111	-0.877	0.381	
SNR:Surp	-0.265	0.098	-2.694	0.007	**
SNR:Ons_W	-0.083	0.071	-1.165	0.243	
Comprehension score GLMM					
Fixed Effect	Estimate	Std. Error	t-value	p-value	
Intercept	1.046	0.111	9.402	$< 2 \times 10^{-16}$	***
Ons_A	-0.015	0.062	-0.244	0.807	
Sgram	0.128	0.063	2.031	0.042	*
Fea	0.055	0.060	0.919	0.358	
Surp	0.122	0.061	1.996	0.046	*
Ons_W	0.044	0.051	0.871	0.384	
SNR	-0.830	0.070	-11.932	$< 2 \times 10^{-16}$	***
SNR:Ons_A	0.033	0.055	0.598	0.550	
SNR:Sgram	-0.051	0.049	-1.022	0.307	
SNR:Fea	0.124	0.058	2.137	0.033	*
SNR:Surp	0.024	0.053	0.462	0.644	
SNR:Ons_W	0.038	0.050	-0.777	0.437	

1064

1065

1066

S3 Table. Post hoc contrasts for the LME model that calculated the relationship between lexical surprisal

1068   and envelope tracking in each SNR.

Interaction variable statistics					
SNR	Surp:SNR Slope	Std. Error	z-ratio	p-value	
Quiet	0.029	2.709×10^{-3}	10.519	< 0.001	***
+3 dB	0.028	2.701×10^{-3}	10.336	< 0.001	***
-3 dB	0.024	2.730×10^{-3}	8.787	< 0.001	***
-6 dB	0.019	2.709×10^{-3}	6.814	< 0.001	***
-9 dB	0.013	2.736×10^{-3}	4.615	< 0.001	***
Comparisons between interaction variables					
Contrast	Estimate	Std. Error	z ratio	p-value	
Quiet – + 3 dB	2.17×10^{-4}	3.53×10^{-3}	0.062	1.000	
Quiet – -3 dB	4.767×10^{-3}	3.50×10^{-3}	1.363	0.651	
Quiet – -6 dB	9.900×10^{-3}	3.51×10^{-3}	2.817	0.039	*
Quiet – -9 dB	1.600×10^{-2}	3.50×10^{-3}	4.570	< 0.001	***
-3 dB – + 3 dB	-4.459×10^{-3}	3.53×10^{-3}	-1.290	0.698	
-3 dB – -6 dB	5.133×10^{-3}	3.52×10^{-3}	1.458	0.590	
-3 dB – -9 dB	1.123×10^{-2}	3.50×10^{-3}	3.211	0.012	*
-6 dB – + 3 dB	-9.683×10^{-3}	3.53×10^{-3}	-2.732	0.049	*
-6 dB – -9 dB	6.097×10^{-3}	3.54×10^{-3}	1.729	0.416	
-9 dB – + 3 dB	-1.578×10^{-2}	3.53×10^{-3}	-4.468	< 0.001	***

1069
1070