Inferring interactions from microbiome data

Román Zapién-Campos[1], Florence Bansept[2], and Arne Traulsen[1*]

[1]Max Planck Institute for Evolutionary Biology, Plön, Germany

[2]Aix Marseille Université, CNRS, Laboratoire de Chimie Bactérienne (UMR7283), IMM, IM2B, Turing Center for Living Systems, Marseille, France

*correspondence: traulsen@evolbio.mpg.de

March 30, 2023

Parameter inference of high-dimensional data is challenging and microbiome time series data is no exception. Methods aimed at predicting from point estimates exist, but often even fail to recover the true parameters from simulated data. Computational methods to robustly infer and quantify the uncertainty in model parameters are needed. Here, we propose a computational workflow addressing such challenges – allowing us to compare mechanistic models and identify the values and the certainty of inferred parameters. This approach allows us to infer which kind of interactions occur in the microbial community. In contrast to point-estimate inference, the distribution for the parameters, our outcome, reflects their uncertainty. To achieve this, we consider as many equations for the statistical moments of the microbiome as parameters. Our inference workflow, which builds upon a mechanistic foundation of microscopic processes,

can take into account that commonly metagenomic datasets only provide information on

relative abundances and hosts' ensembles. With our framework, we move from qualita-

tive prediction to quantifying the likelihood of certain interaction types in microbiomes.

Keywords: microbiome, interactions, Bayesian inference, parameters, dynamical equations.

Numerous studies have shown how important the microbiome is for their hosts, ranging from de-

velopment to health [1, 2]. The promise of manipulating the microbiome relies on having understood

the ecological and evolutionary processes operating on it [3]. Although metagenomics studies have

widely characterized microbiome samples [4], their connection to mathematical models and eco-

evolutionary theories lags behind. Part of the gap is explained by an intrinsic difficulty in analysing

microbiome data [5], in particular the inverse problem of robustly inferring parameters – and thus

interactions between microbes – from data (Fig.1 A). Pioneering studies have achieved point ("best

guess") estimates that allow accurate predictions [6], but fall short in estimating the true parameters

from simulations [5]. This apparent contradiction has been discussed by Cao et al. [5]. Others have

suggested to use Bayesian inference – where probabilities are assigned to parameter values – to go

beyond point-estimates [7]. Here, we present a computational workflow that starts from defining

microscopic transition rates – describing ecological and evolutionary events – and works with macro-

scopic statistical moments of the microbiome composition. Our Bayesian inference workflow, which

naturally bypasses known limitations of point-estimate inference [5], is sufficiently flexible to test

models as diverse as microbiome samples while quantifying the parameter uncertainty stemming

from data limitations (Fig.1 C) – including its extrinsic noise. Two classical ecological models are

used to illustrate its application on datasets describing absolute or relative abundances of microbes

– namely, logistic growth and Lotka-Volterra models. The inference workflow outlined here bridges

a gap between microbiome data and theoretical modelling.

Developing an inference workflow

46 We propose a parameter inference workflow grounded on a mechanistic description of the dynamics

47 of a microbiome (Fig.1 C). First, we write down microscopic transition rates T describing changes

48 in the microbiome composition of one host, the vector $\mathbf{n}$, given the parameter set $\boldsymbol{\theta}$. Note that

Figure 1: **Parameter inference workflow proposed and microbiome data properties**. (**A**) Models serve as a link between parameters and data. Either to simulate biological processes or to infer parameters from data. (**B**) Longitudinal sampling of the same hosts or an ensemble of them are used to obtain datasets. (**C**) Workflow from microscopic rates of a model and experimental data to inference of parameters by Approximate Bayesian Computation. The microscopic rates describe possible eco-evolutionary events, leading to macroscopic patterns (statistical moments). Datasets describe absolute abundances (counts) or relative abundances (frequencies) of microbes. To quantify the probability of parameter values in the model, prior knowledge is updated to a posterior distribution based on the agreement of the model with the data.

49   these rates can be functions of time t as well,

$$T(\mathbf{n} \to \mathbf{n}', t) = f(\mathbf{n}, \boldsymbol{\theta}, t). \tag{1}$$

50   Now, instead of tracking the microbiome composition $\mathbf{n}$ in a single host, we can describe how the
51   probability of observing a microbiome composition $\mathbf{n}$ in an ensemble of hosts, $P(\mathbf{n}, t)$, changes with
52   time,

$$\frac{\partial P(\mathbf{n}, t)}{\partial t} = \underbrace{\sum_{\mathbf{n}'} P(\mathbf{n}', t) T(\mathbf{n}' \to \mathbf{n}, t)}_{\text{probability influx to } \mathbf{n}} - \underbrace{\sum_{\mathbf{n}'} P(\mathbf{n}, t) T(\mathbf{n} \to \mathbf{n}', t)}_{\text{probability outflux from } \mathbf{n}}. \tag{2}$$

In this expression, called the master equation [8], the influx and outflux terms indicate an increase

or decrease in the probability of composition $\mathbf{n}$. This dynamics depends on the ecological and

evolutionary processes contained in the microscopic rates.

Using the master equation, we can derive equations for the statistical moments of the microbiome

composition in an ensemble of hosts – namely, the product of the master equation by a variable of

interest (g_k, where k is the focal microbial type) summed over all possible microbiome compositions,

$$\frac{d\langle g_k \rangle}{dt} = \sum_{\mathbf{n}} g_k \frac{\partial P(\mathbf{n}, t)}{\partial t}. \tag{3}$$

For example, computing the average abundance of microbial type i implies setting $g_k = n_i$, whereas

for the co-moment of microbial types i and j, $g_k = n_i n_j$. To obtain a system of equations of these

moments that fully determines the parameters, we have to derive at least as many equations for them

as parameters in the model. For example, in a Lotka-Volterra model with S microbial types, there

are S growth rates and S^2 intra- and inter-specific interactions, amounting to $S + S^2$ parameters.

Thus, we would need $S + S^2$ equations. We can use the S equations for the first moments $\langle n_k \rangle$,

S for the second moments $\langle n_k^2 \rangle$, and $S(S-1)$ for the co-moments $\langle n_k n_l \rangle$ and covariances $\langle n_k, n_l \rangle$

(see Sup. Methods A). Note that each equation can depend on the vector of other moments, i.e.,

$\langle g_k \rangle = f(\langle \mathbf{g} \rangle, \boldsymbol{\theta}, t)$. A step by step derivation from microscopic rates up to second order moments

for a logistic growth and the Lotka-Volterra models can be found in the Sup. Methods A. These

models include conventional ecological events, such as growth, death, immigration, and direct and

indirect interactions.

We now have the elements to infer the parameters $\boldsymbol{\theta}$ from data. The idea of Approximate Bayesian

Computation (ABC) is to identify feasible parameters by comparing the data to model predictions

[7]. Specifically, for a given set $\boldsymbol{\theta}$, a distance metric between the numerical solution of the equations

for the moments, $\langle g_k \rangle$, and the equivalent moments from data, $\bar{g}_k$, is estimated, e.g.,

$$\sum_k \sum_i |\langle g_k \rangle(\boldsymbol{\theta}, t_i) - \bar{g}_k(t_i)|, \tag{4}$$

for the Euclidean distance. If this distance is smaller than a threshold ε, the set is considered valid.

By testing sets sampled according to an expectation – prior distribution – and recording those below

the threshold ε, a posterior distribution of the parameters reflecting the uncertainty of the inference

can be obtained (Fig. 1 C). With a smaller threshold ε this posterior can become the new prior

and the process can be iterated to improve it. This is called Approximate Bayesian Computation -

Sequential Monte Carlo (ABC-SMC). We discuss how to choose prior distributions of the parameters

in the Sup. Methods C.

Properties of microbiome data

Given a microbiome dataset, all statistical moments $\bar{g}_k$ can be estimated from raw data. Concretely,

it can be done by averaging the variable of interest g_k, over all replicates in each specific time point

(Fig. 1 C).

Microbiome data is nowadays typically produced by metagenome sequencing. Conventionally, for

technical reasons metagenomics only quantifies the *relative abundance* of each microbial type in a

sample (Fig. 1 C) [9]. More recently, some studies have measured absolute numbers of culturable

[10] and non-culturable microbes in samples [11]. We call these counts *absolute abundances*.

Our former equations only track moments of absolute abundance, $\langle g_k \rangle$. As Gloor et al. [9]

show, inferring parameters from relative abundance (x_k) data using them would lead to spurious

correlations (Fig. 2 A-B). To find equivalent expressions for the relative abundance, first, we propose

$n_\Sigma \equiv \sum_j n_j$ to be defined as a scaling factor and an equation for its first moment, $\langle n_\Sigma \rangle$. Then, a

transformation to moments of relative abundances, $\langle \gamma_k \rangle$, is given by,

$$\frac{d\langle \gamma_k \rangle}{dt} = f(\langle \boldsymbol{\gamma} \rangle, \langle n_\Sigma \rangle, \boldsymbol{\theta}, t). \tag{5}$$

Because relative abundance datasets lack information about the scaling factor, its initial condition,

$\bar{n}_\Sigma(0)$, must be inferred as a free parameter, one parameter more than for absolute abundance data.

Note that because $\sum_k x_k = 1$, the number of equations for the types decreases but the number of

parameters per type remains. A detailed derivation of transformations for a logistic growth and the

Lotka-Volterra models is shown in the Sup. Methods A.

Apart from tracking the microbiome of the same hosts over time, as in animal gut studies,

commonly, various hosts are pulled together to produce a single time series (Fig. 1 B). This is the

case when hosts are sacrificed while sampling as in experimental studies of *D. melanogaster*, *C.

elegans*, and Hydra [10]. In contrast to deterministic models, our workflow can deal with hosts

pulled together thanks to its account of the stochastic demography. More concretely – akin to the

concept of biological replicates – if the parameter values and initial conditions are the same in each

host sampled, we can account for their emerging demographic differences.

Figure 2: **Inferring true parameters from simulated data**. (**A**-**B**) Time series comparison between simulations (dots, derived from only 4 replicates) and equations for the statistical moments (lines) of absolute (n_k) and relative abundance (x_k). Two models with three microbial types ($S = 3$) were tested, (**A**) logistic growth with immigration and death ($3S + 1 = 9$ parameters) and (**B**) Lotka-Volterra ($S + S^2 = 12$ parameters). Inferred parameter posteriors are compared to the true parameters (dashed lines, Sup. Methods B) and priors (black distributions, Sup. Method C). All microbial types shared the same priors. (**C**) The inferred interactions for the Lotka-Volterra model resembled the true interactions, qualitatively (arrow heads) and quantitatively (arrow thickness), with various certainties (grayscale, defined by the coefficient of variation of the posterior). (**D**) For both simulated datasets, the most probable model was identified correctly (a.u. = arbitrary time units are determined by the rates in the model, see Appendix B).

Inference from simulated and empirical data

We tested our inference workflow in two ways. Firstly, we recovered the correct parameters from simulated data (Fig. 2). Our approach proved successful in cases with and without inter-specific interactions, namely, data simulated from logistic growth and Lotka-Volterra models. In both cases, our approach identified the correct model while in addition estimating their parameters and certainty.

Secondly, we inferred parameters of an empirical time series of absolute abundance in a reduced mice microbiome – OMM[12] [12] (Fig. 3). The posteriors suggested the growth rates of *Bacteroides caecimuris*, *Muribaculum intestinale*, *Flavonifactor plautii*, *Akkermansia muciniphila*, *Turicimonas muris* to be most certain, with average doubling times ranging from hours to days. Meanwhile, average death and immigration rates were less certain, ranging from ≈ 4 to 11.5 cells per second. Naturally, the certainties obtained from empirical data (Fig. 3) are smaller than those from simulations (Fig. 2), highlighting the limits of the model tested and inference from, noisy, empirical data. However, in each case we obtain a set of parameters – capturing interactions between microbes – with some level of certainty. To exemplify the utility of our outcome, here, our results point to selection as the ecological driver of the OMM[12] dynamics, despite a possible compatibility of this data with a neutral hypothesis once it has reached steady state [13]. In this case, neutrality would imply that the parameter posteriors overlap between microbial types, which is not the case (Fig. 3).

Discussion

The work presented here is motivated by the goal to understand how the microbes in a microbiome interact and the need to quantify the uncertainty of parameter estimation from microbiome data. Although inference methods for point estimation have been used [6], several issues limit their quantitative application, restricting them to recreate qualitative patterns of data [5]. A major issue is the indeterminacy of models with more parameters than equations [6, 5, 14]. We suggest a solution to this issue by deriving equations for the statistical moments of the microbiome composition – at least as many as parameters. In fact, our approach is driven by a mechanistic spirit, where microscopic rates must be written down first. As opposed to approaches where analytic solutions – or expensive stochastic simulations – are needed, here, a numerical solution is sufficient to quantify the distance

Figure 3: **Inferring parameters from empirical data**. The parameters of a logistic growth with immigration and death model were inferred from a mouse dataset. The Oligo-Mouse-Microbiota (OMM[12]) dataset [12] tracks a 12-species defined mice microbiome ($S = 12$), where the absolute abundances of the same individuals were sampled 11 times over 99 days. (**A**) We analysed the first 21 days where 4 replicates are available throughout and assumed logistic growth with parameters for growth, death and immigration. (**B**) Of the $3S + 1 = 37$ parameters inferred, we show only the posteriors of the five most certain ones (defined by the coefficient of variation of the posterior). All microbial types shared the same uniform priors (black distributions, Sup. Methods C). (**C**) The parameters inferred for each taxa varied widely with various certainties. For the shared carrying capacity we found an average $N \approx 1.47 \cdot 10^7$ bacterial cells, with an uncertainty of $\pm 1.33 \cdot 10^5$ cells (CV ≈ 0.0089).

to data [15]. This allows our workflow to handle very diverse models, where model comparison [15]

in a Bayesian sense is possible.

The workflow is not limited by the properties of microbiome data [9]. As we have shown, analyzing

datasets describing the relative abundance of microbial types – even if the total absolute abundance

is dynamic [14] – is possible. Such is the nature of metagenomic sequencing data – the most

common method to characterize microbiomes [5]. In addition, by tracking statistical moments of the

microbiome, our approach naturally accounts for the diverse types of experimental samplings, such

as those where ensembles of hosts are used to obtain a single time series. Concretely, compared to

other methods, we track the demographic variation between hosts explicitly, and assign the remaining

variation to external noise.

Although by design, our approach deals with longitudinal (time series) data, analyzing single time

points (snapshot data) is possible. For example, if data is assumed to be at steady state, the inference

method's aim is to find parameters making the dynamical equations for the moments equal to zero.

This does not mean that the moments are zero, but that their rate of change is. Nevertheless, as our

results illustrate, given the various sources of uncertainty, time series data leads to better parameter

inference, in particular, those time intervals of "high activity" where many changes occur [5]. As

Cao et al. [5] proposed, several of these intervals could be analysed simultaneously to improve the

inference.

Bayesian inference can suffer the curse-of-dimensionality in large and diverse systems [15]. By

tracking numerous statistical moments readily solved numerically, we believe our approach combined

with data of sufficiently high quality can overcome this to some extent – exploring the parameter

space thoroughly in a reasonable time. We implemented an Approximate Bayesian Computation

with Sequential Monte Carlo in our workflow using tools from the Python package pyABC [16]

(Sup. Methods C), on which further optimizations could greatly improve its wider application

[7]. As proof-of-principle, we applied our workflow to a simulated dataset and recovered the true

parameter values. We also applied it to a reduced microbiome in mice [12], where we estimated

values and certainties of parameters describing logistic growth.

In summary, we presented a Bayesian inference workflow bridging microbiome data to theoretical

modelling. By inferring datasets of microbial absolute and relative abundances, we showed its

robustness – identifying likely interactions and certainty of parameters in simulated and empirical

data. Because mechanistic rates serve as stepping stones of the workflow, similar microscopic models

could replace the two classical ecological models that we illustrated – including experimentally
informed models.

Acknowledgements

169 We thank the *Evolutionary Theory Department* in the MPI Plön and the *Collaborative Research*
170 *Centre 1182: Origins and Functions of Metaorganisms* for the fruitful discussions. Finally, we thank
171 the Max Planck Society and the CRC 1182 for the funding provided.

Author contributions

173 RZC and AT developed the concept. RZC and FB worked on the methods. RZC wrote the first
174 draft; all authors reviewed and approved the final version.

Availability of data and software

176 The data generated and software used for the analyses are available in GitHub
177 (https://github.com/romanzapien/microbiome-inference.git).

References

[1] Daniela Esser, Janina Lange, Georgios Marinos, Michael Sieber, Lena Best, Daniela Prasse,
Jay Bathia, Malte C Rühlemann, Kathrin Boersch, Cornelia Jaspers, et al. Functions of the
microbiota for the physiology of animal metaorganisms. *Journal of innate immunity*, 11(5):
393–404, 2019.

[2] Jack A Gilbert, Martin J Blaser, J Gregory Caporaso, Janet K Jansson, Susan V Lynch, and Rob
Knight. Current understanding of the human microbiome. *Nature medicine*, 24(4):392–400,
2018.

[3] Michael A Fischbach. Microbiome: focus on causation and mechanism. *Cell*, 174(4):785–790,
2018.

[4] Luke R Thompson, Jon G Sanders, Daniel McDonald, Amnon Amir, Joshua Ladau, Kenneth J
Locey, Robert J Prill, Anupriya Tripathi, Sean M Gibbons, Gail Ackermann, et al. A communal
catalogue reveals earth's multiscale microbial diversity. *Nature*, 551(7681):457–463, 2017.

[5] Hong-Tai Cao, Travis E Gibson, Amir Bashan, and Yang-Yu Liu. Inferring human microbial
dynamics from temporal metagenomics data: Pitfalls and lessons. *BioEssays*, 39(2):1600188,
2017.

[6] Richard R Stein, Vanni Bucci, Nora C Toussaint, Charlie G Buffie, Gunnar Rätsch, Eric G
Pamer, Chris Sander, and Joao B Xavier. Ecological modeling from time-series inference:
insight into dynamics and stability of intestinal microbiota. *PLoS computational biology*, 9
(12):e1003388, 2013.

[7] Tina Toni, David Welch, Natalja Strelkowa, Andreas Ipsen, and Michael PH Stumpf. Approxi-
mate bayesian computation scheme for parameter inference and model selection in dynamical
systems. *Journal of the Royal Society Interface*, 6(31):187–202, 2009.

[8] C. W. Gardiner. *Handbook of Stochastic Methods*. Springer, NY, Berlin, third edition, 2004.

[9] Gregory B Gloor, Jean M Macklaim, Vera Pawlowsky-Glahn, and Juan J Egozcue. Microbiome
datasets are compositional: and this is not optional. *Frontiers in microbiology*, 8:2224, 2017.

[10] Tanita Wein, Tal Dagan, Sebastian Fraune, Thomas CG Bosch, Thorsten BH Reusch, and
Nils F Hülter. Carrying capacity and colonization dynamics of curvibacter in the hydra host
habitat. *Frontiers in microbiology*, 9:443, 2018.

[11] Gianluca Galazzo, Niels Van Best, Birke J Benedikter, Kevin Janssen, Liene Bervoets, Christel
Driessen, Melissa Oomen, Mayk Lucchesi, Pascalle H van Eijck, Heike EF Becker, et al. How
to count our microbes? the effect of different quantitative microbiome profiling approaches.
Frontiers in cellular and infection microbiology, 10:403, 2020.

[12] Claudia Eberl, Diana Ring, Philipp C Münch, Markus Beutler, Marijana Basic, Emma Caroline
Slack, Martin Schwarzer, Dagmar Srutkova, Anna Lange, Julia S Frick, et al. Reproducible
colonization of germ-free mice with the oligo-mouse-microbiota in different animal facilities.
Frontiers in microbiology, 10:2999, 2020.

[13] M. Sieber, L. Pita, N. Weiland-Brauer, P. Dirksen, J. Wang, B. Mortzfeld, S. Franzenburg, R. A. Schmitz, J.F. Baines, S. Fraune, U. Hentschel, H. Schulenburg, T. C. G. Bosch, and A. Traulsen. Neutrality in the metaorganism. *PLoS Biology*, 17(6):e3000298, 2019.

[14] Tyler A Joseph, Liat Shenhav, Joao B Xavier, Eran Halperin, and Itsik Pe'er. Compositional lotka-volterra describes microbial dynamics in the simplex. *PLoS computational biology*, 16(5): e1007917, 2020.

[15] Mikael Sunnåker, Alberto Giovanni Busetto, Elina Numminen, Jukka Corander, Matthieu Foll, and Christophe Dessimoz. Approximate bayesian computation. *PLoS computational biology*, 9 (1):e1002803, 2013.

[16] Yannik Schälte, Emmanuel Klinger, Emad Alamoudi, and Jan Hasenauer. pyabc: Efficient and robust easy-to-use approximate bayesian computation. *Journal of Open Source Software*, 7 (74):4304, 2022.

[17] Omri Allouche and Ronen Kadmon. A general framework for neutral models of community dynamics. *Ecology letters*, 12(12):1287–1297, 2009.

[18] Christian Kuehn. Moment closure – a brief review. *Control of self-organizing nonlinear systems*, pages 253–271, 2016.

[19] Beth Gibson, Daniel J Wilson, Edward Feil, and Adam Eyre-Walker. The distribution of bacterial doubling times in the wild. *Proceedings of the Royal Society B: Biological Sciences*, 285(1880): 20180789, 2018.

Appendix

A. Derivation of dynamical equations for the microbiome moments

The change in a community is described by the master equation,

$$\frac{\partial P(\mathbf{n}, t)}{\partial t} = \underbrace{\sum_i P(\mathbf{n} + \mathbf{e}_i, t)T(\mathbf{n} + \mathbf{e}_i \to \mathbf{n})}_{\text{influx from states } n_i + 1} + \underbrace{\sum_i P(\mathbf{n} - \mathbf{e}_i, t)T(\mathbf{n} - \mathbf{e}_i \to \mathbf{n})}_{\text{influx from states } n_i - 1}$$
$$- \underbrace{\sum_i P(\mathbf{n}, t)T(\mathbf{n} \to \mathbf{n} + \mathbf{e}_i)}_{\text{outflux to states } n_i + 1} - \underbrace{\sum_i P(\mathbf{n}, t)T(\mathbf{n} \to \mathbf{n} - \mathbf{e}_i)}_{\text{outflux to states } n_i - 1}, \tag{6}$$

where $\mathbf{n}$ is the vector of absolute abundances, and $\mathbf{e}_i$ is a vector with one in the *i-th* entry and zero otherwise.

Dynamical equations for the statistical moments can be obtained from the master equation by multiplication and subsequent summation. E.g., for the first moment, we have

$$\frac{d\langle n_k \rangle}{dt} = \sum_{\mathbf{n}} n_k \frac{\partial P(\mathbf{n}, t)}{\partial t} = \dots \sum_{n_k=0}^{\infty} \dots n_k \frac{\partial P(\mathbf{n}, t)}{\partial t}, \tag{7}$$

where for convenience, we make summations more explicit. For the second moment, we have

$$\frac{d\langle n_k^2 \rangle}{dt} = \sum_{\mathbf{n}} n_k^2 \frac{\partial P(\mathbf{n}, t)}{\partial t} = \dots \sum_{n_k=0}^{\infty} \dots n_k^2 \frac{\partial P(\mathbf{n}, t)}{\partial t}, \tag{8}$$

and for the co-moments,

$$\frac{d\langle n_k n_l \rangle}{dt} = \sum_{\mathbf{n}} n_k n_l \frac{\partial P(\mathbf{n}, t)}{\partial t} = \dots \sum_{n_k=0}^{\infty} \sum_{n_l=0}^{\infty} \dots n_k n_l \frac{\partial P(\mathbf{n}, t)}{\partial t}. \tag{9}$$

The upper sum limit is changed to a finite number for models with a finite carrying capacity.

A.1. Logistic growth with immigration and death

Similarly to Allouche and Kadmon [17], let us define the microscopic transition rates,

$$T(\mathbf{n} \to \mathbf{n} + \mathbf{e}_i) = (f_i n_i + m_i) \frac{N - \sum_j n_j}{N} \tag{10a}$$

$$T(\mathbf{n} \to \mathbf{n} - \mathbf{e}_i) = \phi_i \frac{n_i}{N}, \tag{10b}$$

where N is the shared carrying capacity, f_i is the maximum growth rate, and ϕ_i and m_i are the death and immigration rates for each type i.

Now, we illustrate how to derive dynamical equations for the moments. Let us start with the first moment,

$$
\begin{aligned}
\frac{d\langle n_k \rangle}{dt} = \ldots \sum_{n_k=0}^{N-1} \ldots & n_k P(\mathbf{n}+\mathbf{e}_k,t) T(\mathbf{n}+\mathbf{e}_k \to \mathbf{n}) \\
+ \ldots \sum_{n_k=1}^{N} \ldots & n_k P(\mathbf{n}-\mathbf{e}_k,t) T(\mathbf{n}-\mathbf{e}_k \to \mathbf{n}) \\
- \ldots \sum_{n_k=0}^{N} \ldots & n_k P(\mathbf{n},t) T(\mathbf{n} \to \mathbf{n}+\mathbf{e}_k) \\
- \ldots \sum_{n_k=0}^{N} \ldots & n_k P(\mathbf{n},t) T(\mathbf{n} \to \mathbf{n}-\mathbf{e}_k) \\
+ \ldots \sum_{n_k=0}^{N}\sum_{n_i=0}^{N-1} \ldots & n_k \sum_{i \neq k} P(\mathbf{n}+\mathbf{e}_i,t) T(\mathbf{n}+\mathbf{e}_i \to \mathbf{n}) \\
+ \ldots \sum_{n_k=0}^{N}\sum_{n_i=1}^{N} \ldots & n_k \sum_{i \neq k} P(\mathbf{n}-\mathbf{e}_i,t) T(\mathbf{n}-\mathbf{e}_i \to \mathbf{n}) \\
- \ldots \sum_{n_k=0}^{N}\sum_{n_i=0}^{N} \ldots & n_k \sum_{i \neq k} P(\mathbf{n},t) T(\mathbf{n} \to \mathbf{n}+\mathbf{e}_i) \\
- \ldots \sum_{n_k=0}^{N}\sum_{n_i=0}^{N} \ldots & n_k \sum_{i \neq k} P(\mathbf{n},t) T(\mathbf{n} \to \mathbf{n}-\mathbf{e}_i),
\end{aligned}
\tag{11}
$$

where the first four lines describe birth or death of a microbe of type k and the last four lines describe birth or death of a microbe of type i. Note that by definition $P(\mathbf{n}+\mathbf{e}_i,t)|_{n_i=N} = 0$ and $P(\mathbf{n}-\mathbf{e}_i,t)|_{n_i=0} = 0$, so their summation indices go up to $n_i = N-1$, or start from $n_i = 1$, respectively.

255 After the appropriate transformations of variables and re-indexing, we obtain

$$
\begin{aligned}
\frac{d\langle n_k \rangle}{dt} = {} & \ldots \sum_{n_k=0}^{N} \ldots (n_k - 1) P(\mathbf{n}, t) T(\mathbf{n} \to \mathbf{n} - \mathbf{e}_k) \\
& + \ldots \sum_{n_k=0}^{N} \ldots (n_k + 1) P(\mathbf{n}, t) T(\mathbf{n} \to \mathbf{n} + \mathbf{e}_k) \\
& - \ldots \sum_{n_k=0}^{N} \ldots n_k P(\mathbf{n}, t) T(\mathbf{n} \to \mathbf{n} + \mathbf{e}_k) \\
& - \ldots \sum_{n_k=0}^{N} \ldots n_k P(\mathbf{n}, t) T(\mathbf{n} \to \mathbf{n} - \mathbf{e}_k) \\
& + \ldots \sum_{n_k=0}^{N} \sum_{n_i=0}^{N} \ldots n_k \sum_{i \neq k} P(\mathbf{n}, t) T(\mathbf{n} \to \mathbf{n} - \mathbf{e}_i) \\
& + \ldots \sum_{n_k=0}^{N} \sum_{n_i=0}^{N} \ldots n_k \sum_{i \neq k} P(\mathbf{n}, t) T(\mathbf{n} \to \mathbf{n} + \mathbf{e}_i) \\
& - \ldots \sum_{n_k=0}^{N} \sum_{n_i=0}^{N} \ldots n_k \sum_{i \neq k} P(\mathbf{n}, t) T(\mathbf{n} \to \mathbf{n} + \mathbf{e}_i) \\
& - \ldots \sum_{n_k=0}^{N} \sum_{n_i=0}^{N} \ldots n_k \sum_{i \neq k} P(\mathbf{n}, t) T(\mathbf{n} \to \mathbf{n} - \mathbf{e}_i),
\end{aligned}
\tag{12}
$$

256 Note that the last four terms reduce to zero, and that $T(\mathbf{n} \to \mathbf{n} - \mathbf{e}_k)|_{n_k=0} = 0$ and $T(\mathbf{n} \to$
257 $\mathbf{n} + \mathbf{e}_k)|_{n_k=N} = 0$, which allows including $n_k = 0$ and $n_k = N$ in the summations. Simplifying, we
258 find

$$
\frac{d\langle n_k \rangle}{dt} = \ldots \sum_{n_k=0}^{N} \ldots P(\mathbf{n}, t) \left(T(\mathbf{n} \to \mathbf{n} + \mathbf{e}_k) - T(\mathbf{n} \to \mathbf{n} - \mathbf{e}_k) \right),
\tag{13}
$$

259 and substituting $T(\mathbf{n} \to \mathbf{n} + \mathbf{e}_i)$ and $T(\mathbf{n} \to \mathbf{n} - \mathbf{e}_i)$ from Eqs. (10) leads to

$$
\frac{d\langle n_k \rangle}{dt} = f_k \left(\langle n_k \rangle - \sum_j \frac{\langle n_k n_j \rangle}{N} \right) + m_k \left(1 - \sum_j \frac{\langle n_j \rangle}{N} \right) - \frac{\phi_k}{N} \langle n_k \rangle.
\tag{14}
$$

260 For other moments and models similar derivations can be done.

261 For the second moment, we find

$$
\begin{aligned}
\frac{d\langle n_k^2 \rangle}{dt} = {} & \ldots \sum_{n_k=0}^{N} \ldots (1 + 2n_k) P(\mathbf{n}, t) T(\mathbf{n} \to \mathbf{n} + \mathbf{e}_k) \\
& + \ldots \sum_{n_k=0}^{N} \ldots (1 - 2n_k) P(\mathbf{n}, t) T(\mathbf{n} \to \mathbf{n} - \mathbf{e}_k),
\end{aligned}
\tag{15}
$$

262 which after substituting $T(\mathbf{n} \to \mathbf{n} + \mathbf{e}_i)$ and $T(\mathbf{n} \to \mathbf{n} - \mathbf{e}_i)$ from Eqs. (10) reduces to

$$
\begin{aligned}
\frac{d\langle n_k^2 \rangle}{dt} =& f_k \left(\langle n_k \rangle - \sum_j \frac{\langle n_k n_j \rangle}{N} + 2 \left(\langle n_k^2 \rangle - \sum_j \frac{\langle n_k^2 n_j \rangle}{N} \right) \right) \\
&+ m_k \left(1 - \sum_j \frac{\langle n_j \rangle}{N} + 2 \left(\langle n_k \rangle - \sum_j \frac{\langle n_k n_j \rangle}{N} \right) \right) \\
&+ \frac{\phi_k}{N} \left(\langle n_k \rangle - 2 \langle n_k^2 \rangle \right).
\end{aligned}
\tag{16}
$$

263 For the co-moments, we find

$$
\begin{aligned}
\frac{d\langle n_k n_l \rangle}{dt} =& \ldots \sum_{n_k=0}^{N} \sum_{n_l=0}^{N} \ldots n_l P(\mathbf{n}, t) T(\mathbf{n} \to \mathbf{n} + \mathbf{e}_k) \\
&+ \ldots \sum_{n_k=0}^{N} \sum_{n_l=0}^{N} \ldots n_k P(\mathbf{n}, t) T(\mathbf{n} \to \mathbf{n} + \mathbf{e}_l) \\
&- \ldots \sum_{n_k=0}^{N} \sum_{n_l=0}^{N} \ldots n_l P(\mathbf{n}, t) T(\mathbf{n} \to \mathbf{n} - \mathbf{e}_k) \\
&- \ldots \sum_{n_k=0}^{N} \sum_{n_l=0}^{N} \ldots n_k P(\mathbf{n}, t) T(\mathbf{n} \to \mathbf{n} - \mathbf{e}_l),
\end{aligned}
\tag{17}
$$

264 which after substituting $T(\mathbf{n} \to \mathbf{n} + \mathbf{e}_i)$ and $T(\mathbf{n} \to \mathbf{n} - \mathbf{e}_i)$ from Eqs. (10) leads to

$$
\begin{aligned}
\frac{d\langle n_k n_l \rangle}{dt} =& (f_k + f_l) \left(\langle n_k n_l \rangle - \sum_j \frac{\langle n_k n_l n_j \rangle}{N} \right) \\
&+ m_k \left(\langle n_l \rangle - \sum_j \frac{\langle n_l n_j \rangle}{N} \right) + m_l \left(\langle n_k \rangle - \sum_j \frac{\langle n_k n_j \rangle}{N} \right) \\
&- \frac{\phi_k + \phi_l}{N} \langle n_k n_l \rangle.
\end{aligned}
\tag{18}
$$

Because each equation depends on higher moments, it is not possible to solve this system without
additional assumptions. However, one can find approximate expressions, where lower moments
replace higher moments. For example, $\langle n_k^2 n_j \rangle$ and $\langle n_k n_l n_j \rangle$ are approximated as functions of the
lower moments: $\langle n_k^2 \rangle$, $\langle n_k n_l \rangle$, and $\langle n_j \rangle$. Concretely, we can approximate e.g. $\langle n_k^2 n_j \rangle \approx \langle n_k^2 \rangle \langle n_j \rangle$.
This technique, called moment closure approximation, leads to a closed system of ODEs and we use
it in our approach. Kuehn [18] makes a thorough review on this technique.

A.2. Lotka-Volterra

Let us define the transition rates,

$$T(\mathbf{n} \to \mathbf{n} + \mathbf{e}_i) = n_i \left(f_i + \sum_j A_{i,j} n_j \right) \tag{19a}$$

$$T(\mathbf{n} \to \mathbf{n} - \mathbf{e}_i) = n_i \sum_j B_{i,j} n_j, \tag{19b}$$

where A and B are positively defined matrices, satisfying $A_{i,j} = 0$ if $B_{i,j} > 0$, and $B_{i,j} = 0$ if $A_{i,j} > 0$. Ecologically, while interactions in A promote growth, those in B lead to death. Finally, f_i is the intrinsic growth rate.

For the first moment, similarly to Eq. (13), we have

$$\frac{d\langle n_k \rangle}{dt} = \ldots \sum_{n_k=0}^{\infty} \ldots P(\mathbf{n}, t)(T(\mathbf{n} \to \mathbf{n} + \mathbf{e}_k) - T(\mathbf{n} \to \mathbf{n} - \mathbf{e}_k)), \tag{20}$$

which after substituting $T(\mathbf{n} \to \mathbf{n} + \mathbf{e}_i)$ and $T(\mathbf{n} \to \mathbf{n} - \mathbf{e}_i)$ from Eqs. (19),

$$\frac{d\langle n_k \rangle}{dt} = f_k \langle n_k \rangle + \sum_j (A_{k,j} - B_{k,j}) \langle n_k n_j \rangle, \tag{21}$$

which takes the form of the conventional, detemininistic Lotka-Volterra equations for the abundance with growth rate f_k and interaction matrix $A_{k,j} - B_{k,j}$.

For the second moment, similarly to Eq. (15)

$$\frac{d\langle n_k^2 \rangle}{dt} = \ldots \sum_{n_k=0}^{\infty} \ldots (1 + 2n_k) P(\mathbf{n}, t) T(\mathbf{n} \to \mathbf{n} + \mathbf{e}_k)$$
$$+ \ldots \sum_{n_k=0}^{\infty} \ldots (1 - 2n_k) P(\mathbf{n}, t) T(\mathbf{n} \to \mathbf{n} - \mathbf{e}_k), \tag{22}$$

which after substituting $T(\mathbf{n} \to \mathbf{n} + \mathbf{e}_i)$ and $T(\mathbf{n} \to \mathbf{n} - \mathbf{e}_i)$ from Eqs. (19) leads to

$$\frac{d\langle n_k^2 \rangle}{dt} = f_k(\langle n_k \rangle + 2\langle n_k^2 \rangle)$$
$$+ \sum_j (A_{k,j} + B_{k,j}) \langle n_k n_j \rangle + 2 \sum_j (A_{k,j} - B_{k,j}) \langle n_k^2 n_j \rangle. \tag{23}$$

283 For the co-moments, similarly to Eq. (17), we derived

$$
\begin{aligned}
\frac{d\langle n_k n_l \rangle}{dt} = & \dots \sum_{n_k=0}^{\infty} \sum_{n_l=0}^{\infty} \dots n_l P(\mathbf{n}, t) T(\mathbf{n} \to \mathbf{n} + \mathbf{e}_k) \\
& + \dots \sum_{n_k=0}^{\infty} \sum_{n_l=0}^{\infty} \dots n_k P(\mathbf{n}, t) T(\mathbf{n} \to \mathbf{n} + \mathbf{e}_l) \\
& - \dots \sum_{n_k=0}^{\infty} \sum_{n_l=0}^{\infty} \dots n_l P(\mathbf{n}, t) T(\mathbf{n} \to \mathbf{n} - \mathbf{e}_k) \\
& - \dots \sum_{n_k=0}^{\infty} \sum_{n_l=0}^{\infty} \dots n_k P(\mathbf{n}, t) T(\mathbf{n} \to \mathbf{n} - \mathbf{e}_l),
\end{aligned}
\tag{24}
$$

284 which after substituting $T(\mathbf{n} \to \mathbf{n} + \mathbf{e}_i)$ and $T(\mathbf{n} \to \mathbf{n} - \mathbf{e}_i)$ from Eqs. (19) reduces to

$$
\frac{d\langle n_k n_l \rangle}{dt} = (f_k + f_l)\langle n_k n_l \rangle + \sum_j (A_{k,j} - B_{k,j} + A_{l,j} - B_{l,j})\langle n_k n_l n_j \rangle.
\tag{25}
$$

A.3. From absolute to relative abundance

286 To focus on relative abundance data, we define the relative abundance as

$$
x_k = \frac{n_k}{\sum_j n_j},
\tag{26}
$$

287 and

$$
n_\Sigma = \sum_j n_j,
\tag{27}
$$

288 to serve as a scaling factor. Thus,

$$
n_k = x_k n_\Sigma.
\tag{28}
$$

289   Let us find the transformation to relative abundances for the first moment. Using the definition
290 of the covariance $\langle x_k, n_\Sigma \rangle = \langle x_k n_\Sigma \rangle - \langle x_k \rangle \langle n_\Sigma \rangle$, such that $\langle x_k n_\Sigma \rangle = \langle x_k \rangle \langle n_\Sigma \rangle + \langle x_k, n_\Sigma \rangle$, we have

$$
\begin{aligned}
\frac{d\langle n_k \rangle}{dt} &= \frac{d\langle x_k n_\Sigma \rangle}{dt} \\
&= \langle n_\Sigma \rangle \frac{d\langle x_k \rangle}{dt} + \langle x_k \rangle \frac{d\langle n_\Sigma \rangle}{dt} + \frac{d\langle x_k, n_\Sigma \rangle}{dt}.
\end{aligned}
\tag{29}
$$

291 Rearranging, the transformation is given by

$$
\frac{d\langle x_k \rangle}{dt} = \frac{1}{\langle n_\Sigma \rangle} \left(\frac{d\langle n_k \rangle}{dt} - \langle x_k \rangle \frac{d\langle n_\Sigma \rangle}{dt} - \frac{d\langle x_k, n_\Sigma \rangle}{dt} \right).
\tag{30}
$$

292   For second order moments, we use that $\langle x_k x_l n_\Sigma^2 \rangle = \langle x_k x_l \rangle \langle n_\Sigma^2 \rangle + \langle x_k x_l, n_\Sigma^2 \rangle$,

$$
\begin{aligned}
\frac{d\langle n_k n_l \rangle}{dt} &= \frac{d\langle x_k x_l n_\Sigma^2 \rangle}{dt} \\
&= \langle n_\Sigma^2 \rangle \frac{d\langle x_k x_l \rangle}{dt} + \langle x_k x_l \rangle \frac{d\langle n_\Sigma^2 \rangle}{dt} + \frac{d\langle x_k x_l, n_\Sigma^2 \rangle}{dt}.
\end{aligned}
\tag{31}
$$

293   Rearranging, the transformations are given by

$$\frac{d\langle x_k x_l\rangle}{dt} = \frac{1}{\langle n_\Sigma^2\rangle}\left(\frac{d\langle n_k n_l\rangle}{dt} - \langle x_k x_l\rangle\frac{d\langle n_\Sigma^2\rangle}{dt} - \frac{d\langle x_k x_l, n_\Sigma^2\rangle}{dt}\right),\tag{32}$$

294   and

$$\frac{d\langle x_k^2\rangle}{dt} = \frac{1}{\langle n_\Sigma^2\rangle}\left(\frac{d\langle n_k^2\rangle}{dt} - \langle x_k^2\rangle\frac{d\langle n_\Sigma^2\rangle}{dt} - \frac{d\langle x_k^2, n_\Sigma^2\rangle}{dt}\right),\tag{33}$$

295   where the differential equations for $\langle n_\Sigma\rangle$ and $\langle n_\Sigma^2\rangle$ are given by

$$\frac{d\langle n_\Sigma\rangle}{dt} = \frac{d\langle \sum_j n_j\rangle}{dt} = \sum_j \frac{d\langle n_j\rangle}{dt},\tag{34}$$

296   and

$$\frac{d\langle n_\Sigma^2\rangle}{dt} = \frac{d\langle (\sum_j n_j)^2\rangle}{dt} = \sum_i \sum_j \frac{d\langle n_i n_j\rangle}{dt}.\tag{35}$$

297   A close look at the covariances shows their contribution is negligible. To see this, let us write

$$
\begin{aligned}
\frac{d\langle x_k, n_\Sigma\rangle}{dt} =\ & \dots \sum_{n_k=0}^{\infty}\dots \left(\frac{n_k}{n_\Sigma} - \left\langle\frac{n_k}{n_\Sigma}\right\rangle\right)(n_\Sigma - \langle n_\Sigma\rangle)\,P(\mathbf{n}+\mathbf{e}_k, t)T(\mathbf{n}+\mathbf{e}_k \to \mathbf{n})\\
&+ \dots \sum_{n_k=0}^{\infty}\dots \left(\frac{n_k}{n_\Sigma} - \left\langle\frac{n_k}{n_\Sigma}\right\rangle\right)(n_\Sigma - \langle n_\Sigma\rangle)\,P(\mathbf{n}-\mathbf{e}_k, t)T(\mathbf{n}-\mathbf{e}_k \to \mathbf{n})\\
&- \dots \sum_{n_k=0}^{\infty}\dots \left(\frac{n_k}{n_\Sigma} - \left\langle\frac{n_k}{n_\Sigma}\right\rangle\right)(n_\Sigma - \langle n_\Sigma\rangle)\,P(\mathbf{n}, t)T(\mathbf{n} \to \mathbf{n}+\mathbf{e}_k)\\
&- \dots \sum_{n_k=0}^{\infty}\dots \left(\frac{n_k}{n_\Sigma} - \left\langle\frac{n_k}{n_\Sigma}\right\rangle\right)(n_\Sigma - \langle n_\Sigma\rangle)\,P(\mathbf{n}, t)T(\mathbf{n} \to \mathbf{n}-\mathbf{e}_k)\\
&+ \dots \sum_{n_k=0}^{\infty}\sum_{n_i=0}^{\infty}\dots \left(\frac{n_k}{n_\Sigma} - \left\langle\frac{n_k}{n_\Sigma}\right\rangle\right)(n_\Sigma - \langle n_\Sigma\rangle)\sum_{i\neq k} P(\mathbf{n}+\mathbf{e}_i, t)T(\mathbf{n}+\mathbf{e}_i \to \mathbf{n})\\
&+ \dots \sum_{n_k=0}^{\infty}\sum_{n_i=1}^{\infty}\dots \left(\frac{n_k}{n_\Sigma} - \left\langle\frac{n_k}{n_\Sigma}\right\rangle\right)(n_\Sigma - \langle n_\Sigma\rangle)\sum_{i\neq k} P(\mathbf{n}-\mathbf{e}_i, t)T(\mathbf{n}-\mathbf{e}_i \to \mathbf{n})\\
&- \dots \sum_{n_k=0}^{\infty}\sum_{n_i=0}^{\infty}\dots \left(\frac{n_k}{n_\Sigma} - \left\langle\frac{n_k}{n_\Sigma}\right\rangle\right)(n_\Sigma - \langle n_\Sigma\rangle)\sum_{i\neq k} P(\mathbf{n}, t)T(\mathbf{n} \to \mathbf{n}+\mathbf{e}_i)\\
&- \dots \sum_{n_k=0}^{\infty}\sum_{n_i=0}^{\infty}\dots \left(\frac{n_k}{n_\Sigma} - \left\langle\frac{n_k}{n_\Sigma}\right\rangle\right)(n_\Sigma - \langle n_\Sigma\rangle)\sum_{i\neq k} P(\mathbf{n}, t)T(\mathbf{n} \to \mathbf{n}-\mathbf{e}_i),
\end{aligned}
\tag{36}
$$

298 after the appropriate transformations of variable and re-indexing, we find

$$
\begin{aligned}
\frac{d\langle x_k, n_\Sigma \rangle}{dt} = & \ldots \sum_{n_k=0}^{\infty} \ldots \left(\frac{n_k - 1}{n_\Sigma - 1} - \left\langle \frac{n_k}{n_\Sigma} \right\rangle \right) (n_\Sigma - 1 - \langle n_\Sigma \rangle) P(\mathbf{n}, t) T(\mathbf{n} \to \mathbf{n} - \mathbf{e}_k) \\
& + \ldots \sum_{n_k=0}^{\infty} \ldots \left(\frac{n_k + 1}{n_\Sigma + 1} - \left\langle \frac{n_k}{n_\Sigma} \right\rangle \right) (n_\Sigma + 1 - \langle n_\Sigma \rangle) P(\mathbf{n}, t) T(\mathbf{n} \to \mathbf{n} + \mathbf{e}_k) \\
& - \ldots \sum_{n_k=0}^{N} \ldots \left(\frac{n_k}{n_\Sigma} - \left\langle \frac{n_k}{n_\Sigma} \right\rangle \right) (n_\Sigma - \langle n_\Sigma \rangle) P(\mathbf{n}, t) T(\mathbf{n} \to \mathbf{n} + \mathbf{e}_k) \\
& - \ldots \sum_{n_k=0}^{N} \ldots \left(\frac{n_k}{n_\Sigma} - \left\langle \frac{n_k}{n_\Sigma} \right\rangle \right) (n_\Sigma - \langle n_\Sigma \rangle) P(\mathbf{n}, t) T(\mathbf{n} \to \mathbf{n} - \mathbf{e}_k) \\
& + \ldots \sum_{n_k=0}^{\infty} \sum_{n_i=0}^{\infty} \ldots \left(\frac{n_k}{n_\Sigma - 1} - \left\langle \frac{n_k}{n_\Sigma} \right\rangle \right) (n_\Sigma - 1 - \langle n_\Sigma \rangle) \sum_{i \neq k} P(\mathbf{n}, t) T(\mathbf{n} \to \mathbf{n} - \mathbf{e}_i) \\
& + \ldots \sum_{n_k=0}^{\infty} \sum_{n_i=1}^{\infty} \ldots \left(\frac{n_k}{n_\Sigma + 1} - \left\langle \frac{n_k}{n_\Sigma} \right\rangle \right) (n_\Sigma + 1 - \langle n_\Sigma \rangle) \sum_{i \neq k} P(\mathbf{n}, t) T(\mathbf{n} \to \mathbf{n} + \mathbf{e}_i) \\
& - \ldots \sum_{n_k=0}^{\infty} \sum_{n_i=0}^{\infty} \ldots \left(\frac{n_k}{n_\Sigma} - \left\langle \frac{n_k}{n_\Sigma} \right\rangle \right) (n_\Sigma - \langle n_\Sigma \rangle) \sum_{i \neq k} P(\mathbf{n}, t) T(\mathbf{n} \to \mathbf{n} + \mathbf{e}_i) \\
& - \ldots \sum_{n_k=0}^{\infty} \sum_{n_i=0}^{\infty} \ldots \left(\frac{n_k}{n_\Sigma} - \left\langle \frac{n_k}{n_\Sigma} \right\rangle \right) (n_\Sigma - \langle n_\Sigma \rangle) \sum_{i \neq k} P(\mathbf{n}, t) T(\mathbf{n} \to \mathbf{n} - \mathbf{e}_i).
\end{aligned}
\tag{37}
$$

299 Note that if $n_\Sigma \gg 1$, $n_\Sigma \pm 1 \approx n_\Sigma$, then, if either $n_k \gg 1$, such that $n_k \pm 1 \approx n_k$ or $\frac{n_\Sigma - \langle n_\Sigma \rangle}{n_\Sigma} \approx 0$,
300 the terms from the previous equation simplify, leading to

$$
\frac{d\langle x_k, n_\Sigma \rangle}{dt} \approx 0.
\tag{38}
$$

301 Similar arguments lead to conclude that,

$$
\frac{d\langle x_k x_l, n_\Sigma^2 \rangle}{dt} \approx 0,
\tag{39}
$$

302 and

$$
\frac{d\langle x_k^2, n_\Sigma^2 \rangle}{dt} \approx 0.
\tag{40}
$$

303 These approximations of the covariances are sensible in microbiomes where $n_\Sigma, n_k \gg 1$ is often the
304 case. Moreover, in the infinite population limit, covariances must be zero.

305   Putting all together, the first moment of relative abundance in large populations is given by

$$\frac{d\langle x_k \rangle}{dt} = \frac{1}{\langle n_\Sigma \rangle} \left(\frac{d\langle n_k \rangle}{dt} - \langle x_k \rangle \sum_j \frac{d\langle n_j \rangle}{dt} \right)$$
$$= \frac{1}{\langle n_\Sigma \rangle} \left(\frac{d\langle n_k \rangle}{dt} - \langle x_k \rangle \frac{d\langle n_\Sigma \rangle}{dt} \right),$$

(41)

306   while for the second moments of relative abundance

$$\frac{d\langle x_k x_l \rangle}{dt} = \frac{1}{\langle n_\Sigma^2 \rangle} \left(\frac{d\langle n_k n_l \rangle}{dt} - \langle x_k x_l \rangle \sum_i \sum_j \frac{d\langle n_i n_j \rangle}{dt} \right)$$
$$= \frac{1}{\langle n_\Sigma^2 \rangle} \left(\frac{d\langle n_k n_l \rangle}{dt} - \langle x_k x_l \rangle \frac{d\langle n_\Sigma^2 \rangle}{dt} \right),$$

(42)

307   and

$$\frac{d\langle x_k^2 \rangle}{dt} = \frac{1}{\langle n_\Sigma^2 \rangle} \left(\frac{d\langle n_k^2 \rangle}{dt} - \langle x_k^2 \rangle \sum_i \sum_j \frac{d\langle n_i n_j \rangle}{dt} \right)$$
$$= \frac{1}{\langle n_\Sigma^2 \rangle} \left(\frac{d\langle n_k^2 \rangle}{dt} - \langle x_k^2 \rangle \frac{d\langle n_\Sigma^2 \rangle}{dt} \right).$$

(43)

308   Finally, to solve these equations in terms of relative abundance, the change of variable $n_i = x_i n_\Sigma$

309   is needed all along. As Joseph et al. [14], we see that the second term of each equation serves as

310   "correction factor" due to the fact that relative abundances must add up to one at all times.

 B. True parameters in simulations

	growth f_i	death ϕ_i	immigration m_i	shared carrying capacity N	initial population size
$i=1$	0.5	100	2000	10^5	2800
$i=2$	0.8	130	4000	10^5	1400
$i=3$	1.4	160	3000	10^5	700

Table 1: **Parameters in simulated logistic growth with immigration and death (Fig. 2 A).** The growth and death rates as well as the immigration parameters were only chosen for illustration, thus, time units are arbitrary.

		interaction $I_{i,j}$			
	growth f_i	$j=1$	$j=2$	$j=3$	initial population size
$i=1$	0.9	$-4.8 \cdot 10^{-5}$	$2.7 \cdot 10^{-5}$	$-5 \cdot 10^{-5}$	700
$i=2$	1.7	$-2.9 \cdot 10^{-5}$	$-6.3 \cdot 10^{-5}$	$3.3 \cdot 10^{-5}$	7000
$i=3$	1.3	$2 \cdot 10^{-5}$	$-4.1 \cdot 10^{-5}$	$-5.4 \cdot 10^{-5}$	12000

Table 2: **Parameters in simulated Lotka-Volterra (Fig. 2 B).** The interaction parameters as well as the growth rate were only chosen for illustration, thus, time units are arbitrary.

 # C. Inference settings

parameter	units	prior
growth f_i	time^{-1}	$\mathcal{U}(0,2)$
death ϕ_i	cells $\cdot$ time^{-1}	$\mathcal{U}(0,200)$
immigration m_i	cells $\cdot$ time^{-1}	$\mathcal{N}(3000,1000)$
shared carrying capacity N	cells	$\mathcal{N}(10^5,10^4)$

Table 3: **Priors for simulated logistic growth with immigration and death (Fig. 2 A).** A combination of un-informative (uniform) and informative (normal) priors were used for illustration. These were wide to test the ability of the inference workflow to find the true parameters in simulations (Table 1). $\mathcal{U}(a,b)$ indicates a uniform distribution in the range from a to b. $\mathcal{N}(a,b)$ indicates a normal distribution with mean a and standard deviation b.

parameter	units	prior
growth f_i	time^{-1}	$\mathcal{N}(1.5,0.5)$
intra-specific $I_{i,i}$	(cells $\cdot$ time)$^{-1}$	$\mathcal{U}(-10^{-4},0)$
inter-specific $I_{i,j}$	(cells $\cdot$ time)$^{-1}$	$\mathcal{N}(0,5 \cdot 10^{-5})$

Table 4: **Priors for simulated Lotka-Volterra (Fig. 2 B).** A combination of uninformative (uniform) and informative (normal) priors were used for illustration. These were wide to test the ability of the inference workflow to find the true parameters in simulations (Table 2). $\mathcal{U}(a,b)$ indicates a uniform distribution in the range from a to b. $\mathcal{N}(a,b)$ indicates a normal distribution with mean a and standard deviation b.

parameter	units	prior	motivation
growth f_i	day^{-1}	$\mathcal{U}(0.2, 48)$	from 5 days to 30 minutes doubling time [19]
death ϕ_i	cells $\cdot$ day^{-1}	$\mathcal{U}(0, 10^6)$	from 0 to ≈ 11.5 cells/second*
immigration m_i	cells $\cdot$ day^{-1}	$\mathcal{U}(0, 10^6)$	from 0 to ≈ 11.5 cells/second*
shared carrying capacity N	cells	$\mathcal{U}(1.4 \cdot 10^7, 1.6 \cdot 10^7)$	maximum number of cells in data [12]

Table 5: **Priors for logistic growth with immigration and death in empirical mouse data (Fig. 3).** Available evidence and back-of-the-envelope calculations (marked by *) were used to propose wide priors. $\mathcal{U}(a, b)$ indicates a uniform distribution in the range from a to b. $\mathcal{N}(a, b)$ indicates a normal distribution with mean a and standard deviation b.

	simulations (Fig. 2)		mouse data (Fig. 3)
	logistic	Lotka-Volterra	logistic
numeric integration (numpy)			
method	LSODA		LSODA
ABC-SMC (pyABC)			
calibration samples (to get first ε)	2000	2000	2000
* max. number of generations	30	30	10
accepted samples per generation	AdaptivePopulationSize (CV = 0.4)		APS (CV = 0.3)
samples in first generation	750	750	2000
min. samples per generation	500	500	320
max. samples per generation	1000	1000	640
mismatch threshold (ε) update	QuantileEpsilon (best 10%)		QE (10%)
strategy to sample parameters	MultivariateNormalTransition		MNT
* mismatch threshold (ε) min.	$1.25 \cdot 10^8$	$6.6 \cdot 10^7$	*NA*
* min. ε change btw. generations	*NA*	*NA*	10^{13}

Table 6: **Settings for Approximate Bayesian Computation - Sequential Monte Carlo (ABC-SMC) code.** These settings were chosen to decrease the computing time, but still robustly minimize the distance between data and model. We used tools from the Python package pyABC [16], mainly *ABCSMC*. * The maximum number of generations, mismatch threshold (ε) minimum, and minimum ε change between generations, are all stopping criteria. LSODA is a numerical solver capable of selectively adapting to the stiffness of a system of differential equations.