1

3 Title: *FSHβ* links photoperiodic signalling to seasonal reproduction in

4 Japanese quail.

6 Short Title: Annual timing in birds

8 Authors: Gaurav Majumdar[1], Timothy A Liddle[2], Calum Stewart[2], Christopher J Marshall[2],

9 Maureen Bain[2], Tyler J Stevenson[2]*

10
11 Affiliations: [1]Department of Zoology, Science campus, University of Allahabad, India,
12 [2]School of Biodiversity, One Health and Veterinary Medicine University of
13 Glasgow, Glasgow, United Kingdom.
14

15
16
17
*Corresponding author:
Tyler John Stevenson
School of Biodiversity, One Health, and Veterinary Medicine
University of Glasgow
Glasgow, United Kingdom
e. tyler.stevenson@glasgow.ac.uk
p. 01413305793
25

26 Key words: Neuroendocrine, Photoperiod, Reproduction, Endogenous, Pituitary,
27
28 This manuscript includes:
29 Main Text
30 Figure 1 to 3
31 Supporting Information Tables S1-S12
32 Supporting Information Figures S1-S8
33
34

Abstract

Annual cycles in daylength provide an initial predictive environmental cue that plants and animals use to time seasonal biology. Seasonal changes in photoperiodic information acts to entrain endogenous programs in physiology to optimize an animal's fitness. Attempts to identify the neural and molecular substrates of photoperiodic time measurement in birds have, to date, focussed on blunt changes in light exposure during a restricted period of photoinducibility. The objectives of these studies were first to characterise a molecular seasonal clock in Japanese quail and second, to identify the key transcripts involved in endogenously generated interval timing that underlies photosensitivity in birds. We hypothesized that the mediobasal hypothalamus (MBH) provides the neuroendocrine control of photoperiod-induced changes in reproductive physiology, and that the pars distalis of the pituitary gland contains an endogenous internal timer for the short photoperiod dependent development of reproductive photosensitivity. Here we report distinct seasonal waveforms of transcript expression in the MBH, and pituitary gland and discovered the patterns were not synchronized across tissues. Follicle-stimulating hormone-β (*FSHβ*) expression increased during the simulated spring equinox, prior to photoinduced increases in prolactin, thyrotropin-stimulating hormone-β and testicular growth. Diurnal analyses of transcript expression showed sustained elevated levels of *FSHβ* under conditions of the spring equinox, compared to autumnal equinox, short (<12L) and long (>12L) photoperiods. *FSHβ* expression increased in quail held in non-stimulatory short photoperiod, indicative of the initiation of an endogenously programmed interval timer. These data identify that FSHβ establishes a state of photosensitivity for the external coincidence timing of seasonal physiology. The independent regulation of FSHβ expression provides an alternative pathway through which other supplementary environmental cues, such as temperature, can fine tune seasonal reproductive maturation and involution.

Introduction

Seasonal rhythms in reproduction are ubiquitous in plants and animals. In birds, the annual change in daylength, referred to as photoperiod, provides an initial predictive environmental cue to time seasonal physiology and behavior (Ball, 1993). Temperature, nutrient availability and social cues act as supplementary cues that function to fine tune the timing of breeding (Wingfield and Farner, 1980). Seasonal timing of reproductive physiology and breeding requires the integration of both environmental cues and endogenously generated mechanisms (Gwinner, 1986; Wingfield, 2008; Helm & Stevenson, 2014). Even in the absence of seasonal fluctuations in daylength, temperature, and food availability, endogenous circannual cycles in migration (Gwinner and Dittami, 1990), hibernation (Pengelley and Fisher, 1957), and reproduction (Woodfill et al., 1994; Lincoln et al., 2006) are maintained with remarkable temporal precision. Other endogenous timing mechanisms include interval timers that are programmed to establish a physiological state in anticipation of the next season, such as flowering in plants (Duncan et al., 2015) and the photorefractory state in rodents (Prendergast et al. 2001). The anatomical and cellular basis of endogenous programs that time seasonal transitions in biology remain poorly characterized, but current evidence indicates that, in mammals, circannual time may reside in pituitary lactotropes (Lincoln et al., 2006) and thyrotropes (Wood et al., 2020).

In most long-lived species (e.g., >2 yrs.), the annual change in photoperiod acts to entrain endogenous annual programs to time transitions in physiological state to seasons (Bradshaw and Holzapfel, 2007). In many temperate-zone birds, exposure to long days (>12hr), induces a photostimulated state in which gonadal development occurs in male and female birds. Short photoperiods (e.g., <12hr) can induce gonadal involution in both sexes leading to a reproductively regressed state (Dawson et al., 2001). Birds become reproductively sensitive to stimulatory long photoperiods only after experiencing short

photoperiods for at least ten days, in which a photosensitive state is established (Dawson

et al., 2001).

In most birds, reptiles, and amphibians, annual changes in daylength are detected

by photoreceptors located in the mediobasal hypothalamus (MBH) (Perez et al., 2019).

Stimulatory long photoperiods trigger a molecular cascade that starts with the upregulation

of thyrotropin-stimulating hormone beta subunit (*TSHβ*) in the pars tuberalis of the pituitary

gland and results in gonadal maturation (Nakao et al., 2008). The current gap in our

knowledge is how short days (i.e., <12hr) induce gonadal involution, and how prolonged

exposure to short days stimulates endogenous programs that sensitises the brain to

respond, at a molecular level, to stimulatory long days (Follett and Sharp, 1969). Previous

studies in European starlings (*Sturnus vulgaris*) demonstrated that exposure to short days

increased Gonadotropin-releasing hormone (*GNRH*) expression in the preoptic area

(Stevenson et al., 2009; Stevenson et al., 2012a). As both GnRH and gonadal growth

increased in the absence of stimulatory long day lengths, another cellular pathway must

also be involved in the endogenous development of reproductive physiology in birds.

There were three objectives of the present work. First, we aimed to characterise the

photoperiod-induced seasonal molecular clock in the MBH and pituitary gland in Japanese

quail. Then, we examined the daily waveform of multiple transcripts in the MBH and

pituitary in birds from stimulatory long photoperiod (16L:8D), inhibitory short photoperiod

(8L:16D) and the two equinoxes. The last objective was to determine if *FSHβ* expression

in the pituitary gland was upregulated after prolonged exposure to short photoperiods. We

theorized multiple neuroendocrine regions including the preoptic area, mediobasal

hypothalamus and pituitary cells are independently involved in the timing of seasonal

transitions in physiology. Specifically, we hypothesized that pituitary gonadotropes

establish a state of photosensitivity to stimulatory long day photoperiods and thus act as

calendar cells that provide endogenous timing of gonadal growth.

Results

Molecular characterisation of the photoperiod-induced seasonal clock

To obtain a comprehensive understanding of the seasonal molecular changes in the MBH and pituitary gland, we collected MBH and pituitary gland samples from Japanese quail using an experimental paradigm that aimed to maximize resolution (i.e., high sampling frequency), high-dimensionality (i.e., advanced nucleic acid sequencing) and robust statistical power (i.e., large sample sizes). Our experimental design simulated the photoperiodic regulation of seasonal physiology of Japanese quail, using sequential changes of an autumnal decrease, followed by a spring increase in daylength and measured testes volume, body mass and abdominal fat (Fig.1a). As anticipated (Follett and Sharp, 1969; Robinson and Follett, 1982), light phases that exceed the critical day length (i.e., >12hr) resulted in robust gonadal growth. Increases in body mass and abdominal fat deposition were delayed until the spring increase in day lengths reached 10hrs (10v) (Fig. 1b,c). EdgeR analyses, of MBH sequences obtained using Minion, identified 1481 transcripts were differentially expressed (P<0.05) (Table S1; Table S2), and BioDare2.0 established 398 have rhythmic patterns (Fig. 1e; Fig. S1; Table S3). DAVID gene ontology analyses indicated that gonadotropin-releasing hormone receptor and Wnt signalling pathways were consistently identified as the predominant cellular mechanism recruited during each photoperiodic transition (Table S2). Increased proopiomelanocortin (*POMC*) expression coincided with body mass growth and abdominal fat accumulation (Fig. 1f). Thyroid hormone catabolism enzyme deiodinase type-3 (*DIO3*) increased after prolonged exposure to non-stimulatory photoperiods and was only therefore transiently elevated from 8L to 10v (10v; Fig. 1g).

Next, we used vimentin immunoreactivity to examine changes in tanycytes morphology in relation to changes in deiodinase transcript expression. Highest levels of

vimentin immunoreactivity in the median eminence coincided with the peak in *DIO3*

expression (Fig. 1h) suggesting the localized removal of active thyroid hormone is limited

to a short phase and occurred prior to stimulatory photoperiods (Fig. 1i). The photoperiod-

induced change in vimentin was anatomically localized to the median eminence, as the

area of immunoreactivity in the dorsal 3^{rd}V ependymal layer did not change with seasonal

transitions in reproduction (Fig. S2).

Next, we used MinION to sequence transcripts in the pituitary gland across the

seasonal transitions in reproduction. EdgeR analyses identified 3090 transcripts were

differentially expressed in the pituitary gland (Table S1; Table S4). DAVID gene ontology

analyses identified that gonadotropin-releasing hormone receptor and epidermal growth

factor pathways are consistently observed across photoperiodic transitions (Table S4).

BioDare2.0 established that 130 transcripts were rhythmically expressed (Fig. 1j; Table

S5; Fig. S1). A remarkable 96.5% (384/398) of MBH transcripts show a photoperiod-

induced spiked patterned and only 12/14 remaining transcripts displayed sine waveforms.

Conversely, 100% of pituitary transcripts conformed to sine (124/130) or cosine (6/130)

waveforms. The predominant rhythmic patterns of expression likely reflect endogenous

long-term molecular programs, characteristic of calendar cell function.

Lastly, we examined *GNRH* expression to delineate photoperiod-induced changes

in the neuroendocrine control of reproductive physiology. *GNRH* expression in the preoptic

area was temporarily decreased during the 8L short photoperiod indicating another

nucleus specific cellular timer is present in this brain region (Fig. S2). Overall, the pituitary

showed a distinct transcriptomic profile compared to the MBH suggesting independence in

the representation of seasonal photoperiodic timing. The spring photoperiodic transition

from 8L to 12v resulted in a significant increase in *FSHβ* (Fig. 1k) without a change in

tanycyte restructuring (Fig. 1h). Subsequent transition to 14v and then 16v resulted in the

upregulation of prolactin (*PRL*) (Fig. 1l) and thyrotropin-stimulating hormone-β (*TSHβ*),

respectively (Fig. 1m). These data establish that multiple cells in the pituitary code

seasonal photoperiodic time and *FSHβ* shows an endogenous increase in expression prior

to reproductively stimulatory photoperiods.

Weighted gene co-expression network analyses (WGCNA) were conducted to

discover gene co-expression modules, and then examine whether any of the resulting

module eigengenes co-vary with photoperiod or physiological measures. The eigengene

dendrogram of sequencing from individual animals was plotted and a heatmap of

physiological factors was organised (Fig. S3). The scale-free topology and mean sample

independence was assessed to determine a soft-threshold of 5 for both MBH and the

pituitary gland sequencing datasets. 10 modules were identified for the MBH, and the

pituitary gene set was grouped into 22 modules (Fig. S3). Of these modules there were 6

significant module-trait relationships in the MBH. There was one module with a significant

negative correlation with fat score (Fig. S3; Table S6). 44 transcripts were identified to be

significant in the negative relation for fat score. The other 5 were identified to be negatively

related and included photoperiod, body mass, fat score, testes width, and testes volume

(Fig. S3; Table S6). Overall, there were 23 transcripts that were significant and overlapped

with photoperiod, testes width and testes volume. The other module found 70 transcripts

for both body mass and fat score. Despite several modules showing trends toward

significance, only one module for body mass was positively related in the pituitary gland

(Fig. S3; Table S6). There were 206 transcripts identified to be significantly positively

related to body mass.

To ascertain common molecular mechanisms involved in the transcriptional

regulation of photoperiodically regulated transcripts, transcription factor enrichment

analysis was conducted on significant MBH (Table S7) and pituitary gland (Table S8)

transcripts. Association plots show no overlap in DNA binding motifs between MBH and

pituitary transcripts (Fig. S4) suggesting tissue-specific transcription binding factor

regulation. Within the pituitary gland, several common transcription factors, such as the

Jun proto-oncogene, Spi1 proto-oncogene and myocyte enhancer factor 2 (*MEF2a*) might

be actively involved in the photoperiodic regulation of transcript expression. These findings

indicate multiple transcription factors are likely recruited to control tissue-specific, and cell-

specific transcript expression and seasonal life history transitions in physiology.

Increased *FSHβ* expression is programmed during the spring equinox.

To establish whether increased pituitary *FSHβ* during the spring 12hr transition

reflects constitutively elevated expression or is driven by the sampling 'time of day', we

collected tissue samples every 3 hours from quail that transitioned to the autumnal

equinox (i.e., 12a), the short photoperiod (8L), spring equinox (12v) and long photoperiod

(16L) seasonal phases (Fig. 2a). Photoinduced changes in testes volume confirmed

seasonal reproductive condition (Fig. 2b). Circadian clock genes aryl hydrocarbon receptor

nuclear translocator like (*ARNTL1*) and period 3 (*PER3*) exhibit robust anti-phase daily

waveforms in expression, in the pituitary gland (Fig. 2c,d) (Table S9). Only *ARNTL1*, but

not *PER3* nor *DIO2*, had a rhythmic waveform in the MBH (Table S9; Fig. S5). Consistent

with the previous study, *FSHβ* expression was higher at the spring equinox compared to

all other photoperiod groups (Fig. 2e). *PRL* expression was higher in long photoperiod

(16L) compared to the two equinox and short photoperiod (8L) (Fig. 2f). *FSHβ* did not

display a daily rhythm, which is likely due to the absence of D-box and E-box motifs in the

FSHβ promoter (Fig. 2g). *FSHβ* promoter does contain many DNA motifs that are targeted

by several transcription factors that are responsive to hormonal and nutrient pathways

indicating multiple upstream regulators are recruited to drive transcription (Table S10).

These data support the conjecture that a long- term programmed increase in *FSHβ* occurs

under spring non-stimulatory photoperiod, and it is not driven by short-term daily photic

cues.

FSHβ expression establishes endogenously programmed photosensitivity.

To identify if *FSHβ* expression is driven by an endogenously programmed mechanism or in response to the gradual increase in light, adult quail were exposed to the 8L, 10v or 12v light schedules or kept in 8L for an additional 4 weeks (8Lext) (Fig. S6). The 8Lext treatment permitted confirmation whether *FSHβ* expression would increase in that photoperiod, and therefore reflect an interval timing mechanism. *FSHβ* expression increased 19-fold after 4 additional weeks of 8L suggesting that endogenous drivers initiate transcription despite no change in daylength (P<0.05) (Fig. 3a). But the dominant stimulator of *FSHβ* expression was the transition to 10v and 12v photoperiod, which both expressed significantly increased levels compared to 8L (P<0.001). These data demonstrate that both photoperiod and endogenous timing mechanisms drive *FSHβ* expression in the pars distalis. It is likely that an additive function of endogenous timing and the spring increase in photoperiod drive *FSHβ* expression.

As Opn5 was identified in the pituitary transcriptome, we then assessed its transcript expression across the photoperiod treatments. We discovered that Opn5 expression patterns paralleled *FSHβ* suggesting the potential for direct light detection by Opn5 and subsequent regulation of *FSHβ* expression. Based on *FSHβ* promoter analyses (Fig. 2g), we examined the expression of myocyte enhancer factor 2 (*MEF2a*) expression as a potential upstream regulator of *FSHβ* expression. *MEF2a* remained relatively constant suggesting this transcription factor binding protein is not the primary driver of *FSHβ* expression (Fig. S6). Similarly, *DNMT3a* expression did not change across photoperiod treatments (Fig. S6) suggesting that epigenetic modifications (i.e., DNA methylation) may not provide the endogenous programmed change leading to constitutive *FSHβ* expression. The precise molecular change upstream from *FSHβ* transcription remains to be identified. In the MBH, *DIO3* displayed a rapid reduction in expression after

transfer to 10v and was found to be significantly reduced in 12v photoperiod (P<0.001)

(Fig. 3c). There was a significant difference in MBH *DIO2* expression (Fig. S6), but this

observation was driven by a decrease in 8ext. Interestingly, *GNRH* in the preoptic area

was found to significantly increase after extended exposure to short photoperiods (Fig.

3D). These data indicate that endogenous switches in *FSHβ,* and possibly *GNRH*

expression, in response to short photoperiods may reflect multiple independent cellular

timers that establish a physiological state of photosensitivity.

Discussion

This report used the well-characterised photoperiodic manipulation of the Japanese quail

avian photoperiodic response using a laboratory-based light-schedule that accurately

replicated findings from birds held in semi-natural conditions (Robinson and Follett, 1982).

The data reported herein demonstrate that photoperiods less that 12hrs light induce

gonadal involution. Prolonged exposure to short photoperiods (i.e., 8L) were found to

significantly increase *DIO3* expression and vimentin immunoreactivity in the median

eminence. Increased *DIO3* expression and innervation of tanycytes occurred after gonadal

regression suggesting that another unidentified mechanism is involved in the initiation of

the termination in the breeding state. The gradual increase in photoperiods during the

spring transition was found to be associated with a marked increase in *FSHβ* expression in

the pars distalis while lower levels of *TSHβ* expression in the pars tuberalis were

maintained. As photoperiods increase there is a steady elevation in *FSHβ* expression, but

vimentin immunoreactivity in the median eminence did not decline until after the critical

daylength for photostimulation (i.e., 12L:12D) thus the release of FSH is prevented as

daylengths are below the critical threshold. Previous reports established that *TSHβ*

expression is significantly increased during the period of photoinducibility in quail (Nakao

et al., 2008). Although the present study did not directly examine photoinduction, *TSHβ*

expression was consistently elevated in long day photoperiod (i.e., 16L). The patterns of

expression suggest that stimulatory daylengths longer than 12hr induce thyrotropes to

increase *TSHβ* leading to a cascade of molecular events in the MBH that permit GnRH to

stimulate gonadotropes to release FSHβ and initiate gonadal development (Nakao et al.,

2008; Yoshimura et al., 2003; Yamamura et al., 2004). Note that other pituitary cell types,

somatotropes and corticotropes do not appear to show any molecular switches across the

photoperiodic phases. These findings uncover a two-component mechanism for the

cellular basis of the external coincidence model for the avian photoperiodic response (Fig.

S7). Increase *FSHβ* expression establishes a state of photosensity to stimulatory day

length, and *TSHβ* thryotropes in the pars tuberalis monitor daylength and when light

stimulation occurs during a period of photoinducibility, initiate gonadal development. The

two-component model is exciting as it accommodates evidence for endogenous growth of

quail gonads in the absence of photostimulation (Follett and Sharp, 1969). Moreover, a

TSH-independent programmed change in *FSHβ* expression addresses how seasonal

rhythms in tropical, non-photoperiod birds can be regulated (Gwinner and Dittami, 1990).

The photoperiodic induction of gonadal growth in quail is dependent on circadian

timing mechanisms (Follett and Sharp, 1969). However, only a few proximal promoters of

photoperiodic genes contain D-box elements required for circadian timing input and

include eyes absent-3 (*EYA3*) and *TSHβ*, but not *FSHβ* (Liddle et al., 2022). Similarly, E-

box elements are only identified in the proximal promoter of *EYA3*. The presence of E- and

D-boxes provides a clear molecular mechanism by which the circadian clock can control

the long photoperiod induced expression of these highly photoperiodic genes. Conversely,

FSHβ expression did not show diurnal variation and instead maintained constitutive

expression across long (16L:8D), short (8L:16D), and the 'equinox' (12L:12D)

photoperiodic conditions. We used a broad, unbiased bioinformatic approach to identify

putative transcriptional bindings sites that may regulate *FSHβ* expression. We identified

several potential transcriptional binding proteins and *MEF2* motifs were observed across multiple promoters of genes for transcripts in the pars distalis and pars nervosa of the pituitary gland. However, functional analyses are necessary to establish a causal link between these newly identified signalling pathways (e.g., MEF2) and the seasonal regulation of transcript expression.

The high-dimensionality and high-frequency analyses of seasonal transition in physiology used in this study, facilitated the ability to uncover that photoperiod differentially regulates two endocrine systems: reproduction and energy balance. *POMC* has well described roles in the neuroendocrine regulation of food intake and body mass (Yeo and Heisler, 2012). We found that *POMC* expression increased in response to short photoperiods and was associated with delayed body mass and adipose tissue growth. Interestingly, there was a gradual increase in body mass and adipose tissue mass during the transition for short photoperiod to the spring equinox despite elevated levels of *POMC* expression (i.e., adipose). *POMC* levels did not decrease until after exposure to stimulatory long photoperiods. Given the consistent photoperiod-induced change in *POMC* expression across animals (Helfer and Stevenson, 2020), these data provide significant insight into the temporal regulation of the central, and peripheral control of seasonal energy balance. As the closely related European quail (*Coturnix coturnix*) are migratory (Dorst, 1956; Bertin et al., 2007), the increased fattening observed early in the spring transition may reflect a conserved seasonal physiological response to ensure energy stores are provided for migration.

The integration of environmental cues to time breeding in birds varies between male and female birds (Ball and Ketterson, 2008; Tolla and Stevenson 2022). In most temperate breeding males, full reproductive development can be achieved in response to photoperiod cues. The robust change in gonadal growth and involution provides a powerful approach to identify the key neuroendocrine mechanisms that govern the avian photoperiodic

response. Despite ovarian changes in response to photoperiodic manipulations, female

birds generally require other supplemental cues (e.g., temperature, social cues) to attain

full reproductive development (Wingfield, 1980). In female, white-crowned sparrows

(*Zonotrichia leucophrys*), increased photoperiod induces ovarian development to a pre-

breeding state (Farner et al., 1966), and supplementary cues, such as temperature, can

modify ovarian growth (Wingfield et al., 1996; Wingfield et al., 2003). In Corsica, two

populations of great tits (*Parus major*) differ in egg laying date by up to one-month despite

males from both regions displaying similar timing in reproductive development (Caro et al.,

2006). As the initial predictive environmental cue (i.e., photoperiod) times reproduction

similarly in both male and female birds, the data provided in this paper provides key

insights into the fundamental mechanisms that govern transitions in the hypothalamo-

pituitary control of reproduction. However, studies that seek to understand how

supplementary cues (e.g., temperature) are integrated to fine tune the timing of

reproduction will require a focus on female birds.

In conclusion, these studies provide a comprehensive transcriptome dataset that

can facilitate ecological studies that seek to uncover the molecular substrates which

environmental cues, such as temperature, impact phenological timing and mistiming in

birds (Visser and Gienapp, 2019). The observation for photoperiod-independent regulation

of *FSHβ* expression provides a new cellular mechanism in which supplementary

environmental cues, such as temperature, can regulate the timing of seasonal

reproduction. For example, the marked population differences in Great tit laying dates in

Corsica, despite similar daylength cues, might be driven by local temperatures cues acting

on *FSHβ* expression to advance, or delay follicular maturation. Overall, the data indicate a

multi-cellular, multi-neural interval timing mechanism resides in the brain and has

significant implications for understanding species-specific seasonal transitions in life-

histories.

Materials and Methods

Data and code availability statement
All raw data are available in Table S1. Raw sequencing data is available in Gene
expression omnibus database GSE241775 and BioProject PRJNA1009845. R code used
in available in Table S11.

Animals
All Japanese quail were provided by Moonridge Farms, Exeter United Kingdom
(moonridgefarms.co.uk). Chicks were raised under constant light and constant heat lamp
conditions. 5-week-old male birds were delivered to the Poultry facilities at the University
of Glasgow, Cochno Farm in September 2019 and 2020. Both male and female birds
respond to changes in photoperiod (Ball and Ketterson, 2008; Farner et al., 1966). Only
males were used in the present studies as the robust change in gonadal volume provides
a powerful approach to maintain strong statistical power with fewer animals. Food (50:50
mix of Johnston and Jeff, quail mix & Farm Gate Layers, poultry layers supplemented with
grit) and tap water was provided ad libitum. All procedures were in accordance with the
National Centre for the Replacement, Refinement and Reduction of Animals in Research
ARRIVE guidelines (https://www.nc3rs.org.uk/revision-arrive-guidelines). All procedures
were approved by the Animal Welfare and Ethics Review Board at the University of
Glasgow and conducted under the Home Office Project Licence PP5701950.

Study 1 – Photoperiod-induced transition in seasonal life history states
Male quail (N=108) were housed in a summer-like long-day (LD) photoperiod (16L:8D). To
mimic the autumnal decline and subsequent spring increase in the annual photoperiodic
cycle, birds were exposure to a sequential change in day length from 16L (16a), to 14L
(14a), to 12L (12a), to 10L (10a), to 8L, then back to 10L (10v), to 12L (12v), to 14L (14v)
and lastly 16L (16v)(Fig.1a). Each photoperiod treatment lasted for 2 weeks to minimize
the impact of photoperiodic history effects (Stevenson et al., 2012a). At the end of each
photoperiodic treatment a subset of quail (n=12) body mass was used as a measure to
pseudo randomly select birds for tissue collection and served to reduce the potential for
unintentional bias. Birds were killed by cervical dislocation followed by jugular cut. A
jugular blood sample was collected in 50ul heparinized tubes (Workhardt, UK) and stored
at -20°C. Brain and pituitary stalk were rapidly dissected, frozen on powdered dry ice and
stored at -80°C. Testes were dissected and weighed to the nearest 0.001g using the
Sartorius microbalance (Sartorius, Germany). The length and width of the testes were
measured using callipers and volumes were calculated using the equation for a spheroid
$(4/3 \times 3.14 \times [L/2] \times [W/2]2)$ (King et al., 1997). Body mass was measured using an Ohaus
microscale to the nearest 0.1g. Fat score was assessed using the common scale
developed by Wingfield and Farner (1978). The scale range is 0 to 5, which 0 is no visible
fat and 5 indicates bulging fat bodies are present.

Study 2 – Daily rhythms in molecular profiles during solstices and equinoxes
To investigate the daily molecular representation of summer-like and winter-like solstices
and the autumnal and spring equinoxes, Japanese quail (N=188) were subjected to the
same photoperiodic treatments described in Study 1. A subset of quail was pseudo
randomly selected after the autumnal 12L:12D (n=47), short day 8L:16D (n=48), spring
12L:12D (n=46) and long day 16L:8D (n=47) treatment conditions. For each of the four
photoperiodic treatment conditions, five-six birds were collected shortly after lights on
(Zeitgeber time (zt) 0), and then every three hours for twenty-four-hour period. This

398 resulted in a high frequency daily sampling period that included zt0, zt3, zt6, zt9, zt12,
399 zt15, zt18, and zt21. Brain, pituitary gland, and liver was extracted and stored at -80°C.
400 Testes mass was determined as described above.
401
Study 3 – Endogenous programming of FSHβ in the quail pituitary gland
Male quail (N=24) were housed in LD (16L:8D) photoperiod for 2 days, and then day
length was decreased to 12hr for one day, and then 8L for six weeks. A subset of quail
was killed by decapitation followed by exsanguination and established the photoregressed
8L group (n=6). A subset of birds (n=6) was maintained in short day photoperiods for four
more weeks (8Lext). This group of birds provided the ability to examine whether an
endogenous increase in *FSHβ* expression would occur in constant short day photoperiod
condition. The other twelve birds were transitioned to the 10L:14D (10v) light treatment for
2 weeks and another subset of birds were collected (n=6). The last subset of birds was
transitioned to 12L:12D (12v) for two weeks and were then killed (n=6). For all birds, the
brain and pituitary glands were dissected and immediately frozen in dry ice and then
placed at -80°C. This experimental design provided the ability to examine endogenous
changes in pituitary cell function via the 8Lext group, and the photoinduced increase in
photosensitivity (i.e., 10v and 12v).
416
Hypothalamic and pituitary dissection
During brain extraction the pituitary stalk was severed. The procedure leaves pituitary
gland components of the pars intermedia, pars distalis and pars nervosa resting in the
sphenoid bone. To isolate the anterior hypothalamus/preoptic area and the mediobasal
hypothalamus, we used a brain matrix and coordinates based on previously published
anatomical locations (Stevenson et al., 2012b; Nakao et al., 2008, respectively). Brains
were placed ventral surface in an upward direction. For the anterior hypothalamus/preoptic
area a 2mm diameter from 2mm brain slice was collected. The rostral edge of the optic
nerve was identified and then a 1mm cut in the rostral and 1mm cut in the caudal direction
was performed. Brain slices were checked to confirm the presence of the tractus
septomesencephalicus in the rostral section and the decussation supraoptica dorsalis in
the caudal section. These anatomical regions reliably capture the GnRH neuronal
population in birds (Stevenson and Ball, 2009). The MBH was isolated from a 2mm punch
from a 3mm brain slice that spanned the decessation supraoptica dorsalis to the nervus
oculomotorius (Nakao et al., 2008).
432
Ribonucleic acid extraction and qPCR
RNA was extracted from anterior hypothalamus/preoptic area, mediobasal hypothalamus
and pituitary gland tissue using QIAGEN RNesay Plus Mini Kit (Manchester, UK). Nucleic
acid concentration and 260/280/230 values were determined by spectrophotometry
(Nanodrop, Thermoscientific). cDNA was synthesized from 100ng RNA using SuperScript
III (Invitrogen), and samples stored at –20°C until quantitative PCR (qPCR) was
performed. qPCRs were conducted using SYBR Green Real-time PCR master mix with 5ul
cDNA and 10ul SYBR and primer mix. All samples were run in duplicate. PCR primer
sequences and annealing temperatures are described in Table S12. qPCRs for mRNA
expression in tissue were performed using Stratagene MX3000. qPCR conditions were an
initial denature step at 95°C for 10 min. Then, 40 cycles of a denature at 95°C for 30
seconds, a primer specific annealing temperature for 30 seconds, and then an extension
phase at 72°C for 30 seconds. The qPCR reaction was terminated at 95°C for one-minute.
A melt curve assay was included to confirm specificity of the reactions. The efficiency and
cycle thresholds for each reaction was calculated using PCR Miner (Zhao and Fernald,
2005). All samples were assessed based on the Minimum Information for Publication of
Quantitative Real-Time PCR guidelines (0.7-1.0; Bustin et al., 2009). Actin and

glyceraldehyde 3-phosphate dehydrogenase were used as the reference transcripts. The
most stable reference transcript was used to calculate fold change in target gene
expression.
453
Minion transcriptome sequencing
RNA for transcriptome sequencing was extracted using QIAGEN RNeasy Plus Mini Kit
(Manchester, UK). RNA concentration and 260/280/230 values were determined by
Nanodrop spectrophotometer. Isolated RNA reliably has RIN values >9.0 for both the
mediobasal hypothalamus and pituitary gland. RNA was synthesized into cDNA using
Oxford Nanopore Direct cDNA Native Barcoding (SQK-DCS109 and EXP-NBD104) and
followed the manufacturers protocol. A total of 6 Spot-ON Flow cells (R9 version FLO-
MIN106D) were used for each tissue. A single quail was randomly selected from each
treatment group so that a single Flow cell had n=9 samples giving a total of N=56 quail for
mediobasal hypothalamus and pituitary stalk transcriptome sequencing. Transcriptome
sequencing was conducted using MinION Mk1B (MN26760, Oxford Nanopore
Technologies). Sequencing was performed by MinKNOW version 20.10.3 and Core 4.1.2.
The parameters for each sequencing assay were kept to 48hrs, -180mV voltage and fast5
files saved in a single folder for downstream bioinformatic analyses.
468
Transcriptome Analyses
The transcriptome data analysis pipeline is outlined in Supplementary Data Fig. S8. All
bioinformatic steps were conducted using R Studio and run in a Conda environment. First,
fast5 files were demultiplex and basecalled by Guppy 4.2.1. Then Porechop v0.2.4 was
conducted to remove adapters from reads followed by Filtlong v0.2.0 to filter long reads
with minimum 25 bases and mean q weight of 9. Transcripts were aligned to the Japanese
quail reference genome and transcriptome using Minimap2 v2.17 (Li, 2018). Transcript
expression levels were determined using Salmon v0.14.2 and EdgeR v3.24.3 for
normalisation and differential expression (Patro et al., 2017). DAVID was conducted to
identify functional pathways active during the transitions across photoperiodic states
(Table S2 and Table S4) (Dennis, 2003).
480
BioDare2.0 analyses of significant differentially expressed genes
To identify seasonal rhythmic expression of transcripts, we selected differentially
expressed genes identified by EdgeR (P<0.05). Data were analysed using non-linear
regression for rhythmicity using the online resource BioDare 2.0 (Zielinksi et al., 2014)
(biodare2.ed.ac.uk). The empirical JTK_CYCLE method was used for detection of
rhythmicity and the classic BD2 waveform set was used for comparison testing. The type
of transcript rhythmicity was confirmed as (e.g., sine/cos/arcsine) or non-rhythmic (spike)
expression. Rhythmicity was determined by a Benjamini-Hochberg controlled p-value (BH
corrected p < 0.1). Data for heatmaps were clustered using PAM clustering from the
cluster package (29). Heatmaps were created using the Complexheatmaps package (Gu
et al., 2016). Heat maps generated from statistically significant transcripts are presented in
Fig.1 and Table S3 and Table S5.
493
Enrichment factor identification in promoters of differentially expressed genes
To explore potential upstream molecular pathways involved in the regulation of
differentially expressed genes, we conducted a transcription factor analyses. The aim was
to identify which transcription factor or factors are responsible for observed changes in
gene expression and whether, if any overlap occurs across tissues. Transcription factor
enrichment analysis was achieved using ChIP-X Enrichment Analysis 3 (ChEA3) (Keenan
et al., 2019). We used a conservative approach and only used transcripts detected as

significant by BioDare 2.0 analysis (BH corrected p < 0.05). Enriched transcription factors
were ranked using the ENCODE database and presented in Table S7 and Table S8.
503
Weighted gene co-expression network analyses
Co-expression networks were established using WGCNA package in R (Langfelder and
Horvath, 2008). Raw data from pituitary and MBH sequencing was filtered to remove lowly
expressed transcripts identified using EdgeR. Data was assessed for outliers and values
were excluded. The data also was assessed for scale independence and mean
connectivity, and a power threshold of 5 was selected. The WGCNA package was used to
construct a weighted gene network, with a merging threshold of 0.25. Module-trait
relationship associations were used to identify relationships with measured physiological
data. Data for the analyses are provided in Table S6.
513
Daily waveform analyses of transcript expression
To establish rhythmicity in daily waveform expression of MBH and pituitary transcripts, we
conduct cosinor analyses (Cornelissen, 2014). 2-ΔΔ Cycling time (Ct) values obtained for
the genes investigated in Study 2 were subjected to cosinor analyses based on unimodal
cosinor regression $[y = A + (B \dot{c} \cos(2\pi(x - C)/24))]$, where A. B, and C denote the
mean level (mesor), amplitude, and acrophase of the rhythm, respectively. The
significance of regression analysis determined at $p < 0.05$ was calculated using the
number of samples, R2 values, and numbers of predictors (mesor, amplitude, and
acrophase) (Singh et al., 2013). Data are compiled in Table S9.
523
Bioinformatic analyses of FSHβ promoter
To identify potential links between transcription factors identified using bioinformatic tests,
and transcriptome data, we examined binding motifs in the FSHβ promoter. The upstream
promotor sequence of 3500 bp (-3500 to 0) of Japanese quail FSHβ was obtained from
Ensemble (http://www.ensembl.org/index.html). This promotor sequence was analysed by
CiiDER transcription binding factor (TBFs) analysis tool (Gearing et al., 2019) against
JASPER core database. DNA motifs in the promoter were unique for 470 transcription
binding factors. The top 80 transcription binding factors were then subjected to PANTHER
gene ontology enrichment analyses (Mi et al., 2013). Several pathways were discovered
and included hormone responsive, epigenetic and responsive to nutrients (Table S10).
534
Immunocytochemistry and histological analyses
Snap frozen brains from experiment 1 were sectioned at 20µm using a cryostat (CM1850,
Leica). The tissues sections were collected in supercharged slides (631-0108, VWR) and
stored in -80c till processed for staining. We used mouse monoclonal antibody
(OAAEE00561, Aviva systems) raised against the Human vimentin gene (NCBI Reference
Sequence: NP_003371.2). The antibody has been shown to specifically bind with vimentin
expressed in avian cells (https://www.avivasysbio.com/vim- antibody-oaee00561.html).
Human and quail vimentin peptide show high similarity of 86.96% similarity. *In silico*
analyses using BLAST confirmed that the antibody sequence is specific to vimentin. The
next closest protein had 68% similarity (i.e., desmin) which is expressed in cardiac and
skeletal muscle. We used Goat Anti-mouse Alexa Flour488 (A11001, Invitrogen) for the
secondary antibody. As negative control procedures, we performed omission of primary
antibody and omission of secondary antibody.
Immunocytochemistry was performed using the standard immunofluorescence
protocol (Majumdar et al., 2015) with minor modifications. Briefly, sections (brain) in slides
were first enclosed in margin using ImmEdge pen (H-4000, Vector Labs). The sections
were then first post fixed in 10% neutral buffered formalin (5735, ThermoScientific) for 4
hours. After fixing, the sections were washed (3 times; 5 min each) with TBS (phosphate

buffer saline with 0.2% triton). Then they were blocked in 20% Bovine serum albumin (BSA) in TBS for 1 hour at room temperature (RT). Subsequently the blocking solution was removed by pipetting and the sections were incubated with primary antibody (1:300 dilution) for 2 hours at RT and finally overnight at 4°C. The next day, sections were first washed with TBS (3 times; 5 min each) and then incubated with secondary antibody (1:200 dilution) for 2 hours at RT. Finally, the sections were again washed with TBS (3 times; 5 min each) and mounted in Fluromount-G mounting media with DAPI (004959-52, Invitrogen). The dried slides were visualized using Leica DM4000B fluorescence microscope equipped with Leica DFC310 FX camera. Leica Application Suite (LAS) software was used for image acquisition. All the images were taken at constant exposure for the FITC channel at 10X and 20X magnification.

For analysis and quantification of % area, ImageJ version 1.53j was used. For this, 20X images were first converted into greyscale images (8 bit) and a threshold applied. The threshold was determined using the triangle method on multiple randomly selected images and applied for all the images. The scale of measurement in ImageJ was then set to CM in 300pixels/cm scale. A region of interest in the median eminence and dorsal 3rdV ependymal layer was specified as 300*500 pixels (1*1.67 cm scaled units) and area fraction was measured. At least 3 images from each animal were measured and averaged for each bird. A total of 27 quail were used and distributed evenly across photoperiod treatments (n=3).

Statistical analyses and figure presentation
GLM tests were conducted to test for statistical significance. One-way ANOVA with Bonferroni correction was applied to testes mass, body mass, fat score, vimentin immunoreactivity and qPCR analyses in Study 1. Two-way ANOVAs with photoperiod and zeitgebers main effects were conducted on qPCR analyses in Study 2. One-way ANOVAs were conducted for qPCR data in Study 3. qPCR data were log-transformed if violation of normality was detected. Significance was determined at $P<0.05$. Figures were generated using AdobeIllustrator and BioRender was used to create images in panels Fig1d,n; Fig3e and Fig. S7.

Acknowledgements: The authors thank Elisabetta Tolla, Christopher Elcombe, Ana Monteiro and David Hamilton for their assistance. The authors thanks Professor Gregory Ball and Neil Evans for comments on a previous version of the paper. Funding: the work was funded by a Leverhulme Trust Research Leader to TJS.

References

Ball GF (1993) **The neural integration of environmental information by seasonally breeding birds.** *Am. Zool.* **33**:185–199.

Ball GF, Ketterson ED (2008) **Sex differences in the response to environmental cues regulating seasonal reproduction in birds.** *Philos Trans R Soc Lond B Biol Sci.* **363**:231-246.

Bertin A, Houdelier C, Richard-Yris MA, Guyomarc'h C, Lumineau S (2009) **Stable**

individual profiles of daily timing of migratory restlessness in European quail.

Chronobiol Int. **24**:253-267.

Bradshaw WE, Holzapfel CM (2007) **Evolution of animal photoperiodism.** *Annual Rev.*

Ecol. Evol. System. **38:**1-25.

Bustin SA, Benes V, Garson JA, Hellemans J, Huggett J, Kubista M, Mueller R, Nolan T,

Pfaffl MW, Shipley GL, Vandesompele J, Wittwer CT (2009) **The MIQE guidelines:**

minimum information for publication of quantitative real-time PCR experiments.

Clin. Chem. **55**:611-622.

Caro SP, Lambrechts MM, Chastel O, Sharp PJ, Thomas DW, Balthazart J (2006)

Simultaneous pituitary–gonadal recrudescence in two Corsican populations of

male blue tits with asynchronous breeding dates. *Horm. Behav.* **50**:347–360.

Cornelissen G (2014) **Cosinor-based rhythmometry.** *Theoretical Biology Medical*

Modelling **11**:16.

Dawson A, King VM, Bentley GE, Ball GF (2001) **Photoperiodic control of seasonality**

in birds *J Biol Rhythms* **16**:365-380.

Dennis G (2003) **DAVID: Database for Annotation, Visualization, and Integrated**

Discovery. *Genome Biol* **4**:R60.

Dorst J (1956) **La migration des oiseaux** *Payot Paris* p 422.

Duncan S, Holm S, Questa J, Irwin J, Grant A, Dean C (2015) **Seasonal shift in timing of**

vernalization as an adaption to extreme winter. *eLife* **4**:e06620.

Farner DS, Follett BK, King JR, Morton ML (1966) **A quantitative examination of ovarian**

growth in white-crowned sparrow. *Biol. Bull.* **130**:67–75.

Follett BK, Sharp PJ (1969) **Circadian rhythmicity in photoperiodically induced**

gonadotropin release and gonadal growth in the quail. *Nature* **223**:968-971.

Gearing LJ, Cumming HE, Chapman R, Finkel AM, Woodhouse IB, Luu K, Gould JA,

Forster SC, Hertzog PJ (2019) **CiiiDER: A tool for predicting and analysing**

transcription factor binding sites. *PLoS One* **14**:e0215495.

Gu Z, Eils R, Schlesner M (2016) **Complex heatmaps reveals patterns and correlations**

in multidimensional genomic data. *Bioinformatics* **32**:2847-2849.

Gwinner E (1986) ***Circannual Rhythms: endogenous annual clocks in the***

organization of seasonal processes. Springer-Verlag.

Gwinner E, Dittami J (1990) **Endogenous reproductive rhythms in a tropical bird.**

Science **249**:906-908.

Helfer G, Stevenson TJ (2020) **Pleiotropic effects of proopiomelanocortin and VGF**

nerve growth factor inducible neuropeptides for the long-term regulation of

energy balance *Mol Cell Endocrinol.* **514**:110876

Helm B, Stevenson TJ (2014) **Circannual rhythms: history, present challenges, future**

directions. In *Annual, lunar and tidal clocks*. pp. 203-225.

Keenan AB, Torre D, Lachmann A, Leong AK, Wojciechowicz ML, Utti V, Jagodnik KM,

Kropiwnicki E, Wang Z, Ma'ayan A (2019) ChEA3: **Transcription factor enrichment**

analysis by orthogonal omics integration. *Nucleic Acids Res* **47**:W212–W224.

King VM, Bentley GE, Follett BK (1997) **A direct comparison of photoperiodic time**

measurement and the circadian system in European starlings and Japanese

quail. *J. Biol. Rhythms* **12:**431-442.

Langfelder P, Horvath S (2008) **WGCNA: an R package for weighted correlation**

network analysis. *BMC Bioinformatics* **9**: 559.

Li H (2018) **Minimap2: pairwise alignment for nucleotide sequences.** *Bioinformatic*

34:3094-3100.

Liddle TA, Stevenson TJ, Majumdar G (2022) **Photoperiodic regulation of avian physiology: from external coincidence to seasonal reproduction.** *J. Exp. Zool. Ecol. Inte. Physio.* **337**:890-901.

Lincoln GA, Clarke IJ, Hut RA, Hazlerigg DG (2006) **Characterizing a mammalian circannual pacemaker.** *Science*, **314**:1941-1944.

Maechler M (2019) **Finding groups in data": Cluster analysis extended Rousseeuw et al.** R package version, 2(0).

Majumdar G, Yadav G, Rani S, Kumar V (2015) **Bird eyes distinguish summer from winter: Retinal response to acute photoperiod change in the night-migratory redheaded bunting.** *J Chem Neuroanat.* **68**:55-60.

Mi H, Muruganujan A, Thomas PD (2013) **PANTHER in 2013: modeling the evolution of gene function, and other gene attributes, in the context of phylogenetic trees.** *Nucleic Acids Res* **41**:D377–D386.

Nakao N, Ono H, Yamamura T, Anraku T, Takagi T, Higashi K, Yasuo S, Katou Y, Kageyama S, Uno Y, Kasukawa T, Iigo M, Sharp PJ, Iwasawa A, Suzuki Y, Sugano S, Niimi T, Mizutani M, Namikawa T, Ebihara S, Ueda HR, Yoshimura T (2008) **Thyrotrophin in the pars tuberalis triggers photoperiodic response.** *Nature* **452**:317-322.

Patro R, Duggal G, Love MI, Irizarry R, Kingsford C (2017) **Salmon: fast and bias-aware quantification of transcript expression using dual-phase inference**. *Nat. Methods.* **14**:417-419.

Pengelley ET, Fisher KC (1957) **Onset and cessation of hibernation under constant temperature and light in the golden-mantled ground squirrel (Citellus lateralis).** *Nature* **180:**1371-1372.

Perez JH, Tolla E, Dunn IC, Meddle SL, Stevenson TJ (2019) **A comparative perspective on extra-retinal photoreceptor.** *Trends Endocrinol. Metab.* **30**:39-52.

Prendergast BJ, Kriegsfeld LJ, Nelson RJ (2001) **Photoperiodic polyphenisms in**

rodents: neuroendocrine mechanisms, costs, and functions. *Q Rev Biol* **76**:293-

325.

Robinson JE, Follett BK (1982) **Photoperiodism in Japanese quail: the termination of**

seasonal breeding by photorefractoriness. *Proc R Soc B.* **215**:95-116.

Singh D, Rani S, Kumar V (2013) **Daily expression of six clock genes in central and**

peripheral tissues of a night-migratory songbird: evidence for tissue-specific

circadian timing. *Chronobiol Int.* **30**:1208-17.

Stevenson TJ, Ball GF (2009) **Anatomical localization of the effects of reproductive**

state, castration, and social milieu on cells immunoreactive for gonadotropin-

releasing hormone 1 in male European starlings. *J Comp. Neurol.* **517**:146-155.

Stevenson TJ, Lynch KS, Lamba P, Ball GF, Bernard DJ (2009) **Cloning of**

gonadotropin-releasing hormone I complementary DNAs in songbirds facilitates

dissection of mechanisms mediating seasonal changes in reproduction.

Endocrinol. **150**:1826-1833.

Stevenson TJ, Hahn TP, Ball GF (2012a) **Variation in gonadotrophin-releasing**

hormone-1 gene expression in the preoptic area predicts transitions in seasonal

reproductive state. *J. Neuroendocrinol.* **24**:267-274.

Stevenson TJ, Replogle K, Drnevisch J, Clayton DF, Ball GF (2012b) **High throughput**

analysis reveals dissociable gene expression profiles in two independent

neuronal systems involved in the regulation of social behavior. *BMC Neurosci.*

13:126.

Tolla E, Stevenson TJ (2020) **Sex differences and the neuroendocrine regulation of**

seasonal reproduction by supplementary environmental cues *Inte. Comp. Biol.*

60:1506-1516.

Visser ME, Gienapp P (2019) **Evolutionary and demographic consequences of**

phenological mismatches. *Nature Ecol Evol.* **3**:879-885.

Wingfield JC, Farner DS (1978) **The endocrinology of a naturally breeding population**

of the white-crowned sparrow (Zonotrichia leucophyrs gambelii). *Physiol. Zool.*

51:188-205.

Wingfield JC, Farner DS (1980) **Control of seasonal reproduction in temperate-zone**

birds. *Prog. Reprod. Biol.* **5**:62–101.

Wingfield JC (1980) **Fine temporal adjustment of reproductive functions.** In: Epple A,

Stetson MH, editors. *Avian endocrinology.* Academic Press; New York, NY: pp. 367–

389

Wingfield JC, Hahn TP, Wada M, Astheimer LB, Schoech S (1996) **Interrelationship of**

day length and temperature on the control of gonadal development, body mass

and fat score in white-crowned sparrows, Zonotrichia leucophrys gambelii. *Gen.*

Comp. Endocrinol. **101**:242–255.

Wingfield JC, Hahn TP, Maney DL, Schoech SJ, Wada M, Morton ML (2003) **Effects of**

temperature on photoperiodically induced reproductive development, circulating

plasma luteinizing hormone and thyroid hormones, body mass, fat deposition

and molt in mountain white-crowned sparrows, Zonotrichia leucophrys oriantha.

Gen. Comp. Endocrinol. **131**:143–158.

Wingfield JC (2008) **Organization of vertebrate annual cycles: implications for control**

mechanisms. *Philos. Trans. R. Soc. B. Biol. Sci.* **363**:425–4.

Wood SH, Hindle MM, Mizoro Y, Cheng Y, Saer BRC, Miedzinska K, Christian HC, Begley

N, McNeilly J, McNeilly AS, Meddle SL, Burt DW, Loudon ASI (2020) **Circadian clock**

mechanism driving mammalian photoperiodism. *Nat Commun* **11**:4291.

Woodfill CJI, Wayne NL, Moenter SM, Karsch FJ (1994) **Photoperiodic synchronization**

of a circannual reproductive rhythm in sheep: identification of season-specific

time cues. *Biol. Reprod.* **50**:965-976.

Yamamura T, Hirunagi K, Ebihara S, Yoshimura T. (2004) **Seasonal morphological**

changes in the neuro-glial interaction between Gonadotropin-releasing hormone

nerve terminals and glial endfeet in Japanese quail. *Endocrinol.* **145**:4264-4267.

Yeo GH, Heisler LK (2012) **Unraveling the brain regulation of appetite: lessons from**

genetics. *Nat. Neurosci.* **15**:1343-1349.

Yoshimura T, Yasuo S, Watanabe M, Iigo M, Yamamura T, Hirunagi K, Ebihara S. (2003)

Light-induced hormone conversion of T4 to T3 regulates photoperiodic response

of gonads in birds. *Nature* **426**:178-181.

Zhao S, Fernald RD (2005) **Comprehensive algorithm for quantitative real-time**

polymerase chain reaction. *J. Comput. Biol.* **12**:1045-62.

Zielinski T, Moore AM, Troup E, Halliday KJ, Millar AJ (2014) **Strengths and Limitations**

of Period Estimation Methods for Circadian Data. *PLoS ONE* **9**:e96462.

Fig. 1 - Vernal increase in pituitary *FSHβ* precedes molecular switches in MBH and testes growth.

a, Schematic representation of the simulated annual rhythm in photoperiod. Quail were collected in 16hr light, 8hr dark photoperiod and then every 2 weeks the photoperiod was decreased by 2hrs to 14hr, 12hr, 10hr and then a 8L short photoperiod. Photoperiod was then increased to mimic the vernal transition and birds were collected at 10hr, 12hr, 14hr and 16hr light photoperiods. Testis volume confirmed critical day length (i.e. >12hr) induced growth.
b, Body mass and **c,** abdomenal fat deposition increased until the autumnal equinox (12a), and then increased during the vernal photoperiod transitions. **d,** Diagram highlighting hypothalamic preoptic area (POA), mediobasal hypothalamus (MBH) and pituitary gland. Tanycytes in the MBH gate GnRH release into the pituitary. **e,** Heat-map of RNA-seq of MBH punches identifies distinct waves of transcripts as quail transition across photoperiodic conditions. **f-g,** qPCR assays for proopiomelanocortin (*POMC*) and deiodinase type-3 (*DIO3*) confirmed restricted activation during 10v-8L and 8L-10v phases, respectively. **h-i,** vimentin immunoreactivity in the median eminence (ME) shows tanycyte morphology growth is limited to 10a, 8L, and 10v photoperiods. **j,** Heat-map illustrating photoperiodic transitions in pitutiary transcripts. **k-m,** confirmed that *FSHβ* is elevated under non-stimulatory photoperiods followed by increased prolactin (*PRL*) in 14v and thyrotropin stimulating hormone-β (*TSHβ*) in 16v. **n,** diagram summarizing that long photoperiods increase *GNRH* synthesis and release into the pituitary gland to stimulate *FSHβ* and induce testis growth. Transition to autumnal equinox phases results in reduced *FSHβ* expression and regressed testis. Prolonged exposure to short photoperiods inhibits *GNRH* expression, triggers tanycyte extension, maintains low *FSHβ* and regressed testis. Vernal transitions in photoperiod to the equinox results in resumption of *GNRH* and elevated *FSHβ* expression without testis growth. Data are mean +/- SEM, and residual dot plot. **a-c, f,g,k,m** One-way ANOVA with Bonferonni corrected Tukey's test for multiple comparisons. **h, l** One-way ANOVA with Tukey tests for significant pairwise comparison. Letters denote significant difference between photoperiod phase. Raw data available in Table S1.

Figure 2 - *FSHβ* is constitutively expressed during the vernal equinox.

a, schematic representation of four photoperiod treatment groups with arrows to indicate the daily sampling time. **b,** testes volume remained in a regressed non-functional state in autumnal equinox (12a), short photoperiod (8L) and vernal equinox (12v). Photoperiods that exceeded the critical day length (i.e.,>12hr) induced testes growth. **c,** Pituitary circadian clock gene *ARNTL1* maintained daily rhythmic expression waveforms across all photoperiods (P<0.001) , no significant difference between photoperiod treatment (*P*=0.42) **d,** *PER3* displayed a daily waveform across 12a, 8L, 12v and 16L groups (*P*<0.001) and was anti-phase compared to *ARNTL1*. No significance difference between photoperiod treatments (*P*=0.31) **e,** *FSHβ* expression was significantly higher in 12v compared to long photoperiods (16L; *P*<0.001), autumnal equinox (12a; *P*<0.001) and short photoperiod (8L; *P*<0.001), but was not rhythmic (*P*=0.66). **f,** Similarly, *PRL* was high in 16L compared to 12a (*P*<0.001), and 8L (*P*<0.001), there was no significant daily rhythm (*P*=0.52). **g,** *FSHβ* promoter was devoid of circadian gene binding D-box and E-box motifs but contains a series of hormone and nutrient responsive motifs. **b-f** Two-way ANOVA followed by Tukey's pairwise tests, rhythmic analyses were conducted using GraphPad Prism. ‡ indicates significant photoperiod treatment effect; # denotes significant time of day effect. Data are mean +/- SEM, and residual dot plot (a-d). Rhythmic analyses are presented in Table S7 and raw data are available in Table S1.

Figure 3 - Endogenous and light-induced *FSHβ* expression.

a, *FSHβ* expression increased during the photoinduced transition from 8L to 10v, and 12v. *FSHβ* also showned a smaller, yet significant increase in expression after prolonged exposure to 8L. Y-axis is presented in log-scale due to the significant increase in *FSHβ* expression in 10v and 12v. **b,** *OPN5* was detected in the pituitary gland and showed a significant increase in expression in the transition to 10v and 12v similar to *FSHβ* expression. **c,** *DIO3* was significantly reduced in 12v quail compared to all other treatment groups. **d,** *GNRH* expression remained constant during the transition from 8L to 12v. However, continued exposure to 8L was observed to increase *GNRH* expression. Data are mean +/- SEM, and residual dot plot (a-d). One-way ANOVA with Tukey's test for multiple comparisons. Letters denote significant difference between photoperiod phase. Raw data available in Table S1. **e,** Schematic representation of the endogenous and light-dependent increase in pituitary cell types during the transition from 8L to stimulatory 16v light treatments. Increased color indicates increased transcript expression.