Title: *FSHβ* links photoperiodic signalling to seasonal reproduction in Japanese quail.

Short Title: Annual timing in birds

Authors: Gaurav Majumdar, Timothy A Liddle, Calum Stewart, Christopher J Marshall, Maureen Bain, Tyler John Stevenson*

Affiliations: Institute of Biodiversity, One Health and Veterinary Medicine University of Glasgow, Glasgow, United Kingdom.

*Corresponding author:
Tyler John Stevenson
School of Biodiversity, One Health, and Veterinary Medicine
University of Glasgow
Glasgow, United Kingdom
e. tyler.stevenson@glasgow.ac.uk
p. 01413305793

Key words: Neuroendocrine, Photoperiod, Reproduction, Circannual, Pituitary,

This manuscript includes:
Main Text
Figure 1 to 3
Supporting Information Tables S1-S10
Supporting Information Figures S1-S6

Abstract

Endogenous programs in avian reproductive seasonal life history transitions have been known for over 30 years. Attempts to identify the neural and molecular substrates of photoperiodic time measurement have, to date, focussed on blunt changes in light exposure during a restricted period of photoinducibility. Here we report mediobasal hypothalamus (MBH) and pituitary gland specific transcript expression cascades associated with photoperiod-induced seasonal life history transitions. We identified multiple waves of transcript expression that were not synchronized across brain regions. Follicle-stimulating hormone-β (FSHβ) expression increased during the simulated vernal equinox, prior to photoinduced increases in prolactin, thyrotropin-stimulating hormone-β and testicular growth. Analyses showed sustained elevated levels of daily FSHβ expression in the pituitary under conditions of the vernal, compared to autumnal equinox, short (<12L) and long (>12L) photoperiods. FSHβ expression increased in Quail held in non-stimulatory photoperiod, indicative of an endogenous programmed change in transcription. These data identify that FSHβ provides an endogenous program for the photoperiod-dependent external coincidence timing for seasonal transitions in reproduction. FSHβ may prime the pituitary for subsequent stimulation by triiodothyronine-mediated changes in tanycytes during the photoinducible phase. Overall, the data indicate a multi-cellular, multi-neural interval timing mechanism resides in the brain to control seasonal life-history transitions in birds.

Introduction

Temperate and equatorial species display endogenous cycles in physiology and behavior that serve to anticipate future seasonal environments (Gwinner, 1986; Wingfield, 2008; Helm & Stevenson, 2014). Even in the absence of seasonal fluctuations in temperature, daylength and food availability, endogenous circannual programs in migration (Gwinner and Dittami, 1990), hibernation (Pengelley and Fisher, 1957), and reproduction (Woodfill et al., 1994; Lincoln et al., 2006) are maintained with remarkable temporal precision. The anatomical and cellular basis of circannual rhythms remain poorly characterized, but current evidence indicates that, in mammals, interval timing may reside in pituitary lactotropes (Lincoln et al., 2006) or thyrotropes (Wood et al., 2020). In most long-lived species (e.g., >2 yrs.), the annual change in daylength, referred to as photoperiod, acts to entrain endogenous circannual calendar cells to time physiological state to seasons (Bradshaw and Holzapfel, 2007). However, our understanding of how the brain integrates photoperiodic cues and endogenous rhythms to define the transitions between seasonal life-history states remains elusive.

In most birds, reptiles, and amphibians, annual changes in daylength are detected by photoreceptors located in the mediobasal hypothalamus (MBH) (Perez et al., 2019). In Japanese quail (*Coturnix japonica*), days that exceed a 'critical length' that coincides with a period of photoinducibility (Follett and Sharp, 1969) trigger a molecular cascade that starts with the upregulation of thyrotropin-stimulating hormone beta subunit (*TSHβ*) in the pars tuberalis of the pituitary gland and results in gonadal maturation (Nakao et al., 2008). The current gap in our knowledge is how short days (i.e., <12hr) induce endogenous programs in gonadal growth (Follett and Sharp, 1969) and how prolonged exposure to short days sensitises the brain to respond, at a molecular level, to stimulatory long days. Previous studies in European starlings (*Sturnus vulgaris*) demonstrated that exposure to short days increased Gonadotropin-releasing hormone (*GNRH*) expression in the preoptic

area (Stevenson et al., 2009; Stevenson et al., 2012a). As both GnRH and gonadal growth

increased in the absence of stimulatory long day lengths, another cellular pathway must

underlie the endogenous development of reproductive physiology in birds. We theorized

multiple neuroendocrine regions including the preoptic area, mediobasal hypothalamus

and pituitary cells are independently involved in the timing of seasonal transitions in

physiology. Specifically, we hypothesized that pituitary gonadotropes establish a state of

photosensitivity to stimulatory long day photoperiods and thus act as calendar cells that

provide endogenous timing of gonadal growth.

Results

To address this problem, we simulated the photoperiodic regulation of seasonal

physiology of Japanese quail, using sequential changes of an autumnal decrease,

followed by a vernal increase in daylength and measured testes volume, body mass and

abdominal fat. As anticipated (Follett and Sharp, 1969; Robinson and Follett, 1982), light

phases that exceed the critical day length (i.e., >12hr) resulted in robust gonadal growth

(Fig. 1a). Increases in body mass and abdominal fat deposition were delayed until the

vernal increase in day lengths reached 10hrs (10v) (Fig. 1b,c). EdgeR analyses, of MBH

sequences obtained using Minion, identified 1481 transcripts were differentially expressed

(P<0.05) (Table S2), and BioDare2.0 established 398 have rhythmic patterns (Fig. 1e;

Table S3). DAVID gene ontology analyses indicated that gonadotropin-releasing hormone

receptor and Wnt signalling pathways were consistently identified as the predominant

cellular mechanism recruited during each photoperiodic transition (Table S2). Increased

proopiomelanocortin (*POMC*) expression coincided with body mass growth and abdominal

fat accumulation (Fig. 1f). Thyroid hormone catabolism enzyme deiodinase type-3 (*DIO3*)

increased after prolonged exposure to non-stimulatory photoperiods and was only

therefore transiently elevated from 8L to 10v (10v; Fig. 1g). Highest levels of vimentin

immunoreactivity in the median eminence coincided with the peak in *DIO3* expression (Fig.

1h) suggesting the localized removal of active thyroid hormone is limited to a short phase

and occurred prior to stimulatory photoperiods (Fig. 1i). The photoperiod-induced change

in vimentin was anatomically localized to the median eminence, as the area of

immunoreactivity in the dorsal 3^{rd}V ependymal layer did not change with seasonal

transitions in reproduction (Fig. S1). Next, we used MinION to sequence transcripts in the

pituitary gland across the seasonal transitions in reproduction. EdgeR analyses identified

3090 transcripts were differentially expressed in the pituitary gland (Table S1). DAVID

gene ontology analyses identified that gonadotropin-releasing hormone receptor and

epidermal growth factor pathways are consistently observed across photoperiodic

transitions (Table S4). BioDare2.0 established that 130 transcripts were rhythmically

expressed (Fig. 1j; Table S5). A remarkable 96.5% (384/398) of MBH transcripts show a

photoperiod-induced spiked patterned and only 12/14 remaining transcripts displayed sine

waveforms. Conversely, 100% of pituitary transcripts conformed to sine (124/130) or

cosine (6/130) waveforms. The predominant rhythmic patterns of expression likely reflect

endogenous long-term molecular programs, characteristic of calendar cell function. *GNRH*

expression in the preoptic area was temporarily decreased during the 8L short photoperiod

indicating another nucleus specific cellular timer is present in this brain region (Fig. S1).

Overall, the pituitary showed a distinct transcriptomic profile compared to the MBH

suggesting independence in the representation of seasonal photoperiodic timing. The

vernal photoperiodic transition from 8L to 12v resulted in a significant increase in *FSHβ*

(Fig. 1k) without a change in tanycyte restructuring (Fig. 1h). Subsequent transition to 14v

and then 16v resulted in the upregulation of prolactin (*PRL*) (Fig. 1l) and thyrotropin-

stimulating hormone-β (*TSHβ*), respectively (Fig. 1m). These data establish that multiple

cells in the pituitary code seasonal photoperiodic time and *FSHβ* shows an endogenous

increase in expression prior to reproductively stimulatory photoperiods.

133   To ascertain common molecular mechanisms involved in the transcriptional

134 regulation of photoperiodically regulated transcripts, transcription factor enrichment

135 analysis was conducted on significant MBH (Table S6) and pituitary gland (Table S7)

136 transcripts. Association plots show no overlap in DNA binding motifs between MBH and

137 pituitary transcripts (Fig. S2) suggesting tissue-specific transcription binding factor

138 regulation. Within the pituitary gland, several common transcription factors, such as the

139 Jun proto-oncogene, Spi1 proto-oncogene and myocyte enhancer factor 2 (*MEF2a*) might

140 be actively involved in the photoperiodic regulation of transcript expression. These findings

141 indicate multiple transcription factors are likely recruited to control tissue-specific, and cell-

142 specific transcript expression and seasonal life history transitions in physiology.

143   To establish whether increased pituitary *FSHβ* during the vernal 12hr transition

144 reflects constitutively elevated expression or is driven by the sampling 'time of day', we

145 collected tissue samples every 3 hours from quail that transitioned to the autumnal

146 equinox (i.e., 12a), the short photoperiod (8L), vernal equinox (12v) and long photoperiod

147 (16L) seasonal phases (Fig. 2a). Photoinduced changes in testes volume confirmed

148 seasonal reproductive condition (Fig. 2b). Circadian clock genes aryl hydrocarbon receptor

149 nuclear translocator like (*ARNTL1*) and period 3 (*PER3*) exhibit robust anti-phase daily

150 waveforms in expression, in the pituitary gland (Fig. 2c,d) (Table S8). Only *ARNTL1*, but

151 not *PER3* nor *DIO2*, had a rhythmic waveform in the MBH (Table S8; Fig. S3). Consistent

152 with the previous study, *FSHβ* expression was higher at the vernal equinox compared to

153 all other photoperiod groups (Fig. 2e). *PRL* expression was higher in long photoperiod

154 (16L) compared to the two equinox and short photoperiod (8L) (Fig. 2f). *FSHβ* did not

155 display a daily rhythm (Table S8), which is likely due to the absence of D-box and E-box

156 motifs in the *FSHβ* promoter (Fig. 2g). *FSHβ* promoter does contain many DNA motifs that

157 are targeted by several transcription factors that are responsive to hormonal and nutrient

158 pathways indicating multiple upstream regulators are recruited to drive transcription. These

data support the conjecture that a long- term programmed increase in *FSHβ* occurs under

vernal non-stimulatory photoperiod, and it is not driven by short-term daily photic cues.

To identify if *FSHβ* expression is driven by an endogenously programmed

mechanism or in response to the gradual increase in light, adult quail were exposed to the

8L, 10v or 12v light schedules or kept in 8L for an additional 4 weeks (8Lext). The 8Lext

treatment permitted confirmation whether *FSHβ* expression would increase in that

photoperiod, and therefore reflect an internal timing mechanism. *FSHβ* expression

increased 19-fold after 4 additional weeks of 8L suggesting that endogenous drivers

initiate transcription despite no change in daylength ($P<0.05$) (Fig. 3a). But the dominant

stimulator of *FSHβ* expression was the transition to 10v and 12v photoperiod, which both

expressed significantly increased levels compared to 8L ($P<0.001$). As Opn5 was

identified in the pituitary transcriptome (Table S1), we then assessed its transcript

expression across the photoperiod treatments. We discovered that Opn5 expression

patterns paralleled *FSHβ* suggesting the potential for direct light detection by Opn5 and

subsequent regulation of *FSHβ* expression. Based on *FSHβ* promoter analyses (Fig. 2g),

we examined the expression of myocyte enhancer factor 2 (*MEF2a*) expression as a

potential upstream regulator of *FSHβ* expression. *MEF2a* remained relatively constant

suggesting this transcription factor binding protein is not the primary driver of *FSHβ*

expression (Fig. S4). Similarly, *DNMT3a* expression did not change across photoperiod

treatments (Fig. S4) suggesting that epigenetic modifications (i.e., DNA methylation) may

not provide the endogenous programmed change leading to constitutive *FSHβ* expression.

The precise molecular change upstream from *FSHβ* transcription remains to be identified.

In the MBH, *DIO3* displayed a rapid reduction in expression after transfer to 10v and was

found to be significantly reduced in 12v photoperiod ($P<0.001$) (Fig. 3c). There was a

significant difference in MBH *DIO2* expression (Fig. S4), but this observation was driven by

a decrease in 8ext. Interestingly, *GNRH* in the preoptic area was found to significantly

increase after extended exposure to short photoperiods (Fig. 3D). These data indicate that

endogenous switches in *FSHβ,* and possibly *GNRH* expression, in response to short

photoperiods may reflect multiple independent cellular timers that establish a physiological

state of photosensitivity.

Discussion

The data reported herein demonstrate that that multiple independent interval timers

are required for the control of seasonal life-history transitions in reproductive physiology

(Fig. 3e). The two cell types primarily implicated are thyrotropes and gonadotropes. Short

winter days initiate a gradual increase in *FSHβ* expression in the pars distalis and maintain

low *TSHβ* in the pars tuberalis. As photoperiods increase there is a steady elevation in

FSHβ expression, however, the release of FSH is prevented as daylengths are below the

critical threshold (i.e., 12L:12D) (Gwinner, 1986). Stimulatory daylengths longer than 12hr

trigger thyrotropes to increase *TSHβ* leading to a cascade of molecular events in the MBH

that permit GnRH to stimulate gonadotropes to release FSHβ and initiate gonadal

development (Nakao et al., 2008; Yoshimura et al., 2003; Yamamura et al., 2004). Note

that other pituitary cell types, somatotropes and corticotropes do not appear to show any

molecular switches across the photoperiodic phases. These findings uncover a two-

component mechanism for the cellular basis of the external coincidence model for the

avian photoperiodic response (Fig. S5). The endogenously programmed increase in *FSHβ*

establishes a state of photosensitivity to stimulatory day length, and *TSHβ* thryotropes in

the pars tuberalis monitor daylength and when light stimulation occurs during a period of

photoinducibility, initiate gonadal development. The two-component model is exciting as it

accommodates evidence for endogenous growth of quail gonads in the absence of

photostimulation (Follett and Sharp, 1969). Moreover, a TSH-independent programmed

change in *FSHβ* expression addresses how seasonal rhythms in tropical, non-photoperiod

birds can be regulated (Gwinner and Dittami, 1990).

The high-dimensionality and high-frequency analyses of seasonal transition in

physiology used in this study, also uncovered that photoperiod differentially regulates two

endocrine systems: reproduction and energy balance. *POMC* has well described roles in

the neuroendocrine regulation of food intake and body mass (Yeo and Heisler, 2012). Our

findings reveal that photoperiod induced seasonal transitions in reproduction and body

mass are not parallel. Based on these results we propose that a separate light-dependent

mechanism independently regulates reproduction and energy balance via the *GNRH* and

POMC neuronal systems, respectively. This has significant implications for understanding

species-specific seasonal transitions in life-histories. The comprehensive transcriptome

dataset generated in this study will facilitate ecological studies that seek to uncover the

molecular substrates which environmental cues, such as temperature, impact phenological

timing and mistiming in birds (Visser and Gienapp, 2019).

Materials and Methods

Data and code availability statement
All raw data are available in Extended data Table 1. R code used in Study 1 is freely
available and indicated in the methods reference list. The full R code used is available
upon request.

Animals
All Japanese quail were provided by Moonridge Farms, Exeter United Kingdom
(moonridgefarms.co.uk). Chicks were raised under constant light and constant heat lamp
conditions. From 5 weeks of age, birds were held in the Poultry facilities at the University
of Glasgow, Cochno Farm for the duration of the photoperiodic experiments. Food (50:50
mix of Johnston and Jeff, quail mix & Farm Gate Layers, poultry layers supplemented with
grit) and tap water was provided ad libitum. All procedures were in accordance with the
National Centre for the Replacement, Refinement and Reduction of Animals in Research
ARRIVE guidelines (https://www.nc3rs.org.uk/revision-arrive-guidelines). All procedures
were approved by the Animal Welfare and Ethics Review Board at the University of
Glasgow and conducted under the Home Office Project Licence PP5701950.

Study 1 – Photoperiod-induced transition in seasonal life history states
Male quail (N=108) were housed in a summer-like long-day (LD) photoperiod (16L:8D). To
mimic the autumnal decline and subsequent vernal increase in the annual photoperiodic

cycle, birds were exposure to a sequential change in day length from 16L, to 14L, to 12L, to 10L, to 8L, then back to 10L, to 12L, to 14L and lastly 16L (Fig.1a). Each photoperiod treatment lasted for 2 weeks to minimize the impact of photoperiodic history effects (Stevenson et al., 2012a). At the end of each photoperiodic treatment a subset of quail (n=12) was pseudo randomly selected for tissue collection. Birds were killed by cervical dislocation followed by jugular cut. A jugular blood sample was collected in 50ul heparin (Workhardt, UK) and stored at -20°C. Brain and pituitary stalk were rapidly dissected, frozen on powdered dry ice and stored at -80°C. Testes were dissected and weighed to the nearest 0.001g using the Sartorius microbalance (Sartorius, Germany). The length and width of the testes were measured using callipers and volumes were calculated using the equation for a spheroid ($4/3 \times 3.14 \times [L/2] \times [W/2]2$) (King et al., 1997). Body mass was measured using an Ohaus microscale to the nearest 0.1g. Fat score was assessed using the common scale developed by Wingfield and Farner (1978). The scale range is 0 to 5, which 0 is no visible fat and 5 is bulging fat bodies that exceed the abdominal fat pads.

Study 2 – Daily rhythms in molecular profiles during solstices and equinoxes
To investigate the daily molecular representation of summer-like and winter-like solstices and the autumnal and vernal equinoxes, Japanese quail (N=188) were subjected to the same photoperiodic treatments described in Study 1. A subset of quail was pseudo randomly selected after the autumnal 12L:12D (n=47), short day 8L:16D (n=48), vernal 12L:12D (n=46) and long day 16L:8D (n=47) treatment conditions. For each of the four photoperiodic treatment conditions, five-six birds were collected shortly after lights on (Zeitgeber time (zt) 0), and then every three hours for twenty-four-hour period. This resulted in a high frequency daily sampling period that included zt0, zt3, zt6, zt9, zt12, zt15, zt18, and zt21. Brain, pituitary gland, and liver was extracted and stored at -80°C. Testes mass was determined as described above.

Study 3 – Endogenous programming of FSHβ in the quail pituitary gland
Male quail (N=24) were housed in LD photoperiod for 2 days, and then day length was decreased to 12hr for one day, and then 8L for six weeks. A subset of quail was killed by Home Office approved Schedule 1 methods and established the photoregressed 8L group (n=6). To examine endogenous increase in FSHβ expression, a subset of birds (n=6) was maintained in SD photoperiods for four more weeks (8Lext). The other twelve birds were transitioned to the 10L:14D (10v) light treatment for 2 weeks and another subset of birds were collected (n=6). The last subset of birds was transitioned to 12L:12D (12v) for two weeks and were then killed (n=6). For all birds, the brain and pituitary glands were dissected and immediately frozen in dry ice and then placed at -80°C. This experimental design provided the ability to examine endogenous changes in pituitary cell function via the 8Lext group, and the photoinduced increase in photosensitivity (i.e., 10v and 12v).

Hypothalamic and pituitary dissection
During brain extraction the pituitary stalk was severed. The procedure leaves pituitary gland components of the pars intermedia, pars distalis and pars nervosa resting in the sphenoid bone. To isolate the anterior hypothalamus/preoptic area and the mediobasal hypothalamus, we used a brain matrix and coordinates based on previously published anatomical locations (Stevenson et al., 2012b; Nakao et al., 2008, respectively). Brains were placed ventral surface in an upward direction. For the anterior hypothalamus/preoptic area a 2mm diameter from 2mm brain slice was collected. The rostral edge of the optic nerve was identified and then a 1mm cut in the rostral and 1mm cut in the caudal direction was performed. Brain slices were checked to confirm the presence of the tractus septomesencephalicus in the rostral section and the decussation supraoptica dorsalis in the caudal section. These anatomical regions reliably capture the GnRH neuronal

298 population in birds (Stevenson and Ball, 2009). The MBH was isolated from a 2mm punch
299 from a 3mm brain slice that spanned the decessation supraoptica dorsalis to the nervus
300 oculomotorius (Nakao et al., 2008).
301
Ribonucleic acid extraction and qPCR
RNA was extracted from anterior hypothalamus/preoptic area, mediobasal hypothalamus
and pituitary gland tissue using QIAGEN RNesay Plus Mini Kit (Manchester, UK). Nucleic
acid concentration and 260/280/230 values were determined by spectrophotometry
(Nanodrop, Thermoscientific). cDNA was synthesized from 100ng RNA using SuperScript
III (Invitrogen), and samples stored at −20°C until quantitative PCR (qPCR) was
performed. qPCRs were conducted using SYBR Green Real-time PCR master mix with 5ul
cDNA and 10ul SYBR and primer mix. PCR primer sequences and annealing
temperatures are described in Extended Data Table S10. qPCRs for mRNA expression in
tissue were performed using Stratagene MX3000. qPCR conditions were an initial
denature step at 95°C for 10 min. Then, 40 cycles of a denature at 95°C for 30 seconds, a
primer specific annealing temperature for 30 seconds, and then an extension phase at
72°C for 30 seconds. The qPCR reaction was terminated at 95°C for one-minute. A melt
curve assay was included to confirm specificity of the reactions. The efficiency and cycle
thresholds for each reaction was calculated using PCR Miner (Zhao and Fernald, 2005).
All samples were assessed based on the Minimum Information for Publication of
Quantitative Real-Time PCR guidelines (Bustin et al., 2009).
319
Minion transcriptome sequencing
RNA for transcriptome sequencing was extracted using QIAGEN RNesay Plus Mini Kit
(Manchester, UK). RNA concentration and 260/280/230 values were determined by
Nanodrop spectrophotometer. RNA was synthesized into cDNA using Oxford Nanopore
Direct cDNA Native Barcoding (SQK-DCS109 and EXP-NBD104) and followed the
manufacturers protocol. A total of 6 Spot-ON Flow cells (R9 version FLO-MIN106D) were
used for each tissue. A single quail was randomly selected from each treatment group so
that a single Flow cell had n=9 samples giving a total of N=56 quail for mediobasal
hypothalamus and pituitary stalk transcriptome sequencing. Transcriptome sequencing
was conducted using MinION Mk1B (MN26760, Oxford Nanopore Technologies).
Sequencing was performed by MinKNOW version 20.10.3 and Core 4.1.2. The parameters
for each sequencing assay were kept to 48hrs, -180mV voltage and fast5 files saved in a
single folder for downstream bioinformatic analyses.
333
Transcriptome Analyses
The transcriptome data analysis pipeline is outlined in Extended Data Figure S6. All
bioinformatic steps were conducted using R Studio and run in a Conda environment. First,
fast5 files were demultiplex and basecalled by Guppy 4.2.1. Then Porechop v0.2.4 was
conducted to remove adapters from reads followed by Filtlong v0.2.0 to filter long reads
with minimum 25 bases and mean q weight of 9. Transcripts were aligned to the Japanese
quail reference genome and transcriptome using Minimap2 v2.17 (Li, 2018). Transcript
expression levels were determined using Salmon v0.14.2 and EdgeR v3.24.3 for
normalisation and differential expression (Patro et al., 2017). DAVID was conducted to
identify functional pathways active during the transitions across photoperiodic states
(Extended Data Table 2 and Table 4) (Dennis, 2003).
345
BioDare2.0 analyses of significant differentially expressed genes
To identify seasonal rhythmic expression of transcripts, we selected differentially
expressed genes identified by EdgeR (P<0.05). Data were analysed using non-linear
regression for rhythmicity using the online resource BioDare 2.0 (Zielinksi et al., 2014)

(biodare2.ed.ac.uk). The empirical JTK_CYCLE method was used for detection of
rhythmicity and the classic BD2 waveform set was used for comparison testing. The type
of transcript rhythmicity was confirmed as (e.g., sine/cos/arcsine) or non-rhythmic (spike)
expression. Rhythmicity was determined by a Benjamini-Hochberg controlled p-value (BH
corrected p < 0.1). Data for heatmaps were clustered using PAM clustering from the
cluster package (29). Heatmaps were created using the Complexheatmaps package (Gu
et al., 2016). Heat maps generated from statistically significant transcripts are presented in
Fig1 and Extended Data Table 3 and Table 5.
358
Enrichment factor identification in promoters of differentially expressed genes
To explore potential upstream molecular pathways involved in the regulation of
differentially expressed genes, we conducted a transcription factor analyses. The aim was
to identify which transcription factor or factors are responsible for observed changes in
gene expression and whether, if any overlap occurs across tissues. Transcription factor
enrichment analysis was achieved using ChIP-X Enrichment Analysis 3 (ChEA3) (Keenan
et al., 2019). We used a conservative approach and only used transcripts detected as
significant by BioDare 2.0 analysis (BH corrected p < 0.05). Enriched transcription factors
were ranked using the ENCODE database and presented in Extended Data Table 6 and
Table 7.
369
Daily waveform analyses of transcript expression
To establish rhythmicity in daily waveform expression of MBH and pituitary transcripts, we
conduct cosinor analyses (Cornelissen, 2014). 2-ΔΔ Cycling time (Ct) values obtained for
the genes investigated in Study 2 were subjected to cosinor analyses based on unimodal
cosinor regression $[y = A + (B\dot{c}cos(2\pi(x - C)/24))]$, where A. B, and C denote the mean
level (mesor), amplitude, and acrophase of the rhythm, respectively. The significance of
regression analysis determined at p < 0.05 was calculated using the number of samples,
R2 values, and numbers of predictors (mesor, amplitude, and acrophase) (Singh et al.,
2013). Data are compiled in Extended Data Table 8.
379
Bioinformatic analyses of FSHβ promoter
To identify potential links between transcription factors identified using bioinformatic tests,
and transcriptome data, we examined binding motifs in the FSHβ promoter. The upstream
promotor sequence of 3500 bp (-3500 to 0) of Japanese quail FSHβ was obtained from
Ensemble (http://www.ensembl.org/index.html). This promotor sequence was analysed by
CiiDER transcription binding factor (TBFs) analysis tool (Gearing et al., 2019) against
JASPER core database. DNA motifs in the promoter were unique for 470 transcription
binding factors. The top 80 transcription binding factors were then subjected to PANTHER
gene ontology enrichment analyses (Mi et al., 2013). Several pathways were discovered
and included hormone responsive, epigenetic and responsive to nutrients (Extended Data
Table 9).
391
Immunocytochemistry and histological analyses
Snap frozen brains from experiment 1 were sectioned at 20µm using a cryostat (CM1850,
Leica). The tissues sections were collected in supercharged slides (631-0108, VWR) and
stored in -80c till processed for staining. We used mouse monoclonal antibody
(OAAEE00561, Aviva systems) raised against the Human vimentin gene (NCBI Reference
Sequence: NP_003371.2). The antibody has been shown to specifically bind with vimentin
expressed in avian cells (https://www.avivasysbio.com/vim- antibody-oaee00561.html).
Human and quail vimentin peptide show high similarity of 86.96% similarity. We used Goat
Anti-mouse Alexa Flour488 (A11001, Invitrogen) for the secondary antibody. As negative

control procedures, we performed omission of primary antibody and omission of secondary antibody.

Immunocytochemistry was performed using the standard immunofluorescence protocol (Majumdar et al., 2015) with minor modifications. Briefly, sections (brain) in slides were first enclosed in margin using ImmEdge pen (H-4000, Vector Labs). The sections were then first post fixed in 10% neutral buffered formalin (5735, ThermoScientific) for 4 hours. After fixing, the sections were washed (3 times; 5 min each) with TBS (phosphate buffer saline with 0.2% triton). Then they were blocked in 20% Bovine serum albumin (BSA) in TBS for 1 hour at room temperature (RT). Subsequently the blocking solution was removed by pipetting and the sections were incubated with primary antibody (1:300 dilution) for 2 hours at RT and finally overnight at 4°C. The next day, sections were first washed with TBS (3 times; 5 min each) and then incubated with secondary antibody (1:200 dilution) for 2 hours at RT. Finally, the sections were again washed with TBS (3 times; 5 min each) and mounted in Fluromount-G mounting media with DAPI (004959-52, Invitrogen). The dried slides were visualized using Leica DM4000B fluorescence microscope equipped with Leica DFC310 FX camera. Leica Application Suite (LAS) software was used for image acquisition. All the images were taken at constant exposure for the FITC channel at 10X and 20X magnification.

For analysis and quantification of % area, ImageJ version 1.53j was used. For this, 20X images were first converted into greyscale images (8 bit) and a threshold applied. The threshold was determined using the triangle method on multiple randomly selected images and applied for all the images. The scale of measurement in ImageJ was then set to CM in 300pixels/cm scale. A region of interest in the median eminence and dorsal 3rdV ependymal layer was specified as 300*500 pixels (1*1.67 cm scaled units) and area fraction was measured. At least 3 images from each animal were measured and averaged for each bird. A total of 27 quail were used and distributed evenly across photoperiod treatments (n=3).

Statistical analyses and figure presentation
GLM tests were conducted to test for statistical significance. One-way ANOVA with Bonferroni correction was applied to testes mass, body mass, fat score, vimentin immunoreactivity and qPCR analyses in Study 1. Two-way ANOVAs with photoperiod and zeitgebers main effects were conducted on qPCR analyses in Study 2. One-way ANOVAs were conducted for qPCR data in Study 3. qPCR data were log-transformed if violation of normality was detected. Significance was determined at $P<0.05$. Figures were generated using AdobeIllustrator and BioRender was used to create images in panels Fig1d,n; Fig3e and Extended Data Figure 6.

Acknowledgements: The authors thank Elisabetta Tolla, Christopher Elcombe, Ana Monteiro and David Hamilton for their assistance. The authors thanks Professor Gregory Ball and Neil Evans for comments on a previous version of the paper. Funding: the work was funded by a Leverhulme Trust Research Leader to TJS.

References

Bradshaw WE, Holzapfel CM (2007) **Evolution of animal photoperiodism.** *Annual Rev.*

Ecol. Evol. System. **38:**1-25.

Bustin SA, Benes V, Garson JA, Hellemans J, Huggett J, Kubista M, Mueller R, Nolan T,
Pfaffl MW, Shipley GL, Vandesompele J, Wittwer CT (2009) **The MIQE guidelines:**
minimum information for publication of quantitative real-time PCR experiments.
Clin. Chem. **55**:611-622.

Cornelissen G (2014) **Cosinor-based rhythmometry.** *Theoretical Biology Medical*
Modelling **11**:16.

Dennis G (2003) **DAVID: Database for Annotation, Visualization, and Integrated**
Discovery. *Genome Biol* **4**:R60.

Follett BK, Sharp PJ (1969) **Circadian rhythmicity in photoperiodically induced**
gonadotropin release and gonadal growth in the quail. *Nature* **223**:968-971.

Gearing LJ, Cumming HE, Chapman R, Finkel AM, Woodhouse IB, Luu K, Gould JA,
Forster SC, Hertzog PJ (2019) **CiiiDER: A tool for predicting and analysing**
transcription factor binding sites. *PLoS One* **14**:e0215495.

Gu Z, Eils R, Schlesner M (2016) **Complex heatmaps reveals patterns and correlations**
in multidimensional genomic data. *Bioinformatics* **32**:2847-2849.

Gwinner E (1986) ***Circannual Rhythms: endogenous annual clocks in the***
organization of seasonal processes. Springer-Verlag.

Gwinner E, Dittami J (1990) **Endogenous reproductive rhythms in a tropical bird.**
Science **249**:906-908.

Helm B, Stevenson TJ (2014) **Circannual rhythms: history, present challenges, future**
directions. In *Annual, lunar and tidal clocks*. pp. 203-225.

Keenan AB, Torre D, Lachmann A, Leong AK, Wojciechowicz ML, Utti V, Jagodnik KM,
Kropiwnicki E, Wang Z, Ma'ayan A (2019) ChEA3: **Transcription factor enrichment**
analysis by orthogonal omics integration. *Nucleic Acids Res* **47**:W212–W224.

King VM, Bentley GE, Follett BK (1997) **A direct comparison of photoperiodic time**
measurement and the circadian system in European starlings and Japanese
quail. *J. Biol. Rhythms* **12:**431-442.

Li H (2018) **Minimap2: pairwise alignment for nucleotide sequences.** *Bioinformatic*
34:3094-3100.

Lincoln GA, Clarke IJ, Hut RA, Hazlerigg DG (2006) **Characterizing a mammalian**
circannual pacemaker. *Science*, **314**:1941-1944.

Maechler M (2019) **Finding groups in data": Cluster analysis extended Rousseeuw et**
al. R package version, 2(0).

Majumdar G, Yadav G, Rani S, Kumar V (2015) **Bird eyes distinguish summer from**
winter: Retinal response to acute photoperiod change in the night-migratory
redheaded bunting. *J Chem Neuroanat.* **68**:55-60.

Mi H, Muruganujan A, Thomas PD (2013) **PANTHER in 2013: modeling the evolution of**
gene function, and other gene attributes, in the context of phylogenetic trees.
Nucleic Acids Res **41**:D377–D386.

Nakao N, Ono H, Yamamura T, Anraku T, Takagi T, Higashi K, Yasuo S, Katou Y,
Kageyama S, Uno Y, Kasukawa T, Iigo M, Sharp PJ, Iwasawa A, Suzuki Y, Sugano S,
Niimi T, Mizutani M, Namikawa T, Ebihara S, Ueda HR, Yoshimura T (2008)
Thyrotrophin in the pars tuberalis triggers photoperiodic response. *Nature*
452:317-322.

Patro R, Duggal G, Love MI, Irizarry R, Kingsford C (2017) **Salmon: fast and bias-aware**
quantification of transcript expression using dual-phase inference. *Nat. Methods.*
14:417-419.

Pengelley ET, Fisher KC (1957) **Onset and cessation of hibernation under constant**
temperature and light in the golden-mantled ground squirrel (Citellus lateralis).
Nature **180:**1371-1372.

Perez JH, Tolla E, Dunn IC, Meddle SL, Stevenson TJ (2019) **A comparative**
perspective on extra-retinal photoreceptor. *Trends Endocrinol. Metab.* **30**:39-52.

Robinson JE, Follett BK (1982) **Photoperiodism in Japanese quail: the termination of**
seasonal breeding by photorefractoriness. *Proc R Soc B.* **215**:95-116.

Singh D, Rani S, Kumar V (2013) **Daily expression of six clock genes in central and**
peripheral tissues of a night-migratory songbird: evidence for tissue-specific
circadian timing. *Chronobiol Int.* **30**:1208-17.

Stevenson TJ, Ball GF (2009) **Anatomical localization of the effects of reproductive**
state, castration, and social milieu on cells immunoreactive for gonadotropin-
releasing hormone 1 in male European starlings. *J Comp. Neurol.* **517**:146-155.

Stevenson TJ, Lynch KS, Lamba P, Ball GF, Bernard DJ (2009) **Cloning of**
gonadotropin-releasing hormone I complementary DNAs in songbirds facilitates
dissection of mechanisms mediating seasonal changes in reproduction.
Endocrinol. **150**:1826-1833.

Stevenson TJ, Hahn TP, Ball GF (2012a) **Variation in gonadotrophin-releasing**
hormone-1 gene expression in the preoptic area predicts transitions in seasonal
reproductive state. *J. Neuroendocrinol.* **24**:267-274.

Stevenson TJ, Replogle K, Drnevisch J, Clayton DF, Ball GF (2012b) **High throughput**
analysis reveals dissociable gene expression profiles in two independent
neuronal systems involved in the regulation of social behavior. *BMC Neurosci.*
13:126.

Visser ME, Gienapp P (2019) **Evolutionary and demographic consequences of**
phenological mismatches. *Nature Ecol Evol.* **3**:879-885.

Wingfield JC, Farner DS (1978) **The endocrinology of a naturally breeding population**
of the white-crowned sparrow (Zonotrichia leucophyrs gambelii). *Physiol. Zool.*
51:188-205.

Wingfield JC (2008) **Organization of vertebrate annual cycles: implications for control**

mechanisms. *Philos. Trans. R. Soc. B. Biol. Sci.* **363**:425–4.

Wood SH, Hindle MM, Mizoro Y, Cheng Y, Saer BRC, Miedzinska K, Christian HC, Begley

N, McNeilly J, McNeilly AS, Meddle SL, Burt DW, Loudon ASI (2020) **Circadian clock**

mechanism driving mammalian photoperiodism. *Nat Commun* **11**:4291.

Woodfill CJI, Wayne NL, Moenter SM, Karsch FJ (1994) **Photoperiodic synchronization**

of a circannual reproductive rhythm in sheep: identification of season-specific

time cues. *Biol. Reprod.* **50**:965-976.

Yamamura T, Hirunagi K, Ebihara S, Yoshimura T. (2004) **Seasonal morphological**

changes in the neuro-glial interaction between Gonadotropin-releasing hormone

nerve terminals and glial endfeet in Japanese quail. *Endocrinol*. **145**:4264-4267.

Yeo GH, Heisler LK (2012) **Unraveling the brain regulation of appetite: lessons from**

genetics. *Nat. Neurosci.* **15**:1343-1349.

Yoshimura T, Yasuo S, Watanabe M, Iigo M, Yamamura T, Hirunagi K, Ebihara S. (2003)

Light-induced hormone conversion of T4 to T3 regulates photoperiodic response

of gonads in birds. *Nature* **426**:178-181.

Zhao S, Fernald RD (2005) **Comprehensive algorithm for quantitative real-time**

polymerase chain reaction. *J. Comput. Biol.* **12**:1045-62.

Zielinski T, Moore AM, Troup E, Halliday KJ, Millar AJ (2014) **Strengths and Limitations**

of Period Estimation Methods for Circadian Data. *PLoS ONE* **9**:e96462.

Fig. 1 - Vernal increase in pituitary *FSHβ* precedes molecular switches in MBH and testes growth.

a, Schematic representation of the simulated annual rhythm in photoperiod. Quail were collected in 16hr light, 8hr dark photoperiod and then every 2 weeks the photoperiod was decreased by 2hrs to 14hr, 12hr, 10hr and then a 8L short photoperiod. Photoperiod was then increased to mimic the vernal transition and birds were collected at 10hr, 12hr, 14hr and 16hr light photoperiods. Testis volume confirmed critical day length (i.e. >12hr) induced growth. **b,** Body mass and **c,** abdomenal fat deposition increased until the autumnal equinox (12a), and then increased during the vernal photoperiod transitions. **d,** Diagram highlighting hypothalamic preoptic area (POA), mediobasal hypothalamus (MBH) and pituitary gland. Tanycytes in the MBH gate GnRH release into the pituitary. **e,** Heat-map of RNA-seq of MBH punches identifies distinct waves of transcripts as quail transition across photoperiodic conditions. **f-g,** qPCR assays for proopiomelanocortin (*POMC*) and deiodinase type-3 (*DIO3*) confirmed restricted activation during 10v-8L and 8L-10v phases, respectively. **h-i,** vimentin immunoreactivity in the median eminence (ME) shows tanycyte morphology growth is limited to 10a, 8L, and 10v photoperiods. **j,** Heat-map illustrating photoperiodic transitions in pituitary transcripts. **k-m,** confirmed that *FSHβ* is elevated under non-stimulatory photoperiods followed by increased prolactin (*PRL*) in 14v and thyrotropin stimulating hormone-β (*TSHβ*) in 16v. **n,** diagram summarizing that long photoperiods increase *GNRH* synthesis and release into the pituitary gland to stimulate *FSHβ* and induce testis growth. Transition to autumnal equinox phases results in reduced *FSHβ* expression and regressed testis. Prolonged exposure to short photoperiods inhibits *GNRH* expression, triggers tanycyte extension, maintains low *FSHβ* and regressed testis. Vernal transitions in photoperiod to the equinox results in resumption of *GNRH* and elevated *FSHβ* expression without testis growth. Data are mean +/- SEM, and residual dot plot. **a-c, f,g,k,m** One-way ANOVA with Bonferonni corrected Tukey's test for multiple comparisons. **h, l** One-way ANOVA with Tukey tests for significant pairwise comparison. Letters denote significant difference between photoperiod phase. Raw data available in Table S1.

Figure 2 - *FSHβ* is constitutively expressed during the vernal equinox.

a, schematic representation of four photoperiod treatment groups with arrows to indicate the daily sampling time. **b,** testes volume remained in a regressed non-functional state in autumnal equinox (12a), short photoperiod (8L) and vernal equinox (12v). Photoperiods that exceeded the critical day length (i.e.,>12hr) induced testes growth. **c,** Pituitary circadian clock gene *ARNTL1* maintained daily rhythmic expression waveforms across all photoperiods (P<0.001) , no significant difference between photoperiod treatment (*P*=0.42) **d,** *PER3* displayed a daily waveform across 12a, 8L, 12v and 16L groups (*P*<0.001) and was anti-phase compared to *ARNTL1*. No significance difference between photoperiod treatments (*P*=0.31) **e,** *FSHβ* expression was significantly higher in 12v compared to long photoperiods (16L; *P*<0.001), autumnal equinox (12a; *P*<0.001) and short photoperiod (8L; *P*<0.001), but was not rhythmic (*P*=0.66). **f,** Similarly, *PRL* was high in 16L compared to 12a (*P*<0.001), and 8L (*P*<0.001), there was no significant daily rhythm (*P*=0.52). **g,** *FSHβ* promoter was devoid of circadian gene binding D-box and E-box motifs but contains a series of hormone and nutrient responsive motifs. **b-f** Two-way ANOVA followed by Tukey's pairwise tests, rhythmic analyses were conducted using GraphPad Prism. ǂ indicates significant photoperiod treatment effect; # denotes significant time of day effect. Data are mean +/- SEM, and residual dot plot (a-d). Rhythmic analyses are presented in Table S7 and raw data are available in Table S1.

Figure 3 - Endogenous and light-induced *FSHβ* expression.

a, *FSHβ* expression increased during the photoinduced transition from 8L to 10v, and 12v. *FSHβ* also showned a smaller, yet significant increase in expression after prolonged exposure to 8L. Y-axis is presented in log-scale due to the significant increase in *FSHβ* expression in 10v and 12v. **b,** *OPN5* was detected in the pituitary gland and showed a significant increase in expression in the transition to 10v and 12v similar to *FSHβ* expression. **c,** *DIO3* was significantly reduced in 12v quail compared to all other treatment groups. **d,** *GNRH* expression remained constant during the transition from 8L to 12v. However, continued exposure to 8L was observed to increase *GNRH* expression. Data are mean +/- SEM, and residual dot plot (a-d). One-way ANOVA with Tukey's test for multiple comparisons. Letters denote significant difference between photoperiod phase. Raw data available in Table S1. **e,** Schematic representation of the endogenous and light-dependent increase in pituitary cell types during the transition from 8L to stimulatory 16v light treatments. Increased color indicates increased transcript expression.