Title: Sex-specific effect of antenatal Zika virus infection on murine fetal growth, placental

nutrient transporters, and nutrient sensor signaling pathways

Authors:

Daniela Pereira-Carvalho[1], Alessandra Cristina Chagas Valim[1], Cherley Borba Vieira

Andrade[1], Enrrico Bloise[2], Ariane Fontes Dias[1], Veronica Muller Oliveira Nascimento[1], Felipe

Lopes Brum da Silveira[3], Inácio Medeiros[3], Sharton Vinicius Antunes Coelho[4], Luciana Barros

Arruda[4], Adriane Regina Todeschini[1], Wagner Barbosa Dias[1], Tania Maria Ortiga-Carvalho[1]

[1] Instituto de Biofísica Carlos Chagas Filho, Universidade Federal do Rio de Janeiro, Rio de

Janeiro, Brasil.

[2] Departamento de Morfologia, Universidade Federal de Minas Gerais, Belo Horizonte, Brasil.

[3] Instituto do cérebro. Universidade Federal do Rio Grande do Norte, Brasil.

[4] Instituto de Microbiologia Paulo de Góes, Universidade Federal do Rio de Janeiro, Rio de

Janeiro, Brasil.

KEYWORDS: Fetal growth restriction, glucose transport, placenta, ZIKV, OGT, gestational

infection

Abstract:

Maternal Zika virus (ZIKV) infection during pregnancy can associate with severe intrauterine

growth restriction (IUGR), placental damage, and metabolism disturbance, as well as newborn

neurological abnormalities. Here, we investigated whether maternal ZIKV infection affects

placental nutrient transporters and nutrient-sensitive pathways. Immunocompetent (C57BL/6)

mice were injected with Low (10^3 PFU-ZIKV$_{PE243}$) and High (5×10^7 PFU-ZIKV$_{PE243}$) ZIKV

titers at gestational day (GD) 12.5, for tissue collection at GD18.5 (term). Feto-placental

growth of male fetuses was dramatically affected by ZIKV, whereas no differences were

observed in female fetuses. ZIKV promoted increased expression of glucose transporter type 1

(*Slc2a1*/Glut1) and decreased levels of glucose-6-phosphate in female placentas, with no
differences in amino-acid transport potential. In contrast, glucose transport in male placentas
was not affected by ZIKV, whilst a decreased placental protein expression of sodium-coupled
neutral amino acid 2 (Snat2) was detected in the male low-dose ZIKV-infected group. There
were also sex-dependent differences in the hexosamine biosynthesis pathway (HBP) and O-
GlcNAcylation in ZIKV infected pregnancies, showing that ZIKV can cause disturbance in the
nutrient handling in the placental tissue. Our findings thus identify relevant molecular
alterations in the placenta caused by maternal ZIKV infection related to nutrient transport and
availability. Notably, our results suggest that female and male placentas adopt different
strategies to cope with the altered metabolic state caused by ZIKV. This may have relevance
for understanding the effects of congenital Zika syndrome and could potentially assist future
therapeutic strategies.

Introduction:

Zika virus (ZIKV) is an Aedes-borne arbovirus belonging to the *Flaviviridae* family of
positive, single-stranded, enveloped RNA viruses that has emerged as a major global health
concern in the past decade. ZIKV infection in pregnant women has been linked to mild to
severe neonatal abnormalities, such as microcephaly and intracranial calcifications, as well as
pregnancy loss, intrauterine growth restriction (IUGR), and preterm labor, among other
complications (Oliveira Melo *et al.*, 2016). The ZIKV pandemic is currently thought to be
"controlled"; however, more than 60% of the world's population live in regions with suitable
environmental conditions for ZIKV spread, with neither a licensed vaccine nor an antiviral
drug (Messina *et al.*, 2016) available for prevention and treatment of the disease.

The mechanisms underlying antenatal ZIKV commitment of fetal growth and
developmental trajectories have only been partially elucidated. Similar to other flaviviruses,
ZIKV requires a variety of intricate interactions with host factors for successful infection, such
as reprogramming of cellular metabolism underpinning ZIKV pathogenesis (Thaker *et al.*,
2019; Chen *et al.*, 2020; Yau *et al.*, 2021). In the placenta, ZIKV is able to cause lipid
reprogramming, mitochondrial dysfunction, and inflammatory immune imbalance,
contributing to placental damage (Chen *et al.*, 2020; Andrade *et al.*, 2021; Creisher *et al.*, 2022).
At the functional level, placental damage can impair essential placental barrier mechanisms
that support fetal growth and development, such as unidirectional glucose and amino acid
transfer from mother to conceptus (Winterhager e Gellhaus, 2017).

Glucose (GLUT) and sodium-coupled neutral amino acid (SNAT) transporters are key placental nutrient transporters - highly regulated by environmental factors such as oxygen, nutrients and hormones (Coan *et al.*, 2010; Higgins *et al.*, 2016). Alterations in extracellular glucose and insulin-like growth factor I (IGF-I) levels can cause changes in the activity and expression of glucose and amino acid transporters (Lager e Powell, 2012). When deregulated, GLUT and SNAT transporters have been linked to pregnancy complications, such as IUGR, and the development of neurological and cardiometabolic diseases later in life (Gaccioli e Lager, 2016; Vaughan *et al.*, 2021; Li *et al.*, 2022). In the context of infection, *plasmodium falciparum* (the etiological agent of placental malaria) infestation has been shown to decrease GLUT1 in the term human placenta (Chandrasiri *et al.*, 2014). Additionally, ZIKV infection was linked with an increase in glucose uptake and GLUT3 expression human in first-trimester cytotrophoblast cells (Vota *et al.*, 2021). Demonstrating that placental nutrient uptake and metabolism is impacted by common infective agents.

In this connection, the hexosamine biosynthesis pathway (HBP) is a neglected anabolic branch of glucose metabolism that uses glucose and glutamine to generate UDP-GlcNAc as the final product. Glutamine-fructose amino transferase (GFAT) is HBP's rate-limiting enzyme, whereas UDP-GlcNAc is considered a central metabolic sensor since it requires inputs from carbohydrate, amino acid, nucleotide and fatty acid metabolism. UDP-GlcNAc is the donor substrate of O-GlcNAc transferase (OGT), the enzyme responsible for the addition of O-GlcNAc to serine/threonine residues of target intracellular proteins, while O-GlcNAcase (OGA) is the enzyme responsible for its removal (Hart *et al.*, 2011). O-GlcNAcylation is a highly dynamic and ubiquitous posttranslational modification that modulates protein function in a nutrient-sensitive way, linking cell signaling to metabolism (Hart *et al.*, 2011). Cellular UDP-GlcNAc levels and global O-GlcNAcylation are coordinated and are highly responsive to glucose availability, making O-GlcNAcylation well suited nutrient-sensor (Hart *et al.*, 2019). HBP and O-GlcNAcylation are central in controlling immune cell function, consequently having key regulatory effects on immune responses against infections (Quik *et al.*, 2020). The molecular mechanisms involving O-GlcNAcylation were investigated using different virus infection models, including HPV (Zeng *et al.*, 2016; Shi *et al.*, 2022), HBV (Wang, X. *et al.*, 2020), influenza A virus (Wang, Q. *et al.*, 2020), and HSV (Angelova *et al.*, 2015). Although these studies demonstrated that viruses are able to modulate O-GlcNAcylation, impacting on autophagy mechanisms and viral replication, the effects of the Zika virus remain unexplored.

In addition, OGT and OGA are highly expressed in the placenta, and key placental proteins are OGT targets. Placental OGT expression has been linked to long-term metabolic

and neurodevelopmental programming (Howerton *et al.*, 2013; Howerton e Bale, 2014; Nugent
et al., 2018; Moore *et al.*, 2021). For instance, studies in mice showed that aberrant O-
GlcNAcylation causes reduced placental vasculature development, insulin tolerance, and
changes in body weight (Yang *et al.*, 2015; Nugent *et al.*, 2018; Moore *et al.*, 2021). An OGT
knockout mouse model showed important changes in the expression patterns of hypothalamic
genes during intrauterine development (Howerton *et al.*, 2013; Howerton e Bale, 2014).

A previous study by our group showed that maternal exposure to ZIKV affects placental
function, including placental ultrastructure and ABC transporter protein expression, even in the
absence of vertical transmission and that these effects are dependent on viral infective titers
and maternal immune status (Andrade *et al.*, 2021). Despite their importance to fetal growth
and development and their role in response to infections, there is limited information about
GLUT and SNAT transporter expression profiles and OGT nutrient-sensing pathway activity
in the placenta of ZIKV-infected dams. Therefore, in the present study, we hypothesize that
maternal exposure to ZIKV affects fetal-placental growth, the expression of placental nutrient
transporters and nutrient distribution, as well as placental O-GlcNAcylation, and specific
components of the HBP, in a sex-specific manner.

Materials and methods:

Virus

The ZIKV strain PE243 (ZIKVPE243; GenBank ref. number KX197192) was kindly
provided by Dr Ernesto T. Marques (University of Pittsburgh, PA). Viruses were propagated
in C6/36 cells and titrated by plaque assays in Vero cells, as previously described (Coelho, et
al., 2017). Conditioned medium from noninfected C6/36 cells cultured under the same
conditions was used as a control.

Experimental design

Female C57BL/6 mice aged between 8 and 10 weeks were mated with male mice , and
on the day after mating, copulation was confirmed by visualization of the vaginal plug and
considered gestational day 0.5 (GD0.5). Dam's weight gain was monitored for confirmation of
pregnancy. At GD 12.5, dams with a weight gain greater than 3 g were considered pregnant
and randomly distributed into the following experimental groups: control (received an

intravenous injection (i.v.) of supernatant from noninfected C6/36 cells, n=15), Low Dose
(received an i.v. with 10^3 plaque-forming units (PFU) of ZIKVPE243, n=15), and High Dose
(received an i.v. with $5x10^7$ PFU of ZIKVPE243, n=15) ZIKV. The i.v. injections were
performed at GD12.5, and virus preparation was undertaken as previously described (Coelho
et al., 2017; Andrade *et al.*, 2021). At 18.5, all animals were weighed and euthanized with 5%
isoflurane by inhalation. All placental/fetal units were dissected, collected and weighed. Fetal
tails were kept for sex determination by detection of the *Sx* gene using specific primers
(Forward: 5'-GATGATTTGAGTGGAAATGTGAGGTA-3'; Reverse: 3'-
GAATACAAATATCCGTACGTGGTACAT-5') (Davalieva *et al.*, 2006).

Placental specimens were stored in liquid nitrogen for further qPCR and western
blotting analyses. For the MALDI-IMS analysis, samples were frozen in acetone cooled with
dry ice and subsequently kept in a -80 freezer. For morphology and protein
expression/localization analysis, the tissues were fixed overnight in buffered paraformaldehyde
(4%, Sigma–Aldrich, Brazil). The experimental design is illustrated in Fig 1A. Animals were
kept in a controlled temperature room (23°C) with a light/dark cycle of 12 hours and had access
to water and food *ad libitum*. All procedures were approved by the Animal Care Committee
(CEUA-036/16, 128/22, A7/20-036-16) of the Federal University of Rio de Janeiro.

Histological and immunohistochemistry analysis

Previously fixed placental fragments were subjected to dehydration (increasing ethanol
concentrations; ISOFAR, Brazil) and diaphanization with xylol (ISOFAR, Brazil). The
fragments were embedded in paraffin (Histopar, Easypath, Brazil) and sectioned (5 µm
thickness) using a Rotatory Microtome CUT 5062 (Slee Medical GmbH, Germany). Sections
were submitted to the following techniques and performed as previously described (Fontes *et*
al., 2019):

<u>Periodic acid-Schiff (PAS) staining</u>

Histological sections were submitted to diaphanization with three xylol baths and
hydrated with three decreasing concentrations of ethanol. The sections were oxidized for 15
min. with 0.5% periodic acid (Sigma–Aldrich, USA), washed in flowing water and incubated
for 10 min. with Schiff's reagent (Merck Millipore, Germany) at room temperature.

<u>Immunochemistry of nutrient transporters in the placenta</u>

The sections were submitted to diaphanization with three xylol baths and hydrated with three decreasing concentrations of ethanol. Subsequently, sections were exposed for 30 min to 3% hydrogen peroxide, followed by microwave antigenic recovery in Tris-EDTA (pH=9) and sodium citrate (pH=6) buffers (15 min. for Tris-EDTA buffer and 8 min. for citrate buffer). Sections were washed in PBS+ 0.2% Tween and incubated in 3% BSA in PBS for 1 hour to block nonspecific binding sites. Sections were then incubated with the primary antibody (Supplementary Table 1) overnight at 4°C. On the day after, sections were incubated with the biotin-conjugated secondary antibody (SPD-060, Spring Bioscience, USA) for 1 h at room temperature. Slides were incubated with streptavidin (SPD-060 – Spring Bioscience, USA) for 1 h and with 3,3-diaminobenzidine (DAB) (SPD-060 – Spring Bioscience, USA).

After PAS and immunochemistry procedures, all sections were counterstained with hematoxylin, followed by dehydration in alcohol and xylol. The sections were then mounted with coverslips using Entellan (Merck, Germany). Digital images were obtained using a high-resolution Olympus DP72 (Olympus Corporation, Japan) camera coupled to an Olympus BX53 light microscope (Olympus Corporation, Japan). The PAS-stained sections were used to analyze the area of each placenta of the rodent placental zones, labyrinth (Lz) and junctional (Jz) zones by free-drawing using ImageJ software (National Institutes of Health, USA). For immunochemistry quantification of Glut1 and Snat2, fifteen digital images (40X) from different fields were captured in each placenta zone and analyzed with Image-Pro Plus, version 5.0 software (Media Cybernetics, USA) mask tool. At each staining, a total of 150 digital images were analyzed per group (75 Lz images and 75 Jz images). Hence, 450 images were analyzed by target protein including all experimental groups. The percentage of viable tissue area was considered upon the exclusion of negative spaces. All negative controls were performed with the omission of the primary antibody. Analysis were performed as previously described (Andrade *et al.*, 2021). All analyses were conducted blind to the group.

RT–qPCR

Placental RNA was extracted using TRIzol following the manufacturer's protocol (TRIzol Reagent; Life Technologies, USA). The concentration of RNA was assessed using a NanoPhotometer (Implen, Munchen, Germany). The quantification and RNA purity were evaluated by reading in a nanophotometer, considering the relations A260/280 ~2.0 and

A260/230 ~2.0-2.2. (Montreal Biotech Inc., Canada). A total of 1 μg per sample was reverse

transcribed using the High-Capacity cDNA Reverse Transcription Kit (Applied Biosystems,

USA) according to the manufacturer's instructions. qPCR was performed to evaluate mRNA

expression quantification using gene-specific primer pairs (Supplementary Table 2) and

EVAGREEN (Solis Byodine, USA). The standard thermal cycling protocol was conducted as

follows: combined initial denaturation at 50 °C (2 min) and 95° C (10 min), followed by 40

cycles of denaturation at 95 °C (15 s), annealing at 60 °C (30 s) and extension at 72 °C (45 s),

using QIAquant 96 2plex (QIAGEN, Germany). Relative expression was calculated using the

2-ΔΔCt method (Livak e Schmittgen, 2001), and genes of interest were normalized to the

geometric mean expression of 3 reference genes (*Ywhaz, PPib, ß-actin*).

Western blotting

Placental protein extraction was performed on frozen tissue homogenized using

commercial RIPA lysis buffer (Thermo Scientific, US) supplemented with phosphatase

inhibitors (5 mM Na4P2O7, 50 mM NaF, 5 mM Na3VO4), protease inhibitor mixture (Sigma–

Aldrich), and OGA inhibitor (PugNAc, 1 M). The protein concentration was determined using

the bicinchoninic acid protein assay (Thermo Fisher Scientific, EUA). Lysates were separated

by sodium dodecyl sulfate–polyacrylamide gel electrophoresis (SDS–PAGE) and transferred

onto nitrocellulose membranes (Bio-Rad Laboratories, US) using a semidry technique (semidry

blotter, Invitrogen). The membrane washing steps were performed with Tris-buffered saline

tween (TBST) after the following blocking and incubation steps. Blocking was performed with

3% BSA in phosphate-buffered saline (1x) with 0.1% Tween (1X Tris-buffered saline, 0.1%

Tween - TBS-T) for 1 h at room temperature. The membrane was incubated with primary

antibody (Supplementary Table 1) in 3% BSA + TBS-T overnight. The next day, the

membranes were washed and incubated with rabbit or mouse secondary antibodies diluted in

TBST for 60 minutes. ß-actin was used as a loading control. Protein bands were visualized

using Scientific SuperSignal West Femto enhanced chemiluminescence (ECL) substrate

(Thermo Scientific, US) and captured with ImageQuant LAS 4000.

MALDI-IMS

The analysis of specific biomolecules enriched in murine placentas exposed to ZIKV

was performed using matrix-assisted laser desorption/ionization imaging mass spectrometry

(MALDI-IMS). We used the mass/charge ratio (m/z) to analyze our biomolecules of interest: glucose-6-phosphate (m/z= 259.0232), glutamine (m/z= 145.0618) and UDP-GlCNAc (m/z= 606.3824).

The presence of the biomolecules was analyzed in 5 placentas (randomly picked) per experimental group. Cryosectioning of previously frozen placentas was performed with 10-mm-thick sections using a cryostat (Leica CM1860-UV, Leica Biosystems, Nussloch, Germany). Then, the slices were added to glass slides coated with ITO (indium tin oxide) and conditioned at -80°C until the time of analysis. The slides were removed from -80°C and allowed to reach room temperature in the desiccator until the time of deposition of the matrix. Subsequently, deposition of the matrix was carried out. The matrix solution used was 10 mg/mL 9-aminoacridine (Sigma) in 80% methanol and 20% water. The application was performed with spray with an application flow of 800 µL/h and N2 pressure of 12.5 psi for 20 minutes. The analysis was performed with a Solarix XR mass spectrometer (FT-ICR, Bruker) using negative polarity, mass range from 75 to 1200 m/z, laser focus medium mode and power make laser at 35%. Two hundred shots were fired per field. The data obtained in the spectrometry were normalized into TIC (total ion count).

Statistics

Statistical analyses were performed using GraphPad Prism version 8 (GraphPad, La Jolla, CA, USA). A D'Agostino & Pearson normality test was used to evaluate normal distribution. Data for each sex were analyzed using one-way ANOVA followed by Tukey's test. Outliers were identified using a Grubbs test. Differences were considered significant when $p<0.05$. Litter parameters were evaluated using the mean value of all fetuses and/or placentae in the litter per dam and not the individual conceptus (Bloise *et al.*, 2012; Connor *et al.*, 2020). The number of samples per group for each analysis is shown in each figure and described in the legends of figures and footnotes of tables.

<u>Results:</u>

Antenatal ZIKV infection alters fetal-placental growth in a sex-specific manner

Pregnant mice were inoculated at GD12.5 with a Low Dose (LD, 10^3 PFU) or High Dose (HD, $5x10^7$ PFU) of ZIKVPE243 or with conditioned medium from noninfected C6/36

cells as controls (Fig 1A). To investigate whether maternal ZIKV infection affects placental
efficiency and fetal biometry at term, fetuses and placentas were weighed at GD 18.5
(considered term in C57BL/6). No significant difference was observed in the weight of female
fetuses and their corresponding placentas, as well as in their placental efficiency (Fig 1B). The
fetal head weight of females in the LD group was significantly reduced (-16.02%, p=0.041)
compared to corresponding control group (Fig 1B). In contrast, fetal-placental growth was
dramatically reduced in male fetuses. The fetal weight was significantly reduced in both
infected groups (LD: -13.8%, p=0.002; HD: -20.5%, p=0.032) compared to control male
fetuses (Fig 1D). Additionally, placentas from the infected groups exhibited reduced weight
(LD: -20,92% p=0.033, HD: -15,21%, p=0.001) compared to the correspondent control group,
with no significant difference in placental efficiency (Fig 1D). We therefore investigated
whether maternal ZIKV infection would cause placental remodeling, affecting the two main
functionally distinguishable regions of the rodent placenta. There were no differences in the
areas of the Lz and Jz or in the total placental area when the ZIKV-infected groups were
compared to the control group in either female (Fig 1C) or male (Fig 1E) placentas.

283
284 **Maternal ZIKV infection induces changes in placental amino acid and glucose transport**
285

To investigate whether changes in fetal and placental growth may be related to
alterations in placental transport of amino acids and glucose, we evaluated the expression of
key glucose and amino acid transporters and the levels of glucose-6-phosphate and glutamine
in placental tissue. In particular, the mRNA expression of key transporters for glucose (*Slc2a1*
and *Slc2a3*) and amino acids (*Slc38a1* and *Slc38a2*) was quantified in the placenta by RT–
qPCR. Additionally, the protein expression of the transporters Glut1 and Snat2 was quantified
by immunohistochemistry in both the placenta Lz and Jz areas. Levels of glucose-6-phosphate
and glutamine was analyzed by MALDI-MSI.
In placentas from female fetuses, these analyses showed that the expression of *Slc2a1*
mRNA in the HD group was 62% greater (p=0.047) than in the control group, while this
difference was not observed for the LD group (Fig 2A). No differences were observed in *Slc2a3*
mRNA expression between the experimental groups (Fig 2A). In addition, no differences were
found in *Slc38a1* and *Slc38a2* mRNA expression in female placentas (Fig 2B). At the protein
expression level, the expression of Glut1 in the Lz of female placentas was greater in the ZIKV-
infected groups (LD: +73.40%, p=0.007; HD: +73.38%, p=0.007) than in the control group
(Fig 2C). However, Glut1 expression did not vary among groups in the Jz region of female

placentas (Supplementary Fig 1A). We evaluated the protein expression of Snat2 in female
placental tissue and found no differences between the Zika-infected and control groups in either
the Lz (Fig 2D) or Jz (Supplementary Fig 1B) regions. Interestingly, levels of glucose-6-
phosphate was reduced in the ZIKV-infected groups compared to the control placentas (Fig
2E), whereas the levels of glutamine in the female placenta was similar between groups (Fig
2E).

The same parameters were evaluated in placentas from males, but the results showed
different responses. The mRNA expression of key glucose transporters, *Slc2a1* and *Slc2a3*,
was not significantly different between the experimental groups (Fig 3A). Additionally, the
mRNA expression of the amino acid transporters *Slc38a1* and *Slc38a2* was not significantly
different between the ZIKV-infected and control groups (Fig 3B). In addition, the protein
expression of Glut1 in male placentas was not different between the experimental groups in
either Lz (Fig 3C) or Jz (Supplementary Fig 1C). While Snat2 protein expression in the Lz of
male placentas was lower in the LD group than in the control group (-63.66%, p=0.025), no
differences were observed in the comparison with the HD group (Fig 3D). We found no
differences between the experimental groups when analyzing Snat2 protein expression in the
Jz of male placentas (Supplementary Fig 1D). Meanwhile, the presence of essential molecules
in male placentas, glucose-6-phosphate and glutamine placenta were similar between the
ZIKV-infected and control groups (Fig 3E).

Maternal ZIKV infection alters OGT expression and O-GlcNAcylation in female placentas

To gain further information about the changes in fetal growth and nutrient transport and
distribution, we evaluated pathways sensitive to nutrient availability. For this, we analyzed the
mRNA and protein expression of components of the HBP and O-GlcNAcylation. We also
analyzed the presence of UDP-GlcNAc, the final product of the HBP.

The analysis of *Gfpt1* mRNA expression showed no significant difference in female
placentas (Fig 4A). However, we found that the protein expression of Gfat1 was lower in the
HD (-33.73%, p=0.038) group than in the control group, with no significant changes in the LD
group (Fig 4B). Interestingly, analysis of the presence of UDP-GlcNAc showed that there was
a lower presence of this product in both ZIKV-infected groups in female placentas (Fig 4D).
In addition, *Ogt* mRNA expression was lower in the HD group than in the control group (-
42.17%, p=0.038), but the comparison with the LD group did not present significant changes

(Fig 4D). Additionally, no differences were observed in *Oga* mRNA expression between the experimental groups (Fig 4D). While the protein expression of OGT and OGA was not different between the groups, the O-GlcNAcylation levels were significantly lower in the LD (-35.12%, p=0.045) and HD (-38.09%, p=0.029) ZIKV-infected groups than in the control group in female placentas (Fig 4E).

In male placentas, the relative *Gfpt1* mRNA expression showed no significant difference between the ZIKV-infected and control groups (Fig 5A). However, Gfat1 protein expression in the HD group was greater than that in the control group (+32.54%, p=0.039), with no significant changes in the LD group (Fig 5B). In addition, the analysis of UDP-GlcNAc in the placenta of males was similar between groups (Fig 5C). Then, we analyzed the mRNA expression of the enzymes related to O-GlcNAcylation. The *Ogt* mRNA HD group was greater than the LD (+49.41%, p=0.038) and control (+48.03%, p=0.031) groups. The *Oga* mRNA expression was also greater in the HD group than the LD (+52.67%, p= 0.030) and control (+16.52%, p=0.048) groups (Fig 5D). The protein expression of OGT and OGA as well as O-GlcNAcylation levels were not different between the experimental groups, showing that O-GlcNAcylation was not affected by ZIKV in males (Fig 5E).

Discussion:

In the present study, we investigated the effect of maternal ZIKV infection on placental makers related to nutrient transport and availability using a mouse model of impaired fetal growth and survival with no vertical virus transmission (Andrade *et al.*, 2021). Herein, we showed that maternal ZIKV infection during pregnancy impacts the litter and their placentas in a sex-specific manner, with variations in nutrient transport and O-GlcNAcylation mechanisms, suggesting that females and males adopt different strategies to cope with the altered placental metabolic state caused by ZIKV infection.

The fetal-placental growth in the female litter was slightly affected by maternal infection. However, fetal-placental growth in males was dramatically affected by ZIKV, independently of viral titers. The reduction in the weight of fetuses and placentas is a known impact of ZIKV infection (Nem De Oliveira Souza *et al.*, 2018; Paul *et al.*, 2018; Szaba *et al.*, 2018; Suzukawa *et al.*, 2020). In fact, studies have demonstrated the potential correlation between fetal and placental weight in mice, as fetal growth relies on a variety of placental functions, such as nutrient transport and hormonal production, that can be affected by the reduction in this organ's size, impacting fetal growth (Nardozza *et al.*, 2017). It is important to

mention that fetal growth restriction during pregnancy enhances the risk for preterm birth, fetal

mortality, and the development of chronic diseases later in life through developmental

programming (Sferruzzi-Perri e Camm, 2016; Nardozza *et al.*, 2017; Burton e Jauniaux, 2018).

Moreover, the capability of ZIKV to induce sex-specific responses has been shown in

different models in the literature (Sherer *et al.*, 2021; Andrade *et al.*, 2022). Interestingly, our

study shows that these differences start to be modulated during intrauterine life. The

mechanisms by which the placenta supports fetal growth are likely differ between female and

male fetuses within the litter, even in normal physiological mouse pregnancies (Salazar-Petres

et al., 2022). However, the mechanisms by which these processes are affected by maternal

infection have yet to be determined.

Using the present experimental model, our group had previously shown that ZIKV

infection impacted proliferation and apoptosis processes in the placental tissue (Andrade *et al.*,

2021). In addition, in a mouse model of infection with poly (I:C), a synthetic double-stranded

RNA viral mimic, it was shown that the infection decreased proliferation in placental Lz,

whereas apoptosis in Jz was increased (Monteiro *et al.*, 2022). While these studies show that

viruses are able to impair placental remodeling, our results did not show differences in placental

Lz and Jz areas in either sex.

In our study, the expression of the glucose transporter Glut1 was greater in the exchange

(Lz) zone of the infected groups only in female placentas. Glut1 (*Slc2a1*) is the primary

placental glucose transporter, as it is the predominant isoform abundantly expressed during

early pregnancy and at term in humans and rodents (Sakata *et al.*, 1995; Das *et al.*, 1998).

Alterations in the expression of Glut1 (*Slc2a1*) were observed in models with reduced

availability of oxygen and nutrients (Coan *et al.*, 2010; Higgins *et al.*, 2016), as well as in

gestational disorders (Luscher *et al.*, 2017; Yao *et al.*, 2017; Borges *et al.*, 2019; Yang *et al.*,

2022), and in response to pharmacological treatment (Langdown e Sugden, 2001). Therefore,

placental Glut1 (*Slc2a1*) seems to have great plasticity and plays a role in responses to

alterations in the maternal milieu. Interestingly, the presence of glucose-6-phosphate was

reduced in the infected groups. There are several positive regulators of Glut expression in the

placenta, and low glucose concentration is one of them (Sibiak *et al.*, 2022). In addition, ZIKV

infection was linked with an increase in glucose uptake and GLUT3 expression in first-

trimester cytotrophoblast cells (Vota *et al.*, 2021). These results together may indicate that

more glucose is being transported to the female fetuses and are consistent with the maintenance

of fetal growth observed in females. Overall, our study shows that the placental increase in

Glut1 expression is an adaptation of the female placenta in response to maternal infection in
this infection model.

In males, no changes related to glucose transport and glucose-6-phosphate distribution
were observed. Nevertheless, the protein expression of Snat2 was lower in the transport zone
of the LD group. Low placental Snat2 expression causes fetal growth restriction (Vaughan *et*
al., 2021). Therefore, there is a possible relationship between reduced Snat2 expression and the
reduced fetal growth observed in the fetuses of this group. However, further studies must be
performed to confirm this hypothesis.

The HBP is a metabolic pathway that generates the "sensing molecule" UDP-GlcNAc.
An essential role in the sensing mechanism of the HBP, especially in the context of glucose
availability, is assigned to the enzyme catalyzing the first and rate-limiting step of this pathway,
fructose-6-phosphate transaminase (GFAT) (Chiaradonna *et al.*, 2018). In female placentas,
the protein expression of GFAT1 was reduced in the HD group. In addition, the presence of
UDP-GlcNAc was reduced in both ZIKV-infected groups. Consistent with these findings, the
O-GlcNAcylation levels were reduced in the female placentas from the ZIKV-infected groups.
Together, these results indicate reduced flow of nutrients into the HBP in female placentas.
Additionally, considering the reduced glucose-6-phosphate and UDP-GlcNAc as well as O-
GlcNAcylation, there is an indication of nutrient deprivation in the placental tissue. ZIKV
infection triggers several adaptations to cellular metabolism (Thaker *et al.*, 2019; Chen *et al.*,
2020; Yau *et al.*, 2021). Specifically, regarding glucose metabolism, Thaker et al showed that
ZIKV infection reprograms glucose metabolism, changing glucose utilization, which is linked
to whether cell death occurs (Thaker *et al.*, 2019). Specifically, in placental tissue, these
changes can affect diverse placental functions related to fetal growth and development and
interfere with placental intrinsic defense mechanisms (Zhang *et al.*, 2015).

The reduction in placental O-GlcNAcylation extends beyond nutrient availability.
Diverse studies have shown that protein O-GlcNAcylation regulates innate immune cell
function, such as the NF-κB pathway (Dela Justina *et al.*, 2017; Quik *et al.*, 2020; Dong *et al.*,
2022). Moreover, altered placental OGT expression can impact long-term metabolic and
neurodevelopmental programming. The reduced OGT expression in a placental-specific OGT
knockout mouse model altered the expression patterns of important hypothalamic genes
involved in endocrine and anti-inflammatory signaling during development (Howerton e Bale,
2014). These alterations may mediate changes in placental endocrine function and differences
in the transmission of important signals to fetal development. These are important mechanisms
to be highlighted and understood, as children exposed *in utero* to ZIKV have greater risk of

neurodevelopmental abnormalities in the first 18 months of life (Mulkey *et al.*, 2020). These
findings reinforce the importance of monitoring the long-term neurological development of all
newborns with exposure to ZIKV. While the placental HBP and O-GlcNAcylation are clearly
altered in females, there were slight changes observed in males, again reinforcing the sex-
specific characteristic of the response to ZIKV in this tissue. The specific sex difference
observed in our work should be investigated since O-GlcNAcylation is identified as a potential
mechanism related to sex-dependent neurodevelopment changes (Howerton *et al.*, 2013), and,
as already mentioned, the relationship between infection by ZIKV and possible damage to
several processes related to the nervous system is recurrent (Trus *et al.*, 2019; Andrade *et al.*,
2022), but little explored in a sex-specific way.

In conclusion, our study highlights the importance of the placenta as a key organ in
mediating gestational infection and points to relevant molecular alterations caused by maternal
ZIKV infection. Collectively, our results suggest that female and male placentas adopt different
strategies to cope with the altered metabolic state caused by ZIKV infection and that the
alteration of placental nutrient transport and O-GlcNAcylation may be linked to the known
effects of ZIKV infection. Our findings contribute to the understanding of alterations that could
assist future preventive and therapeutic strategies and sex-specific approaches of individuals
who had their mothers infected.

Figure legends:

Figure 1. Impact of ZIKV infection on fetal-placental growth. (A) Experimental design.
(B) Fetal weight, fetal head weight, placental weight, and placental efficiency in female fetuses.
(C) Quantification of the areas of the Lz, Jz, and total placental area in females and
representative PAS-stained images showing the delimitation of placental areas. **(D)** Fetal
weight, fetal head weight, placental weight, and placental efficiency in male fetuses. **(E)**
Quantification of the areas of the Lz, Jz, and total placental area in males and representative
PAS-stained images showing the delimitation of placental areas. Data are displayed as the mean
+ SEM. and analyzed by one-way ANOVA with Tukey post hoc comparisons. (n=15). *p <
0.05, **p < 0.005.

Figure 2. Nutrient transport in control and ZIKV-infected female placentas. (A) Slc2a1
and Slc2a3 mRNA expression by qPCR (n=8-10). **(B)** Slc38a1 and Slc38a2 mRNA expression
by qPCR (n=8-10). **(C)** Quantification and representative photomicrographs of Glut1

expression in the placental transport zone by immunohistochemistry (n=5). **(D)** Quantification

and representative photomicrographs of Snat2 expression in the placental transport zone by

immunohistochemistry (n=5). **(E)** Matrix-assisted laser desorption/ionization mass

spectrometry imaging analysis showing the distribution of glucose-6-phosphate by 259.022

m/z localization in placental tissue and the distribution of glutamine by 145.061 m/z

localization in placental tissue (n=5). Data are displayed as the mean + SEM. and analyzed by

one-way ANOVA with Tukey post hoc comparisons. $*p < 0.05$, $**p < 0.005$.

Figure 3. Nutrient transport in control and ZIKV-infected male placentas. (A) *Slc2a1* and

Slc2a3 mRNA expression by qPCR (n=8-10). **(B)** *Slc38a1* and *Slc38a2* mRNA expression by

qPCR (n=8-10). **(C)** Quantification and representative photomicrographs of Glut1 expression

in the placental transport zone by immunohistochemistry (n=5). **(D)** Quantification and

representative photomicrographs of Snat2 expression in the placental transport zone by

immunohistochemistry (n=5). **(E)** Matrix-assisted laser desorption/ionization mass

spectrometry imaging analysis showing the distribution of glucose-6-phosphate by 259.022

m/z localization in placental tissue and the distribution of glutamine by 145.061 m/z

localization in placental tissue (n=5). Data are displayed as the mean + SEM. and analyzed by

one-way ANOVA with Tukey post hoc comparisons. $*p < 0.05$.

Supplementary figure 1. Nutrient transport in the junctional zone of control and ZIKV-

infected placentas. (A) Quantification and representative photomicrographs of Glut1

expression in the placental junctional zone by immunohistochemistry in female placentas

(n=5). **(B)** Quantification and representative photomicrographs of Snat2 expression in the

placental junctional zone by immunohistochemistry in female placentas (n=5). **(C)**

Quantification and representative photomicrographs of Glut1 expression in the placental

junctional zone by immunohistochemistry in male placentas (n=5). **(D)** Quantification and

representative photomicrographs of Snat2 expression in the placental junctional zone by

immunohistochemistry in male placentas (n=5). Data are displayed as the mean + SEM. and

analyzed by one-way ANOVA with Tukey post hoc comparisons.

Figure 4. HBP and O-GlcNAcylation in control and ZIKV-infected female placentas. (A)

Gfpt1 mRNA expression by qPCR (n=8-10). **(B)** Relative abundance of GFAT1 and GFAT2

(n=5-8). **(C)** Matrix-assisted laser desorption/ionization mass spectrometry imaging analysis

showing the distribution of UDP-GlcNAc by 606.0742 m/z localization in placental tissue

(n=5). **(D)** *Ogt* and *Oga* mRNA expression by qPCR (n=8-9). **(E)** Relative abundance of OGT,

OGA and O-GlcNAcylation (n=6-8). Data are displayed as the mean + SEM. and analyzed by

one-way ANOVA with Tukey post hoc comparisons. $*p < 0.05$, $**p < 0.005$

Figure 5. HBP and O-GlcNAcylation in control and ZIKV-infected male placentas. (A)

Gfpt1 mRNA expression by qPCR (n=6-8). **(B)** Relative abundance of GFAT1 and GFAT2

(n=7-8). (**C**) Matrix-assisted laser desorption/ionization mass spectrometry imaging analysis

showing the distribution of UDP-GlcNAc by 606.0742 m/z localization in placental tissue

(n=5). **(D)** *Ogt* and *Oga* mRNA expression by qPCR (n=7-8). **(E)** Relative abundance of OGT,

OGA and O-GlcNAcylation (n=6). Data are displayed as the mean + SEM. and analyzed by

one-way ANOVA with Tukey post hoc comparisons. $*p < 0.05$.

Supplementary table 1. List of primary antibodies used in this study.

Primary antibody	Host	Manufacturer, catalog number	Dilution
Actina	*Mouse*	Sigma–Aldrich, A5441	1/2000
Glut1	*Rabbit*	Invitrogen, PA5-16793	1/2000
Snat2	*Rabbit*	Bioss, bs-12125R	1/100
Gfat1	*Rabbit*	Santa Cruz Biotechnology, SC- 134894	1/1000
Gfat2	*Rabbit*	Santa Cruz Biotechnology, SC- 134710	1/1000
OGT	*Mouse*	Santa Cruz Biotechnology, (F-12), SC-74546	1/2000
OGA	*Rabbit*	Santa Cruz Biotechnology, 345	1/2000
O-GlcNAc	*Mouse*	Santa Cruz Biotechnology, RL-2, SC59624	1/1000

519   **Supplementary table 2.** List of primers used for quantitative RT–PCR

Gene	Genebank ID	Forward Primer (5′–3′)	Reverse Primer (5′–3′)
Ywhaz	NM_001253806.1	GAAAAGTTCTTGATCCCCAATGC	TGTGACTGGTCCACAATTCCTT
PPib	NM_011149.2	GAGACTTCACCAGGGG	CTGTCTGTCTTGGTGCTCTC
Bactin	NM_007393.5	GCAGGAGTACGATGAGTCCG	ACGCAGCTCAGTAACAGTCC
Slc2a1	NM_011400.3	CCAGCTGGGAATCGTCGTT	CAAGTCTGCATTGCCCATGAT
Slc2a3	XM_036165983.1	CTCTTCAGGTCACCCAACTACGT	CCGCGTCCTTGAAGATTCC
Slc381	NM_001166458.1	GGACGGAGATAAAGGCACTC	CAGAGGGATGCTGATCAAGG
Slc382	NM_175121.4	ACCTTTGGTGATCAAGGCAT	AGGACCAGATAGTCACCGTT
Uggt1	NM_198899.2	GCCTTACCGTGCCTGTTTTC	GTCTTCTGCTAAGAACTCACTGG
Gfpt2	NM_013529.3	TTTGCCAAGTGCCAGAATGC	GAGACAGTCCACTGTGTGGG
Oga	NM_023799.5	AAGCTTCTACCTGGAATTGA	GCATGAATGTTATCCCAGAT
Ogt	NM_139144.4	ACTTCCAGTGTCGAAGGCTG	AGGCTTCTGCTAGAAGGGGA

521   **Competing interests:** The authors have declared that no competing interests exist.
522
Funding: The research was undertaken in our laboratories and supported, in whole or in part, by the
following agencies: Bill & Melinda Gates Foundation [05/2013; OPP1107597]. D.P.-C. was supported
by Programa Institucional de Internacionalização (PRINT) (grant number: 88887.508140/2020-00) and
Coordenação de Aperfeiçoamento de Pessoal de Nível Superior—Brasil (CAPES—001). A.C.C.V.,
A.F.D., and V.M.O.N. were supported by CAPES—001. E.B. is supported by The National Council for
Scientific and Technological Development (Conselho Nacional de Desenvolvimento Científico e
Tecnológico [CNPq]: 10578/2020-5). S.V.A.C. was supported by Fundação de Amparo à Pesquisa do
Estado do Rio de Janeiro (FAPERJ). L.B.A is supported by Rede Corona-ômica BR MCTI/FINEP
affiliated to Rede Vírus/MCTI (FINEP = 01.20.0029.000462/20), CNPq 404096/2020-4); Carlos
Chagas Filho Research Support Foundation (FAPERJ; LBA E-26/201.206/2021, E-26/210.371/2019).
A.R.T. is supported by FAPERJ and CNPq. W.B.D. is supported by FAPERJ (E-26/210.549/2019-
APQ1). T.M.O.-C is supported by CNPq 306525/2019-4) and FAPERJ (CNE/E26/292798/2018).
535

References:

ANDRADE, C. B. V. et al. ZIKV Disrupts Placental Ultrastructure and Drug Transporter
Expression in Mice. **Front Immunol,** v. 12, p. 680246, 2021.

ANDRADE, T. A. et al. In Utero Exposure to Zika Virus Results in sex-Specific Memory
Deficits and Neurological Alterations in Adult Mice. **ASN Neuro,** v. 14, p.
17590914221121257, Jan-Dec 2022.

ANGELOVA, M. et al. Inhibition of O-Linked N-Acetylglucosamine Transferase Reduces
Replication of Herpes Simplex Virus and Human Cytomegalovirus. **J Virol,** v. 89, n. 16, p.
8474-83, Aug 2015.

BLOISE, E. et al. Impaired placental nutrient transport in mice generated by in vitro
fertilization. **Endocrinology,** v. 153, n. 7, p. 3457-67, Jul 2012.

BORGES, M. H. et al. Human placental GLUT1 glucose transporter expression and the fetal
insulin-like growth factor axis in pregnancies complicated by diabetes. **Biochim Biophys
Acta Mol Basis Dis,** v. 1865, n. 9, p. 2411-2419, Sep 1 2019.

BURTON, G. J.; JAUNIAUX, E. Pathophysiology of placental-derived fetal growth restriction.
Am J Obstet Gynecol, v. 218, n. 2S, p. S745-S761, Feb 2018.

CHANDRASIRI, U. P. et al. Insight into the pathogenesis of fetal growth restriction in
placental malaria: decreased placental glucose transporter isoform 1 expression. **J Infect
Dis,** v. 209, n. 10, p. 1663-7, May 15 2014.

CHEN, Q. et al. Metabolic reprogramming by Zika virus provokes inflammation in human
placenta. **Nat Commun,** v. 11, n. 1, p. 2967, Jun 11 2020.

CHIARADONNA, F.; RICCIARDIELLO, F.; PALORINI, R. The Nutrient-Sensing Hexosamine
Biosynthetic Pathway as the Hub of Cancer Metabolic Rewiring. **Cells,** v. 7, n. 6, Jun 2
2018.

COAN, P. M. et al. Adaptations in placental phenotype support fetal growth during
undernutrition of pregnant mice. **J Physiol,** v. 588, n. Pt 3, p. 527-38, Feb 1 2010.

COELHO, S. V. A. et al. Development of standard methods for Zika virus propagation,
titration, and purification. **J Virol Methods,** v. 246, p. 65-74, Aug 2017.

CONNOR, K. L. et al. Maternal malnutrition impacts placental morphology and transporter
expression: an origin for poor offspring growth. **J Nutr Biochem,** v. 78, p. 108329, Apr 2020.

CREISHER, P. S. et al. Downregulation of transcriptional activity, increased inflammation,
and damage in the placenta following in utero Zika virus infection is associated with adverse
pregnancy outcomes. **Front Virol,** v. 2, Feb 2022.

DAS, U. G. et al. Time-dependent physiological regulation of rodent and ovine placental
glucose transporter (GLUT-1) protein. **Am J Physiol,** v. 274, n. 2, p. R339-47, Feb 1998.

DAVALIEVA, K. et al. Non-invasive fetal sex determination using real-time PCR. **J Matern
Fetal Neonatal Med,** v. 19, n. 6, p. 337-42, Jun 2006.

DELA JUSTINA, V. et al. Increased O-Linked N-Acetylglucosamine Modification of NF-
KappaB and Augmented Cytokine Production in the Placentas from Hyperglycemic Rats.
Inflammation, v. 40, n. 5, p. 1773-1781, Oct 2017.

DONG, H.; LIU, Z.; WEN, H. Protein O-GlcNAcylation Regulates Innate Immune Cell
Function. **Front Immunol,** v. 13, p. 805018, 2022.

FONTES, K. N. et al. Dysregulation of placental ABC transporters in a murine model of
malaria-induced preterm labor. **Sci Rep,** v. 9, n. 1, p. 11488, Aug 7 2019.

GACCIOLI, F.; LAGER, S. Placental Nutrient Transport and Intrauterine Growth Restriction.
Front Physiol, v. 7, p. 40, 2016.

HART, B.; MORGAN, E.; ALEJANDRO, E. U. Nutrient sensor signaling pathways and
cellular stress in fetal growth restriction. **J Mol Endocrinol,** v. 62, n. 2, p. R155-R165, Feb 1
2019.

HART, G. W. et al. Cross talk between O-GlcNAcylation and phosphorylation: roles in
signaling, transcription, and chronic disease. **Annu Rev Biochem,** v. 80, p. 825-58, 2011.

HIGGINS, J. S. et al. Placental phenotype and resource allocation to fetal growth are
modified by the timing and degree of hypoxia during mouse pregnancy. **J Physiol,** v. 594, n.
5, p. 1341-56, Mar 1 2016.

HOWERTON, C. L.; BALE, T. L. Targeted placental deletion of OGT recapitulates the
prenatal stress phenotype including hypothalamic mitochondrial dysfunction. **Proc Natl
Acad Sci U S A,** v. 111, n. 26, p. 9639-44, Jul 1 2014.

HOWERTON, C. L. et al. O-GlcNAc transferase (OGT) as a placental biomarker of maternal
stress and reprogramming of CNS gene transcription in development. **Proc Natl Acad Sci U
S A,** v. 110, n. 13, p. 5169-74, Mar 26 2013.

LAGER, S.; POWELL, T. L. Regulation of nutrient transport across the placenta. **J
Pregnancy,** v. 2012, p. 179827, 2012.

LANGDOWN, M. L.; SUGDEN, M. C. Enhanced placental GLUT1 and GLUT3 expression in
dexamethasone-induced fetal growth retardation. **Mol Cell Endocrinol,** v. 185, n. 1-2, p.
109-17, Dec 20 2001.

LI, P. et al. Intrauterine Growth Restriction Induces Adulthood Chronic Metabolic Disorder in Cardiac and Skeletal Muscles. **Front Nutr,** v. 9, p. 929943, 2022.

LIVAK, K. J.; SCHMITTGEN, T. D. Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. **Methods,** v. 25, n. 4, p. 402-8, Dec 2001.

LUSCHER, B. P. et al. Placental glucose transporter (GLUT)-1 is down-regulated in preeclampsia. **Placenta,** v. 55, p. 94-99, Jul 2017.

MESSINA, J. P. et al. Mapping global environmental suitability for Zika virus. **Elife,** v. 5, Apr 19 2016.

MONTEIRO, V. R. S. et al. Mid-pregnancy poly(I:C) viral mimic disrupts placental ABC transporter expression and leads to long-term offspring motor and cognitive dysfunction. **Sci Rep,** v. 12, n. 1, p. 10262, Jun 17 2022.

MOORE, M. et al. Disruption of O-Linked N-Acetylglucosamine Signaling in Placenta Induces Insulin Sensitivity in Female Offspring. **Int J Mol Sci,** v. 22, n. 13, Jun 28 2021.

MULKEY, S. B. et al. Neurodevelopmental Abnormalities in Children With In Utero Zika Virus Exposure Without Congenital Zika Syndrome. **JAMA Pediatr,** v. 174, n. 3, p. 269-276, Mar 1 2020.

NARDOZZA, L. M. et al. Fetal growth restriction: current knowledge. **Arch Gynecol Obstet,** v. 295, n. 5, p. 1061-1077, May 2017.

NEM DE OLIVEIRA SOUZA, I. et al. Acute and chronic neurological consequences of early-life Zika virus infection in mice. **Sci Transl Med,** v. 10, n. 444, Jun 6 2018.

NUGENT, B. M. et al. Placental H3K27me3 establishes female resilience to prenatal insults. **Nat Commun,** v. 9, n. 1, p. 2555, Jul 2 2018.

OLIVEIRA MELO, A. S. et al. Zika virus intrauterine infection causes fetal brain abnormality and microcephaly: tip of the iceberg? **Ultrasound Obstet Gynecol,** v. 47, n. 1, p. 6-7, Jan 2016.

PAUL, A. M. et al. Congenital Zika Virus Infection in Immunocompetent Mice Causes Postnatal Growth Impediment and Neurobehavioral Deficits. **Front Microbiol,** v. 9, p. 2028, 2018.

QUIK, M.; HOKKE, C. H.; EVERTS, B. The role of O-GlcNAcylation in immunity against infections. **Immunology,** v. 161, n. 3, p. 175-185, Nov 2020.

SAKATA, M. et al. Increase in human placental glucose transporter-1 during pregnancy. **Eur J Endocrinol,** v. 132, n. 2, p. 206-12, Feb 1995.

SALAZAR-PETRES, E. et al. Placental structure, function, and mitochondrial phenotype relate to fetal size in each fetal sex in micedagger. **Biol Reprod,** v. 106, n. 6, p. 1292-1311, Jun 13 2022.

SFERRUZZI-PERRI, A. N.; CAMM, E. J. The Programming Power of the Placenta. **Front Physiol,** v. 7, p. 33, 2016.

682 SHERER, M. L. et al. A Rat Model of Prenatal Zika Virus Infection and Associated Long-
683 Term Outcomes. **Viruses,** v. 13, n. 11, Nov 18 2021.
684
685 SHI, Y. et al. O-GlcNAcylation stabilizes the autophagy-initiating kinase ULK1 by inhibiting
686 chaperone-mediated autophagy upon HPV infection. **J Biol Chem,** v. 298, n. 9, p. 102341,
687 Sep 2022.
688
689 SIBIAK, R. et al. Fetomaternal Expression of Glucose Transporters (GLUTs)-Biochemical,
690 Cellular and Clinical Aspects. **Nutrients,** v. 14, n. 10, May 12 2022.
691
692 SUZUKAWA, A. A. et al. Downregulation of IGF2 expression in third trimester placental
693 tissues from Zika virus infected women in Brazil. **J Infect,** v. 81, n. 5, p. 766-775, Nov 2020.
694
695 SZABA, F. M. et al. Zika virus infection in immunocompetent pregnant mice causes fetal
696 damage and placental pathology in the absence of fetal infection. **PLoS Pathog,** v. 14, n. 4,
697 p. e1006994, Apr 2018.
698
699 THAKER, S. K. et al. Differential Metabolic Reprogramming by Zika Virus Promotes Cell
700 Death in Human versus Mosquito Cells. **Cell Metab,** v. 29, n. 5, p. 1206-1216 e4, May 7
701 2019.
702
703 TRUS, I. et al. Subclinical in utero Zika virus infection is associated with interferon alpha
704 sequelae and sex-specific molecular brain pathology in asymptomatic porcine offspring.
705 **PLoS Pathog,** v. 15, n. 11, p. e1008038, Nov 2019.
706
707 VAUGHAN, O. R. et al. Placenta-specific Slc38a2/SNAT2 knockdown causes fetal growth
708 restriction in mice. **Clin Sci (Lond),** v. 135, n. 17, p. 2049-2066, Sep 17 2021.
709
710 VOTA, D. et al. Zika virus infection of first trimester trophoblast cells affects cell migration,
711 metabolism and immune homeostasis control. **J Cell Physiol,** v. 236, n. 7, p. 4913-4925, Jul
712 2021.
713
714 WANG, Q. et al. O-GlcNAc transferase promotes influenza A virus-induced cytokine storm
715 by targeting interferon regulatory factor-5. **Sci Adv,** v. 6, n. 16, p. eaaz7086, Apr 2020.
716
717 WANG, X. et al. O-GlcNAcylation modulates HBV replication through regulating cellular
718 autophagy at multiple levels. **FASEB J,** v. 34, n. 11, p. 14473-14489, Nov 2020.
719
720 WINTERHAGER, E.; GELLHAUS, A. Transplacental Nutrient Transport Mechanisms of
721 Intrauterine Growth Restriction in Rodent Models and Humans. **Front Physiol,** v. 8, p. 951,
722 2017.
723
724 YANG, M. et al. Dysregulated GLUT1 may be involved in the pathogenesis of preeclampsia
725 by impairing decidualization. **Mol Cell Endocrinol,** v. 540, p. 111509, Jan 15 2022.
726
727 YANG, Y. R. et al. O-GlcNAc cycling enzymes control vascular development of the placenta
728 by modulating the levels of HIF-1alpha. **Placenta,** v. 36, n. 10, p. 1063-8, Oct 2015.
729
730 YAO, G. et al. GDM-Induced Macrosomia Is Reversed by Cav-1 via AMPK-Mediated Fatty
731 Acid Transport and GLUT1-Mediated Glucose Transport in Placenta. **PLoS One,** v. 12, n. 1,
732 p. e0170490, 2017.
733
734 YAU, C. et al. Dysregulated metabolism underpins Zika-virus-infection-associated
735 impairment in fetal development. **Cell Rep,** v. 37, n. 11, p. 110118, Dec 14 2021.
736

737   ZENG, Q. et al. O-linked GlcNAcylation elevated by HPV E6 mediates viral oncogenesis.
738   **Proc Natl Acad Sci U S A,** v. 113, n. 33, p. 9333-8, Aug 16 2016.
739
740   ZHANG, S. et al. Placental adaptations in growth restriction. **Nutrients,** v. 7, n. 1, p. 360-89,
741   Jan 8 2015.
742
743