The positioning mechanics of microtubule asters in *Drosophila* embryo explants

Jorge de-Carvalho[1], Sham Tlili[2,#], Timothy E. Saunders[2,3,4,*], Ivo A. Telley[1,*]

[1] Instituto Gulbenkian de Ciência, Fundação Calouste Gulbenkian, Oeiras, Portugal

[2] Mechanobiology Institute and Department of Biological Sciences, National University of Singapore, Singapore

[3] Institute of Molecular and Cellular Biology, A*Star, Proteos, Singapore

[4] Centre for Mechanochemical Cell Biology, Warwick Medical School, University of Warwick, UK

[*] For correspondence: timothy.saunders@warwick.ac.uk and itelley@igc.gulbenkian.pt

[#] Present address: Aix-Marseille Université | AMU · Institut de Biologie du Développement de Marseille-Luminy (UMR 7288 IBDML)

Keywords

Centrosome, cytoskeleton, ex vivo, microtubules, active systems

Abstract:

Microtubule asters are essential in localizing the action of microtubules in processes including mitosis and cell positioning. In large cells, such as the one-cell sea urchin embryo, aster dynamics are dominated by hydrodynamic pulling forces. However, in systems with more densely positioned nuclei such as the early *Drosophila* embryo, which packs around 6000 nuclei within the syncytium in a crystalline-like order, it is unclear what processes dominate aster dynamics. Here, we take advantage of a cell cycle regulation *Drosophila* mutant to generate embryos with multiple asters, independent from nuclei. We use an *ex vivo* assay to further simplify the system to explore the forces generated by and between asters. Through live imaging, drug and mechanical perturbations, and theoretical modelling, we demonstrate that these asters likely generate an effective pushing force over short distances. Such a pushing force can position asters in a crystalline like order, consistent with experiment. This work demonstrates that the mechanism of aster interactions is likely system size dependent.

Significance Statement:

Using cytosolic explants from *Drosophila* syncytial embryos combined with quantitative microscopy and perturbations, de-Carvalho et al. reveal the mechanical forces separating microtubule asters. Aster separation drives precise nuclear positioning in multinucleated embryo cells, a vital process for tissue formation and gene expression during subsequent embryo development.

Introduction

Eukaryotes assemble a cytoskeletal structure of microtubules (MTs) called an 'aster', which is involved in critical cell functions including intracellular positioning and organelle transport. In metazoan cells, the aster acquires a radial shape in which MTs are focused by the centrosome and emanate towards the cell periphery (Wilson, 1986). Microtubules grow at the centrosome and at microtubule-based nucleation sites (Ishihara et al., 2016). The centrosome is also found at the two focus points of the mitotic spindle – the poles – linking the aster structurally and mechanically to the spindle, thus lending it a decisive role during cell division (Hinchcliffe and Sluder, 2001; Hoffmann, 2021). Positioning of the spindle by astral microtubules contacting the cell cortex determines the cell wall cleavage location (Field et al., 2015; Mitchison et al., 2012; Rappaport, 1969, 1961). During fertilization the maternal and paternal genomes are united by the assembly and migration of the sperm aster, which facilitates transport of the female towards the male pronucleus (Chambers, 1939; Hamaguchi and Hiramoto, 1986; Riparbelli et al., 2000). Where the sperm aster positions inside the egg will roughly define the geometry of the first embryonic cleavage (Albertson, 1984; Hamaguchi and Hiramoto, 1980; Hirano and Ishikawa, 1979). Interestingly, in these large egg cells the aster migration can also occur without contacting the cell periphery (Wühr et al., 2010). Uninuclear cells contain one or, at most, two asters depending on the cell cycle stage.

The mechanics of aster movement has been a matter of long discourse, so far without any consensus on a unified biophysical model (Deshpande and Telley, 2021). Disagreement remains likely because the experimental insight stems from different model organisms and cells which, by nature, have evolved divergent mechanisms of aster positioning. One fundamental question, whether asters position themselves by pulling on or by pushing at the rigid cell periphery, remains extensively debated (Garzon-Coral et al., 2016; Grill and Hyman, 2005; Meaders et al., 2020; Sulerud et al., 2020). Recently, an adaptation of the pulling model for cortex contact–free asters in large cells has gained new momentum (Hamaguchi and Hiramoto, 1980; Simone et al., 2018; Tanimoto et al., 2018; Xie et al., 2022). In this model, the net force moving the aster originates from a balance of viscous drag forces of moving organelles along astral microtubules with difference length. Overall geometric asymmetry of the aster is a key point of this model (Tanimoto et al., 2016).

An interesting yet understudied case of higher complexity is the positioning of multiple nuclei and their associated asters in multinucleated cells (coenocytes). In the syncytial embryo of

insects, such as the fruit fly, the genome is rapidly proliferated and distributed without cell cleavage, leading to a single large cell with hundreds of nuclei (Donoughe and Extavour, 2016; Foe and Alberts, 1983; Sommer and Tautz, 1991). The positioning of these syncytial nuclei depends on the centrosome-nucleated microtubule aster (de-Carvalho et al., 2022; Megraw et al., 1999; Telley et al., 2012) and microtubule-associated crosslinking proteins (Deshpande et al., 2021). Separation of more than two asters in the absence of intermittent cell cortex boundary formation poses an additional challenge to the cell. Besides pulling or pushing on the cell cortex, a net force is required to evenly and stably separate each aster from its neighbors, of which there are on average six (Kanesaki et al., 2011). Recent work from us has shown that short-ranged aster repulsion can position nuclei in a regular order within the syncytium (de-Carvalho et al., 2022). However, experimental testing of these forces is still missing.

Here, by exploiting embryonic explants (Telley et al., 2013), which reduces complexity, and a cell cycle regulation mutant (Freeman et al., 1986; Lee et al., 2003) to uncouple microtubule organization from nuclear division, we have studied the dynamics of aster-aster interaction bottom-up. In this system, we uncover the physical principles of separation for simple aster arrays, reveal the positional autonomy of single asters, and derive the microtubule-associated mechanical separation potential. Our work reveals the underlying local biophysical interactions between the structures that order multiple components within a rapidly proliferating system.

Results

Asters are radial microtubule structures nucleated by the centrosome, which acts as microtubule organizing center (MTOC) as part of the mitotic spindle pole. In the early *Drosophila* embryo, the MTOC is physically connected to the nucleus throughout the cell cycle. To uncouple nucleus and aster associated forces we took advantage of *giant nuclei (gnu)* mutant *Drosophila* embryos. In these embryos, DNA endoreplication occurs without mitosis as the cell cycle is arrested in interphase. This results in the embryo having only one or just a few polyploid nuclei. Yet, centrosome maturation and duplication continue in *gnu* embryos (Freeman et al., 1986; Freeman and Glover, 1987; Lee et al., 2003) (Fig. 1A). We produced embryo explants from *gnu* mutant embryos (Telley et al., 2013) (Fig. 1B) and studied the positioning properties of a small number of microtubule asters in quasi-2D spaces in the absence of nuclei.

Single aster positioning is consistent with radially symmetric forces

Initially, we focused on explants containing a single aster. A single aster consistently moved away from the explant boundary and eventually adopted a steady position (Fig. 1C). In a series

of experiments, we deposited extract and waited 30 min, after which we measured the shortest distance b of the centrosome from the boundary at steady state. Initial aster position within the aster is random, so we expected to observe some aster positions that were already near the center. Thus, the distance between the aster and the boundary, b, varied between $R/2$ and R, R being the explant radius (Suppl. Fig. 1A). Deviation from precise centering ($b=R$) may be due to yolk or lipid droplets (green circles in Fig. 1C) forming exclusion zones. Next, we analyzed the kinematics of single asters located near the boundary after cytosol deposition. Typically, they stayed for up to 10 min (Fig. 1D, phase I), but they always eventually migrated (Video 1, left). Single asters moved rapidly after separation from the boundary, with a maximum velocity of 0.05 ± 0.02 μm/s (Fig. 1D, phase II), at around 20% of its final distance from the explant boundary (Fig. 1E), before linearly decelerating (Fig. 1D, phase III) and stopping 30 μm from the boundary at most (Suppl. Fig. 1B).

We next measured the radial intensity profile of single asters in explants as a proxy for aster size and microtubule length (Fig. 1F, inset). Away from the MTOC, the distribution was well approximated with a mono-exponential decay with decay length of ~12 μm (Fig. 1F). This value is in excellent agreement with the size of asters associated to telophase and early interphase nuclei of wildtype embryo explants (Telley et al., 2012). The signal drops to background level at ~30 μm, suggesting that only very few microtubules are longer. However, we noticed fewer microtubules oriented outwards when the aster was near the boundary (Suppl. Fig. 1C), suggesting that most existing microtubules or those growing from the centrosome buckle and orient along the boundary, or they depolymerize rather than stabilize at short length. Indeed, in some samples we observed splay of microtubules near the explant boundary (Fig. 1G), suggesting that microtubules push against the boundary.

In summary, single asters display three distinct dynamic phases: first, a very slow phase (at least along the radial axis) as they separate from the edge; second, a rapid motion away from the explant boundary; and finally, a gradually slowing down as they migrate towards the explant center. The initial phase of separation appears to be due to splay of microtubules near the boundary edge. Random fluctuations within the cytoplasm may be sufficient to release the aster eventually from the edge (transition from Phase I to Phase II, Fig. 1D). These observations are consistent with the asters generating a repulsive potential, that has a short-ranged cut-off.

Lipid droplet movements in extract are consistent with a repulsive aster potential

Embryos contain high amounts of lipid droplets, which serve as fiduciary markers in our explants to study hydrodynamic flow. We next examined the movement of these droplets as the aster moved through the explant (green droplets in Video 1). We plotted particle tracks relative to the aster position of the migrating aster (Fig. 2A). As the aster moved away from the boundary, the lipid droplets streamed around the aster. Strikingly, we observed a circular exclusion zone for droplets of ~10 μm diameter, which maintained during aster migration (Fig. 2A). Relative to aster movement, collateral droplets moved backwards (Fig. 2B). Next, we quantified the lipid droplet speed with and without an aster present (Fig. 2C). As an aster moved through the lipid droplets, their speed increased, before reducing once the aster came to a halt. This further supports that the aster is acting directly against the lipid droplets. Finally, we measured the root-mean-square (RMS) displacement of the lipid droplets with and without an aster present in the sample (Fig. 2D). In the absence of an aster, the lipid droplets moved in a manner consistent with diffusive-like (random) behavior ($x_{rms} \sim t^{1/2}$). In the presence of an aster, lipid droplets moved faster, and appeared to move in a more directed manner ($x_{rms} \sim t^{2/3}$). Such behavior is consistent with the aster having a repulsive force potential that can act on the surrounding lipid droplets and other boundary constraints. It is challenging to meld these observations with an effective pulling potential.

Aster-aster interactions are consistent with a repulsive potential involving short-ranged effective slipping

Some explants contained a pair of centrosomal asters that separated and adopted a steady-state inter-aster distance (Fig. 3A and Video 1, right). Inter-aster interaction, if existing, must balance with the force involved in moving asters away from the boundary. If the forces between the asters and the droplet boundary are equal, we would expect the asters to be positioned a distance R apart (Methods). Interestingly, the two asters instead approximately partitioned the available space ($d = b_1 = b_2 = 2R/3$) (Suppl. Fig. 1D). This implies that the force associated with aster-boundary interactions is larger than aster-aster directed forces.

Given the above observation, we predicted that the dynamics between aster pairs will differ from the single aster dynamics (Fig. 1D). Using live imaging of aster doublets, we noticed that the peak separation velocity was always near half the final aster separation distance (Fig. 3B), independent of final separation distance. This suggests that the resulting magnitude of the forces are similar in the initial phase of separation and the eventual reaching of the equilibrium position. The dynamic profile is quite distinct from the single aster scenario.

Given this eccentric movement of two asters, the aster separation could be driven by overlap and sliding of astral microtubules (Baker et al., 1993; Deshpande et al., 2021; Lv et al., 2018), or by mutual contact leading to repulsion by microtubules of both asters. Thus, we quantified the microtubule intensity between the separating asters (Fig. 3C, left) and generated kymographs of the microtubule fluorescence intensity along the separation axis (Fig. 3C, right). The intensity at half the separation distance decayed exponentially (Fig. 3D), consistent with models of dynamic microtubule length distribution (Howard, 2001; Jeune-Smith and Hess, 2010). When aster separation ceased there was almost no detectable microtubule signal between the asters.

For a viscous material, the velocity, v, of a submerged object depends on the applied force F: $v \approx \gamma F$, where γ is the effective viscous drag coefficient. Naïvely interpreting the microtubule distribution as the resulting force profile does not match with the observed separation velocity profile. However, multiplying the microtubule distribution by an effective *slipping* term, $f_{slip} = f_0 \frac{x^2}{x_0^2 + x^2}$ ($x_0 \approx 15 \ \mu m$), results in an excellent fit to our observed aster separation velocities (Fig. 3E). We attribute this effective slipping to molecular friction between microtubules (Forth et al., 2014).

Overall, we see that aster-aster and aster-boundary dynamics both appear to involve repulsive interactions with effective slipping at very short distances, though the aster-aster interactions are weaker than those between the asters and the boundary. The two asters may mechanically interact via crosslinking of microtubule overlaps (Bieling et al., 2010; Deshpande et al., 2021; Lv et al., 2018; Subramanian et al., 2013; Wijeratne and Subramanian, 2018), while astral microtubules may simply hit against the boundary interface, which acts as an immovable hard wall.

Perturbations of aster interaction are consistent with a microtubule-mediated repulsive force potential

To further explore the nature of microtubule aster interaction, we performed a series of inhibitory treatments to chemically perturb the interaction. Since small-molecule inhibitors for candidate molecular motors have no effect in *Drosophila* (Firestone et al., 2012; Maliga et al., 2002), we targeted microtubules and ATPases in general. We generated explants with two asters during separation and pulse-injected a defined volume of 200 μM colchicine, which causes acute depolymerization of microtubules. Upon injection the asters stopped separating and sometimes inverted their direction of motion (Fig. 4A, Video 2, right). We then tested whether active

molecular machinery was required for aster repulsion by inhibiting ATP consumption. We injected a series of concentrations of sodium azide into explants that contained a separating pair of asters (Video 2, middle). Adding sodium azide decreased the initial recoil velocity (dashed lines in Fig. 4A) and resulted in a considerable reduction in aster separation distance. However, even at very high concentrations of sodium azide, we still observed residual motion, suggesting that both actively driven microtubule-mediated separation and passive physical contact driven separation occur.

From our observations thus far, we conclude that aster separation is caused by a net repulsive force between asters and between aster and boundary. However, either overall pushing or pulling can cause the separation dynamics and centering (Grill et al., 2003; Laan et al., 2012; Zhu et al., 2010). Pulling within the cytoplasm requires aster asymmetry (Kimura and Kimura, 2011; Tanimoto et al., 2016). Thus, we performed targeted UV photo-ablation experiments in explants and generated ellipse-shaped ablations positioned asymmetrically around one aster, affecting microtubules on the left side more than on the right side of the aster (Fig. 4B, Video 3). If pulling on the boundary (Grill et al., 2003) or hydrodynamic effects from vesicle transport along microtubules (Tanimoto et al., 2016) drives aster motion, we expect a displacement to the right (positive) after ablation. Conversely, if the net force applied on microtubules favors pushing on the MTOC, we expect a displacement to the left (negative). Indeed, asters consistently moved to the left, supporting a dominating effect of microtubule-driven pushing (Fig. 6C). As a control, we performed the same perturbation in explants that were injected with the microtubule inhibitor colchicine (Fig. 6D). Under this condition, asters moved very slowly to the right (positive), which is consistent with a weak hydrodynamic effect from other contractile sources (*e.g.* actomyosin) (Keren, 2016). We conclude that a single aster moves and positions within *Drosophila* explants by microtubule-dependent pushing force.

To challenge these conclusions, we performed two types of UV ablation in explants containing two asters (Fig. 4E): 1) light pulses emitted along an ellipse around both asters to destroy microtubules in the periphery; 2) light pulses emitted along a line between the two asters to destroy microtubules between asters. If forces are attractive, then ablation type 1 will stop separation while ablation type 2 will lead to an acceleration. If forces are repulsive, we predict the opposite response. We found a slight acceleration for peripheral ablation (Video 4) and a strong deceleration with recovery for central ablation (Fig. 4F). Separation recovered likely because of fast re-growth of microtubules after ablation (in the range of μm/min) (Rogers et al.,

2002). In summary, the dynamic behavior of asters in our explants is consistent with a model of radially symmetric microtubule-based repulsion.

Simple repulsive model of aster interactions can replicate the observed aster behavior ex vivo

Next, we set out to formulate a physical model of aster repulsion from the experimental insights. We considered the asters as generating a radially symmetric repulsive potential, embedded within a 2D circular environment (Fig. 5A). The repulsive potential is taken to be exponentially decaying with length scale 15 μm (based on Fig. 1F). We also include a "slippage" term when asters are close to the boundary or each other, of the form $\frac{x^2}{x^2+x_0^2}$, where $x_0 = 18\mu m$ (aster-boundary) and $x_0 = 25\mu m$ (aster-aster), representing the apparently weaker effective force for aster-aster interactions. To match our observations on one- and two-aster cases, the repulsion from the explant boundary was 40% larger than the aster-aster repulsion (Methods). We also include a noise term due to the inherent stochasticity in the system.

We first tested whether this 2D model could replicate the observed dynamics and positioning in the one-aster case. We saw that the model can reproduce the observed dynamics of a single aster, so long as a "slipping" term is included (Fig. 5B). Likewise, we can solve the 1D equation of motion stochastically for the one-aster case (Methods). In this case, we found that the final aster position scales with droplet size up to around 50μm (Fig. 5C).

We next introduced a second aster into the 2D model. We had to account for interactions between aster-boundary and aster-aster. The two-aster scenario was defined by allowing a single aster to divide, then following the positions of each sister aster. We can reproduce the experimentally observed aster dynamics, with a more symmetric speed profile as compared with the one-aster case (Fig. 5D). Again, we can solve the equivalent 1D equation of motion stochastically and we found that this provides a good approximation of the aster positions (compare Fig. 5E and Fig. 5F).

So far, we have shown that our model can closely match experimental observations in the one- and two-aster scenarios. What about systems with greater than two asters? We previously showed that multi-aster samples form crystalline structures, *e.g.,* equilateral triangles with three asters and square-like distributions with four asters (de-Carvalho et al., 2022). Can our model replicate these observations? We ran our dynamic simulations with three of four initial asters randomly placed, until the asters reached equilibrium positions. In the three-aster scenario, most simulations resulted in the asters distributed such that they (approximately) formed the vertices

of an equilibrium triangle. Subsequently, the asters were distributed with angle 60º between each other (Fig. 5G). Likewise, most four-aster simulations resulted in the asters (approximately) forming the vertices of a square, with angle distribution 90º (Fig. 5H). These results are similar to those observed experimentally (de-Carvalho et al., 2022). Consistent with experiment, in a small subset of simulations the four asters formed a triangle with the fourth aster positioned away from the other three (upper right inset, Fig. 5H). In conclusion, our simple repulsive model can reproduce both the observed equilibrium aster distributions and the quantified aster dynamics in a range of scenarios.

Discussion

We have characterized the mechanics of microtubule aster positioning using an *ex vivo* model of the cellular context where, naturally, hundreds of these cytoskeletal structures co-exist and define the regular positioning of nuclei in a multinucleated cell. We generated single embryo explants from mutant *Drosophila* syncytial embryos in which mitosis is inhibited but centrosomes duplicate and divide, each giving rise to a radial microtubule array. This experimental reductionist approach of generating a proliferative aster system *ex vivo* has several advantages (de-Carvalho et al., 2022). First and foremost, it enables the study of the mechanics of single aster positioning (the "atomic" structure) under boundary conditions, and the canonical interaction between two structures. Our experiments support a mechanical model where asters generate a radially symmetric, repulsive force potential. The repulsive interaction towards the boundary is about twice as strong as compared to the repulsion between two asters. The range of action between asters is finite and productive for positioning only up to ~45 µm. We note that two asters positioned close (<3 µm) to each other, for example after centrosome disengagement, show dynamics of weak attraction. We highlight that, by design, our experiments in embryo explants resolve the canonical aster–aster interaction whereas additional cell cortex interactions and more complex boundary conditions imposed by the cell membrane may occur in the intact embryo (Foe et al., 2000; Postner et al., 1992; Winkler et al., 2015).

There has been substantial discussion over the nature of aster force potentials (Garzon-Coral et al., 2016; Meaders et al., 2020; Minc et al., 2011; Mitchison et al., 2012; Pelletier et al., 2020; Sulerud et al., 2020; Tanimoto et al., 2018): are they repulsive or attractive; what range do they act over; and are there different regimes of action depending on temporal or spatial constraints? Here, we show that asters derived from the syncytial embryo of *Drosophila* display a short-range repulsive potential. At very short distances, this potential tends to zero (or arguably even

attractive), likely due to microtubules being unable to form linearly. A simple model can replicate our observations without requiring additional assumptions. Despite this being a highly active system, it displays long ranged and reproducible structural order. It is worth noting that the length scales here (typically 3–10 μm between asters in the embryo) are substantially smaller than those in other model systems used to explore aster dynamics, such as the sea urchin (>50 μm). In that system, effective pulling forces generated by hydrodynamic processes dominate the aster positioning (Tanimoto et al., 2018). Therefore, our results suggest that the mechanism driving aster positioning may well be system size dependent, *i.e.,* there is not a 'one size fits all' model of aster positioning.

What is the mechanism underlying the "slippage" at close distances, both for aster-aster and aster-boundary interactions? We observe splay in the microtubule distribution near the boundary. If microtubules are generating a mechanical pushing force (*i.e.* like a rod being pushed against a wall), such splay would be expected. Microtubule polymerization at a boundary generates forces against the boundary in the range that can lead to buckling (Holy et al., 1997; Howard, 2001). Conversely, it is challenging to reconcile this observation with astral microtubules (and linker proteins) generating a local pulling force (Grill et al., 2003). Between two asters, slippage may be attributed to molecular friction between microtubules (Forth et al., 2014). Single molecule experiments showed that antiparallel microtubule crosslinking, and their sliding by molecular motor activity or entropic effects, results in viscoelastic properties of the microtubule pair as a mechanical element (Shimamoto et al., 2015). At the level of two asters, hundreds of such elements in parallel accumulate to a "softened" repulsion, overcoming any opposing forces on each aster (boundary constraints, viscosity of cytoplasm), leading to aster separation.

We emphasize that the molecular mechanism underlying the generation of the pushing forces remains unclear, in part due to the lack of suitable reagents for targeted perturbation of molecular motors in *Drosophila*. However, our finding that aster separation occurs despite inhibition of ATPases (*e.g.*, motors) is both interesting and intriguing. It is unlikely due to mutual contact between microtubules growing from opposite asters since the probability for such encounters to happen and to be mechanically effective is extremely low. In our view, this "passive" repulsion force is likely caused by entropy driven microtubule sliding by ATP independent crosslinkers (Forth et al., 2014). If possible, targeted inhibition of candidate microtubule associated proteins should give further insight.

Materials and Methods

Fly strains

Flies expressing a fluorescent reporter for microtubules and the centrosome were generated by recombination of the genotypes w*; pUbq>RFP::β2-Tubulin; + (Inoue et al., 2004) and w[1118]; pUbq>Spd2::GFP; + (provided by M. Bettencourt Dias, IGC, Portugal). Two different mutants of giant nucleus (*gnu*), namely w*; +; gnu[305]/TM3 (discontinued stock no. 3321; Bloomington) and w*; +; gnu[Z3-3770A]/TM3 (discontinued stock no. 38440; Bloomington), were each balanced with w[1118]; CyO/Sco; MKRS/TM6B (stock no. 3703, Bloomington). Above-described recombined line on the 2nd chromosome were individually crossed with gnu mutants and kept as balanced stocks. Finally, trans-heterozygous were generated for gnu[305]/gnu[Z3-3770A] mutants, whereby only flies homozygous for the fluorescent reporters on the 2nd chromosome were selected for increased signal collection during live microscopy. These trans-heterozygotes laid fertilized eggs which undergo several embryonic rounds of chromatin replication and centrosome duplication, allowing for the study and quantification of asters at the embryo cortex.

Embryo collection and sample preparation

We followed established procedures of fly husbandry (Schubiger and Edgar, 1994), keeping flies at 25°C under 50-60% humidity. For embryo collections, young adult flies were transferred to a cage coupled to an apple juice agar plate. After 2–3 rounds of egg laying synchronization, developing embryos were collected every 30–60 minutes. In the case of *gnu* mutants, embryos were collected at different time intervals, ranging from 30 min up to 4h. Embryos were dechorionated by short immersion in 7% sodium hypochlorite solution (VWR). After extensive rinsing with water, embryos were aligned and immobilized in a thin strip of heptane glue placed on 22x22mm coverslips, and covered with halocarbon oil (Voltalef 10S, Arkema).

Microscopy

Time-lapse acquisitions were conducted on a Nikon Eclipse Ti-E microscope equipped with a Yokogawa CSU-W Spinning Disk confocal scanner and a piezoelectric stage (737.2SL, Physik Instrumente). For embryo imaging, 15 µm (31 planes) Z-series stacks were acquired every 15s (wildtype, if not states else) or 30s (*gnu* mutant), using a Plan Fluor 40x 1.3NA oil immersion objective, the 488nm and 561nm laser lines, and an Andor Zyla 4.2 sCMOS camera to acquire images. For explants up to 100µm in diameter, we used a Plan Apo VC 60x 1.2NA water immersion objective with 2x post-magnification and an Andor iXon3 888 EMCCD camera. When needed, the Andor Zyla 4.2 sCMOS camera was selected to acquire a 2x wider field of view with the same spatial resolution or, alternatively, the Apo λ S LWD 40x 1.15NA water

immersion objective. For acquisition in explants, the frame rate was 15s for *gnu* mutant 30 s for wildtype embryo explants.

Single embryo explant assay

Embryo extractions were performed as previously described (de-Carvalho et al., 2018; Telley et al., 2013). Briefly, cytosol from wild-type embryos was extracted by puncturing the vitelline membrane with a sharp glass micropipette and flow control by operating a bi-directional syringe pump. Small explants of cytosol (in the picolitre range) were deposited on poly-L-lysine coated glass surface under halocarbon oil. Time-lapse acquisitions typically started in late interphase or prophase. In the case of *gnu* mutant embryos, most extractions were performed when few centrosomes (between 5 and 40) were visible at the anterior-lateral cortex. Repeated use of the same extraction micropipette is not recommended. Explants from wildtype embryos initially containing a single nucleus were selected for time-lapse imaging of subsequent mitotic divisions. Explants from *gnu* mutants initially containing a single free aster near oil interface or two free asters close by were selected for time-lapse imaging of aster separation. All experiments were conducted at 25±1 °C.

Pharmacological perturbation of embryo explants

Pharmacological perturbations were performed by adding different drugs (colchicine at 0.2 mM, sodium azide at 0.5, 10, or 100 mM) diluted in cytoplasm-compatible buffer (50 mM HEPES, pH 7.8, 100 mM KCl, 1 mM $MgCl_2$). Solutions were directly administrated to the explants using a fine pipette (pulled using a Narishige PC-100 Puller with a: 2-step (69% + 55%) heating protocol and with 4 mm drop length) connected to an Eppendorf FemtoJet® 4i pump. The final drug dilution in the explants was approximately 1:10 (solution:cytosol). Buffer injections were conducted as control.

Laser ablation system

The laser ablation system was implemented by I.A.T. on the confocal spinning-disk microscope described above. A Crylas FTSS-355-Q pulsed laser emitting 355 nm, 1.1 ns pulses, 15µJ pulse energy at 1 KHz was aligned with a beam expander (16x), a scan head (SCANcube 7, Scanlab, Germany) coupled to an f-theta lens (f=56 mm, anti-reflection coating for 340–370 nm, SCANLAB AG, Germany). The focus point of the f-theta lens was aligned to be parfocal to the focal plane of the objective, using a tube lens (f=200 mm, Ø=30 mm, 355 nm AR coated, OWIS, Germany) and a dichroic mirror (T387 DCLP, Chroma) in the upper stage filter wheel. Any scattered light was blocked at the emission side with a RazorEdge LP 355 dichroic mirror OD6

@ 355nm (Chroma). The system was controlled with homemade journals for Metamorph software (Molecular Devices Inc.). The optimal laser power was set to ensure microtubule ablation while avoiding thermal expansion of cytoplasm, with post-ablation microtubule signal recovery matching known polymerization dynamics. This combination of conditions proved to be efficient at ablating target structures beyond fluorophore bleaching. In explants containing a single aster, astral microtubules were asymmetrically ablated by positioning an ellipsoid off-center (21.7 by 10.8 μm, 4 times, 15 s interval, 0.54 μm step, laser power: 25%) (Fig. 4B). In explants containing two asters, astral microtubules were ablated using an ellipsoid (21.7 by 10.8 μm, 3 times, 15 s interval, 0.54 μm step, laser power: 10–15%) roughly centered at the mid-point between the two asters, while interpolar microtubules were ablated using linear ablations (21.7 μm, 3 times, 15 s interval 0.54 μm step, laser power: 10–15%) perpendicular to the axis connecting the asters (Fig. 4F).

Simple one-dimensional model of 1- and 2-aster positioning

If we consider each aster to generate a force potential that follows its microtubule distribution, then $F(x) = F_0 e^{-\frac{x}{\lambda}}$ where x is the radial distance from the droplet edge and λ is the decay length of the microtubule distribution. For the 1-aster scenario, the steady-state solution ($F=0$) corresponds to the aster positioning at the center of the drop (assuming the drop size is small enough that there is non-zero force at the boundaries). For the 2-aster scenario, assuming asters push on each other equally to the boundary, we have $F_1(x_1) = F_0 e^{-\frac{x_1}{\lambda}} - F_0 e^{-(x_2-x_1)/\lambda}$ and $F_2(x_2) = -F_0 e^{-\frac{(L-x_2)}{\lambda}} + F_0 e^{-(x_2-x)/\lambda}$, where $0<x_1<x_2<L$ (droplet diameter). The net force is zero when $x_1=L/4$ and $x_2=3L/4$.

Two-dimensional dynamic model of aster interactions

The cytoplasm is viscous. For a viscous material, the velocity, v, of an object is dependent on the applied force F: $v \approx \gamma F$, where γ is the effective viscous drag coefficient. In our simple dynamic model implemented in Matlab® we consider $\gamma = 1$ and isolated asters with a circularly-symmetric force potential described by $f(r) = f_{slip}(r) \times \rho_{MT}(r)$, where r is the distance from the aster center (centrosome), $f_{slip} = f_0 \frac{r^2}{x_0^2+r^2}$ ($x_0 \approx 15\,\mu m$) and $\rho_{MT}(r)$ represents the distribution of microtubules from the aster. We incorporate f_{slip} to account for the reduced apparent microtubule force generation at short distances. For simplicity, we take the same characteristic distance x_0 for both aster-boundary and aster-aster interaction. To account for boundary conditions, we introduce a mirror charge outside the circle for each aster.

For single asters, we only consider interactions between the wall and aster. We take $\rho_{MT}(r) = e^{-r/\lambda}$, with $\lambda = 15\mu m$ and $f_0 = 0.007$ and r is the perpendicular aster-wall separation. We also include a 'noise' term, $\delta f = 0.0005$. So, $\vec{f}(r) = \hat{r}f_0\frac{r^2}{x_0^2+r^2}e^{-r/\lambda} + \hat{r}_{ran}\delta f$ where $\hat{r}$ is the unit vector between aster and wall, and $\hat{r}_{ran}$ is a random unit vector generated at each time iteration. For two asters, the force is given by $\vec{f}(r) = \hat{r}f_{aster-wall}(r) + \hat{x}f_{aster-aster}(x)+\hat{r}_{ran}\delta f$, where x is the distance between the two asters and $\hat{x}$ the unit vector between them. $f_{aster-wall}(r)$ is the same as for the one aster scenario. We take $f_{aster-aster}(x) = f_1\frac{x^2}{x_0^2+x^2}e^{-x/\lambda_{aster}}$, $f_1 = 0.005$ and $\lambda_{aster} = 12\mu m$.

Considering the aster-aster separation, we assumed the aster pair initially separated by 2 µm and centered within the *in silico* explant space. For the single aster case, we initialized the aster 1µm from the boundary. We always considered a system of radius 40µm. Simulations were run until the aster position reached a steady-state and angles between asters were measured at the last time point.

One dimensional dynamic model of aster interactions
We solved the above equations of motion in one-dimension using Matlab's built in stochastic PDE solver (sde), Fig. 5C and Fig. 5F. For each condition we ran 3000 simulations. For the two-aster scenario, we considered asters placed initially either side of the midpoint ($x=L/2$).

Analysis of free asters in explants – distance distributions
Distance between asters and from aster to the boundary were obtained in explants at steady state, *i.e.* where asters did not move anymore (usually 30–45 min after explant deposition). The inter-aster distance was determined as Euclidean distance in 3D. We defined the *boundary distance* (b, b_1, b_2) as the shortest distance from the aster to the interface between glass, oil and cytosol, determined manually using the FIJI measurement tools (at a precision of ±0.5 µm). To determine the explant boundary on the glass (approximated with a circle of radius R), maximum intensity projections of both fluorescence emission channels was assessed to trace the interface between the glass, oil and cytosol. For larger explants with high aspect ratio – a quasi-2D situation – the definition of *boundary distance* served as good approximation for a boundary in two dimensions. However, in small explants where the aspect ratio is not as high, two asters sometimes aligned considerably in the third dimension. In these cases, the definition for *boundary distance* led to

an underestimation of the maximum projected inter-aster distance $M = 2R - b_1 - b_2$; it becomes a geometric problem in 3D and the longest dimension is not necessarily in the plane of the glass-explant interface. This is evident for some data points in small explants (yellow dots in Fig. 5E).

Analysis of free asters in explants – dynamics

The coordinates of free asters were obtained by applying a Gaussian blur filter (radius: 1–2 pixels) and using the plugin TrackMate v3.5.1 of FIJI ImageJ (Schindelin et al., 2012; Tinevez et al., 2017). The coordinates of detected spots were imported into MATLAB® for assignment and distance calculation similarly as mentioned above. The instant relative velocity was calculated using the formula: $v_i = \frac{d_{i+1} - d_{i-1}}{t_{i+1} - t_{i-1}}$, where d is the 3D Euclidian distance and t is time in the flanking time points of the measure point. For unperturbed experiments, data was normalized to the maximum distance achieved in the separated phase to correct for scaling effect during splitting dynamics (Fig. 1E).

To analyze the lipid droplets, we performed a similar analysis using FIJI TrackMateJ (Schindelin et al., 2012; Tinevez et al., 2017). Seven extracts were analyzed with an aster present, with over 100 individual tracks of lipid droplets. RMS distance was then extracted across the entire time course of imaging. Similar analysis was performed in extracts without an aster. Curves (Fig. 2D) were fitted using the 'fit' function in MATLAB®, with r^2 = 0.98 ($b \cdot t^{\frac{1}{2}}$ for no-aster data), r^2 = 0.96 ($a_1 \cdot t^{\frac{1}{2}}$ for 1-aster data), and r^2 = 0.99 ($a_2 \cdot t^{\frac{2}{3}}$ for 1-aster data). Fitting the general function $a \cdot t^c$ gives a best fit for $c = 0.66$ for the one aster data.

Microtubule profile quantification

For single asters (Fig. 1F), we quantified the microtubule intensity using the intensity of the RFP::β-Tubulin signal. Taking the point when asters were either 5 μm or 20 μm from the explant boundary, we used FIJI to measure the microtubule intensity along a 10 μm straight line from the edge and through the aster. The line had a width of 2 μm. For each experiment, we normalized the total intensity by the maximum measured value and then binned the data in 0.2 μm bins. Hence, the recorded intensity does not reach one, and the mean intensity only reaches a maximum around 0.8 as the maximum value does not occur at the same position. Similar analysis was performed for the scenario with two asters (Fig. 3D). In this case, the centroids of the asters were used to define a straight line along which the microtubule intensity was measured throughout the process of aster separation. From this straight line between the asters, we also generated the kymograph shown in Fig. 3C right.

Analysis of free asters in explants – perturbations with drugs and UV ablation

For comparison between control and perturbation experiments, data was time-aligned to the perturbation time-point (t = 0) and plotted as average ± s.d. from at least three replicates for each condition. The change of inter-aster distance during the first 3 s after drug injection was estimated by linear regression assuming normally distributed noise, and the confidence interval of the estimated slope served as test statistic for differences between control and perturbation. Differences in final, steady-state inter-aster distance were tested by comparing the pools of distances from the last 3 s (=12 frames), using Wilcoxon signed-rank test. A significance level of 0.05 was defined prior to testing. In the case of UV ablations, the position of the aster five frames before ablation was defined as coordinate origin. The two main axes of the ellipsoid, along which the pulsed ablation was performed, defined the cartesian coordinate system. A displacement vector of the current aster position relative to the origin was calculated for each time point. The mean and standard deviation of axial (Δx) and lateral (Δy) displacement was plotted in time (Fig. 4C–D).

Acknowledgements

We thank members of the I.A.T. and T.E.S. labs for fruitful discussions, and Virgile Viasnoff, Gianluca Grenci, Tetsuya Hiraiwa, Jacques Prost and Thomas Surrey for constructive comments and feedback. We thank Sophie Theis for generating Figure 2B. We thank the staff of the Fly Facility, the Advanced Imaging Facility (AIF) and the Technical Support Service at the Instituto Gulbenkian de Ciência (IGC). Transgenic fly stocks were obtained from Bloomington Drosophila Stock Center (NIH). We acknowledge financial support from the Human Frontiers Science Program (HFSP), awarded to I.A.T. and T.E.S. and supporting J.C. and S.T. (RGY0083/2016). I.A.T acknowledges Fundação Calouste Gulbenkian (FCG), Fundação para Ciência e a Tecnologia for grant IF/00082/2013); the European Commission grant FP7-PEOPLE-2013-CIG (Nº 818743); LISBOA-01-0145-FEDER-007654 supporting IGC's core operation; LISBOA-01-7460145-FEDER-022170 (Congento) supporting the Fly Facility; PPBI-POCI-01-0145-FEDER-022122 supporting the AIF, all co-financed by FCT (Portugal) and Lisboa Regional Operational Program (Lisboa2020) under the PORTUGAL2020 Partnership Agreement (European Regional Development Fund). T.E.S. acknowledges funding from BBSRC (grant no. BB/W006944/1).

Author contributions

J.C., T.E.S. and I.A.T. conceived the study; I.A.T. and J.C. designed the experiments; J.C. carried out all experiments; T.E.S. performed the quantification of aster dynamics; S.T. and T.E.S. developed and implemented the dynamic aster repulsion model. I.A.T. designed and assembled the experimental setup; T.E.S. and I.A.T. wrote the manuscript. T.E.S. and I.A.T. jointly supervised the project.

Competing interests

The authors have no competing financial or non-financial interests.

Data availability

All original data is available upon request.

Code availability

All the codes developed in this study leading to the results from experiments or modelling are made available upon request.

References

Albertson DG. 1984. Formation of the first cleavage spindle in nematode embryos. *Dev Biol* 101:61–72. doi:10.1016/0012-1606(84)90117-9

Baker J, Theurkauf W, Schubiger G. 1993. Dynamic changes in microtubule configuration correlate with nuclear migration in the preblastoderm Drosophila embryo. *J Cell Biol* 122:113–121. doi:10.1083/jcb.122.1.113

Bieling P, Telley IA, Surrey T. 2010. A Minimal Midzone Protein Module Controls Formation and Length of Antiparallel Microtubule Overlaps. *Cell* 142:420–432. doi:10.1016/j.cell.2010.06.033

Chambers EL. 1939. The Movement of the Egg Nucleus in Relation to the Sperm Aster in the Echinoderm Egg. *J Exp Biol* 16:409–424. doi:10.1242/jeb.16.4.409

de-Carvalho J, Deshpande O, Nabais C, Telley IA. 2018. A cell-free system of Drosophila egg explants supporting native mitotic cycles. *Methods Cell Biol* 144:233–257. doi:10.1016/bs.mcb.2018.03.011

de-Carvalho J, Tlili S, Hufnagel L, Saunders TE, Telley IA. 2022. Aster repulsion drives short-ranged ordering in the Drosophila syncytial blastoderm. *Development* 149. doi:10.1242/dev.199997

Deshpande O, de-Carvalho J, Vieira DV, Telley IA. 2021. Astral microtubule cross-linking safeguards uniform nuclear distribution in the Drosophila syncytium. *J Cell Biol* 221:e202007209. doi:10.1083/jcb.202007209

Deshpande O, Telley IA. 2021. Nuclear positioning during development: Pushing, pulling and flowing. *Semin Cell Dev Biol* 120:in press. doi:10.1016/j.semcdb.2021.09.020

Donoughe S, Extavour CG. 2016. Embryonic development of the cricket Gryllus bimaculatus. *Dev Biol* 411:140–156. doi:10.1016/j.ydbio.2015.04.009

Field CM, Groen AC, Nguyen PA, Mitchison TJ. 2015. Spindle-to-cortex communication in cleaving, polyspermic Xenopus eggs. *Mol Biol Cell* 26:3628–3640. doi:10.1091/mbc.e15-04-0233

Firestone AJ, Weinger JS, Maldonado M, Barlan K, Langston LD, O'Donnell M, Gelfand VI, Kapoor TM, Chen JK. 2012. Small-molecule inhibitors of the AAA+ ATPase motor cytoplasmic dynein. *Nature* 484:125–129. doi:10.1038/nature10936

Foe VE, Alberts BM. 1983. Studies of nuclear and cytoplasmic behaviour during the five mitotic cycles that precede gastrulation in Drosophila embryogenesis. *J Cell Sci* 61:31–70.

Foe VE, Field CM, Odell GM. 2000. Microtubules and mitotic cycle phase modulate spatiotemporal distributions of F-actin and myosin II in Drosophila syncytial blastoderm embryos. *Development* 127:1767–1787. doi:10.1242/dev.127.9.1767

Forth S, Hsia K-C, Shimamoto Y, Kapoor TM. 2014. Asymmetric Friction of Nonmotor MAPs Can Lead to Their Directional Motion in Active Microtubule Networks. *Cell* 157:420–432. doi:10.1016/j.cell.2014.02.018

Freeman M, Glover DM. 1987. The gnu mutation of Drosophila causes inappropriate DNA synthesis in unfertilized and fertilized eggs. *Gene Dev* 1:924–930. doi:10.1101/gad.1.9.924

Freeman M, Nüsslein-Volhard C, Glover DM. 1986. The dissociation of nuclear and centrosomal division in gnu, a mutation causing giant nuclei in Drosophila. *Cell* 46:457–468. doi:10.1016/0092-8674(86)90666-5

Garzon-Coral C, Fantana HA, Howard J. 2016. A force-generating machinery maintains the spindle at the cell center during mitosis. *Sci New York N Y* 352:1124–7. doi:10.1126/science.aad9745

Grill SW, Howard J, Schäffer E, Stelzer EH, Hyman AA. 2003. The Distribution of Active Force Generators Controls Mitotic Spindle Position. *Science* 301:518–521. doi:10.1126/science.1086560

Grill SW, Hyman AA. 2005. Spindle Positioning by Cortical Pulling Forces. *Dev Cell* 8:461–465. doi:10.1016/j.devcel.2005.03.014

Hamaguchi M, Hiramoto Y. 1980. Fertilization process in the heart-urchin, Clypeaster Japonicus observed with a differential interference microscope. *Dev Growth Differ* 22:517–530. doi:10.1111/j.1440-169x.1980.00517.x

Hamaguchi MS, Hiramoto Y. 1986. Analysis of the Role of Astral Rays in Pronuclear Migration in Sand Dollar Eggs by the Colcemid-UV Method. *Dev Growth Differ* 28:143–156. doi:10.1111/j.1440-169x.1986.00143.x

Hinchcliffe EH, Sluder G. 2001. "It Takes Two to Tango": understanding how centrosome duplication is regulated throughout the cell cycle. *Gene Dev* 15:1167–1181. doi:10.1101/gad.894001

Hirano K, Ishikawa M. 1979. Induction of aster formation and cleavage in eggs of the sea urchin HEMICENTROTUS PULCHERRIMUS by injection of sperm components. *Dev Growth Differ* 21:473–481. doi:10.1111/j.1440-169x.1979.00473.x

Hoffmann I. 2021. Centrosomes in mitotic spindle assembly and orientation. *Curr Opin Struc Biol* 66:193–198. doi:10.1016/j.sbi.2020.11.003

Holy TE, Dogterom M, Yurke B, Leibler S. 1997. Assembly and positioning of microtubule asters in microfabricated chambers. *Proc National Acad Sci* 94:6228–6231. doi:10.1073/pnas.94.12.6228

Howard J. 2001. Mechanics of motor proteins and cytoskeleton. Sinauer Associates.

Inoue YH, Savoian MS, Suzuki T, Máthé E, Yamamoto M-T, Glover DM. 2004. Mutations in orbit/mast reveal that the central spindle is comprised of two microtubule populations, those that initiate cleavage and those that propagate furrow ingression. *J Cell Biology* 166:49–60. doi:10.1083/jcb.200402052

Ishihara K, Korolev KS, Mitchison TJ. 2016. Physical basis of large microtubule aster growth. *Elife* 5:e19145. doi:10.7554/elife.19145

Jeune-Smith Y, Hess H. 2010. Engineering the length distribution of microtubules polymerized in vitro. *Soft Matter* 6:1778–1784. doi:10.1039/b919488f

Kanesaki T, Edwards CM, Schwarz US, Grosshans J. 2011. Dynamic ordering of nuclei in syncytial embryos: a quantitative analysis of the role of cytoskeletal networks. *Integr Biol* 3:1112–1119. doi:10.1039/c1ib00059d

Keren K. 2016. Non Equilibrium Steady State Dynamics of Contractile Actin Networks. *Biophys J* 110:174a. doi:10.1016/j.bpj.2015.11.972

Kimura K, Kimura A. 2011. Intracellular organelles mediate cytoplasmic pulling force for centrosome centration in the Caenorhabditis elegans early embryo. *Proc National Acad Sci* 108:137–142. doi:10.1073/pnas.1013275108

Laan L, Pavin N, Husson J, Romet-Lemonne G, van Duijn M, López MP, Vale RD, Jülicher F, Reck-Peterson SL, Dogterom M. 2012. Cortical Dynein Controls Microtubule Dynamics to Generate Pulling Forces that Position Microtubule Asters. *Cell* 148:502–514. doi:10.1016/j.cell.2012.01.007

Lee LA, Hoewyk DV, Orr-Weaver TL. 2003. The Drosophila cell cycle kinase PAN GU forms an active complex with PLUTONIUM and GNU to regulate embryonic divisions. *Gene Dev* 17:2979–2991. doi:10.1101/gad.1132603

Lv Z, Rosenbaum J, Aspelmeier T, Großhans J. 2018. A 'molecular guillotine' reveals the interphase function of Kinesin-5. *J Cell Sci* 131:jcs210583. doi:10.1242/jcs.210583

Maliga Z, Kapoor TM, Mitchison TJ. 2002. Evidence that Monastrol Is an Allosteric Inhibitor of the Mitotic Kinesin Eg5. *Chem Biol* 9:989–996. doi:10.1016/s1074-5521(02)00212-0

Meaders JL, Matos SN de, Burgess DR. 2020. A Pushing Mechanism for Microtubule Aster Positioning in a Large Cell Type. *Cell Reports* 33:108213. doi:10.1016/j.celrep.2020.108213

Megraw T, Li K, Kao L, Kaufman T. 1999. The centrosomin protein is required for centrosome assembly and function during cleavage in Drosophila. *Development* 126:2829–39.

Minc N, Burgess D, Chang F. 2011. Influence of Cell Geometry on Division-Plane Positioning. *Cell* 144:414–426. doi:10.1016/j.cell.2011.01.016

Mitchison T, Wühr M, Nguyen P, Ishihara K, Groen A, Field CM. 2012. Growth, interaction, and positioning of microtubule asters in extremely large vertebrate embryo cells. *Cytoskeleton* 69:738–750. doi:10.1002/cm.21050

Pelletier JF, Field CM, Fuerthauer S, Sonnett M, Mitchison TJ. 2020. Co-movement of astral microtubules, organelles and F-actin by dynein and actomyosin forces in frog egg cytoplasm. *Elife* 9:e60047. doi:10.7554/elife.60047

Postner MA, Miller KG, Wieschaus EF. 1992. Maternal effect mutations of the sponge locus affect actin cytoskeletal rearrangements in Drosophila melanogaster embryos. *J Cell Biol* 119:1205–1218. doi:10.1083/jcb.119.5.1205

Rappaport R. 1969. Aster-equatorial surface relations and furrow establishment. *J Exp Zool* 171:59–67. doi:10.1002/jez.1401710109

Rappaport R. 1961. Experiments concerning the cleavage stimulus in sand dollar eggs. *J Exp Zool* 148:81–89. doi:10.1002/jez.1401480107

Riparbelli MG, Callaini G, Glover DM. 2000. Failure of pronuclear migration and repeated divisions of polar body nuclei associated with MTOC defects in polo eggs of Drosophila. *J Cell Sci* 113 (Pt 18):3341–50.

Rogers SL, Rogers GC, Sharp DJ, Vale RD. 2002. Drosophila EB1 is important for proper assembly, dynamics, and positioning of the mitotic spindle. *J Cell Biology* 158:873–884. doi:10.1083/jcb.200202032

Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S, Rueden C, Saalfeld S, Schmid B, Tinevez J-Y, White DJ, Hartenstein V, Eliceiri K, Tomancak P, Cardona A. 2012. Fiji: an open-source platform for biological-image analysis. *Nat Methods* 9:676–682. doi:10.1038/nmeth.2019

Schubiger G, Edgar B. 1994. Using inhibitors to study embryogenesis. *Methods Cell Biol* 44:697–713.

Shimamoto Y, Forth S, Kapoor TM. 2015. Measuring Pushing and Braking Forces Generated by Ensembles of Kinesin-5 Crosslinking Two Microtubules. *Dev Cell* 34:669–681. doi:10.1016/j.devcel.2015.08.017

Simone AD, Spahr A, Busso C, Gönczy P. 2018. Uncovering the balance of forces driving microtubule aster migration in C. elegans zygotes. *Nat Commun* 9:938. doi:10.1038/s41467-018-03118-x

Sommer R, Tautz D. 1991. Asynchronous mitotic domains during blastoderm formation inMusca domestica L. (Diptera). *Roux's Archives Dev Biology* 199:373–376. doi:10.1007/bf01705931

Subramanian R, Ti S-C, Tan L, Darst SA, Kapoor TM. 2013. Marking and Measuring Single Microtubules by PRC1 and Kinesin-4. *Cell* 154:377–390. doi:10.1016/j.cell.2013.06.021

Sulerud T, Sami AB, Li G, Kloxin A, Oakey J, Gatlin J. 2020. Microtubule-dependent pushing forces contribute to long-distance aster movement and centration in Xenopus laevis egg extracts. *Mol Biol Cell* 31:2791–2802. doi:10.1091/mbc.e20-01-0088

Tanimoto H, Kimura A, Minc N. 2016. Shape–motion relationships of centering microtubule asters. *J Cell Biology* 212:777–787. doi:10.1083/jcb.201510064

Tanimoto H, Sallé J, Dodin L, Minc N. 2018. Physical forces determining the persistency and centring precision of microtubule asters. *Nat Phys* 14:848–854. doi:10.1038/s41567-018-0154-4

Telley IA, Gáspár I, Ephrussi A, Surrey T. 2013. A single Drosophila embryo extract for the study of mitosis ex vivo. *Nat Protoc* 8:310–324. doi:10.1038/nprot.2013.003

Telley IA, Gáspár I, Ephrussi A, Surrey T. 2012. Aster migration determines the length scale of nuclear separation in the Drosophila syncytial embryo. *J Cell Biol* 197:887–895. doi:10.1083/jcb.201204019

Tinevez J-Y, Perry N, Schindelin J, Hoopes GM, Reynolds GD, Laplantine E, Bednarek SY, Shorte SL, Eliceiri KW. 2017. TrackMate: An open and extensible platform for single-particle tracking. *Methods* 115:80–90. doi:10.1016/j.ymeth.2016.09.016

Wijeratne S, Subramanian R. 2018. Geometry of antiparallel microtubule bundles regulates relative sliding and stalling by PRC1 and Kif4A. *Elife* 7:e32595. doi:10.7554/elife.32595

Wilson EB. 1986. The Cell in Development and Inheritance, Columbia University biological series: Columbia University. Macmillan.

Winkler F, Gummalla M, Künneke L, Lv Z, Zippelius A, Aspelmeier T, Grosshans J. 2015. Fluctuation Analysis of Centrosomes Reveals a Cortical Function of Kinesin-1. *Biophys J* 109:856–868. doi:10.1016/j.bpj.2015.07.044

Wühr M, Tan ES, Parker SK, Detrich HW, Mitchison TJ. 2010. A Model for Cleavage Plane Determination in Early Amphibian and Fish Embryos. *Curr Biol* 20:2040–2045. doi:10.1016/j.cub.2010.10.024

Xie J, Najafi J, Borgne RL, Verbavatz J-M, Durieu C, Sallé J, Minc N. 2022. Contribution of cytoplasm viscoelastic properties to mitotic spindle positioning. *Proc National Acad Sci* 119:e2115593119. doi:10.1073/pnas.2115593119

Zhu J, Burakov A, Rodionov V, Mogilner A. 2010. Finding the Cell Center by a Balance of Dynein and Myosin Pulling and Microtubule Pushing: A Computational Study. *Mol Biol Cell* 21:4418–4427. doi:10.1091/mbc.e10-07-0627

Figure 1 – Embryo explant assay enables the kinematic study of individual asters. (A)
Maximum intensity Z-projection of a *gnu* mutant embryo expressing RFP::β-Tubulin (magenta)
and Spd2::GFP (green). Scale bar, 50 μm. **(B)** Schematic of cytosol extraction from a *gnu* mutant
Drosophila embryo and *ex vivo* explant formation (de-Carvalho et al., 2018). **(C)** Maximum
intensity Z-projections of a single aster (arrowhead) moving away from the boundary of an

explant (yellow dashed circle) produced from *gnu* mutant embryos expressing RFP::β-Tubulin (magenta) and Spd2::GFP (green). Yellow dashed circles represent the explant boundary. Scale bar, 20 μm. **(D)** Migration velocity as a function of time, where t=0 is defined as the time when the aster lies midway between the explant edge and the final position of the aster. Solid line represents average over all measurements (n=7). **(E)** Average migration velocity of single asters away from the explant boundary (n=7). Distance normalized by the final, steady-state distance for each aster. **(F)** Normalized intensity of astral microtubules as schematically outlines in the inset. The black line is a mono-exponential fit to the data excluding the first two data points (red), representing the centrosome, and the dashed lines mark ±1 s.d. The decay length is 11.8 ± 0.5 μm (mean ± s.e.m.), and the intensity drops to background level at ~40 μm. Inset: Single Z-plane image of an explant from a *gnu* mutant embryo expressing RFP::β-Tubulin (magenta) and Spd2::GFP (green), containing a single aster. The dashed white line and the circular arrow represent the radial maximum intensity projection of the microtubule signal from the centrosome towards the periphery aiming at measuring aster size. The yellow dashed circle represents the explant boundary. Scale bar, 10 μm. **(G)** Maximum intensity Z-projection of a 3D image stack of a small explant containing one aster that exemplifies microtubule buckling and splay near the explant boundary (yellow dashed circle). Scale bar, 5 μm.

Figure 2 – Yolk particle tracking suggests particle displacement by repulsion. (A) Velocity field of yolk droplets around the single aster as it escapes from the explant boundary. Image rotation and frame matching were performed to overlay all experiments such that the aster, at each time point is positioned at (0,0) and moves in the direction (0,-1). Track color coding denotes displacement angle, where turquoise corresponds to 0° and red to 90°. **(B)** Averaged droplet velocity from seven experiments relative to the aster position. The arrow length denotes the average droplet speed and color the orientation of movement relative to the aster (color bar). The blue arrow highlights the effective direction of aster motion (the velocity calculations were done in the aster frame of reference). **(C)** Velocity profile of droplets with (black, n=7 experiments) and without an aster present (magenta, n= 6 experiments). We only consider droplets within a distance of 20 μm from the aster, where t=0 is the time point at which the aster starts moving away from the explant boundary (n>30 tracks for each condition). **(D)** Root-mean-square (RMS) displacement of lipid droplets in the explants. Droplet movement analyzed with (black, n= 7 experiments) and without (magenta, n=6 experiments) an aster present. Error bars s.e.m. and lines represent fits to models shown in legend.

Figure 3 – Aster-aster separation in explants depends on microtubule distribution and interactions. (A) Maximum intensity Z-projections of two aster separating an explant produced from *gnu* mutant embryos expressing RFP::β-Tubulin (magenta) and Spd2::GFP (green). Yellow dashed circles represent the explant boundary. Scale bar, 20 μm. **(B)** Aster separation velocity as a function of normalized separation distance (n=9). For each experiment, distance is normalized by the final, steady-state separation distance. **(C)** Left: Colourmap of normalized microtubule density between two separating asters. Right: Kymograph of microtubule intensity between the asters during separation. Scale bars, 2 min (horizontal) 5 μm (vertical). **(D)** Normalized microtubule intensity at the midpoint perpendicular axis between asters in function of aster separation distance. Open markers denote average values and error bars the standard deviation. Solid line represents the fitting to exponential decay (n=7), fitting performed using MATLAB *fit* function. **(E)** Fitting to average separation velocity (circles) considering

microtubule intensity and a microtubule slipping term (inefficient repulsion). Microtubule density was either fitted beforehand (solid line in **D**) or directly included (dashed line).

Figure 4 – Aster positioning and separation is determined by a dominant microtubule-dependent pushing force. (A) Aster separation dynamics upon injection of buffer (control, n=3), 0.5 mM (n=3), 10 mM (n=4), 100mM (n=3) sodium azide, or 0.2 mM (n=3) colchicine.

The * symbol denotes significance at p<0.05. Gray or colored areas around average curves denote ± 1 s.d. **(B)** Schematic of single aster eccentric circular UV laser ablation (magenta dashed line); this ablation aims at shortening astral microtubules on the left side of the aster. t=0 min denotes ablation time. **(C–D)** Aster displacement before and after eccentric circular ablation in explants unperturbed (**C**, n=8) or treated with colchicine (**D**, n=8). Arrows represent average displacement magnitude and direction, and vertical and horizontal grey bars denote ±1 s.d. of displacement in x and y, respectively. **(E)** Explants containing two asters were perturbed by (1) ellipse ablation around both asters during separation ("peripheral ablation"); (2) linear ablation between two asters ("central ablation"). **(F)** Change of inter-aster distance upon laser ablation (time = 0) as described in **E**. Upon peripheral ablation, separating asters maintained their movement and sometimes slightly accelerated, while central ablation caused movement towards each other.

A

Aster

r

Aster-wall Potential

MT Slippage
Net potential
MT potential

r (μm)

Aster   Aster
r₁   r₁₂   r₂

Aster-Aster Potential

Net potential

r₁₂

B Simulation of 1-aster positioning

migration velocity [μm/s]

normalised aster-boundary distance

C Data & model prediction:
1-aster final positioning

shortest boundary distance **b** [μm]

Y = R
Y = ½ R
Y = 0.74 R

radius **R** [μm]

D Simulation of 2-aster separation

separation velocity [μm/s]

normalised inter-aster distance

E Experiment: 2 aster separation

inter-aster distance d [μm]

maximum projected separation
2R−b₁−b₂ [μm]

F Model: 2-aster separation

inter-aster distance **dₐ** [μm]

Y = 2**R**
Y = **R**
Y = ½ **R**

R [μm]

G Dynamic model simulation
of three asters

probability

with potential
without potential

angle [°]

H Dynamic model simulation
of four asters

probability

angle [°]

Figure 5 – Model. (A) Schematic of 2D model of aster dynamics. Bottom graphs represent cartoons of the potentials between aster-boundary and aster-aster. **(B)** Fit of 2D model to the single aster velocity profile shown in Fig. 1E. The average model fit (black curve) is calculated from 10 simulation runs (gray lines). Error bars are experimental error (n=7). **(C)** 1D stochastic model of single aster dynamics (red line = mean, blue shaded region $\pm$1s.d.), compared to experimentally observed distribution of aster position (black dots). **(D)** As in (B) but for the two-aster scenario. **(E)** Scatter plot of inter-aster distance (d_a, see inset) as a function of the maximum projected separation, calculated from the explant diameter ($2R$) and the boundary distance of each of the two asters, b_1 and b_2 (n=54). The blue dashed line represents the estimated upper limit of the interaction distance between two asters (~45 μm). Red dots represent cases where the two asters were likely positioned far apart during explant generation. The yellow dots are cases of small explants where projection of the 3D volume leads to overestimation of b_1 and/or b_2. **(F)** 1D stochastic model of two-aster dynamics where black circles denote final aster positions from simulation. **(G)** Angle distribution from aster positions in a dynamic model simulation with three asters. The simulation evolved from initially random positions, and asters robustly moved towards a triangular configuration. The peak at 60º represents equal distances between the three asters. In the absence of a repulsion potential the regularity is lost (blue line). **(H)** Angle distribution from aster positions in a dynamic model simulation with four asters. The two insets show the temporal evolution of position and the final configuration marked with dashed lines. The majority of simulations (17/20) resulted in a regular square (top left inset) with 3/20 resulting in a "Y" configuration (top right inset).

Supplementary Fig. 1 – (A) Scatter plot of shortest distance to explant boundary (b_1 and b_2) as a function of the radius R in explants containing two asters (n=54). The magenta line represents the linear regression. Black dashed lines represent half and full radius distance which are the geometric constraints in the system. **(B)** Trajectories of aster distance to the explant boundary from independent experiments. **(C)** Average microtubule density (black line, inferred from RFP::β-Tubulin signal) along shortest distance to explant boundary, normalised by the maximum intensity within each experiment. Grey traces are individual experiments (n=7). **(D)** Scattered plot of inter-aster distance (d_a) as a function of the radius (R) of explants containing two asters (n=54). Most measured data points fall between the dashed lines denoting the explant radius ($Y = R$) and half of the radius ($Y = \frac{1}{2}R$). The magenta line represents the linear regression.

Video Legends

Video 1: Maximum intensity Z-projection from a 3D time-lapse movie of explants generated from *gnu* mutant embryos expressing RFP::β-Tubulin (magenta) and Spd2::GFP (green). The left explant contains a single aster moving away from the explant boundary, the right explant contains two separating asters. Time in min:sec, scale bar 10 μm. Frame rate is 4 frames/min.

Video 2: Maximum intensity Z-projection from a 3D time-lapse movie of explants generated from a *gnu* mutant embryo expressing RFP::β-Tubulin (magenta) and Spd2::GFP (green), containing two separating asters, after pulse injection of solutions: control with buffer (left), 10 mM sodium azide (middle) and 0.2 mM of colchicine (right). Time in min:sec, scale bar 10 μm. Frame rate is 4 frames/min.

Video 3: Maximum intensity Z-projection from a 3D time-lapse movie of explants generated from a *gnu* mutant embryo expressing RFP::β-Tubulin (magenta) and Spd2::GFP (green) containing a single aster. The aster was allowed to equilibrate followed by an asymmetric elliptic ablation (yellow line at times 00:15 to 01:00) performed in control explants (no injection) and in explants supplemented with 0.2mM of colchicine. Time in min:sec, scale bar 10 μm. Frame rate is 4 frames/min.

Video 4: Maximum intensity Z-projection from a 3D time-lapse movie of an explant containing two separating asters from a *gnu* mutant embryo expressing RFP::β-Tubulin (magenta) and Spd2::GFP (green). The elliptic ablation (yellow line from 00:15 to 00:45) was performed when asters were ~7 μm apart. Time in min:sec, scale bar 10 μm. Frame rate is 4 frames/min.