Considerations on brain age predictions from repeatedly sampled data across time

Max Korbmacher[1,2,3], Meng-Yun Wang[3,4], Rune Eikeland[3,4], Ralph Buchert[5], Ole A. Andreassen[2,6], Thomas Espeseth[7], Esten Leonardsen[2,7], Lars T. Westlye[2,7], Ivan I. Maximov[1,2], Karsten Specht[3,4,8]

Department of Health and Functioning, Western Norway University of Applied Sciences, Bergen, Norway

Norwegian Centre for Mental Disorders Research (NORMENT), Oslo University Hospital & Institute of Clinical Medicine, University of Oslo, Oslo, Norway

Mohn Medical Imaging and Visualisation Center (MMIV), Bergen, Norway

Department of Biological and Medical Psychology, University of Bergen, Bergen, Norway

Department of Diagnostic and Interventional Radiology and Nuclear Medicine, University Medical Center Hamburg-Eppendorf, Martinistr. 52, 20246, Hamburg, Germany

KG Jebsen Centre for Neurodevelopmental Disorders, University of Oslo, Oslo, Norway

Department of Psychology, University of Oslo, Oslo, Norway

Department of Education, UiT The Arctic University of Norway, Tromsø, Norway

Running title

Associations of scan quality and field strength with longitudinal brain age

Keywords

Brain age, densly sampled MRI, magnetic resonance imaging, T1-weighted, scan quality, field strength

Acknowledgements

This study was financed by the Research Council of Norway (#276044 and #223273); South-Eastern Norway Regional Health Authority (#2022080); and the European Union's Horizon2020 Research and Innovation Programme (CoMorMent project; Grant #847776).

Author contributions

Max Korbmacher: Study design, Software, Formal analysis, Visualizations, Project administration, Writing—original draft, Writing—review & editing. Meng-Yun Wang: Writing—review & editing. Rune Eikeland: Writing—review & editing. Ralph Buchert: Writing—review & editing. Ole A. Andreassen: Writing—review & editing, Funding acquisition. Thomas Espeseth: Writing—review & editing, Funding acquisition. Esten Leonardsen: Writing—review & editing. Lars T. Westlye: Writing—review & editing, Funding acquisition. Ivan I. Maximov: Writing—review & editing. Karsten Specht: Writing—review & editing, Funding acquisition.

Conflicts of interest

OOA has received a speaker's honorarium from Lundbeck and is a cosultant to Coretechs.ai.

Code and data availability

Data processing pipeline and weights for the trained convolutional neural network can be found at https://github.com/estenhl/pyment-public. Processed tabular data and analysis code are made available at https://github.com/MaxKorbmacher/BBSC_Brain Age.

Abstract

Introduction. Brain age, the estimation of a person's age from magnetic resonance imaging (MRI)

parameters, has been used as a general indicator of health. The marker requires however further

validation for application in clinical contexts. Here, we show how brain age predictions perform for

for the same individual at various time points and validate our findings with age-matched healthy

controls.

Methods. We used densely sampled T1-weighted MRI data from four individuals (from two

datasets) to observe how brain age corresponds to age and is influenced by acquision and quality

parameters. For validation, we used two cross-sectional datasets. Brain age was predicted by a pre-

trained deep learning model.

Results. We find small within-subject correlations between age and brain age. We also find

evidence for the influence of field strength on brain age which replicated in the cross-sectional

validation data, and inconclusive effects of scan quality.

Conclusion. The absence of maturation effects for the age range in the presented sample, brain age

model-bias (including training age distribution and field strength) and model error are potential

reasons for small relationships between age and brain age in longitudinal data. Future brain age

models should account for differences in field strength and intra-individual differences.

Background: What is brain age and what is it good for?

Brain age refers to the estimation of a person's age from magnetic resonance imaging (MRI)

parameters (Franke & Gaser, 2019). This has been done using either neural networks on 3D data

(Leonardsen et al., 2022) or tabular data containing region-averaged metrics (Korbmacher et al.,

2023; Vidal-Pineiro et al., 2021). Brain age becomes particularly interesting when assuming that

lifespan changes in the brain follow normative patterns and that deviations from such patterns might

be indicative of disease or disease development (Marquand et al., 2019; Kaufmann et al., 2019). An

elevated predicted compared to chronological age in adults may be indicative of psychiatric,

neurodegenerative, and neurological disorders (Kaufmann et al., 2019) and poorer health, for

example measured by various cardiometabolic risk factors (Beck et al., 2022; Korbmacher et al.,

2022). Hence, brain age is a promising developing biomarker of general brain health (Franke &

Gaser, 2019).

However, revealing connections between brain age and structural and functional brain architecture

is needed to fully understand the biological underpinnings of brain age and its potential clinical

implications (Vidal-Pineiro et al., 2021). Furthermore, large cross-sectional samples are often used,

which could obscure effects of predictive power of brain age by confounders, in particular,

differences in MRI acquisition (Jirsaraie et al., 2022). Hence, contributions of individual differences

to brain age estimates require a closer examination. With the aim of assessing the effects of

automated MRI scan quality control (QC) metrics on brain age predictions, we used a pre-trained

deep neural network model (Leonardsen et al., 2022) to predict brain ages from densely sampled T1-

weighted MRI data from three individuals (BBSC1-3) scanned in total N_{BBSC} = 103 times over a

one-year interval (Wang et al., 2022), and an independent data set including one individual (FTHP1)

scanned N_{FTHP} = 557 times over a three-year interval. We first observed within-subject prediction

error and correlations between chronological and predicted age, revealing small, non-significant

correlations and larger prediction errors than previously shown in between-subjects analyses. We

then tested associations of QC metrics on brain age using linear random intercept models showing

potential associations between QC parameters and brain age as well as associations between

acquisition parameters and brain age. We validate the findings in cross sectional data and

investigate differences in the variability in predictions between longitudinal and cross-sectional

datasets.

Results and Discussion

Weak correlation between brain age and age

Crude within-subject correlations between age and brain age revealed differing directionalities of

slopes across subjects, with only the FTHP1 correlation being statistically significant (r = 0.38, 95%

CI [0.24; 0.51], p < .001; **Figure 1**).

Figure 1. Intra-individual correlations between brain age and chronological age at 3T for BBSC1-3

and FTHP1

Dot colour was grey, with overlapping dots presented as darker.

This is likely due to the small age range and short inter-scan-intervals, as illustrated by differences

in model-innate error for the different subjects (**Table 1**) compared to error statistics across age

groups (MAE_{test} = 2.47, $MAE_{external}$ = 3.90, as presented in Leonardsen et al., 2022).

Table 1. Age, predicted age, brain age gap (BAG), and prediction error by subject and field strength

Subject	Field Strength	N Observations	Mean Age	SD Age	Mean Prediction	SD Prediction	Mean BAG	SD BAG	MAE	RMSE
BBSC1	3T	38	30.66	0.38	28.13	1.25	-2.52	1.20	2.55	2.79
BBSC2	3T	40	28.09	0.38	24.37	0.95	-3.72	1.03	3.72	3.85
BBSC3	3T	25	40.66	0.28	30.87	1.37	-9.79	1.46	9.79	9.89
FTHP1	3T	153	49.86	0.54	45.71	3.70	-4.15	3.52	4.31	5.44
FTHP1	1.5T	394	49.64	0.46	48.39	2.52	-1.25	2.54	2.15	2.83

The presented data refer to the longitudinal, densely sampled data of few individuals.

BAG = brain age gap, MAE = mean absolute error, RMSE = root mean squared error. BAG is calculated as the

difference between predicted age and age.

Additionally, the ages of BBSC1-3 fall into some of the least represented parts of the training data

age distribution in the underlying model for the brain age predictions (see Leonardsen et al., 2022)

which might contribute to explaining some of the prediction differences beyond model error and

intra-individual age range across scanning sessions.

Yet, when using age-matched healthy controls from the cross-sectional TOP and NCNG samples

using BBSC and FTHP longitudinal participants' mean ages ± five years (presented in **Table 1**),

correlations between age and brain age estimates were significant for age matches (**Table 2**).

Table 2. Correlations between age-matching cross-sectional sub-samples' ages and brain age

estimates

Matched subject	Field Strength	$N_{subjects}$	Pearson's r [95% CI]*	Mean Age	SD Age	Mean Prediction	SD Prediction	Mean BAG	SD BAG	MAE	RMSE
BBSC1	3T	279	0.56 [0.47, 0.64]	30.64	2.74	28.34	4.10	-2.30	3.42	3.33	4.12
BBSC2	3T	269	0.62 [0.54, 0.69]	28.81	2.83	26.75	3.96	-2.05	3.13	3.02	3.74
BBSC3	3T	248	0.44 [0.34, 0.54]	40.71	2.95	37.86	5.21	-2.85	4.71	4.52	5.50
FTHP1	3T	113	0.71 [0.60, 0.79]	48.60	3.04	44.68	5.93	-3.91	4.34	4.59	5.84
FTHP1	1.5T	49	0.79 [0.65, 0.88]	49.61	3.22	51.98	4.40	2.38	2.71	2.91	3.58

Matched subject refers to the longitudinally sample subjects presented in **Table 1**. Mean ages for the respective subjects

with an interval of five years were used to sample from the cross-sectional validation set consisting of 1.5T and 3T data

from TOP and NCNG samples. BAG = brain age gap, MAE = mean absolute error, RMSE = root mean squared error.

BAG is calculated as the difference between predicted age and age. *All $p < .001$.

Interestingly, we also find systematically underestimated brain ages across subjects (**Figure 1**) with

the underestimations being stronger for a field strength of 3T than 1.5T for FTHP1 (**Table 1**), and

in age-matched cross-sectional data (**Table 2**). While longitudinal brain age predictions were closer

related with age at 3T MRI ($r_{partial}$ = 0.38, 95% CI [0.24, 0.51], $p < .001$) than at 1.5T MRI ($r_{partial}$ =

0.06, 95% CI [-0.04, 0.16], $p = .239$; **Figure 2**), the prediction error was smaller at 1.5T (**Table 1**),

with these findings being robust to exclusions of back-to-back repeat scans acquired in the same

session without repositioning of the head (**Supplement 1**). When using the out-of-sample test sets

TOP and NCNG cross-sectional data, we find highly corresponding relationships between age and

brain age at 1.5T ($r = 0.98$, 95% CI [0.97, 0.98], $p < .001$) and 3T ($r = 0.92$, 95% CI [0.91, 0.93], p

$< .001$), but higher prediction error at 3T for age matched subjects (**Table 2**) and overall (**Table 3**).

Table 3. Age, predicted age, brain age gap (BAG), and prediction error by cross-sectional data set

and field strength

Dataset	Field Strength	N$_{subjects}$	Mean Age	SD Age	Mean Prediction	SD Prediction	Mean BAG	SD BAG	MAE	RMSE
TOP GE750	3T	543	34.15	11.54	31.37	11.11	-2.78	4.10	3.74	4.96
TOP HDxt	3T	313	30.81	8.15	29.65	8.76	-1.16	3.78	3.05	3.95
TOP all	3T	856	32.93	10.55	30.74	10.34	-2.19	4.06	3.49	4.61
NCNG	1.5T	209	54.66	14.50	56.02	14.50	1.36	3.28	2.83	3.55

The presented data refer to the cross-sectional data used as a comparison to the longitudinal data presented in **Table 1**.

TOP 3T data were obtained at two scanners: GE750 and HDtx. BAG = brain age gap, MAE = mean absolute error,

RMSE = root mean squared error. BAG is calculated as the difference between predicted age and age.

Figure 2. Intra-individual correlations between brain age and chronological age at 1.5T and 3T for

FTHP1

Dot colour was grey, with overlapping dots presented darker.

This emphasises the importance of treating predictions for age groups which are underrepresented

in the training sample and differences in field strength with care. In that sense, the observed within-

subjects variability associated with acquisition- or scanner-specific effects might be used to estimate

the minimum size of true within-subject changes (e.g., due to disease) to be detected with a given

power. Previous findings outlined the influence of scanner site on brain age predictions and scan

quality (Jirsaraie et al., 2022; Leonardsen et al., 2022) indicated by the Euler number (Rosen et al.,

2018). Lower quality scans lead to lower prediction errors. We hence hypothesise that there might

be additional reasons for inaccuracies in brain age predictions caused by factors beyond the

characteristics of the brain age model, in particular scan quality and acquisition parameters.

Scan quality and acquisition: possible reasons for inaccurate brain age predictions?

We used linear random intercept models at the participant level to examine associations of

individual quality control (QC) metrics (see **Figure 3**; **Materials and methods**) and brain age,

while controlling for age in BBSC1-3. Entropy-focus criterion (EFC, β_{std} = -0.489, p_{Holm} < .001) and

the foreground-background energy ratio (FBER, β_{std} = 0.456, p_{Holm} < .001) were significant

predictors of brain age. In a seperate analysis of FTHP1 (scanned at different sites using different

scanning parameters) we included scanner site, field strength, and slice thickness as random factors,

rendering none of the QC metrics significant after correcting for multiple testing (p_{Holm} = 1).

Figure 3. Standardized quality control metrics at 3T per subject

For an overview of scan quality control metrics at 1.5T (only applicable for FTHP1) see **Supplement 2**.

Follow-up analyses in FTHP1 focussed on examining acquisition parameters. We observed

individual fixed effects of field strength, manufacturer and slice thickness in one model each, while

keeping scanner site and the other acquisition parameters as random effects at the level of the

intercept, revealing only significant associations of field strength (β = -1.141, $p_{\text{Holm}} < .001$) with

brain age.

For validation, we replicate this finding in healthy controls from the TOP and NCNG (see

Materials and Methods section). BAG was predicted by field strength (β = -2.518, $p < .001$) when

controlling for scanner site, with $\text{Mean}_{\text{BAG-1.5T}}$ = 1.357±3.285 and $\text{Mean}_{\text{BAG-3T}}$ = -2.19±4.06 using

the entire out-of-sample test data. When age-matching FTHP1 and including only the $N = 162$

participants aged 50±5 years ($N = 49$ scanned at 1.5T), the effect of field strength appears stronger

(β = -7.40, $p < .001$), with $\text{Mean}_{\text{BAG-1.5T}}$ = 2.38±2.71 and $\text{Mean}_{\text{BAG-3T}}$ = -3.92±4.35. In that case, also

correlations between age and brain age are stronger at 1.5T compared to 3T (**Table 2**). This was

also true when using the entire cross-sectional data (combining TOP and NCNG data), yet with

smaller correlational differences when comparing 1.5T ($r = 0.98$, 95% CI [0.97, 0.98], $p < .001$) to

3T ($r = 0.92$, 95% CI [0.91, 0.93], $p = .004$).

While our findings indicate an association between QC parameters EFC and FBER and brain age in

all BBSC subjects when controlling for age and constant scanning parameters and scanner site, no

QC parameters were significantly associated with brain age after adjustments for multiple

comparisons in FTHP1. Based on that, one could speculate that scan quality impacts brain age

predictions when particpant ages are sampled from under-represented age groups within the

prediction model. For example, Jirsaraie et al. (2022) showed neural networks' reliability of brain

age predictions was lowest add the ends of the age distributions across scanning sites, and

predictions were less consistent when image quality was low. Furthermore, QC metrics might be

sensitive to individual differences, and vary across scanner sites. FTHP1 results also suggest a

strong effect of field strength on brain age. This indicates overall that brain age estimates are

potentially dependent on intra-individual variables in addition to acquisition parameters and other

scanner site specific covariates. While we cannot generalise from the obtained single-subject results

(FTHP1) on field strength, the additional analyses on external datasets support the effect of field

strength congruent with Jirsaraie et al.'s (2022) findings of lower prediction errors at 1.5T

compared to 3T. This was expressed in our analyses as generally higher brain age estimates at 1.5T

compared to 3T, and higher prediction errors at 3T in both cross-sectional and longitudinal data.

Finally, we show that prediction error in longitudinal data can be much higher than anticipated from

cross sectional estimates, without the presence of mental or physical disorder (see BBSC3 in **Table**

1, compare **Tables 2-3**).

A potential approach for future brain age modelling could be to employ multiple, more specific

models which are better tuned to individual differences, developmental trajectories, and scan

quality. Such models could for example be trained on data with a smaller age range and a single

field strength. Dependent on these parameters, brain age predictions can then be made by a model

selected based on the available scan and group the individual belongs to.

Conclusion

Variability in brain age predictions complicate the metric's clinical usage, for example, as a (pre-)

diagnostic tool. We presented small correlations between age and brain age when repeatedly

sampling T1-weighted MRI data from the same individual in a short period of time (1-3 years).

Reasons might lay in the absence of maturation effects for the age range in the presented sample,

brain age model-bias (including a bimodal or trimodal age training distribution) and model error.

While limited, our results suggests an influence of field strength and mixed evidence for scan

quality on brain age. Individual differences and the processing of such in the brain age model, might

lead to variability in associations between brain age and QC metrics. The presented testing of an

established brain age model on multiple single-subject short-timespan retesting data is a stricter test

than the usual use-case and does not invalidate MRI group differences. However, intra-individual

differences contributing to brain age require further attention in order to advance brain age as a

clinical tool.

Materials and Methods

Participants

We used two datasets for the analyses which had received ethics approval with all participants

consenting formally previously (Opfer et al., 2022; Wang et al., 2022, 2023). The first dataset was

the Bergen Breakfast Scanning Club (BBSC) dataset (Wang et al., 2022, 2023), including three

male subjects (BBSC2:start-age$_{BBSC2}$ = 27, BBSC1:start-age$_{BBSC1}$ = 30, and BBSC3:start-age$_{BBSC3}$ =

40) who were scanned over the period of circa one year with a summer break in the middle of the

scanning period (Wang et al., 2022). This resulted in a total number of N$_{BBSC}$ = 103 scans, relatively

equally distributed across subjects (N$_{BBSC1}$ = 38, N$_{BBSC2}$ = 40, N$_{BBSC3}$ = 25). The second dataset was

the frequently travelling human phantom (FTHP) MRI dataset (Opfer et al., 2022), including one

male subject (FTHP1:start-age$_{FTHP}$ = 48) with 157 imaging sessions at 116 locations, resulting in a

total of N$_{FTHP}$ = 557 MRI volumes. Of these, we excluded N = 6 volumes based on errors in the

processing pipeline, resulting in a final sample for the main analyses of N_{FTHP} = 551. For quality

control (**Supplement 1**), we removed another N_{FTHP} = 25 volumes which were repeat-sequences run

at the same scanner and time without changing head position or acquisition parameters, resulting in

a final sample for the supplemental analyses of N_{FTHP} = 526.

Finally, as additional validation data, we selected healty controls from two of the cross-sectional

out-of-sample test datasets described in Leonardsen et al. (2022): locally collected data (TOP;

Tønnesen et al., 2018) and the Norwegian Cognitive NeuroGenetics sample (NCNG; Espeseth et

al., 2012), as these provided most MRI scans on healthy controls. Together these datasets include a

total of N = 209 scans of healthy controls at 1.5T (Mean$_{age}$ = 54.66±15.51), and N = 856 scans of

healthy controls at 3T (Mean$_{age}$ = 32.93±10.55).

Image acquisition and preprocessing

T1-weighted volumes of BBSC1-3 were acquired with a spin echo sequence (TE=2.95ms, TR =

6.88ms, FA = 12°, TI = 450, 188 slices, slice thickness = 1mm, in-plane resolution = 1mm×1mm,

FOV = 256mm, isotropic voxel size = 1mm^3) at a 3T GE system with 32-channel head coil (see

Wang et al., 2022, 2023). T1-weighted volumes of FTHP1 were acquired at different scanners with

various different scanning parameters (see Opfer et al., 2022 or

https://www.kaggle.com/datasets/ukeppendorf/frequently-traveling-human-phantom-fthp-dataset).

All imaging sites involved in the scanning of FTHP1 were informed that the scan was acquired for

the purpose of MRI-based volumetry. Furthermore, all FTHP sites were asked to use acquisition

parameters in accordance with the ADNI recommendations for magnetization prepared rapid

gradient-echo (MP-RAGE) MRI for volumetric analyses. Thus, the range of FTHP acquisition

parameters is representative of MRI-based volumetry in everyday clinical routine at non-academic

sites. However, the scan quality might be higher than during average clinical assessments, as only

few scans were affected by motion artifacts (relatively young healthy subject). TOP data (Tønnesen

et al., 2018) including only healthy controls were acquired at 3T on a GE 3T Signa HDxT (TR =

7.8ms, TE = 2.956ms, FA = 12°; one subset with HNS coil, one subset with 8HRBRAIN coil), and

a GE 3T Discovery GE750 (TR = 8.16ms, TE = 3.18ms, FA = 12°). NCNG data (Espeseth et al.,

2012) were acquired at a 1.5T Siemens Avanto scanner using two 3D MP-RAGE T1-weighted

sequences (TR = 2400 ms, TE = 3.61 ms, TI = 1000 ms, FA = 8°, with 160 sagital slices (1.3 x 1.3

x 1.2 mm)).

Before prediction, the volumes were automatically processed using Freesurfer version 5.3 (Fischl,

2012) and FSL version 6.0 (Jenkinson et al., 2012; Smith et al., 2004), both being widely used

open-source software packages (see for overview of advantages and disadvantages compared to

other packages: Man et al., 2015) which were validated in clinical and non-clinical samples (Clerx

et al., 2015; Fischl, 2012; Jenkinson et al., 2012; Smith et al., 2004). The processing procedure

included skull-stripping as part of Freesurfer's recon-all pipeline, linearly orienting to MNI152

space (6 degrees of freedom) using FSL's linear registration, and excess border removal. While

linear registration in FSL is sensitive to atrophy and high levels of noise (Dadar et al., 2018), this

does not apply for the current quality controlled data including only healthy controls. As

Freesurfer's skull stripping algorithm can include errors (Falkovskiy et al., 2016; Waters et al.,

2019), the images were manually checked for accuracy. A step-by-step processing tutorial including

necessary code can be found at https://github.com/estenhl/pyment-public.

Brain age estimation

We applied a fully convolutional neural network (Gong et al., 2021; Peng et al., 2021) trained on

53,542 minimally processed magnetic resonance imaging T1-weighted whole-brain images from

individuals aged 3-95 collected at a variety of scanning sites (both 1.5 and 3T field strength),

(SFCN-reg detailed in Leonardsen et al., 2022) to estimate participants' ages directly from the MRI

using Python v3.9.13. The model was tested in both clinical and non-clinical samples (Leonardsen

et al., 2022) and presented high accuracy and test-retest reliability compared to other brain age

models (Dörfel et al., 2023).

Quality control metrics

Quality control (QC) metrics were extracted for each T1-weighted volume by using the automated

MRIQC tool version 22.0.6 (Esteban et al., 2017). Of these metrics, we used those which are

calculated for the whole brain or volume, being **(1)** noise measures: contrast-to-noise ratio (CNR),

signal-to-noise ratio (SNR), coefficient of joint variation of grey and white matter (CJV), **(2)**

measures based on information theory entropy-focus criterion (EFC) and foreground-background

energy ratio (FBER), **(3)** white-matter to maximum intensity (WM2MAX), and **(4)** other measures:

full-width half-maximum (FWHM).

Statistical analyses

All statistical analyses were conducted using R (v4.1.2). First, correlations of brain age with

chronological age and additionally commonly used error metrics for brain age predictions (mean

absolute error and root mean squared error) were assessed on a participant level. We further

investigated associations between brain age and age in FTHP1 (from the Frequently Travelling

Human Phantom dataset) when partialling out scanner site and field strength, as these were

expected to influence prediction accuracy.

Further analyses focussed on associations between quality control (QC) metrics and brain age as

well as acquisition parameters and brain age. We decided to observe each single independent

variable of interest in a dedicated model, as model diagnostics indicated potential multicollinearity

in models including multiple QC metrics. Furthermore, random effect models were chosen due to

the possibility to account for variances being dependent on different grouping variables, such as ID,

scanner site, field strength, and slice thickness.

Hence, linear random intercept models at the particiapant level were used to examine associations

of individual QC metrics and brain age, while controlling for age in the BBSC dataset, by running

one model for each QC metric. Similarly, for dataset 2, we predicted each QC metric as a fixed

effect in addition to the fixed effect of age in a single model. However, we used different random

effects, namely, scanner site, field strength, and slice thickness, as dataset 2 contained only FTHP1.

We also examined single individual acquisition parameters in the FTHP dataset (including only one

subject FTHP1) as fixed effects in addition to the fixed age effect. Those acquisition parameters of

interest were field strength, manufacturer, and slice thickness. Acquisition parameters not used as

fixed effects were used as random effect at the level of the intercept in addition to scanner site. All

p-values were adjusted for multiple testing using Holm correction, marked with p_{Holm}. Standardised

β-values (β_{std}) for predictors were used for comparability across β-weights by scaling QC metrics

for each subject individually.

Finally, as a validation step, we estimated brain ages for healthy controls in NCNG and TOP

datasets and correlated the estimates with age for the entire sample, subjects which were age-

matched to the longitudinal, densely sampled individuals mean age ± five years. This provided a

baseline understanding for differences in inter and intra subject brain age variability. In a second

step, brain age gap (BAG) was examined by field strength and scanner site in the validation sample.

References

Beck, D., de Lange, A. M. G., Pedersen, M. L., Alnæs, D., Maximov, I. I., Voldsbekk, I., ... &

Westlye, L. T. (2022). Cardiometabolic risk factors associated with brain age and accelerate

brain ageing. *Human brain mapping*, *43*(2), 700-720. https://doi.org/10.1002/hbm.25680

Clerx, L., Gronenschild, H. B. M., Echavarri, C., Aalten, P., & IL Jacobs, H. (2015). Can FreeSurfer

compete with manual volumetric measurements in Alzheimer's disease?. *Current Alzheimer*

Research, *12*(4), 358-367. https://doi.org/10.2174/1567205012666150324174813

Dadar, M., Fonov, V. S., Collins, D. L., & Alzheimer's Disease Neuroimaging Initiative. (2018). A

comparison of publicly available linear MRI stereotaxic registration techniques.

Neuroimage, *174*, 191-200. https://doi.org/10.1016/j.neuroimage.2018.03.025

Dörfel, R. P., Arenas-Gomez, J. M., Fisher, P. M., Ganz, M., Knudsen, G. M., Svensson, J., &

Plaven-Sigray, P. (2023). Prediction of brain age using structural magnetic resonance

imaging: A comparison of accuracy and test-retest reliability of publicly available software

packages. *BioRxiv*, 2023-01. https://doi.org/10.1101/2023.01.26.525514

Espeseth, T., Christoforou, A., Lundervold, A. J., Steen, V. M., Le Hellard, S., & Reinvang, I.

(2012). Imaging and cognitive genetics: the Norwegian Cognitive NeuroGenetics sample.

Twin Research and Human Genetics, *15*(3), 442-452. https://doi.org/10.1017/thg.2012.8

Esteban, O., Birman, D., Schaer, M., Koyejo, O. O., Poldrack, R. A., & Gorgolewski, K. J. (2017).

MRIQC: Advancing the automatic prediction of image quality in MRI from unseen sites.

PloS one, *12*(9), e0184661. https://doi.org/10.1371/journal.pone.0184661

Falkovskiy, P., Maréchal, B., Yan, S., Jin, Z., Quian, T., O'Brien, K., ... & Roche, A. (2016).

Quantitative comparison of MP2RAGE skull-stripping strategies. In *ISMRM 2016, ISMRM*

24rd Annual Meeting & Exhibition, SMRT 25th Annual Meeting.

Fischl, B. (2012). FreeSurfer. *Neuroimage*, *62*(2), 774-781.

https://doi.org/10.1016/j.neuroimage.2012.01.021

Franke, K., & Gaser, C. (2019). Ten years of BrainAGE as a neuroimaging biomarker of brain

aging: what insights have we gained?. *Frontiers in neurology*, 789.

https://doi.org/10.3389/fneur.2019.00789

Gong, W., Beckmann, C. F., Vedaldi, A., Smith, S. M., & Peng, H. (2021). Optimising a simple

fully convolutional network for accurate brain age prediction in the pac 2019 challenge.

Frontiers in Psychiatry, *12*, 627996. https://doi.org/10.3389/fpsyt.2021.627996

Jenkinson, M., Beckmann, C. F., Behrens, T. E., Woolrich, M. W., & Smith, S. M. (2012). FSL.

Neuroimage, *62*(2), 782-790. https://doi.org/10.1016/j.neuroimage.2011.09.015

Jirsaraie, R. J., Kaufmann, T., Bashyam, V., Erus, G., Luby, J. L., Westlye, L. T., ... & Sotiras, A.

(2022). Benchmarking the generalizability of brain age models: Challenges posed by

scanner variance and prediction bias. *Human Brain Mapping*.
https://doi.org/10.1002/hbm.26144

Kaufmann, T., van der Meer, D., Doan, N. T., Schwarz, E., Lund, M. J., Agartz, I., ... & Westlye, L.
T. (2019). Common brain disorders are associated with heritable patterns of apparent aging
of the brain. *Nature neuroscience*, *22*(10), 1617-1623. https://doi.org/10.1038/s41593-019-
0471-7

Korbmacher, M., de Lange, A. M., van der Meer, D., Beck, D., Eikefjord, E., Lundervold, A.,
Andreassen, O. A., Westlye, L. T., & Maximov, I. I. (2023). Brain-wide associations
between white matter and age highlight the role of fornix microstructure in brain ageing.
Human brain mapping. (in-press). https://doi.org/10.1002/hbm.26333

Korbmacher, M., Gurholt, T. P., de Lange, A. M., van der Meer, D., Beck, D., Eikefjord, E., ... &
Maximov, I. I. (2022). Bio-psycho-social factors' associations with brain age: a large-scale
UK Biobank diffusion study of 35,749 participants. *MedRxiv*.
https://doi.org/10.1101/2022.12.12.22283360

Leonardsen, E. H., Peng, H., Kaufmann, T., Agartz, I., Andreassen, O. A., Celius, E. G., ... & Wang,
Y. (2022). Deep neural networks learn general and clinically relevant representations of the
ageing brain. *NeuroImage*, *256*, 119210. https://doi.org/10.1016/j.neuroimage.2022.119210

Man, M. Y., Ong, M. S., Mohamad, M. S., Deris, S., Sulong, G., Yunus, J., & Harun, F. K. C.
(2015). A review on the bioinformatics tools for neuroimaging. *The Malaysian journal of
medical sciences: MJMS*, *22*, 9.

Marquand, A. F., Kia, S. M., Zabihi, M., Wolfers, T., Buitelaar, J. K., & Beckmann, C. F. (2019).
Conceptualizing mental disorders as deviations from normative functioning. *Molecular
psychiatry*, *24*(10), 1415-1424. https://doi.org/10.1038/s41380-019-0441-1

Opfer, R., Krüger, J., Spies, L., Ostwaldt, A. C., Kitzler, H. H., Schippling, S., & Buchert, R.
(2022). Automatic segmentation of the thalamus using a massively trained 3D convolutional
neural network: higher sensitivity for the detection of reduced thalamus volume by
improved inter-scanner stability. *European Radiology*, 1-10. https://doi.org/10.1007/s00330-
022-09170-y

Peng, H., Gong, W., Beckmann, C. F., Vedaldi, A., & Smith, S. M. (2021). Accurate brain age
prediction with lightweight deep neural networks. *Medical image analysis*, *68*, 101871.
https://doi.org/10.1016/j.media.2020.101871

Rosen, A. F., Roalf, D. R., Ruparel, K., Blake, J., Seelaus, K., Villa, L. P., ... & Satterthwaite, T. D.
(2018). Quantitative assessment of structural image quality. *Neuroimage*, *169*, 407-418.
https://doi.org/10.1016/j.neuroimage.2017.12.059

Smith, S. M., Jenkinson, M., Woolrich, M. W., Beckmann, C. F., Behrens, T. E., Johansen-Berg, H.,
    ... & Matthews, P. M. (2004). Advances in functional and structural MR image analysis and
    implementation as FSL. *Neuroimage*, *23*, S208-S219.
    https://doi.org/10.1016/j.neuroimage.2004.07.051

Tønnesen, S., Kaufmann, T., Doan, N. T., Alnæs, D., Córdova-Palomera, A., Meer, D. V. D., ... &
    Westlye, L. T. (2018). White matter aberrations and age-related trajectories in patients with
    schizophrenia and bipolar disorder revealed by diffusion tensor imaging. *Scientific reports*,
    8(1), 14129. https://doi.org/10.1038/s41598-018-32355-9

Vidal-Pineiro, D., Wang, Y., Krogsrud, S. K., Amlien, I. K., Baaré, W. F., Bartres-Faz, D., ... & Fjell,
    A. (2021). Individual variations in 'brain age'relate to early-life factors more than to
    longitudinal brain change. *Elife*, *10*, e69995. https://doi.org/10.7554/eLife.69995

Wagen, A. Z., Coath, W., Keshavan, A., James, S. N., Parker, T. D., Lane, C. A., ... & Schott, J. M.
    (2022). Life course, genetic, and neuropathological associations with brain age in the 1946
    British Birth Cohort: a population-based study. *The Lancet Healthy Longevity*, *3*(9), e607-
    e616. https://doi.org/10.1016/S2666-7568(22)00167-2

Wang, M. Y., Korbmacher, M., Eikeland, R., & Specht, K. (2022). Deep brain imaging of three
    participants across 1 year: The Bergen breakfast scanning club project. *Frontiers in Human
    Neuroscience*, *16*. https://doi.org/10.3389%2Ffnhum.2022.1021503

Wang, M. Y., Korbmacher, M., Eikeland, R., & Specht, K. (2023). The Bergen Breakfast Scanning
    Club dataset: a deep brain imaging dataset. *bioRxiv*. doi:10.1101/2023.05.30.542072

Waters, A. B., Mace, R. A., Sawyer, K. S., & Gansler, D. A. (2019). Identifying errors in Freesurfer
    automated skull stripping and the incremental utility of manual intervention. *Brain imaging
    and behavior*, *13*, 1281-1291. https://doi.org/10.1007/s11682-018-9951-8