Combined effects of climate warming and environmentally relevant concentrations of pharmaceutical active compounds on a freshwater community

Claire Duchet[1, 2✉], Kateřina Grabicová [3], Vojtech Kolar [1,2], Olga Lepšová[4], Andras Csercsa[1],

Barbora Zdvihalova[1], Tomáš Randák [3], David Boukal[1,2]

[1]Department of Ecosystem Biology, Faculty of Science, University of South Bohemia,

Branišovská 1760, 37005 České Budějovice, Czech Republic

[2]Institute of Entomology, Biology Centre of the Czech Academy of Sciences, Branišovská

31, 37005 České Budějovice, Czech Republic

[3]Laboratory of Environmental Chemistry and Biochemistry, Research Institute of Fish

Culture and Hydrobiology, University of South Bohemia, Zátiší 728/II, 389 25 Vodňany,

Czech Republic

[4]Department of Botany, Faculty of Science, University of South Bohemia, Branišovská 1760,

37005 České Budějovice, Czech Republic

✉Corresponding author: Claire Duchet

E-mail address: cduchet@prf.jcu.cz

Department of Ecosystem Biology, Faculty of Science, University of South Bohemia

Branišovská 1760, 37005 České Budějovice, Czech Republic

Tel. +420 387 772 244 (office); Fax: +420 387 772 368

Abstract

Predicting the combined effects of stressors on ecosystems is a pressing concern due to increasing anthropogenic pressures on the Earth's biota. Climate warming and chemical pollution are two major stressors affecting freshwater biota. The latter is becoming more common due to the widespread use of mixtures of chemical compounds and the low removal efficiency of water treatment plants, especially in the case of pharmaceutical active compounds (PhACs).

To evaluate the effects of warming and PhACs, we conducted a full factorial experiment (with/without PhAC mixture and with/without warming) in heated outdoor mesocosms with a common community of macroinvertebrates and plankton taxa from different trophic levels. The experiment was conducted twice, in winter and in summer. We repeatedly measured PhAC concentrations, environmental parameters, zooplankton density, and aquatic insect emergence. The summer experiment showed much stronger effects of stressors, dominated by temperature effects, on community structure and temporal dynamics of individual taxa. Both warming and PhACs altered invertebrate community composition in the summer experiment, with contrasting effects observed between insect emergence and phytoplankton and zooplankton responses. Our results suggest that PhACs at environmentally relevant concentrations can alter the effects of climate warming on freshwater biota, with individual-level responses such as delayed or accelerated development and altered phenology of key taxa.

Keywords: warming, pharmaceutical active compound mixture, freshwater invertebrates, community-levels, trophic interactions

Introduction

Freshwater ecosystems are severely impacted by numerous stressors, leading to strong declines in species abundance and biomass (Sánchez-Bayo and Wyckhuys, 2019) (Hallmann et al., 2017). The main causes of species declines are habitat loss due to intensive agriculture and urbanisation, pollution, increases in pathogens and invasive species, and climate change (Sánchez-Bayo and Wyckhuys, 2019). Climate warming and chemical pollution are two ubiquitous stressors (Loos et al., 2013; Parmesan, 2006) that lead to rapid environmental changes and pose a major threat to freshwater biodiversity and ecosystem functioning (Vörösmarty et al., 2010).

Pharmaceutically active compounds (PhACs) represent an emerging class of chemical pollutants. Their importance has increased due to the growing human population and recent advances in pharmaceutical research (Fekadu et al., 2019). Due to their widespread use and low removal efficiency in wastewater treatment plants, PhACs are found in aquatic systems worldwide (Brodin et al., 2014; Koba et al., 2018; Loos et al., 2013; Wilkinson et al., 2022). PhACs bioaccumulate in fish (Du et al., 2014) and benthic macroinvertebrates (Grabicova et al., 2015). At environmentally relevant concentrations (ng.L^{-1} to low µg.L^{-1}), PhACs can affect the behaviour of aquatic ectotherms (Brodin et al., 2014; Saaristo et al., 2019), alter feeding rates (Bláha et al., 2019), and disrupt reproduction (Fursdon et al., 2019). These responses directly affect individual fitness (Fursdon et al., 2019) and can trigger indirect effects such as altered trophic interactions (Bláha et al., 2019). Altered species phenologies and trophic interactions can also redirect the flow of PhACs in food webs through bioaccumulation (Lagesson et al., 2016) and biomagnification (Zenker et al., 2014), increasing their potential toxicity to higher trophic levels including fish and humans.

PhACs are usually detected as complex mixtures in the environment (Grabicova et al.,

2015; Koba et al., 2018; Loos et al., 2013). However, their effects on aquatic ecosystems are

primarily understood through single-species laboratory bioassays, often conducted at

concentrations higher than those commonly found in the environment (Richards et al., 2004).

More empirical data are therefore needed to understand the ecotoxicology of PhAC mixtures

under environmentally realistic conditions and the quantitative consequences of PhACs on

ecological interactions and whole communities (Backhaus, 2014).

Average temperatures are predicted to increase by +1.4–4.4°C by 2100 due to a global

warming (Masson-Delmotte et al., 2021). Freshwater ecosystems dominated by ectotherms are

expected to be particularly impacted by warming (Parmesan and Yohe, 2003; Woodward et al.,

2010). These organisms have a limited dispersal ability and therefore have to deal directly with

changing environments (Woodward et al., 2010). These changes often result in reduced body

sizes, altered species interactions, shifts in species distribution ranges and phenology, and

cascading impacts on community dynamics and ecosystem functioning (Boukal et al., 2019;

Post, 2013). Due to the strong correlation between water surface and air temperatures, climate

warming will strongly affect communities in shallow freshwaters. Thermal stress often prompts

organisms to seek refuge in colder waters at the bottom of water bodies (Balogová and

Gvoždík, 2015), but even these microhabitats may be close to the upper thermal tolerance limits

of many species during heat waves (Stoks et al., 2017).

Despite the widespread distribution of PhACs in freshwater ecosystems and the reality

of climate warming, we currently lack evidence of the combined effects of PhACs and warming

at the population and community level (Heye et al., 2019). Typically, studies have only

examined the combined effects of temperature and single toxicants on individual organisms

and have not considered how PhACs may interact with other stressors at the community level.

The effects of multiple stressors are complex and can be synergistic (greater than the sum of

the individual effects), antagonistic (due to opposing effects of two or more stressors) or

additive (sum of the individual effects; Sala et al., 2000), making predictions about the effects

of multiple stressors difficult (Spaak et al., 2017). However, recent advances in predicting the

interactions between warming and other stressors have shown that impacts on freshwater

ecosystems are usually best explained by the effect of the stronger stressor alone (dominance

null model; Morris et al., 2022).

Warming can modulate the effects of pollutants on freshwater ecosystems, both directly

and indirectly. While many pollutants become more toxic at higher temperatures, warming may

also accelerate their degradation, thereby reducing the exposure of biota to PhACs (Noyes et

al., 2009; Op de Beeck et al., 2017). Population-level responses to the combined stressors can

vary dramatically due to differences in environmental sensitivity (Venâncio et al., 2016),

tolerance of individuals (Baert et al., 2016), ecological trade-offs and local adaptation patterns

(Bennett and Lenski, 2007; Kneitel and Chase, 2004). Furthermore, the fate of species

embedded in communities depends on direct and indirect changes in intraspecific and

interspecific interactions (Post, 2013; Relyea, 2009). These include effects mediated by indirect

traits and population densities and include top-down and bottom-up effects (Abrams, 1996;

Brogan and Relyea, 2015; Guedes et al., 2016; Rodrigues et al., 2018). Persistent pollutants

can also alter community structure and affect or enhance various aspects of ecosystem

functioning (Duchet et al., 2018; Sánchez-Bayo, 2021).

To address these knowledge gaps, we conducted a full-factorial experiment to

investigate the combined effects of warming and a mixture of PhACs on freshwater

communities in outdoor mesocosms. Our experimental community consisted of common

macroinvertebrate and planktonic taxa from different trophic levels and feeding guilds,

including predators, filter feeders, grazers, detritivores, and primary producers. We exposed

the community to a single-pulse cocktail of 15 PhACs from major drug categories, including

cardiovascular, psychoactive, antihistaminics, and antibiotics, at concentrations commonly

found in surface waters (Fedorova et al., 2022; Grabicova et al., 2015; Koba et al., 2018;

Švecová et al., 2021), in combination with warming. We predicted an interactive effect between

the PhAC cocktail and warming on community composition and abundance, resulting in

reduced grazing pressure and an increase in phytoplankton biomass cascading through the food

web. Specifically, we predicted that high temperatures above therma optima (25°C for most

taxa in the community) in summer would have a strong and negative effect on the community,

whereas in winter, warming would have a mild but positive effect. The combined effects of

PhACs and warming on predators may indirectly affect filter feeder biomass by top-down

control, and thus indirectly increase phytoplankton biomass due to reduced grazing pressure

(Belden et al., 2007; Polazzo et al., 2022).

133   2. Material and Methods

2.1. Mesocosms

The mesocosm complex is located at the University of South Bohemia (České Budějovice,

Czech Republic; 48°58'36.6"N 14°26'41.8"E, 381 m above sea level). It consists of 32 circular

outdoor mesocosms (inner diameter 120 cm, water volume 1.13 m^3 each) set into the ground

and equipped with a custom-built, remote-controlled heating system. The 28 mesocosms used

in this study were filled 6 weeks before the start of the experiment with tap water filtered

through a granulated activated carbon filter and enriched with 100 g of dry leaves of common

alder (*Alnus glutinosa*) and 8 g of fish food (Pondstick granules, Apetit®) to add nutrients. The

water in each mesocosm was gently mixed with an air stone to avoid thermal stratification,

except in winter when mixing was stopped in all treatments between 12 January 2021 and 03

February 2021 to allow the water surface in the unheated mesocosms to freeze naturally.

The experimental community included common macroinvertebrate and plankton taxa

from different trophic levels and feeding guilds (predators, filter feeders and grazers,

detritivores, and primary producers; Table S1.1). During the maturation period, the mesocosms

were populated with phytoplankton and zooplankton from a nearby fishpond one month before

the start of the experiment. The water with phytoplankton and zooplankton was thoroughly

mixed in a barrel before a 2.5-litre aliquot was added to each mesocosm. Two weeks later, the

mesocosms were gradually inoculated with macroinvertebrates from nearby fishponds, ditches

and sand pits: water column predators (*Notonecta glauca* backswimmers, 9 per mesocosm and

Aeshna or *Anax* dragonfly larvae, 6 individuals per mesocosm; hereafter *Aeshna* and *Anax*),

benthic predators (libellulid dragonfly larvae, 3 per mesocosm), benthic detritivores (*Asellus*

aquaticus, ca. 100 per mesocosm), grazers (*Cloeon* cf. *dipterum*, mayfly larvae, ca. 200 per

mesocosm), omnivorous scrapers (*Planorbarius corneus* pond snails, 9 individuals per

mesocosm), omnivorous piercers (*Sigara falleni*, water boatmen, 50 individuals per

mesocosm), and phytophilous predators occupying artificial vegetation and mesocosm walls

(damselfly larvae, ca. 50 per mesocosm). The predators were added to the mesocosms only one

week before the start of the experiment to allow the community to settle in.

Three artificial submerged macrophytes made of strips of green plastic mesh (5×100 cm)

attached to a granite stone were added to each mesocosm to provide additional microhabitat.

Two plastic strips (6 cm wide, 50 cm long) hanging vertically in each mesocosm were used to

incubate periphyton that had been pre-cultured for four weeks under natural conditions in one

of the fishponds where the macroinvertebrates were also collected. Finally, a coarse (20x20

cm; 20 mm mesh) and a fine (20x20 cm; 0.5 mm mesh) litter bag containing 9.5 g of dried

alder (*Alnus glutinosa*) leaves were added in each mesocosm to distinguish the contribution of

detritivores and microbial decomposers, respectively, to the decomposition process (Fig. 1).

Figure 1. Schematic view of the mesocosm set-up

2.1. Experimental design

Seven mesocosms were each assigned to one of four treatments: (1) not heated, without PhACs, (2) not heated, treated with PhACs, (3) heated, treated with PhACs, and (4) heated, with PhACs. Each mesocosm was subjected to the same treatment in both experiments. Heating was maintained at +4°C above ambient throughout the experiment (see temperature profiles for each experiment in S2). The PhAC mixture was a cocktail of 15 compounds from the major drug categories (cardiovascular: telmisartan, valsartan, metoprolol, atenolol; psychoactive: carbamazepine, lamotrigine, venlafaxine, citalopram, tramadol; antihistaminic: cetirizine, fexofenadine; antibiotic: sulfamethoxazole, trimethoprim, clarithromycin, clindamycin; Table S3.1), at concentrations commonly detected in surface waters in the Czech Republic (Fedorova et al., 2022; Grabicova et al., 2015; Švecová et al., 2021). The total added PhACs were 20 ng.L^{-1} for citalopram and fexofenadine, 50 ng.L^{-1} for atenolol and clindamycin, 100 ng.L^{-1} for clarithromycin, sulfamethoxazole, trimethoprim and venlafaxine, 150 ng.L^{-1} for valsartan, 200 ng.L^{-1} for cetirizine, 250 ng.L^{-1} for carbamazepine and metoprolol and 500 ng.L^{-1} for

186 lamotrigine, telmisartan and tramadol (total concentration: 2890 ng.L^{-1}). The PhAC mixture

187 was administered as a single pulse exposure.

188  We conducted the experiment twice, from 24 September 2020 to 12 March 2021

(hereafter 'winter experiment') and from 2 June to 2 August 2021 (hereafter "summer

experiment'; Figures S4.1 and S4.2). We refer to the start of each experiment (just before the

PhACs were added and the warming started) as Day 0. The first three samplings for all the

above parameters were done on Day 0 before the PhAC mixture was added on Day 2 and on

Day 7. From then on, samples were taken monthly in winter and fortnightly in summer, until

the end of the experiment.

2.3. Sampling

In both experiments, we repeatedly measured PhAC concentrations, environmental parameters

(dissolved O$_2$, conductivity, pH, turbidity, and temperature with a YSI 556 MPS probe (®YSI,

USA)), chlorophyll-a concentration, zooplankton density and aquatic insect emergence (the

latter occurred only in summer; traps were emptied three times per week).

201  Zooplankton was sampled with a tube sampler (length 1.2 m, inner diameter 7 cm)

equipped with a one-way valve at the bottom. Water column samples were taken from three

regularly spaced locations within each mesocosm to reduce the effects of plankton patchiness

(Arnold et al., 1991; Stephenson et al., 1984). The resulting bulk sample (3 × 3L, depending

on the water level in the mesocosms: min = 2.5L, max = 3.2L in summer; min = 2.8L, max =

3.5L in winter) was collected in a 20L bucket. One litre of the 9L of the bulk sample was

filtered through a plankton net with 80-μm mesh size. The retained organisms (zooplankton

and some other pelagic invertebrates) were transferred to a 120 mL plastic vial and preserved

with formaldehyde (4%). Fixed specimens were counted under a stereomicroscope (Olympus

SZX2-ILLT stereomicroscope; Olympus, Tokyo, Japan) and identified to the lowest feasible

taxonomic level using identification keys (Amoros, 1984; Johannsen, 1937). Biomass of each

taxon was estimated from the literature (Table S5) or from our own measurements on the

macroinvertebrates.

On each sampling day, water samples (6 mL) were collected in each mesocosm, filtered

through a 0.2-µm filter and stored at -20°C until the PhAC compound concentrations were

analysed with an in-line solid-phase extraction liquid chromatography with tandem mass

spectrometry (in-line SPE/LC-MS/MS, triple quadrupole mass spectrometer Quantiva, Thermo

Fisher Scientific, USA; see Text S3 for details).

Dissolved organic carbon (DOC) and a measure of nitrogen and phosphorus

concentration were measured at Day 0 and at the end of each experiment. For technical reasons,

total nitrogen (TN) and total phosphorus (TP) were measured during the winter experiment,

while dissolved nitrogen (DN) and soluble reactive phosphorus (SRP) were measured during

the winter experiment.

Phytoplankton samples were collected from the surface water layer of 0-1 m in 500 ml

PET bottles and fixed with Lugol solution immediately after sampling. After complete

sedimentation in the laboratory, the water was decanted, and the sample transferred to a Bürker

counting chamber. The algal cells were counted using an optical microscope (OLYMPUS

CX23) at 400× magnification. The results are given in cell number.mL^{-1}. Taxa were identified

to species or genus level (where possible) according to (Kaštovský et al., 2018a, 2018b). We

used the currently accepted phytoplankton nomenclature based on the AlgaeBase (Guiry and

Guiry, 2023).

During the summer experiment, a pyramid-shaped floating emergence trap (35 × 35 cm

base dimensions) with a 250 mL collection bottle of soapy water (Cadmus et al., 2016) was

installed in each mesocosm on Day 0. Emerging insects were collected every 2-3 days and

preserved in 70% ethanol (mayflies, chironomids) or frozen (odonates) for later identification

and analysis of PhAC content.

At the end of the experiment, macroinvertebrates were destructively sampled and

periphyton stripes were collected and analysed to quantify algal growth. The contents of the

litter bags were dried at 60°C for 48 hours before being weighed to quantify the litter

decomposition process by invertebrate detritivores and microorganisms.

We focus here only on the temporal dynamics of the pelagic community during the

experiment and include in it only the predatory insects feeding on zooplankton. The whole

community data collected at the end of each experiment are subject to a separate study (see

summary of response variables collected during the experiments, Table S6).

2.3. Statistical analyses

Temporal changes in environmental parameters (pH, dissolved O_2, conductivity, turbidity and

water level) and chlorophyll-a concentrations during the experiment were tested by a

redundancy analysis (RDA on centred data) separately for the summer and the winter

experiment. Treatment was used as a categorical explanatory variable and environmental

variables were used as a response. Day of the experiment was used as a covariate. Statistical

significance of the RDAs was tested by Monte Carlo permutation tests (999 unrestricted

permutations under the reduced model in split-plot hierarchical design).

Differences in community composition of algal groups and species between treatments

were visualised by principal component analyses (PCAs) with centred and log-transformed

data. Only species found in more than three samples were analysed.

Changes in the planktonic community structure were analysed using the principal

response curve (PRC) method (Van den Brink and Braak, 1999) using the most practical level

of taxonomic identification for each group. Interaction of treatment and day was used as the

explanatory variable and community composition as a response. Day of the experiment was

used as a covariate. Before analysis, the biomass of each taxon was log-transformed. Monte

Carlo permutation tests were performed on each sampling date to identify the dates on which

the treatments differed significantly. In all analyses, the permutations were set in a hierarchical

split-plot design and thus permuted for each mesocosm individually. All multivariate and PRC

analyses were conducted using CANOCO 5 (Braak and Smilauer, 2012).

We ran generalised linear mixed models (GLMMs; implemented in packages *glmmTMB*

and *sjplot*) with the appropriate distribution and link functions. A log-link Gamma GLMM was

used for the analyses of the treatment effects on nutrients (dissolved organic carbon, nitrogen

and phosphorus concentrations), while binomial GLMMs were used for the analyses of

treatment effects on cumulative emergence of predators in summer and on predator survival in

winter, respectively. These analyses used only data measured at the end of each experiment.

Probability of successful emergence of odonates in the summer experiment was defined as the

ratio of the estimated number of individuals emerged by Day 56 to the size of the inoculum.

Similarly, survival probability of odonates and *Notonecta* in the winter experiment was defined

as the ratio of the estimated number of individuals collected on Day 169 to the size of the

inoculum. Model selection approach with the corrected Akaike information criterion was used

to identify the most parsimonious model for each univariate response.

Finally, to disentangle the direct and indirect effects of the PhAC mixture and warming

on interactions within the aquatic community, we tested simple networks of cause-effect

relationships describing a trophic cascade and predator-prey relationships using confirmatory

path analysis (piecewise SEM; (Shipley and Douma, 2020), implemented in *piecewiseSEM*

package (Lefcheck, 2016)). This approach can identify mechanistic pathways underlying the

effects of different stressors on aquatic ecosystems (Schmidt et al., 2022). Using *a priori*

knowledge, we built a complete model of cause-effect relationships based on hypothetical

pathways supported by literature (Table S1.1). PhACs and warming were assumed to directly

reduce the biomass of filter feeders (= cladocerans in our mesocosms) and thus indirectly

increase phytoplankton biomass due to reduced grazing pressure (Belden et al., 2007). We also

assessed the effects of PhACs and warming on predators (odonates in summer, and odonates

and *Notonecta* in winter), which indirectly affect filter feeder biomass by top-down control.

PhACs and warming were used as predictors. Phytoplankton (chlorophyll-*a* concentration),

filter feeder biomass and predator biomass were used as response variables. For predator

biomass, we used cumulative emergence of odonates in summer and the survival of *Notonecta*

and odonate larvae in winter, as there were very few odonate larvae left at the end of the

summer experiment and no adults emerged during the winter experiment. We used Fisher's C

statistics to assess the overall fit of each SEM model, with *p* >0.05 defined as indicating a good

fit for each model. All these analyses were done in R version 4.1.2 (RStudio, 2020; R Core

Team, 2018).

3. Results

3.1. PhAC concentrations

Concentrations of the PhAC mixture gradually decreased over time to ~25% of the initial

concentration in the heated mesocosms and to ~30% in the non-heated mesocosms in the

summer experiment (Fig. 2A). Metabolite concentrations increased slightly in the first month

and then remained stable until the end of the experiment (Fig. 2B). In the winter experiment,

the PhACs were also degraded during the course of the experiment (Fig. 2C). In contrast to the

summer experiment, however, the metabolite concentrations increased greatly during the first

two months of the experiment, reaching 50% of the initial concentration of the parent

compounds, and then remained stable (Fig. 2D). Concentrations of all PhACs and their

metabolites in both experiments were below detection limits in all samples taken from the non-

treated mesocosms.

Figure 2. Total concentrations of PhAC main compounds (A, C) and their metabolites (B, D) in the

winter (A, B) and summer (C, D) experiments in heated (orange) and non-heated (blue) mesocosms.

Symbols with error bars = treatment-specific mean values ± SD (n = 7); mean values connected by

colour lines. Data include only treated mesocosms.

3.2. Environmental conditions

Environmental conditions in the summer experiment differed significantly between treatments

in the summer (RDA: pseudo-F = 4.9, p = 0.002, explained variation = 7.4%) and winter (RDA:

pseudo-F = 53.7, p = 0.001, explained variation = 43.1%) experiments. In the summer

experiment, non-heated mesocosms were characterised by higher water levels, higher turbidity,

lower conductivity and lower chlorophyll-*a* concentrations (the latter mainly in the non-treated

treatment) compared to heated mesocosms (Fig. 3A, Table S7.1). In the winter experiment, the

treatments did not differ in chlorophyll-*a* concentration, dissolved oxygen, turbidity and pH,

but non-heated mesocosms were characterised by higher water levels and lower conductivity

compared to the heated ones (Fig. 3B, Table S7.2).

Figure 3. Ordination diagram (RDA) showing differences between the treatments in environmental parameters

and chlorophyll-*a* concentration during the summer (A) and winter (B) experiment. Treatments: CLT = non-

heated, non-treated; CHT = heated, non-treated; TLT = non-heated, treated; THT = heated and treated.

3.3. Effect of the stressors on phytoplankton and nutrients

Community composition showed no significant differences in the major algal groups between

treatments in both the summer and winter experiments (PCA, Fig. S8.1). Species composition

was more homogeneous and PCA showed no specific patterns in the summer experiment (Fig.

S8.1AB). However, during the winter experiment, Chrysophyta seemed to be more abundant

in the non-heated controls, while Streptophyta and Cryptophyta were comparatively more

abundant in the non-heated, treated treatment compared to other treatments. Nevertheless, there

was considerable overlap in the samples (Fig. S8.1C). In terms of species composition,

temperature was the main driver of differences in the winter experiment (second axis in Fig.

S8.1D). The Bacillariophyta *Achnanthes* sp. and the Chlorophyta *Oocystis* spp. and

Scenedesmus sp. tended to be more abundant in the heated controls, while Chrysophyta tended

to be more abundant in the non-heated controls. The Bacillariophyta *Synedra* sp., *Gomphonema*

spp., *Navicula* sp. and *Ulnaria* were more abundant in the non-heated, treated treatment, while

none of the species were more abundant in the heated treatment with added PhACs than in the

other treatments (Fig. S8.1D).

The most parsimonious models identified partly disparate effects of the stressors on

nutrients in the summer and winter experiments. In the summer experiment, the final

concentrations of DOC, DN and SRP were lower in the heated mesocosms than in the non-

heated ones and the added PhAC mixture had a small negative effect on DOC and a significant

negative effect on DN and SRP (Fig. S8.2ACE). In the winter experiment, the final

concentrations of DOC, TN and TP were also lower in the heated mesocosms than in the non-

heated ones, while no effects of PhACs on nutrients could be detected (Fig. S8.2BDF). In both

experiments, we did not detect any synergistic effects of the two stressors on nutrients (Tables

S8.1 and S8.2).

3.3. Effect of the stressors on zooplankton community dynamics

The dynamics of the zooplankton community and its responses to the stressors also differed

markedly between the summer and winter experiments. In the summer experiment, the biomass

of the main zooplankton groups differed significantly between treatments (PRC: F = 0.7, p =

0.003; Fig. 4A). More specifically, differences between treatments appeared immediately after

Day 0 and continued to increase until Day 35, with differences primarily caused by warming

rather than PhACs and mainly affecting ostracods and *Daphnia* (Fig. 4A). Ostracod biomass

increased while *Daphnia* biomass decreased in the heated mesocosms compared to the non-

heated mesocosms, while the other groups did not contribute to the differences between

treatments (Fig. 4A). That is, *Daphnia* biomass decreased in all treatments, but the decline was

steeper in the heated mesocosms (Fig. S9.1). On the other hand, copepod biomass tended to

increase towards the end of the experiment in all treatments, with an intermediate peak in

biomass around Day 35, which appeared to be higher in the treatments without PhAC addition

(Fig. S9.2). In contrast, zooplankton composition did not differ significantly between

treatments in the winter experiment ($F = 0.3$, $p = 0.41$; Fig. 4B).

375
376
377

Figure 4. Principal response curves (PRC) with species weights (vertical bars) indicating treatment-

specific effects on zooplankton during the summer (A) and winter (B) experiment. Treatments: non-

heated, without added PhACs (light blue); yellow line: heated, non-heated, without added PhACs

(yellow); non-heated, with added PhACs (dark blue); heated, with added PhACs (orange). Days = days

after exposure.

3.3. Effect of the stressors on insect predator survival and emergence

The effects of the two stressors on predatory insects differed between taxa and experiments

(Fig. 5).

Figure 5. Estimates of treatment-specific probability of emergence of insect predators in the summer

experiment (A: anisopteran larvae (*Aeshna*), B: zygopteran larvae) and probability of survival of insect

predators in the winter experiment (C: zygopteran larvae, D: *Notonecta* backswimmers). Model

estimates of the most parsimonious models, shown as mean values with 95% confidence intervals based

on fixed effects.

The effects of warming and PhACs on the cumulative probability of emergence in the summer

experiment and on the survival probability in the winter experiment differed between taxa

(Tables S10.1 and S10.2). In the summer experiment, both warming and PhACs increased the

cumulative number of anisopteran (*Aeshna*) larvae that emerged, with an additive effect on the

predictor scale (Fig. 5A). In contrast, warming had a negative effect on damselfly emergence

in the summer experiment (Fig. 5B). In the winter experiment, survival of anisopteran (*Anax*)

larvae was similar in all treatments, while damselfly larvae survived better when exposed to

PhACs only or heating only, with an antagonistic effect of both stressors resulting in similar

survival of the larvae in the treatments without both stressors and with both stressors(Fig. 5C).

Finally, *Notonecta* survived less when exposed to warming, but PhACs had no effect on their

survival in the winter experiment (Fig. 5D).

3.3. Effect of the stressors on the pelagic food web

The effects of the two stressors on relationships in the pelagic food web differed markedly

between the summer and winter experiments (Fig. 6 and Tables S11.1 and 11.2). In the summer

experiment, chlorophyll-*a* concentration as a proxy of phytoplankton biomass was directly and

negatively affected by PhACs (standardised path coefficient = -0.0001; Fig. 6A and Table

S11.1) and directly and positively affected by warming (standardised path coefficient = +0.08).

Both stressors had a direct and negative effect on cladocerans as the main filter-feeding group

(standardised path coefficients = -0.0001 for PhAC and -0.14 for warming). However, the

relationship between primary producers and filter feeders was positive and not significant

(standardised path coefficient = -0.09). These results do not support the hypothesis that a

decrease in filter-feeding biomass due to the stressors leads to a trophic cascade and thus to

higher algal biomass. The effects of stressors on the predators were positive but not significant

(direct effect, standardised path coefficients = +0.0001 for PhAC and +0.02 for warming);

however, predators caused a direct decrease in filter feeder biomass (standardised path

coefficient = -0.21; Fig. 6A).

In contrast, the PhAC mixture had no direct or indirect significant effect on any

component of the simplified pelagic food web in the winter experiment, while only the

phytoplankton biomass was directly and positively affected by warming during winter

(standardized path coefficient = 0.28; Fig. 6B and Table S11.2).

432   **Figure 6.** Path diagram of the selected structural equation model based on average data collected during

433   the summer experiment, and last sampling data during the winter experiment. Orange, blue, and dashed

434   arrows respectively indicate significantly ($p < 0.05$) negative, significantly positive, and non-significant

435   ($p > 0.05$) relationships between variables. Proportions of variation explained by the model for each

436   response variable given by R^2 values. See Tables S5 and S6 for details.

4. Discussion

To the best of our knowledge, this is the first study to examine the combined toxic effects of a mixture of PhACs at environmentally relevant concentrations and warming on freshwater communities. In the winter experiment, we observed very limited effects, while in the summer experiment, warming acted as the stronger stressor and amplified the effects of PhACs on the freshwater community. This led to a PhAC- and temperature-mediated trophic cascade in the planktonic food web, with warming negatively affecting zooplankton and promoting phytoplankton development. Although no effects of PhACs and warming on secondary consumers were observed, our results suggest that low and environmentally realistic concentrations of PhACs may be detrimental to the aquatic community at higher temperatures as a resulting of climate warming. This was evidenced by a significant negative effect of the combined stressors on filter feeder biomass, which even reversed the expected result based on the effects of the individual stressors.

4.1. Effects of warming on the freshwater community

Temperature was the stronger stress factor for the freshwater community in our study, especially in summer. During the summer experiment, the highest temperatures in the heated mesocosms reached up to 29.5°C (compared to 25.5°C in the non-heated mesocosms) and remained above 25°C for almost three weeks in July.

The effects of global warming on freshwater ecosystems have been well studied recently and are expected to have significant impacts on the pelagic communities. For example, warming can stimulate phytoplankton growth in temperate and shallow lakes (Zhang et al., 2019), leading to altered primary productivity, nutrient recycling, and higher trophic productivity (Mooij et al., 2008). However, warming may also promote the development of bacteria that can increase competition for resources with phytoplankton (Joint et al., 2002). The

increased stratification caused by warming may alter plankton structure and dynamics,

favouring small-bodied and buoyancy-regulating species, ultimately leading to changes in

primary productivity, nutrient cycling, and higher trophic levels (Winder and Hunter, 2008).

Warming is not expected to have a significant effect on the total phytoplankton biomass, but

the associated phenomenon of brownification can significantly reduce local phytoplankton

biodiversity due to the dominance of mixotrophic algae that displace other phytoplankton taxa

(Urrutia-Cordero et al., 2017). In our study, we observed an increase in phytoplankton biomass

(measured by chlorophyll-*a* concentration) under warming in both the summer and winter

experiments. However, by the end of the summer experiment, the phytoplankton community

in the heated mesocosms shifted towards a dominance of cyanobacteria and diatoms, while

green algae continued to dominate the non-heated mesocosms.

Moreover, the effects of warming on phytoplankton are context-specific and may be

offset by the grazing effects of filter feeders such as *Daphnia* and other zooplankton taxa

(Kazanjian et al., 2018; Velthuis et al., 2017). Invasive species may also colonize lakes in

response to warming, but the long-term consequences of this disturbance may differ from short-

term observations (Dziuba et al., 2020). In our summer experiment, the zooplankton

community was initially dominated by *Daphnia* and other large cladocerans and gradually

shifted to the dominance by smaller zooplankton species in response to warmer temperatures.

Finally, the presence of top predators may also exacerbate the effects of climate warming

on phytoplankton in shallow aquatic ecosystems by reducing grazing pressure on zooplankton

(Hansson et al. 2013; Šorf et al., 2015; Velthuis et al., 2017; Yvon-Durocher et al., 2011). Our

observations corroborate these studies, as we found that phytoplankton biomass was

significantly higher in the warmer mesocosms, and we attribute this pattern to a lower

zooplankton grazing pressure caused by both predation by odonate larvae and higher

temperatures. Overall, our results reiterate that the complex and non-linear responses of

freshwater communities to environmental stressors should be considered when predicting

future biodiversity patterns (Polazzo et al., 2022).

4.2. Effects of PhACs on the freshwater community

PhACs are emerging contaminants that can have toxic effects on freshwater communities (Fent

et al., 2006). The toxicity of PhACs can be even more pronounced when present in mixtures,

causing significant structural and functional changes in microbial and algal communities and

affecting the survival and behaviour of organisms from zooplankton to macroinvertebrates to

fish (Brodin et al., 2014). We observed no significant effects of PhACs in the winter

experiment. However, in the summer experiment, the PhAC mixture alone had a negative effect

on phytoplankton biomass (chlorophyll-*a* concentration) as well as on nutrients (DN, SRP).

The PhAC mixture also negatively affected predator survival, which may have attenuated the

indirect effect of lower phytoplankton biomass on cladoceran herbivores.

Several previous studies have reported negative effects of PhAC mixtures on freshwater

organisms, but these effects were typically observed at concentrations much higher than those

used in our study. A laboratory study showed that a mixture of 10 PhACs (antidepressants,

antihistaminics, antibiotics, anticonvulsants and analgesics) can be toxic to algae

(*Pseudokirchneriella subcapitata*), daphniids (*Ceriodaphnia dubia*) and zebrafish (*Danio*

rerio) at concentrations much higher (up to 15,000 times higher for the zebrafish) than the

environmental concentrations of the individual compounds (Watanabe et al., 2016). In outdoor

microcosms, high concentrations of a PhAC mixture (analgesics, antidepressants and

antibiotics; >60 mg.L^{-1} for each compound) resulted in elevated fish mortality and a decline in

zooplankton and phytoplankton diversity (Richards et al., 2004). However, David et al. (2020)

did not observe strong effects of a mixture of environmentally relevant concentrations of five

PhACs (antiepileptics, analgesics, and antihypertensives) on three-spined stickleback

populations in mesocosms.

In the field, an antibiotic mixture increased mortality of biofilm and decreased

extracellular leucine-aminopeptidase and alkaline phosphatase, resulting in significant

structural and functional changes in microbial attached communities (Proia et al., 2013), while

algal biomass and biofilm respiration were suppressed by 18% and 40%, respectively, and

photosynthesis by 88% in a mixed PhAC treatment (Rosi-Marshall et al., 2013). The effects on

phytoplankton observed in the summer experiment corroborate the results of these two studies.

4.3. Interactions between warming and PhACs

The interactions between temperature and contaminants pose a major challenge for ecological

risk assessment, especially with global warming (Zhang et al., 2019). The effects of warming

on contaminant toxicity have been extensively studied. However, the complex impacts of

warming and chemical stressors on shallow freshwater ecosystems can be influenced by

additional factors such as the type of contaminants and community composition, including the

number of trophic levels and their feeding behaviour. For example, heat waves in combination

with pesticide exposure can have antagonistic or synergistic effects, depending on the type of

pesticide used (Polazzo et al., 2022).

The combined effects of the PhAC mixture and warming in our study could not be

predicted from the individual effects of each stressor, especially in the summer experiment.

The PhAC mixture combined with warming strongly reduced phytoplankton biomass and its

control by primary consumers (zooplankton filter feeders) directly and indirectly due to

increased predation pressure from higher trophic levels. This trophic cascade also led to a

change in phytoplankton composition.

5. Conclusions and perspective

Although several studies of simplified food webs and communities have included both warming and chemical stressors (Knillmann et al., 2013; Wijewardene et al., 2021), current understanding of the joint effects of complex mixtures of PhACs and warming on freshwater communities is very limited. This hinders our ability to distinguish direct and indirect effects of each stressor and identify emergent phenomena that cannot be inferred from observations based on single stressors. Indeed, the current bias towards single species and population-level experiments has resulted in a knowledge gap for relevant community and ecosystem-level endpoints that prevents the exploration of important indirect effects of interacting multiple stressors that can compromise food web stability (Polazzo et al., 2022).

Our results contribute to filling these gaps and have implications for decision making to mitigate the effects of stressors on freshwater ecosystems. In our study, the PhAC mixture combined with warming strongly reduced phytoplankton biomass and its control by zooplankton. PhACs alone caused little or no impact at the population or community level, but when they were present in combination, they altered the aquatic community in the summer experiment, most likely when high temperatures increased close to or above the upper limits of the thermal niche of many taxa from phytoplankton to the insect predators. Moreover, we found that the effects of warming and PhACs differed strongly between seasons. Future research can extend our study by considering larger spatio-temporal scales and increasing the resolution and scope of the food web interactions. This would help us predict long-term effects of the current levels of anthropogenic stressors on freshwater communities and ecosystem functioning.

6. Acknowledgements

This study was supported by the Czech Science Foundation (project number GA20-16111S).

562   We thank Vladimíra Dekanová, Kateřina Bláhová, Samuel Dijoux, Fanny, Marius Wieczek,

563   Fanny Verfaillie and Markus Haber for assistance during mesocosm sampling.

7. References

Abrams, P.A., 1996. Dynamics and Interactions in Food Webs with Adaptive Foragers, in:

Polis, G.A., Winemiller, K.O. (Eds.), Food Webs: Integration of Patterns & Dynamics.

Springer US, Boston, MA, pp. 113–121. https://doi.org/10.1007/978-1-4615-7007-3_11

Amoros, C., 1984. Introduction pratique à la systématique des organismes des eaux

continentales françaises - 5. Crustacés Cladocères. Publ. Société Linn. Lyon 53, 72–107.

https://doi.org/10.3406/linly.1984.10627

Arnold, D., Hill, I.R., Matthiessen, P., Stephenson, R., 1991. Guidance document on testing

procedures for pesticides in freshwater mesocosms. SETAC Eur. Bruss. Belg.

Backhaus, T., 2014. Medicines, shaken and stirred: a critical review on the ecotoxicology of

pharmaceutical mixtures. Philos. Trans. R. Soc. B Biol. Sci. 369, 20130585.

https://doi.org/10.1098/rstb.2013.0585

Baert, J.M., Janssen, C.R., Sabbe, K., De Laender, F., 2016. Per capita interactions and stress

tolerance drive stress-induced changes in biodiversity effects on ecosystem functions. Nat.

Commun. 7, 12486. https://doi.org/10.1038/ncomms12486

Balogová, M., Gvoždík, L., 2015. Can Newts Cope with the Heat? Disparate Thermoregulatory

Strategies of Two Sympatric Species in Water. PLOS ONE 10, e0128155.

https://doi.org/10.1371/journal.pone.0128155

Belden, J.B., Gilliom, R.J., Lydy, M.J., 2007. How well can we predict the toxicity of pesticide

mixtures to aquatic life? Integr. Environ. Assess. Manag. 3, 364–372.

https://doi.org/10.1002/ieam.5630030307

Bennett, A.F., Lenski, R.E., 2007. An experimental test of evolutionary trade-offs during

temperature   adaptation.   Proc.   Natl.   Acad.   Sci.   104,   8649–8654.

https://doi.org/10.1073/pnas.0702117104

Bláha, M., Grabicova, K., Shaliutina, O., Kubec, J., Randák, T., Zlabek, V., Buřič, M., Veselý,

L., 2019. Foraging behaviour of top predators mediated by pollution of psychoactive

pharmaceuticals and effects on ecosystem stability. Sci. Total Environ. 662, 655–661.

https://doi.org/10.1016/j.scitotenv.2019.01.295

Boukal, D.S., Bideault, A., Carreira, B.M., Sentis, A., 2019. Species interactions under climate

change: connecting kinetic effects of temperature on individuals to community dynamics. Curr.

Opin.   Insect   Sci.,   Global   change   biology • Molecular   Physiology   35,   88–95.

https://doi.org/10.1016/j.cois.2019.06.014

Braak, C.J.F. ter, Smilauer, P., 2012. Canoco reference manual and user's guide: software for

ordination, version 5.0.

Brodin, T., Piovano, S., Fick, J., Klaminder, J., Heynen, M., Jonsson, M., 2014. Ecological

effects of pharmaceuticals in aquatic systems—impacts through behavioural alterations.

Philos. Trans. R. Soc. B Biol. Sci. 369, 20130580. https://doi.org/10.1098/rstb.2013.0580

Brogan, W.R., III, Relyea, R.A., 2015. Submerged Macrophytes Mitigate Direct and Indirect

Insecticide   Effects   in   Freshwater   Communities.   PLOS   ONE   10,   e0126677.

https://doi.org/10.1371/journal.pone.0126677

Cadmus, P., Pomeranz, J.P.F., Kraus, J.M., 2016. Low-cost floating emergence net and bottle

trap: comparison   of   two   designs.   J.   Freshw.   Ecol.   31,   653–658.

https://doi.org/10.1080/02705060.2016.1217944

David, V., Joachim, S., Catteau, A., Nott, K., Ronkart, S., Robert, C., Gillard, N., Bado-Nilles,

A., Chadili, E., Palluel, O., Turies, C., Julian, N., Castiglione, J., Dedourge-Geffard, O., Hani,

Y., Geffard, A., Porcher, J.-M., Beaudouin, R., 2020. Effects of chronic exposure to a

pharmaceutical mixture on the three-spined stickleback (gasterosteus aculeatus) population dynamics in lotic mesocosms. Aquat. Toxicol. 224, 105499. https://doi.org/10.1016/j.aquatox.2020.105499

Du, B., Haddad, S.P., Luek, A., Scott, W.C., Saari, G.N., Kristofco, L.A., Connors, K.A., Rash, C., Rasmussen, J.B., Chambliss, C.K., Brooks, B.W., 2014. Bioaccumulation and trophic dilution of human pharmaceuticals across trophic positions of an effluent-dependent wadeable stream. Philos. Trans. R. Soc. B Biol. Sci. 369, 20140058. https://doi.org/10.1098/rstb.2014.0058

Duchet, C., Moraru, G.M., Spencer, M., Saurav, K., Bertrand, C., Fayolle, S., Gershberg Hayoon, A., Shapir, R., Steindler, L., Blaustein, L., 2018. Pesticide-mediated trophic cascade and an ecological trap for mosquitoes. Ecosphere 9, e02179. https://doi.org/10.1002/ecs2.2179

Dziuba, M.K., Herdegen-Radwan, M., Pluta, E., Wejnerowski, Ł., Szczuciński, W., Cerbin, S., 2020. Temperature increase altered Daphnia community structure in artificially heated lakes: a potential scenario for a warmer future. Sci. Rep. 10, 13956. https://doi.org/10.1038/s41598-020-70294-6

Fedorova, G., Grabic, R., Grabicová, K., Turek, J., Van Nguyen, T., Randak, T., Brooks, B.W., Zlabek, V., 2022. Water reuse for aquaculture: Comparative removal efficacy and aquatic hazard reduction of pharmaceuticals by a pond treatment system during a one year study. J. Hazard. Mater. 421, 126712. https://doi.org/10.1016/j.jhazmat.2021.126712

Fekadu, S., Alemayehu, E., Dewil, R., Van der Bruggen, B., 2019. Pharmaceuticals in freshwater aquatic environments: A comparison of the African and European challenge. Sci. Total Environ. 654, 324–337. https://doi.org/10.1016/j.scitotenv.2018.11.072

Fent, K., Weston, A.A., Caminada, D., 2006. Ecotoxicology of human pharmaceuticals. Aquat. Toxicol. 76, 122–159. https://doi.org/10.1016/j.aquatox.2005.09.009

Fursdon, J.B., Martin, J.M., Bertram, M.G., Lehtonen, T.K., Wong, B.B.M., 2019. The

pharmaceutical pollutant fluoxetine alters reproductive behaviour in a fish independent of

predation risk. Sci. Total Environ. 650, 642–652.

https://doi.org/10.1016/j.scitotenv.2018.09.046

Grabicova, K., Grabic, R., Blaha, M., Kumar, V., Cerveny, D., Fedorova, G., Randak, T., 2015.

Presence of pharmaceuticals in benthic fauna living in a small stream affected by effluent from

a municipal sewage treatment plant. Water Res. 72, 145–153.

https://doi.org/10.1016/j.watres.2014.09.018

Guedes, R.N.C., Smagghe, G., Stark, J.D., Desneux, N., 2016. Pesticide-Induced Stress in

Arthropod Pests for Optimized Integrated Pest Management Programs. Annu. Rev. Entomol.

61, 43–62. https://doi.org/10.1146/annurev-ento-010715-023646

Guiry, M.D., Guiry, G.M., 2023. AlgaeBase. World-wide electronic publication, National

University of Ireland, Galway [WWW Document]. algaebase.org. URL

https://www.algaebase.org/about/ (accessed 3.16.23).

Hallmann, C.A., Sorg, M., Jongejans, E., Siepel, H., Hofland, N., Schwan, H., Stenmans, W.,

Müller, A., Sumser, H., Hörren, T., Goulson, D., de Kroon, H., 2017. More than 75 percent

decline over 27 years in total flying insect biomass in protected areas. PLOS ONE 12,

e0185809. https://doi.org/10.1371/journal.pone.0185809

Hansson, L.-A., Nicolle, A., Granéli, W., Hallgren, P., Kritzberg, E., Persson, A., Björk, J.,

Nilsson, P.A., Brönmark, C., 2013. Food-chain length alters community responses to global

change in aquatic systems. Nat. Clim. Change 3, 228–233.

https://doi.org/10.1038/nclimate1689

Heye, K., Lotz, T., Wick, A., Oehlmann, J., 2019. Interactive effects of biotic and abiotic

environmental stressors on carbamazepine toxicity in the non-biting midge Chironomus

riparius. Water Res. 156, 92–101. https://doi.org/10.1016/j.watres.2019.03.007

Johannsen, O.A., 1937. Aquatic diptera. Cornell University.

Joint, I., Henriksen, P., Fonnes, G.A., Bourne, D., Thingstad, T.F., Riemann, B., 2002.

Competition for inorganic nutrients between phytoplankton and bacterioplankton in nutrient

manipulated mesocosms. Aquat. Microb. Ecol. 29, 145–159.

https://doi.org/10.3354/ame029145

Kaštovský, J., Hauer, T., Geriš, R., Hutňan Chattová, B., Juráň, J., Lepšová-Skácelová, O.,

Pitelková, P., Pusztai, M., Škaloud, P., Šťastný, J., Čapková, K., Bohunická, M., Mühlsteinová,

R., 2018a. Atlas sinic a řas ČR 1. powerprint.

Kaštovský, J., Hauer, T., Geriš, R., Hutňan Chattová, B., Juráň, J., Lepšová-Skácelová, O.,

Pitelková, P., Pusztai, M., Škaloud, P., Šťastný, J., Čapková, K., Bohunická, M., Mühlsteinová,

R., 2018b. Atlas sinic a řas ČR 2. powerprint.

Kazanjian, G., Velthuis, M., Aben, R., Stephan, S., Peeters, E.T.H.M., Frenken, T., Touwen,

J., Xue, F., Kosten, S., Van de Waal, D.B., de Senerpont Domis, L.N., van Donk, E., Hilt, S.,

2018. Impacts of warming on top-down and bottom-up controls of periphyton production. Sci.

Rep. 8, 9901. https://doi.org/10.1038/s41598-018-26348-x

Kneitel, J.M., Chase, J.M., 2004. Trade-offs in community ecology: linking spatial scales and

species coexistence. Ecol. Lett. 7, 69–80. https://doi.org/10.1046/j.1461-0248.2003.00551.x

Knillmann, S., Stampfli, N.C., Noskov, Y.A., Beketov, M.A., Liess, M., 2013. Elevated

temperature prolongs long-term effects of a pesticide on Daphnia spp. due to altered

competition in zooplankton communities. Glob. Change Biol. 19, 1598–1609.

https://doi.org/10.1111/gcb.12151

Koba, O., Grabicova, K., Cerveny, D., Turek, J., Kolarova, J., Randak, T., Zlabek, V., Grabic,

R., 2018. Transport of pharmaceuticals and their metabolites between water and sediments as

a further potential exposure for aquatic organisms. J. Hazard. Mater. 342, 401–407.

https://doi.org/10.1016/j.jhazmat.2017.08.039

Lagesson, A., Fahlman, J., Brodin, T., Fick, J., Jonsson, M., Byström, P., Klaminder, J., 2016.

Bioaccumulation of five pharmaceuticals at multiple trophic levels in an aquatic food web -

Insights from a field experiment. Sci. Total Environ. 568, 208–215.

https://doi.org/10.1016/j.scitotenv.2016.05.206

Lefcheck, J.S., 2016. piecewiseSEM: Piecewise structural equation modelling in r for ecology,

evolution, and systematics. Methods Ecol. Evol. 7, 573–579. https://doi.org/10.1111/2041-

210X.12512

Loos, R., Carvalho, R., António, D.C., Comero, S., Locoro, G., Tavazzi, S., Paracchini, B.,

Ghiani, M., Lettieri, T., Blaha, L., Jarosova, B., Voorspoels, S., Servaes, K., Haglund, P., Fick,

J., Lindberg, R.H., Schwesig, D., Gawlik, B.M., 2013. EU-wide monitoring survey on

emerging polar organic contaminants in wastewater treatment plant effluents. Water Res. 47,

6475–6487. https://doi.org/10.1016/j.watres.2013.08.024

Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S.L., Péan, C., Berger, S., Caud, N., Chen,

Y., Goldfarb, L., Gomis, M.I., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J.B.R.,

Maycock, T.K., Waterfield, T., Yelekçi, Ö., Yu, R., Zhou, B. (Eds.), 2021. Climate Change

2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment

Report of the Intergovernmental Panel on Climate Change. Cambridge University Press,

Cambridge, United Kingdom and New York, NY, USA.

https://doi.org/10.1017/9781009157896

Mooij, W.M., De Senerpont Domis, L.N., Hülsmann, S., 2008. The impact of climate warming

on water temperature, timing of hatching and young-of-the-year growth of fish in shallow lakes

in the Netherlands. J. Sea Res., Dynamics of Fish and Fishers 60, 32–43.

https://doi.org/10.1016/j.seares.2008.03.002

Morris, O.F., Loewen, C.J.G., Woodward, G., Schäfer, R.B., Piggott, J.J., Vinebrooke, R.D.,

Jackson, M.C., 2022. Local stressors mask the effects of warming in freshwater ecosystems.

Ecol. Lett. 25, 2540–2551. https://doi.org/10.1111/ele.14108

Noyes, P.D., McElwee, M.K., Miller, H.D., Clark, B.W., Van Tiem, L.A., Walcott, K.C.,

Erwin, K.N., Levin, E.D., 2009. The toxicology of climate change: Environmental

contaminants in a warming world. Environ. Int. 35, 971–986.

https://doi.org/10.1016/j.envint.2009.02.006

Op de Beeck, L., Verheyen, J., Olsen, K., Stoks, R., 2017. Negative effects of pesticides under

global warming can be counteracted by a higher degradation rate and thermal adaptation. J.

Appl. Ecol. 54, 1847–1855. https://doi.org/10.1111/1365-2664.12919

Parmesan, C., 2006. Ecological and Evolutionary Responses to Recent Climate Change. Annu.

Rev. Ecol. Evol. Syst. 37, 637–669.

https://doi.org/10.1146/annurev.ecolsys.37.091305.110100

Parmesan, C., Yohe, G., 2003. A globally coherent fingerprint of climate change impacts across

natural systems. Nature 421, 37–42. https://doi.org/10.1038/nature01286

Polazzo, F., Roth, S.K., Hermann, M., Mangold-Döring, A., Rico, A., Sobek, A., Van den

Brink, P.J., Jackson, M.C., 2022. Combined effects of heatwaves and micropollutants on

freshwater ecosystems: Towards an integrated assessment of extreme events in multiple

stressors research. Glob. Change Biol. 28, 1248–1267. https://doi.org/10.1111/gcb.15971

Post, E., 2013. Ecology of Climate Change: The Importance of Biotic Interactions, Ecology of

Climate Change. Princeton University Press. https://doi.org/10.1515/9781400846139

Proia, L., Lupini, G., Osorio, V., Pérez, S., Barceló, D., Schwartz, T., Amalfitano, S., Fazi, S.,

Romaní, A.M., Sabater, S., 2013. Response of biofilm bacterial communities to antibiotic

pollutants in a Mediterranean river. Chemosphere 92, 1126–1135.

https://doi.org/10.1016/j.chemosphere.2013.01.063

Relyea, R.A., 2009. A cocktail of contaminants: how mixtures of pesticides at low

concentrations affect aquatic communities. Oecologia 159, 363–376.

https://doi.org/10.1007/s00442-008-1213-9

Richards, S.M., Wilson, C.J., Johnson, D.J., Castle, D.M., Lam, M., Mabury, S.A., Sibley,

P.K., Solomon, K.R., 2004. EFFECTS OF PHARMACEUTICAL MIXTURES IN AQUATIC

MICROCOSMS. Environ. Toxicol. Chem. 23, 1035. https://doi.org/10.1897/02-616

Rodrigues, A.C.M., Machado, A.L., Bordalo, M.D., Saro, L., Simão, F.C.P., Rocha, R.J.M.,

Golovko, O., Žlábek, V., Barata, C., Soares, A.M.V.M., Pestana, J.L.T., 2018. Invasive Species

Mediate Insecticide Effects on Community and Ecosystem Functioning. Environ. Sci. Technol.

52, 4889–4900. https://doi.org/10.1021/acs.est.8b00193

Rosi-Marshall, E.J., Kincaid, D.W., Bechtold, H.A., Royer, T.V., Rojas, M., Kelly, J.J., 2013.

Pharmaceuticals suppress algal growth and microbial respiration and alter bacterial

communities in stream biofilms. Ecol. Appl. 23, 583–593. https://doi.org/10.1890/12-0491.1

RStudio, T., 2020. RStudio: integrated development for R. Rstudio Team PBC Boston MA

URL Httpwww Rstudio Com.

Saaristo, M., Lagesson, A., Bertram, M.G., Fick, J., Klaminder, J., Johnstone, C.P., Wong,

B.B.M., Brodin, T., 2019. Behavioural effects of psychoactive pharmaceutical exposure on

European perch (Perca fluviatilis) in a multi-stressor environment. Sci. Total Environ. 655,

1311–1320. https://doi.org/10.1016/j.scitotenv.2018.11.228

Sala, O.E., Stuart Chapin, F., Iii, Armesto, J.J., Berlow, E., Bloomfield, J., Dirzo, R., Huber-

Sanwald, E., Huenneke, L.F., Jackson, R.B., Kinzig, A., Leemans, R., Lodge, D.M., Mooney,

H.A., Oesterheld, M., Poff, N.L., Sykes, M.T., Walker, B.H., Walker, M., Wall, D.H., 2000.

Global Biodiversity Scenarios for the Year 2100. Science 287, 1770–1774.

https://doi.org/10.1126/science.287.5459.1770

Sánchez-Bayo, F., 2021. Indirect Effect of Pesticides on Insects and Other Arthropods. Toxics

9, 177. https://doi.org/10.3390/toxics9080177

Sánchez-Bayo, F., Wyckhuys, K.A.G., 2019. Worldwide decline of the entomofauna: A review

of its drivers. Biol. Conserv. 232, 8–27. https://doi.org/10.1016/j.biocon.2019.01.020

Schmidt, T.S., Miller, J.L., Mahler, B.J., Van Metre, P.C., Nowell, L.H., Sandstrom, M.W.,

Carlisle, D.M., Moran, P.W., Bradley, P.M., 2022. Ecological consequences of neonicotinoid

mixtures in streams. Sci. Adv. 8, eabj8182. https://doi.org/10.1126/sciadv.abj8182

Shipley, B., Douma, J.C., 2020. Generalized AIC and chi-squared statistics for path models

consistent with directed acyclic graphs. Ecology 101, e02960. https://doi.org/10.1002/ecy.2960

Spaak, J.W., Baert, J.M., Baird, D.J., Eisenhauer, N., Maltby, L., Pomati, F., Radchuk, V.,

Rohr, J.R., Van den Brink, P.J., De Laender, F., 2017. Shifts of community composition and

population density substantially affect ecosystem function despite invariant richness. Ecol.

Lett. 20, 1315–1324. https://doi.org/10.1111/ele.12828

Stephenson, G.L., Hamilton, P., Kaushik, N.K., Robinson, J.B., Solomon, K.R., 1984. Spatial

distribution of plankton in enclosures of three sizes. Can. J. Fish. Aquat. Sci. 41, 1048–1054.

Stoks, R., Verheyen, J., Van Dievel, M., Tüzün, N., 2017. Daily temperature variation and

extreme high temperatures drive performance and biotic interactions in a warming world. Curr.

Opin. Insect Sci., Global change biology * Molecular physiology 23, 35–42.

https://doi.org/10.1016/j.cois.2017.06.008

Švecová, H., Grabic, R., Grabicová, K., Vojs Staňová, A., Fedorova, G., Cerveny, D., Turek,

J., Randák, T., Brooks, B.W., 2021. De facto reuse at the watershed scale: Seasonal changes,

population contributions, instream flows and water quality hazards of human pharmaceuticals.

Environ. Pollut. 268, 115888. https://doi.org/10.1016/j.envpol.2020.115888

Team, R.C., 2018. R: A language and environment for statistical computing. 2014. R Found.

Stat. Comput. Vienna Austria.

Urrutia-Cordero, P., Ekvall, M.K., Ratcovich, J., Soares, M., Wilken, S., Zhang, H., Hansson,

L.-A., 2017. Phytoplankton diversity loss along a gradient of future warming and

brownification in freshwater mesocosms. Freshw. Biol. 62, 1869–1878.

https://doi.org/10.1111/fwb.13027

Van den Brink, P.J., Braak, C.J.F.T., 1999. Principal response curves: Analysis of time-

dependent multivariate responses of biological community to stress. Environ. Toxicol. Chem.

18, 138–148. https://doi.org/10.1002/etc.5620180207

Velthuis, M., de Senerpont Domis, L.N., Frenken, T., Stephan, S., Kazanjian, G., Aben, R.,

Hilt, S., Kosten, S., van Donk, E., Van de Waal, D.B., 2017. Warming advances top-down

control and reduces producer biomass in a freshwater plankton community. Ecosphere 8,

e01651. https://doi.org/10.1002/ecs2.1651

Venâncio, C., Ribeiro, R., Soares, A., Lopes, I., 2016. Multiple Stressor Differential

Tolerances: Possible Implications at the Population Level. PLOS ONE 11, e0151847.

https://doi.org/10.1371/journal.pone.0151847

Vörösmarty, C.J., McIntyre, P.B., Gessner, M.O., Dudgeon, D., Prusevich, A., Green, P.,

Glidden, S., Bunn, S.E., Sullivan, C.A., Liermann, C.R., Davies, P.M., 2010. Global threats to

human water security and river biodiversity. Nature 467, 555–561.

https://doi.org/10.1038/nature09440

Watanabe, H., Tamura, I., Abe, R., Takanobu, H., Nakamura, A., Suzuki, T., Hirose, A.,

Nishimura, T., Tatarazako, N., 2016. Chronic toxicity of an environmentally relevant mixture

of pharmaceuticals to three aquatic organisms (alga, daphnid, and fish). Environ. Toxicol.

Chem. 35, 996–1006. https://doi.org/10.1002/etc.3285

Wijewardene, L., Wu, N., Qu, Y., Guo, K., Messyasz, B., Lorenz, S., Riis, T., Ulrich, U.,

Fohrer, N., 2021. Influences of pesticides, nutrients, and local environmental variables on

phytoplankton communities in lentic small water bodies in a German lowland agricultural area.

Sci. Total Environ. 780, 146481. https://doi.org/10.1016/j.scitotenv.2021.146481

Wilkinson, J.L., Boxall, A.B.A., Kolpin, D.W., Leung, K.M.Y., Lai, R.W.S., Galbán-Malagón,

C., Adell, A.D., Mondon, J., Metian, M., Marchant, R.A., Bouzas-Monroy, A., Cuni-Sanchez,

A., Coors, A., Carriquiriborde, P., Rojo, M., Gordon, C., Cara, M., Moermond, M., Luarte, T.,

Petrosyan, V., Perikhanyan, Y., Mahon, C.S., McGurk, C.J., Hofmann, T., Kormoker, T.,

Iniguez, V., Guzman-Otazo, J., Tavares, J.L., Gildasio De Figueiredo, F., Razzolini, M.T.P.,

Dougnon, V., Gbaguidi, G., Traoré, O., Blais, J.M., Kimpe, L.E., Wong, M., Wong, D.,

Ntchantcho, R., Pizarro, J., Ying, G.-G., Chen, C.-E., Páez, M., Martínez-Lara, J., Otamonga,

J.-P., Poté, J., Ifo, S.A., Wilson, P., Echeverría-Sáenz, S., Udikovic-Kolic, N., Milakovic, M.,

Fatta-Kassinos, D., Ioannou-Ttofa, L., Belušová, V., Vymazal, J., Cárdenas-Bustamante, M.,

Kassa, B.A., Garric, J., Chaumot, A., Gibba, P., Kunchulia, I., Seidensticker, S., Lyberatos, G.,

Halldórsson, H.P., Melling, M., Shashidhar, T., Lamba, M., Nastiti, A., Supriatin, A., Pourang,

N., Abedini, A., Abdullah, O., Gharbia, S.S., Pilla, F., Chefetz, B., Topaz, T., Yao, K.M.,

Aubakirova, B., Beisenova, R., Olaka, L., Mulu, J.K., Chatanga, P., Ntuli, V., Blama, N.T.,

Sherif, S., Aris, A.Z., Looi, L.J., Niang, M., Traore, S.T., Oldenkamp, R., Ogunbanwo, O.,

Ashfaq, M., Iqbal, M., Abdeen, Z., O'Dea, A., Morales-Saldaña, J.M., Custodio, M., de la

Cruz, H., Navarrete, I., Carvalho, F., Gogra, A.B., Koroma, B.M., Cerkvenik-Flajs, V.,

Gombač, M., Thwala, M., Choi, K., Kang, H., Ladu, J.L.C., Rico, A., Amerasinghe, P., Sobek,

A., Horlitz, G., Zenker, A.K., King, A.C., Jiang, J.-J., Kariuki, R., Tumbo, M., Tezel, U., Onay,

T.T., Lejju, J.B., Vystavna, Y., Vergeles, Y., Heinzen, H., Pérez-Parada, A., Sims, D.B., Figy,

M., Good, D., Teta, C., 2022. Pharmaceutical pollution of the world's rivers. Proc. Natl. Acad.

Sci. 119, e2113947119. https://doi.org/10.1073/pnas.2113947119

Winder, M., Hunter, D.A., 2008. Temporal organization of phytoplankton communities linked

to physical forcing. Oecologia 156, 179–192. https://doi.org/10.1007/s00442-008-0964-7

Woodward, G., Perkins, D.M., Brown, L.E., 2010. Climate change and freshwater ecosystems:

impacts across multiple levels of organization. Philos. Trans. R. Soc. B Biol. Sci. 365, 2093–

2106. https://doi.org/10.1098/rstb.2010.0055

Zenker, A., Cicero, M.R., Prestinaci, F., Bottoni, P., Carere, M., 2014. Bioaccumulation and

biomagnification potential of pharmaceuticals with a focus to the aquatic environment. J.

Environ. Manage. 133, 378–387. https://doi.org/10.1016/j.jenvman.2013.12.017

Zhang, H., Huo, S., Yeager, K.M., He, Z., Xi, B., Li, X., Ma, C., Wu, F., 2019. Phytoplankton

response to climate changes and anthropogenic activities recorded by sedimentary pigments in

a shallow eutrophied lake. Sci. Total Environ. 647, 1398–1409.

https://doi.org/10.1016/j.scitotenv.2018.08.081

841