Combined effects of climate warming and pharmaceuticals on a tri-trophic freshwater food web

Claire Duchet[1, 2✉], Kateřina Grabicová[3], Vojtech Kolar[1,2], Olga Lepšová[4], Helena Švecová[3], Andras Csercsa[1], Barbora Zdvihalová[1], Tomáš Randák[3], David S. Boukal[1,2]

[1] Department of Ecosystem Biology, Faculty of Science, University of South Bohemia, Branišovská 1760, 37005 České Budějovice, Czech Republic

[2] Institute of Entomology, Biology Centre of the Czech Academy of Sciences, Branišovská 31, 37005 České Budějovice, Czech Republic

[3] South Bohemian Research Center of Aquaculture and Biodiversity of Hydrocenoses, Faculty of Fisheries and Protection of Waters, University of South Bohemia in České Budějovice, Zátiší 728/II, CZ-389 25 Vodňany, Czech Republic

[4] Department of Botany, Faculty of Science, University of South Bohemia, Branišovská 1760, 37005 České Budějovice, Czech Republic

[✉]Corresponding author: Claire Duchet

E-mail address: cduchet@prf.jcu.cz

Department of Ecosystem Biology, Faculty of Science, University of South Bohemia

Branišovská 1760, 37005 České Budějovice, Czech Republic

Tel. +420 387 772 244 (office); Fax: +420 387 772 368

Abstract

Multiple anthropogenic stressors influence the functioning of ponds and lakes, but their combined effects are often little understood. We ran two mesocosm experiments to evaluate the effects of warming (+4°C above ambient) and environmentally relevant concentrations of a mixture of commonly used pharmaceuticals, an emerging class of chemical contaminants, on tri-trophic food webs representative of pelagic communities in ponds and other small standing waters. We quantified the main and interactive effects of warming and pharmaceuticals on each trophic level and attributed them to the direct effects of both stressors and the indirect effects arising through biotic interactions. Warming and pharmaceuticals had stronger effects in the summer experiment, altering zooplankton community composition and causing delayed or accelerated emergence of top insect predators. In summer, both stressors and top predators reduced filter-feeding zooplankton biomass, while warming and pharmaceuticals had opposing effects on phytoplankton. In the winter experiment, the effects were much weaker and primarily limited to a positive effect of warming on phytoplankton biomass. Overall, we show that pharmaceuticals can exacerbate the effects of climate warming in freshwater ecosystems, especially during the warm season. Our results demonstrate the utility of community-level studies across different seasons for the risk assessment of multiple emerging stressors in freshwater ecosystems.

Keywords: warming, pharmaceutical mixture, aquatic invertebrates, community dynamics, mesocosm, multiple stressors, stressor interactions, zooplankton

Introduction

Freshwater ecosystems face severe impacts from multiple stressors, including habitat loss, chemical pollution, pathogens, invasive species, and climate change (IPBES, 2019). Among these stressors, climate warming and chemical pollution pose major threats to freshwater biodiversity and ecosystem functioning (Vörösmarty et al., 2010). Climate change is predicted to increase average global annual temperatures by up to +4.4°C by 2100, with stronger warming at high latitudes and during cold seasons (Masson-Delmotte et al., 2021). Freshwater ecosystems, dominated by ectotherms, are highly susceptible to warming (Parmesan and Yohe, 2003; Woodward et al., 2010) that can lead to reduced body sizes, altered species interactions, and shifts in species distribution ranges and phenology, with cascading impacts on ecosystem functioning (Boukal et al., 2019; Post, 2013).

Among chemicals, pharmaceuticals represent a prominent class of emerging pollutants (Fekadu et al., 2019). Their widespread use and inadequate removal in wastewater treatment plants contribute to global contamination of aquatic systems (Wilkinson et al., 2022). At environmentally relevant concentrations (ng.L^{-1} to low µg.L^{-1}), pharmaceuticals can disrupt the behavior (Brodin et al., 2014; Saaristo et al., 2019) and reproduction of aquatic ectotherms (Fursdon et al., 2019), affecting individual fitness and triggering indirect effects like altered trophic interactions (Fursdon et al., 2019; Bláha et al., 2019). Although pharmaceuticals can bioaccumulate in benthic macroinvertebrates (Grabicova et al., 2015), leading to exposure of higher trophic levels (Lagesson et al., 2016) like fish (Brodin et al., 2014; Du et al., 2014; Grabicova et al., 2017), they are usually diluted through the trophic level and rarely biomagnified (Grabicova et al., 2015; Xie et al., 2017).

The effects of pharmaceuticals on aquatic systems have primarily been studied using single-species laboratory bioassays, often at higher concentrations than those found in the

environment (Richards et al., 2004). Furthermore, pharmaceuticals typically occur as

complex mixtures in the environment (Grabicova et al., 2015; Loos et al., 2013).

Understanding their ecotoxicology under environmentally realistic conditions and

quantitative consequences on the biota therefore requires new data (Backhaus, 2014).

Moreover, the combined effects of pharmaceuticals and warming on freshwater biota and

ecosystem functioning remain largely unknown (Heye et al., 2019) despite extensive research

on their individual effects. While the dominant stressor alone often explains the main impacts

of multiple stressors on a freshwater ecosystem (the so-called dominance null model; Morris

et al., 2022), interactions between stressors can result in synergistic, antagonistic, or additive

effects, making predictions challenging (Spaak et al., 2017).

Warming can modulate the effects of pollutants on freshwater ecosystems both directly

and indirectly. While many pollutants become more toxic at higher temperatures, warming

may also accelerate their degradation, thereby reducing the exposure of biota to them (Noyes

et al., 2009; Op de Beeck et al., 2017). Population-level responses to the combined stressors

can vary dramatically due to differences in environmental sensitivity (Venâncio et al., 2016),

tolerance of individuals (Baert et al., 2016), ecological trade-offs and local adaptation

patterns (Bennett and Lenski, 2007; Kneitel and Chase, 2004). Additionally, the fate of

species embedded in food webs depends on direct and indirect changes in intraspecific and

interspecific interactions (Post, 2013; Relyea, 2009), including trait- and density-mediated

indirect effects such as top-down and bottom-up effects (Brogan and Relyea, 2015; Guedes et

al., 2016; Rodrigues et al., 2018). Finally, persistent pollutants can alter community structure

and affect or enhance various aspects of ecosystem functioning (Duchet et al., 2018;

Sánchez-Bayo, 2021).

To address the knowledge gap on the combined effects of pharmaceuticals and

warming on freshwater communities, we conducted a full-factorial experiment using outdoor

mesocosms. By simulating realistic conditions, we aimed to provide insights into the

ecological consequences of these stressors in a changing environment. We included key

functional groups representative of the communities in pond and shallow lake ecosystems and

ran the experiment during both summer and winter to account for seasonal turnover in the

communities. Our pelagic communities included three main trophic levels and we therefore

expected (1) a strong negative effect of warming on zooplankton and none or a positive effect

on the top predators (Hansson et al., 2013), leading to an increase in phytoplankton biomass

(Belden et al., 2007). We also expected that (2) warming will have stronger negative effects

during summer and milder, more positive effects during winter, especially on phytoplankton

and zooplankton growth.

2. Material and Methods

2.1. Mesocosms

The mesocosm facility at the University of South Bohemia (České Budějovice, Czech

Republic; 48°58'36.6" N 14°26'41.8" E, 381 m a.s.l.) comprises 32 circular outdoor

mesocosms (120 cm inner diameter, 1.13 m3 water volume each) with a custom-built heating

system (Hennlich®, Schädling, Austria; Fig. 1). The 28 mesocosms used in this study were

filled 6 weeks before the experiment with tap water filtered through a granulated activated

carbon filter, enriched with 100 g of common alder leaves (*Alnus glutinosa*), and 8 g of fish

food (Pondstick granules, Apetit®) to add nutrients. Water in each mesocosm was gently

mixed with an air stone, except from 12 January to 3 February 2021, when mixing was

paused to allow natural freezing in the unheated mesocosms (Fig. S1).

The experimental community included common macroinvertebrate and plankton taxa

representing different trophic levels and feeding guilds (predators, filter feeders, grazers,

detritivores, and primary producers; Table S2). Phytoplankton and zooplankton from a

nearby fishpond were added to the mesocosms one month before the experiment, after

collecting water samples for xenobiotic analysis (see detailed methods in S3, Tables S3.2 and

S3.3). The plankton inoculum was mixed in a barrel and then added to the mesocosms (2.5-

litre aliquot each). Macroinvertebrates from nearby fishponds, ditches, and sand pits were

then sequentially introduced into the mesocosms after two weeks: benthic detritivores

(*Asellus aquaticus* - ca. 100 per mesocosm), grazers (*Cloeon* cf. *dipterum* mayfly larvae - ca.

200 per mesocosm), omnivorous scrapers (*Planorbarius corneus* pond snails - 9 per

mesocosm), omnivorous piercers (*Sigara falleni* water boatman - 50 per mesocosm), water

column predators (*Notonecta glauca* backswimmers - 9 individuals per mesocosm, *Aeshna* or

Anax dragonfly larvae - 6 per mesocosm; hereafter Aeshna and Anax), benthic predators

(libellulid dragonfly larvae - 3 per mesocosm), and phytophilous predators occupying

artificial vegetation and mesocosm walls (damselfly larvae - ca. 50 per mesocosm). The

predators were added to the mesocosms one week before the start.

Each mesocosm received three artificial submerged macrophytes made of four strips of

green plastic mesh (5×100 cm) attached to a granite stone to provide additional microhabitat.

Two polypropylene strips (6 cm wide, 50 cm long) hanging vertically in each mesocosm

were used to incubate pre-cultured periphyton from one of the fishponds where the

macroinvertebrates were collected. One coarse (20x20 cm; 20 mm mesh) and one fine (20x20

cm; 0.5 mm mesh) litter bag containing 9.5 g of dried alder (*Alnus glutinosa*) leaves were

added to each mesocosm to distinguish the contribution of detritivores and microbial

decomposers to the decomposition process (Fig. 1).

Figure 1. Schematic view of the mesocosm set-up

2.1. Experimental design

Seven mesocosms were assigned to each of the four treatments: (1) unheated without

pharmaceuticals, (2) unheated with pharmaceuticals, (3) heated without pharmaceuticals, and

(4) heated with pharmaceuticals. Each mesocosm received the same treatment in both

experiments. Heating was maintained at +4°C above ambient throughout each experiment

(mean ± SD differences in daily mean temperatures between treatments: winter experiment,

4.03 ± 0.04°C; summer experiment, 4.06 ± 0.06°C; see S1), consistent with the SSP5-8.5

projection model for 2081–2100 (Masson-Delmotte et al., 2021).

The pharmaceutical mixture included 15 compounds from four major drug categories

(cardiovascular: telmisartan, valsartan, metoprolol, atenolol; psychoactive: carbamazepine,

lamotrigine, venlafaxine, citalopram, tramadol; antihistaminic: cetirizine, fexofenadine;

antibiotic: sulfamethoxazole, trimethoprim, clarithromycin, clindamycin; Table S3.1) at

concentrations commonly detected in surface waters in the Czech Republic (Fedorova et al.,

2022; Švecová et al., 2021). The pharmaceutical mixture was administered as a single pulse

exposure (total concentration: 2890 ng.L^{-1}; see S3 for more details).

We conducted two experiments: the 'winter experiment' from 24 September 2020 to 12

March 2021 and the 'summer experiment' from 2 June to 2 August 2021 (Figures S4.1 and

S4.2). Day 0 marked the start of each experiment before adding the pharmaceuticals and

initiating warming. The initial three samplings for all parameters were performed on Day 0,

followed by Day 2 and Day 7. Subsequent samples were collected monthly in winter and

fortnightly in summer.

2.3. Sampling

This section covers the methods for assessing the temporal changes in the pelagic

environment and biota including predatory insects feeding on zooplankton. Complete

community data collected at the end of each experiment will be analyzed elsewhere. Table

S4.1 provides a summary of all response variables collected during the experiments.

In both experiments, we measured pharmaceutical concentrations, environmental

parameters (dissolved O_2, conductivity, pH, turbidity, water temperature, and chlorophyll-a

concentration) using a portable multiparameter water quality meter (YSI 556 MPS probe;

YSI®, USA). On each sampling day, one water sample (6 mL) was collected in each

mesocosm, filtered through a 0.2-µm syringe filter (regenerated cellulose) and stored at -

20°C until the pharmaceutical compound concentrations were analysed with an in-line solid-

phase extraction liquid chromatography with tandem mass spectrometry (in-line SPE/LC-

MS/MS, triple quadrupole mass spectrometer Quantiva, Thermo Fisher Scientific, USA; see

S3 for details).

Dissolved organic carbon (DOC), nitrogen, and phosphorus concentrations were

measured at Day 0 and at the end of each experiment. For technical reasons, total nitrogen

(TN) and total phosphorus (TP) were measured during the winter experiment, while dissolved

nitrogen (DN) and soluble reactive phosphorus (SRP) were measured during the summer

experiment. At the end of each experiment, phytoplankton samples were collected from the

surface water layer (ca. 10 cm) in 500 ml PET bottles and fixed with Lugol solution

immediately after sampling before identification and counting in the lab (see S4.1 for details).

During both experiments, we repeatedly sampled zooplankton with a tube sampler (1.2

m length, 7 cm inner diameter) equipped with a one-way bottom valve. On each sampling

date, three samples were taken from different locations within each mesocosm to minimize

plankton patchiness effects (Stephenson et al., 1984). The resulting bulk sample

(approximately 9 L depending on water level, ranging between 7.5–9.6 L in summer and 8.4–

10.5 L in winter) was collected in a 20 L bucket. From this, one litre was filtered through an

80-μm mesh plankton net. Retained organisms (zooplankton and some other pelagic

invertebrates) were preserved in 4% formaldehyde in 120 mL plastic vials. Counting and

identification were done under a stereomicroscope (Olympus SZX2-ILLT stereomicroscope;

Olympus, Tokyo, Japan) to the lowest feasible taxonomic level using identification keys

(Amoros, 1984; Johannsen, 1937).

During the summer experiment, we also monitored aquatic insect emergence with

pyramid-shaped floating emergence traps (35 × 35 cm base dimensions) with a 250 mL

collection bottle of soapy water (Cadmus et al., 2016). One trap was installed in each

mesocosm on Day 0. Emerging insects were collected every 2–3 days and preserved in 70%

ethanol (mayflies, chironomids) or frozen (odonates) for later identification and analysis of

pharmaceutical content.

Biomass of each taxon was estimated by multiplying total abundance (for aquatic

insects) or density (zooplankton) by the average body size based on literature data and our

measurements (Table S4.2).

2.3. Statistical analyses

We employed multiple methods (detailed in S4.2) to assess the effects of the

pharmaceutical mixture and warming on the aquatic community. In brief, redundancy

analysis (RDA) was used to examine temporal changes in environmental parameters and

chlorophyll-a concentrations. Principal component analyses (PCAs) were used to visualize

differences in algal groups and taxa between treatments. Principal response curve (PRC) was

used to analyse changes in zooplankton community structure. We then ran generalized linear

models (GLMs) to assess treatment effects on nutrients, predator emergence and survival and

generalized linear mixed models (GLMMs) to assess treatment effects on *Daphnia* and

copepod biomasses. Model selection approach with the corrected Akaike information

criterion was used to identify the most parsimonious model for each univariate response.

Finally, we used confirmatory path analysis (piecewise SEM) to understand the direct and

indirect effects of the pharmaceutical mixture and warming on the community. Hypothetical

pathways were based on *a priori* knowledge to build a complete model of cause-effect

relationships. Fisher's C statistics were used to assess the overall fit of each SEM model.

Multivariate analyses were performed using CANOCO 5 (ter Braak and Smilauer, 2012), and

univariate analyses were done in R version 4.1.2 (R Core Team, 2018).

3. Results

3.1. Pharmaceutical concentrations

During the summer experiment, total pharmaceutical concentration gradually decreased over

time to about 25% in heated mesocosms and 30% in non-heated mesocosms (Fig. 2A).

Meanwhile, metabolite concentrations increased during the first month and remained stable

thereafter (Fig. 2C). In the winter experiment, pharmaceuticals were degraded over time (Fig.

2B). Contrarily, metabolite concentrations in the winter experiment increased significantly

during the first two months, reaching double the summer concentrations, and remained stable

afterward (Fig. 2D). Pharmaceuticals and their metabolites were below detection limits in

samples from non-treated mesocosms in both experiments.

Figure 2. Total concentrations of pharmaceutical (PhAC) main compounds (A, C) and their

metabolites (B, D) in the winter (A, B) and summer (C, D) experiments in heated (orange) and non-

heated (blue) mesocosms treated with pharmaceuticals. Symbols with error bars = treatment-specific

mean values ± SD ($n = 7$); mean values in the same treatment connected by colour lines.

3.2. Environmental conditions

Environmental conditions significantly differed between treatments in both experiments

(summer RDA: pseudo-F = 4.9, p = 0.002, explained variation = 7.4%; winter RDA: pseudo-

F = 53.7, p = 0.001, explained variation = 43.1%). In the summer experiment, non-heated

mesocosms had higher water levels, higher turbidity, lower conductivity, and lower

chlorophyll-a concentrations (mainly those without pharmaceuticals) compared to heated

mesocosms (Fig. 3A, Table S5.1). In the winter experiment, treatments did not differ in

chlorophyll-a concentration, dissolved oxygen, turbidity, and pH, but non-heated mesocosms

had higher water levels and lower conductivity compared to the heated ones (Fig. 3B, Table

S5.2).

Figure 3. Ordination diagram (RDA) showing differences between the treatments in environmental

parameters and chlorophyll-a concentration during the summer (A) and winter (B) experiment.

Treatments: CLT = non-heated, without pharmaceuticals; CHT = heated, without pharmaceuticals;

TLT = non-heated, with pharmaceuticals; THT = heated, with pharmaceuticals.

3.3. Effect of the stressors on nutrients and phytoplankton

The most parsimonious models showed disparate effects of the stressors on nutrients in the

summer and winter experiments. In summer, heated mesocosms had lower final

concentrations of DOC, DN, and SRP compared to non-heated ones. The added

pharmaceutical mixture had a small negative effect on DOC and a significant negative effect

on DN and SRP (Fig. S6.1A, C, and E). In winter, heated mesocosms also had lower final

concentrations of DOC, TN, and TP compared to non-heated ones, but no effects of

pharmaceuticals on nutrients were detected (Fig. S6.1B, D, and F). No synergistic effects of

the two stressors on nutrients were detected in both experiments (Tables S6.1 and S6.2).

There were some differences in the composition of major algal groups between

treatments (PCA, Fig. S6.2), although nonsignificant (RDA, Table S6.3). In summer,

Cyanobacteria dominated in the heated mesocosms, while Chrysophyta were prevalent in the

non-heated mesocosms (Fig. S6.2A-B). In winter, Chrysophyta were more prevalent in the

non-heated controls, while Streptophyta were more prevalent in the non-heated, treated

treatment (Fig. S6.2C). Warming appeared to be the main driver of taxonomic composition in

the winter experiment (PCA F2, Fig. S6.2D). Notably, chlorophytes (*Oocystis* spp. and

Scenedesmus sp.) were more prevalent in the heated mesocosms, while bacillariophytes

(*Synedra* sp., *Gomphonema* spp., *Navicula* sp. and *Ulnaria* sp.) and chrysophytes were

relatively more abundant in the non-heated mesocosms (Fig. S6.2D).

3.4. Effect of the stressors on zooplankton community dynamics

The dynamics and composition of the zooplankton community and its responses to the

stressors differed markedly between the summer and winter experiments (Fig. 4). In summer,

zooplankton biomass showed significant differences between treatments (PRC: $F = 0.7$, $p =$

0.003; Fig. 4A). These differences were mainly driven by warming, with noticeable effects

on ostracods and *Daphnia* (Fig. 4A). Ostracod biomass increased in heated mesocosms, while

Daphnia biomass decreased compared to non-heated mesocosms (Figs 4A and S7.1b, Table

S7.3). However, copepod biomass tended to increase in all treatments, with a higher increase

in the treated mesocosms (Fig. S7.2a, Table S7.3). In winter, there were no significant

differences in zooplankton biomass and composition between treatments ($F = 0.3$, $p = 0.41$;

Fig. 4B, Figs S7.1c and S7.2b; Table S7.4).

Figure 4. Principal response curves (PRC) with species weights (vertical bars) indicating treatment-specific effects on zooplankton during the summer (A) and winter (B) experiment. Days = days after exposure. Treatments: CLT = non-heated, without pharmaceuticals (light blue); CHT = heated, without pharmaceuticals (yellow); TLT = non-heated, with pharmaceuticals (dark blue); THT = heated, with pharmaceuticals (orange). Days = days after exposure.

3.5. Effect of the stressors on predatory insect survival and emergence

The effects of both stressors on emergence and survival of predatory insects differed between taxa and experiments (Fig. 5; Tables S8.1 and S8.2). In summer, both warming and pharmaceuticals increased the emergence of anisopteran (*Aeshna*) larvae, with an additive effect on the predictor scale (Fig. 5A). However, warming had a negative effect on damselfly emergence (Fig. 5B). In winter, the survival of anisopteran (*Anax*) larvae was similar in all treatments, while damselfly larvae showed better survival when exposed to pharmaceuticals only or heating only, resulting in similar survival in treatments without both stressors and

with both stressors (Fig. 5C). *Notonecta* survival was reduced when exposed to warming in

winter, but pharmaceuticals had no effect on their survival (Fig. 5D).

Figure 5. Estimates of treatment-specific probability of emergence of insect predators in the summer

experiment (A: anisopteran larvae (*Aeshna*), B: zygopteran larvae) and probability of survival of

insect predators in the winter experiment (C: zygopteran larvae, D: *Notonecta* backswimmers).

Treatments: CLT = non-heated, without pharmaceuticals; CHT = heated, without pharmaceuticals;

TLT = non-heated, with pharmaceuticals; THT = heated, with pharmaceuticals. Model estimates are

based on the most parsimonious models and include 95% confidence intervals based on fixed effects.

3.6. Effect of the stressors on the pelagic food web

The effects of the stressors on the pelagic food web differed between summer and winter

experiments (Fig. 6 and Tables S9.1 and S9.2). In summer, pharmaceuticals negatively

affected the phytoplankton (standardised path coefficient < -0.01; Fig. 6A and Table S9.1),

while warming had a positive direct effect (standardised path coefficient = +0.08). Both

stressors had a direct and negative impact on filter-feeding cladocerans (standardised path

coefficients < -0.01 for pharmaceuticals and = -0.14 for warming). However, the relationship

between primary producers and filter feeders was not significant (standardised path

coefficient = -0.09), contradicting the trophic cascade hypothesis. The effects of stressors on

predators were positive; it was significant for warming (direct effect, standardised path

coefficients = +0.02) but not for pharmaceuticals (direct effect, standardised path coefficients

< 0.01). Finally, predators directly caused a decrease in filter feeder biomass (standardised

path coefficient = -0.21; Fig. 6A).

In contrast, in winter, the pharmaceutical mixture had no significant direct or indirect

effect on any component of the simplified pelagic food web. Only phytoplankton biomass

was directly and positively affected by warming during winter (standardized path coefficient

= 0.28; Fig. 6B and Table S9.2).

Fisher's C = 2.55, p = 0.28, df = 2   Fisher's C = 5.20, p = 0.079, df = 2

Figure 6. Path diagram of the selected structural equation model based on average data collected

during the summer experiment, and last sampling data during the winter experiment. PhAC =

pharmaceuticals. Orange, blue, and dashed arrows respectively indicate significantly ($p < 0.05$)

negative, significantly positive, and non-significant ($p > 0.05$) relationships between variables.

Proportions of variation explained by the model for each response variable given by R^2 values. See

Tables S9.1 and S9.2 for details.

4. Discussion

We conducted the first study on the combined effects of a complex mixture of

pharmaceuticals and warming on freshwater communities. While previous studies focused on

simplified food webs (Wijewardene et al., 2021), we examined the consequences of these

stressors on three trophic levels, including phytoplankton, zooplankton, and aquatic insects as

top predators.

As expected, we observed only limited effects of the stressors on the community in the

winter experiment, whereas in summer, warming acted as the stronger stressor in line with

the dominance null model (Morris et al., 2022), and its effects amplified the effects of

pharmaceuticals on the freshwater community. Overall, both stressors had a negative effect

on zooplankton (mainly composed by filter feeders in our experiment) through direct and

indirect effects, promoting phytoplankton development. Our results demonstrate that the

effects of environmentally relevant pharmaceutical mixtures should be considered together

with warming when studying possible future responses of complex freshwater communities

to environmental stressors.

4.1. Effects of warming on the freshwater food web

Warming had the stronger impact on the freshwater food web in our experiments, especially

during summer. During the summer experiment, the daily mean temperature in the heated

mesocosms reached up to 29.5°C (compared to 25.5°C in the non-heated mesocosms) and

remained above 25°C for almost three weeks in July. Such temperatures are comparable to

the optimum growth temperatures of the larvae of many odonates (Suhling et al., 2015) and

green algae (25–35 °C range; Bouterfas et al., 2002) but well above the reported optima for

Daphnia (16–22 °C range; Bruijning et al., 2018).

Warming may indirectly impact community dynamics through increased evaporation,

potentially leading to eutrophication and shifts in stoichiometry (Ptacnik et al., 2010). In our

summer experiment, water levels decreased faster in the heated mesocosms due to higher

evaporation. However, our experimental designs prevented hypoxic and anoxic conditions,

and no significant stoichiometry shift in available nutrients was observed at the end of the

experiment.

Warming can also favour fast-growing copiotrophs (Mooij et al., 2008; Ren et al.,

2017; Zhang et al., 2019) and toxin-producing cyanobacteria (Huisman et al., 2018),

intensifying resource competition with phytoplankton (Joint et al., 2002). Moreover, it can

reduce phytoplankton diversity, leading to shifts in community composition (Urrutia-Cordero

et al., 2017). In our summer experiment, phytoplankton biomass (indicated by chlorophyll-a

concentrations) increased, with cyanobacteria becoming more prevalent in the heated

mesocosms towards the end of the summer experiment, while non-heated mesocosms

remained dominated by green algae. These findings align with Hansson et al. (2020), who

showed that a +4°C temperature increase led to a shift in the phytoplankton community,

favouring cyanobacteria.

Zooplankton grazing can modulate the effects of warming on phytoplankton (Velthuis

et al., 2017). For example, Yvon-Durocher et al. (2011) found that a +4°C temperature

increase in an outdoor mesocosm experiment led to a decrease in phytoplankton biomass,

favouring smaller organisms within the phytoplankton assemblage, while zooplankton

biomass remained unaffected unlike in our experiment.

In tri-trophic systems, the effects of warming should vary predictably between trophic

levels (Hansson et al., 2013). Top predators, by consuming zooplankton and reducing grazing

pressure on phytoplankton, often play a crucial role in modulating the impacts of climate

warming on the community (Hansson et al., 2013; Šorf et al., 2015; Velthuis et al., 2017).

Consequently, warming is expected to have positive effects on top predators and

phytoplankton but negative effects on zooplankton (Hansson et al., 2013 but see Yvon-

Durocher et al., 2011). For example, He et al. (2018) observed interactive effects of warming

(+3°C above ambient) and predation (by the fish *Aristichthys nobilis*) in an outdoor

mesocosm experiment, leading to higher phytoplankton biomass and lower biomasses of

Daphnia and total zooplankton under the combined stressors.

Results from our summer experiment support and further extend these findings. We

observed a shift in the zooplankton community composition under warming, with ostracods

dominating in response to higher temperatures. Predator survival and emergence also

increased under warming. Moreover, predation rate of odonate larvae increases with warmer

temperatures (e.g. Sentis et al., 2015), resulting in accelerated development and earlier

emergence (Suhling et al., 2015). We conclude that these mechanisms caused higher

predation pressure on the zooplankton, which reduced its grazing pressure and led to higher

phytoplankton biomass.

4.2. Effects of pharmaceuticals on the freshwater food web

We observed no significant effects of pharmaceuticals in the winter experiment. However,

the pharmaceutical mixture alone had a negative effect on nutrients, phytoplankton biomass

and *Daphnia* biomass during summer.

Pharmaceutical mixtures can significantly impact microbial and algal communities and

ectotherms from zooplankton to fish (Brodin et al., 2014). However, the negative effects

reported in some studies were based on higher concentrations than those used in our study,

potentially overestimating environmental consequences. For instance, a mixture of 10

pharmaceuticals was toxic to algae, daphniids, and zebrafish at concentrations up to 15,000

times higher than environmental levels (Watanabe et al., 2016). High concentrations of a

pharmaceutical mixture (analgesics, antidepressants and antibiotics; >60 mg.L^{-1} for each

compound) in outdoor microcosms caused fish mortality and reduced zooplankton and

phytoplankton diversity (Richards et al., 2004). Conversely, David et al. (2020) found no

strong effects on a set of biomarkers and population dynamics of three-spined sticklebacks

(*Gasterosteus aculeatus*) at environmentally relevant concentrations of five pharmaceuticals

(antiepileptics, analgesics, and antihypertensives).

In previous *in-situ* studies, antibiotic mixtures induced significant structural and

functional changes in microbial attached communities (Proia et al., 2013), while a mixture of

pharmaceuticals decreased algal biomass and photosynthesis (Rosi-Marshall et al., 2013).

The decline in phytoplankton biomass exposed to pharmaceuticals in our summer experiment

supports these findings.

4.3. Interactions between warming and pharmaceuticals

While warming had stronger effects in our experiments, the combined impact of the

pharmaceutical mixture and warming was not predictable from individual stressor effects,

particularly in the summer experiment. The combination of pharmaceuticals and warming

significantly reduced phytoplankton biomass, and its control by filter-feeding zooplankton directly and indirectly via increased insect predator predation. This led to a change in phytoplankton composition, as expected in warming conditions (Kratina et al., 2012; Lewandowska et al., 2014). The qualitative effects of pharmaceuticals on top predators and zooplankton resembled those of warming in a tri-trophic system (Hansson et al., 2013), but their impact on phytoplankton differed.

These findings highlight the importance of community-level studies to assess stressor effects (Attrill and Depledge, 1997; Brooks and Crowe, 2019; Smith et al., 2009) and avoid underestimating their impacts (Jentsch et al., 2011). Relying solely on individual response variables for ecosystem function can be misleading in identifying underlying drivers of ecosystem changes (Bracken and Stachowicz, 2006; Hooper and Vitousek, 1998)

The interactions between temperature and contaminants present challenges for ecological risk assessment, particularly with global warming (Zhang et al., 2019). While the effects of warming on contaminant toxicity are well-studied, the impacts on shallow freshwater ecosystems can be complex, influenced by factors like contaminant type, community composition, and trophic interactions. For instance, heat waves combined with pesticide exposures can result in antagonistic or synergistic effects depending on the pesticides involved (Polazzo et al., 2022). Additionally, systems near their tolerance limit for one stressor may exhibit a lower threshold to withstand subsequent stress events (Hewitt et al., 2016).

Finally, the outcomes of our summer and winter experiments highlight the challenge of differentiating stress timing from seasonal species turnover. Although the communities had similar functional groups, differences in species composition, although subtle, made it challenging to disentangle the effects of seasonality, stressors, and community composition. However, we consider the taxonomic differences to be less significant than the overall

seasonal stage of the community, with winter communities being dormant or exhibiting low

metabolic activity compared to the highly active summer communities (see Kratina et al.,

2012).

Brooks and Crowe (2019) suggested considering temporal variation in both the cause

and effect to enhance understanding and management of multiple stressors. With increasing

unpredictability in the future, accounting for the intensity and duration of stress events

becomes crucial as they play a significant role in ecosystems' responses (Easterling et al.,

2000; Jentsch et al., 2007; Walter et al., 2013). Although our study did not directly assess

seasonal variation, previous research indicates that climate-driven multiple stressors may

vary seasonally (Godbold and Solan, 2013; Kratina et al., 2012). Additionally, climate

change may alter terrestrial runoff patterns (Easterling et al., 2000; Masson-Delmotte et al.,

2021), affecting local stressor arrival in aquatic systems (Adams, 2005; Schiedek et al.,

2007). Further research is therefore required to explore the consequences of these changes.

5. Conclusions and perspective

Our findings demonstrate that low concentrations of pharmaceuticals alone had minimal

impact, but when combined with warming, they significantly altered the pelagic food web,

especially during summer when temperatures were close to or above the upper thermal limits

for many species. Additionally, we observed distinct effects of warming and pharmaceuticals

between seasons.

Our study highlights that the responses of freshwater communities to environmental

stressors are complex and involve both direct effects on individual species and indirect

effects through trophic interactions. These complexities are vital to consider when predicting

future changes in biodiversity and ecosystem functioning (Hansson et al., 2013; Polazzo et

al., 2022). The lack of community-level studies limits our understanding of ecosystem-level

responses, especially when studying multiple interacting stressors. Although distinguishing between the effects of multiple stressors and understanding emergent phenomena is challenging, future research should consider larger spatio-temporal scales and a more comprehensive understanding of food web interactions. This will help us better predict the impacts of warming and emerging contaminants like pharmaceuticals and aid decision-making to mitigate their effects on freshwater ecosystems.

6. Acknowledgements

This study was supported by the Czech Science Foundation (project number GA20-16111S) and by the Ministry of Education, Youth and Sports of the Czech Republic (project CENAKVA, LM2023038). We thank Vladimíra Dekanová, Kateřina Bláhová, Bruno Carreira, Samuel Dijoux, Marius Wieczek and Fanny Verfaillie for assistance during the mesocosm experiment.

7. References

Adams, S.M., 2005. Assessing cause and effect of multiple stressors on marine systems. Mar. Pollut. Bull., 51, 649–657. https://doi.org/10.1016/j.marpolbul.2004.11.040

Amoros, C., 1984. Introduction pratique à la systématique des organismes des eaux continentales françaises - 5. Crustacés Cladocères. Publ. Société Linn. Lyon 53, 72–107. https://doi.org/10.3406/linly.1984.10627

Attrill, M.J., Depledge, M.H., 1997. Community and population indicators of ecosystem health: targeting links between levels of biological organisation. Aquat. Toxicol. 38, 183–197. https://doi.org/10.1016/S0166-445X(96)00839-9

Backhaus, T., 2014. Medicines, shaken and stirred: a critical review on the ecotoxicology of

pharmaceutical mixtures. Philos. Trans. R. Soc. B Biol. Sci. 369, 20130585.

https://doi.org/10.1098/rstb.2013.0585

Baert, J.M., Janssen, C.R., Sabbe, K., De Laender, F., 2016. Per capita interactions and stress

tolerance drive stress-induced changes in biodiversity effects on ecosystem functions. Nat.

Commun. 7, 12486. https://doi.org/10.1038/ncomms12486

Belden, J.B., Gilliom, R.J., Lydy, M.J., 2007. How well can we predict the toxicity of

pesticide mixtures to aquatic life? Integr. Environ. Assess. Manag. 3, 364–372.

https://doi.org/10.1002/ieam.5630030307

Bennett, A.F., Lenski, R.E., 2007. An experimental test of evolutionary trade-offs during

temperature adaptation. Proc. Natl. Acad. Sci. 104, 8649–8654.

https://doi.org/10.1073/pnas.0702117104

Bláha, M., Grabicova, K., Shaliutina, O., Kubec, J., Randák, T., Zlabek, V., Buřič, M.,

Veselý, L., 2019. Foraging behaviour of top predators mediated by pollution of

psychoactive pharmaceuticals and effects on ecosystem stability. Sci. Total Environ. 662,

655–661. https://doi.org/10.1016/j.scitotenv.2019.01.295

Boukal, D.S., Bideault, A., Carreira, B.M., Sentis, A., 2019. Species interactions under

climate change: connecting kinetic effects of temperature on individuals to community

dynamics. Curr. Opin. Insect Sci., Global change biology • Molecular Physiology 35, 88–

95. https://doi.org/10.1016/j.cois.2019.06.014

Bouterfas, R., Belkoura, M., Dauta, A., 2002. Light and temperature effects on the growth

rate of three freshwater [2pt] algae isolated from a eutrophic lake. Hydrobiologia 489,

207–217. https://doi.org/10.1023/A:1023241006464

Bracken, M.E.S., Stachowicz, J.J., 2006. Seaweed Diversity Enhances Nitrogen Uptake Via

Complementary Use of Nitrate and Ammonium. Ecology 87, 2397–2403.

https://doi.org/10.1890/0012-9658(2006)87[2397:SDENUV]2.0.CO;2

Brodin, T., Piovano, S., Fick, J., Klaminder, J., Heynen, M., Jonsson, M., 2014. Ecological

effects of pharmaceuticals in aquatic systems—impacts through behavioural alterations.

Philos. Trans. R. Soc. B Biol. Sci. 369, 20130580. https://doi.org/10.1098/rstb.2013.0580

Brogan, W.R., III, Relyea, R.A., 2015. Submerged Macrophytes Mitigate Direct and Indirect

Insecticide Effects in Freshwater Communities. PLOS ONE 10, e0126677.

https://doi.org/10.1371/journal.pone.0126677

Brooks, P.R., Crowe, T.P., 2019. Combined Effects of Multiple Stressors: New Insights Into

the Influence of Timing and Sequence. Front. Ecol. Evol. 7.

Bruijning, M., ten Berge, A.C.M., Jongejans, E., 2018. Population-level responses to

temperature, density and clonal differences in Daphnia magna as revealed by integral

projection modelling. Funct. Ecol. 32, 2407–2422. https://doi.org/10.1111/1365-

2435.13192

Cadmus, P., Pomeranz, J.P.F., Kraus, J.M., 2016. Low-cost floating emergence net and bottle

trap: comparison of two designs. J. Freshw. Ecol. 31, 653–658.

https://doi.org/10.1080/02705060.2016.1217944

David, V., Joachim, S., Catteau, A., Nott, K., Ronkart, S., Robert, C., Gillard, N., Bado-

Nilles, A., Chadili, E., Palluel, O., Turies, C., Julian, N., Castiglione, J., Dedourge-

Geffard, O., Hani, Y., Geffard, A., Porcher, J.-M., Beaudouin, R., 2020. Effects of chronic

exposure to a pharmaceutical mixture on the three-spined stickleback (gasterosteus

aculeatus) population dynamics in lotic mesocosms. Aquat. Toxicol. 224, 105499.

https://doi.org/10.1016/j.aquatox.2020.105499

Du, B., Haddad, S.P., Luek, A., Scott, W.C., Saari, G.N., Kristofco, L.A., Connors, K.A.,

Rash, C., Rasmussen, J.B., Chambliss, C.K., Brooks, B.W., 2014. Bioaccumulation and

trophic dilution of human pharmaceuticals across trophic positions of an effluent-

dependent wadeable stream. Philos. Trans. R. Soc. B Biol. Sci. 369, 20140058.

https://doi.org/10.1098/rstb.2014.0058

Duchet, C., Moraru, G.M., Spencer, M., Saurav, K., Bertrand, C., Fayolle, S., Gershberg

Hayoon, A., Shapir, R., Steindler, L., Blaustein, L., 2018. Pesticide-mediated trophic

cascade and an ecological trap for mosquitoes. Ecosphere 9, e02179.

https://doi.org/10.1002/ecs2.2179

Easterling, D.R., Meehl, G.A., Parmesan, C., Changnon, S.A., Karl, T.R., Mearns, L.O.,

2000. Climate Extremes: Observations, Modeling, and Impacts. Science 289, 2068–2074.

https://doi.org/10.1126/science.289.5487.2068

Fedorova, G., Grabic, R., Grabicová, K., Turek, J., Van Nguyen, T., Randak, T., Brooks,

B.W., Zlabek, V., 2022. Water reuse for aquaculture: Comparative removal efficacy and

aquatic hazard reduction of pharmaceuticals by a pond treatment system during a one year

study. J. Hazard. Mater. 421, 126712. https://doi.org/10.1016/j.jhazmat.2021.126712

Fekadu, S., Alemayehu, E., Dewil, R., Van der Bruggen, B., 2019. Pharmaceuticals in

freshwater aquatic environments: A comparison of the African and European challenge.

Sci. Total Environ. 654, 324–337. https://doi.org/10.1016/j.scitotenv.2018.11.072

Fursdon, J.B., Martin, J.M., Bertram, M.G., Lehtonen, T.K., Wong, B.B.M., 2019. The

pharmaceutical pollutant fluoxetine alters reproductive behaviour in a fish independent of

predation risk. Sci. Total Environ. 650, 642–652.

https://doi.org/10.1016/j.scitotenv.2018.09.046

Godbold, J.A., Solan, M., 2013. Long-term effects of warming and ocean acidification are

modified by seasonal variation in species responses and environmental conditions. Philos.

Trans. R. Soc. B Biol. Sci. 368, 20130186. https://doi.org/10.1098/rstb.2013.0186

Grabicova, K., Grabic, R., Blaha, M., Kumar, V., Cerveny, D., Fedorova, G., Randak, T.,

2015. Presence of pharmaceuticals in benthic fauna living in a small stream affected by

effluent from a municipal sewage treatment plant. Water Res. 72, 145–153.

https://doi.org/10.1016/j.watres.2014.09.018

Grabicova, K., Grabic, R., Fedorova, G., Fick, J., Cerveny, D., Kolarova, J., Turek, J.,

Zlabek, V., Randak, T., 2017. Bioaccumulation of psychoactive pharmaceuticals in fish in

an effluent dominated stream. Water Res. 124, 654–662.

https://doi.org/10.1016/j.watres.2017.08.018

Guedes, R.N.C., Smagghe, G., Stark, J.D., Desneux, N., 2016. Pesticide-Induced Stress in

Arthropod Pests for Optimized Integrated Pest Management Programs. Annu. Rev.

Entomol. 61, 43–62. https://doi.org/10.1146/annurev-ento-010715-023646

Hansson, L., Ekvall, M.K., He, L., Li, Z., Svensson, M., Urrutia☐Cordero, P., Zhang, H.,

2020. Different climate scenarios alter dominance patterns among aquatic primary

producers in temperate systems. Limnol. Oceanogr. 65, 2328–2336.

https://doi.org/10.1002/lno.11455

Hansson, L.-A., Nicolle, A., Granéli, W., Hallgren, P., Kritzberg, E., Persson, A., Björk, J.,

Nilsson, P.A., Brönmark, C., 2013. Food-chain length alters community responses to

global change in aquatic systems. Nat. Clim. Change 3, 228–233.

https://doi.org/10.1038/nclimate1689

He, H., Jin, H., Jeppesen, E., Li, K., Liu, Z., Zhang, Y., 2018. Fish-mediated plankton

responses to increased temperature in subtropical aquatic mesocosm ecosystems:

Implications for lake management. Water Res. 144, 304–311.

https://doi.org/10.1016/j.watres.2018.07.055

Hewitt, J.E., Ellis, J.I., Thrush, S.F., 2016. Multiple stressors, nonlinear effects and the

implications of climate change impacts on marine coastal ecosystems. Glob. Change Biol.

22, 2665–2675. https://doi.org/10.1111/gcb.13176

Heye, K., Lotz, T., Wick, A., Oehlmann, J., 2019. Interactive effects of biotic and abiotic

environmental stressors on carbamazepine toxicity in the non-biting midge Chironomus

riparius. Water Res. 156, 92–101. https://doi.org/10.1016/j.watres.2019.03.007

Hooper, D.U., Vitousek, P.M., 1998. Effects of Plant Composition and Diversity on Nutrient

Cycling. Ecol. Monogr. 68, 121–149. https://doi.org/10.1890/0012-

9615(1998)068[0121:EOPCAD]2.0.CO;2

Huisman, J., Codd, G.A., Paerl, H.W., Ibelings, B.W., Verspagen, J.M.H., Visser, P.M.,

2018. Cyanobacterial blooms. Nat. Rev. Microbiol. 16, 471–483.

https://doi.org/10.1038/s41579-018-0040-1

IPBES, 2019. Global assessment report on biodiversity and ecosystem services of the

Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services.

Zenodo. https://doi.org/10.5281/zenodo.6417333

Jentsch, A., Kreyling, J., Beierkuhnlein, C., 2007. A new generation of climate-change

experiments: events, not trends. Front. Ecol. Environ. 5, 365–374.

https://doi.org/10.1890/1540-9295(2007)5[365:ANGOCE]2.0.CO;2

Jentsch, A., Kreyling, J., Elmer, M., Gellesch, E., Glaser, B., Grant, K., Hein, R., Lara, M.,

Mirzae, H., Nadler, S.E., Nagy, L., Otieno, D., Pritsch, K., Rascher, U., Schädler, M.,

Schloter, M., Singh, B.K., Stadler, J., Walter, J., … Beierkuhnlein, C., 2011. Climate

extremes initiate ecosystem-regulating functions while maintaining productivity. J. Ecol.

99, 689–702. https://doi.org/10.1111/j.1365-2745.2011.01817.x

Johannsen, O.A., 1937. Aquatic diptera. Cornell University.

Joint, I., Henriksen, P., Fonnes, G.A., Bourne, D., Thingstad, T.F., Riemann, B., 2002.

Competition for inorganic nutrients between phytoplankton and bacterioplankton in

nutrient manipulated mesocosms. Aquat. Microb. Ecol. 29, 145–159.

https://doi.org/10.3354/ame029145

Kneitel, J.M., Chase, J.M., 2004. Trade-offs in community ecology: linking spatial scales and

species coexistence. Ecol. Lett. 7, 69–80. https://doi.org/10.1046/j.1461-

0248.2003.00551.x

Kratina, P., Greig, H.S., Thompson, P.L., Carvalho-Pereira, T.S.A., Shurin, J.B., 2012.

Warming modifies trophic cascades and eutrophication in experimental freshwater

communities. Ecology 93, 1421–1430. https://doi.org/10.1890/11-1595.1

Lagesson, A., Fahlman, J., Brodin, T., Fick, J., Jonsson, M., Byström, P., Klaminder, J.,

2016. Bioaccumulation of five pharmaceuticals at multiple trophic levels in an aquatic

food web - Insights from a field experiment. Sci. Total Environ. 568, 208–215.

https://doi.org/10.1016/j.scitotenv.2016.05.206

Lewandowska, A.M., Hillebrand, H., Lengfellner, K., Sommer, U., 2014. Temperature

effects on phytoplankton diversity — The zooplankton link. J. Sea Res. 85, 359–364.

https://doi.org/10.1016/j.seares.2013.07.003

Loos, R., Carvalho, R., António, D.C., Comero, S., Locoro, G., Tavazzi, S., Paracchini, B.,

Ghiani, M., Lettieri, T., Blaha, L., Jarosova, B., Voorspoels, S., Servaes, K., Haglund, P.,

Fick, J., Lindberg, R.H., Schwesig, D., Gawlik, B.M., 2013. EU-wide monitoring survey

on emerging polar organic contaminants in wastewater treatment plant effluents. Water

Res. 47, 6475–6487. https://doi.org/10.1016/j.watres.2013.08.024

Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S.L., Péan, C., Berger, S., Caud, N.,

Chen, Y., Goldfarb, L., Gomis, M.I., Huang, M., Leitzell, K., Lonnoy, E., Matthews,

J.B.R., Maycock, T.K., Waterfield, T., Yelekçi, Ö., Yu, R., Zhou, B. (Eds.), 2021. Climate

Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth

Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge

University Press, Cambridge, United Kingdom and New York, NY, USA.

https://doi.org/10.1017/9781009157896

Mooij, W.M., De Senerpont Domis, L.N., Hülsmann, S., 2008. The impact of climate

warming on water temperature, timing of hatching and young-of-the-year growth of fish in

shallow lakes in the Netherlands. J. Sea Res., Dynamics of Fish and Fishers 60, 32–43.

https://doi.org/10.1016/j.seares.2008.03.002

Morris, O.F., Loewen, C.J.G., Woodward, G., Schäfer, R.B., Piggott, J.J., Vinebrooke, R.D.,

Jackson, M.C., 2022. Local stressors mask the effects of warming in freshwater

ecosystems. Ecol. Lett. 25, 2540–2551. https://doi.org/10.1111/ele.14108

Noyes, P.D., McElwee, M.K., Miller, H.D., Clark, B.W., Van Tiem, L.A., Walcott, K.C.,

Erwin, K.N., Levin, E.D., 2009. The toxicology of climate change: Environmental

contaminants in a warming world. Environ. Int. 35, 971–986.

https://doi.org/10.1016/j.envint.2009.02.006

Op de Beeck, L., Verheyen, J., Olsen, K., Stoks, R., 2017. Negative effects of pesticides

under global warming can be counteracted by a higher degradation rate and thermal

adaptation. J. Appl. Ecol. 54, 1847–1855. https://doi.org/10.1111/1365-2664.12919

Parmesan, C., Yohe, G., 2003. A globally coherent fingerprint of climate change impacts

across natural systems. Nature 421, 37–42. https://doi.org/10.1038/nature01286

Polazzo, F., Roth, S.K., Hermann, M., Mangold␣Döring, A., Rico, A., Sobek, A., Van den

Brink, P.J., Jackson, M.C., 2022. Combined effects of heatwaves and micropollutants on

freshwater ecosystems: Towards an integrated assessment of extreme events in multiple

stressors research. Glob. Change Biol. 28, 1248–1267. https://doi.org/10.1111/gcb.15971

Post, E., 2013. Ecology of Climate Change: The Importance of Biotic Interactions, Ecology

of Climate Change. Princeton University Press. https://doi.org/10.1515/9781400846139

Proia, L., Lupini, G., Osorio, V., Pérez, S., Barceló, D., Schwartz, T., Amalfitano, S., Fazi,

S., Romaní, A.M., Sabater, S., 2013. Response of biofilm bacterial communities to

antibiotic pollutants in a Mediterranean river. Chemosphere 92, 1126–1135.

https://doi.org/10.1016/j.chemosphere.2013.01.063

Ptacnik, R., Moorthi, S.D., Hillebrand, H., 2010. Chapter 1 - Hutchinson Reversed, or Why

There Need to Be So Many Species, in: Woodward, G. (Ed.), Advances in Ecological

Research, Integrative Ecology: From Molecules to Ecosystems. Academic Press, pp. 1–43.

https://doi.org/10.1016/B978-0-12-385005-8.00001-0

R Core Team, 2018. R: A language and environment for statistical computing. 2014. R

Found. Stat. Comput. Vienna Austria.

Relyea, R.A., 2009. A cocktail of contaminants: how mixtures of pesticides at low

concentrations affect aquatic communities. Oecologia 159, 363–376.

https://doi.org/10.1007/s00442-008-1213-9

Ren, L., He, D., Chen, Z., Jeppesen, E., Lauridsen, T.L., Søndergaard, M., Liu, Z., Wu, Q.L.,

2017. Warming and nutrient enrichment in combination increase stochasticity and beta

diversity of bacterioplankton assemblages across freshwater mesocosms. ISME J. 11, 613–

625. https://doi.org/10.1038/ismej.2016.159

Richards, S.M., Wilson, C.J., Johnson, D.J., Castle, D.M., Lam, M., Mabury, S.A., Sibley,

P.K., Solomon, K.R., 2004. Effects of Pharmaceutical Mixtures in Aquatic Microcosms.

Environ. Toxicol. Chem. 23, 1035. https://doi.org/10.1897/02-616

Rodrigues, A.C.M., Machado, A.L., Bordalo, M.D., Saro, L., Simão, F.C.P., Rocha, R.J.M.,

Golovko, O., Žlábek, V., Barata, C., Soares, A.M.V.M., Pestana, J.L.T., 2018. Invasive

Species Mediate Insecticide Effects on Community and Ecosystem Functioning. Environ.

Sci. Technol. 52, 4889–4900. https://doi.org/10.1021/acs.est.8b00193

Rosi-Marshall, E.J., Kincaid, D.W., Bechtold, H.A., Royer, T.V., Rojas, M., Kelly, J.J., 2013.

Pharmaceuticals suppress algal growth and microbial respiration and alter bacterial

communities in stream biofilms. Ecol. Appl. 23, 583–593. https://doi.org/10.1890/12-

0491.1

Saaristo, M., Lagesson, A., Bertram, M.G., Fick, J., Klaminder, J., Johnstone, C.P., Wong,

B.B.M., Brodin, T., 2019. Behavioural effects of psychoactive pharmaceutical exposure on

European perch (Perca fluviatilis) in a multi-stressor environment. Sci. Total Environ. 655,

1311–1320. https://doi.org/10.1016/j.scitotenv.2018.11.228

Sánchez-Bayo, F., 2021. Indirect Effect of Pesticides on Insects and Other Arthropods.

Toxics 9, 177. https://doi.org/10.3390/toxics9080177

Schiedek, D., Sundelin, B., Readman, J.W., Macdonald, R.W., 2007. Interactions between

climate change and contaminants. Mar. Pollut. Bull. 54, 1845–1856.

https://doi.org/10.1016/j.marpolbul.2007.09.020

Sentis, A., Morisson, J., Boukal, D.S., 2015. Thermal acclimation modulates the impacts of

temperature and enrichment on trophic interaction strengths and population dynamics.

Glob. Change Biol. 21, 3290–3298. https://doi.org/10.1111/gcb.12931

Smith, M.D., Knapp, A.K., Collins, S.L., 2009. A framework for assessing ecosystem

dynamics in response to chronic resource alterations induced by global change. Ecology

90, 3279–3289. https://doi.org/10.1890/08-1815.1

Šorf, M., Davidson, T.A., Brucet, S., Menezes, R.F., Søndergaard, M., Lauridsen, T.L.,

Landkildehus, F., Liboriussen, L., Jeppesen, E., 2015. Zooplankton response to climate

warming: a mesocosm experiment at contrasting temperatures and nutrient levels.

Hydrobiologia 742, 185–203. https://doi.org/10.1007/s10750-014-1985-3

Spaak, J.W., Baert, J.M., Baird, D.J., Eisenhauer, N., Maltby, L., Pomati, F., Radchuk, V.,

Rohr, J.R., Van den Brink, P.J., De Laender, F., 2017. Shifts of community composition

and population density substantially affect ecosystem function despite invariant richness.

Ecol. Lett. 20, 1315–1324. https://doi.org/10.1111/ele.12828

Stephenson, G.L., Hamilton, P., Kaushik, N.K., Robinson, J.B., Solomon, K.R., 1984. Spatial

distribution of plankton in enclosures of three sizes. Can. J. Fish. Aquat. Sci. 41, 1048–

1054.

Suhling, F., Suhling, I., Richter, O., 2015. Temperature response of growth of larval

dragonflies – an overview. Int. J. Odonatol. 18, 15–30.

https://doi.org/10.1080/13887890.2015.1009392

Švecová, H., Grabic, R., Grabicová, K., Vojs Staňová, A., Fedorova, G., Cerveny, D., Turek,

J., Randák, T., Brooks, B.W., 2021. De facto reuse at the watershed scale: Seasonal

changes, population contributions, instream flows and water quality hazards of human

pharmaceuticals. Environ. Pollut. 268, 115888.

https://doi.org/10.1016/j.envpol.2020.115888

ter Braak, C.J.F., Smilauer, P., 2012. Canoco reference manual and user's guide: software for

ordination, version 5.0.

Urrutia-Cordero, P., Ekvall, M.K., Ratcovich, J., Soares, M., Wilken, S., Zhang, H., Hansson,

L.-A., 2017. Phytoplankton diversity loss along a gradient of future warming and

brownification in freshwater mesocosms. Freshw. Biol. 62, 1869–1878.

https://doi.org/10.1111/fwb.13027

Velthuis, M., de Senerpont Domis, L.N., Frenken, T., Stephan, S., Kazanjian, G., Aben, R.,

Hilt, S., Kosten, S., van Donk, E., Van de Waal, D.B., 2017. Warming advances top-down

control and reduces producer biomass in a freshwater plankton community. Ecosphere 8,

e01651. https://doi.org/10.1002/ecs2.1651

Venâncio, C., Ribeiro, R., Soares, A., Lopes, I., 2016. Multiple Stressor Differential

Tolerances: Possible Implications at the Population Level. PLOS ONE 11, e0151847.

https://doi.org/10.1371/journal.pone.0151847

Vörösmarty, C.J., McIntyre, P.B., Gessner, M.O., Dudgeon, D., Prusevich, A., Green, P.,

Glidden, S., Bunn, S.E., Sullivan, C.A., Liermann, C.R., Davies, P.M., 2010. Global

threats to human water security and river biodiversity. Nature 467, 555–561.

https://doi.org/10.1038/nature09440

Walter, J., Jentsch, A., Beierkuhnlein, C., Kreyling, J., 2013. Ecological stress memory and

cross stress tolerance in plants in the face of climate extremes. Environ. Exp. Bot., Cross-

stress tolerance and stress "memory" in plants 94, 3–8.

https://doi.org/10.1016/j.envexpbot.2012.02.009

Watanabe, H., Tamura, I., Abe, R., Takanobu, H., Nakamura, A., Suzuki, T., Hirose, A.,

Nishimura, T., Tatarazako, N., 2016. Chronic toxicity of an environmentally relevant

mixture of pharmaceuticals to three aquatic organisms (alga, daphnid, and fish). Environ.

Toxicol. Chem. 35, 996–1006. https://doi.org/10.1002/etc.3285

Wijewardene, L., Wu, N., Qu, Y., Guo, K., Messyasz, B., Lorenz, S., Riis, T., Ulrich, U.,

Fohrer, N., 2021. Influences of pesticides, nutrients, and local environmental variables on

phytoplankton communities in lentic small water bodies in a German lowland agricultural

area. Sci. Total Environ. 780, 146481. https://doi.org/10.1016/j.scitotenv.2021.146481

Wilkinson, J.L., Boxall, A.B.A., Kolpin, D.W., Leung, K.M.Y., Lai, R.W.S., Galbán-

Malagón, C., Adell, A.D., Mondon, J., Metian, M., Marchant, R.A., Bouzas-Monroy, A.,

Cuni-Sanchez, A., Coors, A., Carriquiriborde, P., Rojo, M., Gordon, C., Cara, M.,

Moermond, M., Luarte, T., … Teta, C., 2022. Pharmaceutical pollution of the world's

rivers. Proc. Natl. Acad. Sci. 119, e2113947119. https://doi.org/10.1073/pnas.2113947119

Woodward, G., Perkins, D.M., Brown, L.E., 2010. Climate change and freshwater

ecosystems: impacts across multiple levels of organization. Philos. Trans. R. Soc. B Biol.

Sci. 365, 2093–2106. https://doi.org/10.1098/rstb.2010.0055

Xie, Z., Lu, G., Yan, Z., Liu, J., Wang, P., Wang, Y., 2017. Bioaccumulation and trophic

transfer of pharmaceuticals in food webs from a large freshwater lake. Environ. Pollut.

222, 356–366. https://doi.org/10.1016/j.envpol.2016.12.026

Yvon-Durocher, G., Montoya, J.M., Trimmer, M., Woodward, G., 2011. Warming alters the

size spectrum and shifts the distribution of biomass in freshwater ecosystems. Glob.

Change Biol. 17, 1681–1694. https://doi.org/10.1111/j.1365-2486.2010.02321.x

Zhang, H., Huo, S., Yeager, K.M., He, Z., Xi, B., Li, X., Ma, C., Wu, F., 2019.

Phytoplankton response to climate changes and anthropogenic activities recorded by

sedimentary pigments in a shallow eutrophied lake. Sci. Total Environ. 647, 1398–1409.

https://doi.org/10.1016/j.scitotenv.2018.08.081

789