**Preservation of co-expression defines the primary tissue fidelity of human neural organoids**

Jonathan M. Werner[1], Jesse Gillis[1,2, *]

[1]The Stanley Institute for Cognitive Genomics, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY 11724, USA

[2]Physiology Department and Donnelly Centre for Cellular and Biomolecular Research, University of Toronto, Toronto, ON, Canada

*Corresponding author

**Abstract:**

Human neural organoid models offer an exciting opportunity for studying often inaccessible human-specific brain development; however, it remains unclear how precisely organoids recapitulate fetal/primary tissue biology. Here, we characterize field-wide replicability and biological fidelity through a meta-analysis of single-cell RNA-sequencing data for first and second trimester human primary brain (2.95 million cells, 51 datasets) and neural organoids (1.63 million cells, 130 datasets). We quantify the degree to which primary tissue cell-type marker expression and co-expression are recapitulated in organoids across 12 different protocol types. By quantifying gene-level preservation of primary tissue co-expression, we show neural organoids lie on a spectrum ranging from virtually no signal to co-expression near indistinguishable from primary tissue data, demonstrating high fidelity is within the scope of current methods. Additionally, we show neural organoids preserve the cell-type specific co-

expression of developing rather than adult cells, confirming organoids are an appropriate

model for primary tissue development. Overall, quantifying the preservation of primary tissue

co-expression is a powerful tool for uncovering unifying axes of variation across

heterogeneous neural organoid experiments.

**Introduction:**

Pluripotent stem cells create self-organized multi-cellular structures, termed organoids,

when cultured in a 3D *in vitro* environment[1,2]. The advantage of organoid models over 2D cell

culture counterparts is their ability to generate structures that resemble endogenous tissues

both in the differentiated cell-types produced and their 3D spatial organization[3,4]. The ability to

model organogenesis in a controlled *in vitro* environment creates opportunities to study

previously inaccessible developmental tissues from both humans and a range of model

organisms[5,6,7]. As such, organoids are genetically accessible[8] and environmentally

perturbable[9] models enabling the study of molecular, cellular, and developmental mechanisms

behind tissue construction. However, the applicability of studies in organoids to *in vivo* biology

hinges on how well these *in vitro* models recapitulate primary tissue developmental processes,

which remains an open question.

Quantifying the degree to which organoid systems replicate primary tissue biological

processes is a critical step toward understanding the strengths and limitations of these *in vitro*

models[10–14]. However, studies that perform such primary tissue/organoid comparisons are

inherently confounded by batch[15] (*in vivo* vs *in vitro*), making it difficult to disentangle batch

effects from underlying primary tissue and organoid biology. Meta-analytic approaches across

many primary tissue and organoid datasets offer a route around these confounds, enabling the

discovery of replicable primary tissue and organoid signatures independent of batch, which

can then be interrogated for how well organoids recapitulate primary tissue biology[16–18]. An

important biological signature for this purpose is gene co-expression[19]. Genes that are

functionally related tend to be expressed together, resulting in correlated gene expression

dynamics that can define functionally relevant gene modules[19]. Gene co-expression

relationships represent a shared genomic space that can be aggregated across experiments

(e.g.,[20]) in either *in vivo* or *in vitro* systems, thus providing a useful framework for quantifying

functional similarities and differences. Excitingly, coupling meta-analytic comparisons of

primary tissue and organoid co-expression with single-cell RNA-sequencing data (scRNA-seq)

stands to deliver cell-type specific quantifications of organoids' current capacity for producing

functionally equivalent cell-types to primary tissues[21,22].

Among organoid systems, human neural organoids are particularly well suited for meta-

analytic evaluation due to well-described broad cell-type annotations and their known lineage

relationships[23], the wide variety of differentiation protocols in use[24], and the increasing amount

of single-cell primary brain tissue and neural organoid data publicly available. In particular, the

diversity of differentiation protocols for human neural organoids poses a unique challenge for

organoid quality control that can be met by meta-analytic approaches. Neural organoids can

either be undirected[25] (multiple brain region identities) or directed (specific brain region

identity) with an increasing number of protocols striving to produce a wider variety of region-

specific organoids[11,26–37]. Meta-analytic primary tissue/organoid comparisons across

differentiation protocols stand to derive generalizable quality control metrics applicable to any

differentiation protocol, fulfilling a currently unmet need for unified quality control metrics

across heterogeneous neural organoids.

Prior comparisons between primary brain tissues and neural organoids demonstrated

that organoids have the capacity to produce diverse cell-types that capture both regional and

temporal variation similar to primary tissue data as assayed through transcriptomic[10,11,13,16,17,]

[38], epigenomic[39,40], electrophysiologic[41], and proteomic studies[42]. At the morphological level,

neural organoids can produce cellular organizations structurally similar to various *in vivo* brain

regions, including cortical layers[43] and hippocampus[27], as well as modeling known inter-

regional interactions like neuromuscular junctions[34] and interneuron migration[29]. Additionally,

several prior studies have compared primary tissue/organoid co-expression and concluded

that neural organoids recapitulate primary brain tissue co-expression[5,13,39], but these

assessments are highly targeted to study-specific properties, limiting potential generalization or

potential assessment across the field. Typically, only a single organoid differentiation protocol

is used in these assessments and it remains unclear whether organoids across different

protocols will produce similar results. This lack of breadth also affects the use of primary tissue

data used as a reference, with the primary tissue datasets utilized being treated as gold-

standard datasets with little consideration for the extent one primary tissue reference may

generalize to another. While prior meta-analytic comparisons of primary tissue/organoid co-

expression have been performed[17], these were done at the bulk level (lack cell-type resolution)

and included a small number of cortical organoid protocols, limiting the biological resolution

and generalizability of these findings.

In this study, we perform a meta-analytic assessment of primary brain tissue (2.95

million cells, 50 datasets, Fig. 1A) and neural organoid (1.63 million cells, 130 datasets, 12

protocols, Fig. 1B) scRNA-seq datasets, constructing robust primary tissue cell-type specific

markers and co-expression to query how well neural organoids recapitulate primary tissue cell-

type specific biology. We sample primary brain tissue data over the first and second trimesters

and across 15 different developmentally defined brain regions, extracting lists of cell-type

markers that define broad primary tissue cell-type identity regardless of temporal, regional, or

technical variation (Fig. 1A). We derive co-expression networks from individual primary tissue

and organoid datasets as well as aggregate co-expression networks across datasets (Fig. 1C).

From these networks, we assess the strength of co-expression within primary tissue cell-type

marker sets as well as the preservation of co-expression patterns between primary tissue and

organoid data (Fig. 1D-E). We also provide an R package to download our primary tissue

reference co-expression network to assay new neural organoid data using simple, meaningful,

and fast statistics (Fig. 1F). By constructing robust primary tissue cell-type representations

through meta-analytic approaches, we demonstrate the preservation of primary tissue cell-type

co-expression provides both specific and generalizable characterization of the primary tissue

fidelity of human neural organoids.

**Results:**

**Meta-analytic framework for primary tissue/organoid comparisons**

We reason that, if they exist, primary tissue cell-type specific signals robust to temporal,

regional, and technical variation will constitute *in vivo* standards applicable to any organoid

dataset regardless of time in culture or differentiation protocol. We first show it is possible to

learn sets of marker genes that define broad primary tissue cell-types (Fig. 1A, Supp. Table 1)

across timepoints (gestational weeks GW5-GW25) and brain regions (15 developmentally

defined brain regions) through a meta-analytic differential expression framework (Fig. 1A, Fig.

2A-B). We then compare co-expression within these marker sets between primary tissue and

organoid data to quantify the degree organoids preserve primary tissue cell-type specific co-

expression. An important aspect of our analysis is our cross-validation of primary tissue

differential expression and co-expression. We employ a leave-one-out cross-validation

approach when learning robust differentially expressed marker genes from our annotated

primary tissue datasets (2,174,934 cells, 37 datasets) and we interrogate co-expression of our

primary tissue marker genes within a large cohort of unannotated primary tissue datasets

(776,343 cells, 14 datasets). This approach ensures we are extracting primary tissue markers

and co-expression relationships independent of temporal, regional, and technical variation, a

powerful approach for deriving broad primary tissue signatures appropriate for comparison to a

wide range of organoid datasets.

**Cross-temporal and -regional primary tissue cell-type markers**

To learn markers that define broad primary tissue cell-types (see methods), we apply

the MetaMarkers[44] framework to our cross-temporal and -regional annotated primary tissue

datasets (Fig. 2A-B). MetaMarkers uses robust differential expression statistic thresholds (log2

fold-change >= 4 and FDR-adjusted p-value <= 0.05) for determining whether a gene is

differentially expressed (DE) within individual datasets, then ranks all genes via the strength of

their recurrent DE across datasets (see methods). We test the generalizability of our primary

tissue MetaMarker gene sets in predicting primary cell-types by employing a leave-one-out

primary tissue cross-validation (Fig. 2A-B). We construct an aggregate expression predictor in

the left-out dataset using MetaMarkers learned from the remaining datasets (see methods),

quantifying how well the MetaMarker gene sets predict the left-out cell-type annotations with

the area-under-the-receiver-operating-characteristic curve statistic (AUROC, Fig. 2B-C). The

AUROC is the probability of correctly prioritizing a true positive (e.g., cell of the right type)

above a negative, (e.g., cell of the wrong type), given some predictor of the positive class, in

this case, aggregate cell-type marker expression.

Starting with just the top 10 primary tissue MetaMarkers per cell-type, we achieve a

mean AUROC across all primary tissue datasets of 0.944 ± 0.0280 SD, 0.864 ± 0.0796 SD,

0.873 ± 0.0676 SD, 0.937 ± 0.0669 SD, 0.879 ± 0.0535 SD, and 0.931 ± 0.0737 SD, for

dividing progenitors, neural progenitors, intermediate progenitors, GABAergic neurons,

glutamatergic neurons, and non-neuronal cell-types respectively (Fig. 2C). These extremely

high performances demonstrate that even a small number of meta-analytically derived primary

tissue cell-type markers have high utility in predicting primary tissue cell-type annotations

regardless of temporal and regional variability. For all following analysis, we take the top 100

MetaMarkers per cell-type as robust representations of our 6 broad primary tissue cell-type

annotations (average AUROC >= 0.90 except for intermediate progenitors: 0.897 ± 0.0777

SD), with the 100 MetaMarkers achieving modest increases in performance over the top 10

MetaMarkers for all cell-types except GABAergic cells (Fig. 2C, mean AUROC for 100

GABAergic MetaMarkers: 0.922 ± 0.0777 SD). When comparing MetaMarkers to markers

derived from individual primary tissue datasets, we find the MetaMarkers are consistently top

performers in predicting primary tissue annotations (Fig. 2D), with MetaMarkers producing the

top results for intermediate progenitors, glutamatergic neurons, and GABAergic neurons

(Supp. Fig. 1), as well as comparable performance to top individual datasets for dividing

progenitors, neural progenitors, and non-neuronal cell-types (Supp. Fig. 1).

We explore the primary tissue MetaMarker sets further by computing the average

expression of the top 100 MetaMarkers for our 6 annotated cell-types across all cells within our

37 annotated primary tissue datasets (Fig. 2E), continuing our leave-one-out approach. Each

annotated primary tissue cell-type expresses the corresponding matched MetaMarker set over

all other MetaMarker sets, with the exception of some off-target expression for the neural

progenitor MetaMarkers in astrocytes (aggregated over all datasets Fig. 2E, individual datasets

Supp. Fig. 1B). This demonstrates our MetaMarker gene sets act as robust cell-type markers

in aggregate across all first and second trimester timepoints (Fig. 2E, Supp. Fig. 1B).

Additionally, we investigate the expression of the top 100 MetaMarker gene sets across

annotated primary brain regions, demonstrating each primary tissue cell-type maximally

expresses the corresponding primary tissue MetaMarker set across all annotated brain regions

(Supp. Fig. 2A-B). Overall, we are able to meta-analytically extract cell-type markers that

define broad primary tissue cell-types independent of temporal and regional variation.

**Broad primary tissue cell-type markers capture organoid temporal variation**

After extracting meta-analytic cell-type markers that capture broad primary tissue

temporal and regional variation, we can test how well these markers also capture organoid

temporal and regional (protocol) variation. We start with a large-scale temporal organoid

atlas[38] derived from a forebrain differentiation protocol containing timepoints ranging from 23

days to 6 months in culture. When comparing primary tissue and organoid data along a

temporal axis, one might expect younger primary tissue expression data to be a better

reference for younger organoid cell-types (better able to predict cell-types) and vice-versa for

older primary and organoid data (Supp. Fig. 3A). We test this relationship using the same

AUROC quantification as in Figure 1C, but now using the top 100 primary tissue cell-type

markers per primary tissue dataset to predict organoid cell-type annotations across all

organoid timepoints (Supp. Fig. 3B, see methods).

We observe highly consistent performance across all primary tissue datasets (GW5 –

GW25) when predicting organoid cell-types regardless of the organoid timepoint (Supp. Fig.

3B). The average difference in AUROC scores when predicting organoid cell-types using either

our youngest (GW5) or oldest (GW25) primary data is $0.000382 \pm 0.0357$ SD, $0.141 \pm 0.192$

SD, $0.139 \pm 0.0317$ SD, $0.00171 \pm 0.113$ SD and $0.119 \pm 0.216$ SD for dividing progenitors,

neural progenitors, glutamatergic neurons GABAergic neurons, and non-neuronal cells

respectively (No annotated intermediate progenitors in the GW25 primary tissue dataset). This

demonstrates strikingly consistent performance across distant primary tissue timepoints,

highlighting that broad primary tissue cell-type signatures are applicable as reference for

organoid cell-types regardless of the primary tissue or organoid timepoint. The one exception

is for neural progenitors, where there seemingly is a temporal shift in performance with

younger primary tissue datasets predicting younger organoid annotations over older organoid

annotations and vice-versa for older primary tissue/organoid data (Supp. Fig. 3B). However, a

subset of the young GW6-8 primary tissue datasets report sharp increases in performance

predicting older organoid timepoints in opposition to other GW6-8 primary tissue datasets,

suggesting variance in performance is driven by intersections between the quality of individual

organoid and primary tissue datasets rather than overarching temporal variability. Importantly,

our lists of top 100 primary tissue MetaMarkers perform comparably to marker sets from

individual primary tissue datasets, with less variance in performance across the organoid

timepoints for the differentiated cell-types (mean AUROC variance across organoid timepoints

for individual primary tissue datasets vs. primary MetaMarker variance; glutamatergic: 0.0147,

0.00672, GABAergic: 0.00487, 0.00201, non-neuronal: 0.00733, 0.00647, Supp. Fig. 3B). This

demonstrates our meta-analytic primary tissue cell-type markers robustly capture organoid

temporal variation.

**Broad primary tissue cell-type markers capture organoid protocol variation**

We assess whether our primary tissue MetaMarker gene sets capture organoid

variation outside the annotated forebrain temporal organoid atlas by performing principal-

component analysis (PCA) across all organoid datasets, representing data from 12 different

differentiation protocols. Our lists of 100 primary tissue MetaMarkers are consistently heavily

weighted in the first PC across organoid datasets (Supp. Fig. 3C-D). While a large portion of

PC1-weighted genes are dividing progenitor MetaMarkers (representing cell-cycle signal),

markers for non-dividing fetal cell-types also comprise those genes consistently heavily

weighted in PC1 across organoid datasets (Supp. Fig. 3C-D).

**Aggregate organoid co-expression weakly captures primary tissue co-expression**

Our primary tissue MetaMarkers that capture both primary tissue and organoid

temporal/regional variation enable assessments of cell-type specific co-expression between

arbitrary primary tissue and organoid datasets. One normally would need matched cell-type

annotations across datasets to compare cell-type specific biology, but here we couple our

meta-analytically derived cell-type markers with gene co-expression quantifications, which do

not rely on cell-type annotations, to extract cell-type specific co-expression from any given

scRNA-seq dataset. Practically, if organoids are producing cell-types functionally identical to

primary tissue cell-types, we would expect near identical co-expression relationships within our

primary tissue MetaMarker gene sets across primary tissue and organoid datasets.

We first explore marker set co-expression within our unannotated primary tissue

datasets, which were not included in deriving our primary tissue MetaMarker sets. The

aggregate (Fig. 3A, see methods) unannotated primary tissue co-expression network nearly

perfectly constructs cell-type specific co-expression modules when hierarchically clustering the

co-expression of our top 100 primary tissue MetaMarker gene sets (Fig. 3B). Turning to the

aggregate organoid co-expression network, while some cell-type co-expression structure

exists, it is much weaker than the unannotated primary tissue co-expression with less well-

defined intra-gene set co-expression relationships (Fig. 3B). We quantify this through the

Adjusted Rands Index (ARI) metric, comparing the MetaMarker clustering through co-

expression in any given network to the perfect clustering of MetaMarker gene sets by cell-type.

We perform this quantification for both the aggregate co-expression networks (diamond,

triangle, and square special characters, Supp. Fig. 4A) and for all individual primary tissue and

organoid co-expression networks (boxplots, Supp. Fig. 4A). Individual organoid networks

perform worse than individual primary tissue networks on average, with the aggregate

organoid network additionally underperforming compared to the aggregate primary tissue

networks, though within the range of individual primary tissue networks (Supp. Fig. 4A). In

aggregate, organoid co-expression weakly captures broad primary tissue cell-type specific co-

expression. This is potentially explained through the directed nature of the vast majority of

organoid datasets we investigate, which may more accurately produce particular lineages

(excitatory or inhibitory neurons as an example) rather than the comprehensive cell-

types/lineages present within primary tissue data. We explore cell-type specific co-expression

within individual datasets further in the following analysis.

**Organoid datasets vary in primary tissue cell-type marker set co-expression**

Having broadly assessed co-expression across our MetaMarker gene sets, we then

asked how well do organoids recapitulate primary tissue co-expression within each cell-type

specific MetaMarker gene set. We score intra-gene set co-expression strength through a

simple machine learning framework[45,46], which quantifies whether genes in a given set are

more strongly co-expressed with each other compared to the rest of the genome (Fig. 3C).

Co-expression module scores across the annotated and unannotated primary tissue

datasets are largely comparable with the exception of a sharp decrease in intermediate

progenitor performance for the unannotated primary tissue datasets (Fig. 3D). Six out of the

fourteen unannotated datasets are sampled from either the ganglionic eminences or the

hypothalamus, potentially explaining this decrease in performance and suggesting our

intermediate progenitor MetaMarkers are enriched for signal from cortical areas. In contrast,

performance is much more variable across the individual organoid datasets for all cell-types

except the dividing progenitors, ranging from no signal (AUROC <= 0.50) to comparable

results with primary tissue networks (Fig. 3D).

Importantly, the variation among organoid datasets for co-expression of the

differentiated cell-types is likely influenced via the compositional variation in cell-types

produced across directed and undirected differentiation protocols. A protocol that aims to

produce a directed excitatory lineage organoid is not expected to produce inhibitory cell-types

and thus should not necessarily exhibit strong inhibitory neuron co-expression. That is indeed

the case when comparing our co-expression module scores by organoid protocols (Supp. Fig.

4B). For example, the dorsal patterned forebrain organoid protocol produces stronger

excitatory module co-expression compared to inhibitory module co-expression.

In contrast, undirected organoid protocols are expected to produce a variety of

lineages/cell-types comparable to those present in primary tissue samples and should exhibit

consistent strong co-expression across cell-types. Instead, we report the undirected organoid

protocols (cortical, cerebral) as the more variable protocols for producing strong cell-type co-

expression (Supp. Fig. 4B). Visualizing the top and bottom performing cerebral organoid co-

expression networks for glutamatergic co-expression reveals the extent of this variability, in

comparison to the top performing primary tissue co-expression network (Fig. 3E). The top

performing organoid network produces near identical intra- and inter-cell type co-expression

relationships to the primary tissue dataset (Fig. 3E). Contrastingly, the bottom performing

organoid co-expression network exhibits extensive off-target inter-cell type co-expression and

extremely poor intra-cell type co-expression, essentially failing to recapitulate primary tissue

cell-type co-expression (Fig. 3E). While variability in co-expression performance may reflect

compositional differences among directed organoids, the dramatic range in performance

across the undirected organoid datasets reveals extensive variability in fidelity to primary

tissue.

**Organoid datasets vary in preserving gene-level primary tissue co-expression**

We take our primary tissue/organoid co-expression comparisons a step further and ask

how well individual organoid datasets preserve gene-level primary tissue co-expression

relationships. For any given individual gene, we quantify whether that gene's top co-expressed

partners are preserved in one co-expression network compared to another (Fig. 4A). We use

the aggregate co-expression network from the annotated primary tissue datasets as our

reference co-expression network and test how well individual co-expression networks, either

primary tissue or organoid, perform in preserving primary tissue gene-level co-expression

patterns (Fig. 4A, top 10 co-expressed neighbors). We start by quantifying the preserved co-

expression of genes within our primary tissue MetaMarker gene sets, using the average

preserved co-expression AUROC as a measure of preserved co-expression for any given

gene set (Fig. 4A). Across our 6 annotated primary tissue cell-types, primary tissue co-

expression networks deliver consistently high performance for preserved co-expression scores

of our primary tissue MetaMarker gene sets (Fig. 4B, mean preserved co-expression score

across cell-types and primary tissue datasets: annotated 0.970 ± 0.0241 SD, unannotated

0.963 ± 0.00940 SD). This indicates that across the highly temporally and regionally diverse

primary tissue data, the co-expression relationships of our MetaMarker gene sets are

incredibly highly preserved, again reflecting the temporally and regionally robust nature of our

primary tissue cell-type markers.

In contrast, individual organoid datasets vary substantially in preserved co-expression scores across our primary tissue MetaMarker gene sets (Fig. 4B). As before with our quantification of intra-gene set co-expression, compositional variation across directed organoid protocols may influence the variation in performance for differentiated cell-types, especially for excitatory and inhibitory neurons (Supp. Fig. 5). To explore the effects of likely compositional variation, we compared preserved co-expression scores of organoids grown in a vertical shaker versus an orbital shaker[47], where the original authors reported either a GABAergic or glutamatergic character for organoids grown in either a vertical or orbital shaker respectively. We report that organoids grown in an orbital shaker produce higher preserved primary tissue co-expression scores for intermediate progenitors and glutamatergic cell-types whereas organoids grown in a vertical shaker produce higher scores for GABAergic cell-types, in agreement with the authors original observations (3 replicates each, glutamatergic, intermediate progenitor, GABAergic; Orbital: 0.896 ± 0.0105 SD, 0.795 ± 0.00146 SD, 0.665 ± 0.0302 SD. Vertical: 0.644 ± 0.0126 SD, 0.686 ± 0.0167 SD, 0.762 ± 0.00589 SD). However, regardless of putative compositional variation across organoid protocols and/or treatments, we demonstrate that organoids from all sampled protocols consistently fail to preserve glutamatergic or GABAergic primary tissue co-expression at a level comparable to primary tissue (Supp. Fig. 5, Preserved Co-expression score ~0.90 or higher). This suggests a persistent remaining biological gap in the fidelity of organoid neurons in reference to primary tissue neurons. Organoids across protocols additionally exhibit near-zero preservation of non-neuronal primary tissue co-expression, suggesting organoids generally do not produce or produce extremely dysregulated non-neuronal cell-types (Fig. 4B, Supp. Fig. 5).

Additionally, we report extensive variability in the preservation of primary tissue

progenitor co-expression across organoid datasets. Again, regardless of the differentiation

protocol, organoids persistently fail to achieve comparable preserved co-expression of the

neural and intermediate progenitor MetaMarkers in reference to primary tissue data (Fig. 4B,

Supp. Fig. 5). In contrast to the differentiated cell-types, where compositional variation of

lineages across differentiation protocols may explain performance, progenitors are present

within every neural organoid. Variability in preserved co-expression of progenitor Meta-

Markers is likely a stronger reflection of the fidelity to primary tissue of any given organoid

dataset. Interestingly, a vascularized organoid protocol produces the highest preserved co-

expression of the neural and intermediate progenitors as well as the glutamatergic and

GABAergic cell-types. This suggests that vascularized organoids are particularly adept at

producing cell-types with high fidelity to primary tissue, but also that the preservation of co-

expression is associated across cell-types. We quantify this by computing correlations of

preserved co-expression scores between the 6 MetaMarker gene sets across all organoid

datasets and find significantly positive correlations (FDR-adjusted p-value < .001) across all

comparisons with the exception of the non-neuronal cell-type (Fig. 4C, non-neuronal FDR-

adjusted p-values range from 0.650 to 0.731). This indicates preserved primary tissue co-

expression is a global feature of organoid datasets. For example, if an organoid is producing

neural progenitors that preserve primary tissue co-expression, that organoid is likely producing

other cell-types that preserve primary tissue co-expression.

Neural organoids are commonly employed for the study of diverse disease mechanisms

through various perturbations. We tested the relevance of our preserved co-expression scores

for quantifying primary tissue fidelity across normal and perturbed organoids. A subset of our

organoid datasets come from studies that performed diverse perturbations (22q11.2 deletion,

SMARCB1 knockdown, exposure to Alzheimer's serum, SETBP1 point mutations, amyotrophic

lateral sclerosis patient-derived organoids). We compare the MetaMarker preserved co-

expression scores between normal and perturbed organoids and find only a single significant

difference across all cell-type MetaMarker sets (Intermediate Progenitor normal vs. mutant

preserved co-expression score FDR-adjusted p-value: 0.0295, Supp. Fig. 6A). This

demonstrates our broad primary tissue cell-type co-expression signatures are also applicable

for comparison with organoids in perturbation experiments.

**Fidelity of finer resolution cell-types through preserved co-expression**

While our broad cell-type annotations are useful for unifying meta-analysis across

heterogeneous primary tissue and organoid datasets, it is also of interest the degree neural

organoids are capable of producing primary tissue cell-types at a finer resolution. As our

approach for quantifying the preservation of co-expression is derived from a genome-wide co-

expression network of primary neural tissue, we can also putatively assess preserved co-

expression of more specific cell-type markers. We investigate preserved co-expression of

more specific cell-type markers by utilizing marker genes derived from a morphogen screen in

neural organoids that reported the production of extensive neural cell-type diversity[48]. As

examples of protocol specific trends, we show the dorsal patterned forebrain organoid

preserves co-expression of telencephalic excitatory neuron markers over markers for mib-brain

and thalamic excitatory neurons as well as dopaminergic mid-brain neurons (Fig. 4D).

Similarly, the ventral mid-brain organoid protocol, which reported production of dopaminergic

neurons, preserves co-expression of dopaminergic neuron markers over excitatory neuron

markers on average (Fig. 4D). Extending across all the organoid datasets, we demonstrate

preserved co-expression of fine resolution cell-types exhibit high correlations with the

preservation of our broader class-level markers for several Glutamatergic and GABAergic cell-

types (Fig. 4E). In summary, our results show that disruption of co-expression at one level of

cell-type hierarchy captures disruption at finer levels, suggesting a single score for organoid

fidelity can capture shared variation. More generally, our quantification for preserved co-

expression in organoids can also be applied to the study of finer resolution cell-types to study

variation from the shared baseline.

**Genome-wide preservation of co-expression reveal consistent organoid deficits**

After revealing cell-type specific variation for preserving primary tissue co-expression

within organoids, our co-expression networks additionally allow genome-wide assessments of

preserved co-expression. We extend our analysis via GO terms to quantify preserved primary

tissue co-expression within organoids across the whole genome. GO terms with significantly

preserved primary tissue co-expression (see methods) in organoids are mostly related to basic

cellular functions like response to DNA damage and protein translation, as well as GO terms

related to neurodevelopment (Fig. 4F). GO terms that significantly lack preservation of primary

tissue co-expression are largely related to angiogenesis or immune function (Fig. 4F),

concordant with the fact that organoids lack vasculature and an immune system. These results

demonstrate quantifications of preserved co-expression can capture known biological deficits

in neural organoids.

While GO terms are useful for partitioning the genome into functional units for

comparison, our co-expression networks also enable assessments of preserved co-expression

for individual genes. As a particular use-case, we search for genes with exceptionally high

preserved primary tissue co-expression across primary tissue datasets that also have poor

preserved primary tissue co-expression across organoid datasets. We only consider genes

that have some measurable expression in every organoid and primary tissue dataset and

compute the average preserved co-expression AUROC for each gene across the organoid and

primary tissue datasets (Supp. Fig. 6B). The top 10 enriched GO terms for genes (76 in total)

with high primary tissue (average AUROC >= 0.99) and low organoid (average AUROC < 0.70)

preserved co-expression are related to extra-cellular matrix (ECM) and vascular

characterizations (Fig. 4G). The poor conservation of genes related to vasculature can be

explained by the absence of vascularization in the vast majority of our organoid datasets. The

subset of these 76 genes in the ECM GO terms are CAV1, CAV2, COL4A1, CTSK, ENG,

LAMB1, LAMC1, NID1, NID2, DDR2, and VWA1. Notably, these genes produce collagen and

laminins, components of Matrigel, the artificial ECM typically included in organoid cultures.

These results highlight preserved primary tissue co-expression of ECM-related genes as a

particularly consistent deficit across neural organoids, suggesting that investigations into the

signaling between artificial ECM and cells in organoid cultures may be a route forward for

general improvements of organoid fidelity.

In summary, we interrogate co-expression in organoids at multiple levels, revealing

organoids vary in preserving primary tissue co-expression at gene-, cell-type, and whole

genome resolutions through the use of a robust aggregate primary tissue co-expression

network. We demonstrate the applicability of our approach for quantifying primary tissue fidelity

in organoids against a variety of use-cases, such as comparing different culture conditions

(vertical vs orbital shaking), comparing normal and perturbed organoids, and investigating

preserved co-expression of individual genes and fine resolution cell-type markers.

**Temporal variation in organoid preservation of primary tissue co-expression**

We score preserved co-expression in organoids using the aggregate primary tissue co-

expression network (GW5-25), which by design aims to capture signal robust to temporal

variation. To investigate temporal trends in organoid co-expression, we employ a similar

approach as when predicting organoid cell-type annotations (Supp. Fig. 3), this time

quantifying the preservation of primary tissue co-expression for the top 100 cell-type markers

per individual primary tissue dataset across all organoid timepoints (Fig. 5A-B). We uncover a

broad temporal shift in the preservation of primary tissue co-expression within organoids

across all cell-types, with younger organoids (23 days – 1.5 months) as the top performers for

mostly first trimester primary tissue co-expression transitioning to older organoids (2 – 6

months) as top performers for mostly second trimester primary tissue co-expression (Fig. 5B).

This temporal shift is broadly consistent across the cell-types, beginning around GW9-10 (Fig.

5B). Our approach in predicting organoid annotations in Figure 2 is based on aggregate

marker expression and did not produce temporally variable results, whereas our approach

here comparing preserved co-expression of the same marker genes does produce temporally

variable results. This indicates that the co-expression relationships of genes rather than their

expression levels better capture temporal variation in developing systems.

**Organoids preserve developing brain co-expression over adult brain co-expression**

We demonstrate temporal variation in developing brain co-expression relationships is

captured by organoids, but only from the single forebrain organoid protocol used in the

temporal organoid atlas. In order to extend analysis across all our organoid datasets and

assess broad temporal variation in co-expression, we next investigate the preserved co-

expression within organoids of both developing and adult brain co-expression relationships.

We construct an aggregate adult co-expression network from a medial temporal gyrus

scRNA-seq dataset[49]. We compare the preserved co-expression scores of organoids for either

developing or adult glutamatergic, GABAergic, and non-neuronal cell-types. Organoids

unanimously preserve developing brain co-expression over adult co-expression (Supp. Fig.

6C) for glutamatergic and GABAergic cell-types with equally poor performance for the non-

neuronal cells, again suggesting organoids generally fail to produce non-neuronal cell-types.

We extend this analysis genome-wide and place organoids in context between developing and

adult data by computing the average preservation of co-expression AUROC across all genes

for organoid, developing, and adult co-expression using the annotated primary developing

brain tissue network as the reference. The adult co-expression network produces a global

preserved developing brain co-expression score of 0.591, indicating very poor performance

across the genome in preserving developing co-expression relationships (Supp. Fig. 6D).

Organoids vary substantially in their global preservation of developing brain co-expression with

some organoid datasets performing comparably to the adult data. This result is largely

influenced by the number of cells present within individual organoid datasets (Supp. Fig. 6D,

corr 0.647, p-value < .001), suggesting a cell-sampling limitation for uncovering developing

brain co-expression within organoids. However, organoid datasets report more variable global

preserved co-expression scores compared to down-sampled developing brain data (Supp. Fig.

6D), indicating a remaining gap between primary developing brain tissue and organoid data

not explained through cell number sampling alone.

We further explore the applicability of our preserved co-expression quantifications for

investigating temporal variation through a study that tested the limits of neuronal maturation in

organoids. This study generated data from human cortical organoids either transplanted or not

into developing rat brains to test the limits of maturation organoids can achieve *in vitro*[50]. We

compare the preservation of developing and adult co-expression between these age-matched

non-transplanted and transplanted human cortical organoids. We report that while the non-

transplanted organoids preserve developing co-expression over adult for glutamatergic and

GABAergic markers (Supp. Fig. 6E, non-transplanted glutamatergic and GABAergic mean

developing brain AUROCs: 0.798 ± 0.0278 SD, 0.698 ± 0.0208 SD. Non-transplanted

glutamatergic and GABAergic mean adult AUROCs: 0.672 ± 0.0234 SD, 0.585 ± 0.0291 SD),

the transplanted organoids have increased preservation of adult co-expression for

glutamatergic markers (Supp. Fig. 6E, transplanted glutamatergic mean developing brain

AUROCs: 0.759 ± 0.00909 SD. Transplanted glutamatergic mean adult AUROCs: 0.850 ±

0.0332 SD). This indicates the transplanted human organoids are adopting adult human

glutamatergic co-expression, concordant with the original authors' conclusions of increased

maturation in transplanted organoids. The transplanted organoids additionally report increased

preservation of both developing and adult non-neuronal marker co-expression, in agreement

with the original authors' observations of oligodendrocytes within transplanted organoids. By

recapitulating known maturation dynamics in organoid models, we demonstrate the broad

applicability of preserved co-expression quantifications for investigating a range of biological

phenomenon in neural organoids.

**Variability in organoid co-expression is driven by marker gene expression**

We investigate the impact of various technical features in our analysis on our co-

expression results by assessing their correlation with our co-expression module scores and

preserved co-expression AUROCs, focusing on technical features like sequencing depth,

number of cells, etc. An important technical consideration for our analysis is ensuring all

datasets have an identical gene namespace for meaningful comparisons of expression data.

We fit all datasets to the GO gene universe, dropping gene annotations not in GO or zero-

padding missing GO annotations in individual datasets. Excessive zero-padding of genes

within our MetaMarker gene sets may artificially lower co-expression module scores or

preserved co-expression scores, though we find this relationship to be relatively weak with little

impact on score variance (Supp. Fig. 7, $R^2$ for co-expression module scores and zero-padding:

0.00267, 0.0165, 0.126, 0.0261, 0.0354, 0.00451, $R^2$ for preserved co-expression and zero-

padding: 0.0665, 0.322, 0.151, 0.0307, 0.0411, 0.00203 for neural prog., dividing prog.,

intermediate prog., glutamatergic, GABAergic, and non-neuronal cell-types respectively).

Sequencing depth is also similarly found to have little impact on our co-expression module

scores or preserved co-expression scores (Supp. Fig. 7). Rather, the features strongly related

to performance are the number of cells in a dataset and the strength of marker set expression

(Supp. Fig. 7, range of significant positive (p-value < .05) correlations between marker set

expression or cell number and co-expression module scores or preserved co-expression

scores: 0.204 – 0.809).

**Preservation of primary tissue co-expression as a generalizable quality control metric**

As a general summary, our approach for quantifying preserved primary tissue co-

expression across numerous organoid protocols reveal the axes on which organoids lie for

recapitulating primary tissue co-expression relationships at gene, cell-type, and whole-genome

resolutions. These assessments provide powerful quality control information, identifying which

genes and/or cell-types organoids can or cannot currently model on par with primary tissue

data. We make our methods accessible through an R package to aid in future organoid studies

and protocol development, providing means for rapidly constructing co-expression networks

from scRNA-seq data (Fig. 6A) as well as querying preserved co-expression of users' data

with our aggregate primary tissue brain co-expression network (Fig. 6A). Additionally, we make

the results of our meta-analysis across primary tissue and organoid datasets available for

users to place their data in reference to a field-wide collection (Fig. 6B).

**Discussion**

Through the use of meta-analytic differential expression and co-expression, we are able to

provide cell-type specific measurements of human neural organoids' current capacity to

replicate primary tissue biology. We extracted broad cell-type markers that define primary brain

tissue cell-types across a large temporal axis (GW5 – 25) and across numerous heterogenous

brain regions to act as a generalizable primary tissue reference for organoids that also vary

temporally and regionally (by protocol). By quantifying intra-marker set co-expression and the

preservation of co-expression across networks, we revealed human neural organoids lie on a

spectrum of near-zero to near-identical recapitulation of primary tissue cell-type specific co-

expression in comparison to primary tissue data. We made our aggregate primary tissue

reference data and methods for measuring preserved co-expression publicly available as an R

package to aid in the quality control and protocol development of future human neural

organoids.

Prior work comparing primary brain tissue and neural organoid systems demonstrated

organoids can produce cell-types[11,12] and morphological structures[27,43] similar to primary

tissues and are capable of modeling temporal[13,38,40] and regional[3,12,28,29] primary tissue

variation. Multiple lines of evidence support these findings such as assessments of

cytoarchitecture and cell-type proportions[3,11,16,23], whole transcriptome and marker gene

expression correlations[10,12], and comparisons of co-expression modules[5,13,17,39]. Our meta-

analytic approach is able to quantify these field-wide observations within a generalizable

framework, recapitulating that organoids model broad primary tissue biology with our specific

approach offering several key advancements for primary tissue/organoid comparisons. First,

we derive quantifications of preserved primary tissue co-expression that can be extended from

individual genes to the entire genome and, second, we place organoid co-expression in

reference to robust meta-analytic primary tissue performance providing a general benchmark

for protocol development and quality control across heterogeneous organoid systems.

Certainly, while comparisons between primary tissue and organoid systems at a high-

resolution of cell-type annotation are of interest, our results centered on broad cell-types at the

cell-class level constitute a critical foundation for these more fine-tuned investigations of

organoids. Cell-type specification within the brain involves complex spatial and temporal

mechanisms[51] to produce the high cellular heterogeneity we observe, with the exact resolution

of meaningful cell-type annotations still being actively debated and posing a general

conceptual challenge within the field of single-cell genomics[52]. We focus here on establishing

methods for assessing consistent and accurate production of primary tissue cell-types at the

class-level within organoids as a critical actionable first step towards increasing primary tissue

fidelity across variable organoid differentiation protocols. While we prioritize broad cell-type

comparisons, we also display the flexibility of our approach by scoring the preserved co-

expression of finer resolution cell-type markers. This demonstrates our quantifications of

preserved co-expression are applicable to a variety of cell-type annotation resolutions.

One exciting application for the use of neural organoid systems is the study of a wide-range

of human neurological diseases using human *in vitro* models[53,54], which critically depends on

the *in vivo* fidelity of cell-types produced in organoids. Neural organoids have been used to

model and investigate human disorders of neurodevelopmental[3,55], neuropsychiatric[56–58], and

neurodegenerative[59–61] nature, as well as infectious diseases[28,62,63]. It is essential that

organoid systems model *in vivo* cell-types with extreme fidelity to fully realize the therapeutic

potential of human organoids and ensure findings in these *in vitro* models are not specific to

potential artifactual or inaccurate *in vitro* biology. While our results demonstrate that high

primary tissue fidelity in organoids is currently methodologically possible, we also report a high

degree of variability across organoids and studies/protocols indicating a remaining

methodological gap. The broad applicability of our meta-analytic approach offers the potential

for benchmarking primary tissue fidelity across numerous organoid protocols, aiding in

increasing the quality of neural organoids for use in a wide-range of human health-related

translational investigations.

The generalizable and flexible nature of our analysis is well suited to aid in the

development of organoid differentiation protocols and the general quality control of neural

organoids. Our results demonstrate the type of experiments possible through comparing

preserved co-expression across organoid experimental variables, such as the differences in

preserved co-expression between organoids grown in vertical or orbital shakers, as well as

between transplanted or non-transplanted organoids. Importantly, our broad sampling across

organoid protocols enabled clear identification of promising avenues for increasing organoid

primary tissue fidelity. The strong performance across cell-types for the vascularized protocol

we assessed suggests vascularized protocols as a route forward for global increases in

primary fidelity. Additionally, our findings of specific ECM-related genes with consistent poorly

preserved primary tissue co-expression in organoids suggests investigations into the

interactions between Matrigel or other ECM-substrates and organoids may lead to general

protocol adjustments for increasing primary tissue fidelity[64]. Looking beyond neural organoids,

our framework for quantifying preserved co-expression can be applied to other organoid

systems granted there is sufficient annotated primary tissue data to act as a reference.

**Methods**

**Dataset download and scRNA-seq pre-processing**

Links for all downloaded data (GEO accession numbers, data repositories, etc.) are

provided in Supp. Table 1. All scRNA-seq data was processed using the Seurat v4.2.0 R

package[65]. Data made available in 10XGenomics format (barcodes.tsv.gz, features.tsv.gz,

matrix.mtx.gz) were converted into Seurat objects using the Read10X() and

CreateSeuratObject() Seurat functions. Data made available as expression matrices were

converted into sparse matrices and then converted into Seurat objects using the

CreateSeuratObject() function. Ensembl gene IDs were converted into gene names using the

biomaRt v2.52.0[66] package.

Where metadata was made available, we separated data by batch (Age, Donor, Cell

line, etc.) for our final total of 130 organoid and 51 primary tissue datasets (Supp. Table 1). We

processed and analyzed each batch independently without integration. We used consistent

thresholds for filtering cells across all datasets, keeping cells that had less than 50% of reads

mapping to mitochondrial genes and had between 200 and 6000 detected genes. Several

datasets provided annotations for potential doublets; we excluded all cells labeled as doublets

when annotations were made available. All data made available with raw expression counts

were CPM normalized with NormalizeData(normalization.method = 'RC', scale.factor = 1e6),

otherwise normalizations were kept as author supplied.

For primary tissue and organoid data made available with cell-type annotations, we

provide our mapping between author provided annotations and our broad cell-type annotations

in Supp. Table 2. Vascular annotated cell-types were excluded.

**Primary tissue MetaMarker generation and cross-validation**

MetaMarkers were computed using the MetaMarkers v0.0.1[44] R package, which

requires shared cell-type and gene annotations across datasets to derive a ranked list of

MetaMarkers. Gene markers for individual datasets were first computed using the

compute_markers() function on the CPM normalized expression data for our annotated

primary tissue datasets (Supp. Table. 1). A ranked list of MetaMarkers was then computed

using the make_meta_markers() function using all 37 individual annotated primary tissue

dataset marker lists. Genes are first ranked through their recurrent differential expression (the

number of datasets that gene was called as DE using a threshold of log2 FC >= 4 and FDR p-

value <= .05) and then through the averaged differential expression statistics of each gene

across individual datasets. When we take the top 100 markers per individual dataset as in Fig.

2D, Fig. 5, Supp. Fig. 1A, and Supp. Fig. 3B, we rank markers for each dataset by their

AUROC statistic as computed with the compute_markers() MetaMarkers function.

For the cross-validation of our primary tissue MetaMarkers, we excluded a single

annotated primary tissue dataset, computed MetaMarkers from the remaining 36 annotated

primary tissue datasets, and then used those MetaMarkers to predict the cell-type annotations

of the left-out dataset. We construct an aggregate expression predictor to quantify the

predictive strength a list of genes has, in this case our MetaMarker lists, in predicting cell-type

annotations. Taking any arbitrary number of genes (10, 20, 50, 100, 250, or 500 MetaMarkers),

we sum the expression counts for those genes within each cell and then rank all cells by this

aggregate expression vector. We compute an AUROC using this ranking and the cell-type

annotations for a particular cell-type through the Mann-Whitney U test. Formally:

$$AUROC = \frac{U}{n_0 * n_1}$$

643      where U is the Mann-Whitney U test statistic, $n_0$ is the number of positives (cells with a

644    given cell-type annotation) and $n_1$ is the number of negatives (cells without that cell-type

645    annotation).

$$U = R_0 - \frac{n_0(n_0 + 1)}{2}$$

where $R_0$ is the sum of the positive ranks.

As an example, if there are 10 genes that are perfect glutamatergic markers (only

glutamatergic cells express these genes), then ranking cells by the summed expression of

these genes will place all glutamatergic cells (positives) in front of all other cells (negatives),

producing an AUROC of 1. The violin plots in Supp. Fig. 1B and in Figure 2E visualize our

aggregate expression approach, where datapoints per cell-type are the aggregated expression

counts for the given top 100 MetaMarkers across all cells per dataset (Supp. Fig. 1B) or

aggregated across all datasets (Fig. 2E). We also compared the aggregate expression of the

Neural Progenitor MetaMarkers across author provided cell-type annotations included in our

broad Non-neuronal annotation, revealing the off-target expression of Neural Progenitor

MetaMarkers is specific to annotated astrocytes (Supp. Fig. 1B).

For Supp. Fig. 1A, we took the top 100 cell-type markers per individual primary tissue

dataset (x-axis) and used those genes to predict cell-type annotations as described above for

all other annotated primary tissue datasets, reported as the AUROC boxplot distributions. The

MetaMarker distribution was computed using a leave-one-out approach as described above.

We ranked the individual primary tissue datasets by their median AUROC performance per

cell-type to derive the distributions of ranks presented in Figure 2D, excluding the dividing

progenitor data as performance was highly consistent across all primary tissue datasets.

**Cross-regional primary tissue MetaMarker expression**

We investigated the aggregate expression of our top 100 MetaMarkers per cell-type

across annotated brain regions separately for the annotated first-trimester and second-

trimester primary tissue atlases due to differing regional annotations. MetaMarkers were

computed with a leave-one-out approach as described above using all 37 of the annotated

primary tissue datasets. For the heatmaps in Supp. Fig. 2, rows represent the annotated cells

present within the given dataset, columns represent the aggregated expression for the top 100

given cell-type MetaMarkers and each annotated region present. We average the aggregated

expression for each cell-type per region and then normalize each region (column) by the

maximum average expression value across the cell-types. A value of 1 indicates that cell-type

is the one maximally expressing the given MetaMarker set for that brain region. The heatmaps

are ordered by cell-type and region and are not clustered.

**Organoid PCA**

PCA analysis was performed using the Seurat function RunPCA() with the top 2000

variable features, determined using the Seurat function

FindVariableFeatures(selection.method = 'vst', nfeatures = 2000). For each organoid dataset,

we took the eigenvector for the first principal component, computed the absolute value, and

then divided by the maximum value to compute a normalized vector between 0 and 1. We

visualized the normalized eigenvectors for each organoid dataset in Supp. Fig. 3C, keeping

primary tissue MetaMarker genes that were detected in the top 2000 variable genes of at least

10 organoid datasets. Genes missing from any given dataset's top 2000 variable genes were

given a value of 0. The heatmap was produced using the ComplexHeatmap v2.12.1[67] package

and was hierarchically clustered using the ward.D2 method for both rows and columns.

**Generating co-expression networks from scRNA-seq data**

To generate a shared gene annotation space across all datasets, we fit each dataset to

the GO gene universe before computing co-expression matrices. Using human GO

annotations (sourced 2023-01-01 using the org.Hs.eg.db v3.15.0[68] and AnnotationDbi

v1.58.0[69] R packages), we excluded gene expression from a dataset if the gene annotation

was not present in GO and we zero-padded missing GO genes for each dataset.

We compute a gene-by-gene co-expression matrix per dataset using the spearman

correlation coefficient computed across all cells in a given dataset. We then rank the

correlation coefficients in the gene-by-gene matrix and divide by the maximum rank to obtain a

rank-standardized co-expression matrix. All results reported using individual dataset co-

expression networks (Fig. 3D-E, Fig. 4B, Figs. 5-6, Supp. Figs. 4-7) were obtained using the

rank-standardized co-expression networks.

We compute the aggregated co-expression networks by taking the average of the rank

standardized co-expression networks for each gene-gene index.

**Hierarchical clustering of primary tissue MetaMarkers by co-expression**

707  We visualize the co-expression of primary tissue MetaMarker genes using the

708 ComplexHeatmap package and the ward.D2 algorithm for hierarchical clustering. We use the

709 fossil v0.4.0 package[70] to compute the adjusted Rands Index with the adj.rand.index() function.

710 To compute the adjusted Rands Index, we calculate a consensus clustering of MetaMarkers

711 per co-expression network across 100 k-means clusterings (using the arguments row_km = 6,

712 column_km = 6, row_km_repeats = 100, column_km_repeats = 100 within the Heatmap

713 function) to compare to the perfect grouping of MetaMarkers by cell-type.

714  For the heatmaps in Fig. 3E, genes are ordered within each MetaMarker gene set by

715 their average intra-gene set co-expression.

**717 Co-expression module learning analysis**

718  EGAD v1.24.0[45] is a machine learning framework that quantifies the strength of co-

719 expression within an arbitrary gene-set compared to the rest of the genome with an AUROC

720 quantification (Fig. 3C). We compute co-expression module AUROCs for all GO gene-sets

721 (between 10 and 1000 genes per GO term) and our top 100 primary tissue MetaMarker gene-

722 sets for each individual primary tissue and organoid co-expression network as well as the

723 aggregated annotated, unannotated and organoid networks. For the annotated primary tissue

724 co-expression networks, we employ a leave-one-out approach, learning MetaMarkers from 36

725 of the annotated datasets and computing co-expression module AUROCs for these

726 MetaMarkers in the left-out dataset's co-expression network. We compute co-expression

727 module AUROCs using the EGAD run_GBA() function with default parameters. In Figure 3D,

728 the 'All GO terms' distributions report the average co-expression module AUROC across all

729 GO terms for each individual network.

**Preservation of co-expression**

To compute our preservation of co-expression AUROC, we take the top 10 co-expressed partners for gene A in a reference co-expression network as our positive gene annotations. In a test co-expression network, we rank all genes through their co-expression with gene A and compute an AUROC using this ranking and the positive annotations derived from the reference network. If gene A in the test network has the exact same top 10 co-expressed partners as in the reference network, that would result in an AUROC of 1. To summarize a given gene-set's preserved co-expression, we take the average preserved co-expression AUROC across all genes in that gene set as the preservation of co-expression score for that gene set. We use the aggregated annotated primary tissue co-expression matrix as our reference network.

The preserved co-expression scores for the annotated primary tissue data in Figure 4B were computed with a leave-one-out approach. MetaMarkers and an aggregated co-expression matrix were computed from 36 of the annotated primary tissue datasets and then preserved co-expression scores were computed using the co-expression network of the left-out annotated primary tissue dataset.

**Preservation of fine resolution cell-types**

To define markers for finer resolution cell-types, we utilize the differential expression (DE) statistics computed from a study that performed a morphogen screen in neural organoids and reported extensive neural cell-type diversity[48]. For each cell-type, we rank genes by their adjusted DE p-value and take the top 10 genes per cell-type to compute preserved co-

expression scores. When comparing against our MetaMarker gene sets in Figure 4E, we

ensure no overlap in the top 10 cell-type and top 100 MetaMarker gene sets.

**Preservation of GO term co-expression**

We compute p-values for the preservation of co-expression of GO terms using a mean

sample error approach. Using the aggregated annotated primary tissue co-expression network

as the reference and the aggregated organoid network as the test network, we first compute

the preserved co-expression AUROCs for all individual genes, taking the mean and standard

deviation value as the population mean and population standard deviation. For any given GO

term, we first compute the preserved co-expression score for the term (the average of the

preserved co-expression AUROCs for the genes in the term) and then compute the sample

error for that score with:

$$SE = \frac{SD_{pop}}{\sqrt{n_{GO}}}$$

765    where $SD_{pop}$ is the population standard deviation and $n_{GO}$ is the number of genes in the GO

766    term. We then compute a z-score through:

$$Z_{GO} = \frac{mu_{GO} - mu_{pop}}{SE}$$

767    where $mu_{go}$ is the preserved co-expression score for the GO term and $mu_{pop}$ is the population

768    mean preserved co-expression AUROC. We compute left-sided p-values using the standard

769    normal distribution:

$$p_L = P(X \leq Z_{GO})$$

771   Where X is a normal distribution with mean = 0 and standard deviation = 1. We use the R

772   function pnorm($Z_{GO}$) to compute this p-value.

773   We then compute the right-sided p-value as:

$$p_R = 1 - p_L$$

We adjust p-values using the R function p.adjust(method = 'BH'). We filter for GO terms

that have between 20 and 250 genes per term and use a threshold of FDR-corrected p-value

<= 0.0001 to call significance. Significant left-sided p-values are interpreted as GO terms with

significantly smaller preserved co-expression scores (significantly not preserved) than

expected through sampling error and right-sided p-values are interpreted as GO terms with

significantly larger preserved co-expression scores (significantly preserved) than expected

through sampling error. We use the R package rrvgo to visualize the significant GO terms in

Fig. 4F.

**Computing correlation significance**

We employ a permutation test to compute p-values for any given correlation coefficient.

We permute data-pairs and compute a correlation coefficient, repeating for 10,000 random

permutations to generate a distribution of correlation coefficients under the null hypothesis of

independence. We calculate a two-sided p-value for the original correlation coefficient as the

number of permuted correlation coefficients whose absolute value is greater than or equal to

the absolute value of the original correlation coefficient, divided by 10,000. We adjust p-values

using the R function p.adjust(method = 'BH') and use a FDR-corrected p-value threshold of <=

.05 to call significance.

**Comparing co-expression of normal vs. perturbed organoids**

For both the co-expression module AUROCs and the preserved co-expression scores of normal and perturbed organoids, we test for significant differences per cell-type using the Mann Whitney U test, adjusting p-values with the R function p.adjust(method = 'BH') and using a FDR-corrected p-value threshold of <= .05 to call significance.

**Organoid temporal analysis**

The organoid temporal analysis for both predicting organoid annotations with primary tissue markers (Supp. Fig. 3B) and scoring the preserved co-expression of organoid co-expression using primary tissue networks as reference (Fig. 5) were performed for all pair-wise combinations of the 37 annotated primary tissue datasets and the 26 temporally annotated forebrain organoid datasets. We excluded the GW7-28 annotated primary tissue dataset from the temporal preserved co-expression analysis (Fig. 5) due to the wide temporal range sampled. For predicting organoid annotations with primary tissue markers, we used the top 100 markers per primary tissue dataset to construct aggregate expression predictors in the organoid datasets as described above. The MetaMarkers performance was calculated using MetaMarkers derived from all 37 annotated primary tissue datasets. For scoring preserved co-expression, individual primary tissue networks were used as the reference with individual organoid networks as the test networks. We computed the preserved co-expression scores of the top 100 primary tissue cell-type markers per individual primary dataset for each individual organoid network.

**GO enrichment analysis**

We compute enrichment for GO terms using Fisher's Exact Test as implemented

through the hypergeometric test. We compute raw p-values for GO terms with between 10-

1000 genes and compute FDR-adjusted p-values using p.adjust(method = 'BH'). We only

consider GO sets with between 20 and 500 when choosing the top 10 GO sets in Figure 4G,

ranked by FDR-adjusted p-value.

**R and R packages**

All analysis was carried out in R v4.3.1. Colors with selected using the MetBrewer

v0.2.0 R library. Plots were generated using ggplot2 v3.3.6[71]. Spearman correlation matrices

for co-expression networks were computed using a python v3.6.8 script, implemented in R with

the reticulate v1.26 R package, as well as using functions from the matrixStats v0.62.0 R

library. All code used in generating results and visualizations will be made public at the time of

publication. The preservedCoexp R library is made available at

https://github.com/JonathanMWerner/preservedCoexp. All code used for analysis is made

available at https://github.com/JonathanMWerner/meta_organoid_analysis.

**Author Contributions**

JMW and JG conceived the project. JMW and JG designed analyses. JMW performed

analyses. JMW and JG wrote the manuscript.

**Acknowledgments**

JMW was supported by NSF award no. DGE-1938105. JG and JMW were supported by NIH

grants R01MH113005 and R01LM012736. This material is based upon work supported by the

National Science Foundation Graduate Research Fellowship Program under grant no. DGE-

1938105. Any opinions, findings, and conclusions or recommendations expressed in this

material are those of the author(s) and do not necessarily reflect the views of the National

Science Foundation. We thank Tomasz Nowakowski and members of the Gillis lab for helpful

comments on the manuscript.

## References

1. Eiraku, M. *et al.* Self-Organized Formation of Polarized Cortical Tissues from ESCs and Its Active

Manipulation by Extrinsic Signals. *Cell Stem Cell* **3**, 519–532 (2008).

2. Sato, T. *et al.* Single Lgr5 stem cells build crypt-villus structures in vitro without a mesenchymal

niche. *Nature* **459**, 262–265 (2009).

3. Lancaster, M. A. *et al.* Cerebral organoids model human brain development and microcephaly.

*Nature* **501**, 373–379 (2013).

4. Corrò, C., Novellasdemunt, L. & Li, V. S. W. A brief history of organoids. *Am. J. Physiol.-Cell Physiol.*

**319**, C151–C165 (2020).

5. Pollen, A. A. *et al.* Establishing Cerebral Organoids as Models of Human-Specific Brain Evolution. *Cell*

**176**, 743-756.e17 (2019).

6. Kanton, S. *et al.* Organoid single-cell genomic atlas uncovers human-specific features of brain

development. *Nature* **574**, 418–422 (2019).

7. Benito-Kwiecinski, S. *et al.* An early cell shape transition drives evolutionary expansion of the human

forebrain. *Cell* **184**, 2084-2102.e19 (2021).

8. Fleck, J. S. *et al.* Inferring and perturbing cell fate regulomes in human brain organoids. *Nature* 1–8

(2022) doi:10.1038/s41586-022-05279-8.

9. Sarieva, K. & Mayer, S. The Effects of Environmental Adversities on Human Neocortical

Neurogenesis Modeled in Brain Organoids. *Front. Mol. Biosci.* **8**, (2021).

10.    Camp, J. G. *et al.* Human cerebral organoids recapitulate gene expression programs of fetal

neocortex development. *Proc. Natl. Acad. Sci.* **112**, 15672–15677 (2015).

11.    Velasco, S. *et al.* Individual brain organoids reproducibly form cell diversity of the human

cerebral cortex. *Nature* **570**, 523–527 (2019).

12.    Bhaduri, A. *et al.* Cell stress in cortical organoids impairs molecular subtype specification.

*Nature* **578**, 142–148 (2020).

13.    Gordon, A. *et al.* Long-term maturation of human cortical organoids matches key early

postnatal transitions. *Nat. Neurosci.* **24**, 331–342 (2021).

14.    Feng, W. *et al.* Computational profiling of hiPSC-derived heart organoids reveals chamber

defects associated with NKX2-5 deficiency. *Commun. Biol.* **5**, 1–18 (2022).

15.    Leek, J. T. *et al.* Tackling the widespread and critical impact of batch effects in high-throughput

data. *Nat. Rev. Genet.* **11**, 733–739 (2010).

16.    Tanaka, Y., Cakir, B., Xiang, Y., Sullivan, G. J. & Park, I.-H. Synthetic Analyses of Single-Cell

Transcriptomes from Multiple Brain Organoids and Fetal Brain. *Cell Rep.* **30**, 1682-1689.e3 (2020).

17.  Cheroni, C. *et al.* Benchmarking brain organoid recapitulation of fetal corticogenesis. *Transl.*

*Psychiatry* **12**, 1–16 (2022).

18.  Kim, H. J. *et al.* Comprehensive characterization of fetal and mature retinal cell identity to

assess the fidelity of retinal organoids. *Stem Cell Rep.* **18**, 175–189 (2023).

19.  Zhang, B. & Horvath, S. A General Framework for Weighted Gene Co-Expression Network

Analysis. *Stat. Appl. Genet. Mol. Biol.* **4**, (2005).

20.  Lee, J., Shah, M., Ballouz, S., Crow, M. & Gillis, J. CoCoCoNet: conserved and comparative co-

expression across a diverse set of species. *Nucleic Acids Res.* **48**, W566–W571 (2020).

21.  Crow, M., Paul, A., Ballouz, S., Huang, Z. J. & Gillis, J. Exploiting single-cell expression to

characterize co-expression replicability. *Genome Biol.* **17**, 101 (2016).

22.  Mead, B. E. *et al.* Harnessing single-cell genomics to improve the physiological fidelity of

organoid-derived cell types. *BMC Biol.* **16**, 62 (2018).

23.  Agboola, O. S., Hu, X., Shan, Z., Wu, Y. & Lei, L. Brain organoid: a 3D technology for investigating

cellular composition and interactions in human neurological development and disease models in

vitro. *Stem Cell Res. Ther.* **12**, 430 (2021).

24.  Mayhew, C. N. & Singhania, R. A review of protocols for brain organoids and applications for

disease modeling. *STAR Protoc.* **4**, 101860 (2023).

25.  Lancaster, M. A. & Knoblich, J. A. Generation of cerebral organoids from human pluripotent

stem cells. *Nat. Protoc.* **9**, 2329–2340 (2014).

26.   Muguruma, K., Nishiyama, A., Kawakami, H., Hashimoto, K. & Sasai, Y. Self-Organization of

Polarized Cerebellar Tissue in 3D Culture of Human Pluripotent Stem Cells. *Cell Rep.* **10**, 537–550

(2015).

27.   Sakaguchi, H. *et al.* Generation of functional hippocampal neurons from self-organizing human

embryonic stem cell-derived dorsomedial telencephalic tissue. *Nat. Commun.* **6**, 8896 (2015).

28.   Qian, X. *et al.* Brain-Region-Specific Organoids Using Mini-bioreactors for Modeling ZIKV

Exposure. *Cell* **165**, 1238–1254 (2016).

29.   Xiang, Y. *et al.* Fusion of Regionally Specified hPSC-Derived Organoids Models Human Brain

Development and Interneuron Migration. *Cell Stem Cell* **21**, 383-398.e7 (2017).

30.   Birey, F. *et al.* Assembly of functionally integrated human forebrain spheroids. *Nature* **545**, 54–

59 (2017).

31.   Xiang, Y. *et al.* hESC-Derived Thalamic Organoids Form Reciprocal Projections When Fused with

Cortical Organoids. *Cell Stem Cell* **24**, 487-497.e7 (2019).

32.   Miura, Y. *et al.* Generation of human striatal organoids and cortico-striatal assembloids from

human pluripotent stem cells. *Nat. Biotechnol.* **38**, 1421–1430 (2020).

33.   Eura, N. *et al.* Brainstem Organoids From Human Pluripotent Stem Cells. *Front. Neurosci.* **14**,

(2020).

34.   Andersen, J. *et al.* Generation of Functional Human 3D Cortico-Motor Assembloids. *Cell* **183**,

1913-1929.e26 (2020).

35.   Huang, W.-K. *et al.* Generation of hypothalamic arcuate organoids from human induced

pluripotent stem cells. *Cell Stem Cell* **28**, 1657-1670.e10 (2021).

36.      Nayler, S., Agarwal, D., Curion, F., Bowden, R. & Becker, E. B. E. High-resolution transcriptional

landscape of xeno-free human induced pluripotent stem cell-derived cerebellar organoids. *Sci. Rep.*

**11**, 12959 (2021).

37.      Sozzi, E., Nilsson, F., Kajtez, J., Parmar, M. & Fiorenzano, A. Generation of Human Ventral

Midbrain Organoids Derived from Pluripotent Stem Cells. *Curr. Protoc.* **2**, e555 (2022).

38.      Uzquiano, A. *et al.* Proper acquisition of cell class identity in organoids allows definition of fate

specification programs of the human cerebral cortex. *Cell* **185**, 3770-3788.e27 (2022).

39.      Luo, C. *et al.* Cerebral Organoids Recapitulate Epigenomic Signatures of the Human Fetal Brain.

*Cell Rep.* **17**, 3369–3384 (2016).

40.      Amiri, A. *et al.* Transcriptome and epigenome landscape of human cortical development

modeled in brain organoids. *Science* **362**, eaat6720 (2018).

41.      Fair, S. R. *et al.* Electrophysiological Maturation of Cerebral Organoids Correlates with Dynamic

Morphological and Cellular Development. *Stem Cell Rep.* **15**, 855–868 (2020).

42.      Nascimento, J. M. *et al.* Human Cerebral Organoids and Fetal Brain Tissue Share Proteomic

Similarities. *Front. Cell Dev. Biol.* **7**, 303 (2019).

43.      Qian, X. *et al.* Sliced Human Cortical Organoids for Modeling Distinct Cortical Layer Formation.

*Cell Stem Cell* **26**, 766-781.e9 (2020).

44.      Fischer, S. & Gillis, J. How many markers are needed to robustly determine a cell's type?

*iScience* **24**, 103292 (2021).

45.      Ballouz, S., Weber, M., Pavlidis, P. & Gillis, J. EGAD: ultra-fast functional analysis of gene

networks. *Bioinformatics* **33**, 612–614 (2017).

46.    Skinnider, M. A., Squair, J. W. & Foster, L. J. Evaluating measures of association for single-cell

940           transcriptomics. *Nat. Methods* **16**, 381–386 (2019).

941    47.    Suong, D. N. A. *et al.* Induction of inverted morphology in brain organoids by vertical-mixing

942           bioreactors. *Commun. Biol.* **4**, 1–13 (2021).

943    48.    Amin, N. D. *et al.* Generating human neural diversity with a multiplexed morphogen screen in

944           organoids. 2023.05.31.541819 Preprint at https://doi.org/10.1101/2023.05.31.541819 (2023).

945    49.    Jorstad, N. L. *et al.* Comparative transcriptomics reveals human-specific cortical features.

946           *Science* **382**, eade9516 (2023).

947    50.    Revah, O. *et al.* Maturation and circuit integration of transplanted human cortical organoids.

948           *Nature* **610**, 319–326 (2022).

949    51.    Nowakowski, T. J. *et al.* Spatiotemporal gene expression trajectories reveal developmental

950           hierarchies of the human cortex. *Science* **358**, 1318–1323 (2017).

951    52.    Zeng, H. What is a cell type and how to define it? *Cell* **185**, 2739–2755 (2022).

952    53.    Chen, H. I., Song, H. & Ming, G. Applications of Human Brain Organoids to Clinical Problems.

953           *Dev. Dyn.* **248**, 53–64 (2019).

954    54.    Eichmüller, O. L. & Knoblich, J. A. Human cerebral organoids — a new tool for clinical neurology

955           research. *Nat. Rev. Neurol.* **18**, 661–680 (2022).

956    55.    Mariani, J. *et al.* FOXG1-Dependent Dysregulation of GABA/Glutamate Neuron Differentiation in

957           Autism Spectrum Disorders. *Cell* **162**, 375–390 (2015).

56.    Notaras, M. *et al.* Schizophrenia is defined by cell-specific neuropathology and multiple

neurodevelopmental mechanisms in patient-derived cerebral organoids. *Mol. Psychiatry* **27**, 1416–

1434 (2022).

57.    Stachowiak, E. K. *et al.* Cerebral organoids reveal early cortical maldevelopment in

schizophrenia—computational anatomy and genomics, role of FGFR1. *Transl. Psychiatry* **7**, 1–24

(2017).

58.    Dixon, T. A. & Muotri, A. R. Advancing preclinical models of psychiatric disorders with human

brain organoid cultures. *Mol. Psychiatry* **28**, 83–95 (2023).

59.    Smits, L. M. *et al.* Modeling Parkinson's disease in midbrain-like organoids. *Npj Park. Dis.* **5**, 1–8

(2019).

60.    Chen, X. *et al.* Modeling Sporadic Alzheimer's Disease in Human Brain Organoids under Serum

Exposure. *Adv. Sci.* **8**, 2101462 (2021).

61.    Szebényi, K. *et al.* Human ALS/FTD brain organoid slice cultures display distinct early astrocyte

and targetable neuronal pathology. *Nat. Neurosci.* **24**, 1542–1554 (2021).

62.    Garcez, P. P. *et al.* Zika virus impairs growth in human neurospheres and brain organoids.

*Science* **352**, 816–818 (2016).

63.    Pellegrini, L. *et al.* SARS-CoV-2 Infects the Brain Choroid Plexus and Disrupts the Blood-CSF

Barrier in Human Brain Organoids. *Cell Stem Cell* **27**, 951-961.e5 (2020).

64.    Kozlowski, M. T., Crook, C. J. & Ku, H. T. Towards organoid culture without Matrigel. *Commun.*

*Biol.* **4**, 1–15 (2021).

65.    Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-3587.e29 (2021).

66.    Durinck, S., Spellman, P. T., Birney, E. & Huber, W. Mapping identifiers for the integration of

genomic datasets with the R/Bioconductor package biomaRt. *Nat. Protoc.* **4**, 1184–1191 (2009).

67.    Gu, Z., Eils, R. & Schlesner, M. Complex heatmaps reveal patterns and correlations in

multidimensional genomic data. *Bioinformatics* (2016).

68.    Carlson, M. org.Hs.eg.db: Genome wide annotation for Human. (2019).

69.    Pagès, H., Carlson, M., Falcon, S. & Li, N. AnnotationDbi: Manipulation of SQLite-based

annotations in Bioconductor. (2022).

70.    Vavrek, M. J. fossil: Palaeoecological and Palaeogeographical Analysis Tools. (2020).

71.    Wickham, H. ggplot2: Elegant Graphics for Data Analysis. *Springer-Verl. N. Y.* (2016).

999    **Figure 1**

1000

1001

**Using meta-analysis to quantify preserved primary tissue co-expression in organoids**

**A** Collection of annotated primary tissue brain scRNA-seq datasets, ranging from gestational

week (GW) 5 to 25 and sampling from 15 developmentally defined brain regions. The primary

tissue datasets are annotated at broad cell-type levels (Neural Progenitor, Dividing Progenitor,

Intermediate Progenitor, Glutamatergic, GABAergic, and Non-neuronal) and these annotations

are used to compute MetaMarkers, cell-type markers identified through recurrent differential

expression.

**B** Collection of human neural organoid scRNA-seq datasets, sampling from 12 different

differentiation protocols. Included is an annotated temporal forebrain organoid dataset.

**C** Example of a sparse co-expression network derived from a scRNA-seq data and an

example of an aggregate co-expression network averaged over many scRNA-seq datasets.

The aggregate network enhances the sparse signal from the individual network.

**D** Schematic showing a quantification of intra-marker set co-expression

**E** Schematic showing a quantification for the strength of preserved co-expression between two

co-expression networks, measuring the replication of the top 10 co-expressed partners of an

individual gene across the networks.

**F** Example plot from the preservedCoexp R library, placing cell-type specific preserved co-

expression scores of an example forebrain organoid dataset in reference to scores derived

from primary tissue datasets. Red lines denote the percentile of the organoid cell-type scores

within the primary tissue distributions.

1022

1023

1024

1025

1026

**Figure 2**

**Meta-analytic primary tissue cell-type markers**

**A** Annotated UMAPs of the annotated primary tissue brain scRNA-seq datasets.

**B** Example of our leave-one-out cross-validation approach for learning primary tissue

MetaMarkers and testing the markers' capacity for predicting annotations in the left-out

dataset, quantified with the AUROC statistic.

**C** Meta-analytic primary tissue markers have high performance in predicting primary tissue

cell-type annotations. Boxplot distributions of the AUROC statistic for predicting cell-type

annotations across all leave-one-out combinations of our annotated primary tissue datasets,

with an increasing number of MetaMarkers used for predicting cell-type annotations on the x-

axis.

**D** MetaMarkers have the highest performance in predicting primary tissue cell-type

annotations. Boxplots of marker gene-set performances. Gene-sets are the top 100 cell-type

markers from individual primary tissue datasets compared to the MetaMarker performance.

Performances for each cell-type in individual primary tissue datasets are presented in Supp.

Fig. 1A. Datasets are ordered by their median performance.

**E** Averaged distributions of gene expression for the top 100 MetaMarkers demonstrating clear

cell-type specificity. This is performed with a leave-one-out cross-validation, with individual

dataset distributions reported in Supp. Fig. 1B.

1047

1048

1049

1050

1051

1052

**Figure 3**

**1056   Neural organoids vary in recapitulating primary tissue cell-type marker set co-**

**1057   expression**

**A** Example of a sparse co-expression network derived from a scRNA-seq data and an

example of an aggregate co-expression network averaged over many scRNA-seq datasets.

The aggregate network enhances the sparse signal from the individual network.

**B** Marker gene-sets show clear cell-type clusters via their co-expression relationships in

primary tissue and organoid networks. The aggregated co-expression networks for the

unannotated primary tissue datasets and organoid datasets, showing the hierarchically

clustered co-expression of the primary tissue MetaMarkers for the 6 cell-types.

**C** Schematic for the co-expression module learning framework, measuring the co-expression

strength within an arbitrary gene-set compared to the rest of the genome, quantified with the

AUROC statistic.

**D** Distributions of co-expression module AUROCs for individual annotated primary tissue,

unannotated primary tissue, and organoid datasets for the co-expression strength of the

MetaMarker gene-sets for the 6 cell-types. The grey 'All GO terms' distributions report the

average co-expression module AUROC across all GO terms for each individual dataset. Co-

expression module AUROCs for the aggregate co-expression networks are denoted with the

special characters.

**E** Top primary tissue and top and bottom organoid co-expression networks based on

Glutamatergic co-expression module AUROCs. Genes are ordered within each MetaMarker

gene set by their average intra-gene set co-expression.

1077

1078

1079

 **Figure 4**

1081

**Neural organoids vary in their preservation of primary tissue gene-level co-expression**

**A** Schematic showing the quantification for gene-level preserved co-expression. The

preserved co-expression score for any given gene-set is the average preserved co-expression

AUROC across all genes within that gene set.

**B** Organoids strongly vary in preserved primary tissue cell-type specific co-expression in

comparison to fetal data. Boxplot distributions show the preserved co-expression scores for

the primary tissue MetaMarker gene-sets of the 6 cell-type annotations across all individual

networks.

**C** The majority of cell-types are significantly correlated in preserved co-expression within

organoid networks. Spearman correlation matrix for the preserved co-expression scores for all

6 cell-type annotations across all individual organoid datasets.

**D** Preserved co-expression scores computed from the dorsal patterned forebrain and ventral

midbrain organoid datasets for the top 10 cell-type markers of various neural cell-types.

**E** Scatter plots comparing the preserved co-expression score of the top 100 MetaMarkers

against the top 10 markers (no overlaps in gene sets) for various neural cell-types per

organoid dataset. Spearman correlation coefficients are reported in the bottom right corner.

**F** Scatter plots summarizing the semantic distances of GO terms that are significantly

preserved or non-preserved between the aggregate annotated primary tissue and organoid co-

expression networks.

**G** Organoids globally fail to preserve primary tissue co-expression of ECM and vascular

related genes. Bar plot detailing the top 10 GO terms from a GO enrichment test of the 76

genes with high and low preserved co-expression AUROCs within primary tissue networks and

organoid networks respectively. The preserved co-expression for each individual gene from

primary tissue networks and organoid networks is reported in Supp. Fig. 6B.

1106

1107

1108    **Figure 5**

1109

1110

**Neural organoids capture temporal dynamics in primary tissue co-expression**

**A** Schematic showing two potential outcomes when comparing the preserved co-expression

between primary tissue and organoid data on a temporal axis. There may be a temporal

relationship, with younger organoids recapitulating younger primary tissue co-expression over

older primary tissue co-expression and vice versa for older organoids, or there may be no

temporal relationship.

**B** Organoid co-expression models temporal trends in primary tissue co-expression. Line plots

showing the preserved co-expression scores computed from individual organoid co-expression

networks for cell-type markers of individual primary tissue datasets. Primary tissue datasets on

the x-axis are ordered from youngest to oldest.

1121

1122    **Figure 6**

1123

1124

**The preservedCoexp R package enables fast computation of preserved co-expression**

**A** The preservedCoexp R package can compute co-expression networks and genome-wide

preservation of co-expression in a few minutes even for low-memory computers. Line plots

showing the computational time to either compute co-expression networks or preserved co-

expression as the number of cells or genes increases. Points are the mean value from 10

replicates, with error bars depicting ± 1 standard deviation.

**B** Example plot from the preservedCoexp R package, placing cell-type specific preserved co-

expression scores of an example forebrain organoid dataset in reference to scores derived

from primary tissue datasets or organoid datasets. Red lines denote the percentile of the

forebrain organoid cell-type scores within either the primary tissue distributions or organoid

distributions.

1136

1137