A broad-spectrum family GH13 α-glucosidase from *Marinovum* sp., a member of the *Roseobacter* clade

Jinling Li,[1] Janice Mui,[1] Bruna M. da Silva,[2,3] Douglas E.V. Pires,[2,3,4] David B. Ascher,[2,4] Niccolay Madiedo Soler,[5,6] Ethan D. Goddard-Borger,[5,6] Spencer J. Williams[1]*

[1] School of Chemistry and Bio21 Molecular Science and Biotechnology Institute, University of Melbourne, Parkville, Victoria 3010, Australia.

[2] Computational Biology and Clinical Informatics, Baker Heart and Diabetes Institute, Melbourne, 3004, Victoria, Australia

[3] School of Computing and Information Systems, University of Melbourne, Parkville, Victoria 3010, Australia.

[4] School of Chemistry and Molecular Biosciences, University of Queensland, St Lucia 4072, Queensland, Australia.

[5] ACRF Chemical Biology Division, The Walter and Eliza Hall Institute of Medical Research, Parkville, Victoria 3010, Australia.

[6] Department of Medical Biology, University of Melbourne, Parkville, Victoria 3010, Australia.

E-mail: sjwill@unimelb.edu.au; ORCID: 0000-0001-6341-4364

Abstract

Glycoside hydrolases (GHs) are a diverse group of enzymes that catalyze the hydrolysis of glycosidic bonds. The Carbohydrate-Active enZymes (CAZy) database organizes GHs into families based on sequence data and function, with fewer than 1% of the predicted proteins characterized biochemically. Consideration of genomic context can provide a useful guide to infer possible enzyme activities for proteins of unknown function. We used the MultiGeneBLAST tool to discover a putative gene cluster in *Marinovum* sp., a member of the marine *Roseobacter* clade, that encodes homologues of enzymes belonging to the sulfoquinovose monooxygenase pathway. This putative gene cluster lacks a gene encoding a classical family GH31 sulfoquinovosidase (SQase) candidate, but which instead includes an uncharacterized family GH13 protein (*Ms*GH13). We show that recombinant *Ms*GH13 lacks SQase activity but is a broad spectrum α-glucosidase active on a diverse array of α-linked disaccharides, including: maltose (100%), sucrose (7.9%), nigerose (52%), trehalose (20%), isomaltose (6.4%), and kojibiose (3.8%). Using AlphaFold, a 3D model for the *Ms*GH13 enzyme was constructed that predicted its active site shared close similarity with a narrower specificity α-glucosidase from *Halomonas* sp. H11 of the same GH13 subfamily. This study supports recent findings that bacteria belonging to the *Roseobacter* clade can metabolize SQ without a classical SQase encoded in their genomes.

Keywords: carbohydrates, enzymes, glycosidase, bioinformatics

Introduction

Glycoside hydrolases (GHs), which catalyze the hydrolysis of glycosidic bonds, are a large and diverse group of enzymes. GHs have been classified into families in the Carbohydrate-Active enZymes (CAZy) database (www.cazy.org[1] and www.cazypedia.org[2]). The CAZy database, which consists of around 170 families, organizes GHs into families on the basis of sequence data and function. However, fewer than 1% of the predicted proteins within the CAZy database have been biochemically characterized, and therefore sequence-based predictions of functions are the primary means for assigning possible functions to uncharacterized GH proteins. Often, especially within bacteria, genes encoding specific GHs occur within gene clusters that contain other genes encoding related enzymes that enable the products of GH action to be metabolized. Thus, genetic context can be a powerful ally in predicting the function of uncharacterized GHs.

GH family 31 (GH31) is a large glycoside hydrolase family with >20,000 members that possess diverse activities on α-glycosidic bonds including α-glucosidase, α-glucan lyase, α-xylanase and sulfoquinovosidases (SQases).[3] SQases catalyze hydrolysis of glycosides of sulfoquinovose (SQ; 6-deoxy-6-sulfo-glucose),[4, 5] such as in the ubiquitous plant sulfolipid α-sulfoquinovosyl diacylglyceride (SQDG). SQDG is produced by essentially all photosynthetic organisms, and is one of the most abundant organosulfur compounds in nature.[6, 7] Upon its release, SQ can be utilized as a nutrient by sulfoglycolytic organisms through the pathways of sulfoglycolysis.[8, 9] Thus, SQases act as the gateway enzyme for SQ glycosides to enter sulfoglycolysis. Family GH31 SQases are usually found within gene clusters encoding assorted sulfoglycolysis pathways. Recently, a new SQ assimilation pathway, the sulfoglycolytic sulfoquinovose monooxygenase (sulfo-SMO) pathway, was reported that involves oxidative desulfurization of SQ to release sulfite. Representative sulfo-SMO gene clusters in *Agrobacterium tumefaciens* and *Novosphingobium aromaticivorans* encode family GH31 SQases **(Fig. 1b)**.[10, 11]

By searching using the MultiGeneBLAST tool,[12] we discovered a *Marinovum* sp. (a member of the *Roseobacter* clade bacteria)[13] that contains genes that are homologous to the key sulfo-SMO enzymes, suggesting that it encodes a sulfo-SMO pathway **(Fig. 1a)**. However, this proposed gene cluster lacked a candidate gene encoding a putative family GH31 SQase. Instead, this putative gene cluster contained a gene encoding a GH13 protein (hereafter *Ms*GH13). Family GH13 is the largest of the CAZy families of glycoside hydrolases, and contains approximately 150,000 sequences but

with <800 members that have been biochemically characterized to define their substrate.[14] Activities within family GH13 share some overlap with family GH31, and includes diverse activities on α-glycosidic bonds including α-glucosidase, α-amylase, α-glucosyltransferase, and sucrose α-glucosidase, but no SQases have been described in this family.

In this work we explore the possibility that MsGH13 is a non-classical SQase. We clone and express MsGH13 and screen it against a panel of activated, 4-nitrophenyl α-glycosides. We show that MsGH13 lacks SQase activity, but is active on PNPGlc. Further analysis reveals it to be active on a wide range of different α-glucosidic linkages, namely maltose, sucrose, nigerose, trehalose, isomaltose and kojibiose. We use AlphaFold to present a model for the MsGH13 enzyme and its active site.

Materials and Methods

Bioinformatic identification of *Marinovum* sp. Sulfo-SMO locus

The *A. tumefaciens* sulfoquinovosidase (AAK90105.1, locus tag Atu3285), ABC transporter membrane spanning domains (AAK90106.2 and AAK90107.2, locus tags Atu3284 and Atu3283), ABC transporter substrate binding domain (AAK90108.1, locus tag Atu3282), ABC transporter ATP binding domain (AAK90109.2, locus tag Atu3281), sulfo-SMO transcriptional regulator (AAK90110.1, locus tag Atu3280), SQ monooxygenase (AAK90111.1, locus tag Atu3279), 6-oxo-Glc reductase (AAK90112.1, locus tag Atu3278) and flavin reductase (AAK90113.1, locus tag Atu3277) were submitted separately as queries to the NCBI protein-protein BLAST[15] (BLASTp) (https://blast.ncbi.nlm.nih.gov/). The database used was non-redundant protein sequences (nr), with *A. tumefaciens* (taxid: 358) sequences excluded. Standard algorithm parameters were used, except the maximum target sequences was set to 10,000. The raw BLASTp results were filtered by E-value threshold, with protein sequences retained if E-value ≤ 1.19e-51, the smallest of the maximum E-values obtained from the BLASTp searches of the three enzymes that interact with the SQ moiety (sulfoquinovosidase, SQ monooxygenase and the ABC transporter substrate binding domain). The lists of protein sequences were converted to lists of the corresponding nucleotide sequences. Protein sequences that did not return a corresponding nucleotide sequence were discarded. The nine lists were combined, with duplicates and any remaining *A. tumefaciens* sequences removed to give a list of 858,315 candidate nucleotide sequences. Python scripts were used to filter BLASTp results by E-

value threshold, convert protein sequences to corresponding nucleotide sequences, combine the lists, remove duplicates, and remove any *A. tumefaciens* sequences. These scripts are available on GitHub, along with instructions for their use (https://github.com/jmui-unimelb/Gene-Cluster-Search-Pipeline).[16]

The list of candidate sequences was converted to a MultiGeneBLAST[12] reference library, and searched with the *A. tumefaciens* sulfo-SMO gene cluster as a query (http://multigeneblast.sourceforge.net/). The search was run on a high-performance computing (HPC) cluster to enable parallelization (https://www.bio21.unimelb.edu.au/hpc/index.php). A modified MultiGeneBLAST script is available on GitHub that can be run in parallel on an HPC (https://github.com/jmui-unimelb/Gene-Cluster-Search-Pipeline).[16]

Data were filtered by MultiGeneBLAST score (minimum score 3.0). The MultiGeneBLAST score for a gene cluster is determined by the following equation: score = $h + i \times s$. h is the number of query genes that had homologs (i.e., had a BLAST hit of at least a user-specified sequence coverage and % identity to the query gene; default values used in this study were sequence coverage = 25 and % identity = 30). s is the number of contiguous gene pairs with conserved synteny (i.e., gene order), while i is a factor that determines the weight of synteny in the score (default value: 0.5; default used in this study). A value of 0.5 means that the number of homologous genes is given twice the weight as conservation of gene order.[12] Putative sulfo-SMO gene clusters identified using this workflow were screened for clusters that contained a GH13 enzyme as a candidate SQase.

Cloning, expression and purification of *Ms*GH13

A dsDNA oligonucleotide encoding *Ms*GH13 (MAI18115.1) was synthesized (Integrated DNA Technologies) and cloned into the pET29b(+) vector at the *NdeI* and *XhoI* sites (**Fig. S1**). The resulting plasmid was sequence-verified by Sanger sequencing, then transformed into 'T7 Express' *E. coli* (New England BioLabs) and selected on LB-agar + 50 µg/ml kanamycin at 37 °C for 16 h. A single colony was used to inoculate 10 ml of LB media + 50 µg/ml kanamycin and the cultures were incubated at 37 °C for 16 h. These starter cultures were used to inoculate 1000 ml of S-broth (35 g tryptone, 20 g yeast extract, 5 g NaCl, pH 7.4) + 50 µg/ml kanamycin, which was incubated with shaking (250 rpm) at 37 °C until it reached an A_{600} of 0.7. The culture was cooled to room temperature, isopropyl thiogalactoside (IPTG) added to a final concentration of 400 µM, and

incubation with shaking (200 rpm) continued at 18 °C for 19 h. Cells were harvested by centrifugation at 8,000 g for 20 min at 4 °C, then the pellet was resuspended in 40 ml binding buffer (50 mM NaP$_i$, 300 mM NaCl, 5 mM imidazole, pH 7.5) containing protease inhibitor (Roche cOmplete EDTA-free protease inhibitor cocktail) and lysozyme (0.1 mg/ml) by nutating at 4 °C for 30 min. Benzonase (1 µl) was added to the mixture and lysis was effected by sonication [10 × (15 s on; 45 s off) at 45% amplitude]. The lysate was centrifuged at 18,000 g for 20 min at 4 °C and the supernatant was collected. The supernatants were filtered (0.45 µm) and loaded onto a 1 ml HisTrap IMAC column (GE). The column was washed with 10 ml of binding buffer, then the protein was eluted using elution buffer (50 mM NaP$_i$, 300 mM NaCl, 400 mM imidazole, pH 7.5). Fractions containing product, as judged by SDS-PAGE, were further purified by size exclusion chromatography on a HiPrep 16/60 Sephacryl S-200 HR column (GE) using 50 mM NaP$_i$, 150 mM NaCl, pH 7.5.

Substrate screening of *Ms*GH13

Hydrolytic activity was examined towards a range of 4-nitrophenyl α-glycosides (each 4 mM) in 600 µL reaction mixtures containing 50 nM *Ms*GH13 in 50 mM citric acid/phosphate buffer, 150 mM NaCl, pH 7 at room temperature. Reactions were performed in a cuvette and monitored for release of 4-nitrophenol/4-nitrophenolate using an UV/Vis spectrophotometer at the isosbestic point (λ = 348 nm, ε = 5125 M^{-1} cm^{-1}). The following sugars were examined: α-D-glucopyranoside (PNPGlc), 4-nitrophenyl α-D-sulfoquinovoside (PNPSQ), 4-nitrophenyl α-D-galactopyranoside (PNPGal), 4-nitrophenyl α-D-xylopyranoside (PNPXyl), 4-nitrophenyl α-D-mannopyranoside (PNPMan), 4-nitrophenyl *N*-acetyl-α-D-galactosaminide (PNPGalNAc), 4-nitrophenyl α-D-glucuronide (PNPGlcA) and 4-nitrophenyl α-D-glucopyranoside 6-phosphate (6-P-PNPGlc) (**Fig. S2**).

Hydrolytic activities toward various disaccharides were assessed in 600 µL reaction mixtures containing 73.8 nM *Ms*GH13 in 50 mM citric acid/phosphate buffer, 150 mM NaCl, pH 7 at room temperature. After 5 h, reactions were heat inactivated then applied to an aluminium backed high performance thin layer chromatography (HPTLC) sheet (Supelco) and eluted with ethyl acetate, methanol and water (7:4:2). Products were visualized by spraying with a solution of 0.2% orcinol, 10% H$_2$SO$_4$ and 10% water in 80% ethanol, and heating. The following sugars were assessed as substrates: trehalose [α-D-Glc*p*-(1↔1)-α-D-Glc*p*], kojibiose [α-D-Glc*p*-(1→2)-D-Glc], nigerose [α-D-Glc*p*-(1→3)-D-Glc], maltose [α-D-Glc*p*-(1→4)-D-Glc], isomaltose [α-D-Glc*p*-(1→6)-D-Glc], and

sucrose [β-D-Fru$f$-(2↔1)-α-D-Glc$p$] (**Fig. S3**).

pH dependence of *Ms*GH13 activity

The Michaelis parameter k_{cat}/K_M was measured for PNPGlc hydrolysis using the substrate depletion method in 50 mM citric acid/phosphate buffer, 150 mM NaCl at a range of pH values (4.0, 4.5, 5.0, 6.0, 6.5, 7.0, 7.5, 8.0, 8.3) at room temperature. A concentration of PNPGlc of 0.1 mM was used, being $<K_M/10$. Reactions were monitored at the isosbestic point of 4-nitrophenol (λ = 348 nm, ε = 5125 M^{-1} cm^{-1}). Reactions were initiated by adding *Ms*GH13 to a final concentration of 50 nM and the rate measured continuously using a UV/visible spectrophotometer. k_{cat}/K_M values were calculated using the equation y = (y$_0$ - y$_\infty$)×exp(-k×t) + y$_\infty$, where k_{cat}/K_M = k/[E]. pK_a values were fit to a bell shaped curve using the Prism 9.50 software package (Graphpad Scientific Software), with the equation y = m×(1/(1+[(10^{-x})/(10^{-pKa1})+(10^{-pKa2})/(10^{-x})]))+c. Below pH 4.0 and above pH 8.0 the enzyme was unstable.

Temperature stability

Temperature stability of *Ms*GH13 was assessed by incubation of 50 nM *Ms*GH13 in the assay buffer for 3 h at different temperatures (room temperature, 30˚C, 35˚C, 40˚C, 45˚C, 50˚C, 55˚C). After this time, PNPGlc was added to a final concentration of 0.1 mM, and the reaction rate measured using a UV/Vis spectrometer.

Michaelis-Menten kinetics

Michaelis-Menten kinetics were measured for *Ms*GH13 catalyzed hydrolysis of PNPGlc using a UV/visible spectrophotometer. Release of the chromogenic 4-nitrophenolate was monitored at the isosbestic point (λ = 348 nm, ε = 5125 M^{-1} cm^{-1}). Reactions were conducted in 50 mM citric acid/phosphate, 150 mM NaCl (pH 6.5) at 25 ºC using 50 nM *Ms*GH13 at substrate concentrations ranging from 0.1 to 2 mM. For quantitative analysis of *Ms*GH13 hydrolysis of disaccharides, the reaction was measured by using Colorimetric Detection Kit (Invitrogen^{TM} by Thermo Fisher Scientific). The reaction was performed by using 12.3 μM *Ms*GH13 in 50 mM citric acid/phosphate, 150 mM NaCl buffer (pH 6.5) with various concentrations of the disaccharide substrates. Samples at each concentration were measured after 10 and 20 min incubation to ensure a linear initial rate for

the reaction. The reaction was quenched by heating at 80˚C for 5 minutes. The quenched samples were cooled to room temperature then glucose release was quantified according to the manufacturer's instructions at 30˚C using a UV/visible spectrophotometer. Hydrolysis of maltose, nigerose, trehalose, kojibiose and isomaltose each produce two equivalents of glucose, and the reaction rates were therefore halved. Kinetic parameters (k_{cat}, K_M, k_{cat}/K_M) were calculated using the Prism 9.50 software package (GraphPad Scientific Software) using the Michaelis-Menten equation.

NMR spectroscopy

The stereochemistry of the initially formed product of *Ms*GH13 catalyzed hydrolysis of PNPGlc, was monitored by ^{1}H NMR spectroscopy (500 MHz). *Ms*GH13 (73.8 nM) was added to a solution of PNPGlc (10 mM) in 50 mM citric acid/phosphate buffer, 150 mM NaCl at pD 7.4. ^{1}H NMR (500 MHz) revealed the formation of a new signal at δ 5.15 ppm, $J_{1,2}$ 3.8 Hz, assigned α-glucose. The α-glucose slowly underwent mutarotation and formed a signal at δ 4.56 ppm, $J_{1,2}$ 8.0 Hz, assigned β-glucose.

3D modelling

A 3D model for *Ms*GH13 was constructed using Alphafold.[17, 18] An overlay with *Halomonas* sp. H11 glucosidase (PDB 3WY4) was generated using the *align* function within PyMOL.

Results

MultiGeneBLAST was used to search the non-redundant protein sequence (nr) database for organisms that possess groups of genes encoding proteins of the sulfoglycolytic monooxygenase pathway: SmoA, SmoB, SmoC, SmoD and SmoI. Putative sulfo-SMO gene clusters were filtered and manually screened for the absence of a homolog of a gene encoding the family GH31 SmoG SQase, leading to the identification of *Marinovum* sp. (GenBank: NZGN01000053.1), containing a gene encoding a family GH13 protein, *Ms*GH13.

The gene encoding *Ms*GH13 (MAI18115.1), codon-harmonized for *E. coli*, was synthesized, cloned, expressed and purified to homogeneity from this host. To assess the substrate specificity of *Ms*GH13, it was screened against various PNP glycosides: PNPGlc was a good substrate, but it was inactive against PNPSQ, PNPGal, PNPXyl, PNPMan, PNPGalNAc, PNPGlcA and 6-P-PNPGlc (for structures see **Fig. S2**).

The stereochemical outcome of *Ms*GH31 catalyzed hydrolysis of PNPGlc was assessed by ^{1}H NMR. This revealed the initial formation of α-glucose (δ 5.15 ppm), which slowly underwent mutarotation to give a 1:1.42 ratio of α-glucose and β-glucose (δ 4.56 ppm). This demonstrated that the enzyme is a retaining a α-glucoside hydrolase (**Fig. 2**).

The effects of pH on the hydrolytic activity and the temperature stability of *Ms*GH13 were evaluated using PNPGlc as a substrate. The pH dependence of k_{cat}/K_M gave a bell-shaped curve, with an optimum at pH 6.5, and pK_{a1} and pK_{a2} values of 6.2 ± 0.2 and 7.1 ± 0.2 (**Fig. 3a**). The effect of temperature on *Ms*GH13 activity was assessed at pH 6.5 by incubation for 3 h at various temperatures, then addition of substrate. *Ms*GH13 incubated at 25 ℃ retained activity, enzyme incubated at 30 ℃ retained only 16.7% activity, and enzyme incubated at 35 ℃ and above lost all activity. Kinetic parameters for PNPGlc measured at pH 6.5 and 25 ℃ were k_{cat} = 12.4 s^{-1}, K_M = 0.95 mM and k_{cat}/K_M = 13,000 M^{-1} s^{-1} (**Fig. 3b**).

Hydrolytic activity on trehalose, kojibiose, nigerose, maltose, isomaltose, and sucrose was qualitatively assessed by HPTLC. Samples of each disaccharide were incubated with *Ms*GH13, then separated by HPTLC and visualized by staining with orcinol. Nigerose, trehalose, sucrose and maltose were completely hydrolyzed, while kojibiose and isomaltose were partially hydrolyzed (**Fig. 4a**). The kinetic parameters for hydrolysis of nigerose, sucrose, maltose, trehalose, kojibiose and isomaltose were assessed using a colorimetric glucose detection assay (**Table 1**). The greatest activity

was obtained for maltose (k_{cat} = 135 s^{-1}, K_M = 3.22 mM and k_{cat}/K_M = 42,000 M^{-1} s^{-1}). The relative percentage of k_{cat}/K_M values versus maltose were: nigerose (52%), trehalose (20%), sucrose (7.9%), isomaltose (6.4%) and kojibiose (3.8%).

Discussion

We used MultiGeneBLAST to search the NCBI database for bacteria containing homologues of sulfo-SMO genes, but lacking candidate SQases belonging to family GH31. This search identified *Marinovum* sp., which possessed genes encoding homologues of the *Agrobacterium tumefaciens* sulfoquinovose monooxygenase SmoC (75% identity), flavin reductase SmoA (47% identity), 6-oxo-glucose dehydrogenase SmoB (61% identity), as well as an ABC transporter cassette and an associated substrate binding protein. The *Marinovum* sp. gene cluster contained a gene encoding a member of family GH13, and which was computer annotated as an α-glucosidase. Family GH13 has been divided into 35 subfamilies based on sequence similarity, structure and function.[14] The protein from *Marinovum* sp. belongs to the GH13_23 subfamily, which contains >3200 members, but just 9 members of this subfamily have been characterized. These activities include α-glucosyltransferase (EC 2.4.1.-), oligo-α-1,6-glucosidase (EC 3.2.1.10, including isomaltose), exo-1,3-α-glucanase (3.2.1.84, including nigerose), α-glucosidase (EC 3.2.1.20, including maltose and sucrose), kojibiose hydrolase (EC 3.2.1.216, including kojibiose), and α,α-trehalase (EC 3.2.1.28) activities.[19-22] Based on the genomic context of the annotated GH13 in *Marinovum* sp., we set out to explore whether this might encode an SQase or another enzyme activity.

Recombinant *Ms*GH13 was screened against a range of chromogenic monosaccharide substrates that included head-groups representing activities present within GH13 and GH31, among others. *Ms*GH13 was active upon PNPGlc but was inactive against other α-glycosides including PNPSQ, PNPGal, PNPXyl, PNPMan, PNPGalNAc, PNPGlcA and 6-P-PNPGlc. Thus, *Ms*GH13 is an α-glucosidase, and is not an SQase. ^{1}H NMR analysis of the products from *Ms*GH13 cleavage of PNPGlc revealed initial formation of α-Glc, which mutarotated to give an equilibrium mixture of α- and β-glucose; thus, *Ms*GH13 is a retaining α-glucosidase.

*Ms*GH13 catalyzed the hydrolysis of the following disaccharides: trehalose, kojibiose, nigerose, maltose, isomaltose, and sucrose. The second order rate constants k_{cat}/K_M revealed a 25-fold range of activity, with a 2-fold preference for maltose over nigerose, and greater ratios for the remaining

disaccharide substrates (**Fig. 4b-4g, Table 1**). Thus, *Ms*GH13 is an unusually broad-spectrum α-glucosidase with the ability to cleave 1,1-, 1,2-, 1,3-, 1,4- and 1,6-glucosidic linkages. This ability to cleave diverse linkages is shared by *Thermus thermophilus* HB27 α,α-trehalase, another member of subfamily GH13_23. While this enzyme prefers trehalose, it displays good relative activity on isomaltose (63%) and kojibiose (20%), and modest activities on nigerose (16%), maltose (10%) and sucrose (8%).[22]

Comparison of the AlphaFold 3D structure prediction of *Ms*GH13 to the experimentally determined 3D X-ray structure of *Halomonas* sp. H11 glucosidase (HaG) mutant with maltose[23] (PDB: 3WY4, green), which is also a member of subfamily GH13_23, shows excellent structural similarity with RMSD = 2.24 Å (**Fig. 5**). This overlay reveals conservation of the key catalytic machinery, allowing assignment of catalytic nucleophile (*Ms*GH13/HaG: D201/D202), acid/base (E272/E271), and substrate-binding residues (D61/D62, H104/H105, R199/R200, H336/H332, D337/D333 and R410/R400). Like *Ms*GH13, HaG prefers maltose (100%).[24] However, HaG is much more specific, with good relative activity on sucrose (33%), very low relative hydrolysis activity against isomaltose (1.1%), trehalose (2.6%), nigerose (4.2%), and no detectable activity on kojibiose.[24] These active site residues are also conserved within the broad-spectrum HB27 α,α-trehalase.[22]

This study was initiated based on the genomic context of *Ms*GH13 within a putative sulfo-SMO gene cluster, and the observation that sulfoglycolytic gene clusters typically contain SQase genes of GH family 31. This is believed to allow sulfoglycolytic organisms to use SQ glycosides as a nutrient. However, a recent study of *Roseobacter* clade bacteria demonstrated that *Dinoroseobacter shibae* DFL 12 and *Roseobacter denitrificans* OCh 114 can grow on SQ, and identified many more Roseobacter clade bacterial genomes that contained putative sulfo-SMO metabolic genes (including *Marinovum algicola* DG 898) but also lacked a family GH31 SQase candidate.[25] This study reported that marine phytoplankton and cyanobacteria contain intracellular SQ at micro to millimolar levels, and that SQ can be detected within particulate organic matter (size > 3 μm sampled from the Bohai Sea), suggesting that some bacteria may encounter sufficient environmental SQ to support their growth without a requirement for an SQase.[25]

In conclusion, *Ms*GH13, while possessing a genomic context that is suggestive of possible SQase activity, is really a broad spectrum α-glucosidase with activity on a wide range of α-linked disaccharides. The high similarity of the active sites of the AlphaFold predicted *Ms*GH13 structure

with that of the maltose/sucrose specific HaG α-glucosidase provides little insight into how the active site of the latter can accommodate such a broad range of disaccharides. Possibly, this reflects limitations in the use of AlphaFold to predict structures of complexes, and provides an impetus for structural studies to define in greater detail how disaccharides with other linkages can be accommodated within the active site.

Declarations

Ethical Approval Not applicable

Consent to participate Not applicable

Consent to publish Not applicable

Author contributions

SJW conceived the study with input from EGB and DA. JL, JM, BMdS, DEVP and DBA performed bioinformatics. NMS and EGB conducted molecular biology. JL conducted enzymology. JL and SJW wrote the paper with input and approval of the final draft from the authors.

Funding

S.J.W is supported by the Australian Research Council (DP210100233, DP210100235). D.B.A. is supported by an Investigator Grant from the National Health and Medical Research Council (NHMRC) of Australia (GNT1174405). E.D.G-B. acknowledges support from The Walter and Eliza Hall Institute of Medical Research, National Health and Medical Research Council of Australia (NHMRC) Ideas grant GNT2000517, the Australian Cancer Research Fund; and the Brian M. Davis Charitable Foundation Centenary Fellowship. JL thanks the China Scholarship Council for a PhD scholarship. BMdS is supported by Melbourne Research Scholarship.

Competing interests The authors declare that they have no conflict of interest.

Availability of data and materials

All data generated or analysed during this study are included in this published article and its supplementary information files.

References

(1) Drula, E.; Garron, M. L.; Dogan, S.; Lombard, V.; Henrissat, B.; Terrapon, N. (2022) The carbohydrate-active enzyme database: functions and literature. *Nucleic Acids Res.*, *50*, D571-D577.

(2) The CAZypedia Consortium. (2018) Ten years of CAZypedia: a living encyclopedia of carbohydrate-active enzymes. *Glycobiology*, *28*, 3-8.

(3) Arumapperuma, T.; Li, J.; Hornung, B.; Madiedo Soler, N.; Goddard-Borger, E. D.; Terrapon, N.; Williams, S. J. (2023) A subfamily classification to choreograph the diverse activities within Glycoside Hydrolase family 31 (GH31). *J. Biol. Chem.*, *299*, 103038.

(4) Speciale, G.; Jin, Y.; Davies, G. J.; Williams, S. J.; Goddard-Borger, E. D. (2016) YihQ is a sulfoquinovosidase that cleaves sulfoquinovosyl diacylglyceride sulfolipids. *Nat. Chem. Biol.*, *12*, 215-217.

(5) Abayakoon, P.; Jin, Y.; Lingford, J. P.; Petricevic, M.; John, A.; Ryan, E.; Mui, J. W. Y.; Pires, D. E. V.; Ascher, D. B.; Davies, G. J.; et al. (2018) Structural and Biochemical Insights into the Function and Evolution of Sulfoquinovosidases. *ACS Cent. Sci.*, *4*, 1266-1273.

(6) Harwood, J. L.; Nicholls, R. G. (1979) The plant sulpholipid-- a major component of the sulphur cycle. *Biochem. Soc. Trans.*, *7*, 440-447.

(7) Goddard-Borger, E. D.; Williams, S. J. (2017) Sulfoquinovose in the biosphere: occurrence, metabolism and functions. *Biochem. J.*, *474*, 827-849.

(8) Snow, A. J. D.; Burchill, L.; Sharma, M.; Davies, G. J.; Williams, S. J. (2021) Sulfoglycolysis: catabolic pathways for metabolism of sulfoquinovose. *Chem. Soc. Rev.*, *50*, 13628.

(9) Wei, Y.; Tong, Y.; Zhang, Y. (2022) New mechanisms for bacterial degradation of sulfoquinovose. *Biosci. Rep.*, *42*, BSR20220314.

(10) Sharma, M.; Lingford, J. P.; Petricevic, M.; Snow, A. J. D.; Zhang, Y.; Jarva, M. A.; Mui, J. W.; Scott, N. E.; Saunders, E. C.; Mao, R.; et al. (2022) Oxidative desulfurization pathway for complete catabolism of sulfoquinovose by bacteria. *Proc. Natl. Acad. Sci. U. S. A.*, *119*, e2116022119.

(11) Liu, J.; Wei, Y.; Ma, K.; An, J.; Liu, X.; Liu, Y.; Ang, E. L.; Zhao, H.; Zhang, Y. (2021) Mechanistically Diverse Pathways for Sulfoquinovose Degradation in Bacteria. *ACS Catal.*, *11*, 14740-14750.

(12) Medema, M. H.; Takano, E.; Breitling, R. (2013) Detecting Sequence Homology at the Gene Cluster Level with MultiGeneBlast. *Mol. Biol. Evol.*, *30*, 1218-1223.

(13) Martens, T.; Heidorn, T.; Pukall, R.; Simon, M.; Tindall, B. J.; Brinkhoff, T. (2006) Reclassification of Roseobacter gallaeciensis Ruiz-Ponte et al. 1998 as Phaeobacter gallaeciensis gen. nov., comb. nov., description of Phaeobacter inhibens sp. nov., reclassification of Ruegeria algicola (Lafay et al. 1995) Uchino et al. 1999 as Marinovum algicola gen. nov., comb. nov., and emended descriptions of the genera Roseobacter, Ruegeria and Leisingera. *Int. J. Syst. Evol. Microbiol.*, *56*, 1293-1304.

(14) Stam, M. R.; Danchin, E. G.; Rancurel, C.; Coutinho, P. M.; Henrissat, B. (2006) Dividing the large glycoside hydrolase family 13 into subfamilies: towards improved functional annotations of alpha-amylase-related proteins. *Protein Eng. Des. Sel.*, *19*, 555-562.

(15) Altschul, S. F.; Gish, W.; Miller, W.; Myers, E. W.; Lipman, D. J. (1990) Basic local alignment search tool. *J. Mol. Biol.*, *215*, 403-410.

(16) Mui, J. W. (2021) *Gene cluster search pipeline*. GitHub, https://github.com/jmui-unimelb/Gene-Cluster-Search-Pipeline

(17) Jumper, J.; Evans, R.; Pritzel, A.; Green, T.; Figurnov, M.; Ronneberger, O.; Tunyasuvunakool, K.; Bates, R.; Zidek, A.; Potapenko, A.; et al. (2021) Highly accurate protein structure prediction with AlphaFold. *Nature*, *596*, 583-589.

(18) Varadi, M.; Anyango, S.; Deshpande, M.; Nair, S.; Natassia, C.; Yordanova, G.; Yuan, D.; Stroe, O.; Wood, G.; Laydon, A.; et al. (2022) AlphaFold Protein Structure Database: massively expanding the structural coverage of protein-

sequence space with high-accuracy models. *Nucleic Acids Res.*, *50*, D439-D444.

(19) Sato, T.; Hasegawa, N.; Saito, J.; Umezawa, S.; Honda, Y.; Kino, K.; Kirimura, K. (2012) Purification, characterization, and gene identification of an alpha-glucosyl transfer enzyme, a novel type alpha-glucosidase from Xanthomonas campestris WU-9701. *J. Mol. Catal. B Enzym.*, *80*, 20-27.

(20) Kashiwabara, S.; Matsuki, Y.; Kishimoto, T.; Suzuki, Y. (1998) Clustered proline residues around the active-site cleft in thermostable oligo-1,6-glucosidase of Bacillus flavocaldarius KP1228. *Biosci. Biotech. Biochem.*, *62*, 1093-1102.

(21) Li, W.; Xue, Y.; Li, J.; Yuan, J.; Wang, X.; Fang, W.; Fang, Z.; Xiao, Y. (2016) A cold-adapted and glucose-stimulated type II alpha-glucosidase from a deep-sea bacterium Pseudoalteromonas sp. K8. *Biotechnol. Lett.*, *38*, 345-349.

(22) Alarico, S.; da Costa, M. S.; Empadinhas, N. (2008) Molecular and physiological role of the trehalose-hydrolyzing alpha-glucosidase from Thermus thermophilus HB27. *J. Bacteriol.*, *190*, 2298-2305.

(23) Shen, X.; Saburi, W.; Gai, Z.; Kato, K.; Ojima-Kato, T.; Yu, J.; Komoda, K.; Kido, Y.; Matsui, H.; Mori, H.; et al. (2015) Structural analysis of the alpha-glucosidase HaG provides new insights into substrate specificity and catalytic mechanism. *Acta Crystallogr. D*, *71*, 1382-1391.

(24) Ojima, T.; Saburi, W.; Yamamoto, T.; Kudo, T. (2012) Characterization of Halomonas sp. strain H11 alpha-glucosidase activated by monovalent cations and its application for efficient synthesis of alpha-D-glucosylglycerol. *Appl. Environ. Microbiol.*, *78*, 1836-1845.

(25) Liu, L.; Chen, X. F.; Ye, J. N.; Ma, X. Y.; Han, Y.; He, Y. J.; Tang, K. (2023) Sulfoquinovose is a widespread organosulfur substrate for Roseobacter clade bacteria in the ocean. *ISME J.*, *17*, 393-405.

Table 1: Michaelis-Menten kinetic parameters for *Ms*GH13 from *Marinovum* sp.

Substrate	k_{cat} (s^{-1})	K_M (mM)	k_{cat}/K_M (M^{-1} s^{-1}) × 10^3	Relative activity, (k_{cat}/K_M) %
PNPGlc	12.4±3.3	0.95±0.47	13±7.3	31
trehalose	18.0±0.9	2.13±0.43	8.5±1.8	20
kojibiose	53.2±5.1	34.2±7.5	1.6±0.4	3.8
nigerose	68.3±3	3.1±0.36	22±2.7	52
maltose	135±17	3.22±0.99	42±14	100
isomaltose	0.66±0.03	0.24±0.07	2.7±0.8	6.4
sucrose	61.2±10.6	18.8±7.1	3.3±1.4	7.9

Fig. 3 a) Sulfoglycolytic sulfoquinovose monooxygenase gene clusters from *Marinovum* sp. (MAI181XX.1) and *A. tumefaciens* (also known as *Agrobacterium fabrum*) strain C58. **b)** The sulfo-SMO pathway of *A. tumefaciens* for desulfurization of SQGro.

Fig. 2 Excerpt of ^{1}H-NMR (500 MHz, D$_2$O) of PNPGlc (top) and the reaction course of *Ms*GH13 catalyzed hydrolysis of PNPGlc at 10 min and 12 h. H1 of PNPGlc is present at δ 5.82 ppm. At *t* = 10 min, the initially formed signal at δ 5.15 ppm, $J_{1,2}$ 3.8 Hz, is assigned as H1 of α-glucose (middle). At *t* = 12 hours the α-glucose underwent mutarotation and formed a signal at δ 4.6 ppm, $J_{1,2}$ 8.0 Hz, assigned H1 of β-glucose (bottom).

Fig. 3 Subfamily GH13_23 *Ms*GH13 from *Marinovum* sp. is an α-glucosidase active on PNPGlc. **a)** pH dependence of k_{cat}/K_M using PNPGlc as substrate. **b)** Michaelis-Menten plot for PNPGlc as substrate. (inset) Lineweaver-Burk plot.

Fig. 4 Subfamily GH13_23 *Ms*GH13 from *Marinovum* sp. is an α-glucosidase with activity on sucrose, trehalose, nigerose, maltose, kojibiose and isomaltose. **a)** HPTLC analysis of *Ms*GH13 digests of various disaccharides. Products were visualized with 0.2% orcinol in 10% H_2SO_4, 10% H_2O in ethanol. Michaelis-Menten and (inset) Lineweaver-Burk plots for **b)** sucrose, **c)** trehalose, **d)** nigerose, **e)** maltose, **f)** kojibiose and **g)** isomaltose.

Fig. 5 Predicted structural basis of sugar recognition by *Ms*GH13. **a)** Overlay of *Ms*GH13 (AlphaFold predicted 3D structure, blue) and *Halomonas* sp. H11 glucosidase (HaG) E271Q mutant with maltose (PDB: 3WY4, green).[23] **b)** Surface presentation of *Ms*GH13 predicted structure with active site residues and overlay of maltose from HaG complex. **c)** Comparison of overlay of Alphafold prediction of *Ms*GH13 with Michaelis complex of HaG with maltose. The overlay reveals conservation of the key catalytic machinery, allowing assignment of catalytic nucleophile (*Ms*GH13/HaG: D201/D202), acid/base (E272/Q271 (corresponding to E271), respectively), and substrate-binding residues (D61/D62, H104/H105, R199/R200, H336/H332, D337/D333 and R410/R400, respectively).