Interspecies regulatory landscapes and elements revealed by novel joint systematic

integration of human and mouse blood cell epigenomes

Guanjue Xiang[1,2,3], Xi He[1], Belinda M. Giardine[4], Kathryn J. Isaac[5], Dylan J. Taylor[5], Rajiv C.

McCoy[5], Camden Jansen[4], Cheryl A. Keller[4], Alexander Q. Wixom[4], April Cockburn[4], Amber

Miller[4], Qian Qi[6], Yanghua He[6,7], Yichao Li[6], Jens Lichtenberg[8], Elisabeth F. Heuston[8], Stacie M.

Anderson[9], Jing Luan[10], Marit W. Vermunt[10], Feng Yue[11], Michael E.G. Sauria[12], Michael C.

Schatz[12], James Taylor[5,12], Berthold Göttgens[13], Jim R. Hughes[14], Douglas R. Higgs[14], Mitchell

J. Weiss[6], Yong Cheng[6], Gerd A. Blobel[10], David M. Bodine[8], Yu Zhang[15], Qunhua Li[15,16], Shaun

Mahony[4,16,17], Ross C. Hardison[4,16,17] *

[1]Bioinformatics and Genomics Graduate Program, Huck Institutes of the Life Sciences, The

Pennsylvania State University, University Park, PA 16802

[2]Department of Data Science, Dana-Farber Cancer Institute, Boston, MA 02215

[3]Department of Biostatistics, Harvard T.H. Chan School of Public Health, Boston, MA 02215

[4]Department of Biochemistry and Molecular Biology, The Pennsylvania State University,

University Park, PA 16802

[5]Department of Biology, Johns Hopkins University, Baltimore, MD 21218

[6]Department of Hematology, St. Jude Children's Research Hospital, Memphis, TN 38105

[7]Department of Human Nutrition, Food and Animal Sciences, University of Hawai`i at Mānoa,

Honolulu, HI 96822, USA

[8]Genetics and Molecular Biology Branch, National Human Genome Research Institute,

Bethesda, MD 20892

[9]Flow Cytometry Core, National Human Genome Research Institute, Bethesda, MD 20892

[10]Department of Pediatrics, Children's Hospital of Philadelphia, and Perelman School of

Medicine, University of Pennsylvania, Philadelphia, PA 19104

[11]Department of Biochemistry and Molecular Genetics, Feinberg School of Medicine,

Northwestern University, Evanston, IL 60611

[12]Department of Computer Science, Johns Hopkins University, Baltimore, MD 21218

[13]Welcome and MRC Cambridge Stem Cell Institute, University of Cambridge, Cambridge, UK

[14]MRC Weatherall Institute of Molecular Medicine, Oxford University, Oxford, UK

[15]Department of Statistics, The Pennsylvania State University, University Park, PA 16802

[16]Center for Computational Biology and Bioinformatics, Genome Sciences Institute, Huck

Institutes of the Life Sciences, The Pennsylvania State University, University Park, PA

16802

[17]Center for Eukaryotic Gene Regulation, The Pennsylvania State University, University Park,

PA 16802

* Corresponding author: Ross C. Hardison, Department of Biochemistry and Molecular Biology,

The Pennsylvania State University, 304 Wartik Lab, University Park, PA 16802, Phone: 814-

863-0113; E-mail: rch8@psu.edu

Running Title: Integrated epigenomic profiles between species

Key words: epigenetics, gene regulatory elements, dimensional reduction, regulatory potential

Abstract

Knowledge of locations and activities of *cis*-regulatory elements (CREs) is needed to decipher

basic mechanisms of gene regulation and to understand the impact of genetic variants on

complex traits. Previous studies identified candidate CREs (cCREs) using epigenetic features in

one species, making comparisons difficult between species. In contrast, we conducted an

interspecies study defining epigenetic states and identifying cCREs in blood cell types to

generate regulatory maps that are comparable between species, using integrative modeling of

eight epigenetic features jointly in human and mouse in our **V**ali**d**ated **S**ystematic **I**ntegrati**on**

(VISION) Project. The resulting catalogs of cCREs are useful resources for further studies of

gene regulation in blood cells, indicated by high overlap with known functional elements and

strong enrichment for human genetic variants associated with blood cell phenotypes. The

contribution of each epigenetic state in cCREs to gene regulation, inferred from a multivariate

regression, was used to estimate epigenetic state Regulatory Potential (esRP) scores for each

cCRE in each cell type, which were used to categorize dynamic changes in cCREs. Groups of

cCREs displaying similar patterns of regulatory activity in human and mouse cell types, obtained

by joint clustering on esRP scores, harbored distinctive transcription factor binding motifs that

were similar between species. An interspecies comparison of cCREs revealed both conserved

and species-specific patterns of epigenetic evolution. Finally, we showed that comparisons of

the epigenetic landscape between species can reveal elements with similar roles in regulation,

even in the absence of genomic sequence alignment.

Introduction

The morphology and functions of different cell types are determined by the expression of distinctive sets of genes in each. This differential gene expression is regulated by the interplay of transcription factors (TFs) binding to *cis*-regulatory elements (CREs) in the genomic DNA, such as promoters and enhancers, forging interactions among the CREs and components of transcriptional apparatus and ultimately leading to patterns of gene activation and repression characteristic of each cell type (Maston et al. 2006; Hamamoto and Fukaya 2022). Epigenetic features such as accessibility of DNA and modifications of histone tails in chromatin have pronounced impacts on the ability of TFs to bind to CREs, and furthermore, they serve as a molecular memory of transcription and repression (Strahl and Allis 2000; Ringrose and Paro 2004). Frequently co-occurring sets of chromatin features define epigenetic states, which are associated with gene regulation and expression (Ernst and Kellis 2010; Hoffman et al. 2013; Zhang et al. 2016). Genome-wide assignment of DNA intervals to epigenetic states (annotation) provides a view of the regulatory landscape that can be compared across cell types, which in turn leads to insights into the processes regulating gene expression (Libbrecht et al. 2021).

Comprehensive mapping of CREs within the context of the regulatory landscape in different cell types is needed to achieve a broad understanding of differential gene expression. Maps of candidate CREs (cCREs) provide guidance in understanding how changes in cCREs, including single nucleotide variants and indels, can lead to altered expression (Hardison 2012), and they can inform approaches for activation or repression of specific genes in potential strategies for therapies (Bauer et al. 2013). Indeed, most human genetic variants associated with common traits and diseases are localized in or near cCREs (Hindorff et al. 2009; Maurano et al. 2012; The ENCODE Project Consortium 2012). Thus, knowledge of the activity and epigenetic state of

cCREs in each cell type can facilitate understanding the impact of trait-associated genetic

variants on specific phenotypes. Furthermore, genome editing approaches in somatic cells have

recently been demonstrated to have promise as therapeutic modalities (Frangoul et al. 2021),

and a full set of cCREs annotated by activity and state can help advance similar applications.

The different types of blood cells in humans and mice are particularly tractable systems for

studying many aspects of gene regulation during differentiation. The striking differences among

mature cell types result from progressive differentiation starting from a common hematopoietic

stem cell (HSC) (Kondo et al. 2003). While single cell analyses reveal a pattern of ostensibly

continuous expression change along each hematopoietic lineage (Laurenti and Göttgens 2018),

intermediate populations of multi-lineage progenitor cells with decreasing differentiation

potential have been defined, which provide an overall summary and nomenclature for major

stages in differentiation. These stem, progenitor, and mature cell populations can be isolated

using characteristic cell surface markers (Spangrude et al. 1988; Payne and Crooks 2002),

albeit with many fewer cells in progenitor populations. In addition to the primary blood cells,

several immortalized cell lines provide amenable systems for intensive study of various aspects

of gene regulation during differentiation and maturation of blood cells (Weiss et al. 1997).

The VISION project aims to produce a **V**ali**d**ated **S**ystematic **I**ntegrati**on** of hematopoietic

epigenomes, harvesting extensive epigenetic and transcriptomic datasets from many

investigators and large consortia into concise, systematically integrated summaries of regulatory

landscapes and cCREs (Hardison et al. 2020). We previously published the results of these

analyses for progenitor and mature blood cell types from mouse (Xiang et al. 2020). In the

current study, we generated additional epigenetic datasets and compiled data from human

blood cells to expand the integrative analyses to include data from both human and mouse. The

systematic integrative analysis of epigenetic features across blood cell types was conducted

jointly in both species to learn epigenetic states, generate concise views of epigenetic

landscapes, and predict regulatory elements that are comparable in both species. This joint

modeling enabled further comparisons using approaches that were not dependent on DNA

sequence alignments between species, including a demonstration of the role of orthologous

transcription factors in cell type-specific regulation in both species. An exploration of

comparisons of epigenetic landscapes between species showed that they were informative for

inferring regulatory roles of elements in lineage-specific (i.e., non-aligning) DNA. Together, this

work provides valuable community resources that enable researchers to leverage the extensive

existing epigenomic data into further mechanistic regulatory studies of both individual loci and

genome-wide trends in human and mouse blood cells.

Results

Extracting and annotating epigenetic states by modeling epigenomic information jointly

in human and mouse

A large number of data sets of epigenetic features related to gene regulation and expression

(404 data sets, 216 in human and 188 in mouse; Fig. 1, Supplemental Material "Data generation

and collection", Supplemental Tables S1 and S2) served as the input for our joint integrative

analysis of human and mouse regulatory landscapes across progenitor and mature blood cell

types. The features included chromatin accessibility, which is a hallmark of almost all regulatory

elements, occupancy by the structural protein CTCF, and histone modifications associated with

gene activation or repression. After normalizing and denoising these diverse data sets

(Supplemental Fig. S1), we conducted an iterative joint modeling to discover epigenetic states,

i.e., sets of epigenetic features commonly found together, in a consistent manner for both

human and mouse blood cells (Fig. 2). The joint modeling took advantage of the Bayesian

framework of the Integrative and Discriminative Epigenomic Annotation System, or IDEAS

(Zhang et al. 2016; Zhang and Hardison 2017), to iteratively learn states in both species. The

joint modeling proceeded in four steps: initial training on randomly selected regions in both

species, retaining the 27 epigenetic states that exhibit similar combinatorial patterns of features

in both human and mouse, using these 27 states as prior information to sequentially run the

IDEAS genome segmentation on the human and mouse data sets, and removal of two

heterogenous states (Fig. 2A and Supplemental Figs. S2, S3, S4, and S5). This procedure

ensured that the same set of epigenetic states was learned and applied for both species.

Previously, the segmentation and genome annotation (Libbrecht et al. 2021) method

ChromHMM (Ernst and Kellis 2012) was used to combine data between species by

concatenating the datasets for both human and mouse cell types (Yue et al. 2014). This earlier

approach produced common states between species, but it did not benefit from the positional

information and automated approach to handling missing data that are embedded in IDEAS.

The resulting model with 25 epigenetic states (Fig. 2B) was similar to that obtained from mouse

blood cell data (Xiang et al. 2020). The states captured combinations of epigenetic features

characteristic of regulatory elements such as promoters and enhancers, transcribed regions,

repressed regions marked by either Polycomb (H3K27me3) or heterochromatin (H3K9me3),

including states that differ quantitatively in the contribution of specific features to each state. For

example, H3K4me1 is the predominant component of states E1 and E, but E1 has a lower

contribution of that histone modification. Similar proportions of the genomes of human and

mouse were covered by each state (Fig. 2B).

Assigning all genomic bins in human and mouse to one of the 25 states in each hematopoietic

cell type produced an annotation of blood cell epigenomes that gave a concise view of the

epigenetic landscape and how it changes across cell types, using labels and color conventions

consistently for human and mouse. The value of this concise view can be illustrated in

orthologous genomic intervals containing genes expressed preferentially in different cell

lineages as well as genes that are uniformly expressed (Fig. 2C, D). For example, the gene

SLC4A1/*Slc4a1*, encoding the anion transporter in the erythrocyte plasma membrane, is

expressed in the later stages of erythroid maturation (Dore and Crispino 2011). The epigenetic

state assignments across cell types matched the differential expression pattern, with genomic

intervals in the gene and its flanking regions, including a non-coding gene located upstream (to

its right, *Bloodlinc* in mouse), assigned to states indicative of enhancers (yellow and orange)

and promoters (red) only in erythroid cell types, with indications of stronger activation in the

more mature erythroblasts (region boxed and labeled E in Fig. 2 C, D). A similar pattern was

obtained in both human and mouse. Those genomic intervals assigned to the enhancer- or

promoter-like states contain candidates for regulatory elements, an inference that was

supported by chromatin binding data including occupancy by the transcription factor GATA1 (Xu

et al. 2012; Pimkin et al. 2014) and the co-activator EP300 (ENCODE datasets ENCSR000EGE

and ENCSR982LJQ) in erythroid cells. Similarly, the gene and flanking regions for *GRN*/*Grn*,

encoding the granulin precursor protein that is produced at high levels in granulocytes and

monocytes (Jian et al. 2013), and *ITGA2B*/*Itga2b*, encoding the alpha 2b subunit of integrin that

is abundant in mature megakaryocytes (van Pampus et al. 1992; Pimkin et al. 2014), were

assigned to epigenetic states indicative of enhancers and promoters in the expressing cell types

(boxed regions labeled G and MK, respectively). In contrast, genes expressed in all the blood

cell types, such as *UBTF*/*Ubtf*, were assigned to active promoter states and transcribed states

across the cell types. We conclude that these concise summaries of the epigenetic landscapes

across cell types showed the chromatin signatures for differential or uniform gene expression

and revealed discrete intervals as potential regulatory elements, with the consistent state

assignments often revealing similar epigenetic landscapes of orthologous genes in human and

mouse.

While these resources are useful, some limitations should be kept in mind. For example, IDEAS

used data from similar cell types to improve state assignments in cell types with missing data,

but the effectiveness of this approach may be impacted by the pattern of missing data. In

particular, the epigenetic data on human stem and progenitor cell types were largely limited to

ATAC-seq data, whereas histone modification data and CTCF occupancy were available for the

analogous cell types in mouse (Fig. 1). Thus, the state assignments for epigenomes in human

stem and progenitor cells may be less robust compared to those for similar cell types in mouse.

Another limitation is the broad range of quality in the data sets that cannot be completely

adjusted by normalization, which leads to over- or under-representation of some epigenetic

signals in specific cell types (Supplemental Fig. S5). Despite these limitations, the annotation of

blood cell epigenomes after normalization and joint modeling of epigenetic states produced a

highly informative painting of the activity and regulatory landscapes across the genomes of

human and mouse blood cells.

Candidate *cis*-regulatory elements in human and mouse

We define a candidate *cis*-regulatory element, or cCRE, as a DNA interval with a high signal for

chromatin accessibility in any cell type (Xiang et al. 2020). We utilized a version of the IDEAS

methodology to combine peaks of accessibility across different cell types, running it in the signal

intensity state (IS) mode only on chromatin accessibility signals (Xiang et al. 2021), which helps

counteract excessive expansion of peak calls when combining them (Supplemental Fig. S6).

Employing the same peak-calling procedure to data from human and mouse resulted in 200,342

peaks of chromatin accessibility for human and 96,084 peaks for mouse blood cell types

(Supplemental Table S3). Applying the peak caller MACS3 (Zhang et al. 2008) on the same

human ATAC-seq data generated a larger number of peaks, but those additional peaks tended

to have low signal and less enrichment for overlap with other function-related genomic datasets

(Supplemental Fig. S7).

The ENCODE Project released regulatory element predictions in a broad spectrum of cell types

in the Index of DHSs (Meuleman et al. 2020) and the SCREEN cCRE catalog (The ENCODE

Project Consortium et al. 2020), using data that were largely different from those utilized for the

VISION analyses. Almost all the VISION cCRE calls in human blood cells were included in the

regulatory element predictions from ENCODE (Supplemental Fig. S8A), supporting the quality

of the VISION cCRE calls. Furthermore, as expected from its focus on blood cell types, the

VISION cCRE catalog shows stronger enrichment for regulatory elements active in blood cells

(Supplemental Fig. S8B, Supplemental Table S4).

Enrichment of the cCRE catalog for function-related elements and trait-associated

genetic variants

Having generated catalogs of cCREs along with an assignment of their epigenetic states in

each cell type, we characterized the human cCREs further by connecting them to orthogonal

(not included in VISION predictions) datasets of DNA elements implicated in gene regulation or

in chromatin structure and architecture (termed structure-related) (Fig. 3A, Supplemental Fig.

S9, Supplemental Table S5). About two-thirds (136,664 or 68%) of the VISION human cCREs

overlapped with elements in the broad groups of CRE-related (97,361 cCREs overlapped) and

structure-related (83,327 cCREs overlapped) elements, with 44,024 cCREs overlapping

elements in both categories (Fig. 3A, B). In contrast, ten sets of randomly chosen DNA intervals,

matched in length and GC-content with the human cCRE list, showed much less overlap with

the orthogonal sets of elements (Fig. 3B). Of the CRE-related superset, the enhancer-related

group of datasets contributed the most overlap with VISION cCREs, followed by SuRE peaks,

which measure promoter activity in a massively parallel reporter assay (van Arensbergen et al.

2017), and CpG islands (Fig. 3C). Compared to overlaps with the random matched intervals, the

VISION cCREs were highly enriched for overlap with each group of CRE-related datasets (Fig.

3C). Of the structure-related superset, the set of CTCF occupied segments (OSs) contributed

the most overlap, followed by chromatin loop anchors, again with high enrichment relative to

overlaps with random matched sets (Fig. 3D). Considering the VISION cCREs that intersected

with both structure- and CRE-related elements, major contributors were the cCREs that overlap

with enhancers and CTCF OSs or loop anchors (Supplemental Fig. S10). Furthermore, the

VISION cCREs captured known blood cell CREs (Supplemental Table S4) and CREs

demonstrated to impact a specific target gene in a high throughput analysis (Gasperini et al.

2019) (Fig. 3E). We conclude that the intersections with orthogonal, function- or structure-

related elements lent strong support for the biological significance of the VISION cCRE calls

and added to the annotation of potential functions for each cCRE.

The catalog of VISION human blood cell cCREs showed a remarkable enrichment for genetic

variants associated with blood cell traits, further supporting the utility of the catalog. We initially

observed a strong enrichment by overlap with variants from the NHGRI-EBI GWAS Catalog

(Buniello et al. 2019) associated with blood cell traits (Supplemental Fig. S11). We then

analyzed the enrichments while considering the haplotype structure of human genomes,

whereby association signals measured at assayed genetic markers likely reflect an indirect

effect driven by linkage disequilibrium (LD) with a causal variant (that may or may not have

been genotyped). We employed stratified linkage disequilibrium score regression (sLDSC,

Finucane et al. 2015) to account for LD structure and estimate the proportion of heritability of

each trait explained by a given genomic annotation, quantifying the enrichment of heritability in

587 traits from the UK Biobank (UKBB) GWAS (Ge et al. 2017 and http://www.nealelab.is/uk-

biobank/) within the VISION cCREs relative to the rest of the genome (Supplemental Material

section "Stratified linkage disequilibrium score regression"). These traits encompassed 54

282 "blood count" traits that measure properties including size and counts of specific blood cell

283 types, 60 "blood biochemistry" traits that measure lipid, enzyme, and other molecular

284 concentrations within whole blood samples, and 473 non-blood-related traits, allowing us to

285 assess the specific relevance of the cCREs to regulation of blood-related versus other

286 phenotypes. At a 5% FDR threshold, we discovered 53 traits for which cCREs were significantly

287 enriched in heritability (Fig. 3F). Of these traits, 52 (98%) were blood-related and 50 were blood

288 count traits, representing 93% of all UKBB blood count traits included in our analysis. The

289 remaining 2 significant traits pertained to blood biochemistry, specifically, the male and female

290 glycated hemoglobin concentrations. These metrics and observations together lend support to

291 the VISION cCRE annotation being composed of informative genomic regions associated with

292 regulation of genes involved in development of blood cell traits.

294 **Estimates of regulatory impact of cCREs during differentiation**

295 The epigenetic states assigned to cCREs can reveal those that show changes in apparent

296 activity during differentiation. Inferences about the activity of a cCRE in one or more cell types

297 are based on whether the cCRE was actuated, i.e., was found in a peak of chromatin

298 accessibility, and which epigenetic state was assigned to the actuated cCRE. Those states can

299 be associated with activation (e.g., enhancer-like or promoter-like) or repression (e.g.,

300 associated with polycomb or heterochromatin). In addition to these categorical state

301 assignments, quantitative estimates of the impact of epigenetic states on expression of target

302 genes are useful, e.g., to provide an estimate of differences in inferred activity when the states

303 change. Previous work used signals from single or multiple individual features such as

304 chromatin accessibility or histone modifications in regression modeling to explain gene

305 expression (e.g., Karlić et al. 2010; Dong et al. 2012), and we applied a similar regression

306 modeling using epigenetic states as predictor variables to infer estimates of regulatory impact of

307 each state on gene expression (Xiang et al. 2020).

We used state assignments of cCREs across cell types in a multivariate regression model to

estimate the impact of each state on the expression of local genes (Supplemental Material,

"Estimation of the impact of epigenetic states and cCREs on gene expression"). That impact

was captured as β coefficients, which showed the expected strong positive impact for promoter

and enhancer associated states and negative impacts from heterochromatin and polycomb

states (Fig. 4A). The β coefficients were then used in further analysis, such as estimating the

change in regulatory impact as a cCRE shifts between states during differentiation (difference

matrix to the left of the β coefficient values in Fig. 4A). The β coefficient values also were used

to generate an epigenetic state Regulatory Potential (esRP) score for each cCRE in each cell

type, calculated as the β coefficient values for the epigenetic states assigned to the cCRE

weighted by the coverage of the cCRE by each state (Fig. 4B). These esRP scores were the

basis for visualizing the collection of cCREs and how their regulatory impact changed across

differentiation (Supplemental Fig. S12 and Supplemental movie S1). Comparison of the

integrative esRP scores with signal intensities for single features (ATAC-seq and H3K27ac)

showed all were informative for visualizations, and esRP performed slightly better than the

single features in differentiating cCREs based on locations within gene bodies (Supplemental

Fig. S13).

In addition, we explored the utility of the esRP scores for clustering the cCREs into groups with

similar activity profiles across blood cell types in both human and mouse. Focusing on the esRP

scores in 12 cell types shared between human and mouse along with the average across cell

types, we identified clusters jointly in both species. The clustering proceeded in three steps,

specifically finding robust *k*-means clusters for the combined human and mouse cCREs,

identifying the clusters shared by cCREs in both species, and then further grouping those

shared *k*-means clusters hierarchically to define fifteen joint metaclusters (JmCs) (Supplemental

Fig. S14). Each cCRE in both mouse and human was assigned to one of the fifteen JmCs, and

each JmC was populated with cCREs from both mouse and human.

These JmCs established discrete categories for the cCREs based on the cell type distribution of

their inferred regulatory impact (Fig. 4C). The clusters of cCREs with high esRP scores across

cell types were highly enriched for promoter elements (Supplemental Fig. S15A). The cell type-

restricted clusters of cCREs showed enrichment both for selected enhancer catalogs and for

functional terms associated with those cell types (Supplemental Fig. S15A and B). Furthermore,

clustering of human genes by the JmC assignments of cCREs in a 100kb interval centered on

their TSS (Supplemental Material section "Enrichment of JmCs assigned to cCREs in gene

loci") revealed a strong enrichment for JmCs with high activity in the cell type(s) in which the

genes are expressed (Fig. 4D). Examples include *IFNG* showing enrichment for JmC 12, which

has high esRP scores in T and NK cells, *CSF1R* showing enrichment for JmC 15, which has

high scores in monocytes, and *GATA1* showing enrichment for JmC 10, which has high scores

in erythroid cells and megakaryocytes. Moreover, running sLDSC on cCREs in individual JmCs

showed enrichment for heritability of blood cell-related traits in some specific JmCs

(Supplemental Fig. S16).

As expected from previous work (e.g., Heintzman et al. 2009; Meuleman et al. 2020), similar

metaclusters of cCREs were generated based on single signals from the histone modification

H3K27ac or chromatin accessibility across cell types (Supplemental Fig. S17). Clustering based

any of the three features better resolved individual cell types when larger numbers of clusters

were considered, prior to collapsing the shared robust clusters into JmCs (Supplemental Fig.

S18).

In summary, we show that the β coefficients and esRP scores provide valuable estimates of

regulatory impacts of states and cCREs, respectively. The esRP-driven joint metaclusters

provide refined subsets of cCREs that should be informative for investigating cell type-specific

and general functions of cCREs. We also built self-organizing maps as a complementary

approach to systematic integration of epigenetic features and RNA data across cell types

(Supplementary Figure S10, Jansen et al. 2019).

Motif enrichment in joint metaclusters of human and mouse cCREs

We examined the sets of cCREs in each JmC to ascertain enrichment for transcription factor

binding site (TFBS) motifs because these enriched motifs suggest the families of transcription

factors that play a major role in regulation by each category of cCREs. Furthermore, having sets

of cCREs determined and clustered for comparable blood cell types in human and mouse

provided the opportunity to discover which TFBS motifs were shared between species and

whether any were predominant in only one species.

To find TFBS motifs associated with each JmC, we calculated enrichment for all non-redundant

motifs in the Cis-BP database (Weirauch et al. 2014) using Maelstrom from GimmeMotifs

(Bruse and van Heeringen 2018) (Supplemental Material "Enrichment for transcription factor

binding site motifs in joint metaclusters of cCREs"). The results confirmed previously

established roles of specific TFs in cell lineages and showed little evidence for novel motifs (Fig.

4E). For example, TFBS motifs for the GATA family of transcription factors were enriched in

JmCs 2 and 10, which have high esRP scores in progenitor and mature cells in the erythroid

and megakaryocytic lineages, as expected for the known roles of GATA1 and GATA2 in this

lineage (Blobel and Weiss 2009; Fujiwara et al. 2009). The GATA motif was also enriched in

JmC 14, as expected for the role of GATA3 in natural killer (NK) and T cells (Rothenberg and

Taghon 2005). Furthermore, motifs for the known lymphoid transcription factors TBX21,

TCF7L1, and LEF1 (Chi et al. 2009) were enriched in cCREs with high esRP scores in NK and

T cells (JmCs 9 and 12), and motifs for myeloid-determining transcription factors CEBPA and

CEBPB (Graf and Enver 2009) and the myeloid transcription factor PU.1 (Tenen et al. 1997)

were enriched in cCREs that are active in progenitor cells and monocytes (JmCs 3 and 15).

TFBS motifs for promoter-associated transcription factors such as E2F2 and SP1 (Dynan and

Tjian 1983; Kaczynski et al. 2003) were enriched in broadly active cCREs (JmCs 1 and 4).

These patterns of motif enrichments in the JmCs fit well with the expectations from previous

studies of transcription factor activity across lineages of blood cells, and thus, they lend further

credence to the value of the cCRE calls and the JmC groupings for further studies of regulation

in the blood cell types.

The genome-wide collection of cCREs across many blood cell types in human and mouse

provided an opportunity for an unbiased and large-scale search for indications of transcription

factors that may be active specifically in one species for a shared cell type. Prior studies of

transcription factors have shown homologous transcription factors used in analogous cell types

across species (e.g., Carroll 2008; Noyes et al. 2008; Schmidt et al. 2010; Cheng et al. 2014;

Villar et al. 2014), but it is not clear if there are significant exceptions. In our study, we found that

for the most part, the motif enrichments were quite similar between the human and mouse

cCREs in each JmC. Note that these similarities were not forced by requiring sequence

matches between species; the cCREs were grouped into JmCs based on their pattern of

activity, as reflected in the esRP scores, across cell types, not by requiring homologous

sequences. This similarity between species indicates that the same transcription factors tend to

be active in similar groups of cell types in both mouse and human. An intriguing potential

exception to the sharing of motifs between species was the enrichment of TFBS motifs for

CTCF and ZBTB7A in some JmCs, suggestive of some species selectivity in their binding in the

context of other TFs (Supplemental Figs. S20 and S21). These indications of conditional,

preferential usage of these TFs in human or mouse could serve as the basis for more detailed

studies in the future.

In summary, after grouping the cCREs in both human and mouse by their inferred regulatory

impact across blood cell in a manner agnostic to DNA sequence or occupancy by TFs, the

enrichment for TFBS motifs within those groups recapitulated known activities of TFs both

broadly and in specific cell lineages. The results also showed considerable sharing of inferred

TF activity in both human and mouse.

Evolution of sequence and inferred function of cCREs

The human and mouse cCREs from blood cells were assigned to three distinct evolutionary

categories (Fig. 5A). About one-third of the cCREs were present only in the reference species

(39% for human, 28% for mouse), as inferred from the failure to find a matching orthologous

sequence in whole-genome alignments with the other species. We refer to these as

nonconserved (N) cCREs. Of the two-thirds of cCREs with an orthologous sequence in the

second species, slightly over 30,000 were also identified as cCREs in the second species. The

latter cCREs comprise the set of cCREs conserved in both sequence and inferred function,

which we call SF conserved (SF) cCREs. Almost the same number of cCREs in both species

fall into the SF category; the small difference resulted from interval splits during the search for

orthologous sequences (Supplemental Fig. S22). The degree of chromatin accessibility in

orthologous SF cCREs was positively correlated between the two species (Supplemental Fig.

S23). The remaining cCREs (91,000 in human and 36,000 in mouse) were conserved in

sequence but not in an inferred function as a regulatory element, and we call them S conserved

(S) cCREs. The latter group could result from turnover of regulatory motifs or acquisition of

different functions in the second species.

The distributions of epigenetic states assigned to the blood cell cCREs in each of the three

evolutionary categories were similar between human and mouse, but those distributions differed

among evolutionary categories, with significantly more SF cCREs assigned to promoter-like

states than were S or N cCREs (Supplemental Fig. S24). Indeed, the SF cCREs tended to be

close to or encompass the TSSs of genes, showing a substantial enrichment in overlap with

TSSs compared to the overlap observed for all cCREs (Fig. 5B). Many of the S and N cCREs

were assigned to enhancer-like states (Supplemental Fig. S24D), giving a level of enrichment

for overlap with enhancer datasets comparable to that observed for the full set of cCREs (Fig.

5B).

For both human and mouse, the level of sequence conservation, estimated by the maximum

phyloP score (Pollard et al. 2010), was higher in the collection of cCREs than in sets of

randomly chosen genomic intervals matching the cCREs in length and G+C content (Fig. 5C).

Among the evolutionary categories of cCREs, the distribution of phyloP scores for SF cCREs

was significantly higher than the distribution for S cCREs, which in turn was higher than that for

N cCREs, for both species (Fig. 5C). The whole genome alignments underlying the phyloP

scores are influenced by proximity to the highly conserved coding exons (King et al. 2007), and

the high phyloP scores of the promoter-enriched SF cCREs could reflect both this effect as well

as strong constraint on conserved function (Supplemental Fig. S25). In all three evolutionary

categories, the distribution of phyloP scores was higher for promoter-proximal cCREs than for

distal ones, but the relative levels of inferred conservation were the same for both, i.e., SF>S>N

(Supplemental Fig. S26).

In summary, this partitioning of the cCRE catalogs by conservation of sequence and inferred

function revealed informative categories that differed both in evolutionary trajectories and in

types of functional enrichment.

Conservation of non-coding genomic DNA sequences among species has been used

extensively to predict regulatory elements (Gumucio et al. 1992; Hardison 2000; Pennacchio

and Rubin 2001), but the observation that predicted regulatory elements fall into distinct

evolutionary categories (SF, S, and N) raised the question of whether inter-species DNA

sequence alignments or annotation of epigenetic states would be more effective in finding

elements that were experimentally determined to be active in gene regulation. Recent advances

in massively parallel reporter assays have enabled the testing of large sets of candidate

elements, approaching comprehensive assessment of the predicted elements (Agarwal et al.

2023). We used the set of over 57,000 human genomic elements shown to be active in K562

cells to address this question (Supplemental Material), and we found that requiring alignment to

the mouse genome would miss about 40% of the active elements, whereas requiring presence

in a non-quiescent epigenetic state or one associated with gene activation would cover 87% or

82.5%, respectively, of the active elements (Fig. 5D). Thus, the epigenetic state annotation can

enable a more comprehensive prediction and examination or gene regulatory elements. This

realization motivated a comparison of epigenetic states between human and mouse, as

described in the next section.

Comparison of epigenetic states around orthologous genes in human and mouse

The consistent state assignments from the joint modeling facilitated epigenetic comparisons

between species. Such comparisons are particularly informative for orthologous genes with

similar expression patterns but some differences in their regulatory landscapes. For example,

the orthologous genes *GATA1* in human and *Gata1* in mouse each encode a transcription factor

with a major role in regulating gene expression in erythroid cells, megakaryocytes, and

eosinophils (Ferreira et al. 2005), with a similar pattern of gene expression across blood cell

types in both species (Supplemental Fig. S27). The human and mouse genomic DNA

sequences aligned around these orthologous genes, including their promoters and proximal

enhancers; the alignments continued through the genes downstream of *GATA1*/*Gata1* (Fig. 6A).

An additional, distal regulatory element located upstream of the mouse *Gata1* gene, which was

bound by GATA1 and EP300 (Fig. 6A), was found only in mouse (Valverde-Garduno et al.

2004). The DNA sequences of the upstream interval harboring the mouse regulatory element

did not align between mouse and human except in portions of the *GLOD5*/*Glod5* genes (Fig.

6A). Thus, the interspecies sequence alignments provide limited information about this distal

regulatory element.

This limitation to sequence alignments led us to explore whether comparisons of epigenetic

information would be more informative, utilizing the consistent assignment of epigenetic states

in both human and mouse, which do not rely on DNA sequence alignment. In the large genomic

regions (76kb and 101kb in the two species) encompassing the orthologous human *GATA1* and

mouse *Gata1* genes and surrounding genes, we computed the correlation for each genomic bin

between the epigenetic state assignments across cell types in one species and that in the other

species for all the bins (Supplemental Fig. S28). This local, all-versus-all comparison of the two

loci yielded a matrix of correlation values showing similarities and differences in profiles of

epigenetic states in the two species (Fig. 6B). The conserved promoter and proximal enhancers

of the *GATA1*/*Gata1* genes were highly correlated in epigenetic states across cell types

between the two species, in a region of the matrix that encompassed the aligning DNA

sequences (labeled Px in Fig. 6B). In contrast, whereas the mouse-specific distal regulatory

element did not align with the human DNA sequence, the epigenetic states annotating it

presented high correlations with active epigenetic states in the human *GATA1* locus (labeled D

in Fig. 6B).

The complexity of the correlation matrix (Fig. 6B) indicated that multiple epigenetic trends could

be contributing to the patterns. To systematically reduce the high dimensionality of the matrix to

a set of simpler matrices, we employed nonnegative matrix factorization (NMF) because of its

interpretability (Stein-O'Brien et al. 2018; Lee and Roy 2021). The decomposed matrices from

NMF revealed a set of factors, each of which (represented by each column in the mouse matrix

and each row in the human matrix in Fig. 6C) captures a group of highly correlated elements in

the original matrix that show a pattern distinct from the rest of the elements. The complex

correlation matrix was decomposed into six distinct factors, as determined by the number of

factors at which an "elbow" was found in the BIC score (Supplemental Fig. S29). Each factor

encapsulated a specific epigenetic regulatory machinery or process exhibiting consistent cross-

cell type patterns in both humans and mice (Supplemental Fig. S30). For example, the

correlation matrices reconstructed by using signals from factor 3 exclusively highlighted the

positive regulators for the *GATA1*/*Gata1* gene loci; these regulatory elements were evident in

reconstructed correlation matrices between species (Fig. 6D) and within individual species (Fig.

6E). By applying a *Z*-score approach to identify peak regions in the factor 3 signal vector (with

FDR < 0.1; Supplemental Material), we pinpointed regions in both species showing an

epigenetic regulatory machinery exhibiting positive regulatory dynamics for *GATA1*/*Gata1* gene

loci, particularly in the ERY and MK cell types. In contrast, the correlation matrices

reconstructed from the signals for factor 6 (Fig. 6F and G) highlighted regions marked by the

transcription elongation modification H3K36me3 (epigenetic states colored green, Fig. 6G). The

correlations in the factor 6 elongation signature were observed, as expected, between the

human/mouse orthologous gene pairs *GATA1* and *Gata1* as well as between human *HDAC6*

and mouse *Hdac6* (green rectangles in Fig. 6F). The factor 6 correlations were also observed

between the *GATA1*/*Gata1* and *HDAC6*/*Hdac6* genes (black rectangles in Fig. 6F and G),

showing a common process, specifically transcriptional elongation, at both loci. A similar

analysis for other factors revealed distinct regulatory processes or elements, such as active

promoters (factor 2), exhibiting unique cross-cell type patterns (Supplemental Fig. 30). The

genomic bins with high scores for a given NMF factor in human showed high correlation with

bins with high scores for that same factor in mouse, indicating that the NMF factors capture a

similar set of epigenetic state patterns in each species (Supplemental Fig. S31). The patterns

captured by NMF factors 3 and 6 were robust to the choice of k in the NMF (Supplemental Fig.

S32). Overall, these results underscore this method's capability to objectively highlight

regulatory regions with analogous epigenetic patterns across cell types in both species. This

method could aid in extracting additional information about similar epigenetic patterns between

human and model organisms such as mice, for which only a portion of their genome aligns with

human.

Because some of the NMF factors reflected processes in gene expression and regulation that

occur in many genes, some of the highly correlated regions across species could reflect false

positives. Thus, it is prudent to restrict the current approach to genomic intervals around

orthologous genes to reduce the impact of false discovery. We examined patterns of epigenetic

state correlations across cell types between the human *GATA1* gene locus and three non-

orthologous loci in mouse to investigate the scope of this issue (Supplemental Material). While

genomic bins of high epigenetic state correlation were observed between non-orthologous loci,

the discovery of bins implicated in a cell type-specific process, such as erythroid or

megakaryocytic regulation, could be enhanced by utilizing a broader background model for

computing peaks of NMF signal (Supplemental Fig. S33). With this refined approach to peak

identification, the false discovery rate estimated for epigenetic state comparison between the

human *GATA1* locus and the mouse *Cd4* locus was reduced to 0.1 or less (Supplemental Fig.

S33R). Furthermore, the epigenetic state comparisons between the human *GATA1* locus and

the mouse *Rps19* locus revealed a previously unreported region with hallmarks of erythroid

regulatory elements (Supplemental Fig. S34). These initial results suggest that the genomic

scale of the epigenetic state correlations could be expanded in future work with judicious

attention to reducing false discovery, e.g., by linking the discovered elements to evidence of

conserved synteny between species.

Examination of human genomic elements shown to be active in a lentiMPRA assay (Agarwal et

al. 2023) at 30 loci revealed that the active elements were highly enriched in genomic bins with

high cross cell-type epigenetic state correlation between species (Supplemental Fig. S35). The

enrichment for active elements was further increased in bins with both high epigenetic state

correlation and interspecies sequence conservation, while enrichment was reduced in bins with

only sequence conservation. These results further support the value of the cross cell-type

epigenetic state correlation between species in predicting and interpreting cCREs.

The comparison of epigenetic state profiles across cell types also provided a means to

categorize cCREs between species that did not require a match in the underlying genomic DNA

sequence (Supplemental Figs. S36 and S37). Results from that approach indicated that certain

cCREs were potentially involved in regulation of orthologous genes, even for cCREs with DNA

sequences that did not align between species.

In summary, the IDEAS joint modeling on the input data compiled here and the consistent state

assignments in both mouse and human confirmed and extended previous observations on

known regulatory elements, and they revealed both shared and distinctive candidate regulatory

elements and states between species. Correlations of state profiles between species provided a

comparison of chromatin landscapes even in regions with DNA sequences that were not

conserved between species. Our initial results reported here support continuing the

development of this approach of comparing cross cell-type epigenetic state profiles between

species for functional prediction and interpretation of cCREs.

Discussion

In this paper, the VISION consortium introduces a set of resources describing the regulatory

landscapes of both human and mouse blood cell epigenomes. A key, novel aspect of our work

is that the systematic integrative modeling that generated these resources was conducted jointly

across the data from both species, which enabled robust comparisons between species without

being limited by sequence alignments, allowing comparisons in non-conserved and lineage-

specific genomic regions.

One major resource is the annotation of the epigenetic states across the genomes of progenitor

and mature blood cells of both species. These state maps show the epigenetic landscape in a

compact form, capturing information from the input data on multiple histone modifications, CTCF

occupancy, and chromatin accessibility, and they use a common set of epigenetic states to

reveal the patterns of epigenetic activity associated with gene expression and regulation both

across cell types and between species. A second major resource is a catalog of cCREs

actuated in one or more of the blood cell types in each species. The cCREs are predictions of

discrete DNA segments likely involved in gene regulation, based on the patterns of chromatin

accessibility across cell types, and the epigenetic state annotations suggest the type of activity

for each cCRE in each cell type, such as serving as a promoter or enhancer, participating in

repression, or inactivity. A third major resource is a quantitative estimate of the regulatory

impact of human and mouse cCREs on gene expression in each cell type, i.e., an esRP score,

derived from multivariate regression modeling of the epigenetic states in cCREs as predictors of

gene expression. The esRP scores are a continuous variable capturing not only the integration

of the input epigenetic data, but also the inferred impacts on gene expression. Those impacts

may be manifested as activation or repression during regulation or as transcriptional elongation.

They are useful for many downstream analyses, such as determining informative groups of

cCREs by clustering analysis. These resources along with browsers for visualization and tools

for analysis are provided at our project website, http://usevision.org. Among these tools is

cCRE_db, which records the several dimensions of annotation of the cCREs and provides a

query interface to support custom queries from users.

Our human blood cell cCRE catalog should be valuable for mechanistic interpretations of trait-

related human genetic variants. Human genetic variants associated with traits intrinsic to blood

cells were significantly enriched in the VISION cCRE catalog, whereas variants associated with

a broad diversity of other traits were not enriched. We expect that the extensive annotations in

our cCRE catalog combined with information about TFBS motifs and TF occupancy should lead

to specific, refined hypotheses for mechanisms by which a variant impacts expression, such as

alterations in TF binding, which can be tested experimentally in further work.

The jointly learned state maps and cCRE predictions allowed us to extend previous work on the

evolution of regulatory elements between mouse and human. Several previous studies focused

on transcription factor (TF) occupancy, e.g. examining key TFs in one tissue across multiple

species (Schmidt et al. 2010; Ballester et al. 2014; Villar et al. 2014) or a diverse set of TFs in

multiple cell types and in mouse and human (Cheng et al. 2014; Yue et al. 2014; Denas et al.

2015). Other studies focused on discrete regions of high chromatin accessibility in multiple cell

types and tissues between mouse and human (Stergachis et al. 2014; Vierstra et al. 2014).

These previous studies revealed that only a small fraction of elements was conserved both in

genomic sequence and in inferred function. A notable fraction of elements changed

considerably during mammalian diversification, including turnover of TF binding site motifs and

repurposing of elements (Schmidt et al. 2010; Cheng et al. 2014; Stergachis et al. 2014; Denas

et al. 2015). These prior studies focused primarily on regions of the genome with sequences

that aligned between human and mouse, with the non-aligning regions used to infer that some

elements were lineage-specific and that many were derived from transposable elements and

endogenous retroviruses (Bourque 2009; Rebollo et al. 2012; Jacques et al. 2013; Sundaram et

al. 2014).

Our evolutionary analyses confirmed the previous observations, e.g., finding about one-third of

cCREs are conserved in both sequence and inferred function between human and mouse, and

further showing that this evolutionary category was highly enriched for proximal regulatory

elements. Going beyond the prior studies, our jointly learned epigenetic state maps generated a

representation of multiple epigenetic features, not just TF occupancy or chromatin accessibility,

and they are continuous in bins across genomes of both species. Thus, they provided a basis

for comparisons of the epigenetic profiles between species. These epigenetic comparisons were

a strong complement to genomic sequence alignments, allowing us to find elements with similar

epigenetic profiles even in genomic regions in which the DNA sequence does not align between

species. In the current work, we used both a correlation between profiles of epigenetic states

and joint clusterings of cCREs between species by esRP scores as initial explorations of these

epigenetic comparisons. Previous work compared epigenetic profiles across species, such as

the phylo-HMGP method to find different evolutionary states in multi-species epigenomic data

(Yang et al. 2018) and the LECIF scores to find evidence of conservation from functional

genomic data (Kwon and Ernst 2021). These approaches are powerful but limited to the

genomic regions with DNA sequences that align between the species. In contrast, our approach

of correlating epigenetic states is agnostic to the underlying DNA sequence alignments (or

absence of them), and thus it complements traditional approaches that rely on DNA sequence

alignments to find similar elements. Our inter-species comparisons of loci surrounding pairs of

orthologous genes included both DNA segments that align between human and mouse and

those that do not. Our detection, even in segments of DNA that do not align between species, of

epigenetic similarity indicative of a common role in gene regulation suggests that processes or

structures, such as chromatin interactions, chromatin complexes, or molecular condensates,

may be maintained between species in a manner that is not fully revealed by comparisons of

genome sequences. Hence, further studies of this apparent epigenetic dimension of regulatory

conservation may be productive.

Several innovations were developed to produce the resources introduced here. A major

innovation was to extend the IDEAS framework (Zhang et al. 2016) to jointly learn epigenetic

states and assign them to annotate the epigenomes in human and mouse blood cells. The

IDEAS method employs a Bayesian approach to the modeling to learn the states, which we

utilized to bring in states learned from the data in one species as priors for learning states in the

data from the second species. Another extension of the IDEAS framework was to learn states

based on one feature, specifically ATAC-seq data, defining discrete signal intensity states. This

approach was used for calling cCREs, implemented as the IDEAS-IS method (Xiang et al.

2021). The approach is relatively simple and benefits from joint modeling across the input

datasets. Other methods for predicting cCREs based on chromatin accessibility across many

cell types prevented excessive expansion of the summary calls for overlapping peaks by

employing a centroid determination for the DNase hypersensitive sites (DHS) index (Meuleman

et al. 2020) or by choosing the highest signal peak for the ENCODE cCRE catalog (The

ENCODE Project Consortium et al. 2020). The ENCODE cCRE catalog paired DHS peaks with

individual chromatin modifications or CTCF occupancy, which led to complications when data

on diagnostic features were missing from some cell types. The IDEAS framework used for the

VISION cCRE sets leveraged data in related cell types to ameliorate the impact of missing data.

While the resources introduced here are valuable for many applications, it is prudent to

acknowledge their limitations. First, the quality of the products of integrated analyses are limited

by the quality and completeness of the input, raw data. We endeavored to reduce the impact of

variances in the input data by normalization. The S3V2 procedure (Xiang et al. 2021)

systematically normalized the input data to adjust for differences in signal-to-noise and variance

in signal across the datasets. Some epigenetic features were not determined in some cell types,

and we used the IDEAS method in part because it is able to assign an epigenetic state even in

the context of missing data by learning patterns from local similarities in cell types for which the

data are present (Zhang and Mahony 2019). However, these approaches cannot completely

overcome all issues with variance in input data, and further developments in these directions

(such as Shahraki et al. 2023; Xiang et al. 2023) may help to improve integrative resources.

Second, the resolution of both the epigenetic state assignments and the cCRE inference is

limited to 200 bp, which is the window size we utilized in the IDEAS analyses. Other resources,

such as DHS calls (Meuleman et al. 2020), DNase footprints (Vierstra et al. 2020), and motif

instances (Weirauch et al. 2014), achieve a higher resolution. Indeed, one can use these higher

resolution datasets to derive further information about cCREs, such as families of TFs that are

likely to be binding to them. Regarding esRP scores, a third limitation is that we do not make

explicit assignments for target genes of cCREs. Predictions of a large number of target gene-

cCRE pairs were made in our prior work (Xiang et al. 2020); these assignments cover large

genomic intervals around each gene and are most useful when used with further filtering, such

as restricting cCREs and target genes to the same topologically associated domains. On-going

work is examining other models and approaches for assigning likely target genes to cCREs. A

fourth limitation is that our inference of repression-related cCREs apply only to those with stable

histone modifications. Elements that had been involved in initiation of repression but eventually

were packaged into quiescent chromatin, e.g., via a hit-and-run mechanism (Shah et al. 2019),

would not be detected. A fifth limitation concerns the scale of the studies of epigenetic

conservation by correlations of epigenetic states. Our current approach is limited to individual

examination of specific genetic loci, since we used orthologous genes as the initial anchors, and

it is likely that a direct application to whole chromosomes or genomes would generate high false

discovery. Exploring ways to expand the scale of the analytical approach is a goal of future

research. Finally, the work presented here was restricted to blood cell types. In future work,

extension of the approaches developed in this study to a broader spectrum of cell types would

expand the utility of the resulting resources.

In conclusion, we present several important new resources to enable further and more detailed

studies of gene regulation in human and mouse blood cells both during normal differentiation

and in pathological contexts. The patterns of epigenetic states in cCREs across cell types show

value in developing an understanding of how genetic variants impact blood cell traits and

diseases. Furthermore, the joint modeling between species opens avenues for further

exploration of comparisons of epigenetic landscapes in addition to sequence alignments for

insights into evolution and function of regulatory elements between species.

Methods

Data generation, collation, normalization, and integration

The data sets used as input, including the ones generated for the work reported here (with

methods), are described in Supplemental Material section "Data generation and collection" and

Supplemental Tables S1 and S2. The S3V2 approach (Xiang et al. 2021) was used for

normalization and denoising the data sets prior to integration. The data sets were integrated to

find and assign epigenetic states using IDEAS (Zhang et al. 2016; Zhang and Hardison 2017);

the extension of this approach to joint learning and annotation between species is described in

Supplemental Material sections "Data normalization" and "Joint systematic integration of human

and mouse blood cell epigenomes by IDEAS".

Prediction, annotation, and estimation of regulatory impact of cCREs

The identification of cCREs as peaks of chromatin accessibility employed IDEAS in the signal

intensity state (IS) mode (Xiang et al. 2021). This approach and comparisons with MACS peaks

(Zhang et al. 2008) are described in Supplemental Material section "Prediction of VISION

cCREs using IDEAS-IS". The cCREs are provided in Supplemental Table S3. Annotation of

potential cCRE functions used intersections with orthogonal data sets of elements implicated in

regulation or chromatin structure (Supplemental Table S5). Enrichment of genetic variants

associated with blood cell traits used stratified linkage disequilibrium score regression (sLDSC,

Finucane et al. 2015). The impact of epigenetic states in cCREs on regulation of gene

expression used a multivariate linear regression approach like one described previously (Xiang

et al. 2020). Methods and supplementary results on these analyses are presented in detail in

the Supplemental Material.

Identification of clusters of cCREs based on epigenetic regulatory potential scores

The sets of human and mouse cCREs were placed jointly into groups based on their epigenetic

regulatory potential (esRP) scores using a series of k-means clustering steps, as described in

detail in Supplemental Material and Supplementary Fig. S14. Methods and results for

enrichment of the resulting joint meta-clusters (JmCs) for orthogonal sets of regulatory elements

and SNPs associated with blood cell traits, along with comparisons of clusters based on

chromatin accessibility and H3K27ac signal, are described in Supplemental Material and

Supplementary Figs. S15 - S18. Motifs that were differentially enriched across JmCs were

identified using the Maelstrom tool in the GimmeMotifs suite (v0.17.1) (Bruse and van

Heeringen 2018) and SeqUnwinder (Kakumanu et al. 2017), as described in detail in

Supplemental Material and Supplementary Fig. S21.

Partitioning cCREs to evolutionary categories based on DNA sequence alignments and

cCRE calls between species

The human and mouse cCREs were assigned to three evolutionary categories using the

following procedure. The set of human cCREs was mapped to mouse genome assembly mm10

using the liftOver tool at the UCSC Genome Browser (Hinrichs et al. 2006). Human cCREs that

failed to map to mm10 were grouped as N cCREs. Matches to mouse cCREs for the human

cCREs that could be mapped by liftOver to mm10 were determined using the intersect tool in

BEDTools (Quinlan and Hall 2010). Human cCREs that overlapped with mouse cCREs were

labeled as SF cCREs, while human cCREs that mapped to mm10 but did not match mouse

cCREs were labeled as S cCREs. A similar process was performed on the set of mouse cCREs

using liftOver to map to human genome build GRCh38

Calculation of pairwise correlation coefficients for epigenetic landscapes between

human and mouse

A bin-to-bin pairwise correlation analysis was used to quantify the similarity of epigenetic

landscapes between two DNA regions in human and mouse. For each 200bp bin in one cell

type in one species, the assigned epigenetic state was replaced by a vector of mean signals of

8 epigenetic features in the IDEAS state model. After replacing the states in all 15 matched cell

types (14 analogous cell types and one pseudo-cell type with average values for all cell types)

in the two species, the original two categorical state vectors with 15 elements were converted

into two numeric vectors with 120 numbers (Supplemental Fig. S28). The similarity of cross-cell

type epigenetic landscape between two bins in the two species was defined as the correlation

coefficient between each pair of numeric vectors with 120 numbers. When calculating the

correlation coefficients, we added random noise (mean=0, sd=0.2) to the raw values to avoid

high correlation coefficients created between regions with states that have low signals. The

complex correlation matrix was decomposed into distinctive factors using Nonnegative Matrix

Factorization (Lee and Seung 1999). Methods and supplementary results on these analyses are

presented in detail in the Supplemental Material.

802   Data access

All raw and processed sequencing data generated in this study have been submitted to the

NCBI Gene Expression Omnibus (GEO; https://www.ncbi.nlm.nih.gov/geo/) under accession

number GSE229101 and the NCBI BioProject database

(https://www.ncbi.nlm.nih.gov/bioproject/) under accession number PRJNA952902. Resources

developed in the VISION project are available at the website https://usevision.org; the data can

be viewed via a track hub at the UCSC Genome Browser or any compatible browser by using

this URL: https://usevision.org/data/trackHub/hub.txt or by clicking the track hubs link at

usevision.org. The database cCRE db supports flexible user queries on extensive annotation of

the cCREs, including epigenetic states and esRP scores across cell types, chromatin

accessibility scores across cell types, membership in JmCs, and evolutionary categories. Code

developed for this study is in the Supplemental Material and at these GitHub repositories:

https://github.com/guanjue/Joint_Human_Mouse_IDEAS_State for the joint human-mouse

IDEAS pipeline and https://github.com/usevision/cre_heritability for the sLDSC analysis.

817   Competing interest statement

818   The authors declare no competing interests.

820   Acknowledgments

This work was supported by grants from the National Institutes of Health:

R24DK106766 to RCH, GAB, MJW, YZ, FY, JT, MS, DB, DH, JRH, BG; R01DK054937 to GAB;

R01GM121613 to YZ and SM; R01GM109453 to QL; R35GM133747 to RCM; F31HG012900 to

DJT; R01HG011139; National Science Foundation DBI CAREER 2045500 to SM, and

intramural funds from the National Human Genome Research Institute. We dedicate this paper

to the memory of JT.

References

Agarwal V, Inoue F, Schubach M, Martin BK, Dash PM, Zhang Z, Sohota A, Noble WS,

Yardimci GG, Kircher M et al. 2023. Massively parallel characterization of transcriptional

regulatory elements in three diverse human cell types. *bioRxiv*

doi:10.1101/2023.03.05.531189.

Ballester B, Medina-Rivera A, Schmidt D, Gonzalez-Porta M, Carlucci M, Chen X, Chessman K,

Faure AJ, Funnell AP, Goncalves A et al. 2014. Multi-species, multi-transcription factor

binding highlights conserved control of tissue-specific biological pathways. *eLife* **3**:

e02626.

Bauer DE, Kamran SC, Lessard S, Xu J, Fujiwara Y, Lin C, Shao Z, Canver MC, Smith EC,

Pinello L et al. 2013. An erythroid enhancer of BCL11A subject to genetic variation

determines fetal hemoglobin level. *Science* **342**: 253-257.

Blobel GA, Weiss MJ. 2009. Nuclear Factors that Regulate Erythropoiesis. In *Disorders of*

Hemoglobin: Genetics, Pathophysiology, and Clinical Management, (ed. MH Steinberg,

et al.), pp. 62-85. Cambridge University Press, Cambridge.

Bourque G. 2009. Transposable elements in gene regulation and in the evolution of vertebrate

genomes. *Curr Opin Genet Dev* **19**: 607-612.

Bruse N, van Heeringen SJ. 2018. GimmeMotifs: an analysis framework for transcription factor

846   motif analysis. *bioRxiv* doi:https://doi.org/10.1101/474403.

847   Buniello A, MacArthur JAL, Cerezo M, Harris LW, Hayhurst J, Malangone C, McMahon A,

848   Morales J, Mountjoy E, Sollis E et al. 2019. The NHGRI-EBI GWAS Catalog of published

849   genome-wide association studies, targeted arrays and summary statistics 2019. *Nucleic*

850   *Acids Res* **47**: D1005-D1012.

851   Carroll SB. 2008. Evo-devo and an expanding evolutionary synthesis: a genetic theory of

852   morphological evolution. *Cell* **134**: 25-36.

853   Cheng L, Li Y, Qi Q, Xu P, Feng R, Palmer L, Chen J, Wu R, Yee T, Zhang J et al. 2021.

854   Single-nucleotide-level mapping of DNA regulatory elements that control fetal

855   hemoglobin expression. *Nat Genet* **53**: 869-880.

856   Cheng Y, Ma Z, Kim BH, Wu W, Cayting P, Boyle AP, Sundaram V, Xing X, Dogan N, Li J et al.

857   2014. Principles of regulatory information conservation between mouse and human.

858   *Nature* **515**: 371-375.

859   Chi AW, Bell JJ, Zlotoff DA, Bhandoola A. 2009. Untangling the T branch of the hematopoiesis

860   tree. *Curr Opin Immunol* **21**: 121-126.

861   Corces MR, Buenrostro JD, Wu B, Greenside PG, Chan SM, Koenig JL, Snyder MP, Pritchard

862   JK, Kundaje A, Greenleaf WJ et al. 2016. Lineage-specific and single-cell chromatin

863   accessibility charts human hematopoiesis and leukemia evolution. *Nat Genet* **48**: 1193-

864   1203.

865   Denas O, Sandstrom R, Cheng Y, Beal K, Herrero J, Hardison RC, Taylor J. 2015. Genome-

866   wide comparative analysis reveals human-mouse regulatory landscape and evolution.

867   *BMC Genomics* **16**: 87.

868   Dong X, Greven MC, Kundaje A, Djebali S, Brown JB, Cheng C, Gingeras TR, Gerstein M,

869   Guigo R, Birney E et al. 2012. Modeling gene expression using chromatin features in

870   various cellular contexts. *Genome Biol* **13**: R53.

Dore LC, Crispino JD. 2011. Transcription factor networks in erythroid cell and megakaryocyte

development. *Blood* **118**: 231-239.

Dynan WS, Tjian R. 1983. The promoter-specific transcription factor Sp1 binds to upstream

sequences in the SV40 early promoter. *Cell* **35**: 79-87.

Ernst J, Kellis M. 2010. Discovery and characterization of chromatin states for

systematic annotation of the human genome. *Nat Biotechnol* **28**: 817-825.

Ernst J, Kellis M. 2012. ChromHMM: automating chromatin-state discovery and

characterization. *Nat Methods* **9**: 215-216.

Ferreira R, Ohneda K, Yamamoto M, Philipsen S. 2005. GATA1 function, a paradigm for

transcription factors in hematopoiesis. *Mol Cell Biol* **25**: 1215-1227.

Finucane HK, Bulik-Sullivan B, Gusev A, Trynka G, Reshef Y, Loh PR, Anttila V, Xu H, Zang C,

Farh K et al. 2015. Partitioning heritability by functional annotation using genome-wide

association summary statistics. *Nat Genet* **47**: 1228-1235.

Frangoul H, Altshuler D, Cappellini MD, Chen YS, Domm J, Eustace BK, Foell J, de la Fuente J,

Grupp S, Handgretinger R et al. 2021. CRISPR-Cas9 Gene Editing for Sickle Cell

Disease and beta-Thalassemia. *The New England journal of medicine* **384**: 252-260.

Fujiwara T, O'Geen H, Keles S, Blahnik K, Linnemann AK, Kang YA, Choi K, Farnham PJ,

Bresnick EH. 2009. Discovering hematopoietic mechanisms through genome-wide

analysis of GATA factor chromatin occupancy. *Mol Cell* **36**: 667-681.

Gasperini M, Hill AJ, McFaline-Figueroa JL, Martin B, Kim S, Zhang MD, Jackson D, Leith A,

Schreiber J, Noble WS et al. 2019. A Genome-wide Framework for Mapping Gene

Regulation via Cellular Genetic Screens. *Cell* **176**: 377-390 e319.

Ge T, Chen CY, Neale BM, Sabuncu MR, Smoller JW. 2017. Phenome-wide heritability analysis

of the UK Biobank. *PLoS Genet* **13**: e1006711.

Graf T, Enver T. 2009. Forcing cells to change lineages. *Nature* **462**: 587-594.

Gumucio DL, Heilstedt-Williamson H, Gray TA, Tarle SA, Shelton DA, Tagle D, Slightom J,

Goodman M, Collins FS. 1992. Phylogenetic footprinting reveals a nuclear protein which

binds to silencer sequences in the human g and e globin genes. *Mol Cell Biol* **12**: 4919-

4929.

Hamamoto K, Fukaya T. 2022. Molecular architecture of enhancer-promoter interaction. *Curr*

Opin Cell Biol **74**: 62-70.

Hardison RC. 2000. Conserved noncoding sequences are reliable guides to regulatory

elements. *Trends in Genetics* **16**: 369-372.

Hardison RC. 2012. Genome-wide epigenetic data facilitate understanding of disease

susceptibility association studies. *J Biol Chem* **287**: 30932-30940.

Hardison RC, Zhang Y, Keller CA, Xiang G, Heuston EF, An L, Lichtenberg J, Giardine BM,

Bodine D, Mahony S et al. 2020. Systematic integration of GATA transcription factors

and epigenomes via IDEAS paints the regulatory landscape of hematopoietic cells.

IUBMB Life **72**: 27-38.

Heintzman ND, Hon GC, Hawkins RD, Kheradpour P, Stark A, Harp LF, Ye Z, Lee LK, Stuart

RK, Ching CW et al. 2009. Histone modifications at human enhancers reflect global cell-

type-specific gene expression. *Nature* **459**: 108-112.

Hindorff LA, Sethupathy P, Junkins HA, Ramos EM, Mehta JP, Collins FS, Manolio TA. 2009.

Potential etiologic and functional implications of genome-wide association loci for human

diseases and traits. *Proc Natl Acad Sci U S A* **106**: 9362-9367.

Hinrichs AS, Karolchik D, Baertsch R, Barber GP, Bejerano G, Clawson H, Diekhans M, Furey

TS, Harte RA, Hsu F et al. 2006. The UCSC Genome Browser Database: update 2006.

Nucleic Acids Res **34**: D590-598.

Hoffman MM, Ernst J, Wilder SP, Kundaje A, Harris RS, Libbrecht M, Giardine B, Ellenbogen

PM, Bilmes JA, Birney E et al. 2013. Integrative annotation of chromatin elements from

ENCODE data. *Nucleic Acids Res* **41**: 827-841.

Jacques PE, Jeyakani J, Bourque G. 2013. The majority of primate-specific regulatory

sequences are derived from transposable elements. *PLoS Genet* **9**: e1003504.

Jansen C, Ramirez RN, El-Ali NC, Gomez-Cabrero D, Tegner J, Merkenschlager M, Conesa A,

Mortazavi A. 2019. Building gene regulatory networks from scATAC-seq and scRNA-seq

using Linked Self Organizing Maps. *PLoS Comput Biol* **15**: e1006555.

Jian J, Konopka J, Liu C. 2013. Insights into the role of progranulin in immunity, infection, and

inflammation. *J Leukoc Biol* **93**: 199-208.

Kaczynski J, Cook T, Urrutia R. 2003. Sp1- and Kruppel-like transcription factors. *Genome Biol*

4: 206.

Kakumanu A, Velasco S, Mazzoni E, Mahony S. 2017. Deconvolving sequence features that

discriminate between overlapping regulatory annotations. *PLoS Comput Biol* **13**:

e1005795.

Karlić R, Chung HR, Lasserre J, Vlahovicek K, Vingron M. 2010. Histone modification levels are

predictive for gene expression. *Proc Natl Acad Sci U S A* **107**: 2926-2931.

King DC, Taylor J, Zhang Y, Cheng Y, Lawson HA, Martin J,

ENCODE_groups_for_Transcriptional_Regulation_and_Multispecies_Sequence_Analysi

s, Chiaromonte F, Miller W, Hardison RC. 2007. Finding cis-regulatory elements using

comparative genomics: some lessons from ENCODE data. *Genome Res* **17**: 775-786.

Kondo M, Wagers AJ, Manz MG, Prohaska SS, Scherer DC, Beilhack GF, Shizuru JA,

Weissman IL. 2003. Biology of hematopoietic stem cells and progenitors: implications for

clinical application. *Annu Rev Immunol* **21**: 759-806.

Kwon SB, Ernst J. 2021. Learning a genome-wide score of human-mouse conservation at the

functional genomics level. *Nature communications* **12**: 2495.

Laurenti E, Göttgens B. 2018. From haematopoietic stem cells to complex differentiation

landscapes. *Nature* **553**: 418-426.

Lee DD, Seung HS. 1999. Learning the parts of objects by non-negative matrix factorization.

948   *Nature* **401**: 788-791.

949   Lee DI, Roy S. 2021. GRiNCH: simultaneous smoothing and detection of topological units of

950   genome organization from sparse chromatin contact count matrices with matrix

951   factorization. *Genome Biol* **22**: 164.

952   Libbrecht MW, Chan RCW, Hoffman MM. 2021. Segmentation and genome annotation

953   algorithms for identifying chromatin state and other genomic patterns. *PLoS Comput Biol*

954   **17**: e1009423.

955   Martens JH, Stunnenberg HG. 2013. BLUEPRINT: mapping human blood cell epigenomes.

956   *Haematologica* **98**: 1487-1489.

957   Maston GA, Evans SK, Green MR. 2006. Transcriptional Regulatory Elements in the Human

958   Genome. *Annu Rev Genomics Hum Genet* **7**: 29-59.

959   Maurano MT, Humbert R, Rynes E, Thurman RE, Haugen E, Wang H, Reynolds AP, Sandstrom

960   R, Qu H, Brody J et al. 2012. Systematic localization of common disease-associated

961   variation in regulatory DNA. *Science* **337**: 1190-1195.

962   Meuleman W, Muratov A, Rynes E, Halow J, Lee K, Bates D, Diegel M, Dunn D, Neri F,

963   Teodosiadis A et al. 2020. Index and biological spectrum of human DNase I

964   hypersensitive sites. *Nature* **584**: 244-251.

965   Noyes MB, Christensen RG, Wakabayashi A, Stormo GD, Brodsky MH, Wolfe SA. 2008.

966   Analysis of homeodomain specificities allows the family-wide prediction of preferred

967   recognition sites. *Cell* **133**: 1277-1289.

968   Payne KJ, Crooks GM. 2002. Human hematopoietic lineage commitment. *Immunol Rev* **187**:

969   48-64.

970   Pennacchio LA, Rubin EM. 2001. Genomic strategies to identify mammalian regulatory

971   sequences. *Nat Rev Genet* **2**: 100-109.

Pimkin M, Kossenkov AV, Mishra T, Morrissey CS, Wu W, Keller CA, Blobel GA, Lee D, Beer
MA, Hardison RC et al. 2014. Divergent functions of hematopoietic transcription factors
in lineage priming and differentiation during erythro-megakaryopoiesis. *Genome Res* **24**:
1932-1944.

Pollard KS, Hubisz MJ, Rosenbloom KR, Siepel A. 2010. Detection of nonneutral substitution
rates on mammalian phylogenies. *Genome Res* **20**: 110-121.

Qi Q, Cheng L, Tang X, He Y, Li Y, Yee T, Shrestha D, Feng R, Xu P, Zhou X et al. 2021.
Dynamic CTCF binding directly mediates interactions among cis-regulatory elements
essential for hematopoiesis. *Blood* **137**: 1327-1339.

Quinlan AR, Hall IM. 2010. BEDTools: a flexible suite of utilities for comparing genomic
features. *Bioinformatics* **26**: 841-842.

Rebollo R, Romanish MT, Mager DL. 2012. Transposable elements: an abundant and natural
source of regulatory sequences for host genes. *Annu Rev Genet* **46**: 21-42.

Ringrose L, Paro R. 2004. Epigenetic regulation of cellular memory by the Polycomb and
Trithorax group proteins. *Annu Rev Genet* **38**: 413-443.

Rothenberg EV, Taghon T. 2005. Molecular genetics of T cell development. *Annu Rev Immunol*
23: 601-649.

Schmidt D, Wilson MD, Ballester B, Schwalie PC, Brown GD, Marshall A, Kutter C, Watt S,
Martinez-Jimenez CP, Mackay S et al. 2010. Five-vertebrate ChIP-seq reveals the
evolutionary dynamics of transcription factor binding. *Science* **328**: 1036-1040.

Schwartz S, Zhang Z, Frazer KA, Smit A, Riemer C, Bouck J, Gibbs R, Hardison R, Miller W.
2000. PipMaker-A web server for aligning two genomic DNA sequences. *Genome Res*
10: 577-586.

Shah M, Funnell APW, Quinlan KGR, Crossley M. 2019. Hit and Run Transcriptional
Repressors Are Difficult to Catch in the Act. *Bioessays* **41**: e1900041.

Shahraki MF, Farahbod M, Libbrecht MW. 2023. Robust chromatin state annotation. *bioRxiv*

doi:https://doi.org/10.1101/2023.07.15.549175.

Spangrude GJ, Heimfeld S, Weissman IL. 1988. Purification and characterization of mouse

hematopoietic stem cells. *Science* **241**: 58-62.

Stein-O'Brien GL, Arora R, Culhane AC, Favorov AV, Garmire LX, Greene CS, Goff LA, Li Y,

Ngom A, Ochs MF et al. 2018. Enter the Matrix: Factorization Uncovers Knowledge from

Omics. *Trends Genet* **34**: 790-805.

Stergachis AB, Neph S, Sandstrom R, Haugen E, Reynolds AP, Zhang M, Byron R, Canfield T,

Stelhing-Sun S, Lee K et al. 2014. Conservation of trans-acting circuitry during

mammalian regulatory evolution. *Nature* **515**: 365-370.

Strahl BD, Allis CD. 2000. The language of covalent histone modifications. *Nature* **403**: 41-45.

Stunnenberg HG, International Human Epigenome C, Hirst M. 2016. The International Human

Epigenome Consortium: A Blueprint for Scientific Collaboration and Discovery. *Cell* **167**:

1145-1149.

Sundaram V, Cheng Y, Ma Z, Li D, Xing X, Edge P, Snyder MP, Wang T. 2014. Widespread

contribution of transposable elements to the innovation of gene regulatory networks.

Genome Res **24**: 1963-1976.

Tenen DG, Hromas R, Licht JD, Zhang DE. 1997. Transcription factors, normal myeloid

development, and leukemia. *Blood* **90**: 489-519.

The ENCODE Project Consortium. 2012. An integrated encyclopedia of DNA elements in the

human genome. *Nature* **489**: 57-74.

The ENCODE Project Consortium, Moore JE, Purcaro MJ, Pratt HE, Epstein CB, Shoresh N,

Adrian J, Kawli T, Davis CA, Dobin A et al. 2020. Expanded encyclopaedias of DNA

elements in the human and mouse genomes. *Nature* **583**: 699-710.

Valverde-Garduno V, Guyot B, Anguita E, Hamlett I, Porcher C, Vyas P. 2004. Differences in

the chromatin structure and cis-element organization of the human and mouse GATA1

loci: implications for cis-element identification. *Blood* **104**: 3106-3116.

van Arensbergen J, FitzPatrick VD, de Haas M, Pagie L, Sluimer J, Bussemaker HJ, van

Steensel B. 2017. Genome-wide mapping of autonomous promoter activity in human

cells. *Nat Biotechnol* **35**: 145-153.

van Pampus EC, Denkers IA, van Geel BJ, Huijgens PC, Zevenbergen A, Ossenkoppele GJ,

Langenhuijsen MM. 1992. Expression of adhesion antigens of human bone marrow

megakaryocytes, circulating megakaryocytes and blood platelets. *Eur J Haematol* **49**:

122-127.

Vierstra J, Lazar J, Sandstrom R, Halow J, Lee K, Bates D, Diegel M, Dunn D, Neri F, Haugen

E et al. 2020. Global reference mapping of human transcription factor footprints. *Nature*

583: 729-736.

Vierstra J, Rynes E, Sandstrom R, Zhang M, Canfield T, Hansen RS, Stehling-Sun S, Sabo PJ,

Byron R, Humbert R et al. 2014. Mouse regulatory DNA landscapes reveal global

principles of cis-regulatory evolution. *Science* **346**: 1007-1012.

Villar D, Flicek P, Odom DT. 2014. Evolution of transcription factor binding in metazoans -

mechanisms and functional implications. *Nat Rev Genet* **15**: 221-233.

Weirauch MT, Yang A, Albu M, Cote AG, Montenegro-Montero A, Drewe P, Najafabadi HS,

Lambert SA, Mann I, Cook K et al. 2014. Determination and inference of eukaryotic

transcription factor sequence specificity. *Cell* **158**: 1431-1443.

Weiss MJ, Yu C, Orkin SH. 1997. Erythroid-cell-specific properties of transcription factor GATA-

1 revealed by phenotypic rescue of a gene-targeted cell line. *Mol Cell Biol* **17**: 1642-

1651.

Xiang G, Giardine BM, Mahony S, Zhang Y, Hardison RC. 2021. S3V2-IDEAS: a package for

normalizing, denoising and integrating epigenomic datasets across different cell types.

Bioinformatics **37**: 3011-3013.

Xiang G, Guo Y, Bumcrot D, Sigova A. 2023. JMnorm: a novel Joint Multi-feature normalization

method for integrative and comparative epigenomics. *bioRxiv*

doi:https://doi.org/10.1101/2023.06.14.545004.

Xiang G, Keller CA, Heuston E, Giardine BM, An L, Wixom AQ, Miller A, Cockburn A, Sauria

MEG, Weaver K et al. 2020. An integrative view of the regulatory and transcriptional

landscapes in mouse hematopoiesis. *Genome Res* **30**: 472-484.

Xu J, Shao Z, Glass K, Bauer DE, Pinello L, Van Handel B, Hou S, Stamatoyannopoulos JA,

Mikkola HK, Yuan GC et al. 2012. Combinatorial assembly of developmental stage-

specific enhancers controls gene expression programs during human erythropoiesis.

Dev Cell **23**: 796-811.

Yang Y, Gu Q, Zhang Y, Sasaki T, Crivello J, O'Neill RJ, Gilbert DM, Ma J. 2018. Continuous-

Trait Probabilistic Model for Comparing Multi-species Functional Genomic Data. *Cell*

Syst **7**: 208-218 e211.

Yue F Cheng Y Breschi A Vierstra J Wu W Ryba T Sandstrom R Ma Z Davis C Pope BD et al.

2014. A comparative encyclopedia of DNA elements in the mouse genome. *Nature* **515**:

355-364.

Zhang Y, An L, Yue F, Hardison RC. 2016. Jointly characterizing epigenetic dynamics across

multiple human cell types. *Nucleic Acids Res* **44**: 6721-6731.

Zhang Y, Hardison RC. 2017. Accurate and reproducible functional maps in 127 human cell

types via 2D genome segmentation. *Nucleic Acids Res* **45**: 9823-9836.

Zhang Y, Liu T, Meyer CA, Eeckhoute J, Johnson DS, Bernstein BE, Nussbaum C, Myers RM,

Brown M, Li W et al. 2008. Model-based analysis of ChIP-Seq (MACS). *Genome Biol* **9**:

R137.

Zhang Y, Mahony S. 2019. Direct prediction of regulatory elements from partial data without

imputation. *PLoS Comput Biol* **15**: e1007399.

1073

Figure Legends

Figure 1. Cell types and data sets used for systematic integration of epigenetic features of blood cells. (A) The tree on the left shows the populations of stem, progenitor, and mature blood cells and cell lines in human. The diagram on the right indicates the epigenetic features and transcriptomes for which genome-wide data sets were generated or collected, with distinctive icons for the major sources of data, specifically the Blueprint project (Martens and Stunnenberg 2013; Stunnenberg et al. 2016), Corces et al. (2016), abbreviated CMB, and St. Jude Children's Research Hospital (SJCRH, Cheng et al. 2021; Qi et al. 2021). (B) Cell types and epigenetic data sets in mouse, diagrammed as for panel A. Sources were described in Xiang et al. (2020) and Supplemental Table S1. Abbreviations for blood cells and lines are: HSC = hematopoietic stem cell, MPP = multipotent progenitor cell, LMPP = lymphoid-myeloid primed progenitor cell, CMP = common myeloid progenitor cell, MEP = megakaryocyte-erythrocyte progenitor cell, K562 = a human cancer cell line with some features of early megakaryocytic and erythroid cells, HUDEP = immortalized human umbilical cord blood-derived erythroid progenitor cell lines expressing fetal globin genes (HUDEP1) or adult globin genes (HUDEP2), CD34_E = human erythroid cells generated by differentiation from CD34+ blood cells, ERY = erythroblast, RBC = mature red blood cell, MK = megakaryocyte, GMP = granulocyte monocyte progenitor cell, EOS = eosinophil, MON = monocyte, MONp = primary monocyte, MONc = classical monocyte, NEU = neutrophil, CLP = common lymphoid progenitor cell, B = B cell, NK = natural killer cell, TCD4 = CD4+ T cell, TCD8 = CD8+ T cell, LSK = Lin-Sca1+Kit+ cells from mouse bone marrow containing hematopoietic stem and progenitor cells, HPC7 = immortalized mouse cell line capable of differentiation in vitro into more mature myeloid cells, G1E = immortalized mouse cell line blocked in erythroid maturation by a knockout of the *Gata1* gene and its subline

ER4 that will further differentiate after restoration of *Gata1* function in an estrogen inducible

manner (Weiss et al. 1997), MEL = murine erythroleukemia cell line that can undergo further

maturation upon induction (designated iMEL), CFUE = colony forming unit erythroid, FL =

designates ERY derived from fetal liver, BM = designates ERY derived from adult bone marrow,

CFUMK = colony forming unit megakaryocyte, iMK = immature megakaryocyte, MK_fl =

megakaryocyte derived from fetal liver.

1103

Figure. 2. Genome segmentation and annotation jointly between human and mouse using

IDEAS. (A) Workflow for joint modeling. (1) Initial epigenetic states from 100 randomly selected

regions separately in human and mouse hematopoietic cell types were identified in IDEAS runs.

(2) States that were reproducible and shared in both species were retained. (3a and 3b) The

profile of epigenetic feature contribution to each of the reproducible states was sequentially

refined by applying IDEAS across the full genomes of human and of mouse, updating the state

model after each IDEAS run. (4) Two heterogeneous states were removed to generate the final

joint epigenetic states in the two species. **(B)** The 25 joint epigenetic states for human and

mouse hematopoietic cell types. The average signal of the epigenetic features for each state

are shown in the heatmap. The corresponding state colors, the state labels based on the

function, and the average proportions of the genome covered by each state across cell types

are listed on the right-side of the heatmap. **(C)** Annotation of epigenetic states in a large

genomic interval containing *SLC4A1* and surrounding genes across human blood cell types.

The genomic interval is 210kb, GRCh38 Chr17:44,192,001-44,402,000, with gene annotations

from GENCODE V38. Binding patterns for selected transcription factors are from the VISION

project ChIP-seq tracks (CTCF and GATA1 in adult erythroblasts, signal tracks from MACS,

track heights 100 and 80, respectively) or from the ENCODE data portal (EP300 in K562 cells,

experiment ENCSR000EGE, signal track is fold change over background, track height is 50).

The epigenetic state assigned to each genomic bin in the different cell types is designated by

the color coding shown in panel (B). The replicates in each cell type examined in Blueprint are

labeled by the id for the donor of biosamples. Genes and regulatory regions active primarily in

erythroid (E), granulocytes (G), and megakaryocytes (MK) are marked by gray rectangles. **(D)**

Annotation of epigenetic states in a large genomic interval containing *Slc4a1* and surrounding

genes across mouse blood cell types. The genomic interval is 198kb, mm10

Chr11:102,290,001-102,488,000, with gene annotations from GENCODE VM23. Binding

patterns for selected transcription factors are from the VISION project ChIP-seq tracks (CTCF in

adult erythroblasts, GATA1 and EP300 from the highly erythroid fetal liver, signal tracks from

MACS, track heights 200, 200, and 150, respectively; the EP300 track was made by re-mapping

reads from ENCODE experiment ENCSR982LJQ). The tracks of epigenetic states and

highlighted regions are indicated as in panel (C).

1134

Figure. 3. Overlaps of VISION cCREs with other catalogs and enrichment for variants

associated with blood cell traits. (A) Venn diagram showing intersections of human VISION

cCREs with a combined superset of elements associated with nuclear structure (CTCF OSs,

loop anchors, and TAD boundaries) and with a combined superset of DNA intervals associated

with *cis*-regulatory elements (CREs), including TSSs, CpG islands, peaks from a massively

parallel promoter and enhancer assay, and enhancers predicted from enhancer RNAs, peaks of

binding by EP300, and histone modifications in erythroblasts (see Supplemental Material,

Supplemental Fig. S9, and Supplemental Table S5). **(B)** The proportions of cCREs and

randomly selected, matched sets of intervals in the overlap categories are compared in the bar

graph. For the random sets, the bar shows the mean, and the dots show the values for each of

ten random sets. **(C)** The UpSet plot provides a higher resolution view of intersections of

VISION cCREs with the four groups of CRE-related elements, specifically enhancer-related

(Enh), transcription start sites (TSS), Survey of Regulatory Elements (SuRE), and CpG islands

(CpG). The enrichment for the cCRE overlaps compared to those in randomly selected,

matched sets of intervals are shown in the boxplots below each overlap subset, with dots for the

enrichment relative to individual random sets. **(D)** Overlaps and enrichments of VISION cCREs

for three sets of structure-related elements, specifically CTCF OSs (CT), loop anchors (LA), and

TAD boundary elements. **(E)** Overlaps of VISION cCREs with two sets of experimentally

determined blood cell cCREs. **(F)** Enrichment of SNPs associated with blood cell traits from UK

Biobank in VISION cCREs. Results of the sLDSC analysis of all cCREs are plotted with

enrichment of the cCRE annotation in heritability of each trait on the x-axis, and the significance

of the enrichment on the y-axis. The analysis covers 292 unique traits with GWAS results from

both males and females and 3 traits with results only from males. The vertical dotted line

indicates an enrichment of 1, and the horizontal dotted line delineates the 5% FDR significance

threshold. Points and labels in red represent traits for which there was significant enrichment of

SNPs associated with the VISION cCREs. Traits with a negative enrichment were assigned an

arbitrary enrichment of 0.1 for plotting and appear as the column of points at the bottom left of

the plot. The shape of the point indicates the sex in which the GWAS analysis was performed

for each trait.

1164

Figure 4. Beta coefficients of states, esRP scores of cCREs, joint human-mouse

metaclusters of cCREs based on esRP scores, and enrichment for TFBS motifs. (A) Beta

coefficients and the difference of beta coefficients of the 25 epigenetic states. The vertical

columns on the right show the beta coefficients along with the ID, color, and labels for the 25

joint epigenetic states. The triangular heatmap shows the difference of the beta coefficients

between two states in the right columns. Each value in the triangle heatmap shows the

difference in beta coefficients between the state on top and the state below based on the order

of states in the right columns. **(B)** An example of calculating esRP score for a cCRE in a cell

type based on the beta coefficients of states. For a cCRE covering more than one 200bp bin,

the esRP equals the weighted sum of beta coefficients of states that covers the cCRE, where

the weights are the region covered by different states. **(C)** The average esRP score of all

cCREs in JmCs across blood cell types shared by human and mouse. The right column shows

the number of human cCREs in each JmC. **(D)** The average enrichment of JmCs in 15

homologous gene clusters. The genes are clustered based on the JmCs' enrichments by *k*-

means. **(E)** Motifs enriched in joint metaclusters. The top heatmap shows the enrichment of

motifs in the cCREs in each JmC in human (H) and mouse (M) as a *Z*-score. The logo for each

motif is given to the right of the heat map, labeled by the family of transcription factors that

recognize that motif. The heatmap below is aligned with the motif enrichment heatmap, showing

the mean esRP score for the cCREs in each JmC for all the common cell types examined

between human and mouse. A summary description of the cell types in which the cCREs in

each JmC are more active is given at the bottom.

1186

Figure. 5. Evolutionary and epigenetic comparisons of cCREs. (A) Workflow to partition

blood cell cCREs in human and mouse into three evolutionary categories. N=nonconserved,

S=conserved in sequence but not inferred function, SF=conserved in both sequence and

inferred function as a cCRE, y=yes, n=no. **(B)** Enrichment of SF-conserved human cCREs for

TSSs. The number of elements in seven sets of function-related DNA intervals that overlap with

the 32,422 SF human cCREs was determined, along with the number that overlap with three

subsets (32,422 each) randomly selected from the full set of 200,342 human cCREs. The ratio

of the number of function-related elements overlapping SF-cCREs to the number overlapping a

randomly chosen subset of all cCREs gave the estimate of enrichment plotted in the graph. The

mean for the three determinations of enrichment is indicated by the horizontal line for each set.

Results are also shown for a similar analysis for the S and N cCREs. **(C)** Distribution of phyloP

scores for three evolutionary categories of cCREs in human and mouse. The maximum phyloP

score for each genomic interval was used to represent the score for each cCRE, using genome

sequence alignments of 100 species with human as the reference (phyloP100) and alignments

of 60 species with mouse as the reference (phyloP60). The distribution of phyloP scores for

each group are displayed as a violin plot. All ten random sets had distributions similar to the one

shown. The asterisk (*) over brackets indicates comparison for which the P values for Welch's *t*-

test is less than 2.2×10^{-16}. **(D)** Proportion of human genomic elements active in a massively

parallel reporter assay (MPRA) that align with mouse or are in a state reflecting dynamic

chromatin. A set of 57,061 genomic elements found to be active in a lentivirus MPRA that tested

a close to comprehensive set of predicted regulatory elements in K562 cells (Agarwal et al.

2023) were assessed for their ability to align with the mouse genome (blue bar) or whether the

IDEAS epigenetic state assigned in K562 cells was not quiescent or was in a set of states

associated with gene activation (magenta bars). The results are plotted as percentages of the

total number of MPRA-active elements.

1212

Figure. 6. Epigenetic comparisons of regulatory landscapes and cCREs. (A and B) DNA

sequence alignments and correlations of epigenetic states in human *GATA1* and mouse *Gata1*

genes and flanking genes. **(A)** Dot-plot view of chained blastZ alignments by PipMaker

(Schwartz et al. 2000) between genomic intervals encompassing and surrounding the human

GATA1 (GRCh38 ChrX:48,760,001-48,836,000; 76kb) and mouse *Gata1* (mm10

ChrX:7,919,401-8,020,800; 101.4kb, reverse complement of reference genome) genes. The

axes are annotated with gene locations (GENCODE), predicted *cis*-regulatory elements

(cCREs), and binding patterns for GATA1 and EP300 in erythroid cells. **(B)** Matrix of Pearson's

correlation values between epigenetic states (quantitative contributions of each epigenetic

feature to the assigned state) across 15 cell types analogous for human and mouse. The

correlation is shown for each 200bp bin in one species with all the bins in the other species,

using a red-blue heat map to indicate the value of the correlation. Axes are annotated with

genes and cCREs in each species. **(C)** Decomposition of the correlation matrix (panel **B**) into

six component parts or factors using nonnegative matrix factorization. **(D-G)** Correlation

matrices for genomic intervals encompassing *GATA1*/*Gata1* and flanking genes, reconstructed

using values from NMF factors. **(D and E)** Correlation matrices using values of NMF factor 3

between human and mouse (panel **D**) or within human and within mouse (panel **E**). The red

rectangles highlight the positive regulatory patterns in the *GATA1*/*Gata1* genes (labeled Px),

which exhibit conservation of both DNA sequence and epigenetic state pattern. The orange

rectangles denote the distal positive regulatory region present only in mouse (labeled D), which

shows conservation of epigenetic state pattern without corresponding sequence conservation.

Beneath the correlation matrices in panel **E** are maps of IDEAS epigenetic states across 15 cell

types, followed by a graph of the score and peak calls for NMF factor 3 and annotation of

cCREs (thin black rectangles) and genes. **(F and G)** Correlation matrices using values of NMF

factor 6 between human and mouse (panel **F**) or within human and within mouse (panel **G**). The

green rectangles highlight the correlation of epigenetic state patterns within the same gene,

both across the two species and within each species individually, while the black rectangles

highlight the high correlation observed between the two genes *GATA1* and *HDAC6*.

A.

B.

C. NMF decomposition

Factors 1-6

Human

Low   High
Score

D. NMF factor 3 correlation

E. NMF factor 3: positive regulation ERY & MK

F. NMF factor 6 correlation

G. NMF factor 6: transcription elongation