Sex-specific age-related changes in glymphatic function assessed by resting-state functional

magnetic resonance imaging

Feng Han[1], Xufu Liu[1], Yifan Yang[1], Xiao Liu[1,2] *

Affiliations:

[1] Department of Biomedical Engineering, The Pennsylvania State University, PA, USA;

[2] Institute for Computational and Data Sciences, The Pennsylvania State University, PA, USA

*Correspondence to: xxl213@psu.edu

Summary

The glymphatic system that clears out brain wastes, such as amyloid-β (Aβ) and tau, through cerebrospinal fluid (CSF) flow may play an important role in aging and dementias. However, a lack of non-invasive tools to assess the glymphatic function in humans hindered the understanding of the glymphatic changes in healthy aging. The global infra-slow (<0.1 Hz) brain activity measured by the global mean resting-state fMRI signal (gBOLD) was recently found to be coupled by large CSF movements. This coupling has been used to measure the glymphatic process and found to correlate with various pathologies of Alzheimer's disease (AD), including Aβ pathology. Using resting-state fMRI data from a large group of 719 healthy aging participants, we examined the sex-specific changes of the gBOLD-CSF coupling, as a measure of glymphatic function, over a wide age range between 36-100 years old. We found that this coupling index remains stable before around age 55 and then starts to decline afterward, particularly in females. Menopause may contribute to the accelerated decline in females.

Keywords

Healthy aging humans; glymphatic function; cerebrospinal fluid (CSF) flow; global brain

activity; gender-effect; sleep quality; menopause.

Introduction

Aging is the leading risk factor for cognitive decline and neurodegenerative disorders that are often associated with excessive accumulation of misfolded proteins, including the amyloid-β and tau, in the brain (Hou et al., 2019; Mawuenyega et al., 2010; Peng et al., 2016). Recent studies suggested the aggregation of toxic proteins could be partly attributed to impaired glymphatic clearance at advancing ages (Benveniste et al., 2019; Boland et al., 2018; Jessen et al., 2015). The glymphatic system, as "glia-lymphatic system", constitutes a pathway for brain waste clearance in the central nervous system (Iliff et al., 2012; Jessen et al., 2015). In this clearance pathway, cerebrospinal fluid (CSF) moves from the periarterial space, facilitated by aquaporin-4 (AQP4) channels in astroglial endfeet, into the interstitial space to wash out interstitial solutes, including Aβ and tau, into the perivenous space surrounding deep-draining veins (Jessen et al., 2015; Tarasoff-Conway et al., 2015). The paravascular CSF recirculation and interstitial solute efflux have been found to decrease in aged mice due to widespread loss of perivascular AQP4 polarization and reduced pulsatility of intracortical arterioles (Kress et al., 2014). In humans, the clearance along the glymphatic pathway and downstream meningeal lymphatic vessels, measured by contrast-agent MRI, decreased and delayed in older patients as compared with younger ones (Zhou et al., 2020). However, the invasive nature of these imaging tools has hindered a large-scale study of glymphatic function in healthy aging subjects. As a result, it remains unclear how the glymphatic function changes in aging, which is vital to understanding the mechanisms of age-related neurodegenerative disorders and cognitive decline.

Global infra-slow (< 0.1 Hz) brain activity measured by resting-state fMRI (rsfMRI) was recently

linked to the glymphatic function (Kiviniemi et al., 2016) and used to quantify its changes in

Alzheimer's disease (AD) (Han et al., 2021b) and Parkinson's disease (PD) patients (Han et al.,

2021a). Increasing evidence suggested the brain exhibits highly structured, brain-wide infra-slow

(< 0.1 Hz) activity during the resting state (Gu et al., 2021; Liu et al., 2021; Raut et al., 2021;

Thompson et al., 2014). This global brain activity is evident in neural signals of distinct scales,

ranging from single neuron recordings to whole-brain fMRI, and closely related to transient

arousal modulations (Gu et al., 2021; Liu et al., 2021). The fMRI measure of this activity, i.e., the

global mean rsfMRI blood-oxygenation-level-dependent (gBOLD) signal, is coupled by large CSF

movements (Fultz et al., 2019) and astroglial calcium waves (Wang et al., 2018), suggesting its

potential link to the glymphatic function. The gBOLD is greatly enhanced during sleep (Fukunaga

et al., 2006; Larson-Prior et al., 2009; Olbrich et al., 2009), in accordance with the sleep-enhanced

nature of the glymphatic function (Xie et al., 2013). In contrast, arterial and respiratory pulsations,

which had been traditionally regarded as the main glymphatic drivers (Iliff et al., 2013; Yamada

et al., 2013), actually lack of this attribute with the decreased amplitude during sleep (Boudreau et

al., 2013; Douglas et al., 1982; Snyder et al., 1964). For all these reasons, the strength of gBOLD-

CSF coupling has been proposed as a surrogate measure of the glymphatic function and found to

correlate with various AD pathologies and cognitive decline in PD (Han et al., 2021b, 2021a).

Recently, the disengagement of gBOLD from the default mode network (DMN) regions were

found to account for early, preferential Aβ accumulation in these higher-order brain areas in the

early stage of Aβ pathology (Han et al., 2022). In these early studies, the gBOLD-CSF coupling

also was found to correlate significantly with age and sex (Han et al., 2021b, 2021a), but the related

findings were limited by narrow age ranges and complicated by the inclusion of patient data.

The fMRI-based glymphatic measurement, together with the widely available rsfMRI data, provides us a unique opportunity to study the change of the glymphatic function over a wider age range and in a larger population of healthy aging subjects. In this study, we used rsfMRI data of 719 healthy aging subjects in the Human Connectome Project Aging (HCP-A) (Harms et al., 2018) to study the age-related changes in glymphatic function in a sex-specific way. We found that the fMRI-based glymphatic measure remained relatively stable within the range of age 36-54 and then started to decrease around age mid-50s. Compared with males, females showed a larger and more abrupt decline of the glymphatic function at this transitioning point. In addition, menopause may lead to an accelerated glymphatic decline in females.

Results

Nonlinear age trajectory of the gBOLD-CSF coupling

We used rsfMRI data of 719 healthy subjects (403 females) from the HCP-A project with age ranging from 36 to 100. Following previous procedures (Fultz et al., 2019; Han et al., 2021b, 2021a), we obtained the whole-brain gBOLD signal from the gray matter regions and the bottom-slice CSF signal around the bottom of the cerebellum to measure CSF movements via MR inflow effects (Fultz et al., 2019; Gao et al., 1996; Gao and Liu, 2012) (**Fig. S1A** and **S1B**). Consistent with the previous studies, the averaged cross-correlation function of the two signals displayed a biphasic pattern with a negative peak ($r = $ -0.33) at the +3.2 sec lag (**Fig. S1C**). The gBOLD-CSF correlation at this +3.2 sec time lag was then computed for individual subjects to quantify their

coupling and thus the glymphatic function. The gBOLD-CSF coupling was then averaged within

10 equal-size groups of subjects at different ages, and the resulting age-related trend displayed a

clear non-linear trajectory: it remains relatively stable between ages 36 to 54 and then begins to

decline at around 55 years old (i.e., yrs) (**Fig. 1A**). We then divided the entire cohort into younger

and older groups according to the age boundary (53.9 yrs) between the fourth and fifth groups,

which had the largest drop among pairs of consecutive groups. The gBOLD-CSF coupling is not

correlated ($r = -3.7 \times 10^{-3}$, $p = 0.95$) with age in the younger group whereas this correlation is

significant ($r = 0.14$, $p = 3.9 \times 10^{-3}$) in the older group (**Fig. 1B**).

We examined potential factors mediating this age-coupling association. Neither the coupling index

nor age was significantly correlated with the total Pittsburgh Sleep Quality Index (PSQI) score,

but they were similarly correlated with a few PSQI items, including sleep medication, trouble

sleeping, and sleep hours. Nevertheless, the age-related changes in the gBOLD-CSF coupling

remain largely unchanged with adjusting for these sleep-related measurements (**Fig. 2**). Likewise,

the age trajectory of the gBOLD-CSF remained similar with controlling for other non-sleep factors,

including the head motion assessed by mean framewise displacement (FD) (**Fig. S2**), the brain

volume (**Fig. S3**), and the CSF volume (mainly from the ventricles) (**Fig. S4**), even though these

factors showed a significant dependence on age (all $p < 3.9 \times 10^{-7}$ for linear regression).

Sex-specific differences in age-related changes

The gBOLD–CSF coupling is different between females and males. The mean gBOLD-CSF

strength is significantly ($p = 7.9 \times 10^{-6}$, two-sample t-test) weaker in females than males (**Fig. 3A**),

consistent with the previous finding in an AD cohort (Han et al., 2021b). The age trajectories of

the coupling index are different for the two groups. The gBOLD–CSF coupling decreases more

evidently and abruptly with aging in the females, particularly around 55 yrs, whereas the males

showed a more gradual and steady age-related changes (**Fig. 3B**).

We further tested whether and how the gBOLD-CSF coupling is affected by menopause, which

often occurs before age 55. This analysis was focused on the age range of 36-55, which includes

both post-menopause and other females. The menopause status had no significant effects on the

gBOLD-CSF coupling strength for two subgroups of subjects of similar ages (both $p > 0.13$; two

sample t-test). However, it showed a marginally significant ($p = 0.061$) interaction with age on the

coupling strength. Consistent with this result, the gBOLD-CSF coupling in the postmenopausal

females appeared to decline earlier in the age mid-40s (**Fig. 3C**), making the drop around age 55

less abrupt and significant ($p = 0.058$). The result suggests the potential effect of the menopause

to accelerate the decline of the fMRI-based glymphatic measure.

Discussion

Here we study the sex-specific age-related change in the glymphatic function in a large healthy

aging population based on the gBOLD-CSF coupling measured by rsfMRI. We showed that the

gBOLD-CSF coupling changes with age in a non-linear way: it remains relatively stable from 36

yrs to 54 yrs and then begins to decrease afterwards. Importantly, the decrease at age mid-50s is

larger and more abrupt in females than in males.

Converging evidence has suggested a link between the glymphatic function and the resting-state

global brain activity, often measured by the gBOLD signal of rsfMRI. The gBOLD, once regarded

as noise, measured a brain-wide, low-frequency (< 0.1 Hz) activity linked to transient arousal

modulations, which also has been observed in monkey electrocorticography (ECoG) (Liu et al.,

2015, 2018) and mice spiking data as highly structured patterns (Liu et al., 2021). The low-

frequency rsfMRI signals were first linked to the glymphatic function due to their potential link to

CSF dynamics and vasomotor waves (Kiviniemi et al., 2016). It was found later that it is the global

component of rsfMRI, i.e., gBOLD, that is coupled to CSF movements in a sleep-dependent way

(Fukunaga et al., 2006; Fultz et al., 2019; Helakari et al., 2022; Larson-Prior et al., 2009; Olbrich

et al., 2009). The sleep dependency makes it more suitable for driving the sleep-dependent

glymphatic clearance (Holth et al., 2019; Xie et al., 2013), as compared with the cardiac and

respiratory pulsations that are actually suppressed during sleep (Baust and Bohnert, 1969;

Boudreau et al., 2013; Douglas et al., 1982; Guazzi and Zanchetti, 1965; Snyder et al., 1964).

Nevertheless, the gBOLD is not independent from these physiological drivers but shows strong

correlations with the low-frequency modulation of cardiac and respiratory functions (Birn et al.,

2006; Chang et al., 2009; Gu et al., 2020; Özbay et al., 2019, 2018; Power et al., 2018). Consistent

with these human findings of strong low-frequency physiological modulation, a recent mice study

demonstrated very strong arterial constrictions/dilations during sleep in the same frequency range

(<0.1 Hz) (Turner et al., 2020). Such low-frequency vessel modulations were coupled by pupil

size changes suggestive of transient arousal modulations (Turner et al., 2022), similar to the global

brain activity measured by gBOLD (Liu et al., 2021, 2018; Pais-Roldán et al., 2020; Turchi et al.,

2018). Animal research also suggested that the gBOLD is coupled by large calcium signals of

astrocytes (Pais-Roldán et al., 2020), and the AQP4 channels on the endfeet of these cells are a

key player of the glymphatic system (Iliff et al., 2012; Jessen et al., 2015). To date, the key

evidence linking the gBOLD to the glymphatic function came from a human study showing that

the gBOLD is coupled to large CSF movements (Fultz et al., 2019). Based on this, the gBOLD-

CSF coupling was used to quantify glymphatic function and found correlated with various

pathologies of AD and cognitive decline in PD (Han et al., 2021b, 2021a). The preferential

reduction of gBOLD signal in the higher-order default mode network was found to account for

early, preferential β-amyloid accumulations in the same regions at the early stage of AD (Han et

al., 2022). All these findings established the foundation for using the gBOLD-CSF coupling to

measure the glymphatic function. But a direct proof of their relationship would need future

experiments capable of recording brain signals across distinct spatial and temporal scales. It is also

worth noting that the debate is ongoing regarding specific components of glymphatic theory, e.g.,

the convective flow in the interstitial space and the involvement of AQP4 channels in the process

(Abbott et al., 2018; Hladky and Barrand, 2022). Nevertheless, a consensus view is reached

regarding the existence of periarterial CSF flow (Mestre et al., 2018) and its role in waste clearance

(van Veluw et al., 2020), which is more related to the gBOLD-CSF coupling in the present study.

Brain aging affects the glymphatic function. The CSF inflow of larger tracers or macromoleculars

decreased up to 85% in old wild-type mice as compared to young counterpart (Da Mesquita et al.,

2018; Nedergaard and Goldman, 2020). The decreased glymphatic flow in aged mice has been

partly attributed to dysregulation of astroglial water transport due to the widespread loss of AQP4

polarization (Kress et al., 2014), the decline of CSF pressure (Fleischman et al., 2012), and the

changes of CSF secretion and protein content (Chen et al., 2009). Arterial wall stiffening and

associated reduction of arterial pulsatility (Zieman et al., 2005) may also account for age-related

glymphatic reduction (Iliff et al., 2013). Human studies of glymphatic function remain sparse (Eide

et al., 2018; Zhou et al., 2020). Previous studies using the gBOLD-CSF coupling found a consistent

association between the age and glymphatic function, but only in patients of relatively old ages

(Han et al., 2021a, 2021b). A retrospective study combined contrast-agent MRI data from various

patient groups to study the glymphatic function and its change with age (Zhou et al., 2020). Despite

different methodologies and patient cohorts from the present study, a similar age trajectory was

observed for the glymphatic function: it remains stable before 50 yrs and then begins to decline

since then (Zhou et al., 2020). The age-related glymphatic changes could be critical for the aging-

related risk for neurodegenerative diseases (Hou et al., 2019). The glymphatic dysfunction may

result in inadequate clearance and thus accumulation of toxic proteins, such as Aβ and tau, and

thereby increase the vulnerability to developing cognitive impairments and neurodegenerative

diseases (Jessen et al., 2015; Tarasoff-Conway et al., 2015). Epidemiologic research suggested the

late-onset AD, the most common AD variant, starts around the mid-60s with the prevalence

doubled every 5 years afterwards (Qiu et al., 2009). But the pathophysiological process, including

the accumulation of aggregated Aβ, could begin more than a decade before the dementia (Jack et

al., 2013; Sperling et al., 2014). Together, these findings suggested a timeline consistent with our

finding that the glymphatic function begins to decrease at age mid-50s.

The female sex is another leading risk factor for developing AD (Mielke et al., 2014). It has been

found a significantly weaker glymphatic function, as measured by gBOLD-CSF coupling strength,

in females than in males (Han et al., 2021b). Here we confirmed the finding with a much larger

dataset from healthy populations. Importantly, we further showed that the glymphatic function had

different age trajectories in the two groups with the females displaying a larger and more abrupt

decline at around 55 yrs. In fact, women indeed show larger and faster cognitive declines than men

with aging (Levine et al., 2021; Nooyens et al., 2022). Together with increasing evidence that links

the glymphatic dysfunction and cognitive impairments (Da Mesquita et al., 2018; Iliff et al., 2014;

Zou et al., 2019), the sex-specific glymphatic change with aging may provide a possible

explanation for the sex differences in age-related cognitive decline. The menopause and associated

hormone loss have been suggested to contribute to cognitive decline in females (Brown and

Gervais, 2020; Hachul et al., 2015). Our result is not inconsistent with this notion by showing a

marginally significant ($p = 0.061$) interaction between age and menopause on the coupling metrics.

Among the females of age 36-55, the post-menopausal group appeared to show a decline of

gBOLD-CSF coupling with age, which is absent in the non-menopause group. A limited sample

size and information related to menopause could partially account for statistical non-significance.

But the finding should warrant future studies looking into the menopause effects on the glymphatic

function with a refined design and/or augmented dataset.

Sleep quality appeared not to be a major contributor to the age-related glymphatic changes. The

gBOLD-CSF coupling and age are similarly correlated with a few sleep measures, including the

frequency of using sleep medication and the sleep troubles in the month prior to the experiments.

However, the age-related changes in the gBOLD-CSF coupling remained largely unchanged after

adjusting for these sleep measures. Nevertheless, the change in sleep architecture might be related

to the sex difference seen in the age-related glymphatic changes. The age and sex are known to

have a strong interaction effect on the composition of sleep stages. Aging in males is associated

with a significant increase of light sleep (stages 1&2) but decrease in slow wave sleep (SWS: sleep

stages 3&4), whereas this age-related change is absent in women (Mander et al., 2017; Redline et

al., 2004). It is known that the glymphatic function increases during sleep and anesthesia featuring

strong slow wave activity (SWA) (Hablitz et al., 2019; Ju et al., 2017; Xie et al., 2013), and one

would thus expect an improved glymphatic clearance with a higher percentage of SWS. The

empirical evidence, however, suggested an opposite by showing higher SWS is associated with

lower CSF Aβ42 level (Varga et al., 2016), which is an early indicator of preclinical AD and often

accompanied by cortical Aβ accumulation (Jack et al., 2013; Palmqvist et al., 2017). The paradox

might be explained by an observation about the gBOLD and its coupling with CSF flow. They

were more specifically related to ultra-slow (0.6-1 Hz) component of SWA (often related to K-

complexes) (Özbay et al., 2019) and phasic changes of SWA power (Fultz et al., 2019; Gu et al.,

2022), which could be stronger during the light sleep than SWS. Indeed, the SWA was found to

decrease in subjects with more cortical Aβ and poorer memory consolidation (Mander et al., 2015;

Winer et al., 2020), as well as AD patients (De Gennaro et al., 2017), but the reduction was specific

to its ultra-slow component (0.6-1 Hz) with the delta-band (1-4 Hz) power showing opposite

changes. Based on all these findings, it is possible that the age-related increase in the percentage

of light sleep in males may help to offset some age-related decline in glymphatic function and thus

lead to its slow deterioration as compared with females. However, the test of this hypothesis would

have to be left for future studies, particularly those with assessment of subjects' sleep architecture.

STAR Methods

Participants and study data

This study included 719 healthy human subjects (36~100 yrs; 403 females) who have participated

in all 4 sessions of rsfMRI scanning in the HCP-A project

(https://www.humanconnectome.org/study/hcp-lifespan-aging). For these subjects, we also used

their T1-weighted structural MRI imaging and demographic data, such as the age, sex, and the

menstrual cycle of females. These "typical aging" subjects were healthy for their age without

identified pathological causes of cognitive declines, such as stroke or clinical dementia

(Bookheimer et al., 2019). All participants provided written informed consent, and investigators

at each HCP-A participating site obtained ethical approval from the corresponding institutional

review board (IRB).

The use of de-identified data from the HCP-A and the sharing of analysis results have been

reviewed and approved by the Pennsylvania State University IRB (IRB#: STUDY00008766) and

also strictly followed the National Institute of Mental Health (NIMH) Data Archive-data use

certification (DUC).

Image acquisition and preprocessing

The rsfMRI data were acquired at 3T MR scanners (Siemens Medical Solutions; Siemens,

Erlangen, Germany) with a matched protocol (Harms et al., 2018) across four acquisition sites

including Washington University St. Louis, University of Minnesota, Massachusetts General

Hospital, and University of California, Los Angeles (researchers in Oxford University dedicating

to the data analysis). For each subject, 4 sessions of rsfMRI (including the anterior to posterior

phase encoding (PE) from Day1, i.e., AP1, as well as PA1, AP2, and PA2) were followed by one

T1-weighted structural MRI session (MPRAGE sequence, echo time (TE)= 1.8/3.6/5.4/7.2 ms

[multi-echo], repetition time (TR) = 2,500 ms, field of view (FOV) = 256×256 mm^2, 320×300

matrix, number of slices = 208, voxel size = $0.8 \times 0.8 \times 0.8$ mm^3, flip angle = 8°) (Harms et al.,

2018). The T1-weighted MRI served to provide the whole brain and CSF volume information and

was used for the anatomical segmentation and registration. For rsfMRI acquisition, 488 fMRI

volumes were collected with a multiband gradient-recalled (GRE) echo-planar image (EPI)

sequence (TR/TE=800/37 ms, flip angle=52°, FOV = 208 mm, 104×90 matrices, 72 oblique axial

slices, 2 mm isotropic voxels, multiband acceleration factor of 8).

We referred to the previous study (Han et al., 2021b) in preprocessing the rsfMRI data using the

FSL (version 5.0.9; https://fsl.fmrib.ox.ac.uk/fsl/fslwiki/FSL) (Smith et al., 2004) and AFNI

(version 16.3.05; https://afni.nimh.nih.gov/) (Cox, 1996). The general fMRI preprocessing

procedures included motion correction, skull stripping, spatial smoothing (full width at half

maximum (FWHM) = 4mm), temporal filtering (bandpass, approximately 0.01 to 0.1 Hz), and the

co-registration of each fMRI volume to corresponding T1-weighted structural MRI and then to the

152-brain Montreal Neurological Institute (MNI-152) space. The motion parameters were not

regressed out to avoid attenuating the gBOLD signal (Gu et al., 2020; Han et al., 2021b). The

preprocessing of structural images was performed using FSL. Processing steps included spatial

normalization and skull stripping.

Extract gBOLD and the CSF inflow signals and compute their coupling

We followed the previous studies (Han et al., 2021b, 2021a) to extract the gBOLD signal and CSF

inflow signal. We defined the mask of the gray matter regions based on the Harvard-Oxford

cortical and subcortical structural atlases (https://neurovault.org/collections/262/). We then

transformed the gray-matter mask from the MNI-152 space back to the original space of each

session to avoid spatial blurring from the registration process (Fultz et al., 2019), and spatially

averaged the Z-normalized gray-matter rsfMRI signals to obtain the gBOLD signal. Following the

previous study (Han et al., 2021b), the CSF inflow signal was extracted from the CSF region at

the bottom edge of fMRI, with a similar voxel number of CSF ROI for all subjects/sessions (see

an example in **Fig. S1A**). We extracted the CSF ROI from the preprocessed fMRI signal at the

original individual space referring to the corresponding CSF region below the bottom of the

cerebellum from the high-resolution T1-weighted MRI (see an exemplary time-series of gBOLD

and bottom CSF fMRI in **Fig. S1B**).

The cross-correlation function was calculated on the extracted gBOLD signal and the CSF inflow

signal for each fMRI session from each individual subject. The cross-correlation function

quantified the Pearson's correlation at different time lags. We first averaged all the cross-

correlation functions from the 4 individual fMRI sessions (AP1, PA1, AP2, and PA2) for each

subject, and further averaged the functions across all subjects (**Fig. S1C**). Referring to the previous

studies (Han et al., 2021b, 2021a), we quantified the gBOLD–CSF coupling with the session-mean

cross-correlation at the lag of +3.2 seconds, where the negative peak of the subject-mean cross-

correlation located, for each subject.

The relationship between the gBOLD–CSF coupling and age

To access the association between the gBOLD–CSF coupling and age, we first divided all the

subjects into 10 sub-groups with different ages (based on the deciles), and further calculated and

compared the mean gBOLD–CSF coupling for each sub-group. Moreover, all the subjects were

separated into the younger (age < 53.9 yrs) and older (age ≥ 53.9 yrs) groups. The linear regression

was used to evaluate the association between the ages and the coupling measures for the subjects

in each sub-group.

Several sensitivity analyses were then performed to test whether the age-coupling association

would be driven by the various sleep quality measures (i.e., PSQI measures), the brain volume, as

well as the CSF volume. We included the total PSQI score, as well as 4 different PSQI items,

which were selected due to their strong dependence ($p < 0.05$ for linear regression or ordinal

regression) with the coupling measures or age, covering the components/aspects of sleep

medication, trouble sleeping, and sleep hours in the sensitivity test. In the test, each of these PSQI

measures was first linked to the coupling measures and age, and then regressed out from the

coupling measures to further examine the age-coupling associations studied in **Fig. 1**. Similar age-

coupling association tests were applied on the whole brain volume and CSF volume, respectively.

The whole brain volume was accessed by the volume number of all brain regions excluding the

CSF area from the T1-weight MRI, where the CSF regions mainly from all the ventricles were

extracted to quantify the CSF volume. To test whether the head motion would drive the age-

coupling association, we adjusted the gBOLD-CSF coupling for the head motion from each

rsfMRI acquisition, which was quantified by the session-mean framewise displacement (FD),

following the previous study (Han et al., 2021b) and replicated the analysis in **Fig. 1**. The FD was

calculated as the sum of the absolute value of all 6 translational and rotational realignment

parameters derived from the preprocessing (Power et al., 2012). We did not use motion-censoring

methods (Power et al., 2014, 2012) to avoid the influence from the cross-correlation of the

concatenated timeseries of gBOLD and CSF fMRI signals.

Sex-specific coupling changes with aging

We further compared the gBOLD–CSF coupling strength between the male and female subjects,

as well as the respective trajectory of coupling changing with aging. First, we compared the

coupling measures between sexes with a two-sample t-test. Second, we divided the males or

females into 7 sub-groups (one group per 10 years; starting from 36~45 yrs, i.e., $36 \leq$ age < 46),

respectively, and compared the coupling measures across these sub-groups.

To examine whether the menopause would affect the tendency of the coupling measure changing

with aging, we separated all the females into the "post-menopause" and "other females" groups

(based on the measure of "whether having no period for 12 months" in the "menstrual cycle" data),

divided the "other females" into two stages of the "36~45 yrs" and "46~55 yrs", and compared the

coupling measures between the two stages. Similarly, we also selected the same age range/sub-

groups for the post-menopausal subjects and contrasted the corresponding coupling measures

between the two sub-groups, as well as compared the coupling across the "post-menopause" and

"other females" subjects for each age-stage. Furthermore, we applied the same grouping metric on

the "post-menopause" subjects as the entire group of females above, i.e., 7 groups with an age

duration of 10 years (from 36~45 yrs), and then replicated the analyses for whole females to

observe the trajectory of the coupling changing with aging.

Statistical analysis

The present study used the two-sample t-test for the group comparison of continuous variables, including the coupling difference of neighboring groups and that between males and females. The linear regression was applied to estimate the association between age and gBOLD–CSF coupling for the younger group and older group, respectively. The ordinal regression was used for responses with natural ordering among categories (i.e., the association between the coupling measure or age and these PSQI measures). The cross-correlation function was used to evaluate the relationship between the gBOLD signal and CSF inflow signal at different time lags. We also tested the interaction effects of age and menopause on the coupling measure. In the study, Pearson's correlation was employed to access the inter-subject associations between different variables. A p-value less than 0.05 was considered statistical significance.

Acknowledgements

The HCP-A dataset reported in this study was supported by grants U01AG052564 and U01AG052564-S1 and by the 14 NIH Institutes and Centers that support the NIH Blueprint for Neuroscience Research, by the McDonnell Center for Systems Neuroscience at Washington University, by the Office of the Provost at Washington University, and by the University of Minnesota Medical School. We gratefully acknowledge the efforts of all the individuals who have contributed to the project.

Data and/or research tools used in the preparation of this manuscript were obtained from the

National Institute of Mental Health (NIMH) Data Archive (NDA). NDA is a collaborative

informatics system created by the National Institutes of Health to provide a national resource to

support and accelerate research in mental health. Dataset identifier(s): 10.15154/1528735. This

manuscript reflects the views of the authors and may not reflect the opinions or views of the NIH

or of the Submitters submitting original data to NDA.

Funding:

This work is supported by funding from the following:

National Institutes of Health grant RF1 MH123247-01 (Xiao Liu);

National Institutes of Health grant R01 NS113889 (Xiao Liu).

Author contributions:

Conceptualization: Feng Han, Xiao Liu

Methodology: Feng Han, Xiao Liu

Investigation: Feng Han, Xufu Liu, Yifan Yang, Xiao Liu

Visualization: Feng Han, Yifan Yang, Xiao Liu

Funding acquisition: Xiao Liu

Project administration: Xiao Liu

Supervision: Xiao Liu

Writing – original draft: Feng Han, Xiao Liu

Writing – review & editing: Feng Han, Xufu Liu, Yifan Yang, Xiao Liu

Competing Interests:

The authors report no financial interests or potential conflicts of interest.

Data and Materials Availability:

The multimodal data, including subject characteristics, structural MRI, and rsfMRI are all publicly

available at the NDA website upon the approval of the data use application

(https://nda.nih.gov/get/access-data.html). All the code used in the present study are available from

the corresponding author upon request.

Fig. 1 The change of gBOLD–CSF coupling with age. (**A**) The gBOLD–CSF coupling remains relatively stable before age mid-50s and then begins to decrease since then. The entire cohort was grouped into 10 subgroups of equal size. Each error bar represents one standard error of the mean. (**B**) The gBOLD-CSF coupling was significantly correlated with age for the group of subjects over 53.9 yrs ($r = 0.14$, $p = 3.9 \times 10^{-3}$; gray dots at the right), but not so ($r = -3.7 \times 10^{-3}$, $p = 0.95$; black dots at the left) for the younger group. Each dot represents one subject.

Fig. 2 The age-related changes in gBOLD-CSF coupling remain similar with controlling for sleep measures. The gBOLD-CSF coupling (left) and age (middle) are similarly correlated with a few sleep-related measurements in PSQI, including the sleep medication frequency (2nd row), the frequency of bathroom-use during sleep-time (3rd row), the frequency of having pain during sleep-time (4th row), and sleep hours (5th row), but not the total PSQI score (1st row). However, the

444   age-related changes in gBOLD-CSF coupling remain similar with adjusting for these PSQI

445   measures (right). Each dot represents one subject, and error bars represent the standard error of the

446   mean.

Fig. 3 The age-coupling association showed distinct patterns between males and females. (A) The gBOLD-CSF coupling is significantly weaker in females (red) than in males (blue) (p = 7.9×10^{-6}, two-sample t-test). **(B)** The age-related changes of the gBOLD-CSF coupling were summarized separately for the females and males and showed different patterns of trajectory: gBOLD–CSF coupling in the females showed a steep decline around 55 yrs (p = 1.1×10^{-3} for two consecutive subgroups around that age), in contrast to the slow and gradual decreases in the males. **(C)** The gBOLD-CSF coupling trends in the "Post-menopause" and "Other females" groups. All females above age 55 are post-menopause. Within the age range of 36-55, the menopause status and age showed a marginally significant (p = 0.061) interaction effect on the gBOLD-CSF coupling. The coupling index started to decrease early at age mid-40s and thus its reduction around age mid-50s is less significant (p = 0.058) as compared with the entire female group.

References

Abbott, N.J., Pizzo, M.E., Preston, J.E., Janigro, D., Thorne, R.G., 2018. The role of brain

barriers in fluid movement in the CNS: is there a 'glymphatic' system? Acta Neuropathol.

135, 387–407. https://doi.org/10.1007/s00401-018-1812-4

Baust, W., Bohnert, B., 1969. The regulation of heart rate during sleep. Exp. Brain Res.

https://doi.org/10.1007/BF00235442

Benveniste, H., Liu, X., Koundal, S., Sanggaard, S., Lee, H., Wardlaw, J., 2019. The Glymphatic

System and Waste Clearance with Brain Aging: A Review. Gerontology 65, 106–119.

https://doi.org/10.1159/000490349

Birn, R.M., Diamond, J.B., Smith, M.A., Bandettini, P.A., 2006. Separating respiratory-

variation-related fluctuations from neuronal-activity-related fluctuations in fMRI.

Neuroimage. https://doi.org/10.1016/j.neuroimage.2006.02.048

Boland, B., Yu, W.H., Corti, O., Mollereau, B., Henriques, A., Bezard, E., Pastores, G.M.,

Rubinsztein, D.C., Nixon, R.A., Duchen, M.R., Mallucci, G.R., Kroemer, G., Levine, B.,

Eskelinen, E.-L., Mochel, F., Spedding, M., Louis, C., Martin, O.R., Millan, M.J., 2018.

Promoting the clearance of neurotoxic proteins in neurodegenerative disorders of ageing.

Nat. Rev. Drug Discov. 17, 660–688. https://doi.org/10.1038/nrd.2018.109

Bookheimer, S.Y., Salat, D.H., Terpstra, M., Ances, B.M., Barch, D.M., Buckner, R.L., Burgess,

G.C., Curtiss, S.W., Diaz-Santos, M., Elam, J.S., Fischl, B., Greve, D.N., Hagy, H.A.,

Harms, M.P., Hatch, O.M., Hedden, T., Hodge, C., Japardi, K.C., Kuhn, T.P., Ly, T.K.,

Smith, S.M., Somerville, L.H., Uğurbil, K., van der Kouwe, A., Van Essen, D., Woods,

R.P., Yacoub, E., 2019. The Lifespan Human Connectome Project in Aging: An overview.

Neuroimage 185, 335–348. https://doi.org/10.1016/j.neuroimage.2018.10.009

Boudreau, P., Yeh, W.H., Dumont, G.A., Boivin, D.B., 2013. Circadian variation of heart rate

variability across sleep stages. Sleep. https://doi.org/10.5665/sleep.3230

Brown, A.M.C., Gervais, N.J., 2020. Role of Ovarian Hormones in the Modulation of Sleep in

Females Across the Adult Lifespan. Endocrinology 161.

https://doi.org/10.1210/endocr/bqaa128

Chang, C., Cunningham, J.P., Glover, G.H., 2009. Influence of heart rate on the BOLD signal:

The cardiac response function. Neuroimage.

https://doi.org/10.1016/j.neuroimage.2008.09.029

Chen, R.L., Kassem, N.A., Redzic, Z.B., Chen, C.P.C., Segal, M.B., Preston, J.E., 2009. Age-

related changes in choroid plexus and blood–cerebrospinal fluid barrier function in the

sheep. Exp. Gerontol. 44, 289–296.

https://doi.org/https://doi.org/10.1016/j.exger.2008.12.004

Cox, R.W., 1996. AFNI: Software for analysis and visualization of functional magnetic

resonance neuroimages. Comput. Biomed. Res. https://doi.org/10.1006/cbmr.1996.0014

Da Mesquita, S., Louveau, A., Vaccari, A., Smirnov, I., Cornelison, R.C., Kingsmore, K.M.,

Contarino, C., Onengut-Gumuscu, S., Farber, E., Raper, D., Viar, K.E., Powell, R.D.,

Baker, W., Dabhi, N., Bai, R., Cao, R., Hu, S., Rich, S.S., Munson, J.M., Lopes, M.B.,

Overall, C.C., Acton, S.T., Kipnis, J., 2018. Functional aspects of meningeal lymphatics in

ageing and Alzheimer's disease. Nature 560, 185–191. https://doi.org/10.1038/s41586-018-

0368-8

De Gennaro, L., Gorgoni, M., Reda, F., Lauri, G., Truglia, I., Cordone, S., Scarpelli, S.,

Mangiaruga, A., D'atri, A., Lacidogna, G., Ferrara, M., Marra, C., Rossini, P.M., 2017. The

Fall of Sleep K-Complex in Alzheimer Disease. Sci. Rep. 7, 39688.

https://doi.org/10.1038/srep39688

Douglas, N.J., White, D.P., Pickett, C.K., Weil, J. V., Zwillich, C.W., 1982. Respiration during

sleep in normal man. Thorax. https://doi.org/10.1136/thx.37.11.840

Eide, P.K., Vatnehol, S.A.S., Emblem, K.E., Ringstad, G., 2018. Magnetic resonance imaging

provides evidence of glymphatic drainage from human brain to cervical lymph nodes. Sci.

Rep. 8, 7194. https://doi.org/10.1038/s41598-018-25666-4

Fleischman, D., Berdahl, J.P., Zaydlarova, J., Stinnett, S., Fautsch, M.P., Allingham, R.R., 2012.

Cerebrospinal Fluid Pressure Decreases with Older Age. PLoS One 7, 1–9.

https://doi.org/10.1371/journal.pone.0052664

Fukunaga, M., Horovitz, S.G., van Gelderen, P., de Zwart, J.A., Jansma, J.M., Ikonomidou,

V.N., Chu, R., Deckers, R.H.R., Leopold, D.A., Duyn, J.H., 2006. Large-amplitude,

spatially correlated fluctuations in BOLD fMRI signals during extended rest and early sleep

stages. Magn. Reson. Imaging 24, 979–992. https://doi.org/10.1016/j.mri.2006.04.018

Fultz, N.E., Bonmassar, G., Setsompop, K., Stickgold, R.A., Rosen, B.R., Polimeni, J.R., Lewis,

L.D., 2019. Coupled electrophysiological, hemodynamic, and cerebrospinal fluid

oscillations in human sleep. Science (80-.). https://doi.org/10.1126/science.aax5440

Gao, J.H., Liu, H.L., 2012. Inflow effects on functional MRI. Neuroimage.

https://doi.org/10.1016/j.neuroimage.2011.09.088

Gao, J.H., Miller, I., Lai, S., Xiong, J., Fox, P.T., 1996. Quantitative assessment of blood inflow

effects in functional MRI signals. Magn. Reson. Med.

https://doi.org/10.1002/mrm.1910360219

Gu, Y., Han, F., Sainburg, L.E., Liu, X., 2020. Transient Arousal Modulations Contribute to

Resting-State Functional Connectivity Changes Associated with Head Motion Parameters.

Cereb. Cortex. https://doi.org/10.1093/cercor/bhaa096

Gu, Y., Han, F., Sainburg, L.E., Schade, M.M., Buxton, O.M., Duyn, J.H., Liu, X., 2022. An

orderly sequence of autonomic and neural events at transient arousal changes. Neuroimage.

https://doi.org/10.1016/j.neuroimage.2022.119720

Gu, Y., Sainburg, L.E., Kuang, S., Han, F., Williams, J.W., Liu, Y., Zhang, N., Zhang, X.,

Leopold, D.A., Liu, X., 2021. Brain Activity Fluctuations Propagate as Waves Traversing

the Cortical Hierarchy. Cereb. Cortex 31, 3986–4005.

https://doi.org/10.1093/cercor/bhab064

Guazzi, M., Zanchetti, A., 1965. Carotid sinus and aortic reflexes in the regulation of circulation

during sleep. Science (80-.). https://doi.org/10.1126/science.148.3668.397

Hablitz, L.M., Vinitsky, H.S., Sun, Q., Stæger, F.F., Sigurdsson, B., Mortensen, K.N., Lilius,

T.O., Nedergaard, M., 2019. Increased glymphatic influx is correlated with high EEG delta

power and low heart rate in mice under anesthesia. Sci. Adv. 5, eaav5447.

https://doi.org/10.1126/sciadv.aav5447

Hachul, H., Frange, C., Bezerra, A.G., Hirotsu, C., Pires, G.N., Andersen, M.L., Bittencourt, L.,

Tufik, S., 2015. The effect of menopause on objective sleep parameters: Data from an

epidemiologic study in São Paulo, Brazil. Maturitas 80, 170–178.

https://doi.org/https://doi.org/10.1016/j.maturitas.2014.11.002

Han, F., Brown, G.L., Zhu, Y., Belkin-Rosen, A.E., Lewis, M.M., Du, G., Gu, Y., Eslinger, P.J.,

Mailman, R.B., Huang, X., Liu, X., 2021a. Decoupling of Global Brain Activity and

Cerebrospinal Fluid Flow in Parkinson's Disease Cognitive Decline. Mov. Disord. 36,

2066–2076. https://doi.org/10.1002/mds.28643

Han, F., Chen, J., Belkin-Rosen, A., Gu, Y., Luo, L., Buxton, O.M., Liu, X., 2021b. Reduced

coupling between cerebrospinal fluid flow and global brain activity is linked to Alzheimer

disease–related pathology. PLoS Biol. 19, 1–25.

https://doi.org/10.1371/journal.pbio.3001233

Han, F., Liu, Xufu, Mailman, R.B., Huang, X., Liu, Xiao, 2022. Early β-amyloid accumulation

and hypoconnectivity in the default mode network are related to its disengagement from

global brain activity. bioRxiv. https://doi.org/10.1101/2022.07.24.501309

Harms, M.P., Somerville, L.H., Ances, B.M., Andersson, J., Barch, D.M., Bastiani, M.,

Bookheimer, S.Y., Brown, T.B., Buckner, R.L., Burgess, G.C., Coalson, T.S., Chappell,

M.A., Dapretto, M., Douaud, G., Fischl, B., Glasser, M.F., Greve, D.N., Hodge, C.,

Jamison, K.W., Jbabdi, S., Kandala, S., Li, X., Mair, R.W., Mangia, S., Marcus, D.,

Mascali, D., Moeller, S., Nichols, T.E., Robinson, E.C., Salat, D.H., Smith, S.M.,

Sotiropoulos, S.N., Terpstra, M., Thomas, K.M., Tisdall, M.D., Ugurbil, K., van der

Kouwe, A., Woods, R.P., Zöllei, L., Van Essen, D.C., Yacoub, E., 2018. Extending the

Human Connectome Project across ages: Imaging protocols for the Lifespan Development

and Aging projects. Neuroimage 183, 972–984.

https://doi.org/10.1016/j.neuroimage.2018.09.060

Helakari, H., Korhonen, V., Holst, S.C., Piispala, J., Kallio, M., Väyrynen, T., Huotari, N.,

Raitamaa, L., Tuunanen, J., Kananen, J., Järvelä, M., Tuovinen, T., Raatikainen, V.,

Borchardt, V., Kinnunen, H., Nedergaard, M., Kiviniemi, V., 2022. Human NREM Sleep

Promotes Brain-Wide Vasomotor and Respiratory Pulsations. J. Neurosci. JN-RM-0934-21.

https://doi.org/10.1523/jneurosci.0934-21.2022

Hladky, S.B., Barrand, M.A., 2022. The glymphatic hypothesis: the theory and the evidence,

Fluids and Barriers of the CNS. BioMed Central. https://doi.org/10.1186/s12987-021-

577   00282-z

578 Holth, J.K., Fritschi, S.K., Wang, C., Pedersen, N.P., Cirrito, J.R., Mahan, T.E., Finn, M.B.,

579   Manis, M., Geerling, J.C., Fuller, P.M., Lucey, B.P., Holtzman, D.M., 2019. The sleep-

580   wake cycle regulates brain interstitial fluid tau in mice and CSF tau in humans. Science

581   (80-.). https://doi.org/10.1126/science.aav2546

582 Hou, Y., Dan, X., Babbar, M., Wei, Y., Hasselbalch, S.G., Croteau, D.L., Bohr, V.A., 2019.

583   Ageing as a risk factor for neurodegenerative disease. Nat. Rev. Neurol. 15, 565–581.

584   https://doi.org/10.1038/s41582-019-0244-7

585 Iliff, J.J., Chen, M.J., Plog, B.A., Zeppenfeld, D.M., Soltero, M., Yang, L., Singh, I., Deane, R.,

586   Nedergaard, M., 2014. Impairment of glymphatic pathway function promotes tau pathology

587   after traumatic brain injury. J. Neurosci. https://doi.org/10.1523/JNEUROSCI.3020-14.2014

588 Iliff, J.J., Wang, M., Liao, Y., Plogg, B.A., Peng, W., Gundersen, G.A., Benveniste, H., Vates,

589   G.E., Deane, R., Goldman, S.A., Nagelhus, E.A., Nedergaard, M., 2012. A paravascular

590   pathway facilitates CSF flow through the brain parenchyma and the clearance of interstitial

591   solutes, including amyloid β. Sci. Transl. Med.

592   https://doi.org/10.1126/scitranslmed.3003748

593 Iliff, J.J., Wang, M., Zeppenfeld, D.M., Venkataraman, A., Plog, B.A., Liao, Y., Deane, R.,

594   Nedergaard, M., 2013. Cerebral arterial pulsation drives paravascular CSF-Interstitial fluid

595   exchange in the murine brain. J. Neurosci. https://doi.org/10.1523/JNEUROSCI.1592-

596   13.2013

597 Jack, C.R., Knopman, D.S., Jagust, W.J., Petersen, R.C., Weiner, M.W., Aisen, P.S., Shaw,

598   L.M., Vemuri, P., Wiste, H.J., Weigand, S.D., Lesnick, T.G., Pankratz, V.S., Donohue,

599   M.C., Trojanowski, J.Q., 2013. Tracking pathophysiological processes in Alzheimer's

disease: An updated hypothetical model of dynamic biomarkers. Lancet Neurol. 12, 207–

216. https://doi.org/10.1016/S1474-4422(12)70291-0

Jessen, N.A., Munk, A.S.F., Lundgaard, I., Nedergaard, M., 2015. The Glymphatic System: A

Beginner's Guide. Neurochem. Res. https://doi.org/10.1007/s11064-015-1581-6

Ju, Y.-E.S., Ooms, S.J., Sutphen, C., Macauley, S.L., Zangrilli, M.A., Jerome, G., Fagan, A.M.,

Mignot, E., Zempel, J.M., Claassen, J.A.H.R., Holtzman, D.M., 2017. Slow wave sleep

disruption increases cerebrospinal fluid amyloid-β levels. Brain 140, 2104–2111.

https://doi.org/10.1093/brain/awx148

Kiviniemi, V., Wang, X., Korhonen, V., Keinänen, T., Tuovinen, T., Autio, J., Levan, P.,

Keilholz, S., Zang, Y.F., Hennig, J., Nedergaard, M., 2016. Ultra-fast magnetic resonance

encephalography of physiological brain activity-Glymphatic pulsation mechanisms? J.

Cereb. Blood Flow Metab. https://doi.org/10.1177/0271678X15622047

Kress, B.T., Iliff, J.J., Xia, M., Wang, M., Wei Bs, H.S., Zeppenfeld, D., Xie, L., Hongyi Kang,

B.S., Xu, Q., Liew, J.A., Plog, B.A., Ding, F., PhD, R.D., Nedergaard, M., 2014.

Impairment of paravascular clearance pathways in the aging brain. Ann. Neurol. 76, 845–

861. https://doi.org/10.1002/ana.24271

Larson-Prior, L.J., Zempel, J.M., Nolan, T.S., Prior, F.W., Snyder, A., Raichle, M.E., 2009.

Cortical network functional connectivity in the descent to sleep. Proc. Natl. Acad. Sci. U. S.

A. https://doi.org/10.1073/pnas.0900924106

Levine, D.A., Gross, A.L., Briceño, E.M., Tilton, N., Giordani, B.J., Sussman, J.B., Hayward,

R.A., Burke, J.F., Hingtgen, S., Elkind, M.S.V., Manly, J.J., Gottesman, R.F., Gaskin, D.J.,

Sidney, S., Sacco, R.L., Tom, S.E., Wright, C.B., Yaffe, K., Galecki, A.T., 2021. Sex

Differences in Cognitive Decline among US Adults. JAMA Netw. Open 4, 1–13.

https://doi.org/10.1001/jamanetworkopen.2021.0169

Liu, X., De Zwart, J.A., Schölvinck, M.L., Chang, C., Ye, F.Q., Leopold, D.A., Duyn, J.H.,

2018. Subcortical evidence for a contribution of arousal to fMRI studies of brain activity.

Nat. Commun. https://doi.org/10.1038/s41467-017-02815-3

Liu, X., Leopold, D.A., Yang, Y., 2021. Single-neuron firing cascades underlie global

spontaneous brain events. Proc. Natl. Acad. Sci. U. S. A. 118, 1–10.

https://doi.org/10.1073/pnas.2105395118

Liu, X., Yanagawa, T., Leopold, D.A., Chang, C., Ishida, H., Fujii, N., Duyn, J.H., 2015.

Arousal transitions in sleep, wakefulness, and anesthesia are characterized by an orderly

sequence of cortical events. Neuroimage. https://doi.org/10.1016/j.neuroimage.2015.04.003

Mander, B.A., Marks, S.M., Vogel, J.W., Rao, V., Lu, B., Saletin, J.M., Ancoli-Israel, S., Jagust,

W.J., Walker, M.P., 2015. β-amyloid disrupts human NREM slow waves and related

hippocampus-dependent memory consolidation. Nat. Neurosci. 18, 1051–1057.

https://doi.org/10.1038/nn.4035

Mander, B.A., Winer, J.R., Walker, M.P., 2017. Sleep and Human Aging. Neuron 94, 19–36.

https://doi.org/https://doi.org/10.1016/j.neuron.2017.02.004

Mawuenyega, K.G., Sigurdson, W., Ovod, V., Munsell, L., Kasten, T., Morris, J.C., Yarasheski,

K.E., Bateman, R.J., 2010. Decreased clearance of CNS β-amyloid in Alzheimer's disease.

Science (80-.). 330, 1774. https://doi.org/10.1126/science.1197623

Mestre, H., Tithof, J., Du, T., Song, W., Peng, W., Sweeney, A.M., Olveda, G., Thomas, J.H.,

Nedergaard, M., Kelley, D.H., 2018. Flow of cerebrospinal fluid is driven by arterial

pulsations and is reduced in hypertension. Nat. Commun. https://doi.org/10.1038/s41467-

018-07318-3

Mielke, M.M., Vemuri, P., Rocca, W.A., 2014. Clinical epidemiology of Alzheimer's disease:

Assessing sex and gender differences. Clin. Epidemiol.

https://doi.org/10.2147/CLEP.S37929

Nedergaard, M., Goldman, S.A., 2020. Glymphatic failure as a final common pathway to

dementia. Science (80-.). 370, 50–56. https://doi.org/10.1126/science.abb8739

Nooyens, A.C.J., Wijnhoven, H.A.H., Schaap, L.S., Sialino, L.D., Kok, A.A.L., Visser, M.,

Verschuren, W.M.M., Picavet, H.S.J., van Oostrom, S.H., 2022. Sex Differences in

Cognitive Functioning with Aging in the Netherlands. Gerontology.

https://doi.org/10.1159/000520318

Olbrich, S., Mulert, C., Karch, S., Trenner, M., Leicht, G., Pogarell, O., Hegerl, U., 2009. EEG-

vigilance and BOLD effect during simultaneous EEG/fMRI measurement. Neuroimage.

https://doi.org/10.1016/j.neuroimage.2008.11.014

Özbay, P.S., Chang, C., Picchioni, D., Mandelkow, H., Chappel-Farley, M.G., van Gelderen, P.,

de Zwart, J.A., Duyn, J., 2019. Sympathetic activity contributes to the fMRI signal.

Commun. Biol. https://doi.org/10.1038/s42003-019-0659-0

Özbay, P.S., Chang, C., Picchioni, D., Mandelkow, H., Moehlman, T.M., Chappel-Farley, M.G.,

van Gelderen, P., de Zwart, J.A., Duyn, J.H., 2018. Contribution of systemic vascular

effects to fMRI activity in white matter. Neuroimage.

https://doi.org/10.1016/j.neuroimage.2018.04.045

Pais-Roldán, P., Takahashi, K., Sobczak, F., Chen, Y., Zhao, X., Zeng, H., Jiang, Y., Yu, X.,

2020. Indexing brain state-dependent pupil dynamics with simultaneous fMRI and optical

fiber calcium recording. Proc. Natl. Acad. Sci. U. S. A. 117, 6875–6882.

https://doi.org/10.1073/pnas.1909937117

Palmqvist, S., Schöll, M., Strandberg, O., Mattsson, N., Stomrud, E., Zetterberg, H., Blennow,

K., Landau, S., Jagust, W., Hansson, O., 2017. Earliest accumulation of β-amyloid occurs

within the default-mode network and concurrently affects brain connectivity. Nat. Commun.

https://doi.org/10.1038/s41467-017-01150-x

Peng, W., Achariyar, T.M., Li, B., Liao, Y., Mestre, H., Hitomi, E., Regan, S., Kasper, T., Peng,

S., Ding, F., Benveniste, H., Nedergaard, M., Deane, R., 2016. Suppression of glymphatic

fluid transport in a mouse model of Alzheimer's disease. Neurobiol. Dis. 93, 215–225.

https://doi.org/https://doi.org/10.1016/j.nbd.2016.05.015

Power, J.D., Barnes, K.A., Snyder, A.Z., Schlaggar, B.L., Petersen, S.E., 2012. Spurious but

systematic correlations in functional connectivity MRI networks arise from subject motion.

Neuroimage. https://doi.org/10.1016/j.neuroimage.2011.10.018

Power, J.D., Mitra, A., Laumann, T.O., Snyder, A.Z., Schlaggar, B.L., Petersen, S.E., 2014.

Methods to detect, characterize, and remove motion artifact in resting state fMRI.

Neuroimage 84, 320–341. https://doi.org/10.1016/j.neuroimage.2013.08.048

Power, J.D., Plitt, M., Gotts, S.J., Kundu, P., Voon, V., Bandettini, P.A., Martin, A., 2018.

Ridding fMRI data of motion-related influences: Removal of signals with distinct spatial

and physical bases in multiecho data. Proc. Natl. Acad. Sci. U. S. A.

https://doi.org/10.1073/pnas.1720985115

Qiu, C., Kivipelto, M., von Strauss, E., 2009. Epidemiology of Alzheimer's disease: occurrence,

determinants, and strategies toward intervention. Dialogues Clin. Neurosci. 11, 111–128.

https://doi.org/10.31887/DCNS.2009.11.2/cqiu

Raut, R. V., Snyder, A.Z., Mitra, A., Yellin, D., Fujii, N., Malach, R., Raichle, M.E., 2021.

Global waves synchronize the brain's functional systems with fluctuating arousal. Sci. Adv.

7, 1–16. https://doi.org/10.1126/sciadv.abf2709

Redline, S., Kirchner, H.L., Quan, S.F., Gottlieb, D.J., Kapur, V., Newman, A., 2004. The

Effects of Age, Sex, Ethnicity, and Sleep-Disordered Breathing on Sleep Architecture.

Arch. Intern. Med. 164, 406–418. https://doi.org/10.1001/archinte.164.4.406

Smith, S.M., Jenkinson, M., Woolrich, M.W., Beckmann, C.F., Behrens, T.E.J., Johansen-Berg,

H., Bannister, P.R., De Luca, M., Drobnjak, I., Flitney, D.E., Niazy, R.K., Saunders, J.,

Vickers, J., Zhang, Y., De Stefano, N., Brady, J.M., Matthews, P.M., 2004. Advances in

functional and structural MR image analysis and implementation as FSL. Neuroimage 23,

S208–S219. https://doi.org/https://doi.org/10.1016/j.neuroimage.2004.07.051

Snyder, F., Hobson, J.A., Morrison, D.F., Goldfrank, F., 1964. Changes in respiration, heart rate,

and systolic blood pressure in human sleep. J. Appl. Physiol.

https://doi.org/10.1152/jappl.1964.19.3.417

Sperling, R., Mormino, E., Johnson, K., 2014. The evolution of preclinical Alzheimer's disease:

Implications for prevention trials. Neuron. https://doi.org/10.1016/j.neuron.2014.10.038

Tarasoff-Conway, J.M., Carare, R.O., Osorio, R.S., Glodzik, L., Butler, T., Fieremans, E., Axel,

L., Rusinek, H., Nicholson, C., Zlokovic, B. V., Frangione, B., Blennow, K., Ménard, J.,

Zetterberg, H., Wisniewski, T., De Leon, M.J., 2015. Clearance systems in the brain -

Implications for Alzheimer disease. Nat. Rev. Neurol.

https://doi.org/10.1038/nrneurol.2015.119

Thompson, G.J., Pan, W.-J., Magnuson, M.E., Jaeger, D., Keilholz, S.D., 2014. Quasi-periodic

patterns (QPP): Large-scale dynamics in resting state fMRI that correlate with local

infraslow electrical activity. Neuroimage 84, 1018–1031.

https://doi.org/https://doi.org/10.1016/j.neuroimage.2013.09.029

Turchi, J., Chang, C., Ye, F.Q., Russ, B.E., Yu, D.K., Cortes, C.R., Monosov, I.E., Duyn, J.H.,

Leopold, D.A., 2018. The Basal Forebrain Regulates Global Resting-State fMRI

Fluctuations. Neuron. https://doi.org/10.1016/j.neuron.2018.01.032

Turner, K.L., Gheres, K.W., Drew, P.J., 2022. Relating Pupil Diameter and Blinking to Cortical

Activity and Hemodynamics across Arousal States. Journal of Neuroscience.

https://doi.org/10.1523/JNEUROSCI.1244-22.2022

Turner, K.L., Gheres, K.W., Proctor, E.A., Drew, P.J., 2020. Neurovascular coupling and

bilateral connectivity during NREM and REM sleep. Elife 9, e62071.

https://doi.org/10.7554/eLife.62071

van Veluw, S.J., Hou, S.S., Calvo-Rodriguez, M., Arbel-Ornath, M., Snyder, A.C., Frosch, M.P.,

Greenberg, S.M., Bacskai, B.J., 2020. Vasomotion as a Driving Force for Paravascular

Clearance in the Awake Mouse Brain. Neuron.

https://doi.org/10.1016/j.neuron.2019.10.033

Varga, A.W., Wohlleber, M.E., Giménez, S., Romero, S., Alonso, J.F., Ducca, E.L., Kam, K.,

Lewis, C., Tanzi, E.B., Tweardy, S., Kishi, A., Parekh, A., Fischer, E., Gumb, T., Alcolea,

D., Fortea, J., Lleó, A., Blennow, K., Zetterberg, H., Mosconi, L., Glodzik, L., Pirraglia, E.,

Burschtin, O.E., de Leon, M.J., Rapoport, D.M., Lu, S., Ayappa, I., Osorio, R.S., 2016.

Reduced Slow-Wave Sleep Is Associated with High Cerebrospinal Fluid Aβ42 Levels in

Cognitively Normal Elderly. Sleep 39, 2041–2048. https://doi.org/10.5665/sleep.6240

Wang, M., He, Y., Sejnowski, T.J., Yu, X., 2018. Brain-state dependent astrocytic Ca2+ signals

are coupled to both positive and negative BOLD-fMRI signals. Proc. Natl. Acad. Sci. U. S.

A. 115, E1647–E1656. https://doi.org/10.1073/pnas.1711692115

Winer, J.R., Mander, B.A., Kumar, S., Reed, M., Baker, S.L., Jagust, W.J., Walker, M.P., 2020.

Sleep Disturbance Forecasts β-Amyloid Accumulation across Subsequent Years. Curr. Biol.

30, 4291-4298.e3. https://doi.org/10.1016/j.cub.2020.08.017

Xie, L., Kang, H., Xu, Q., Chen, M.J., Liao, Y., Thiyagarajan, M., O'Donnell, J., Christensen,

D.J., Nicholson, C., Iliff, J.J., Takano, T., Deane, R., Nedergaard, M., 2013. Sleep drives

metabolite clearance from the adult brain. Science (80-.).

https://doi.org/10.1126/science.1241224

Yamada, S., Miyazaki, M., Yamashita, Y., Ouyang, C., Yui, M., Nakahashi, M., Shimizu, S.,

Aoki, I., Morohoshi, Y., McComb, J.G., 2013. Influence of respiration on cerebrospinal

fluid movement using magnetic resonance spin labeling. Fluids Barriers CNS.

https://doi.org/10.1186/2045-8118-10-36

Zhou, Y., Cai, J., Zhang, W., Gong, X., Yan, S., Zhang, K., Luo, Z., Sun, J., Jiang, Q., Lou, M.,

2020. Impairment of the Glymphatic Pathway and Putative Meningeal Lymphatic Vessels

in the Aging Human. Ann. Neurol. 87, 357–369.

https://doi.org/https://doi.org/10.1002/ana.25670

Zieman, S.J., Melenovsky, V., Kass, D.A., 2005. Mechanisms, Pathophysiology, and Therapy of

Arterial Stiffness. Arterioscler. Thromb. Vasc. Biol. 25, 932–943.

https://doi.org/10.1161/01.ATV.0000160548.78317.29

Zou, W., Pu, T., Feng, W., Lu, M., Zheng, Y., Du, R., Xiao, M., Hu, G., 2019. Blocking

meningeal lymphatic drainage aggravates Parkinson's disease-like pathology in mice

overexpressing mutated α-synuclein. Transl. Neurodegener. 8, 7.

https://doi.org/10.1186/s40035-019-0147-y

764   **Supplementary Materials**

765   **Fig. S1** to **S4** for multiple supplementary figures

Fig. S1 gBOLD is coupled with CSF changes in HCP-Aging data. (**A**) **Top**: CSF region at

bottom fMRI slice shown in an example subject; **Bottom**: location of bottom rsfMRI slice marked

in corresponding structural MRI (dashed line). (**B**) gBOLD (blue) and CSF (red) rsfMRI signals

showed a coupled change (black arrows) from a representative example. (**C**) Averaged cross-

correlation function between gBOLD and CSF across 719 subjects. The red dashed line marks the

time lag (+3.2-sec) where the negative peak of the mean cross-correlation occurs. The shaded

regions represent the area within one standard error of the mean.

Fig. S2 The age-coupling association was not affected by head motion during fMRI

acquisition. Similar results as **Fig.1** were found when the mean FD was regressed out from the

gBOLD-CSF coupling for each rsfMRI session.

Fig. S3 The age-coupling association was not dominant by the whole brain volume. Consistent

results as **Fig.1** were found when we regressed out the whole brain volume (ventricles excluded)

from the gBOLD-CSF coupling for each rsfMRI session.

Fig. S4 The age-coupling association was not affected by the whole CSF volume. Similar

results as **Fig.1** were found when the volume of the whole CSF area was regressed out from the

gBOLD-CSF coupling for each rsfMRI session.