Chromosomal instability can favor macrophage-mediated immune response and induce

a broad, vaccination-like anti-tumor IgG response

Brandon H. Hayes[1-3], Mai Wang[1,2], Hui Zhu[1,2], Steven H. Phan[1,2], Jason C. Andrechak[1-3], Alexander H.

Chang[1,2], Lawrence J. Dooling[1,2], Michael P. Tobin[1-3], Tristan Marchena[1,2], Dennis E. Discher[1-3,†]

[1]Molecular and Cell Biophysics Lab, [2]Physical Sciences Oncology Center at Penn, [3]Bioengineering

Graduate Group, University of Pennsylvania, Philadelphia, PA, USA.

†Correspondence: discher@seas.upenn.edu

Abstract

Chromosomal instability (CIN), a state in which cells undergo mitotic aberrations that generate

chromosome copy number variations, generates aneuploidy and is thought to drive cancer

evolution. Although associated with poor prognosis and reduced immune response, CIN

generates aneuploidy-induced stresses that could be exploited for immunotherapies.

Macrophages, particularly, have been understudied in the CIN context. Here, through MPS1

inhibition-induced CIN in poorly immunogenic B16F10 mouse melanoma, we find that CIN-

afflicted cancer cells skew macrophages towards an anti-cancer phenotype while also pushing

them away from a pro-cancer one. We confirm these findings via RNA-sequencing, protein

expression, and short-term tumor studies. These results further translate to *in vivo* efficacy in

suppressing tumor growth: Mice can survive challenges of CIN-afflicted tumors. Long-term

survival, however, is dependent on CD47 expression and IgG opsonization. Mice bearing CIN-

afflicted tumors with wild-type CD47 levels see prolonged survival compared to their

chromosomally stable counterparts, but all succumb. Mice bearing CIN-afflicted CD47 knockout

tumors, however, show 28% long-term survival. When CD47 knockout was further paired with

IgG opsonization, survival rate increased to 97%. Successful rejection and clearance of CIN-

afflicted tumors induced *de novo* anti-cancer IgG antibodies that were multi-epitope and

functionally promoted macrophage-mediated phagocytosis. These *de novo* IgG antibodies could

also suppress *in vitro* tumoroid and *in vivo* tumor growth in a CD47 knockout context. These

results highlight an unexpected therapeutic benefit from CIN when paired with maximal

macrophage anti-cancer activity: an anti-cancer vaccination-like antibody response that can lead

to durable cures and further potentiate cell-mediated acquired immunity.

(250 words; 250 max)

Keywords: macrophages, aneuploidy, phagocytosis, CD47, antibodies

Introduction

Chromosomal instability (CIN) has long been thought to be indicative of poor prognosis and

reduced immune cell activity against tumors (Davoli et al., 2017; Vasudevan et al., 2021). CIN

is a state of high frequency of chromosome mis-segregation, which often generates micronuclei

and can ultimately cause aneuploidy—an abnormal number of chromosomes. CIN-induced

genomic heterogeneity can serve as a tumor promotor (Sheltzer et al., 2017) and allow some

tumor subpopulations to favor aggression, metastatic potential, immune evasion, and resistance

to therapies (Ben-David & Amon, 2019; Chunduri & Storchová, 2019; Vasudevan et al., 2021).

However, early-stage CIN also induces anti-cancer vulnerabilities (Cohen-Sharir et al., 2021;

Vasudevan et al., 2020), such as proliferation deficits (Wang et al., 2021). These early-stage

CIN-afflicted cells have yet to adapt and achieve aneuploidies that favor growth and immune

evasion (Tripathi et al., 2019, Vasudevan et al., 2021). CIN in diploid cells that is caused by

spindle checkpoint disruption by MPS1 inhibition (MPS1i) further induces a senescence-

associated secretory pathway phenotype and upregulation of NF-κB and interferon-mediated

pathways, among others, that drive immune clearance of chromosomally aberrant cells

(Santaguida et al., 2017; Wang et al., 2021). CIN and ploidy changes also inhibit tumor growth

in immunocompetent mice while having little effect in immunocompromised mice (Senovilla et

al., 2012; Boilève et al., 2013), suggesting CIN can somehow increase immunogenicity.

Recent analyses of The Cancer Genome Atlas (TCGA) particularly showed that highly aneuploid

tumors have macrophages that are polarized toward a pro-cancer, M2-like phenotype, among

other pro-cancer immune changes (Davoli et al., 2017; Taylor et al., 2018). MPS1i-treated

cancer cells have also been occasionally found to escape immune-mediated clearance (Wang

et al., 2021), but another study further found that cancer cells respond to CIN ultimately by

triggering IL-6-STAT3 signaling (Hong et al., 2022) that protect from CIN-induced cell death,

minimize interferon-related anti-cancer responses, and allow cells to adapt to CIN- and

aneuploidy-induced stresses. Among these studies, two key observations stood out. First, ploidy

changes tend to increase factors that can promote macrophage-mediated phagocytosis (Chao

et al., 2010; Krysko et al., 2018). Second, pro-survival signaling amidst CIN leads to an increase

of other factors (such as IL6) that induce a pro-cancer, M2-like phenotype (Fernando et al.,

2014). Together, these observations led us to hypothesize that macrophages may play a key
effector role in confronting and influencing CIN- and aneuploidy-afflicted tumors.

Macrophages have become attractive candidates for immunotherapies due to their ability to
phagocytose cancer cells (Alvey et al., 2018; Klichinsky et al., 2020). Generally, phagocytosis
of 'self' cells is inhibited by the key macrophage checkpoint interaction between SIRPα on the
macrophage and CD47 on the target cell (Oldenborg et al., 2000). CD47 is ubiquitously
expressed on all cells, including cancer cells (Willingham et al., 2012).   While tumor cell
engulfment can be driven to some extent via IgG opsonization by using anti-tumor monoclonal
antibodies that bind Fc receptors on macrophages (Alvey et al., 2017; Morrissey et al., 2020;
Suter et al., 2021; Tsai & Discher, 2008), this is generally insufficient to eliminate cancers,
especially solid tumors. Maximal macrophage-mediated phagocytosis is achieved when CD47-
SIRPα signaling is disrupted, in addition to IgG opsonization. However, achieving complete
tumor rejection is still a major challenge for macrophage-oriented therapies (Ingram et al., 2017;
Sockolosky et al., 2016), even after having identified these three key factors. Previous studies
achieved durable cures in immunocompetent mice, with some acquired immunity benefits, but
tumor rejection success is highly variable (Dooling et al., 2022; Hayes et al., 2022). Additionally,
in vivo macrophages often polarize toward tumor-associated macrophage (TAM) phenotypes,
which correlate with poor clinical prognoses (Cerezo-Wallis et al., 2020; Noy & Pollard, 2014).
TAMs also have poor phagocytic function and promote tumor growth and invasion (Alvey et al.,
2017; Georgouli et al., 2019). Nonetheless, CIN could create conditions in which macrophages
are not outcompeted in number by cancer cell proliferation and avoid becoming TAM-like.

Inspired by recent advances in exploiting CIN-related anti-cancer vulnerabilities, we
hypothesized that early-stage CIN in cancer cells may stimulate higher anti-cancer macrophage
activity than their chromosomally stable counterparts (under conditions of maximal
phagocytosis). Our results provide detailed *in vitro* and *in vivo* evidence that early-stage CIN in
cancer favors anti-cancer macrophage activity and consistently leads to durable cures in mice
when under conditions of maximal phagocytosis. Elimination of these CIN-afflicted solid tumors
further drives development of both anti-cancer opsonizing IgGs and enhanced cell-mediated
immunity, both of which help suppress growth against aggressive chromosomally stable tumors.

Results

MPS1i-induces CIN and generates conditions that favor anti-cancer macrophages while

minimizing skewing towards a pro-cancer macrophage phenotype

To investigate the possible effects that CIN may have in mediating macrophage immune

response, we chose to work with the poorly immunogenic B16F10 mouse melanoma model. To

induce CIN in B16F10, we treated cells with the MPS1 inhibitor reversine (Hong et al., 2022;

Kitajiam et al., 2022; Santaguida et al., 2010; Santaguida et al., 2017). Treatment occurred for

24 h, after which the drug was washed out, and cells were allowed to recover for an additional

48 h (**Fig. 1A**). After the recovery period, we quantified CIN in B16F10 cells, characterized its

effects in early-stage tumors, and studied potential effects on macrophages *in vitro* (**Fig. 1A**).

We first confirmed that MPS1i treatment increased the frequency of micronuclei formation in

interphase cells (**Fig. 1B**), as micronuclei are often used as a surrogate for CIN and genome

instability (Cohen-Sharier et al., 2020; Crasta et al., 2012; Harding et al., 2017; Mackenzie et al.,

2017). Regardless of the MPS1i concentration used, we saw substantial increases over the cell

line's baseline level of micronuclei formation. We further verified these results with other MPS1i

inhibitors AZ3146 and BAY 12-17389 (**Fig. S1A**) to confirm the effect was not somehow limited

to just reversine. Upon confirmation of increased micronuclei induction, we then proceeded to

quantitatively confirm and assess copy number variations that resulted from MPS1i treatment

using single-cell RNA-sequencing. For DMSO-treated B16F10 cells, approximately 10% of the

population were considered aneuploid outliers (**Fig. 1C-top; S1B-C**) – cells with copy number

profiles that are 2.5 standard deviations away from the distribution peak (see **Methods**).

However, 34% of the MPS1i-treated B16F10 cells were aneuploid (**Fig. 1C-bottom; S1B-C**).

After confirming that MPS1i indeed induces CIN and quantifiable aneuploidy in B16F10 cells,

we next sought to understand how CIN may affect early-stage immune response and tumor

development. We proceeded to establish tumors in mice with either MPS1i-treated or DMSO-

treated B16F10 (following the same schema outlined in **Fig. 1A**). We isolated tumors from mice

at two timepoints, five and ten days after initial challenge, and then we processed whole-tumors

for bulk RNA-sequencing (**Fig. 1D**). Comparison of tumors comprised of MPS1i-treated and

DMSO-treated cells by differential gene expression showed distinct downregulated transcripts

for numerous genes encoding M2-like (pro-cancer) macrophage polarization markers (**Fig. 1E**).

At day 5 post-challenge, we see that classical M2-like, pro-cancer markers *Arg1*, *Marco*, and

Cd274 are downregulated. However, several M2 markers are upregulated at this timepoint:

Mrc1, *Cd163*, and genes related to the complement component 1a (C1q). By day 10, *Marco*,

Mrc1, *Cd163, Cd274,* and the C1q complex genes are now all downregulated. *Arg1*, however,

returns to either basal or increased expression on day 10. Overall, though, we generally see

globally decreased expression for most markers by day 10. Furthermore, cytokines *Ccl2*, *Ccl4*,

Ccl7, *Ccl22*, and *Il21r*, which associated with an M2-like, pro-cancer macrophage phenotype

(Cerezo-Wallis et al., 2020), were found to be consistently downregulated at both days 5 and 10

post-challenge. Gene set enrichment analysis further revealed that numerous pathways related

cell cycle and proliferation were also downregulated (**Fig. 1F**), consistent with previous studies

(Cohen-Sharir et al., 2020; Wang et al., 2021). These downregulated pathways suggest longer

tumor doubling time since cells have yet to adapt to these CIN-associated proliferation deficits.

The identification of downregulated M2-like macrophage markers in CIN-afflicted tumors next

led us to better assess how early-stage CIN in cancer may influence macrophages specifically.

Previous studies (Cerezo-Wallis et al., 2020; Xian et al., 2021) have found that the secretome

can have profound and rapid polarization effects on bone marrow-derived macrophages

(BMDMs) *in vitro*. We proceeded to perform a similar secretome study, in which conditioned

media from MPS1i-treated or DMSO-treated B16F10 cells was collected and added to previously

differentiated BMDMs. BMDMs were treated with conditioned media for 24 h, after which they

were collected and processed for single-cell RNA-sequencing (**Fig. 1G**). Unsupervised

clustering analysis using uniform manifold approximation and projection (UMAP) nonlinear

dimensionality reduction identified four distinct macrophage population clusters: two clusters (0

and 2) that consisted of approximately 75% of macrophages treated with MPS1i-treated B16F10

secretome and two clusters (1 and 3) that consisted of approximately 75% macrophages treated

with DMSO-treated B16F10 secretome (**Fig. 1H**).

The gene expression profiles of these clusters showed that the two clusters (1 and 3) that

consisted mostly of macrophages treated with DMSO-treated B16F10 secretome had increased

expression of anti-inflammatory, M2-like pro-cancer macrophage polarization markers (**Fig. 1I**).

Consistent with the whole-tumor bulk RNA-seq, we were successfully able to capture increased

expression of *Mrc1* in both clusters 1 and 3 and observed increased expression of *Ccl2*, *Ccl4*,

and *Ccl7* in cluster 3. More importantly though, gene expression revealed that clusters 1 and 3

tended to downregulate many genes associated with a pro-inflammatory, M1-like anti-cancer

response (**Fig. 1J**). Clusters 0 and 2, on the other hand, which consisted mostly of MPS1i-treated

B16F10 secretome, showed little-to-no expression for most M2-like, pro-cancer markers (**Fig.**

1I) and increased expression of pro-inflammatory, anti-cancer markers (**Fig. 1J**). These results

suggest that early-stage CIN in cancer cells engages pro-inflammatory macrophage activity.

Macrophages polarize to an M1-like phenotype and can clear CIN-afflicted tumoroids

Next, we sought to experimentally confirm the polarization effects. Here, we use B16F10 CD47

knockout (KO) cells (**Fig. S2A**) to remove the inhibitory effects of CD47 on phagocytosis and

more easily study macrophage engulfment ability (**Fig. S2B-C**). Moving forward, we chose to

use a B16F10 tumoroid model to better simulate the biophysical properties of the 3D tumor

microenvironment and proliferative capacity of tumor masses (**Fig. 2A**) (Dooling et al., 2022).

We chose this model primarily because 2D phagocytosis assays fail to consider the mechanical

cohesive forces between tumor cells and other biophysical aspects that can affect immune

response (Nia et al., 2020). We have previously found that macrophages need to phagocytose

cooperatively—large numbers of macrophages working together to overcome the mechanical

strength of cohesion between solid tumor cells—to achieve durables responses (Dooling et al.,

2022). CIN can also often induce additional effects on cells in its early stages, such as increases

in cell size due to cytokinesis defects and poly-aneuploidy (Chunduri & Storchová, 2019; Mallin

et al., 2022). Single macrophages may struggle to phagocytose these large targets in a

conventional 2D assay (Champion et al., 2006), even with IgG opsonization, which we

experimentally confirmed with MPS1i-treated cells (**Fig. S2B**). MPS1i treatment, in fact,

increases cell size, as measured by flow cytometry (**Fig. S2C**), although size effects are

confounded by decreased Tyrp1 expression, the antigen targetable for opsonization using anti-

Tyrp1 (**Fig. S2D**). CD47 levels were relatively consistent across treatment conditions and similar

to non-perturbed levels (**Fig. S2E**) for B16F10 sgCtrl. We thus hypothesized that cooperative

phagocytosis (Dooling et al., 2022) may be able to overcome the CIN-induced size increases

and reduced binding of anti-Tyrp1. BMDMs were added to pre-assembled tumoroids at a 3:1

ratio, and this "immuno-tumoroid" co-culture was maintained for five days. We first assessed

surface protein expression of macrophage polarization markers. Consistent with the polarization

trends in our whole-tumor bulk RNA-sequencing and single-cell RNA-sequencing of BMDMs

(**Fig. 1E, 1I-J**), we find that BMDMs co-cultured in tumoroids of MPS1i-treated B16F10 showed

increased expression of M1-like, anti-cancer macrophage markers MHCII and CD86 while

having decreased expression of M2-like, pro-cancer markers CD163 and CD206 (**Fig. 2B-C**).

We next wondered if these polarization studies translated to functional effects on macrophage-

mediated phagocytosis and clearance of tumoroids. We again prepared tumoroids, later adding

BMDMs to these pre-assembled tumoroids at a 3:1 ratio with either anti-Tyrp1 or mouse IgG2a

isotype control for IgG opsonization. In our 3D tumoroid *in vitro assays*, we found that

macrophages can suppress the growth of chromosomally unstable tumoroids and clear them,

surprisingly both with and without anti-Tyrp1 (**Fig. 2D-E**), regardless of MPS1i concentration

used for treatment. In contrast, DMSO-treated B16F10 cells (chromosomally stable with regular

proliferative capacity) require anti-Tyrp1 for macrophages to clear them (**Fig. 2D, S3B**). These

results link the phenotypic studies with functional effects: CIN can induce anti-cancer

macrophage phenotypes to favor macrophage-mediated clearance.

We also attempted to use two other MPS1i inhibitors to see if we achieved similar results:

tumoroids made of AZ3146-treated cells, which BMDMs could only suppress at very high

concentrations (**Fig. S3C-i**), and tumoroids made of BAY 12-17389-treated cells, which were

heavily suppressed and/or cleared similarly to tumoroids comprised of reversine-treated cells

(**Fig. S3C-ii**). Given that AZ3146-treated cells tended to proliferate well, we next sought to verify

if tumoroids made of cells treated with either BAY 12-17389 or reversine would eventually grow.

We observed that tumoroids decreased in size 24 h after plating, most likely due to some cell

death (**Fig. S3D**). Afterwards though, tumoroids generally showed some proliferation, albeit with

deficits. This suggests that the anti-cancer macrophage activity observed may require CIN be

severe enough to induce some proliferation deficits. Otherwise, low-level CIN may be tolerable

for cancer cells, eventually favor their evolution (Vasudevan et al., 2020), and allow cancer cells

to still proliferate more rapidly than phagocytosis kinetics can manage.

Lastly, we sought to confirm both our transcriptomic analyses and tumoroid surface marker

expression before proceeding to long-term *in vivo* experimentation. We proceeded to establish

tumors in mice with either MPS1i-treated or DMSO-treated B16F10 cells (the drug treatment

followed the same schema outlined in **Fig. 1A**). We then isolated tumors five days post-challenge

for immune infiltrate analyses by flow cytometry (**Fig. 2F, S5**). We found that CIN-afflicted tumors

had approximately 3-fold more CD45+ immune cells (**Fig. 2G-i**), 2.5-fold more CD45+ CD11b+

(myeloid) cells (**Fig. 2G-ii**), and a near 6-fold increase in F4/80-positive macrophage (**Fig. 2G-**

iii) compared to their DMSO counterparts. Furthermore, we found that CIN-afflicted tumors

showed approximately 6-fold more CD86^{hi} CD206^{lo} (M1-like, anti-cancer) macrophages

compared to their DMSO counterparts. We should note that these tumors also saw an increase

in CD86^{lo} CD206^{hi} (M2-like, pro-cancer) macrophages compared to DMSO control (**Fig. 2G-iv**).,

but this increase is expected given that there are generally more macrophages infiltrating these

tumors. Despite the increase in M2-like macrophages, these tumors made of MPS1i-treated cells

still have roughly 2-fold more M1-like macrophages (**Fig. 2G-iv**). These results suggest that

although other immune cell types infiltrate tumors, which is consistent with upregulated surface

expression of H2-K^b (**Fig. S4**), macrophage functional activity is indeed positively affected by

CIN-afflicted cancer cells.

MPS1i-induced CIN favors tumor rejection with IgG opsonization and CD47 disruption

While previous studies show that complete CD47 ablation can favor suppression and even

complete rejection of IgG-opsonized B16F10 tumors (Andrechak et al., 2022; Dooling et al.,

2022; Hayes et al., 2022; Kamber et al., 2021), inter-experimental variation is high, particularly

regarding complete tumor rejection and clearance. Furthermore, how CIN could influence these

results is unknown. Therefore, we sought to assess if CIN in B16F10, which thus far has shown

to skew macrophages toward an anti-cancer phenotype and away from pro-cancer one, can

translate to improved efficacy and consistency in macrophage-oriented therapies. For these

subsequent *in vivo* experiments, we first pre-treated B16F10 cells with either MPS1i or DMSO

for 24 h, washed the drug out, and then allowed the cells to recover for an additional 48 h (**Fig.**

3A). After the recovery period elapsed, we established tumors in mice, with either MPS1i-treated

or DMSO-treated B16F10 cells. To further test for CD47-mediated effects, we used either

B16F10 CD47 KO or B16F10 sgCtrl, expressing wild-type CD47 (WT) levels. Starting four days

post-challenge, mice received either anti-Tyrp1 or mIgG2a isotype control for opsonization.

As expected, all tumors comprised of DMSO-treated B16F10 sgCtrl showed exponential growth

and no survivors (**Fig. 3B**), consistent with previous studies (Dooling et al., 2022; Hayes et al.,

2022). These results re-confirm the inhibitory effect that CD47 has on macrophage-mediated

immunity (Ingram et al., 2017; Sockolosky et al., 2016; Willingham et al., 2012), suppressing

macrophage immune response even with anti-Tyrp1 IgG opsonization. Mice with CIN-afflicted

B16F10 sgCtrl tumors ultimately showed exponential growth and no survivors as well, but they

did show increased median survival. Furthermore, median survival increased even further when

mice were treated with anti-Tyrp1, such that all mice were considered partial responders

(survival of 20+ days, one week longer than median survival of tumors comprised of DMSO-

treated B16F10 sgCtrl without anti-Tyrp1 treatment). All tumors comprised of DMSO-treated

B16F10 CD47 KO showed exponential growth and no survivors (**Fig. 3C**), regardless of anti-

Tyrp1 treatment or not. However, CIN-afflicted B16F10 CD47 KO tumors showed more positive

outcomes. Even without anti-Tyrp1, all challenged mice with CIN-afflicted CD47 KO tumors were

either cured completely (28%) or considered partial responders. When paired with anti-Tyrp1

treatment, 97% of mice challenged with CIN-afflicted CD47 KO tumors survive.

Long-term survival results show that challenging mice with both MPS1i-treated and DMSO-

treated B16F10 sgCtrl failed to yield any survivors, regardless of anti-Tyrp1 opsonization or not

(**Fig. 3D-i**). Similarly, we also failed to generate any survivors among mice challenged with

DMSO-treated B16F10 CD47 KO cells, regardless of anti-Tyrp1 opsonization or not (**Fig. 3D-ii**).

This again highlights a challenge in optimizing macrophage-mediated therapies: achieving

consistency in long-term therapeutic outcomes. Mice challenged with MPS1i-treated B16F10

CD47 KO cells, however, were able to survive, both without anti-Tyrp1 (28% survival) and with

anti-Tyrp1 (97% survival) (**Fig. 3D-ii**). To determine if the degree of CIN affected survival, we

also challenged mice with B16F10 CD47 KO cells treated with varying concentrations of

reversine. Regardless of MPS1i concentration, >80% of mice survive when also treated with

anti-Tyrp1 (**Fig. S6**). These results show that, in physiologically relevant microenvironments,

early-stage CIN can favor survival when paired with IgG opsonization and CD47 disruption. This

further suggests that macrophages are key effector cells in achieving survival against CIN-

afflicted tumors, since 97% survival was achieved under conditions of maximal phagocytosis.

Although no mice challenged with CIN-afflicted B16F10 sgCtrl survived, MPS1i increase median

survival significantly when paired with anti-Tyrp1 opsonization (**Fig. 3D-i**). This suggests that

macrophages still display some anti-cancer activity, despite the inhibitory effects of CD47, and

are important effector cells in final therapeutic outcome. To better support the hypothesis that

macrophages are indeed key effector cells in rejecting CIN-afflicted tumors, we established

tumors comprised of wild-type (WT) B16F10 in mice. Although WT tumors are generally

unaffected by anti-CD47 and anti-Tyrp1 (Dooling et al., 2022; Hayes et al., 2022), we

hypothesized that we could eliminate CIN-afflicted WT B16F10 tumors by providing adoptive

transfer of marrow to increase macrophage numbers to compensate for the CD47-mediated

inhibition of endogenous macrophages. Furthermore, we engineered marrow by priming Fcγ

receptors with anti-Tyrp1, initially inhibiting CD47-SIRPα interaction via anti-SIRPα antibody

blockade or combining both (**Fig. 3E-i**). As expected, mice challenged with CIN-afflicted WT

tumors and treated with regular, unprimed marrow only had ~14% survival (**Fig. 3E-ii & iii**). This

result, however, was identical to chromosomally stable WT tumors treated with marrow primed

with both anti-SIRPα and anti-Tyrp1. These results suggest increased macrophages numbers

do provide some minor benefit. As the marrow engineering becomes more rigorous to maximize

phagocytosis, survival improves. 37% of mice challenged with CIN-afflicted WT tumors survived

when treated with marrow that was initially engineered with anti-SIRPα. 50% survived with

marrow primed with anti-Tyrp1. Lastly, 80% survived with anti-Tyrp1-primed and anti-SIRPα-

blocked monocytes (**Fig. 3E-ii & iii**). Altogether, these results support three conclusions: (1) the

increase in survival attributed to anti-SIRPα supports the idea that CD47 modulates therapeutic

outcome even during CIN; (2) the increase in survival attributed to anti-Tyrp1 highlights the

importance of IgG opsonization; (3) the high success from the combination emphasizes

macrophages playing a key role in achieving complete rejection and clearance.

Clearance of CIN-afflicted tumors promotes *de novo* pro-phagocytic & anti-cancer IgG

Macrophages and related phagocytic cells make up innate immunity, a first line of defense

against pathogens and disease, but they are also responsible for initiating acquired immunity.

The activation of the acquired immune response involves two branches of immunity: humoral

(mediated by macromolecules such as antibodies) and cell-mediated immunity (involving T cells,

for example). We hypothesized that mice that survived challenges with CIN-afflicted tumors

would show signs of an acquired immune response due to their high survival rate.

We collected convalescent serum from survivors to quantify *de novo* anti-cancer IgG antibodies

that may have resulted from successful rejection and clearance of CIN-afflicted tumors (**Fig. 4A**).

Sera was collected and then subsequently used in antibody binding testing and Western blotting

to confirm emergence of anti-B16F10 antibodies. We first quantified IgG2a and IgG2b titers in

convalescent sera, both of which have been previously found to engage mouse macrophage

Fcγ receptors (Bruhns, 2012; Nimmerjahn et al., 2010) that are typically required for

macrophage-mediated phagocytosis. B16F10 cells, either Tyrp1-expresing or Tyrp1 KO, were

incubated with sera (convalescent from survivors or naïve from unchallenged mice) and then

counterstained with conjugated antibodies against IgG2a/c and IgG2b (**Fig. 4B-i & ii**). All mice

that survived challenges from CIN-afflicted tumors yielded sera that showed significantly large

increases in IgG2a/c binding against both Tyrp1-positive and Tyrp1 KO B16F10 cells. We

similarly saw a statistically significant increases in IgG2b binding against both Tyrp1-positive and

Tyrp1 KO B16F10 cells using the same sera, although the increases in binding were more

variable. Altogether, we see that surviving CIN-afflicted tumors generates IgG-rich sera, like

previous studies (Dooling et al., 2022; Hayes et al., 2022). The increases in binding observed

against Tyrp1 KO cells also suggest that IgG antibodies target a repertoire of antigens beyond

Tyrp1, which we further qualitatively confirmed via Western blotting (**Fig. 4B-iii**). Convalescent

sera were used to immunoblot against B16F10 lysates, revealing many bands at multiple

molecular weights and more bands than when immunoblotting with naïve sera, supporting our

hypothesis of antigen broadening beyond Tyrp1. Lastly, we also tested IgG2a and IgG2b titers

from additional *in vivo* experiments: survivors from both titrated CIN-afflicted tumors from **Fig.**

S6 and adoptive marrow transfers in **Fig. 3E**. Similarly, convalescent sera from all these mice

show increases in IgG2a/c and IgG2b in both Tyrp1-expressing and Tyrp1 KO B16F10 cells

(**Fig. S7**). These results confirm that regardless of the method used to exploit CIN, induction of

anti-cancer IgG can be expected.

To test where these *de novo* serum antibodies functionally promote macrophage-mediated
phagocytosis, we performed conventional 2D phagocytosis assays in which cancer cell
suspensions were opsonized with sera (or anti-Tyrp1 or mouse IgG2a isotype as controls) (**Fig.**
4C-i). Under conditions of CD47 KO, we see that nearly all unpurified convalescent sera
increased phagocytosis relative to naïve serum and mIgG2a isotype control. Furthermore, this
increase in phagocytosis is identical to that provided by anti-Tyrp1 (both ~5-fold higher than
baseline). We also find that sera continue to promote phagocytosis even in B16F10 CD47/Tyrp1
double KO cells (~3-fold higher than baseline). Convalescent sera are still able to provide
increases in phagocytosis against double KO cells, whereas anti-Tyrp1 expectedly does not
drive phagocytosis due to lack of antigen. This again supports the hypothesis of acquired
immunity with *de novo* IgG antibodies that target B16F10 antigens beyond Tyrp1. Upon
confirming the functional effect of *de novo* IgG in the convalescent sera, we then wondered if
convalescent serum IgG would be able to suppress tumoroid growth, given that this model better
captures both the biophysical microenvironment and proliferative capacity of tumors (Dooling et
al., 2022). Indeed, we found that convalescent serum IgG added simultaneously with
macrophages to B16F10 CD47 KO tumoroids led to either tumoroid elimination or significantly
suppressed growth (**Fig. 4C-ii**), although the efficacy was less potent as anti-Tyrp1. This
suggests that perhaps the polyclonal *de novo* IgGs here still lack the specificity benefits that
accompany a monoclonal antibody such as anti-Tyrp1. Nonetheless, these results demonstrate
induction of a generally potent anti-cancer antibody response B16F10 in a CD47 KO context.

We then proceeded to test the function of convalescent serum *in vivo* by opsonizing B16F10
CD47 KO cells just prior to subcutaneous implantation in naïve mice. For comparison, we also
opsonized B16F10 CD47 KO cells with either anti-Tyrp1 (positive control) or mIgG2a control
(negative control). We found that both convalescent sera and anti-Tyrp1 suppressed tumor
growth by days 11 and 13 relative to mIgG2a control (**Fig. 4D-i**). Interestingly, we found that
convalescent sera showed a trend of suppressing growth more than anti-Tyrp1, although this
was not statistically significant. We continued to monitor all mice for long-term survival, and we
found that this pre-opsonization with both convalescent sera and anti-Tyrp1 eliminated tumors
in challenged mice with near identical cure rates (25% and 22%, respectively) (**Fig. 4D-ii**).

Altogether, we found that the convalescent sera from mice originally challenged with

chromosomally unstable tumors has potent anti-cancer effects *in vitro* and *in vivo*.

Acquired immunity suppresses growth of chromosomally stable tumors and becomes

more effective with ongoing challenges of CIN-afflicted tumors

The anti-cancer IgG antibody development in survivors led us to further hypothesize that we

should see improved median survival and/or survival rate if surviving mice were re-challenged.

We therefore challenged surviving mice with a second injection of either DMSO-treated or

B16F10 CD47 KO cells (**Fig. 4E**). If additional survivors resulted from this experiment, we also

intended to undergo a third challenge, akin to a prime-boost strategy for anti-cancer vaccination.

It should be highlighted that starting from this second challenge, no mice received anti-Tyrp1.

This was done to maximally challenge acquired immunity and to better simulate the possibility

of recurrence post-therapy. Age-matched naïve mice receiving their first challenge responded

similarly to the younger cohorts (**Fig. 3B**). Of the previously cured mice, only a single mouse (of

11 total) survived a challenge with DMSO-treated B16F10 cells (**Fig. 4F**). However, median

survival increased (21 days) compared to their naïve counterparts (14 days), supporting the

initial hypothesis of prolonged survival. Survivors that were re-challenged again with MPS1i-

treated cells, however, showed 100% survival, even in the absence of anti-Tyrp1 (**Fig. 4F**). Age-

matched naïve mice receiving their first challenge of MPS1i-treated cells responded relatively

similarly (a single survivor out of 6 total, 17% survival) to the younger cohort (28% survival). This

complete success rate against a second challenge of CIN-afflicted B16F10 CD47 KO further

supports an acquired immune response, at least against ongoing chromosomally unstable cells.

Mice that failed to reject re-challenge tumors comprised of DMSO-treated B16F10 CD47 KO in

Fig. 4F were euthanized and had their tumors harvested to measure their immune cell infiltrate

by flow cytometry (**Fig. S8**). Although these mice did not survive, we found that these previous

survivors showed roughly a 2-fold increase in the number of immune cells in their tumors (**Fig.**

4G-i), a 2-fold increase in the number of CD3+ CD8+ T cells (**Fig. 4G-ii**), and a near 3-fold

increase in the number of F4/80+ macrophages (**Fig. 4G-iii**) compared to their naïve

counterparts. We further found that of the F4/80+ macrophage infiltrate, previous survivors had

roughly four times as many MHCII-high macrophages (M1-like, anti-cancer). These immune

infiltrate analyses suggest that although the acquired immune response in these previous

survivors was still not potent enough to clear chromosomally stable and regularly proliferating

B16F10 CD47 KO tumors, it did enhance cell-mediated immunity.

Lastly, we performed a third tumor challenge on second challenge survivors with untreated

B16F10 CD47 KO cells (chromosomally stable and regular proliferating). Mice were left

untreated post-challenge (no anti-Tyrp1). ~56% of these mice completely resisted tumor growth.

Of the four mice that developed tumors, two had to be euthanized prematurely due to tumor

rupture but were indeed showing signs of growth suppression. The last two mice consisted of

two long-term partial responders: one whose tumor did not reach terminal burden until 82 days

post-challenge and another who experienced stable tumor regression with almost no regrowth.

Overall, the 56% survival rate in the third challenge, in the absence of anti-Tyrp1, and the partial

responses observed in mice that grew tumors suggest a durable immunological response that

results from facing and clearing ongoing chromosomally unstable tumors, at least in the context

of CD47 disruption.

Discussion

Macrophage-oriented immunotherapies against solid tumors have maximal efficacy when three elements are combined: large numbers of macrophages for cooperativity, IgG opsonization that activates Fc receptors and stimulates macrophage-mediated phagocytosis, and disruption of the CD47 macrophage immune checkpoint (Dooling et al., 2022). However, even with these factors properly applied, complete tumor rejection and clearance is not guaranteed and varies greatly across *in vivo* mouse studies (Andrechak et al., 2022; Dooling et al., 2022; Hayes et al., 2022; Kamer et al., 2021). Here, we show a 97% survival rate of mice challenged with CIN-afflicted B16F10 tumors when maximizing macrophage-mediated activity. This result improves upon previous studies and is more impressive given that B16F10 are typically poorly immunogenic, do not respond to either anti-CD47 or anti-PD-1/PD-L1 monotherapies, and show large variability in cure rates (anywhere from 20-40%) even when macrophages should be maximally phagocytic.

These results suggest that CIN in early stages generates anti-cancer vulnerabilities that favor macrophage-mediated immune response, contingent on conditions of maximal phagocytosis. Immunocompetent mice consistently survive these challenges at high survival rates. These survivors also develop *de novo* anti-cancer IgG, similar to previous studies (Dooling et al., 2022; Hayes et al., 2022), that are pro-phagocytic, multi-epitope, and efficacious *in vivo*. The emergence of these IgGs could be synergistic with clinically relevant CD47 blockade treatments for solid tumors and help address concerns regarding resistance due to antigen loss (Jalil et al., 2020). Mice that are re-challenged with chromosomally stable CD47 KO tumors (and without exogenous anti-Tyrp1 opsonization) shows increased median survival and increased immune infiltrate, further supporting the hypothesis of newly generated anti-cancer acquired immunity. More interestingly, though, we see that all mice re-challenged with CIN-afflicted CD47 KO tumors survive, even in the absence of anti-Tyrp1 opsonization. A third challenge of these two-time survivors with chromosomally stable CD47 KO tumors shows >50% mice survive and improved median survival for non-survivors. These results elucidate two advantages that CIN can provide to better therapeutic outcomes. First, early-stage CIN facilitates survival while generating potent *de novo* IgGs that can drive positive phagocytic feedback to minimize recurrence. Second, ongoing vulnerability-inducing CIN in tumor cells can both strengthen cell-

454   mediated acquired immunity and create an antigen reservoir for the maintenance of long-term
455   humoral immunity.
456
The effects of CIN and aneuploidy in macrophages nonetheless still require more investigation.
While secretome from MPS1i-treated cancer cells has been found to trigger expression of *Arg1*
and *Il6* (Xian et al., 2021), both of which are pro-cancer M2-like macrophage markers (Fernando
et al., 2014; Mujal et al., 2022), our findings suggest that polarization is much more complex.
Here, whole-tumor bulk RNA-sequencing hints at CIN-afflicted tumors having a macrophage
population that is both less anti-inflammatory and M2-like. Single-cell RNA-sequencing of
BMDMs treated with secretome from CIN-afflicted cells further suggests that CIN induces a
microenvironment that can pushes macrophages to a pro-inflammatory, anti-cancer phenotype
while minimizing polarization to a pro-cancer phenotype. Our transcriptomics analyses also align
more with deeper investigation that suggest additional markers are required for macrophage
polarization distinction (Jablonksi et al., 2015). We further confirm these findings by observing
increased surface protein expression of anti-cancer M1-like macrophages *in vitro* 3D tumoroid
co-cultures with CIN-afflicted cells and *in vivo* immune infiltrate experiments. Furthermore,
BMDMs show enhanced clearance of CIN-afflicted cells in 3D tumoroid phagocytosis assays.
Additionally, the aforementioned study (Xian et al., 2021) only found this trend in SKOV3 cells,
whereas their aneuploid fused B16 cells actually show decreased *Arg1* expression, suggesting
possible cell-intrinsic complications. Meta-analyses of aneuploid tumor genomics data from
TCGA also suggest reduced anti-cancer macrophage activity (Davoli et al., 2017; Xian et al.,
2021). Our study, however, distinctly highlights macrophage response at early timepoints of CIN,
during the which cancer cells have not yet adapted to CIN- and aneuploidy-induced stresses.
TCGA studies, on the other hand, are limited to much later timepoints when cells have overcome
to CIN- and aneuploidy-associated stresses and developed their own unique aneuploidies to
drive tumor progression. Additionally, more recent studies find that aneuploidy and CIN paired
with high tumor mutational burdens show increased median survival in patients (Spurr et al.,
2022a; Spurr et al., 2022b), suggesting possible other complexities that favor immune infiltration
and that could explain how CIN can favor macrophage-mediated immune response.
483
484

References

Ahn, J., Xia, T., Rabasa Capote, A., Betancourt, D., Barber, G.N., 2018. Extrinsic Phagocyte-
Dependent STING Signaling Dictates the Immunogenicity of Dying Cells. Cancer Cell 33,
862-873.e5. https://doi.org/10.1016/j.ccell.2018.03.027

Alvey, C.M., Spinler, K.R., Irianto, J., Pfeifer, C.R., Hayes, B., Xia, Y., Cho, S., Dingal, P.C.P.D.,
Hsu, J., Smith, L., Tewari, M., Discher, D.E., 2017. SIRPA-Inhibited, Marrow-Derived
Macrophages Engorge, Accumulate, and Differentiate in Antibody-Targeted Regression of
Solid Tumors. Current Biology 27, 2065-2077.e6. https://doi.org/10.1016/j.cub.2017.06.005

Andrechak, J.C., Dooling, L.J., Hayes, B.H., Kadu, S., Zhang, W., Pan, R., Vashisth, M., Irianto,
J., Alvey, C.M., Discher, D.E., 2022a. Cooperative phagocytosis underlies macrophage
immunotherapy of solid tumours and initiates a broad anti-tumour IgG response (preprint).
Bioengineering. https://doi.org/10.1101/2022.01.01.474150

Andrechak, J.C., Dooling, L.J., Tobin, M.P., Zhang, W., Hayes, B.H., Lee, J.Y., Jin, X., Irianto,
J., Discher, D.E., 2022b. CD47-SIRPα Checkpoint Disruption in Metastases Requires
Tumor-Targeting Antibody for Molecular and Engineered Macrophage Therapies. Cancers
14, 1930. https://doi.org/10.3390/cancers14081930

Ben-David, U., Amon, A., 2020. Context is everything: aneuploidy in cancer. Nat Rev Genet 21,
44–62. https://doi.org/10.1038/s41576-019-0171-x

Boilève, A., Senovilla, L., Vitale, I., Lissa, D., Martins, I., Métivier, D., van den Brink, S., Clevers,
H., Galluzzi, L., Castedo, M., Kroemer, G., 2013. Immunosurveillance against
tetraploidization-induced colon tumorigenesis. Cell Cycle 12, 473–479.
https://doi.org/10.4161/cc.23369

Bruhns, P., 2012. Properties of mouse and human IgG receptors and their contribution to
disease models. Blood 119, 5640–5649. https://doi.org/10.1182/blood-2012-01-380121

Cerezo-Wallis, D., Contreras-Alcalde, M., Troulé, K., Catena, X., Mucientes, C., Calvo, T.G.,
Cañón, E., Tejedo, C., Pennacchi, P.C., Hogan, S., Kölblinger, P., Tejero, H., Chen, A.X.,
Ibarz, N., Graña-Castro, O., Martinez, L., Muñoz, J., Ortiz-Romero, P., Rodriguez-Peralto,
J.L., Gómez-López, G., Al-Shahrour, F., Rabadán, R., Levesque, M.P., Olmeda, D.,
Soengas, M.S., 2020. Midkine rewires the melanoma microenvironment toward a
tolerogenic and immune-resistant state. Nat Med 26, 1865–1877.
https://doi.org/10.1038/s41591-020-1073-3

Champion, J.A., Mitragotri, S., 2006. Role of target geometry in phagocytosis. Proc. Natl. Acad.
Sci. U.S.A. 103, 4930–4934. https://doi.org/10.1073/pnas.0600997103
526
527   Chao, M.P., Jaiswal, S., Weissman-Tsukamoto, R., Alizadeh, A.A., Gentles, A.J., Volkmer, J.,
528   Weiskopf, K., Willingham, S.B., Raveh, T., Park, C.Y., Majeti, R., Weissman, I.L., 2010.
529   Calreticulin Is the Dominant Pro-Phagocytic Signal on Multiple Human Cancers and Is
530   Counterbalanced   by   CD47.   Sci.   Transl.   Med.   2.
531   https://doi.org/10.1126/scitranslmed.3001375
532
533   Chunduri, N.K., Storchová, Z., 2019. The diverse consequences of aneuploidy. Nat Cell Biol 21,
534   54–62. https://doi.org/10.1038/s41556-018-0243-8
535
Cohen-Sharir, Y., McFarland, J.M., Abdusamad, M., Marquis, C., Bernhard, S.V., Kazachkova,
M., Tang, H., Ippolito, M.R., Laue, K., Zerbib, J., Malaby, H.L.H., Jones, A., Stautmeister,
L.-M., Bockaj, I., Wardenaar, R., Lyons, N., Nagaraja, A., Bass, A.J., Spierings, D.C.J.,
Foijer, F., Beroukhim, R., Santaguida, S., Golub, T.R., Stumpff, J., Storchová, Z., Ben-
David, U., 2021. Aneuploidy renders cancer cells vulnerable to mitotic checkpoint inhibition.
Nature 590, 486–491. https://doi.org/10.1038/s41586-020-03114-6
542
543   Crasta, K., Ganem, N.J., Dagher, R., Lantermann, A.B., Ivanova, E.V., Pan, Y., Nezi, L.,
544   Protopopov, A., Chowdhury, D., Pellman, D., 2012. DNA breaks and chromosome
545   pulverization from errors in mitosis. Nature 482, 53–58. https://doi.org/10.1038/nature10802
546
547   Cunha, L.D., Yang, M., Carter, R., Guy, C., Harris, L., Crawford, J.C., Quarato, G., Boada-
548   Romero, E., Kalkavan, H., Johnson, M.D.L., Natarajan, S., Turnis, M.E., Finkelstein, D.,
549   Opferman, J.T., Gawad, C., Green, D.R., 2018. LC3-Associated Phagocytosis in Myeloid
550   Cells   Promotes   Tumor   Immune   Tolerance.   Cell   175,   429-441.e16.
551   https://doi.org/10.1016/j.cell.2018.08.061
552
553   Davoli, T., Uno, H., Wooten, E.C., Elledge, S.J., 2017. Tumor aneuploidy correlates with markers
554   of immune evasion and with reduced response to immunotherapy. Science 355, eaaf8399.
555   https://doi.org/10.1126/science.aaf8399
556
557   Fernando, M.R., Reyes, J.L., Iannuzzi, J., Leung, G., McKay, D.M., 2014. The Pro-Inflammatory
558   Cytokine, Interleukin-6, Enhances the Polarization of Alternatively Activated Macrophages.
559   PLoS ONE 9, e94188. https://doi.org/10.1371/journal.pone.0094188
560
Georgouli, M., Herraiz, C., Crosas-Molist, E., Fanshawe, B., Maiques, O., Perdrix, A., Pandya,
P., Rodriguez-Hernandez, I., Ilieva, K.M., Cantelli, G., Karagiannis, P., Mele, S., Lam, H.,
Josephs, D.H., Matias-Guiu, X., Marti, R.M., Nestle, F.O., Orgaz, J.L., Malanchi, I.,

Fruhwirth, G.O., Karagiannis, S.N., Sanz-Moreno, V., 2019. Regional Activation of Myosin
II in Cancer Cells Drives Tumor Progression via a Secretory Cross-Talk with the Immune
Microenvironment. Cell 176, 757-774.e23. https://doi.org/10.1016/j.cell.2018.12.038
Harding, S.M., Benci, J.L., Irianto, J., Discher, D.E., Minn, A.J., Greenberg, R.A., 2017. Mitotic
progression following DNA damage enables pattern recognition within micronuclei. Nature
548, 466–470. https://doi.org/10.1038/nature23470
570
571 Hayes, B.H., Tsai, R.K., Dooling, L.J., Kadu, S., Lee, J.Y., Pantano, D., Rodriguez, P.L.,
572   Subramanian, S., Shin, J.-W., Discher, D.E., 2020. Macrophages show higher levels of
573   engulfment after disruption of *cis* interactions between CD47 and the checkpoint receptor
574   SIRPα. Journal of Cell Science 133, jcs237800. https://doi.org/10.1242/jcs.237800
575
576 Hayes, B.H., Zhu, H., Andrechak, J.C., Discher, D.E., 2022. Titrating CD47 by mismatch
577   CRISPRi reveals incomplete repression can eliminate IgG-opsonized tumors but CD47
578   heterogeneity limits induction of anti-tumor IgG (preprint). Cancer Biology.
579   https://doi.org/10.1101/2022.09.27.509740
580
581 Ingram, J.R., Blomberg, O.S., Sockolosky, J.T., Ali, L., Schmidt, F.I., Pishesha, N., Espinosa,
582   C., Dougan, S.K., Garcia, K.C., Ploegh, H.L., Dougan, M., 2017. Localized CD47 blockade
583   enhances immunotherapy for murine melanoma. Proc. Natl. Acad. Sci. U.S.A. 114, 10184–
584   10189. https://doi.org/10.1073/pnas.1710776114
585
586 Jablonski, K.A., Amici, S.A., Webb, L.M., Ruiz-Rosado, J. de D., Popovich, P.G., Partida-
587   Sanchez, S., Guerau-de-Arellano, M., 2015. Novel Markers to Delineate Murine M1 and M2
588   Macrophages. PLoS ONE 10, e0145342. https://doi.org/10.1371/journal.pone.0145342
589
590 Jalil, A.R., Andrechak, J.C., Discher, D.E., 2020. Macrophage checkpoint blockade: results from
591   initial clinical trials, binding analyses, and CD47-SIRPα structure–function. Antibody
592   Therapeutics 3, 80–94. https://doi.org/10.1093/abt/tbaa006
593
594 Kamber, R.A., Nishiga, Y., Morton, B., Banuelos, A.M., Barkal, A.A., Vences-Catalán, F., Gu,
595   M., Fernandez, D., Seoane, J.A., Yao, D., Liu, K., Lin, S., Spees, K., Curtis, C., Jerby-Arnon,
596   L., Weissman, I.L., Sage, J., Bassik, M.C., 2021. Inter-cellular CRISPR screens reveal
597   regulators of cancer cell phagocytosis. Nature 597, 549–554.
598   https://doi.org/10.1038/s41586-021-03879-4
599
Kitajima, S., Tani, T., Springer, B.F., Campisi, M., Osaki, T., Haratani, K., Chen, M., Knelson,
E.H., Mahadevan, N.R., Ritter, J., Yoshida, R., Köhler, J., Ogino, A., Nozawa, R.-S.,
Sundararaman, S.K., Thai, T.C., Homme, M., Piel, B., Kivlehan, S., Obua, B.N., Purcell, C.,
Yajima, M., Barbie, T.U., Lizotte, P.H., Jänne, P.A., Paweletz, C.P., Gokhale, P.C., Barbie,

D.A., 2022. MPS1 inhibition primes immunogenicity of KRAS-LKB1 mutant lung cancer.
Cancer Cell 40, 1128-1144.e8. https://doi.org/10.1016/j.ccell.2022.08.015
606
Klichinsky, M., Ruella, M., Shestova, O., Lu, X.M., Best, A., Zeeman, M., Schmierer, M.,
Gabrusiewicz, K., Anderson, N.R., Petty, N.E., Cummins, K.D., Shen, F., Shan, X., Veliz,
K., Blouch, K., Yashiro-Ohtani, Y., Kenderian, S.S., Kim, M.Y., O'Connor, R.S., Wallace,
S.R., Kozlowski, M.S., Marchione, D.M., Shestov, M., Garcia, B.A., June, C.H., Gill, S.,
2020. Human chimeric antigen receptor macrophages for cancer immunotherapy. Nat
Biotechnol 38, 947–953. https://doi.org/10.1038/s41587-020-0462-y
613
614   Krysko, D.V., Ravichandran, K.S., Vandenabeele, P., 2018. Macrophages regulate the
615   clearance   of   living   cells   by   calreticulin.   Nat   Commun   9,   4644.
616   https://doi.org/10.1038/s41467-018-06807-9
617
618   Mackenzie, K.J., Carroll, P., Martin, C.-A., Murina, O., Fluteau, A., Simpson, D.J., Olova, N.,
619   Sutcliffe, H., Rainger, J.K., Leitch, A., Osborn, R.T., Wheeler, A.P., Nowotny, M., Gilbert, N.,
620   Chandra, T., Reijns, M.A.M., Jackson, A.P., 2017. cGAS surveillance of micronuclei links
621   genome   instability   to   innate   immunity.   Nature   548,   461–465.
622   https://doi.org/10.1038/nature23449
623
624   Mallin, M.M., Kim, N., Choudhury, M.I., Lee, S.J., An, S.S., Sun, S.X., Konstantopoulos, K.,
625   Pienta, K.J., Amend, S.R., 2022. Cells in the Polyaneuploid Cancer Cell (PACC) state have
626   increased   metastatic   potential   (preprint).   Cancer   Biology.
627   https://doi.org/10.1101/2022.09.16.508155
628
629   Morrissey, M.A., Kern, N., Vale, R.D., 2020. CD47 Ligation Repositions the Inhibitory Receptor
630   SIRPA to Suppress Integrin Activation and Phagocytosis. Immunity 53, 290-302.e6.
631   https://doi.org/10.1016/j.immuni.2020.07.008
632
633   Mujal, A.M., Combes, A.J., Rao, A.A., Binnewies, M., Samad, B., Tsui, J., Boissonnas, A.,
634   Pollack, J.L., Argüello, R.J., Meng, M.V., Porten, S.P., Ruhland, M.K., Barry, K.C., Chan, V.,
635   Krummel, M.F., 2022. Holistic Characterization of Tumor Monocyte-to-Macrophage
636   Differentiation Integrates Distinct Immune Phenotypes in Kidney Cancer. Cancer
637   Immunology Research 10, 403–419. https://doi.org/10.1158/2326-6066.CIR-21-0588
638
639   Nia, H.T., Munn, L.L., Jain, R.K., 2020. Physical traits of cancer. Science 370, eaaz0868.
640   https://doi.org/10.1126/science.aaz0868
641

642   Nimmerjahn, F., Lux, A., Albert, H., Woigk, M., Lehmann, C., Dudziak, D., Smith, P., Ravetch,
643   J.V., 2010. FcγRIV deletion reveals its central role for IgG2a and IgG2b activity in vivo. Proc.
644   Natl. Acad. Sci. U.S.A. 107, 19396–19401. https://doi.org/10.1073/pnas.1014515107
645

646   Noy, R., Pollard, J.W., 2014. Tumor-Associated Macrophages: From Mechanisms to Therapy.
647   Immunity 41, 49–61. https://doi.org/10.1016/j.immuni.2014.06.010
648

649   Oldenborg, P.-A., Zheleznyak, A., Fang, Y.-F., Lagenaur, C.F., Gresham, H.D., Lindberg, F.P.,
650   2000. Role of CD47 as a Marker of Self on Red Blood Cells. Science 288, 2051–2054.
651   https://doi.org/10.1126/science.288.5473.2051
652

653   Perry, J.S.A., Morioka, S., Medina, C.B., Iker Etchegaray, J., Barron, B., Raymond, M.H., Lucas,
654   C.D., Onengut-Gumuscu, S., Delpire, E., Ravichandran, K.S., 2019. Interpreting an
655   apoptotic corpse as anti-inflammatory involves a chloride sensing pathway. Nat Cell Biol 21,
656   1532–1543. https://doi.org/10.1038/s41556-019-0431-1
657

658   Santaguida, S., Richardson, A., Iyer, D.R., M'Saad, O., Zasadil, L., Knouse, K.A., Wong, Y.L.,
659   Rhind, N., Desai, A., Amon, A., 2017. Chromosome Mis-segregation Generates Cell-Cycle-
660   Arrested Cells with Complex Karyotypes that Are Eliminated by the Immune System.
661   Developmental Cell 41, 638-651.e5. https://doi.org/10.1016/j.devcel.2017.05.022
662

Senovilla, L., Vitale, I., Martins, I., Tailler, M., Pailleret, C., Michaud, M., Galluzzi, L., Adjemian,
S., Kepp, O., Niso-Santano, M., Shen, S., Mariño, G., Criollo, A., Boilève, A., Job, B.,
Ladoire, S., Ghiringhelli, F., Sistigu, A., Yamazaki, T., Rello-Varona, S., Locher, C., Poirier-
Colame, V., Talbot, M., Valent, A., Berardinelli, F., Antoccia, A., Ciccosanti, F., Fimia, G.M.,
Piacentini, M., Fueyo, A., Messina, N.L., Li, M., Chan, C.J., Sigl, V., Pourcher, G.,
Ruckenstuhl, C., Carmona-Gutierrez, D., Lazar, V., Penninger, J.M., Madeo, F., López-Otín,
C., Smyth, M.J., Zitvogel, L., Castedo, M., Kroemer, G., 2012. An Immunosurveillance
Mechanism   Controls   Cancer   Cell   Ploidy.   Science   337,   1678–1684.
https://doi.org/10.1126/science.1224922
672

673   Sockolosky, J.T., Dougan, M., Ingram, J.R., Ho, C.C.M., Kauke, M.J., Almo, S.C., Ploegh, H.L.,
674   Garcia, K.C., 2016. Durable antitumor responses to CD47 blockade require adaptive
675   immune   stimulation.   Proc.   Natl.   Acad.   Sci.   U.S.A.   113.
676   https://doi.org/10.1073/pnas.1604268113
677

Spurr, L.F., Martinez, C.A., Kang, W., Chen, M., Zha, Y., Hseu, R., Gutiontov, S.I., Turchan,
W.T., Lynch, C.M., Pointer, K.B., Chang, P., Murgu, S., Husain, A.N., Cody, B., Vokes, E.E.,
Bestvina, C.M., Patel, J.D., Diehn, M., Gajewski, T.F., Weichselbaum, R.R., Chmura, S.J.,
Pitroda, S.P., 2022a. Highly aneuploid non-small cell lung cancer shows enhanced

responsiveness to concurrent radiation and immune checkpoint blockade. Nat Cancer 3,
1498–1512. https://doi.org/10.1038/s43018-022-00467-x
684

685   Spurr, L.F., Weichselbaum, R.R., Pitroda, S.P., 2022b. Tumor aneuploidy predicts survival
686   following immunotherapy across multiple cancers. Nat Genet 54, 1782–1785.
687   https://doi.org/10.1038/s41588-022-01235-4
688

689   Suter, E.C., Schmid, E.M., Harris, A.R., Voets, E., Francica, B., Fletcher, D.A., 2021.
690   Antibody:CD47 ratio regulates macrophage phagocytosis through competitive receptor
691   phosphorylation. Cell Reports 36, 109587. https://doi.org/10.1016/j.celrep.2021.109587
692

693   Taylor, A.M., Shih, J., Ha, G., Gao, G.F., Zhang, X., Berger, A.C., Schumacher, S.E., Wang, C.,
694   Hu, H., Liu, Jianfang, The Cancer Genome Atlas Reseach Network, Lazar, A.J., Cherniack,
695   A.D., Beroukhim, R., Meyerson, M., 2018. Genomic and Functional Approaches to
696   Understanding Cancer Aneuploidy. Cancer Cell 33, 676-689.e3.
697   https://doi.org/10.1016/j.ccell.2018.03.007
698

699   Tripathi, R., Modur, V., Senovilla, L., Kroemer, G., Komurov, K., 2019. Suppression of tumor
700   antigen presentation during aneuploid tumor evolution contributes to immune evasion.
701   OncoImmunology 8, 1657374. https://doi.org/10.1080/2162402X.2019.1657374
702

703   Tsai, R.K., Discher, D.E., 2008. Inhibition of "self" engulfment through deactivation of myosin-II
704   at the phagocytic synapse between human cells. Journal of Cell Biology 180, 989–1003.
705   https://doi.org/10.1083/jcb.200708043
706

707   Vasudevan, A., Baruah, P.S., Smith, J.C., Wang, Z., Sayles, N.M., Andrews, P., Kendall, J., Leu,
708   J., Chunduri, N.K., Levy, D., Wigler, M., Storchová, Z., Sheltzer, J.M., 2020. Single-
709   Chromosomal Gains Can Function as Metastasis Suppressors and Promoters in Colon
710   Cancer. Developmental Cell 52, 413-428.e6. https://doi.org/10.1016/j.devcel.2020.01.034
711

712   Vasudevan, A., Schukken, K.M., Sausville, E.L., Girish, V., Adebambo, O.A., Sheltzer, J.M.,
713   2021. Aneuploidy as a promoter and suppressor of malignant growth. Nat Rev Cancer 21,
714   89–103. https://doi.org/10.1038/s41568-020-00321-1
715

716   Wang, R.W., Viganò, S., Ben-David, U., Amon, A., Santaguida, S., 2021. Aneuploid senescent
717   cells activate NF-κB to promote their immune clearance by NK cells. EMBO Reports 22.
718   https://doi.org/10.15252/embr.202052032
719

Willingham, S.B., Volkmer, J.-P., Gentles, A.J., Sahoo, D., Dalerba, P., Mitra, S.S., Wang, J.,
Contreras-Trujillo, H., Martin, R., Cohen, J.D., Lovelace, P., Scheeren, F.A., Chao, M.P.,

Weiskopf, K., Tang, C., Volkmer, A.K., Naik, T.J., Storm, T.A., Mosley, A.R., Edris, B.,
Schmid, S.M., Sun, C.K., Chua, M.-S., Murillo, O., Rajendran, P., Cha, A.C., Chin, R.K.,
Kim, D., Adorno, M., Raveh, T., Tseng, D., Jaiswal, S., Enger, P.Ø., Steinberg, G.K., Li, G.,
So, S.K., Majeti, R., Harsh, G.R., van de Rijn, M., Teng, N.N.H., Sunwoo, J.B., Alizadeh,
A.A., Clarke, M.F., Weissman, I.L., 2012. The CD47-signal regulatory protein alpha (SIRPa)
interaction is a therapeutic target for human solid tumors. Proc. Natl. Acad. Sci. U.S.A. 109,
6662–6667. https://doi.org/10.1073/pnas.1121623109
729

730   Xian, S., Dosset, M., Almanza, G., Searles, S., Sahani, P., Waller, T.C., Jepsen, K., Carter, H.,
731   Zanetti, M., 2021. The unfolded protein response links tumor aneuploidy to local immune
732   dysregulation. EMBO Rep 22. https://doi.org/10.15252/embr.202152509
733

Zhou, Y., Fei, M., Zhang, G., Liang, W.-C., Lin, W., Wu, Y., Piskol, R., Ridgway, J., McNamara,
E., Huang, H., Zhang, J., Oh, J., Patel, J.M., Jakubiak, D., Lau, J., Blackwood, B., Bravo,
D.D., Shi, Y., Wang, J., Hu, H.-M., Lee, W.P., Jesudason, R., Sangaraju, D., Modrusan, Z.,
Anderson, K.R., Warming, S., Roose-Girma, M., Yan, M., 2020. Blockade of the Phagocytic
Receptor MerTK on Tumor-Associated Macrophages Enhances P2X7R-Dependent STING
Activation by Tumor-Derived cGAMP. Immunity 52, 357-373.e9.
https://doi.org/10.1016/j.immuni.2020.01.014
741
742

Materials and Methods

Cell culture

B16F10 cells (CRL-6475) were obtained from American Type Culture Collection (ATCC) and

cultured at 37°C and 5% CO_2 in either RPMI-1640 (Gibco 11835-030) or Dulbecco's Modified

Eagle Medium (DMEM, Gibco 10569-010) supplemented with 10% fetal bovine serum (FBS,

Sigma F2442), 100 U/ mL penicillin and 100 µg/mL streptomycin (1% P/S, Gibco 15140-122).

B16F10 cells were maintained in passage in RPMI-1640 but switched to DMEM at least three

days prior to *in vivo* subcutaneous injections. 293T human embryonic kidney (CRL-1573) cells

were also obtained from ATCC and cultured at 37°C and 5% CO_2 in DMEM culture media

supplemented with 10% FBS and 1% P/S. All cell lines were passaged every 2-3 days when a

confluency of ~80% was reached. For trypsinization, cells were washed once with Dulbecco's

phosphate-buffered saline (PBS, Gibco 14190-136) and then detached with 0.05% Trypsin

(Gibco 25300-054) for 5 min at 37°C and 5% CO_2. Trypsin was quenched with an equal volume

of complete culture media.

Lentiviral production and transduction

24 h prior to lentivirus production, 293T cells were plated at a density of $8x10^5$ cells in 2 mL of

DMEM in individual wells of a 6-well plate. On the day of transfection, 1.35 µg of psPAX2, 165

ng of pCMV-VSV-G, and 1.5 µg of lentiviral plasmid were added into a microcentrifuge tube with

7.5 µL of Mirus TransIT-Lenti transfection reagent (Mirus Bio, 6604). Once all plasmids were

pooled together with the transfection reagent, 300 µL of serum-free media was added. The

solution was gently mixed by pipetting and then allowed to incubate for 30 min. After 30 min

elapsed, the solution was gently added dropwise by pipetting to an individual 6-well containing

the 293T cells plated the day prior. We note that we also use B16F10 CD47 and Tyrp1 KO lines

in this study, whose preparation has been previously described (Hayes, et al., 2020). LentiV-

cas9_puro and Lenti_sgRNA_EFS_GFP plasmids (Addgene #108100 and 65656, respectively)

were gifts from Christopher Vakoc. psPAX2 was a gift from Didier Trono (Addgene plasmid

#12260). pVSV-G was a gift from Bob Weinberg (Addgene plasmid #8454). The single guide

RNA (sgRNA) oligonucleotides (CD47, 5′-TCCCCGTAGAGATTACAATG-3′; SIRPα, 5′-

TAATTCTAAGGTCATCTGCG-3′) were designed using the Broad CRISPR algorithm. sgRNAs

were cloned into the sgRNA vector using a BsmBI restriction digest.

Viral production was allowed to continue for 48 h, after which the supernatant was collected from

each well and centrifuged for 5 min at 300 x *g*. The supernatant was collected and then either

added directly to B16F10 cells (0.5 mL of lentivirus-containing supernatant to $2x10^4$ cells in an

individual 6-well) or stored at -70°C for long-term storage. 72 h after transduction, spent media

with lentivirus was aspirated, B16F10 cells were washed with PBS, and fresh media was added.

For selecting successfully lentivirally transduced cells, cells were cultured in fresh media

containing 1 μg/mL of puromycin (Invitrogen A1113803). Cells were kept in puromycin-

containing until a non-transduced control population also treated with puromycin completely

died.

Antibodies

Antibodies used for *in vivo* treatment and blocking and for *in vitro* phagocytosis are as follows:

anti-mouse/human Tyrp1 clone TA99 (BioXCell BE0151), mouse IgG2a isotype control clone

C1.18.4 (BioXCell BE0085), and Ultra-LEAF anti-mouse CD172a (SIRPα) clone P84

(BioLegend 144036). Low-endotoxin and preservative-free antibody preparations were used for

in vivo treatments and *in vitro* phagocytosis experiments. For primary antibody staining of

surface proteins via flow cytometry, the following were used: anti-mouse CD47 clone MIAP301

(BioXCell BE0270) and anti-mouse/human Tyrp1 clone TA99. Secondary antibodies used for

flow cytometry are as follows: Alexa Fluor 647 donkey anti-mouse IgG (ThermoFisher A-31571)

and Alexa Fluor 647 goat anti-rat IgG (ThermoFisher A-21247). All secondary antibody

concentrations used followed the manufacturer's recommendations.

For immune infiltrate analysis, the following BioLegend antibodies were used: Brilliant Violet 650

anti-mouse CD45 clone 30-F11 (103151), Brilliant Violet 785 anti-mouse CD45 clone 30-F11

(103149), APC/Cy7 anti-mouse CD45 clone 30-F11 (103115), APC anti-mouse/human CD11b

clone M1/70 (101212),   PE/Cyanine7 anti-mouse/human CD11b clone M1/70 (101216),

PE/Dazzle 594 anti-mouse Ly6G clone 1A8 (127647), PerCP anti-mouse Ly-6G clone 1A8

(127653), PE anti-mouse F4/80 clone BM8 (123110), Brilliant Violet 605 anti-mouse Ly-6C clone

HK1.4 (128035), APC anti-mouse I-A/I-E clone M5/114.15.2 (107614), Pacific Blue anti-mouse

CD86 clone GL-1 (105022), APC/Cy7 anti-mouse CD86 clone GL-1 (105029), APC anti-mouse

CD206 (MMR) clone C068C2 (141707), Brilliant Violet 421 anti-mouse CD206 clone C068C2

(141717), PE/Cy7 anti-mouse CD163 clone S15049F (156707), APC/Cy7 anti-mouse CD3e

clone 145-2C11 (100329), and Alexa Fluor 647 anti-mouse CD8a clone 53-6.7 (100727).

TruStain FcX PLUS (anti-mouse CD16/32) clone S17011E (156603) was used in all immune

infiltrate experiments to block Fc receptors. For immunogenicity post-MPS1i treatment, APC

anti-mouse H-2kb/H-2Db clone 28-8-6 (114613) was used. For IgG titer in tumor challenge

surviving mice, the following BioLegend antibodies were used: PE anti-mouse IgG2a clone

RMG2a-62 (407108, known to bind IgG2c as well) and APC anti-mouse IgG2b clone RMG2b-1

(406711). Primary antibody used in Western blotting was anti-β-actin clone C4 (Santa Cruz sc

47778). Secondary antibody used in Western blotting was HRP sheep anti-mouse IgG (GE Life

Sciences NA931V).

Mice

C57BL/6 mice (Jackson Laboratory 000664) were 6-12 weeks old at the time of tumor challenges

and for bone marrow harvesting, with the exception of second and third challenge experiments.

Additionally, for re-challenge experiments, age-matched naïve mice were used. All experiments

were performed in accordance with protocols approved by the Institutional Animal Care and Use

Committee (IACUC) of the University of Pennsylvania.

Drug treatments & micronuclei quantification

For MPS1i studies, the following chemical drugs were used: reversine (Cayman Chemical

10004412), AZ3146 (Cayman Chemical 19991), and BAY 12-17389 (Selleck Chemicals S8215).

24 h prior to treatment, B16F10 cells were in either 6-well or 12-well plates. For 6-wells, 20,000

cells were plated per well. For 12-wells, 2,000 cells were plated per well. On the day of treatment,

spent media was aspirated, and fresh media supplemented with 10% FBS, 1% P/S, and either

MPS1i or DMSO vehicle control was added to each well. The concentration used for each

treatment is listed in the Figure legend. The volume of DMSO added was equal to the volume

required for the highest MPS1i concentration for each experiment. All cells were treated for 24

h, after which drug-containing spent media was aspirated. Cells were then washed with a full

volume of PBS for 5 min. PBS was aspirated, and two repeat washes were performed. Cells

were then allowed to recover from MPS1i treatment for an additional 48 h, with a fresh media

replacement 24 h after the initial wash.

For imaging and micronuclei quantification, cells were fixed with 4% formaldehyde for 20 min

after the 48 h recovery period and later imaged on an Olympus IX inverted microscope with a

40x/0.6 NA objective. The Olympus IX microscope was equipped with a Prime sCMOS camera

(Photometrics) and a pE-300 LED illuminator (CoolLED) and was controlled with MicroManager

software v2. At least 200 B16F10 were imaged per individual well for micronuclei quantification.

Bone marrow-derived macrophages (BMDMs)

Bone marrow was harvested from the femurs and tibia of donor mice, lysed with ACK buffer

(Gibco A1049201) to deplete red blood cells, and then cultured on Petri culture dishes for 7 days

in Iscove's Modified Dulbecco's Medium (IMDM, Gibco 12440-053) supplemented with 10%

FBS, 1% P/S, and 20 ng/mL recombinant mouse macrophage colony-stimulating factor (M-CSF,

BioLegend 576406). 72 h after initial plating, one whole volume of fresh IMDM supplemented

10% FBS, 1% P/S, and 20 ng/mL M-CSF. After 7 days of differentiation, spent media was

removed, BMDMs were gently washed once with phosphate-buffered saline (PBS), and fresh

IMDM supplemented with 10% FBS, 1% P/S, and 20 ng/mL M-CSF was added.

Conditioned media treatment of BMDMs

BMDMs that had successfully undergone 7 days of differentiation in 20 ng/mL M-CSF were used.

Spent media was removed, BMDMs were gently washed once with phosphate-buffered saline

(PBS), and fresh IMDM supplemented 10% FBS, 1% P/S, and 20 ng/mL M-CSF was added.

Then, conditioned media from B16F10 cells treated with either MPS1i or DMSO was collected.

Conditioned media was centrifuged for 5 min at 300 x g to remove any cellular debris. The

supernatant was collected and then supplemented to 5% FBS, 1% P/S, and 20 ng/mL M-CSF.

One whole volume of conditioned media was added to BMDMs with fresh media. 24 h after

treatment, BMDMs were detached using 0.05% Trypsin and processed for single-cell RNA-

sequencing.

***In vitro* phagocytosis**

For two-dimensional phagocytosis assays, BMDMs were detached using 0.05% Trypsin and re-

plated in either 6-well or 12-well plates, at a density of 1.8×10^4 cells per cm^2 in IMDM

supplemented 10% FBS, 1% P/S, and 20 ng/mL M-CSF. After 24 h elapsed, BMDMs were

labeled with 0.5 µM CellTracker DeepRed dye (Invitrogen C34565), according to the

manufacturer's protocol. Following staining, BMDMs were washed and incubated in serum-free

IMDM supplemented 0.1% (w/v) BSA and 1% P/S. B16F10 cells were labeled with

carboxyfluorescein diacetate succinimidyl ester (Vybrant CFDA-SE Cell Tracer, Invitrogen

V12883), also according to the manufacturer's protocol. B16F10 cells were detached and

opsonized with 10 µg/mL anti-Tyrp1, with 10 µg/mL mouse IgG2a isotype control antibody, or

5% (v/v) mouse serum in 1% BSA. Opsonization was allowed for 30-45 min on ice. Opsonized

B16F10 suspensions were then added to BMDMs at a ~2:1 ratio and incubated at 37°C and 5%

CO_2 for 2 h. Non-adherent cells were removed by gently washing with PBS. For imaging, cells

were fixed with 4% formaldehyde for 20 min and later imaged on an Olympus IX inverted

microscope with a 40x/0.6 NA objective. The Olympus IX microscope was equipped with a Prime

sCMOS camera (Photometrics) and a pE-300 LED illuminator (CoolLED) and was controlled

with MicroManager software v2. At least 300 macrophages were imaged per individual well for

calculation of phagocytosis.

3D tumoroid formation and phagocytosis

Briefly, non-TC-treated 96-well U-bottom plates were treated with 100 µL of anti-adherence

rinsing solution (StemCell Technologies 07010) for 1 h. The cells were then washed with 100 µL

of complete RPMI 1640 cell culture media. This generated surfaces conducive to generating

tumoroids and preventing cells from adhering to the well bottom during experiments. B16F10

were detached by brief trypsinization, resuspended at a concentration of 1×10^4 cells per mL in

complete RPMI 1640 cell culture media (10% FBS, 1% P/S) with 50 µM β-mercaptoethanol

(Gibco 21985023). 100 µL of this cell suspension was added to each well such that each

tumoroid initially started with approximately 1×10^3 cells. Aggregation of B16F10 cells was

confirmed 24 h later by inspection under microcopy. Upon confirmation of tumoroid formation,

BMDMs were labeled with 0.5 µM CellTracker DeepRed dye (Invitrogen C34565), according to

the manufacturer's protocol. BMDMs were then detached by brief trypsinization and gentle

scraping and resuspended in complete RPMI 1640 cell culture media at a concentration that

would allow for delivery of $3x10^3$ BMDMs to each individual tumoroid culture. The BMDM cell

suspension was also supplemented with 120 ng/mL M-CSF and antibodies (either anti-Tyrp1 or

mouse IgG2a isotype control) such that delivery of 20 μL of this suspension to each individual

tumoroid culture result in final concentrations of 20 ng/mL M-CSF and 20 μg/mL of antibody. For

tumoroid studies in which mouse convalescent serum was used, the BMDM cell suspension was

supplemented with 120 ng/mL M-CSF and serum such that delivery of 20 μL of this suspension

to each individual tumoroid culture resulted in final concentrations of 20 ng/mL M-CSF and a

final mouse serum concentration of 1:200. Tumoroids were imaged on an Olympus IX inverted

microscope with a 4x/0.6 NA objective.

For macrophage polarization tumoroid experiments (Fig. 2B-C), B16F10 tumoroids were

prepared in the same manner as described above, with the exception that no anti-Tyrp1 or

mIgG2a was added. After 5 days had elapsed for the experiment, tumoroids were dissociated

by brief trypsinization and stained with conjugated antibodies targeting MHCII, CD86, CD163,

and CD206, following the manufacturer's protocol. Cells were then run on a BD LSRII (Benton

Dickinson) flow cytometer. Data were analyzed with FCS Express 7 software (De Novo

Software).

***In vivo* tumor models**

B16F10 cells cultured in DMEM growth media were detached by brief trypsinization, washed

twice with PBS, and resuspended at $2x10^6$ cells per mL. Cell suspensions were kept on ice until

injection. All subcutaneous injections were performed on the right flank while mice were

anesthetized under isoflurane. Fur on the injection site was wet slightly with a drop of 70%

ethanol and brushed aside to better visualize the skin. A 100 μL bolus containing $2x10^5$ cells

cancer cells was injected beneath the skin. For treatments, mice received either intravenous or

intraperitoneal injections of anti-Tyrp1 clone TA99 or mouse IgG2a isotype control clone C1.18.4

(250 μg antibody in 100 μL PBS) on days 4, 5, 7, 9, 11, 13, and 15 post-tumor challenge.

Intravenous injections were done via the lateral tail vein. Tumors were monitored by palpation

and measured with digital calipers. The projected area was roughly elliptical and was calculated

as $A = \pi/4 \times L \times W$, where L is the length along the longest axis and W is the width measured

along the perpendicular axis. For our studies, a projected area of 125 mm^2 was considered

terminal burden for survival analyses. Mice were humanely euthanized following IACUC

protocols if tumor size reached 2.0 cm on either axis, if tumor reached a projected area greater

than 200 mm^2 or if a tumor was ulcerated.

Adoptive cell transfers

Fresh bone marrow was harvested as described in the **Bone marrow-derived macrophages**

(BMDMs) methods section. Marrow cells were counted on a hemocytometer and resuspended

to a concentration of 8x10^7 cells per mL in in 5% (v/v) FBS/PBS. To block SIRPα, cells were

then incubated with anti-SIRPα clone P84 (18 µg/mL) for 1 h at room temperature on a rotator.

After the incubation period elapsed, cells were centrifuged at 300 x g for 5 min, washed with

PBS, and then centrifuged once more at 300 x g for 5 min to remove any unbound anti-SIRPα.

Cells were again re-suspended to a concentration of 8x10^7 cells per mL in in 2% (v/v) FBS/PBS,

with or without anti-Tyrp1 clone TA99 (1 mg/mL). Marrow cells were then injected intravenously

(2x10^7 cells in 250 µL per mouse) into tumor-bearing mice. All adoptive transfers were done four

days post-challenge. Control data were adapted from (Dooling et al., 2022) to minimize mice for

experiments, since the cited study establishes proper benchmarks for comparison and also finds

that these control conditions minimally (if at all) improve survival.

Serum collection & IgG titer quantification

Blood was drawn retro-orbitally from mice anesthetized under isoflurane, using heparin- or

EDTA-coated microcapillary tubes. Collected blood was allowed to clot for 1 h at room

temperature in a microcentrifuge tube. The serum was separated from the clot by centrifugation

at 1,500 x g and stored at -20°C for use in flow cytometry and phagocytosis assays.

For IgG titer quantification, B16F10 cells were detached by trypsinization and incubated with 5%

(v/v) mouse serum in 1% BSA. Opsonization was allowed for 30-45 min on ice. After the

incubation period elapsed, cells were centrifuged at 300 x g for 5 min, washed once with PBS,

centrifuged again at 300 x g for 5 min, and then resuspended in 0.1% (w/v) BSA with both PE

anti-mouse IgG2a/c clone RMG2a-62 and APC anti-mouse IgG2b clone RMG2b-1 (see

Antibodies section for more information). Anti-IgG conjugated-antibody incubation occurred for

30-45 min, after which cells were centrifuged at 300 x *g* for 5 min, washed once with PBS,

centrifuged again at 300 x *g* for 5 min, and then resuspended in 5% (v/v) FBS/PBS. Cells were

run on a BD LSRII (Benton Dickinson) flow cytometer.

Western blotting

Lysate was prepared from B16F10 cells using RIPA buffer containing 1X protease inhibitor

cocktail (Millipore Sigma P8340) and boiled in 1X NuPage LDS sample buffer (Invitrogen

NP0007) with 2.5% β-mercaptoethanol. Proteins were separated by electrophoresis in NuPage

4-12% Bis-tris gels run with 1X MOPS buffer (Invitrogen NP0323) and transferred to an iBlot

nitrocellulose membrane (Invitrogen IB301002). The membranes were blocked with 5% (w/v)

non-fat milk in Tris buffered saline (TBS) plus Tween-20 (TBST) for 1 h and stained with 5%

(v/v) mouse serum overnight at 4°C with agitation. The membranes were washed TBST and

incubated with 1:500 secondary antibody conjugated with horseradish peroxidase (GRP) in 5%

(w/v) milk in TBST for 1 h at room temperature with agitation. The membranes were then washed

again three times with TBST. Membranes probed with HRP-conjugated secondary antibody

were developed a 3,3',5,5'-teramethylbenzidine (TMB) substrate (Genscript L002V or Millipore

Sigma T0565). Developed membranes were scanned and then processed with ImageJ.

Immune infiltrate analysis of tumors

For day 5 post-challenge measurements: If mice required anti-Tyrp1 treatment, mice received a

single dose of intravenously delivered anti-Tyrp1 or mouse IgG2a isotype control four days (96

h) post-tumor challenge. 24 later, mice were humanely sacrificed. Otherwise, mice were

sacrificed five days post-tumor challenge. For immune analysis of second challenge non-

survivors: Mice were humanely euthanized when tumor burden reached >150 mm^2.

Tumors from euthanized mice were excised and placed into 5% (v/v) FBS/PBS. Tumors were

then disaggregated using Dispase (Corning 354235) supplemented with 4 mg/mL of collagenase

type IV (Gibco 17104-019) and DNAse I (Millipore Sigma, 101041159001) for 30-45 min (until

noticeable disaggregation) at 37°C, centrifuged for 5 min at 300 x *g*, and resuspended in 1 mL

of ACK lysis buffer for 12 min at room temperature. Samples were centrifuged for 5 min at 300

x *g*, washed once with PBS, and then resuspended in 5% (w/v) BSA/PBS for 20 min. After 20

min elapsed, fluorophore-conjugated antibodies to immune markers were added to each cell

suspension. The following markers were used for analysis: for macrophages, CD45, CD11b,

F4/80, Ly-6C, Ly-6G, CD86, CD206, MHCII; for T cells, CD45, CD3e, CD8a. Antibody binding

occurred for 30 mins while samples were kept on ice and covered from light. Samples were then

centrifuged for 5 min at 300 x *g,* washed once with PBS, and resuspended in FluoroFix Buffer

(BioLegend, 422101) for 1 h at room temperature prior to analysis on a BD LSRII (Benton

Dickinson) flow cytometer. Data were analyzed with FCS Express 7 software (De Novo

Software).

Bulk RNA-sequencing

RNA library was constructed using NEBNext® Ultra™ II RNA Library Prep Kit for Illumina® and

NEBNext® Multiplex Oligos for Illumina (E7770S, E7335) per the manufacturer's instructions.

The library prepared was processed at the Next Generation Sequencing Core at the University

of Pennsylvania (12-160, Translational Research Center) using NovaSeq 6000, 100 cycles

(Illumina). The reads were aligned to mouse reference, mm10 (GENCODE vM23/Ensembl 98)

using star alignment. Cell count matrix was generated and imported to RStudio for downstream

analysis. Package "DESeq2" (v1.32.0) was used for normalization and differential expression

analysis. Package "fgsea" (v1.18.0) was used for gene set enrichment analysis. Additional

exploratory data analysis was then done using either RStudio or Python 3.8.

1009

Single-cell RNA-sequencing

RNA libraries were prepared using the 10X Genomics Chromium Single Cell Gene Expression

kit (v3.1, single index, PN-1000128; PN-1000127; PN-1000213) per the manufacturer's

instructions. The libraries were sequenced at the Next Generation Sequencing Core using

NovaSeq 6000, 100 cycles (Illumina). Raw base call (BCL) files were analyzed using CellRanger

(version 5.0.1) to generate FASTQ files, and the "count" command was used to generate raw

count matrices aligned to mm10 (GENCODE vM23/Ensembl 98). Cells were filtered to make

sure that they expressed a minimum of 1,400 genes with less than 15 percent mitochondrial

content. Data was normalized using the "LogNormalize" method from the Seurat package.

Differential expression analysis was performed using the "FindAllMarkers" command, and the

output was used for gene set enrichment analysis (GSEA).

1021

InferCNV

Count matrix of single cell RNA-sequencing results was used as input for InferCNV object

construction (1.7.1) (InferCNV of the Trinity CTAT Project, see the following for more information:

https://github.com/broadinstitute/inferCNV). Gene position files were created for GRCm38.

Single-cell RNA-sequencing data of DMSO-treated B16F10 were used as reference for copy

number profile construction. Cell types were annotated either manually or using the package

"SingleR" (v1.6.1). For manual annotation, cells were clustered and assigned cell types based

on the expression of cell type specific signature genes (SI.5a). Denoised results from InferCNV

were used as the input ("infercnv.observations.txt"). The averaged copy number of each

chromosome segment was calculated, and the difference between each cell's copy number and

the overall mean at each segment was calculated. The deviation was summed across the entire

chromosome to obtain the distribution of the deviation. Cells sharing an absolute deviation that

is more than 2.5 times standard deviation away from the distribution peak were marked as

outliers for a certain chromosome in question.

1036

Transcriptomic gene sets

For analyzing both bulk and single-cell RNA-sequencing data sets, either hallmark gene sets

(from the Human Molecular Signatures Database) or customized gene sets were used.

Customized gene sets were used exclusively for macrophage-associated analyses and were

made by combining gene sets from the following: Ahn et al., 2018; Cerezo-Wallis et al., 2020;

Cunha et al., 2018; Mujal et al., 2022; Perry et al., 2019; Zhou et al., 2020.

1043

Statistical analysis & curve fitting

Statistical analyses and curve fitting were performed in GraphPad Prism 9.4. Details for each

analysis are provided in the figure legends. Tumor and tumoroid growth curve data (projected

area vs time) were fit to the exponential growth model ($A = A_0 e^{kt}$ for tumors and $A = A_1 e^{k(t-1)}$ for

tumoroids) using non-linear least squares regression with prefactors A_0 or A_1 and k, the

exponential growth rate. For differential gene expression analysis for both bulk RNA-sequencing

and single-cell RNA-sequencing datasets, statistical analyses were done using either RStudio

2022.02.3+492 or Python 3.8.

1052

Code availability

1054 Sequencing data were analyzed using RStudio 2022.02.3+492 and the Seurat package.

1055 Additional analyses after normalization were done on either RStudio or Python 3.8. No new code

1056 central to the conclusions of this study was developed.

1057

Author contributions

1059 Conceptualization: BHH. Formal analysis: BHH, MW, HZ, SP, AC. Funding acquisition: BHH,

1060 JCA, LJD, DED. Investigation & data collection: BHH, MW, HZ, SP, AC, JCA, LJD, MPT, TM.

1061 Methodology: BHH, MW. Resources: DED. Visualization: BHH, MW. Writing: BHH, DED.

1062

Acknowledgements

1064 This work was supported by funding from the following sources: NIH R01 HL124106 (DED), U54

1065 CA193417 (DED), NSF GRFP DGE-1845298 (BHH, JCA, MPT), and NIH F32 CA228285 (LJD).

1066 The authors acknowledge the following University of Pennsylvania core facilities: Cell Center

1067 Stockroom, the Penn Cytomics and Cell Sorting Resource Laboratory, the Penn Genomic and

1068 Sequencing Core, and the Cell & Development Biology Microscopy Core.

1069

Competing interests

1071 The authors declare no competing interests.

1072

Data availability statement

1074 All data are available within the article and its supplementary information. Data can be provided

1075 upon reasonable request from the corresponding author.

1076

1077   **Main Figure Legends**

1078

Figure 1. MPS1i-induced chromosomal instability causes proliferation deficits in B16F10

tumors and generates microenvironment conditions that favor anti-cancer M1-like

macrophages while minimizing skewing to a pro-cancer M2-like phenotype.

(A) Timeline and schematic for treatment of B16F10 mouse melanoma cells with MPS1 inhibitors

(MPS1i). B16F10 cells were treated with MPS1i (reversine) or the equivalent volume of DMSO

vehicle control. Cells were treated for 24 h, after which they were washed twice with PBS and

allowed to recover for an additional 48 h. After the recovery period elapsed, numerous follow-up

experiments were conducted to characterize chromosomal instability, aneuploidy, and effects

on BMDMs (BMDMs).

1088

(B) (i) Representative DNA images of B16F10 cells 72 h after initial treatment, as outlined in (A).

Scale bars = 10 μm. Yellow arrowheads point at micronuclei, which are used as a signature

readout of chromosomal instability. **(ii)** Quantification of the percentage of B16F10 cells with

micronuclei after MPS1i treatment, across a range of different MPS1i concentrations and DMSO

vehicle control (mean ± SEM shown, n = 3 replicates per condition). Statistical significance was

calculated by ordinary one-way ANOVA and Tukey's multiple comparison test (* $p < 0.05$; ** $p <$

0.01; *** $p < 0.001$; **** $p < 0.0001$).

1096

(C) Inferred copy number in DMSO and MPS1i-treated B16F10 cells from single-cell RNA-

sequencing and the InferCNV pipeline. Cells that are considered aneuploid (labeled as outliers

in the InferCNV algorithm) show full-level chromosome gains and/or losses compared to the

consensus copy number profile of the DMSO-treated B16F10 population. Approximately 34% of

MPS1i-treated B16F10 show aneuploidy, as determined by InferCNV, compared to 10% in the

DMSO-treated B16F10 population. Single-cell RNA-sequencing confirms that MPS1i induces

chromosomal instability and aneuploidy in B16F10 cells.

1104

(D) Timeline and schematic for whole-tumor bulk RNA-sequencing. Prior to tumor inoculation,

B16F10 CD47 KO cells were treated with 2.5 μM MPS1i (reversine) or the equivalent volume of

DMSO vehicle control. Cells were treated for 24 h, after which they were washed twice with PBS

and allowed to recover for an additional 48 h. After the recovery period elapsed, all mice were

subcutaneously injected with $5x10^5$ B16F10 cells. Five or ten days after tumor inoculation, mice

were humanely euthanized, and their tumors were excised and disaggregated for bulk RNA-

sequencing analysis.

1112

(E) Heatmap of selected RNA transcripts related to M2 macrophage polarization and pro-tumor

function that were differentially expressed in tumors comprised of MPS1i-treated B16F10

compared to tumors comprised of their DMSO-treated counterpart. Heatmap shows $\log_2$-

transformed transcript reads which were then z-score normalized. Generally, M2 and pro-tumor

macrophage markers are downregulated at day 5 and even further at day 10 compared to DMSO

controls. Tumors analyzed were two comprised of DMSO-treated B16F10 cells five days post-

inoculation, two comprised of MPS1i-treated B16F10 cells five days post-inoculation, and two

comprised of MPS1i-treated B16F10 cells ten days post-inoculation.

1121

(F) Top ten downregulated hallmark gene sets in tumors comprised of MPS1i-treated B16F10

cells (here, we consider Myc targets v1 and Myc targets v2 as one overall group). Cells from

tumors made of MPS1i-treated B16F10 show downregulated transcriptional signatures of cell

cycle and proliferation-related pathways, suggesting overall deficits in proliferation. Gene set

enrichment analysis pathways were obtained from MSigDB. Abbreviations used: CL

homeostasis = cholesterol homeostasis; OXPHOS = oxidative phosphorylation.

1128

1129   **(G)** Schema for conditioned media (secretome) treatment of BMDMs and subsequent

1130   characterization by single-cell RNA-sequencing. Media from DMSO and MPS1i-treated B16F10

1131   cells was collected and added at a 1:1 ratio with fresh media to 7-day differentiated BMDMs

1132   (with 20 ng/mL M-CSF). Treatment was allowed for 24 h, after which BMDMs were collected

1133   and processed for single-cell RNA-sequencing.

1134

1135   **(H) (Left)** UMAP plots of expression profiles for all analyzed BMDMs, treated with the secretome

1136   from DMSO-treated (gray) or MPS1i-treated (red) B16F10 cells. Each circle represents an

1137   individual cell. **(Right)** Same UMAP plots but colors now represent cells clustered together

1138   based on similarity of global gene expression. **(Inset)** Composition of each cluster.

1139

(I) Dot plot showing proportion of cells in each cluster expressing anti-inflammatory and M2-like
polarization-associated genes (pro-cancer gene set overall). Indicated to the right are a heatmap
scale for average gene expression and a dot size reference for what proportion of cells in the
cluster express a gene. Clusters 1 and 3, both of which consist of ~75% of BMDMs treated with
conditioned media from DMSO-treated B16F10, generally show downregulated transcript levels
for genes associated with skewing macrophages to a pro-inflammatory, anti-cancer, M1-like
phenotype. Cluster 3, in particular, shows significant downregulated. On the contrary, clusters 0
and 2, which consist of 75% of BMDMs treated with conditioned media from MPS1i-treated
B16F10, show upregulated expression of many of these anti-cancer genes. Cluster 2, in
particular, shows signatures of high upregulation.

1150

(**J**) Dot plot showing proportion of cells in each cluster expressing pro-inflammatory and M1-like
polarization-associated genes (anti-cancer gene set overall). Indicated to the right are a heatmap
scale for average gene expression and a dot size reference for what proportion of cells in the
cluster express a gene. Clusters 1 and 3, both of which consist of ~75% of BMDMs treated with
conditioned media from DMSO-treated B16F10, generally show upregulated transcript levels for
genes associated with skewing macrophages to an anti-inflammatory, pro-cancer, M2-like
phenotype. On the contrary, clusters 0 and 2, which consist of 75% of BMDMs treated with
conditioned media from MPS1i-treated B16F10, show either little-to-no expression or
downregulation of many of these genes.

Figure 2. Chromosomally unstable tumoroids and tumors in early stages show increased

anti-cancer macrophage polarization and activity.

(A) Timeline and schematic for generating engineered "immuno-tumoroids" for time-lapsed

studies of macrophage-mediated phagocytosis of cancer cells. Tumoroids are formed by plating

and culturing B16F10 cells on non-adhesive surfaces in U-bottom shaped wells. Prior to plating

for tumoroid formation, B16F10 were treated with either DMSO or MPS1i (reversine) as depicted

in Fig. 1A. ~24 h after plating, bone marrow-derived macrophages (BMDMs) with or without

opsonizing anti-Tyrp1 are added to the cohesive B16F10 tumoroids. Immuno-tumoroids are

imaged at the listed timepoints. At day 5, a separate subset of prepared immuno-tumoroids were

collected, dissociated, and stained for measurement of macrophage polarization markers by flow

cytometry.

1171

1172 **(B)** Representative flow cytometry histograms for flow cytometry measurements of macrophage

1173 polarization markers. For M1-like (anti-tumor) markers, we chose MHCII and CD86. For M2-like

1174 (pro-tumor) markers, we chose CD163 and CD206.

1175

(C) Geometric mean fluorescence intensity quantification by flow cytometry of anti-tumor

macrophage markers (MHCII and CD86) and pro-tumor macrophage markers (CD163 and

CD206) in immuno-tumoroid cultures, with either DMSO or MPS1i-treated B16F10 CD47 KO

cells. Generally, immuno-tumoroid cultures with MPS1i-treated B16F10 CD47 KO cells induce

surface expression of M1-like, anti-tumor markers in macrophage while also reducing expression

of M2-like, pro-tumoral markers. This suggests that, in addition to reduced cancer cell

proliferation, chromosomal instability in cancer cells in early stages can prime a

microenvironment conducive to anti-tumor macrophages. All data were collected from five

independent immune-tumoroid culture experiments (one 96-well per replicate). Statistical

significance was calculated by an unpaired two-tailed t-test with Welch's correction (** $p < 0.01$).

1186

1187 **(D)** Representative fluorescence images depicting either growth or repression of B16F10 CD47

1188 KO cells (green) in immuno-tumoroids from days 1-5. BMDMs, shown in magenta, were added

1189 at a 3:1 ratio to initial B16F10 numbers after the day 1 images were acquired. Scale bar = 0.5

1190 mm.

1191

1192   **(E)** Tumoroid growth was measured by calculating the GFP+ area at the indicated timepoints

1193   (mean ± SD, n = 16 total tumoroids from two independent experiments for each condition). All

1194   data were then normalized to average GFP+ area on day 1 of each drug treatment's respective

1195   mouse IgG2a isotype control condition. Overall, macrophages can clear MPS1i-treated B16F10

1196   cells regardless of either MPS1i treatment concentration or IgG opsonization.

1197

(F) Experimental timeline for immune cell infiltration analyses in tumors comprised of either

DMSO or MPS1i-treated B16F10 CD47 KO cells. Prior to tumor inoculation, B16F10 CD47 KO

cells were treated with 2.5 μM MPS1i (reversine) or the equivalent volume of DMSO vehicle

control. Cells were treated for 24 h, after which they were washed twice with PBS and allowed

to recover for an additional 48 h. After the recovery period elapsed, all C57BL/6 mice were

subcutaneously injected with $2x10^5$ B16F10 cells. Four days (96 h) post-challenge, mice were

treated with 250 μg of anti-Tyrp1 or mouse IgG2a isotype. Mice were then euthanized 24 h after

antibody treatment, and their tumors were excised and disaggregated for immune infiltrate

analysis by flow cytometry.

1207

(G) Immune cell infiltrate measurements of B16F10 CD47 KO tumors comprised of DMSO or

MPS1i-treated cells. **(i)** Quantification of the percentage of CD45+ (immune) cells in the excised

tumors, showing that tumors comprised of MPS1i-treated cells show ~3-fold increased tumor

immune cell infiltrate compared to their DMSO counterparts. **(ii)** Quantification of the percentage

of tumor infiltrating myeloid cells in the excised tumors. B16F10 CD47 KO tumors comprised of

MPS1i-treated cells show ~2.5-fold more myeloid cells compared to their DMSO counterparts.

(iii) Quantification of tumor infiltrating F4/80+ macrophages relative to the total number of tumor

cells, with the tumors comprised of MPS1i-treated cells showing a ~6-fold increase compared to

their DMSO counterparts. **(iv)** Quantification of the number of M1-like macrophages

($CD86^{hi}CD206^{lo}$) and M2-like macrophages ($CD86^{lo}CD206^{hi}$) relative to the total number of

macrophages in each tumor condition. Tumors comprised of MPS1i-treated cells have ~6-fold

more $CD86^{hi}CD206^{lo}$ compared to their DMSO counterparts. They also have a similar increase

in $CD86^{lo}CD206^{hi}$ macrophages, but the number of $CD86^{hi}CD206^{lo}$ macrophages was still overall

twice as high. Mean ± SEM shown (n = 8 mice challenged with DMSO-treated B16F10 cells, n

= 5 mice challenged with MPS1i-treated B16F10 cells). Statistical significance was calculated

by an unpaired two-tailed t-test with Welch's correction (ns, not significant; ** $p < 0.01$; **** $p <$

0.0001).

1225

1226

1227

1228

Figure 3. MPS1i-induced chromosomally unstable cancer cells are maximally cleared when IgG-opsonized and depleted of the CD47 macrophage checkpoint.

(**A**) Timeline for *in vitro* treatment B16F10 cells prior to injection in mice and then subsequent therapeutic treatment for tumor-challenged mice. Prior to tumor inoculation, B16F10 CD47 KO cells were treated with 2.5 µM MPS1i (reversine) or the equivalent volume of DMSO vehicle control. Cells were treated for 24 h, after which they were washed twice with PBS and allowed to recover for an additional 48 h. After the recovery period elapsed, all mice were subcutaneously injected with 2×10^5 B16F10 cells. For anti-Tyrp1 and mouse IgG2a isotype control treatments, mice were treated intravenously or intraperitoneally with 250 µg with antibody on days 4, 5, 7, 9, 11, 13, and 15 after tumor challenge.

(**B**) Tumor growth curve of projected tumor area versus days after tumor challenge, with B16F10 sgCtrl cells (expressing WT levels of CD47). Each line represents a separate tumor and is fit with an exponential growth equation: $A = A_0 e^{kt}$. Experimental conditions are as follows: n = 9 mice that were challenged with DMSO-treated B16F10 sgCtrl and subsequently treated with mouse IgG2a control, n = 11 mice that were challenged with DMSO-treated B16F10 sgCtrl and subsequently treated with anti-Tyrp1, n = 6 mice that were challenge with MPS1i-treated B16F10 sgCtrl and subsequently treated with mouse IgG2a control, and n = 7 mice that were challenge with MPS1i-treated B16F10 sgCtrl and subsequently treated with anti-Tyrp1. All data were collected across three independent experiments. Inset bar graphs depict response type for each indicated tumor challenge. A partial response was defined as a mouse that survived at least one week (20+ days) beyond the median survival of the B16F10 sgCtrl cohort treated with mouse IgG2a isotype control (13 days).

(**C**) Tumor growth curve of projected tumor area versus days after tumor challenge, with B16F10 CD47 KO cells. Each line represents a separate tumor and is fit with an exponential growth equation: $A = A_0 e^{kt}$. Complete anti-tumor responses in which a tumor never grew are depicted with the same symbol as their growing counterparts and with solid lines at A = 0. Experimental conditions are as follows: n = 7 mice that were challenged with DMSO-treated B16F10 CD47 KO and subsequently treated with mouse IgG2a control, n = 9 mice that were challenged with DMSO-treated B16F10 CD47 KO and subsequently treated with anti-Tyrp1, n = 14 mice that

were challenge with MPS1i-treated B16F10 CD47 KO and subsequently treated with mouse
IgG2a control, and n = 35 mice that were challenge with MPS1i-treated B16F10 CD47 KO and
subsequently treated with anti-Tyrp1. All data in which mice were challenged with DMSO-treated
cells were collected from three independent experiments. Data for the condition in which mice
were challenged with MPS1i-challenged cells and then given mouse IgG2a control were
collected from four independent experiments. Data for the final condition in which mice were
injected with MPS1i-treated cells and then treated with anti-Tyrp1 was collected from seven
independent experiments. Inset bar graphs depict response type for each indicated tumor
challenge. A partial response was defined as a mouse that survived at least one week (20+
days) beyond the median survival of the B16F10 sgCtrl cohort treated with mouse IgG2a isotype
control (13 days).

1271

1272   **(D)** Survival curves of mice up to 100 days after the tumor challenges in (B) and (C). **(i)** Survival
1273   curves for mice challenged with B16F10 sgCtrl. **(ii)** Survival curves for mice challenged with
1274   B16F10 CD47 KO. **(iii)** Triangular matrix depicting p-values between the different tested *in vivo*
1275   conditions. Statistical significance was determined by the Log-rank (Mantel-Cox) test.

1276

(E) (i) Schematic depicting the different engineering anti-tumor macrophage strategies that can
be used for validating macrophages' role in clearing CIN-afflicted B16F10 cells. Fresh bone
marrow was isolated from the tibia of donor C57BL/6 mice and was then incubated with anti-
SIRPα clone P84 (18 µg/mL) or anti-Tyrp1 (1 µg/mL) only, both together added subsequently,
or neither. Donor marrow (2×10^7 cells) is then injected intravenously into C57BL/6 harboring
B16F10 WT tumors on day 4 post-challenge. Additional anti-Tyrp1 injections, when necessary,
are done on days 5, 7, 9, 11, 13, and 15 post-tumor challenge. **(ii)** Survival curves of mice up to
100 days after the B16F10 WT tumor challenge. All mice were initially challenged with 2×10^5
B16F10 WT cells. **(iii)** Triangular matrix depicting p-values between the different tested *in vivo*
conditions. Statistical significance was determined by the Log-rank (Mantel-Cox) test.

1287

Figure 4. MPS1i-induced chromosomal instability favors induction of pro-phagocytic *de*

***novo* IgG and can lead to durable acquired immunity.**

(A) Schematic illustrating protocol for sera collection from surviving mice from Fig. 3C-D and

follow-up experiments to characterize potential *de novo* anti-cancer IgG antibodies and their

functionality both *in vitro* and *in vivo*. Serum from all mice was collected at least 100 days after

initial tumor challenge.

1294

1295 **(B) (i)** Representative flow cytometry histograms showing that convalescent sera from survivors

1296 in Fig. 3C-D contain IgG2a/c (top) and IgG2b (bottom) that bind to both WT and Tyrp1 KO

1297 B16F10 cells.

1298

(ii) Median fluorescence intensity quantification of IgG2a/c and IgG2b binding from sera from

surviving mice. Convalescent sera show statistically significant increase in both IgG2a/c and

IgG2b titers. Binding to even Tyrp1 KO cells suggests broader recognition of antigens unique to

B16F10. Statistical significance was calculated by an unpaired two-sample Kolmogorov-Smirnov

test (**** $p < 0.0001$). For IgG2a/c quantification: for binding against Tyrp1+ cells, n = 23 distinct

naïve serum samples and n = 28 distinct convalescent serum samples from surviving mice; for

binding against Tyrp1 KO cells, n = 23 distinct naïve serum samples and n = 27 distinct

convalescent serum samples. For IgG2b quantification: for binding against Tyrp1+ cells, n = 14

distinct naïve serum samples and n = 16 distinct convalescent serum samples from surviving

mice; for binding against Tyrp1 KO cells, n = 13 distinct naïve serum samples and n = 17 distinct

convalescent serum samples.

1310

1311 **(iii)** Western blotting of B16F10 lysate with either naïve sera or first challenge survivor sera as

1312 primary probe followed by anti-mouse IgG [H+L] secondary staining. Numerous bands appear

1313 when immunoblotting with convalescent survivor sera (and more than when immunoblotting with

1314 naïve sera), qualitatively confirming binding to numerous antigens and suggesting acquired

1315 immunity beyond Tyrp1.

1316

(C) (i) Phagocytosis of serum-opsonized CD47 KO or CD47/Tyrp1 double KO B16F10 cells by

BMDMs on 2D tissue culture plastic. Additionally, B16F10 cells opsonized with either anti-Tyrp1

or mouse IgG2a were included as controls for comparisons. Serum IgG derived from survivors

has both opsonization and pro-phagocytic functional ability against B16F10. Furthermore,

convalescent sera IgG from survivors is still able to drive engulfment of Tyrp1 KO cells, further

suggesting targeting of antigens beyond Tyrp1. Statistical significance was calculated by two-

way ANOVA and Tukey's multiple comparison test (mean ± SD, n = 21-23 distinct sera samples

collected from survivors for B16F10 CD47 KO phagocytosis per condition and n = 14-23 for

B16F10 CD47/Tyrp1 double KO phagocytosis per condition).

1326

(ii) Convalescent sera from first challenge survivors can repress growth of B16F10 CD47 KO

immuno-tumoroids (with macrophages). Tumoroid growth was measured by calculating the

GFP+ area at the indicated timepoints (mean ± SD, n = 16 total tumoroids from two independent

experiments for each condition, except n = 8 for opsonization with sera from naïve mice).

Statistical significance was calculated by Brown-Forsythe and Welch ANOVA tests with Dunnett

T3 corrections for multiple comparisons (ns, not significant; ** $p < 0.01$; **** $p < 0.0001$). Scale

bars = 0.5 mm.

1334

1335   **(D**) B16F10 CD47 KO cells were pre-opsonized with either convalescent sera from first

1336   challenge survivors, anti-Tyrp1 or mouse IgG2a isotype control. All mice were subcutaneously

1337   injected with $2x10^5$ pre-opsonized B16F10 CD47 KO cells.

1338

(i) Tumor growth curves shown are at early timepoints where growth is still in the linear regime.

Linear fits highlight growth suppression of tumors comprised of B16F10 CD47 KO cells pre-

opsonized with convalescent sera and anti-Tyrp1, compared to the mouse IgG2a isotype

counterparts. Mean ± SEM for all timepoints, with n = 16 mice with tumors pre-opsonized with

convalescent sera (each from a distinct survivor), n = 9 mice with tumors pre-opsonized with

anti-Tyrp1, and n = 14 mice with tumors pre-opsonized with mouse IgG2a isotype control.

Statistical significance was calculated by ordinary one-way ANOVA and Tukey's multiple

comparison test at days 9, 11, and 13 (* $p < 0.05$; **** $p < 0.0001$). Significance represented in

plot legend is representative of all three timepoints.

1348

1349   **(ii)** Survival curves up to 100 days of mice from (D-i) with pre-opsonized tumors. Both

1350   convalescent sera and anti-Tyrp1 provide similar survival benefits, suggesting potent *de novo*

1351   IgG opsonization and anti-cancer function. Statistical significance was determined by the Log-

1352   rank (Mantel-Cox) test (ns, not significant; ** $p < 0.01$; **** $p < 0.0001$).

1353

1354   **(E)** Schematic illustrating the series of experimental tumor challenges to assess acquired

1355   immunity. Survivors from the first challenge (Fig. 3C-D) were again challenged with either DMSO

1356   or MPS1i-treated B16F10 CD47 KO cells. Survivors from this second tumor challenge were once

1357   again challenged, this time with untreated B16F10 CD47 KO.

1358

(F) Survival curves of survivors from Fig. 3C-D for a second tumor challenge experiment. Prior

to tumor inoculation, B16F10 CD47 KO cells were treated with 2.5 µM MPS1i (reversine) or the

equivalent volume of DMSO vehicle control. Cells were treated for 24 h, after which they were

washed twice with PBS and allowed to recover for an additional 48 h. After the recovery period

elapsed, all mice were subcutaneously injected with 2×10^5 B16F10 CD47 KO cells. Experimental

conditions are as follows: n = 7 age-matched naïve mice (never tumor-challenged) injected with

DMSO-treated B16F10 CD47 KO cells, n = 11 surviving mice (from Fig. 3C-D) injected DMSO-

treated B16F10 CD47 KO cells, n = 6 age-matched naïve mice injected with MPS1i-treated

B16F10 CD47 KO cells, and n = 8 surviving mice (from Fig. 3C-D) injected with MPS1i-treated

B16F10 CD47 KO cells. Previous survivors challenged with DMSO-treated B16F10 CD47 KO

cells show increased median survival (21 days) compared to their naïve counterpart (14 days).

All previous survivors that were again challenged with MPS1i-treated B16F10 CD47 KO cells

survive. All mice challenged were from three independent experiments. Statistical significance

was determined by the Log-rank (Mantel-Cox) test (** $p < 0.01$; *** $p < 0.001$; **** $p < 0.0001$).

1373

(G) Non-survivors from the second tumor challenge in (F) were euthanized after tumor size was

>150 mm^2, and their tumors were excised and disaggregated for immune infiltrate analysis by

flow cytometry. **(i)** Quantification of CD45+ (immune) cells in the excised tumors, showing that

first challenge survivors still show ~2.5-fold increased tumor immune cell infiltrate despite

reaching terminal burden. n = 4 mice for age-matched naïve control, n = 5 mice that survived

the first tumor challenge. **(ii)** Quantification of tumor infiltrating CD8a+ cytotoxic T cell relative to

the total number of tumor cells. First challenge survivors show ~2-fold increase in CD8a+ T cells.

n = 4 mice for age-matched naïve control, n = 3 mice that survived the first tumor challenge. **(iii)**

Quantification of tumor infiltrating F4/80+ macrophages relative to the total number of tumor

cells. First challenge survivors show ~3-fold increase in macrophages. n = 4 mice for age-

matched naïve control, n = 5 mice that survived the first tumor challenge. **(iv)** Quantification of

MHCII+ tumor infiltrating F4/80+ macrophages relative to the total number of F4/80

macrophages. First challenge survivors show ~3-fold increase in MHCII+ macrophages. n = 4

mice for age-matched naïve control, n = 5 mice that survived the first tumor challenge. For all

experiments, mean ± SEM shown, and statistical significance was calculated by an unpaired

two-tailed t-test with Welch's correction (* $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$; **** $p < 0.0001$).

1390

(H) (i) Third tumor challenge survival curves of long-term survivors from (G). All mice were

challenged with $2x10^5$ B16F10 CD47 KO cells (n = 9 mice from three independent experiments),

delivered subcutaneously. For benchmarking and statistical comparison, survival curves from

Fig. 3F for naïve mice (n = 7 mice from three independent experiments) and second challenge

(n = 11 mice from three independent experiments) are included. Long-term survivors challenged

a third time show ~70% survival without any additional therapeutic modality, suggesting

significantly improved acquired immune response. Statistical significance was determined by the

Log-rank (Mantel-Cox) test (** $p < 0.01$; *** $p < 0.001$; **** $p < 0.0001$).

1399

(ii) Individual tumor growth curves for third challenge in long-term survivors (n = 9 mice from

three independent experiments) shown in (H-i). In total, 4 mice developed tumors, two of which

had to be euthanized prematurely due to tumor rupture despite not reach a terminal burden of

125 mm^2. These mice are still considered and included as casualties in the survival curve

analysis. The two remaining mice show significantly slower tumor growth than naïve mice

challenged with regular B16F10 CD47 KO (median survival of 14 days) and can be considered

durable partial responders.

1407

1408   **Supplementary Figure Legends**

1409

Supplementary Figure 1. Characterization of MPS1i-induced genome and chromosomal

instability in B16F10 mouse melanoma.

(A) Quantification of micronuclei induced by different MPS1 inhibitors (AZ3146, BAY 12-17389,

and reversine) at different concentrations. Statistical significance was calculated by ordinary

one-way ANOVA and Tukey's multiple comparison test (* $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$;

**** $p < 0.0001$).

(B) Consensus clustering of B16F10 analyzed in single-cell RNA-sequencing. B16F10 cells

were treated with 2.5 µM MPS1i (reversine) or the equivalent volume of DMSO vehicle control.

Cells were treated for 24 h, after which they were washed twice with PBS and allowed to recover

for an additional 48 h. Cells were then collected and processes using 10X Genomics Chromium

Single Cell Gene Expression kit for RNA isolation and library preparation.

(C) (i) UMAP plots of expression profiles for all B16F10 analyzed in single-cell RNA-sequencing.

Top: UMAP plots depict clusters in which B16F10 fall, as determined by consensus clustering.

Bottom: UMAP plots highlight cells with detectable copy number variations (CNVs), as

determined by InferCNV. Red circles indicate individual cells with at least one CNV. Gray circles

represent individual cells with no detectable CNV, suggesting these are chromosomally stable.

(ii) Quantification of the composition (MPS1i-treated or DMSO-treated B16F10) of each cluster.

1427

Supplementary Figure 2. Characterization of MPS1i-induced genome and chromosomal

instability in B16F10 mouse melanoma.

(A) Flow cytometry analysis of CD47 and Tyrp1 expression on B16F10 mouse melanoma cells.

Representative histograms for anti-CD47 (left) and anti-Tyrp1 (right) binding to B16F10 CD47

knockout (KO) and sgRNA ctrl (wild-type CD47 expression). Binding was detected by using

secondary antibodies conjugated with Alexa Fluor 647.

(B) Phagocytosis of B16F10 cells in in a standard phagocytosis assay on two-dimensional tissue

culture plastic, with or without the following: IgG opsonization by anti-Tyrp1, CD47 KO, and

MPS1i-treatment of B16F10. B16F10 cells were treated with MPS1i or DMSO prior to this assay,

following the same protocol described in Fig. 1A. **(i)** Schematic of the 2D phagocytosis assay.

(ii) Representative images of phagocytosis. B16F10 cells (green) were incubated with

opsonizing anti-Tyrp1 or mouse IgG2a control and added to adherent bone marrow-derived

macrophages (magenta). Random fields were imaged and then quantified to determine the

percentage of phagocytic macrophages. Yellow arrowheads denote phagocytic events

(complete engulfment of a target B16F10 cells). **(iii)** Quantification of phagocytic macrophages.

Statistical significance was calculated by three-way ANOVA and Tukey's multiple comparison

test. Mean ± SD shown, n = 3 replicates per condition (** $p < 0.01$; **** $p < 0.0001$).

(C) Quantification of cell size measurements by flow cytometry in both MPS1i- and DMSO-

treated cells. Statistical significance was calculated by three-way ANOVA and Tukey's multiple

comparison test. Mean ± SD shown, n = 3 replicates per condition (* $p < 0.05$; ** $p < 0.01$; *** p

< 0.001).

(D) Quantification of Tyrp1 expression by flow cytometry in both MPS1i- and DMSO-treated

cells. Statistical significance was calculated by three-way ANOVA and Tukey's multiple

comparison test. Mean ± SD shown, n = 3 replicates per condition (**** $p < 0.0001$).

(E) Quantification of CD47 expression by flow cytometry in both MPS1i- and DMSO-treated

B16F10 sgCtrl cells. Statistical significance was calculated by three-way ANOVA and Tukey's

multiple comparison test. Mean ± SD shown, n = 3 replicates per condition (**** $p < 0.0001$).

1455

Supplementary Figure 3. Macrophages readily clear CIN-afflicted tumoroids but only if

CIN is accompanied by proliferation deficits.

(A) Timeline and schematic for generating engineered "immuno-tumoroids" for time-lapsed

studies of macrophage-mediated phagocytosis of cancer cells. Tumoroids are formed by plating

and culturing B16F10 cells on non-adhesive surfaces in U-bottom shaped wells. Prior to plating

for tumoroid formation, B16F10 were treated with either MPS1i or DMSO (AZ3146, BAY 12-

17389 or reversine). Tumoroid growth was measured by calculating the GFP+ area at the

indicated timepoints (mean ± SD, n = 16 total tumoroids from two independent experiments for

each condition). All data were then normalized to average GFP+ area on day 1 of each drug

treatment's respective mouse IgG2a isotype control condition. Immuno-tumoroids are imaged at

the listed timepoints. **(B)** To assess macrophage-mediated clearance, ~24 h after plating, bone

marrow-derived macrophages (BMDMs) with or without opsonizing anti-Tyrp1 are added to the

cohesive B16F10 tumoroids. Tumoroid growth curves for DMSO-treated B16F10 CD47

knockout (KO) cells. **(C)** Tumoroid growth curves (with BMDMs) for MPS1i-treated B16F10

CD47 KO cells (treated with varying concentrations of MPS1i): **(i)** AZ3146, **(ii)** BAY 12-17389.

(D) Tumoroid growth curves (without BMDMs) for MPS1i-treated B16F10 CD47 KO cells (treated

with varying concentrations of MPS1i): **(i)** BAY 12-17389, **(ii)** reversine. Both BAY 12-17389 and

reversine induce proliferation deficits, which may play an essential role in mediating clearance

by macrophages that could not be observed in AZ3146-treated cells.

1475

Supplementary Figure 4. MPS1i-induced chromosomal instability upregulates MHC-1

class molecules on B16F10, suggesting increased antigen presentation.

Flow cytometry analysis and quantification H-2K^b expression on B16F10 mouse melanoma cells.

B16F10 cells were treated with MPS1i or the equivalent volume of DMSO vehicle control.

Concentrations used: 2.5 µM AZ3146, 1.0 µM BAY 12-17380, and 2.5 µM reversine. Cells were

treated for 24 h, after which they were washed twice with PBS and allowed to recover for an

additional 48 h. Binding was detected by using primary antibodies conjugated with Alexa Fluor

647. Statistical significance was calculated by Brown-Forsythe and Welch ANOVA and Dunnett's

T3 multiple comparison test. Mean ± SEM shown, n = 3 independent replicates (* $p < 0.05$; ** p

< 0.01; *** $p < 0.001$; **** $p < 0.0001$).

1486

Supplementary Figure 5. Flow cytometry gating strategy for identification & quantification

of macrophage infiltrate and characterization in CIN-afflicted and chromosomally stable

B16F10 tumors.

Representative flow cytometry gating strategy for *in vivo* B16F10 CD47 KO tumor immune

infiltrate five days after initial challenge. Tumors were comprised of cells treated with either **(i)**

DMSO or **(ii)** MPS1i (2.5 µM), following the timeline schema in Fig. 2F. Singlets were separated

from debris, doublets, and aggregates by FSC-A vs FSC-H and SSC-A vs SSC-H gates. Debris

and dead cells were further removed by FSC-A vs SSC-A gating. Overall immune cell infiltrate

was determined by CD45+ expression. Macrophages were then further identified based on

Ly6G- and F4/80+ expression. Upon identification of macrophages, cells were further

characterized for polarization based on CD86 (M1-like marker) and CD206 (M2-like marker)

expression.

1499

Supplementary Figure 6. MPS1i-induced CIN favors clearance when paired with CD47 KO

and IgG opsonization, regardless of the degree of CIN.

Tumor growth curve of projected tumor area versus days after tumor challenge, with B16F10

CD47 KO cells. Each line represents a separate tumor and is fit with an exponential growth

equation: $A = A_0 e^{kt}$. Complete anti-tumor responses in which a tumor never grew are depicted

with the same symbol as their growing counterparts and with solid lines at A = 0.

(A) Tumor growth curve data in which there was no complete response, from Fig. 3C. n = 7 mice

that were challenged with DMSO-treated B16F10 CD47 KO and subsequently treated with

mouse IgG2a control, n = 9 mice that were challenged with DMSO-treated B16F10 CD47 KO

and subsequently treated with anti-Tyrp1.

(B) Tumor growth curve data from mice challenges with CIN-afflicted B16F10, but with varying

concentration of MPS1i used to cause different degrees of instability. All mice here were

challenged with B16F10 CD47 KO and treated with anti-Tyrp1 for conditions of maximal

phagocytosis and to better compare to the ~97% cure rate in Fig. 3C, 4D-ii. n = 6 mice challenged

with cells with each tested MPS1i concentration: 0.1, 0.5, 1.0 or 2.5 µM.

(C) Survival curves of mice up to 100 days after the tumor challenges in (A) and (B).

1516

Supplementary Figure 7. Survivors challenged with CIN-afflicted tumors generate anti-

cancer IgG, regardless of the degree of CIN.

Median fluorescence intensity quantification of IgG2a/c and IgG2b binding from sera from

surviving mice with different degrees of chromosomal instability by titrating reversine

concentration used to treat B16F10 CD47 KO cells before injection. Convalescent sera are

collected from survivors from Fig. S6. Binding of IgG2a/c and IgG2b against **(A)** B16F10 cells

expressing Tyrp1 and **(B)** B16F10 Tyrp1 knockout (KO). n = 4-5 distinct sera samples collected

from survivors.

1525

Supplementary Figure 8. Flow cytometry gating strategy for identification & quantification

of immune infiltrate and characterization in re-challenge experiments.

Representative flow cytometry gating strategy for *in vivo* B16F10 CD47 KO tumor immune

infiltrate from 2nd challenge non-survivors and age-matched naïve controls. Singlets were

separated from debris, doublets, and aggregates by FSC-A vs FSC-H and SSC-A vs SSC-H

gates. Debris and dead cells were further removed by FSC-A vs SSC-A gating. For T cell

quantification: CD45+ cells were gated on and then analyzed for CD3e and CD8a surface protein

expression. For macrophage quantification: myeloid cells were isolated by CD45+ and CD11b+

expression, after which macrophages were isolated based on F4/80 expression.

1535

1536

1537

Figure 1

MPS1i-induced chromosomal instability & deficits in proliferation favor pro-inflammatory macrophages

Figure 2

Macrophages readily clear genomically unstable tumoroids & skew to an M1-like phenotype

A

3D tumoroid phagocytosis assay

Plate DMSO or MPS1i-treated B16s to form tumoroids

Image

Day 0 1 5 10

Add macrophages + anti-Tyrp1 or mIgG2a isotype

Flow cytometry for macrophage polarization markers

non-adhesive surface in 96-well U-bottom

+ macrophages, anti-Tyrp1/mIgG2a

B

M1 markers: MHCII, CD86

M2 markers: CD163, CD206

DMSO / MPS1i

event # — Binding signal

C

MHCII — gMFI (× 1000) — DMSO vs MPS1i **

CD86 — DMSO vs MPS1i *

CD163 — DMSO vs MPS1i **

CD206 — DMSO vs MPS1i **

D

DMSO mIgG2a ctrl / DMSO anti-Tyrp1 / MPS1i mIgG2a ctrl

Day 1, Day 4, Day 7, Day 10

E

Normalized GFP area (to day 1 average area of mIgG2a condition)

DMSO — mIgG2a ctrl / anti-Tyrp1

0.1 µM MPS1i / 0.5 µM MPS1i / 1.0 µM MPS1i / 2.5 µM MPS1i / 5.0 µM MPS1i

Days of tumoroid growth

Genomically unstable tumors show increased immune infiltrate & particularly greater macrophage numbers at early stages of engraftment

F

Inoculate with DMSO or MPS1i-treated B16F10 CD47 KO

Day 0 1 2 3 4 5

+ 250 µg anti-Tyrp1, i.v.

Euthanize & harvest tumor for immune infiltrate measurements

G

B16F10 treatment: DMSO / MPS1i

(i) % CD45+ of all tumor-harvested cells ****

(ii) % CD45+ CD11b+ of all tumor-harvested cells ****

(iii) % of F4/80+ macrophages to all tumor-harvested cells ****

(iv) % of polarized macrophages to all tumor-harvested cells — CD86hi CD206lo ** / CD86lo CD206hi *

Figure 3

Figure 4

Surviving chromosomally unstable tumors leads to a broad *de novo*
pro-phagocytic & anti-cancer IgG response

A Collect convalescent serum from survivors

Opsonize target cells with anti-Tyrp1 or serum

Experimental validation of *de novo* anti-cancer IgG

Quantify IgG binding

Assess functional effect on phagocytosis

Assess effects on *in vivo* growth rate

B (i) IgG2a/c binding
- naive serum
- Conv. serum on B16F10 Tyrp1+
- Conv. serum on B16F10 Tyrp1 KO

IgG2b binding
- naive serum
- Conv. serum on B16F10 Tyrp1+
- Conv. serum on B16F10 Tyrp1 KO

event #

Binding signal

(ii) Binding against B16F10 Tyrp1+ / Binding against B16F10 Tyrp1 KO

Median fluorescence intensity (IgG2a, IgG2b) — ****

naive serum / 1st chal. conv. serum

(iii) naive / convalescent — MW (kDA): 260, 160, 110, 80, 60, 50, 40 (ctrl), 30, 20. ctrl = β-actin

C (i) % phagocytic macrophages: CD47 KO, CD47/Tyrp1 DKO — ****, ns

Legend: mIgG2a ctrl, naive serum, 1st chal. conv. serum, anti-Tyrp1

(ii) mIgG2a / naive / anti-Tyrp1 / Convalescent

GFP+ area (mm²) — 5 days of tumoroid growth — ****, ns, **

D (i) Tumor area (mm²) vs Days post-challenge
- 1st chal. conv. serum
- anti-Tyrp1
- mIgG2a ctrl

*, ****

(ii) % survival vs Days post-challenge — ns, **, ****
- 1st chal. conv. serum — 4/16
- anti-Tyrp1 — 2/9
- mIgG2a ctrl

Rechallenges on survivors further shows acquired immunity & immune response
against both genomically stable and unstable tumors

E First challenge (Fig. 3C) → 2nd challenge on survivors → 3rd challenge on survivors with only untreated B16F10 CD47 KO

F (i) Second challenge survival — % survival vs Days post-challenge

Challenge status	MPS1i	Survivors
Naive	—	0/7
Previously cured	—	1/11
Naive	+	1/6
Previously cured	+	8/8

B16F10 CD47 KO rechallenge

****, ***, ****, **

G Immune infiltrate in 2nd challenge non-survivors

Legend: naive / 1st challenge survivor

(i) % CD45+ of all tumor-harvested cells — *

(ii) % of CD3+ CD8a+ T cells to all tumor-harvested cells — ****

(iii) % of F4/80+ macrophages to all tumor-harvested cells — **

(iv) % of MHCII+ macrophages to all tumor-harvested cells — ***

H (i) Third challenge survival — % survival vs Days post-challenge
- 6/9
- 1/12

(ii) Third challenge growth — Tumor area (mm²) vs Days post-challenge
- tumor rupture, mouse euthanized

Legend: Naive, 2nd challenge, 3rd challenge — **, ****, ****

Supplementary Figure 1

Induction of micronuclei and chromosomal instability in B16F10 by MPS1i

A

AZ3146, BAY 12-17389, Reversine (from Fig. 1B-ii for comparison)

MPS1i [uM]

B, **C (i)**, **(ii)**

Supplementary Figure 2

Characterization & validation of factors affecting phagocytosis

A

CD47 KO validation
- Negative ctrl
- CD47 KO
- sgRNA ctrl

# of events — CD47 protein (AF647)

Tyrp1 validation
- Negative ctrl
- Tyrp1 KO
- CD47 KO
- sgRNA ctrl

# of events — Tyrp1 protein (AF647)

B

(i) Standard 2D phagocytosis assay

Macrophages
Opsonized B16F10

2D tissue culture plastic

(ii) Target: B16F10 CD47 KO

mIgG2a ctrl | anti-Tyrp1

DMSO

MPS1i

Macrophages
Opsonized B16F10

(iii)

% phagocytic macrophages

CD47	WT	WT	KO	KO	WT	WT	KO	KO
MPS1i	—	—	—	—	+	+	+	+
anti-Tyrp1	—	+	—	+	—	+	—	+

C Size measurements

FSC-A median intensity vs MPS1i [uM] (0, 0.1, 0.5, 1.0, 2.5)

D Tyrp1 expression

Tyrp1 MFI vs MPS1i [uM] (0, 0.1, 0.5, 1.0, 2.5)

E CD47 expression in B16F10 sgCtrl

CD47 MFI vs MPS1i [uM] (0, 0.1, 0.5, 1.0, 2.5), CD47 KO

Supplementary Figure 3

MPS1i-induced CIN favors clearance with CD47 ablation and IgG opsonization, regardless of MPS1i concentrtion

Supplementary Figure 4

MPS1i also induces upregulation of MHC-I class molecules

Supplementary Figure 5

Immune infiltrate analysis in MPS1-treated and DMSO-treated tumors

A   **Chromosomally stable tumor (DMSO-treated B16F10)**

B   **Chromosomally unstable tumor (MPS1i-treated B16F10)**

Supplementary Figure 6

MPS1i-induced CIN favors *in vivo* clearance with CD47 ablation and IgG opsonization, regardless of degree of CIN

Supplementary Figure 7

Quantification of IgG binding for convalescent sera
from titrated CIN tumor challenge survivors

Quantification of IgG binding for convalescent sera
from adoptive marrow transfer survivors

Supplementary Figure 8

Flow cytometry analysis of immune infiltrate in 2nd tumor challenge mice