# Prediction of fast subcellular calcium signals based on label-free second harmonic generation microscopy measurements

Tihomir Georgiev[1*]

[1]Department of Experimental Ophthalmology, University Hospital Cologne, University of Cologne, Cologne, Germany

[*]Correspondence to: tihomir.georgiev@mail.de

## Abstract

Calcium ($Ca^{2+}$) sparks were predicted based on the label-free second harmonic generation (SHG) signal of myosin filaments with pix2pix, a conditional adversarial network. Predicted $Ca^{2+}$ sparks and ground truth $Ca^{2+}$ sparks show similar properties. Predicted $Ca^{2+}$ spark frequencies and a predominant location at the periphery of skeletal muscle fibers were in good agreement with ground truth data. High frequencies of predicted $Ca^{2+}$ sparks could be observed at y-shaped structures of myosin filaments. Important image pre-processing was applied in order to enhance the signal of $Ca^{2+}$ sparks used for training. Only open-source software was applied. I provide the image pre-processing algorithm and several algorithms to evaluate the quality of predicted $Ca^{2+}$ sparks. This technique can be extended to other label-free microscopy techniques and other fluorescent indicators developed during the last years. It remains a difficult task to predict the exact location of $Ca^{2+}$ sparks.

## Introduction

$Ca^{2+}$ is an important intracellular signal[1]. $Ca^{2+}$ sparks are fast and small $Ca^{2+}$ signals in different cell types[2,3]. High frequencies of $Ca^{2+}$ sparks can be observed in permeabilized skeletal muscle fibers[4] and during an osmotic treatment in intact skeletal muscle fibers[5]. Machine learning has been applied for the analysis of $Ca^{2+}$ signals recently[6-12]. Generative adversarial networks have been introduced by Goodfellow et al[13] and have been applied for segmentation of $Ca^{2+}$ signals[14]. Machine learning approaches for the analysis of $Ca^{2+}$ sparks are rare[15]. Recently, machine learning has been applied for segmentation and classification of $Ca^{2+}$ sparks[15-17]. Additionally, T cell classification based on intracellular calcium signals has been carried out using machine learning predictions[18].

In light microscopy, in silico labelling of subcellular structures based on bright field

images has been achieved with deep learning[19,20]. Image translations of medical

images have been generated with deep learning for different applications such as

synthesis of post contrast MRI images[21] or translation of MRI images to PET

images[22].

To my knowledge, this paper is the first attempt to translate structural images to

images of $Ca^{2+}$ signals. Second harmonic generation (SHG) microscopy is a

nonlinear optical microscopy technique that allows the label-free visualization of

myosin filaments in skeletal muscle fibers[23,24]. Localized $Ca^{2+}$ signals have been

investigated with respect to the myosin filaments in striated muscle cells previously

[25,26]. In this study, I used the SHG signal of myosin filaments as subcellular structural

information to generate synthetic images of $Ca^{2+}$ signals. The training of the network

was carried out on pairs of images consisting of an SHG measurement and a two

photon microscopy[27] measurement of $Ca^{2+}$ sparks in skeletal muscle fibers. For the

training process of the network, an important image pre-processing of the $Ca^{2+}$

signals was applied in order to enhance the signal of $Ca^{2+}$ sparks compared to the

cellular background. Different strategies have been applied to translate images

previously[28,29]. I applied the pix2pix algorithm[29] for the translation of images which is

based on conditional adversarial networks. In principle this technique can be

extended to other label-free microscopy techniques and other fluorescent indicators

such as dopamine[30-33], acetylcholine[34,35], glutamate[36,37], GABA[38], ATP[12,39],

serotonin[40], norepinephrine[41] and adenosine[42] indicators. The prediction of $Ca^{2+}$

sparks based on subcellular structures is a difficult task since the exact location of

$Ca^{2+}$ sparks is difficult to predict. Additionally, $Ca^{2+}$ sparks are small and fast

subcellular signals in a noisy cellular environment.

Changes in the SHG signal depending on different conformations of myosin

molecules have been described in skeletal muscle fibers and myofibrils[43-45]. An

increase of the SHG signal in skeletal muscle fibers has been shown after an

increase of the intracellular $Ca^{2+}$ concentration[45]. Additionally, localized sarcomere

contractions could be observed in response to $Ca^{2+}$ sparks in cardiac muscle cells[26].

Therefore, I investigated possible subcellular changes of the SHG signal at regions of

$Ca^{2+}$ sparks.

# Results

## Generation of translated images

As a first step, $Ca^{2+}$ sparks were enhanced compared to the background signal of the cell (Fig. 1a). Then, the network was trained on pairs of SHG images and enhanced $Ca^{2+}$ images (Fig. 1b). The trained network was then applied to translate $Ca^{2+}$ images based on SHG images (Fig. 1c). Fig. 1c shows only one $Ca^{2+}$ spark in the translated image. Some translated $Ca^{2+}$ sparks could be seen on every image of the measurement. Other translated $Ca^{2+}$ sparks appeared only on some images of the measurement (Extended Data Fig. 1b, c, f). On several translated images two, three or four $Ca^{2+}$ sparks could be observed (Extended Data Fig. 1a, b, c, f). $Ca^{2+}$ sparks could be found at subcellular structures of the myosin pattern, so called y-shaped structures (Fig. 1d).

I tested the performance of the trained network on three different experiments Exp 1 (experiment 1), Exp 2 (experiment 2), Exp 3 (experiment 3). In Exp 1 measurements of fibers were translated that were not included in the training data. Measurements showing a plane of a skeletal muscle fiber unseen during training were translated in Exp 2. However, measurements of another plane of this fiber (Exp 2) were included in the training data. Exp3 contains two measurements of a different fiber. Other measurements of the same plane of this fiber were included in the training data.

Structural similarity index (SSIM)[46] and mean squared error (SME)[47] were calculated (Fig. 1e and f). The translated images achieved high SSIM values.

87 **Figure 1** *Generation of translated images. a, Pre-processing of Ca²⁺ sparks with enhancement of*

88 *Ca²⁺ sparks compared to the background signal of the fiber. On the left side Ca²⁺ sparks (arrows)*

89 *without enhancement are shown. On the right side Ca²⁺ sparks after enhancement can be seen. Scale*

## Properties of translated Ca$^{2+}$ images

As a next step, I investigated several properties of translated images. The fiber

masks of ground truth and predicted images were similar (Fig. 2a, b, c and d). The

mean fiber intensity was lower in the predicted images, this was particularly the case

in experiment 1 (Exp 1) (Fig. 2e). However, the mean of the fiber mean fluorescence

intensity in the ground truth data of Exp 1 was higher than in the other two

experiments and in the training data set.

Then, I investigated the properties of predicted Ca$^{2+}$ sparks compared to ground truth

Ca$^{2+}$ sparks. I found that predicted Ca$^{2+}$ spark area in Exp 1 was in good agreement

with ground truth Ca$^{2+}$ spark area (Fig. 2f). Interestingly, larger differences could be

observed for the other two experiments. The predicted spark mean intensity was

lower in all experiments (Fig. 2g). This was most pronounced in Exp 1. The predicted

ratio Ca$^{2+}$ spark area/ Ca$^{2+}$ spark edge was larger than this ratio in the ground truth

data for all experiments suggesting that predicted Ca$^{2+}$ sparks are more circular (Fig.

2h).

The predicted Ca$^{2+}$ spark frequency in Exp 1 of 2.52 sparks/(10000*µm$^2$) was in good

agreement with the ground truth Ca$^{2+}$ spark frequency of 3.05 sparks/(10000*µm$^2$)

(Fig. 2i). The predicted Ca$^{2+}$ spark frequency in experiment 3 (Exp 3) of 2.78

sparks/(10000*µm$^2$) was even closer to the corresponding ground truth Ca$^{2+}$ spark

frequency of 2.74 sparks/(10000*µm$^2$). The training data set had a Ca$^{2+}$ spark

frequency of 1.85 sparks/(10000*µm$^2$). However, there was a larger difference in

experiment 2 (Exp 2) between the predicted Ca$^{2+}$ spark frequency of 2.46

sparks/(10000*µm$^2$) and the ground truth Ca$^{2+}$ spark frequency of 4.36

sparks/(10000*µm$^2$).

**Figure 2 Properties of translated images of Ca$^{2+}$ sparks. a**, Ground truth (red) and predicted
(green) fiber mask of a fiber unseen during training of the network (Exp 1). The overlay of both masks
is yellow. Scale bar 20 µm. **b** and **c**, F1 scores and Matthews correlation coefficients for the evaluation
of predicted fiber masks for the three experiments. **d**, **e**, **f**, **g** and **h**, Box plots and violin plots of the
fiber area, fiber mean intensity, spark area, spark mean intensity and spark area/spark edge of the
ground truth (Exp GT) and predicted (Exp P) images in the three experiments and the training data
set. Orange lines are the medians of the data and green triangles are the means. **i**, Ca$^{2+}$ spark
frequencies of the ground truth and predicted images of the three experiments and the training data
set.

## Translation of subcellular structures to fast subcellular signals

As a next step, I investigated specific subcellular regions of the muscle fiber known to
have high frequencies of Ca$^{2+}$ sparks, the fiber periphery[5] and y-shaped structures of
the myosin pattern[25]. Additionally, I investigated the ratio Ca$^{2+}$ spark area over
myosin filaments/spark area.

As expected, most of the predicted and ground truth Ca$^{2+}$ sparks were located at the
fiber periphery (Fig. 3a, b, and c). The ground truth Ca$^{2+}$ spark frequency at the fiber
periphery in Exp 1 showed a lower fraction of Ca$^{2+}$ sparks compared to the predicted
fraction. This difference can be also seen in the ratio Ca$^{2+}$ spark area over fiber
periphery/spark area (Extended Data Fig. 3b). One of the fibers used in Exp 1
showed a high frequency of Ca$^{2+}$ sparks in the centre of the fiber not seen for the
other five fibers (Extended Data Fig. 2). Interestingly, predicted images of this fiber
showed several Ca$^{2+}$ sparks in the fiber centre in close proximity to the ground truth
Ca$^{2+}$ sparks but not at such high frequencies (Extended Data Fig. 2). If this fiber was
excluded from the analysis, there is a good agreement between ground truth data
and predicted data (Extended Data Fig. 3a).

**Figure 3  Translation of subcellular structures to fast subcellular signals. a,** Binary images of
the fiber centre (cyan), periphery of the fiber (blue) and $Ca^{2+}$ sparks (green) based on a predicted $Ca^{2+}$
image of a fiber unseen during training of the algorithm (Exp 1). Additionally, a binary image of a $Ca^{2+}$
spark measured in the ground truth image is presented (red). Scale bar 20 µm. **b,** Overlay of the
predicted $Ca^{2+}$ signal (green) of the fiber shown in **a** and fiber periphery (blue). The arrows indicate
$Ca^{2+}$ sparks. **c,** Percentage of $Ca^{2+}$ sparks at the fiber periphery for ground truth (Exp GT) and
predicted (Exp P) images in the three experiments and the training data set. **d,** Overlay of binary
images (left) of the myosin filaments (red) and a predicted $Ca^{2+}$ spark. The binary SHG signal is based
on a mean SHG signal of the whole measurement. Overlay of the mean SHG signal of the whole
measurement (red) and the predicted $Ca^{2+}$ signal (green) of the same image (right). This fiber was
unseen during the training process of the algorithm (Exp 1). Scale bar 10 µm. **e,** Box plots and violin
plots of the ratios of the $Ca^{2+}$ spark area over myosin filaments and the $Ca^{2+}$ spark area for ground
truth (Exp GT) and predicted (Exp P) images in the three experiments and the training data set.
Orange lines are the medians of the data and green triangles are the means. **f,** Overlay of the mean
SHG signal during the whole measurement (yellow), mask of y-shaped structures (blue), binary image
of a predicted $Ca^{2+}$ spark (green) and binary image of a ground truth $Ca^{2+}$ spark (red). The fiber was
unseen during the training process (Exp 1). Scale bar 5 µm. **g,** $Ca^{2+}$ spark frequencies at y-shaped

Then, I looked at the ratio Ca$^{2+}$ spark area over myosin filaments/spark area (Fig. 3d

and e) and found that this ratio was lower for predictions. Several predicted Ca$^{2+}$

sparks were found at regions without myosin filaments (Exp 1 P and Exp 2 P). Such

Ca$^{2+}$ sparks at myosin free regions can be found in the training data set.

As a next step, I investigated the translation of y-shaped structures of the myosin

filament pattern to Ca$^{2+}$ sparks (Fig. 3f, g and h and Extended Data Fig. 4). In Exp 1,

the predicted Ca$^{2+}$ spark frequency at y-shaped structures of 7.94

sparks/(10000*μm$^2$) was 1.26 times higher than the ground truth Ca$^{2+}$ spark

frequency at y-shaped structures of 6.31 sparks/(10000*μm$^2$). In Exp 2, no Ca$^{2+}$

sparks were predicted at y-shaped structures while ground truth Ca$^{2+}$ frequency at y-

shaped structures was 3.92 sparks/(10000*μm$^2$). In Exp 3, the predicted Ca$^{2+}$ spark

frequency at y-shaped structures of 13.93 sparks/(10000*μm$^2$) was 1.56 times higher

than the ground truth Ca$^{2+}$ spark frequency at y-shaped structures of 8.95

sparks/(10000*μm$^2$).

**Accuracy of Ca$^{2+}$ spark prediction**

Predicted Ca$^{2+}$ sparks seen in all image translations of a measurement were several

times associated with high ground truth Ca$^{2+}$ spark activity in close proximity (Fig. 4a,

b, c and d, Extended Data Movie 1 and 2). The predicted and ground truth Ca$^{2+}$

sparks partly overlapped in these cases. However, the exact overlap was not high.

This is reflected in low Pearson correlation coefficients, Matthews correlation

coefficients and F1 scores (Fig. 4 e, f, g). The predicted Ca$^{2+}$ sparks seen in all

image translations of a measurement were in close proximity or at y-shaped

structures of the myosin filaments (Extended Data Fig. 5).

***Figure 4   Accuracy of Ca²⁺ spark prediction. a**, Image sequence (10 consecutive images) of the*
*ground truth Ca²⁺  signal (red) and the predicted Ca²⁺ signal (green). The yellow arrow indicates a*
*predicted Ca²⁺ spark. In close proximity to the predicted Ca²⁺ spark ground truth Ca²⁺ sparks can be*
*observed. Predicted and ground truth Ca²⁺ sparks overlap partly sometimes. This fiber was unseen*
*during the training process (Exp 1). Scale bar 10 µm. **b**, Image sequence of another fiber. Other*
*measurements of this fiber and this plane were seen during the training process (Exp 3). The yellow*
*arrow indicates a predicted Ca²⁺ spark (green) that is close to several ground truth Ca²⁺ sparks (red)*
*and on several images they partly overlap. The green arrow shows at a predicted Ca²⁺ spark that is*
*not in close proximity to ground truth signals. The red arrow indicates a ground truth Ca²⁺ spark that is*
*not in close proximity to predicted Ca²⁺ sparks. Scale bar 10 µm. **c** and **d**, xyt volumes of the*
*measurements partly shown as image sequences in **a** and **b**. Yellow arrows indicate predicted Ca²⁺*
*sparks in close proximity to many ground truth Ca²⁺ sparks partly overlapping with them. Red arrows*
*show at ground truth Ca²⁺ sparks with no corresponding predicted Ca²⁺ sparks at this location. Green*
*arrow shows at predicted Ca²⁺ sparks with no corresponding ground truth Ca²⁺ sparks at this location.*
*Box volume 34.88 µm, 34.88 µm, 122.18 s in **c** and 34.88 µm, 34.88 µm, 40.18 s in **d**. **e**, Pearson*
*correlation coefficients of the whole images (light blue) and of the overlapping fiber masks of predicted*
*and ground truth images (dark blue) for the three experiments. **f** and **g**, Matthews correlation*
*coefficients and F1 scores calculated based on binary images of ground truth Ca²⁺ sparks and*
*predicted Ca²⁺ sparks for the three experiments.*

Extended Data Fig. 6 and 7 present the results when only one fiber was included in
Exp1.

### Does the algorithm translate subcellular structures or signals?

To investigate possible subcellular changes of the SHG signal at regions of Ca²⁺
sparks, I analysed if there is such correlation (Fig 5a, b and c). If there are Ca²⁺
dependent changes of the SHG signal, it is possible that the algorithm translates
localized signals in SHG images to localized signals in Ca²⁺ images. The Pearson
correlation coefficients for the different experiments are: Exp 1 GT 1nf: 0.0727, Exp 1
P 1nf: -0.4127, Exp 1 GT 1: -0.2998, Exp 1 P 1: -0.4564, Exp 1 GT 2: 0.0818, Exp 1
P 2: 0.6419, Exp 2 GT: 0.0299, Exp 2 P: -0.4498, Exp 3 GT: -0.1464, Exp 3 P: -
0.4887, Training: -0.1896.

Interestingly, Exp 1 P 1nf shows negative Pearson correlation for the predicted data.
The ratio SHG signal (spark)/SHG signal (fiber) is lower for Exp 1 P 1nf compared to
Exp 1 GT 1nf (Fig 5d).

244 The F1 score for the fiber masks in Exp 1 c1nf (cell 1 without a band pass filter) is
245 somewhat lower than the F1 score in Exp 1 c1 and Exp 1 c2 (Extended Data Fig.8).
246 The F1 score for Ca$^{2+}$ sparks is highest in Exp 1c1nf (Extended Data Fig.8).

## Discussion

Several properties of Ca$^{2+}$ signals predicted based on label-free SHG measurements
were in good agreement with the ground truth Ca$^{2+}$ signals. Predictions of fast
subcellular signals based on measurements of label-free techniques have the
potential to make obsolete elaborate staining of cells or one additional channel is
then available for another fluorescent marker. The frequency of Ca$^{2+}$ sparks, the fiber
area and the Ca$^{2+}$ spark area (in Exp 1) could be predicted quite well. Ca$^{2+}$ sparks at
myosin free regions might be nuclear Ca$^{2+}$ sparks[48]. The poor prediction of Ca$^{2+}$
sparks in Exp 2 suggests that a different plane of the same fiber is already a different
scene with different myosin pattern and associated Ca$^{2+}$ sparks signalling pattern. As
expected, most predicted Ca$^{2+}$ sparks were located at the fiber periphery.

Predicted Ca$^{2+}$ sparks seen in all images of a measurement were in several cases
associated with locations of high ground truth Ca$^{2+}$ sparks activity.

The better F1 scores for Ca$^{2+}$ sparks in Exp 1 1nf compared to Exp 1 1 (Extended
Data Fig. 8) might be explained by a different part of the fiber that is easier to predict.
The influence of a different plane for the prediction of Ca$^{2+}$ sparks has been shown in
Exp 2. Although the training data shows a negative Pearson correlation, I think that
results in Fig. 5 suggest that the algorithm translates subcellular structures. In
contrast to the training data, Exp 1 GT 2 and Exp 2 GT show somewhat positive

Pearson correlation. Additionally, a low SHG signal at $Ca^{2+}$ sparks might be
explained by specific positions within the fiber (perinuclear, nuclear signals or signals
at the periphery of the fiber). $Ca^{2+}$ sparks in skeletal muscle fibers might not be
sufficient to increase the SHG signal as it was observed after a $Ca^{2+}$ transient in
skeletal muscle fibers[45]. But this needs further investigation.

Local changes in the sarcomere length may be an important information for the
algorithm[26]. However, I did not investigate such local changes in the sarcomere
length. Additionally, polarization dependent effects of the SHG signal[43-45] might
influence the decision of the algorithm.

The prediction of the exact location of $Ca^{2+}$ sparks based on subcellular structures
remains a difficult task. In further improvement of the algorithm the 3D nature of xyt
volumes should be taken into account instead of translating 2D images. Other
existing algorithms such as U-net[49] could be tested for such a task as it was done in
an image translation study of medical images[21]. It can be expected that data
augmentation of the training data set[12,49,50], pre-processing of the SHG signal,
denoising, larger data sets, powerful computational resources and algorithms
designed for this specific task will further improve the accuracy of such image
translations.

I hope that this work will stimulate further research to improve the translation of fast
subcellular signals based on subcellular structures or other subcellular signals.

## Methods

### Data

The SHG images and two photon microscopy images used to train and test the
algorithm were measured simultaneously in a previous publication[25].

### Image pre-processing

The first step of the image pre-processing was to automatically identify $Ca^{2+}$ sparks.
Similar ideas for automatic detection of $Ca^{2+}$ sparks have been described in a
previous publication[51]. First, a gaussian filter was applied on each xyt measurement.
Then, based on the gaussian signal image the mean intensity was calculated along
the t axis, a fiber mask was generated using the mean image with a threshold of $2*\sigma$,
with $\sigma$ the standard deviation. The mask was labelled and objects smaller than 3000

pixels were excluded. Then, a cleaned binary fiber mask was created. To exclude labelled structures outside the fiber, the $Ca^{2+}$ measurement after application of a gaussian filter was multiplicated by the cleaned binary fiber mask. Binary image that includes the $Ca^{2+}$ sparks was generated. Pixels were assigned a value of 1 if their value is higher than $\mu + 3.6* \sigma$, with $\mu$ the mean and $\sigma$ the standard deviation. $\mu$ and $\sigma$ were measured within the fiber mask. Then objects of the binary images were labelled and objects smaller than 55 pixels were excluded. Then again, a binary image of $Ca^{2+}$ sparks was created based on the cleaned labelled image.

Finally, the $Ca^{2+}$ sparks in the original image were enhanced 10 times compared to the fiber background by reducing the fiber background 10 times. This step was important in order to make it easier for the network to see the $Ca^{2+}$ sparks.

**Translation of $Ca^{2+}$ sparks based on SHG images**

19 measurements of 4 fibers from two mice were used to train the algorithm. The training was carried out with 4123 images containing 256x256 pixels with 818 $Ca^{2+}$ sparks. Thereby, 1031 images containing 512x512 pixels were cropped to 4123 256x256 pixel images.

To test the performance of the trained network, three different experiments Exp 1 (experiment 1), Exp 2 (experiment 2), Exp 3 (experiment 3) were carried out.

Exp 1 consists of 545 512x512 pixel images from 5 measurements of two fibers from two mice (the same animals as used for training but different fibers), which were not included in the training data. 683 ground truth $Ca^{2+}$ sparks are included in this data set.

Exp 2 consists of 196 512x512 pixel images from 4 measurements of a fiber that was included in the training data but showing a different plane. This data set contains 292 ground truth $Ca^{2+}$ sparks.

Exp 3 consists of 98 512x512 pixel images from two measurements of a fiber. Different measurements of this muscle fiber were included at the same plane in the training data. The data set contains 102 ground truth $Ca^{2+}$ sparks.

Fibers were selected if they showed clear $Ca^{2+}$ sparks activity and a band pass filter was placed in the SHG signal detection path[25] and no or little shift of the muscle fiber was observed during measurement. Additionally, 2 measurements were included in

Exp 1 without a band pass filter in front of the SHG detector in order to test if a
contamination of the SHG signal influences the prediction of $Ca^{2+}$ sparks.

Since there is a high variability of $Ca^{2+}$ sparks signalling patterns and because no
augmentation was carried out during the training process, images in Exp 1 and Exp 2
were rotated at 90°, 180°, 270° and 360°. Therefore, the algorithm was tested on
2180 512x512 pixel images in Exp1 and 784 512x512 pixel images in Exp 2.

All images in the three experiments were cropped to 256x256 pixel images as input
for the image translation algorithm.

In Exp 1 and in Exp 2 for the four angles 2307 and 702 $Ca^{2+}$ sparks were predicted
respectively. In Exp 3 103 $Ca^{2+}$ sparks were predicted.

Additional analysis was carried out for Exp 1 (Fig. 5 and Extended Data Fig. 8). Exp 1
was divided in three different data sets. Exp 1 GT 1nf is the ground truth data set of
fiber 1 without a band pass filter in front of the SHG signal detector. Exp 1 GT 1nf
consists of 401 ground truth $Ca^{2+}$ sparks. Exp 1 P 1nf are the predicted images of
Exp 1 GT 1nf with 1628 $Ca^{2+}$ sparks after rotation of images with 4 different angles
as input. Exp 1 GT 1 is the ground truth data set of fiber 1 with a band pass filter in
front of the SHG signal detector. This data set contains 90 $Ca^{2+}$ sparks. The
corresponding prediction Exp 1 P 1 for 4 rotations contains 155 $Ca^{2+}$ sparks. Exp 1
GT 2 is the ground truth data set of fiber 2 from another mouse and contains 192
$Ca^{2+}$ sparks. The corresponding prediction for 4 rotations contains 524 $Ca^{2+}$ sparks.

The code for image translations was taken with minor modifications from
https://www.tensorflow.org/tutorials/generative/pix2pix and is based on the
publication of Isola et al[29]. The structure of the network was not modified. The
network was trained for 35 epochs, batch size of 1.

**Analysis of translated images**

As first analysis step of the translated images, fiber masks of the ground truth and
predicted $Ca^{2+}$ signal were generated in order to determine the fiber area and the
mean fluorescence intensity within the fiber. First, a gaussian filter was applied on the
enhanced $Ca^{2+}$ images. Then, a binary image of the mask was created. Pixels were
assigned the value 1 if their value was higher than 0.00001* σ, with σ the standard
deviation of the corresponding image. In the enhanced ground truth $Ca^{2+}$ image the
$Ca^{2+}$ signal outside the fiber was set to 0 as described previously in the image pre-

processing part. In the predicted $Ca^{2+}$ images some pixels with values higher than 0

could be rarely observed outside the fiber. To exclude these pixels, objects in the

binary images satisfying the condition 0.00001* σ were labelled. Then, objects

smaller than 3000 pixels were excluded and a cleaned binary image was generated.

Predicted $Ca^{2+}$ sparks were automatically identified as described in the image pre-

processing section but with fiber masks generated as described in this section

(analysis of translated images) since predicted images might show some variability of

the generated fibers. Properties of $Ca^{2+}$ sparks ($Ca^{2+}$ spark area, $Ca^{2+}$ spark mean

fluorescence intensity and the ratio $Ca^{2+}$ spark area/ $Ca^{2+}$ sparks edges) were

determined based on the labelled binary images of $Ca^{2+}$ sparks. $Ca^{2+}$ frequencies

(densities) were defined as the following ratio: number of $Ca^{2+}$ sparks in an

experiment/ overall fiber mask area in an experiment.

An algorithm was developed to automatically detect the fiber periphery. First, the

border of the image was detected using ndi.binary_delation of scipy. Then, pixels

touching the image border and being part of the fiber mask were assigned a value of

1. Then, using a fiber erosion (ndi.binary_erosion of scipy) over 40 steps an image

was iteratively created that shows the centre of the fiber. On each of the 40 steps the

images with pixels assigned a value 1 when touching the image border and being

part of the fiber mask were added to the eroded image in order to avoid the centre of

the fiber touching the image border being assigned to the fiber border (512x512 pixel

images Fig. 3a and b). With this eroded image and the fiber mask image an image of

the fiber periphery was created. If there were overlapping pixels of the fiber periphery

and the binary image of $Ca^{2+}$ sparks, the $Ca^{2+}$ spark was counted as a signal at the

fiber periphery.

For the measurement of the $Ca^{2+}$ spark area over myosin filaments, the mean SHG

signal over the whole measurement was used. First, a gaussian filter was applied on

the mean SHG signal. Then, a fiber mask was generated with pixels assigned the

value 1 if they fulfil the condition > σ, with σ the standard deviation of the mean SHG

signal. To create a binary image of the myosin filaments the module filters.rank.mean

of scikit image was applied on the mean SHG signal with a disk size of 5. The images

of the mean SHG signal were multiplied by the fiber mask. The $Ca^{2+}$ spark area over

the myosin filaments was calculated based on the binary image of the mean SHG

signal and the labelled binary image of $Ca^{2+}$ sparks.

393 Y-shaped structures were manually annotated on mean SHG measurements using

394 Fiji[52]. With these binary images of y-shaped structures and the labelled binary

395 images of $Ca^{2+}$ sparks, the $Ca^{2+}$ spark area over y-shaped structures was calculated.

396 If 30% of the $Ca^{2+}$ spark area overlaid y-shaped structures, it was counted as $Ca^{2+}$

397 spark at y-shaped structures and the corresponding $Ca^{2+}$ spark frequencies at y-

398 shaped structures were calculated for the three experiments with the corresponding

399 areas of y-shaped structures.

400 The ratios SHG signal(spark)/SHG signal(fiber) and Calcium signal(spark)/Calcium

401 signal(fiber) in Fig. 5 are the mean SHG signal or $Ca^{2+}$ signal at the region of a $Ca^{2+}$

402 spark divided by the mean SHG signal or mean $Ca^{2+}$ signal of the fiber mask when

403 regions of $Ca^{2+}$ sparks were excluded from the fiber mask.

404 **Statistics**

405 Pearson correlation coefficients and linear regression were calculated with Scipy[53],

406 Matthews correlation coefficients were calculated with Scikit learn[54], structural

407 similarity index (SSIM)[46] and mean squared error[47] (SME) were calculated with Scikit

408 image[55]. Violin plots and box plots were plotted with Matplotlib[56]. Boxes in boxplots

409 extend from Q1 to Q3 (with Q1 lower quartile and Q3 upper quartile), median is

410 indicated by the horizontal line, triangles indicate means, upper and lower whiskers

411 show last data point less than Q3 + 1.5*(Q3-Q1) and first data point greater than Q1-

412 1.5 (Q3-Q1) with (Q3-Q1) the interquartile range and the circles indicate datapoints

413 beyond the whiskers[56]. The F1 score was calculated as described by Moen et al[57].

414 **Software**

415 The following software was used in this paper: Python, Jupyter, Spyder,

416 Tensorflow[58], Tensorflow_io, Keras[59], Scikit learn[54], Scikit image[55], Numpy[60],

417 Matplotlib[56], Scipy[53], Tifffile[61], Os, Pathlib, Time, Datetime, IPython[62], Xlsxwriter,

418 ImageJ[63], Fiji[52], ChimeraX[64], https://www.tensorflow.org/tutorials/generative/pix2pix

419 and https://github.com/embl-bio-it/image-analysis-with-python/blob/master/session-

420 3to5/image_analysis_tutorial_solutions.ipynb.

421 **Code availability**

422 The code and the checkpoint used in this paper is provided at

423 https://github.com/TihomirGeorgiev3/SHGtoSpark_part_1 and

424 https://github.com/TihomirGeorgiev3/SHGtoSpark_part_2

**Data availability**

Data will be provided upon request. Some images will be provided at

https://github.com/TihomirGeorgiev3/SHGtoSpark_part_1 and

https://github.com/TihomirGeorgiev3/SHGtoSpark_part_2. The algorithms can be

tested on these images.

# References

Berridge, M. J., Bootman, M. D. & Roderick, H. L. Calcium signalling: dynamics, homeostasis and remodelling. *Nat Rev Mol Cell Biol* **4**, 517-529 (2003). https://doi.org:10.1038/nrm1155

Cheng, H., Lederer, W. J. & Cannell, M. B. Calcium sparks: elementary events underlying excitation-contraction coupling in heart muscle. *Science* **262**, 740-744 (1993). https://doi.org:10.1126/science.8235594

Cheng, H. & Lederer, W. J. Calcium sparks. *Physiol Rev* **88**, 1491-1545 (2008). https://doi.org:10.1152/physrev.00030.2007

Kirsch, W. G., Uttenweiler, D. & Fink, R. H. Spark- and ember-like elementary Ca2+ release events in skinned fibres of adult mammalian skeletal muscle. *J Physiol* **537**, 379-389 (2001). https://doi.org:10.1111/j.1469-7793.2001.00379.x

Wang, X. *et al.* Uncontrolled calcium sparks act as a dystrophic signal for mammalian skeletal muscle. *Nat Cell Biol* **7**, 525-530 (2005). https://doi.org:10.1038/ncb1254

Theis, L. *et al.* Benchmarking Spike Rate Inference in Population Calcium Imaging. *Neuron* **90**, 471-482 (2016). https://doi.org:10.1016/j.neuron.2016.04.014

Berens, P. *et al.* Community-based benchmarking improves spike rate inference from two-photon calcium imaging data. *PLoS Comput Biol* **14**, e1006157 (2018). https://doi.org:10.1371/journal.pcbi.1006157

Li, X. *et al.* Reinforcing neuron extraction and spike inference in calcium imaging using deep self-supervised denoising. *Nat Methods* **18**, 1395-1400 (2021). https://doi.org:10.1038/s41592-021-01225-0

Rupprecht, P. *et al.* A database and deep learning toolbox for noise-optimized, generalized spike inference from calcium imaging. *Nat Neurosci* **24**, 1324-1337 (2021). https://doi.org:10.1038/s41593-021-00895-5

Lecoq, J. *et al.* Removing independent noise in systems neuroscience data using DeepInterpolation. *Nat Methods* **18**, 1401-1408 (2021). https://doi.org:10.1038/s41592-021-01285-2

Sita, L. *et al.* A deep-learning approach for online cell identification and trace extraction in functional two-photon calcium imaging. *Nat Commun* **13**, 1529 (2022). https://doi.org:10.1038/s41467-022-29180-0

Li, X. *et al.* Real-time denoising enables high-sensitivity fluorescence time-lapse imaging beyond the shot-noise limit. *Nat Biotechnol* **41**, 282-292 (2023). https://doi.org:10.1038/s41587-022-01450-8

Goodfellow, I. *et al.* Generative Adversarial Nets. *Advances in Neural Information Processing Systems* **27** (2014).

Kamran, S. A. *et al.* New open-source software for subcellular segmentation and analysis of spatiotemporal fluorescence signals using deep learning. *iScience* **25**, 104277 (2022). https://doi.org:10.1016/j.isci.2022.104277

Yang, S. *et al.* Calcium Spark Detection and Event-Based Classification of Single Cardiomyocyte Using Deep Learning. *Front Physiol* **12**, 770051 (2021). https://doi.org:10.3389/fphys.2021.770051

Janicek, R. *et al.* Local recovery of cardiac calcium-induced calcium release interrogated by ultra-effective, two-photon uncaging of calcium. *J Physiol* **599**, 3841-3852 (2021). https://doi.org:10.1113/JP281482

Dotti, P. *et al.* A deep learning-based approach for efficient detection and classification of local Ca(2)(+) release events in Full-Frame confocal imaging. *Cell Calcium* **121**, 102893 (2024). https://doi.org:10.1016/j.ceca.2024.102893

This, S., Costantino, S. & Melichar, H. J. Machine learning predictions of T cell antigen specificity from intracellular calcium dynamics. *Sci Adv* **10**, eadk2298 (2024). https://doi.org:10.1126/sciadv.adk2298

19    Christiansen, E. M. *et al.* In Silico Labeling: Predicting Fluorescent Labels in Unlabeled Images.
*Cell* **173**, 792-803 e719 (2018). https://doi.org:10.1016/j.cell.2018.03.040
20    Ounkomol, C., Seshamani, S., Maleckar, M. M., Collman, F. & Johnson, G. R. Label-free
prediction of three-dimensional fluorescence images from transmitted-light microscopy. *Nat*
*Methods* **15**, 917-920 (2018). https://doi.org:10.1038/s41592-018-0111-2
21    Jayachandran Preetha, C. *et al.* Deep-learning-based synthesis of post-contrast T1-weighted
MRI for tumour response assessment in neuro-oncology: a multicentre, retrospective cohort
study. *Lancet Digit Health* **3**, e784-e794 (2021). https://doi.org:10.1016/S2589-
7500(21)00205-3
22    Yan, Y., Lee, H., Somer, E. & Grau, V. in *PRedictive Intelligence in MEdicine.* (eds Islem Rekik,
Gozde Unal, Ehsan Adeli, & Sang Hyun Park) 26-33 (Springer International Publishing) (2018).
23    Both, M. *et al.* Second harmonic imaging of intrinsic signals in muscle fibers in situ. *J Biomed*
*Opt* **9**, 882-892 (2004). https://doi.org:10.1117/1.1783354
24    Plotnikov, S. V., Millard, A. C., Campagnola, P. J. & Mohler, W. A. Characterization of the
myosin-based source for second-harmonic generation from muscle sarcomeres. *Biophys J* **90**,
693-703 (2006). https://doi.org:10.1529/biophysj.105.071555
25    Georgiev, T., Zapiec, B., Forderer, M., Fink, R. H. A. & Vogel, M. Colocalization properties of
elementary Ca(2+) release signals with structures specific to the contractile filaments and the
tubular system of intact mouse skeletal muscle fibers. *J Struct Biol* **192**, 366-375 (2015).
https://doi.org:10.1016/j.jsb.2015.09.018
26    Awasthi, S. *et al.* Multimodal SHG-2PF Imaging of Microdomain Ca2+-Contraction Coupling in
Live Cardiac Myocytes. *Circ Res* **118**, e19-28 (2016).
https://doi.org:10.1161/CIRCRESAHA.115.307919
27    Denk, W., Strickler, J. H. & Webb, W. W. Two-photon laser scanning fluorescence microscopy.
*Science* **248**, 73-76 (1990). https://doi.org:10.1126/science.2321027
28    Gatys, L. A., Ecker, A. S. & Bethge, M. A Neural Algorithm of Artistic Style. *CoRR*
**abs/1508.06576** (2015).
29    Isola, P., Zhu, J.-., Yan, Zhou, T. & Efros, A. A. Image-to-Image Translation with Conditional
Adversarial Networks. *CoRR* **abs/1611.07004** (2016).
30    Sun, F. *et al.* A Genetically Encoded Fluorescent Sensor Enables Rapid and Specific Detection
of Dopamine in Flies, Fish, and Mice. *Cell* **174**, 481-496 e419 (2018).
https://doi.org:10.1016/j.cell.2018.06.042
31    Patriarchi, T. *et al.* Ultrafast neuronal imaging of dopamine dynamics with designed
genetically encoded sensors. *Science* **360** (2018). https://doi.org:10.1126/science.aat4422
32    Patriarchi, T. *et al.* An expanded palette of dopamine sensors for multiplex imaging in vivo.
*Nat Methods* **17**, 1147-1155 (2020). https://doi.org:10.1038/s41592-020-0936-3
33    Sun, F. *et al.* Next-generation GRAB sensors for monitoring dopaminergic activity in vivo. *Nat*
*Methods* **17**, 1156-1166 (2020). https://doi.org:10.1038/s41592-020-00981-9
34    Jing, M. *et al.* A genetically encoded fluorescent acetylcholine indicator for in vitro and in vivo
studies. *Nat Biotechnol* **36**, 726-737 (2018). https://doi.org:10.1038/nbt.4184
35    Jing, M. *et al.* An optimized acetylcholine sensor for monitoring in vivo cholinergic activity.
*Nat Methods* **17**, 1139-1146 (2020). https://doi.org:10.1038/s41592-020-0953-2
36    Helassa, N. *et al.* Ultrafast glutamate sensors resolve high-frequency release at Schaffer
collateral synapses. *Proc Natl Acad Sci U S A* **115**, 5594-5599 (2018).
https://doi.org:10.1073/pnas.1720648115
37    Marvin, J. S. *et al.* Stability, affinity, and chromatic variants of the glutamate sensor iGluSnFR.
*Nat Methods* **15**, 936-939 (2018). https://doi.org:10.1038/s41592-018-0171-3
38    Marvin, J. S. *et al.* A genetically encoded fluorescent sensor for in vivo imaging of GABA. *Nat*
*Methods* **16**, 763-770 (2019). https://doi.org:10.1038/s41592-019-0471-2
39    Wu, Z. *et al.* A sensitive GRAB sensor for detecting extracellular ATP in vitro and in vivo.
*Neuron* **110**, 770-782 e775 (2022). https://doi.org:10.1016/j.neuron.2021.11.027
40    Wan, J. *et al.* A genetically encoded sensor for measuring serotonin dynamics. *Nat Neurosci*
**24**, 746-752 (2021). https://doi.org:10.1038/s41593-021-00823-7

41  Feng, J. *et al.* A Genetically Encoded Fluorescent Sensor for Rapid and Specific In Vivo
Detection of Norepinephrine. *Neuron* **102**, 745-761 e748 (2019).
https://doi.org:10.1016/j.neuron.2019.02.037
42  Peng, W. *et al.* Regulation of sleep homeostasis mediator adenosine by basal forebrain
glutamatergic neurons. *Science* **369** (2020). https://doi.org:10.1126/science.abb0556
43  Nucciotti, V. *et al.* Probing myosin structural conformation in vivo by second-harmonic
generation microscopy. *Proc Natl Acad Sci U S A* **107**, 7763-7768 (2010).
https://doi.org:10.1073/pnas.0914782107
44  Schurmann, S., von Wegner, F., Fink, R. H., Friedrich, O. & Vogel, M. Second harmonic
generation microscopy probes different states of motor protein interaction in myofibrils.
*Biophys J* **99**, 1842-1851 (2010). https://doi.org:10.1016/j.bpj.2010.07.005
45  Forderer, M., Georgiev, T., Mosqueira, M., Fink, R. H. & Vogel, M. Functional second
harmonic generation microscopy probes molecular dynamics with high temporal resolution.
*Biomed Opt Express* **7**, 525-541 (2016). https://doi.org:10.1364/BOE.7.000525
46  Wang, Z., Bovik, A. C., Sheikh, H. R. & Simoncelli, E. P. Image quality assessment: from error
visibility to structural similarity. *IEEE Transactions on Image Processing* **13**, 600-612 (2004).
47  Wang, Z. & Bovik, A. C. Mean squared error: Love it or leave it? A new look at Signal Fidelity
Measures. *IEEE Signal Processing Magazine* **26**, 98-117 (2009).
48  Georgiev, T., Svirin, M., Jaimovich, E. & Fink, R. H. Localized nuclear and perinuclear Ca(2+)
signals in intact mouse skeletal muscle fibers. *Front Physiol* **6**, 263 (2015).
https://doi.org:10.3389/fphys.2015.00263
49  Ronneberger, O., Fischer, P. & Brox, T. in *Medical Image Computing and Computer-Assisted
Intervention -- MICCAI 2015.* (eds Nassir Navab, Joachim Hornegger, William M. Wells, &
Alejandro F. Frangi) 234-241 (Springer International Publishing).
50  Dosovitskiy, A., Springenberg, J. T., Riedmiller, M. & Brox, T. Discriminative Unsupervised
Feature Learning with Convolutional Neural Networks. *Advances in Neural Information
Processing Systems* **27** (2014).
51  Steele, E. M. & Steele, D. S. Automated detection and analysis of Ca(2+) sparks in x-y image
stacks using a thresholding algorithm implemented within the open-source image analysis
platform ImageJ. *Biophys J* **106**, 566-576 (2014). https://doi.org:10.1016/j.bpj.2013.12.040
52  Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat Methods* **9**,
676-682 (2012). https://doi.org:10.1038/nmeth.2019
53  Virtanen, P. *et al.* SciPy 1.0: fundamental algorithms for scientific computing in Python. *Nat
Methods* **17**, 261-272 (2020). https://doi.org:10.1038/s41592-019-0686-2
54  Pedregosa, F. *et al.* Scikit-learn: Machine Learning in Python. *Journal of Machine Learning
Research* **12**, 2825-2830 (2011).
55  van der Walt, S. *et al.* scikit-image: image processing in Python. *PeerJ* **2**, e453 (2014).
https://doi.org:10.7717/peerj.453
56  Hunter, J. D. Matplotlib: A 2D Graphics Environment. *Computing in Science & Engineering*
(2007). https://doi.org:10.1109/MCSE.2007.55
57  Moen, E. *et al.* Deep learning for cellular image analysis. *Nat Methods* **16**, 1233-1246 (2019).
https://doi.org:10.1038/s41592-019-0403-1
58  Abadi, M. *et al.* TensorFlow: Large-Scale Machine Learning on Heterogeneous Systems.
(2015). https://doi.org:10.5281/zenodo.7764425
59  Chollet, F. & others. Keras.  (2015).
60  Harris, C. R. *et al.* Array programming with NumPy. *Nature* **585**, 357-362 (2020).
https://doi.org:10.1038/s41586-020-2649-2
61  Gohlke, C. Tifffile.  (2022). https://doi.org:10.5281/zenodo.6795861
62  Perez, F. & Granger, B. E. IPython: A System for Interactive Scientific Computing. *Computing
in Science & Engineering* **9**, 21-29 (2007).
63  Schneider, C. A., Rasband, W. S. & Eliceiri, K. W. NIH Image to ImageJ: 25 years of image
analysis. *Nat Methods* **9**, 671-675 (2012). https://doi.org:10.1038/nmeth.2089

64    Pettersen, E. F. *et al.* UCSF ChimeraX: Structure visualization for researchers, educators, and
developers. *Protein Sci* **30**, 70-82 (2021). https://doi.org:10.1002/pro.3943

**Extended Data**

***Extended Data Figure 1   Predicted Ca²⁺ signalling patterns. a** and **b**, Translations of two*

*consecutive SHG images in a fiber not included in the training data set (Exp 1). Green arrows show at*

*predicted Ca$^{2+}$ sparks that can be seen on both images. The red arrow indicates a predicted Ca$^{2+}$*

*spark that can be observed only on one of the images. Scale bar 20 µm. **c – f**, Translations of four*

*consecutive SHG images of another fiber in Exp 1. Green arrows indicate predicted Ca$^{2+}$ sparks seen*

*in all images. Red arrows show predicted Ca$^{2+}$ sparks seen only on two images and located more in*

*the centre of the fiber.*

***Extended Data Figure 2   Predicted Ca²⁺ sparks at the centre of the fiber.*** *Binary images of the*

*predicted fiber periphery (blue), predicted fiber centre (cyan), predicted Ca²⁺ sparks (green) and*

*ground truth Ca²⁺ sparks (red) of the four images shown in Extended Data Fig. 1c – f. In **a** and **d**, two*

*predicted Ca²⁺ sparks at the centre of the fiber can be observed. In **a** and **c**, two ground truth Ca²⁺*

*sparks at the centre of the fiber can be seen. The predicted and ground truth Ca²⁺ sparks are located*

*in close proximity. In **c**, a small part of one of the three ground truth Ca²⁺ sparks extends beyond the*

*predicted fiber mask. This was seen extremely rarely since there is a high agreement between*

*predicted and ground truth fiber masks. Scale bar 20 μm.*

**Extended Data Figure 3** **Ca²⁺ sparks at the fiber periphery. a,** *Fraction of Ca²⁺ sparks at the fiber*

*periphery when one of two fibers in Exp1 was excluded from the analysis. The excluded fiber showed*

*an unusually high Ca²⁺ spark frequency at the center of the fiber.* **b,** *Violin plots of the ratio of Ca²⁺*

*spark area over the fiber periphery and Ca²⁺ spark area for the three experiments (GT means ground*

*truth and P prediction) and the training data set. Green triangles are the means.*

**Extended Data Figure 4 Ca²⁺ spark frequencies at y-shaped structures. a – d**, $Ca^{2+}$ spark
frequencies for the three experiments. $Ca^{2+}$ sparks were counted as $Ca^{2+}$ sparks at y-shaped
structures if at least 20% (**a**), 40% (**b**), 50% (**c**) and 60% (**d**) of the $Ca^{2+}$ spark area covered y-shaped
structures.

***Extended Data Figure 5   Ca²⁺ sparks seen in all image translations of a measurement***

***associated with y-shaped structures. a**, SHG measurement of the fiber shown in Fig. 4a. Scale bar*

*10 μm. **b**, Overlay of the predicted Ca²⁺ signal (green) (first image in Fig. 4a) and the SHG signal*

*(red). The arrow indicates a predicted Ca²⁺ spark that could be seen in all image translations of this*

*measurement (149 images) and is located in close proximity to y-shaped structures. **c**, SHG*

*measurement of the fiber shown in Fig. 4b. Scale bar 10 μm. **d**, Overlay of the predicted Ca²⁺ signal*

*(green) (first image in Fig. 4b) and the SHG signal (red). The arrow shows a predicted Ca²⁺ spark that*

*could be seen in all image translations of this measurement (49 images) and is located at y-shaped*

*structures.*

*Extended Data Figure 6 **Properties of generated images with only one fiber in Exp1.** Only one fiber in Exp1 was included in the analyses presented in this figure. This fiber was shown in Fig. 2a, in Fig. 3a, b, d, f and in Fig. 4a, c. 1788 images out of 2180 images in Exp 1 are measurements of this fiber. **a** and **b**, Structural similarity index (SSIM) and mean squared error (MSE) of the three experiments. **c**, Pearson correlation coefficients of the overlapping fiber masks of predicted and ground truth images for the three experiments. **d** and **e**, Matthews correlation coefficients and F1 scores calculated based on ground truth and predicted fiber masks for the three experiments. **f**, Box plots and violin plots of fiber areas of ground truth (GT) and predicted (P) fiber masks for the three experiments. **g**, Box plots and violin plots of fiber mean intensity of ground truth (GT) and predicted (P) fiber masks for the three experiments and the training data set.*

**Extended Data Figure 7   Properties of predicted Ca²⁺ sparks with only one fiber in Exp1.** *Only one fiber in Exp1 was included in the analyses presented in this figure. This fiber was shown in Fig. 2a, in Fig. 3a, b, d, f and in Fig. 4a, c. 1788 images out of 2180 images in Exp 1 are measurements of*

this fiber. *a* and *b*, F1 scores and Matthews correlation coefficients for the three experiments calculated based on binary images of ground truth and predicted $Ca^{2+}$ sparks. *c* and *d*, Box plots and violin plots of $Ca^{2+}$ spark area and $Ca^{2+}$ spark mean intensity for the three experiments (GT means ground truth and P prediction) and the training data set. *e*, $Ca^{2+}$ spark frequencies of the three experiments and the training data set. The ground truth $Ca^{2+}$ spark frequency of Exp1 was 2.77 sparks/(10000*$\mu m^2$). The predicted $Ca^{2+}$ spark frequency of Exp1 was 2.46 sparks/(10000*$\mu m^2$). *f*, Box plots and violin plots of the ratios $Ca^{2+}$ spark area/$Ca^{2+}$ spark edge for the three experiments and the training data set. *g*, $Ca^{2+}$ spark frequencies at y-shaped structures for the three experiments. The ground truth $Ca^{2+}$ spark frequency at y-shaped structures of Exp1 was 6.37 sparks/(10000*$\mu m^2$). The predicted $Ca^{2+}$ spark frequency at y-shaped structures of Exp1 was 8.30 sparks/(10000*$\mu m^2$). *h*, Violin plots of $Ca^{2+}$ spark area over y-shaped structures/$Ca^{2+}$ spark area for the three experiments. *i*, Box plots and violin plots of the ratios $Ca^{2+}$ spark area over myosin filaments/$Ca^{2+}$ spark area for the three experiments and the training data set.

**Extended Data Figure 8   F1 scores for binary images of the Ca²⁺ fiber mask (a) and binary images of Ca²⁺ sparks (b) in Exp 1.** *Fiber 1 (mouse 1) measured without a band pass filter in front of the SHG signal detector (Exp 1 c1nf), fiber 1 (mouse 1) measured with a band pass filter in front of the SHG signal detector (Exp 1 c1) and fiber 2 (mouse 2) with a band pass filter in front of the SHG signal detector (Exp 1 c2).*

**Movie 1**

xyt volume shown in Fig. 4c. Box volume 34.88 µm, 34.88 µm, 122.18 s.

**Movie 2**

xyt volume shown in Fig. 4d. Box volume 34.88 µm, 34.88 µm, 40.18 s.