Prediction of fast subcellular Ca^{2+} signals based on subcellular structures in label-free second harmonic generation microscopy measurements

Tihomir Georgiev[1*]

[1]Department of Experimental Ophthalmology, University Hospital Cologne, University of Cologne, Cologne, Germany

[*]Correspondence to: tihomir.georgiev@mail.de

Abstract

Ca^{2+} sparks were predicted based on the label-free second harmonic generation (SHG) signal of myosin filaments with pix2pix, a conditional adversarial network. Predicted Ca^{2+} sparks and ground truth Ca^{2+} sparks show similar properties. Predicted Ca^{2+} spark frequencies and a predominant location at the periphery of skeletal muscle fibers were in good agreement with ground truth data. High frequencies of predicted Ca^{2+} sparks could be observed at y-shaped structures of myosin filaments. Important image pre-processing was applied in order to enhance the signal of Ca^{2+} sparks used for training. Only freely available software was applied. I provide the image pre-processing algorithm and several algorithms to evaluate the quality of predicted Ca^{2+} sparks. This technique can be extended to other label-free microscopy techniques and other fluorescent indicators developed during the last years. It remains a difficult task to predict the exact location of Ca^{2+} sparks.

Introduction

Ca^{2+} is an important intracellular signal[1]. Ca^{2+} sparks are fast and small Ca^{2+} signals in different cell types[2,3]. High frequencies of Ca^{2+} sparks can be observed in permeabilized skeletal muscle fibers[4] and during an osmotic treatment in intact skeletal muscle fibers[5]. Machine learning has been applied for the analysis of Ca^{2+} signals recently[6-12]. Generative adversarial networks have been introduced by Goodfellow et al[13] and have been applied for segmentation of Ca^{2+} signals[14]. Machine learning approaches for the analysis of Ca^{2+} sparks are rare[15]. Recently, machine learning has been applied for segmentation and classification of Ca^{2+} sparks[15,16].

In light microscopy, in silico labelling of subcellular structures based on bright field images has been achieved with deep learning[17,18]. Image translations of medical images have been generated with deep learning for different applications such as synthesis of post contrast MRI images[19] or translation of MRI images to PET images[20].

To my knowledge, this paper is the first attempt to translate structural images to images of Ca^{2+} signals. Second harmonic generation (SHG) microscopy is a nonlinear optical microscopy technique that allows the label-free visualization of myosin filaments in skeletal muscle fibers[21]. Localized Ca^{2+} signals have been investigated with respect to the myosin filaments in striated muscle cells previously [22,23]. In this study, I used the SHG signal of myosin filaments as subcellular structural information to generate synthetic images of Ca^{2+} signals. The training of the network was carried out on pairs of images consisting of an SHG measurement and a two photon microscopy[24] measurement of Ca^{2+} sparks in skeletal muscle fibers. For the training process of the network, an important image pre-processing of the Ca^{2+} signals was applied in order to enhance the signal of Ca^{2+} sparks compared to the cellular background. Different strategies have been applied to translate images previously[25,26]. I applied the pix2pix algorithm[26] for the translation of images which is based on conditional adversarial networks. In principle this technique can be extended to other label-free microscopy techniques and other fluorescent indicators such as dopamine[27-30], acetylcholine[31,32], glutamate[33,34], GABA[35], ATP[12,36], serotonin[37], norepinephrine[38] and adenosine[39] indicators. The prediction of Ca^{2+} sparks based on subcellular structures is a difficult task since the exact location of Ca^{2+} sparks is difficult to predict. Additionally, Ca^{2+} sparks are small and fast subcellular signals in a noisy cellular environment.

Results

Generation of translated images

As a first step, Ca^{2+} sparks were enhanced compared to the background signal of the cell (Fig. 1a). Then, the network was trained on pairs of SHG images and enhanced Ca^{2+} images (Fig. 1b). The trained network was then applied to translate Ca^{2+} images based on SHG images (Fig. 1c). Fig. 1c shows only one Ca^{2+} spark in the translated image. Some translated Ca^{2+} sparks could be seen on every image of the

measurement. Other translated Ca²⁺ sparks appeared only on some images of the

Figure 1 *Generation of translated images. a, Pre-processing of Ca²⁺ sparks with enhancement of Ca²⁺ sparks compared to the background signal of the fiber. On the left side Ca²⁺ sparks (arrows)*

measurement (Extended Data Fig. 1b, c, f). On several translated images two, three or four Ca^{2+} sparks could be observed (Extended Data Fig. 1a, b, c, f). Ca^{2+} sparks could be found at subcellular structures of the myosin pattern, so called y-shaped structures (Fig. 1d). Structural similarity index (SSIM)[40] and mean squared error (SME)[41] were calculated (Fig. 1e and f). The translated images achieved high SSIM values.

Properties of translated Ca^{2+} images

As a next step, I investigated several properties of translated images. The fiber masks of ground truth and predicted images were similar (Fig. 2a, b, c and d). The mean fiber intensity was lower in the predicted images, this was particularly the case in experiment 1 (Exp 1) (Fig. 2e). However, the mean of the fiber mean fluorescence intensity in the ground truth data of Exp 1 was higher than in the other two experiments and in the training data set.

Then, I investigated the properties of predicted Ca^{2+} sparks compared to ground truth Ca^{2+} sparks. I found that predicted Ca^{2+} spark area in Exp 1 was in good agreement with ground truth Ca^{2+} spark area (Fig. 2f). Interestingly, larger differences could be observed for the other two experiments. The predicted spark mean intensity was lower in all experiments (Fig. 2g). This was most pronounced in Exp 1. The predicted ratio Ca^{2+} spark area/ Ca^{2+} spark edge was larger than this ratio in the ground truth data for all experiments suggesting that predicted Ca^{2+} sparks are more circular (Fig. 2h).

Figure 2 Properties of translated images of Ca^{2+} sparks. a, Ground truth (red) and predicted (green) fiber mask of a fiber unseen during training of the network (Exp 1). The overlay of both masks is yellow. Scale bar 20 µm. **b** and **c**, F1 scores and Matthews correlation coefficients for the evaluation of predicted fiber masks for the three experiments. **d**, **e**, **f**, **g** and **h**, Box plots and violin plots of the fiber area, fiber mean intensity, spark area, spark mean intensity and spark area/spark edge of the ground truth (Exp GT) and predicted (Exp P) images in the three experiments and the training data set. Orange lines are the medians of the data and green triangles are the means. **i**, Ca^{2+} spark frequencies of the ground truth and predicted images of the three experiments and the training data set.

The predicted Ca^{2+} spark frequency in Exp 1 of 2.52 sparks/(10000*µm^2) was in good agreement with the ground truth Ca^{2+} spark frequency of 3.05 sparks/(10000*µm^2) (Fig. 2i). The predicted Ca^{2+} spark frequency in experiment 3 (Exp 3) of 2.78 sparks/(10000*µm^2) was even closer to the corresponding ground truth Ca^{2+} spark frequency of 2.74 sparks/(10000*µm^2). The training data set had a Ca^{2+} spark frequency of 1.85 sparks/(10000*µm^2). However, there was a larger difference in experiment 2 (Exp 2) between the predicted Ca^{2+} spark frequency of 2.46 sparks/(10000*µm^2) and the ground truth Ca^{2+} spark frequency of 4.36 sparks/(10000*µm^2).

Translation of subcellular structures to fast subcellular signals

As a next step, I investigated specific subcellular regions of the muscle fiber known to have high frequencies of Ca^{2+} sparks, the fiber periphery[5] and y-shaped structures of the myosin pattern[22]. Additionally, I investigated the ratio Ca^{2+} spark area over myosin filaments/spark area.

As expected, most of the predicted and ground truth Ca^{2+} sparks were located at the fiber periphery (Fig. 3a, b, and c). The ground truth Ca^{2+} spark frequency at the fiber periphery in Exp 1 showed a lower fraction of Ca^{2+} sparks compared to the predicted fraction. This difference can be also seen in the ratio Ca^{2+} spark area over fiber periphery/spark area (Extended Data Fig. 3b). One of the fibers used in Exp 1 showed a high frequency of Ca^{2+} sparks in the centre of the fiber not seen for the other five fibers (Extended Data Fig. 2). Interestingly, predicted images of this fiber showed several Ca^{2+} sparks in the fiber centre in close proximity to the ground truth Ca^{2+} sparks but not at such high frequencies (Extended Data Fig. 2). If this fiber was

excluded from the analysis, there is a good agreement between ground truth data and predicted data (Extended Data Fig. 3a).

Figure 3 ***Translation of subcellular structures to fast subcellular signals. a,*** *Binary images of the fiber centre (cyan), periphery of the fiber (blue) and Ca²⁺ sparks (green) based on a predicted Ca²⁺ image of a fiber unseen during training of the algorithm (Exp 1). Additionally, a binary image of a Ca²⁺ spark measured in the ground truth image is presented (red). Scale bar 20 µm.* ***b,*** *Overlay of the predicted Ca²⁺ signal (green) of the fiber shown in* ***a*** *and fiber periphery (blue). The arrows indicate Ca²⁺ sparks.* ***c,*** *Percentage of Ca²⁺ sparks at the fiber periphery for ground truth (Exp GT) and predicted (Exp P) images in the three experiments and the training data set.* ***d,*** *Overlay of binary images (left) of the myosin filaments (red) and a predicted Ca²⁺ spark. The binary SHG signal is based on a mean SHG signal of the whole measurement. Overlay of the mean SHG signal of the whole measurement (red) and the predicted Ca²⁺ signal (green) of the same image (right). This fiber was unseen during the training process of the algorithm (Exp 1). Scale bar 10 µm.* ***e,*** *Box plots and violin plots of the ratios of the Ca²⁺ spark area over myosin filaments and the Ca²⁺ spark area for ground truth (Exp GT) and predicted (Exp P) images in the three experiments and the training data set. Orange lines are the medians of the data and green triangles are the means.* ***f,*** *Overlay of the mean*

Then, I looked at the ratio Ca^{2+} spark area over myosin filaments/spark area (Fig. 3d and e) and found that this ratio was lower for predictions. Several predicted Ca^{2+} sparks were found at regions without myosin filaments (Exp 1 P and Exp 2 P). Such Ca^{2+} sparks at myosin free regions can be found in the training data set.

As a next step, I investigated the translation of y-shaped structures of the myosin filament pattern to Ca^{2+} sparks (Fig. 3f, g and h and Extended Data Fig. 4). In Exp 1, the predicted Ca^{2+} spark frequency at y-shaped structures of 7.94 sparks/(10000*µm^2) was 1.26 times higher than the ground truth Ca^{2+} spark frequency at y-shaped structures of 6.31 sparks/(10000*µm^2). In Exp 2, no Ca^{2+} sparks were predicted at y-shaped structures while ground truth Ca^{2+} frequency at y-shaped structures was 3.92 sparks/(10000*µm^2). In Exp 3, the predicted Ca^{2+} spark frequency at y-shaped structures of 13.93 sparks/(10000*µm^2) was 1.56 times higher than the ground truth Ca^{2+} spark frequency at y-shaped structures of 8.95 sparks/(10000*µm^2).

Accuracy of Ca^{2+} spark prediction

Predicted Ca^{2+} sparks seen in all image translations of a measurement were several times associated with high ground truth Ca^{2+} spark activity in close proximity (Fig. 4a, b, c and d, Extended Data Movie 1 and 2). The predicted and ground truth Ca^{2+} sparks partly overlapped in these cases. However, the exact overlap was not high. This is reflected in low Pearson correlation coefficients, Matthews correlation coefficients and F1 scores (Fig. 4 e, f, g). The predicted Ca^{2+} sparks seen in all image translations of a measurement were in close proximity or at y-shaped structures of the myosin filaments (Extended Data Fig. 5).

Figure 4   Accuracy of Ca²⁺ spark prediction. a, *Image sequence (10 consecutive images) of the ground truth Ca²⁺ signal (red) and the predicted Ca²⁺ signal (green). The yellow arrow indicates a predicted Ca²⁺ spark. In close proximity to the predicted Ca²⁺ spark ground truth Ca²⁺ sparks can be observed. Predicted and ground truth Ca²⁺ sparks overlap partly sometimes. This fiber was unseen during the training process (Exp 1). Scale bar 10 μm.* **b**, *Image sequence of another fiber. Other measurements of this fiber and this plane were seen during the training process (Exp 3). The yellow arrow indicates a predicted Ca²⁺ spark (green) that is close to several ground truth Ca²⁺ sparks (red) and on several images they partly overlap. The green arrow shows at a predicted Ca²⁺ spark that is not in close proximity to ground truth signals. The red arrow indicates a ground truth Ca²⁺ spark that is not in close proximity to predicted Ca²⁺ sparks. Scale bar 10 μm.* **c** *and* **d**, *xyt volumes of the measurements partly shown as image sequences in* **a** *and* **b**. *Yellow arrows indicate predicted Ca²⁺ sparks in close proximity to many ground truth Ca²⁺ sparks partly overlapping with them. Red arrows show at ground truth Ca²⁺ sparks with no corresponding predicted Ca²⁺ sparks at this location. Green arrow shows at predicted Ca²⁺ sparks with no corresponding ground truth Ca²⁺ sparks at this location. Box volume 34.88 μm, 34.88 μm, 122.18 s in* **c** *and 34.88 μm, 34.88 μm, 40.18 s in* **d**. **e**, *Pearson correlation coefficients of the whole images (light blue) and of the overlapping fiber masks of predicted and ground truth images (dark blue) for the three experiments.* **f** *and* **g**, *Matthews correlation coefficients and F1 scores calculated based on binary images of ground truth Ca²⁺ sparks and predicted Ca²⁺ sparks for the three experiments.*

Extended Data Fig. 6 and 7 present the results when only one fiber was included in Exp1.

Discussion

Several properties of Ca²⁺ signals predicted based on label-free SHG measurements were in good agreement with the ground truth Ca²⁺ signals. Predictions of fast subcellular signals based on measurements of label-free techniques have the potential to make obsolete elaborate staining of cells or one additional channel is then available for another fluorescent marker. The frequency of Ca²⁺ sparks, the fiber area and the Ca²⁺ spark area (in Exp 1) could be predicted quite well. Ca²⁺ sparks at myosin free regions might be nuclear Ca²⁺ sparks[42]. The poor prediction of Ca²⁺ sparks in Exp 2 suggests that a different plane of the same fiber is already a different scene with different myosin pattern and associated Ca²⁺ sparks signalling pattern. As expected, most predicted Ca²⁺ sparks were located at the fiber periphery.

Predicted Ca²⁺ sparks seen in all images of a measurement were in several cases associated with locations of high ground truth Ca²⁺ sparks activity. The prediction of the exact location of Ca²⁺ sparks based on subcellular structures remains a difficult

task. In further improvement of the algorithm the 3D nature of xyt volumes should be taken into account instead of translating 2D images. Other existing algorithms such as U-net[43] could be tested for such a task as it was done in an image translation study of medical images[19]. It can be expected that data augmentation of the training data set[12,43,44], pre-processing of the SHG signal, denoising, larger data sets, powerful computational resources and algorithms designed for this specific task will further improve the accuracy of such image translations.

I hope that this work will stimulate further research to improve the translation of fast subcellular signals based on subcellular structures or other subcellular signals.

Methods

Data

The SHG images and two photon microscopy images used to train and test the algorithm were measured simultaneously in a previous publication[22].

Image pre-processing

The first step of the image pre-processing was to automatically identify Ca^{2+} sparks. Similar ideas for automatic detection of Ca^{2+} sparks have been described in a previous publication[45]. First, a gaussian filter was applied on each xyt measurement. Then, based on the gaussian signal image the mean intensity was calculated along the t axis, a fiber mask was generated using the mean image with a threshold of $2^*\sigma$, with σ the standard deviation. The mask was labelled and objects smaller than 3000 pixels were excluded. Then, a cleaned binary fiber mask was created. To exclude labelled structures outside the fiber, the Ca^{2+} measurement after application of a gaussian filter was multiplicated by the cleaned binary fiber mask. Binary image that includes the Ca^{2+} sparks was generated. Pixels were assigned a value of 1 if their value is higher than $\mu + 3.6^* \sigma$, with μ the mean and σ the standard deviation. μ and σ were measured within the fiber mask. Then objects of the binary images were labelled and objects smaller than 55 pixels were excluded. Then again, a binary image of Ca^{2+} sparks was created based on the cleaned labelled image.

Finally, the Ca^{2+} sparks in the original image were enhanced 10 times compared to the fiber background by reducing the fiber background 10 times. This step was important in order to make it easier for the network to see the Ca^{2+} sparks.

Translation of Ca^{2+} sparks based on SHG images

19 measurements of 4 fibers from two mice were used to train the algorithm. The training was carried out with 4123 images containing 256x256 pixels with 818 Ca^{2+} sparks. Thereby, 1031 images containing 512x512 pixels were cropped to 4123 256x256 pixel images.

To test the performance of the trained network, three different experiments Exp 1 (experiment 1), Exp 2 (experiment 2), Exp 3 (experiment 3) were carried out.

Exp 1 consisted of 545 512x512 pixel images from 5 measurements of two fibers from two mice, which were not included in the training data. 683 ground truth Ca^{2+} sparks were included in this data set.

Exp 2 consisted of 196 512x512 pixel images from 4 measurements of a fiber that was included in the training data but showing a different plane. This data set contained 292 ground truth Ca^{2+} sparks.

Exp 3 consisted of 98 512x512 pixel images from two measurements of a fiber that was included at the same plane in the training data. The data set contained 102 ground truth Ca^{2+} sparks.

Since there is a high variability of Ca^{2+} sparks signalling patterns and because no augmentation was carried out during the training process, images in Exp 1 and Exp 2 were rotated at 90°, 180°, 270° and 360°. Therefore, the algorithm was tested on 2180 512x512 pixel images in Exp1 and 784 512x512 pixel images in Exp 2.

All images in the three experiments were cropped to 256x256 pixel images as input for the image translation algorithm.

The code for image translations was taken with minor modifications from https://www.tensorflow.org/tutorials/generative/pix2pix and is based on the publication of Isola et al[26]. The structure of the network was not modified. The network was trained for 35 epochs, batch size of 1.

Analysis of translated images

As first analysis step of the translated images, fiber masks of the ground truth and predicted Ca^{2+} signal were generated in order to determine the fiber area and the mean fluorescence intensity within the fiber. First, a gaussian filter was applied on the

enhanced Ca^{2+} images. Then, a binary image of the mask was created. Pixels were assigned the value 1 if their value was higher than 0.00001* σ, with σ the standard deviation of the corresponding image. In the enhanced ground truth Ca^{2+} image the Ca^{2+} signal outside the fiber was set to 0 as described previously in the image pre-processing part. In the predicted Ca^{2+} images some pixels with values higher than 0 could be rarely observed outside the fiber. To exclude these pixels, objects in the binary images satisfying the condition 0.00001* σ were labelled. Then, objects smaller than 3000 pixels were excluded and a cleaned binary image was generated.

Predicted Ca^{2+} sparks were automatically identified as described in the image pre-processing section but with fiber masks generated as described in this section (analysis of translated images) since predicted images might show some variability of the generated fibers. Properties of Ca^{2+} sparks (Ca^{2+} spark area, Ca^{2+} spark mean fluorescence intensity and the ratio Ca^{2+} spark area/ Ca^{2+} sparks edges) were determined based on the labelled binary images of Ca^{2+} sparks. Ca^{2+} frequencies (densities) were defined as the following ratio: number of Ca^{2+} sparks in an experiment/ overall fiber mask area in an experiment.

An algorithm was developed to automatically detect the fiber periphery. First, the border of the image was detected using ndi.binary_delation of scipy. Then, pixels touching the image border and being part of fiber mask were assigned a value of 1. Then, using a fiber erosion (ndi.binary_erosion of scipy) over 40 steps an image was iteratively created that shows the centre of the fiber. On each of the 40 steps the images with pixels assigned a value 1 when touching the image border and being part of the fiber mask were added to the eroded image in order to avoid the centre of the fiber touching the image border being assigned to the fiber border (512x512 pixel images Fig. 3a and b). With this eroded image and the fiber mask image an image of the fiber periphery was created. If there were overlapping pixels of the fiber periphery and the binary image of Ca^{2+} sparks, the Ca^{2+} spark was counted as a signal at the fiber periphery.

For the measurement of the Ca^{2+} spark area over myosin filaments, the mean SHG signal over the whole measurement was used. First, a gaussian filter was applied on the mean SHG signal. Then, a fiber mask was generated with pixels assigned the value 1 if they fulfil the condition > σ, with σ the standard deviation of the mean SHG signal. To create a binary image of the myosin filaments the module filters.rank.mean

of scikit image was applied on the mean SHG signal with a disk size of 5. The images of the mean SHG signal were multiplied by the fiber mask. The Ca^{2+} spark area over the myosin filaments was calculated based on the binary image of the mean SHG signal and the labelled binary image of Ca^{2+} sparks.

Y-shaped structures were manually annotated on mean SHG measurements using Fiji[46]. With these binary images of y-shaped structures and the labelled binary images of Ca^{2+} sparks, the Ca^{2+} spark area over y-shaped structures was calculated. If 30% of the Ca^{2+} spark area overlaid y-shaped structures, it was counted as Ca^{2+} spark at y-shaped structures and the corresponding Ca^{2+} spark frequencies at y-shaped structures were calculated for the three experiments with the corresponding areas of y-shaped structures.

Statistics

Pearson correlation coefficients were calculated with Scipy[47], Matthews correlation coefficients were calculated with Scikit learn[48], structural similarity index (SSIM)[40] and mean squared error[41] (SME) were calculated with Scikit image[49]. Violin plots and box plots were plotted with Matplotlib[50]. Boxes in boxplots extend from Q1 to Q3 (with Q1 lower quartile and Q3 upper quartile), median is indicated by the horizontal line, triangles indicate means, upper and lower whiskers show last data point less than Q3 + 1.5*(Q3-Q1) and first data point greater than Q1- 1.5 (Q3-Q1) with (Q3-Q1) the interquartile range and the circles indicate datapoints beyond the whiskers[50]. The F1 score was calculated as described by Moen et al[51].

Software

The following software was used in this paper: Python, Jupyter, Spyder, Tensorflow[52], Tensorflow_io, Keras[53], Scikit learn[48], Scikit image[49], Numpy[54], Matplotlib[50], Scipy[47], Tifffile[55], Os, Pathlib, Time, Datetime, IPython[56], Xlsxwriter, ImageJ[57], Fiji[46], ChimeraX[58] and https://github.com/embl-bio-it/image-analysis-with-python/blob/master/session-3to5/image_analysis_tutorial_solutions.ipynb.

Code availability

The code and the checkpoint used in this paper will be provided at https://github.com/TihomirGeorgiev3.

Data availability

Data will be provided upon request. Some images will be provided at https://github.com/TihomirGeorgiev3. The algorithms can be tested on these images.

References

Berridge, M. J., Bootman, M. D. & Roderick, H. L. Calcium signalling: dynamics, homeostasis and remodelling. *Nat Rev Mol Cell Biol* **4**, 517-529 (2003). https://doi.org:10.1038/nrm1155

Cheng, H., Lederer, W. J. & Cannell, M. B. Calcium sparks: elementary events underlying excitation-contraction coupling in heart muscle. *Science* **262**, 740-744 (1993). https://doi.org:10.1126/science.8235594

Cheng, H. & Lederer, W. J. Calcium sparks. *Physiol Rev* **88**, 1491-1545 (2008). https://doi.org:10.1152/physrev.00030.2007

Kirsch, W. G., Uttenweiler, D. & Fink, R. H. Spark- and ember-like elementary Ca2+ release events in skinned fibres of adult mammalian skeletal muscle. *J Physiol* **537**, 379-389 (2001). https://doi.org:10.1111/j.1469-7793.2001.00379.x

Wang, X. *et al.* Uncontrolled calcium sparks act as a dystrophic signal for mammalian skeletal muscle. *Nat Cell Biol* **7**, 525-530 (2005). https://doi.org:10.1038/ncb1254

Theis, L. *et al.* Benchmarking Spike Rate Inference in Population Calcium Imaging. *Neuron* **90**, 471-482 (2016). https://doi.org:10.1016/j.neuron.2016.04.014

Berens, P. *et al.* Community-based benchmarking improves spike rate inference from two-photon calcium imaging data. *PLoS Comput Biol* **14**, e1006157 (2018). https://doi.org:10.1371/journal.pcbi.1006157

Li, X. *et al.* Reinforcing neuron extraction and spike inference in calcium imaging using deep self-supervised denoising. *Nat Methods* **18**, 1395-1400 (2021). https://doi.org:10.1038/s41592-021-01225-0

Rupprecht, P. *et al.* A database and deep learning toolbox for noise-optimized, generalized spike inference from calcium imaging. *Nat Neurosci* **24**, 1324-1337 (2021). https://doi.org:10.1038/s41593-021-00895-5

Lecoq, J. *et al.* Removing independent noise in systems neuroscience data using DeepInterpolation. *Nat Methods* **18**, 1401-1408 (2021). https://doi.org:10.1038/s41592-021-01285-2

Sita, L. *et al.* A deep-learning approach for online cell identification and trace extraction in functional two-photon calcium imaging. *Nat Commun* **13**, 1529 (2022). https://doi.org:10.1038/s41467-022-29180-0

Li, X. *et al.* Real-time denoising enables high-sensitivity fluorescence time-lapse imaging beyond the shot-noise limit. *Nat Biotechnol* **41**, 282-292 (2023). https://doi.org:10.1038/s41587-022-01450-8

Goodfellow, I. *et al.* Generative Adversarial Nets. *Advances in Neural Information Processing Systems* **27** (2014).

Kamran, S. A. *et al.* New open-source software for subcellular segmentation and analysis of spatiotemporal fluorescence signals using deep learning. *iScience* **25**, 104277 (2022). https://doi.org:10.1016/j.isci.2022.104277

Yang, S. *et al.* Calcium Spark Detection and Event-Based Classification of Single Cardiomyocyte Using Deep Learning. *Front Physiol* **12**, 770051 (2021). https://doi.org:10.3389/fphys.2021.770051

Janicek, R. *et al.* Local recovery of cardiac calcium-induced calcium release interrogated by ultra-effective, two-photon uncaging of calcium. *J Physiol* **599**, 3841-3852 (2021). https://doi.org:10.1113/JP281482

Christiansen, E. M. *et al.* In Silico Labeling: Predicting Fluorescent Labels in Unlabeled Images. *Cell* **173**, 792-803 e719 (2018). https://doi.org:10.1016/j.cell.2018.03.040

Ounkomol, C., Seshamani, S., Maleckar, M. M., Collman, F. & Johnson, G. R. Label-free prediction of three-dimensional fluorescence images from transmitted-light microscopy. *Nat Methods* **15**, 917-920 (2018). https://doi.org:10.1038/s41592-018-0111-2

Jayachandran Preetha, C. *et al.* Deep-learning-based synthesis of post-contrast T1-weighted MRI for tumour response assessment in neuro-oncology: a multicentre, retrospective cohort

study. *Lancet Digit Health* **3**, e784-e794 (2021). https://doi.org/10.1016/S2589-7500(21)00205-3

Yan, Y., Lee, H., Somer, E. & Grau, V. in *PRedictive Intelligence in MEdicine.* (eds Islem Rekik, Gozde Unal, Ehsan Adeli, & Sang Hyun Park) 26-33 (Springer International Publishing) (2018).

Both, M. *et al.* Second harmonic imaging of intrinsic signals in muscle fibers in situ. *J Biomed Opt* **9**, 882-892 (2004). https://doi.org:10.1117/1.1783354

Georgiev, T., Zapiec, B., Forderer, M., Fink, R. H. A. & Vogel, M. Colocalization properties of elementary Ca(2+) release signals with structures specific to the contractile filaments and the tubular system of intact mouse skeletal muscle fibers. *J Struct Biol* **192**, 366-375 (2015). https://doi.org:10.1016/j.jsb.2015.09.018

Awasthi, S. *et al.* Multimodal SHG-2PF Imaging of Microdomain Ca2+-Contraction Coupling in Live Cardiac Myocytes. *Circ Res* **118**, e19-28 (2016). https://doi.org:10.1161/CIRCRESAHA.115.307919

Denk, W., Strickler, J. H. & Webb, W. W. Two-photon laser scanning fluorescence microscopy. *Science* **248**, 73-76 (1990). https://doi.org:10.1126/science.2321027

Gatys, L. A., Ecker, A. S. & Bethge, M. A Neural Algorithm of Artistic Style. *CoRR* **abs/1508.06576** (2015).

Isola, P., Zhu, J.-., Yan, Zhou, T. & Efros, A. A. Image-to-Image Translation with Conditional Adversarial Networks. *CoRR* **abs/1611.07004** (2016).

Sun, F. *et al.* A Genetically Encoded Fluorescent Sensor Enables Rapid and Specific Detection of Dopamine in Flies, Fish, and Mice. *Cell* **174**, 481-496 e419 (2018). https://doi.org:10.1016/j.cell.2018.06.042

Patriarchi, T. *et al.* Ultrafast neuronal imaging of dopamine dynamics with designed genetically encoded sensors. *Science* **360** (2018). https://doi.org:10.1126/science.aat4422

Patriarchi, T. *et al.* An expanded palette of dopamine sensors for multiplex imaging in vivo. *Nat Methods* **17**, 1147-1155 (2020). https://doi.org:10.1038/s41592-020-0936-3

Sun, F. *et al.* Next-generation GRAB sensors for monitoring dopaminergic activity in vivo. *Nat Methods* **17**, 1156-1166 (2020). https://doi.org:10.1038/s41592-020-00981-9

Jing, M. *et al.* A genetically encoded fluorescent acetylcholine indicator for in vitro and in vivo studies. *Nat Biotechnol* **36**, 726-737 (2018). https://doi.org:10.1038/nbt.4184

Jing, M. *et al.* An optimized acetylcholine sensor for monitoring in vivo cholinergic activity. *Nat Methods* **17**, 1139-1146 (2020). https://doi.org:10.1038/s41592-020-0953-2

Helassa, N. *et al.* Ultrafast glutamate sensors resolve high-frequency release at Schaffer collateral synapses. *Proc Natl Acad Sci U S A* **115**, 5594-5599 (2018). https://doi.org:10.1073/pnas.1720648115

Marvin, J. S. *et al.* Stability, affinity, and chromatic variants of the glutamate sensor iGluSnFR. *Nat Methods* **15**, 936-939 (2018). https://doi.org:10.1038/s41592-018-0171-3

Marvin, J. S. *et al.* A genetically encoded fluorescent sensor for in vivo imaging of GABA. *Nat Methods* **16**, 763-770 (2019). https://doi.org:10.1038/s41592-019-0471-2

Wu, Z. *et al.* A sensitive GRAB sensor for detecting extracellular ATP in vitro and in vivo. *Neuron* **110**, 770-782 e775 (2022). https://doi.org:10.1016/j.neuron.2021.11.027

Wan, J. *et al.* A genetically encoded sensor for measuring serotonin dynamics. *Nat Neurosci* **24**, 746-752 (2021). https://doi.org:10.1038/s41593-021-00823-7

Feng, J. *et al.* A Genetically Encoded Fluorescent Sensor for Rapid and Specific In Vivo Detection of Norepinephrine. *Neuron* **102**, 745-761 e748 (2019). https://doi.org:10.1016/j.neuron.2019.02.037

Peng, W. *et al.* Regulation of sleep homeostasis mediator adenosine by basal forebrain glutamatergic neurons. *Science* **369** (2020). https://doi.org:10.1126/science.abb0556

Wang, Z., Bovik, A. C., Sheikh, H. R. & Simoncelli, E. P. Image quality assessment: from error visibility to structural similarity. *IEEE Transactions on Image Processing* **13**, 600-612 (2004).

Wang, Z. & Bovik, A. C. Mean squared error: Love it or leave it? A new look at Signal Fidelity Measures. *IEEE Signal Processing Magazine* **26**, 98-117 (2009).

Georgiev, T., Svirin, M., Jaimovich, E. & Fink, R. H. Localized nuclear and perinuclear Ca(2+) signals in intact mouse skeletal muscle fibers. *Front Physiol* **6**, 263 (2015). https://doi.org:10.3389/fphys.2015.00263

Ronneberger, O., Fischer, P. & Brox, T. in *Medical Image Computing and Computer-Assisted Intervention -- MICCAI 2015.* (eds Nassir Navab, Joachim Hornegger, William M. Wells, & Alejandro F. Frangi) 234-241 (Springer International Publishing).

Dosovitskiy, A., Springenberg, J. T., Riedmiller, M. & Brox, T. Discriminative Unsupervised Feature Learning with Convolutional Neural Networks. *Advances in Neural Information Processing Systems* **27** (2014).

Steele, E. M. & Steele, D. S. Automated detection and analysis of Ca(2+) sparks in x-y image stacks using a thresholding algorithm implemented within the open-source image analysis platform ImageJ. *Biophys J* **106**, 566-576 (2014). https://doi.org:10.1016/j.bpj.2013.12.040

Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat Methods* **9**, 676-682 (2012). https://doi.org:10.1038/nmeth.2019

Virtanen, P. *et al.* SciPy 1.0: fundamental algorithms for scientific computing in Python. *Nat Methods* **17**, 261-272 (2020). https://doi.org:10.1038/s41592-019-0686-2

Pedregosa, F. *et al.* Scikit-learn: Machine Learning in Python. *Journal of Machine Learning Research* **12**, 2825-2830 (2011).

van der Walt, S. *et al.* scikit-image: image processing in Python. *PeerJ* **2**, e453 (2014). https://doi.org:10.7717/peerj.453

Hunter, J. D. Matplotlib: A 2D Graphics Environment. *Computing in Science & Engineering* (2007). https://doi.org:10.1109/MCSE.2007.55

Moen, E. *et al.* Deep learning for cellular image analysis. *Nat Methods* **16**, 1233-1246 (2019). https://doi.org:10.1038/s41592-019-0403-1

Abadi, M. *et al.* TensorFlow: Large-Scale Machine Learning on Heterogeneous Systems. (2015). https://doi.org:10.5281/zenodo.7764425

Chollet, F. & others. Keras. (2015).

Harris, C. R. *et al.* Array programming with NumPy. *Nature* **585**, 357-362 (2020). https://doi.org:10.1038/s41586-020-2649-2

Tifffile (https://github.com/cgohlke/tifffile, 2022).

Perez, F. & Granger, B. E. IPython: A System for Interactive Scientific Computing. *Computing in Science & Engineering* **9**, 21-29 (2007).

Schneider, C. A., Rasband, W. S. & Eliceiri, K. W. NIH Image to ImageJ: 25 years of image analysis. *Nat Methods* **9**, 671-675 (2012). https://doi.org:10.1038/nmeth.2089

Pettersen, E. F. *et al.* UCSF ChimeraX: Structure visualization for researchers, educators, and developers. *Protein Sci* **30**, 70-82 (2021). https://doi.org:10.1002/pro.3943

Extended Data

Extended Data Figure 1   Predicted Ca²⁺ signalling patterns. a *and **b**, Translations of two consecutive SHG images in a fiber not included in the training data set (Exp 1). Green arrows show at*

predicted Ca^{2+} sparks that can be seen on both images. The red arrow indicates a predicted Ca^{2+} sparks that can be observed only on one of the images. Scale bar 20 µm. **c – f**, Translations of four consecutive SHG images of another fiber in Exp 1. Green arrows indicate predicted Ca^{2+} sparks seen in all images. Red arrows show predicted Ca^{2+} sparks seen only on two images and located more in the centre of the fiber.

Extended Data Figure 2 Predicted Ca²⁺ sparks at the centre of the fiber. *Binary images of the predicted fiber periphery (blue), predicted fiber centre (cyan), predicted Ca²⁺ sparks (green) and ground truth Ca²⁺ sparks (red) of the four images shown in Extended Data Fig. 1c – f. In **a** and **d**, two predicted Ca²⁺ sparks at the centre of the fiber can be observed. In **a** and **c**, two ground truth Ca²⁺ sparks at the centre of the fiber can be seen. The predicted and ground truth Ca²⁺ sparks are located in close proximity. In **c**, a small part of one of the three ground truth Ca²⁺ sparks extends beyond the predicted fiber mask. This was seen extremely rarely since there is a high agreement between predicted and ground truth fiber masks. Scale bar 20 µm.*

Extended Data Figure 3 Ca²⁺ sparks at the fiber periphery. a, *Fraction of Ca²⁺ sparks at the fiber periphery when one of two fibers in Exp1 was excluded from the analysis. The excluded fiber showed an unusually high Ca²⁺ spark frequency at the center of the fiber.* **b**, *Violin plots of the ratio of Ca²⁺ spark area over the fiber periphery and Ca²⁺ spark area for the three experiments (GT means ground truth and P prediction) and the training data set. Green triangles are the means.*

Extended Data Figure 4 **Ca²⁺ spark frequencies at y-shaped structures. a – d**, Ca²⁺ spark frequencies for the three experiments. Ca²⁺ sparks were counted as Ca²⁺ sparks at y-shaped structures if at least 20% (**a**), 40% (**b**), 50% (**c**) and 60% (**d**) of the Ca²⁺ spark area covered y-shaped structures.

Extended Data Figure 5   Ca²⁺ sparks seen in all image translations of a measurement associated with y-shaped structures. a, SHG measurement of the fiber shown in Fig. 4a. Scale bar 10 μm. **b**, Overlay of the predicted Ca²⁺ signal (green) (first image in Fig. 4a) and the SHG signal (red). The arrow indicates a predicted Ca²⁺ spark that could be seen in all image translations of this measurement (149 images) and is located in close proximity to y-shaped structures. **c**, SHG measurement of the fiber shown in Fig. 4b. Scale bar 10 μm. **d**, Overlay of the predicted Ca²⁺ signal (green) (first image in Fig. 4b) and the SHG signal (red). The arrow shows a predicted Ca²⁺ spark that could be seen in all image translations of this measurement (49 images) and is located at y-shaped structures.

Extended Data Figure 6   Properties of generated images with only one fiber in Exp1. *Only one fiber in Exp1 was included in the analyses presented in this figure. This fiber was shown in Fig. 2a, in Fig. 3a, b, d, f and in Fig. 4a, c. 1788 images out of 2180 images in Exp 1 are measurements of this fiber.* ***a*** *and* ***b****, Structural similarity index (SSIM) and mean squared error (MSE) of the three experiments.* ***c****, Pearson correlation coefficients of the overlapping fiber masks of predicted and ground truth images for the three experiments.* ***d*** *and* ***e****, Matthews correlation coefficients and F1 scores calculated based on ground truth and predicted fiber masks for the three experiments.* ***f****, Box plots and violin plots of fiber areas of ground truth (GT) and predicted (P) fiber masks for the three experiments.* ***g****, Box plots and violin plots of fiber mean intensity of ground truth (GT) and predicted (P) fiber masks for the three experiments and the training data set.*

Extended Data Figure 7 Properties of predicted Ca²⁺ sparks with only one fiber in Exp1. *Only one fiber in Exp1 was included in the analyses presented in this figure. This fiber was shown in Fig. 2a, in Fig. 3a, b, d, f and in Fig. 4a, c. 1788 images out of 2180 images in Exp 1 are measurements of*

*this fiber. **a** and **b**, F1 scores and Matthews correlation coefficients for the three experiments calculated based on binary images of ground truth and predicted Ca^{2+} sparks. **c** and **d**, Box plots and violin plots of Ca^{2+} spark area and Ca^{2+} spark mean intensity for the three experiments (GT means ground truth and P prediction) and the training data set. **e**, Ca^{2+} spark frequencies of the three experiments and the training data set. The ground truth Ca^{2+} spark frequency of Exp1 was 2.77 sparks/(10000*μm^2). The predicted Ca^{2+} spark frequency of Exp1 was 2.46 sparks/(10000*μm^2). **f**, Box plots and violin plots of the ratios Ca^{2+} spark area/Ca^{2+} spark edge for the three experiments and the training data set. **g**, Ca^{2+} spark frequencies at y-shaped structures for the three experiments. The ground truth Ca^{2+} spark frequency at y-shaped structures of Exp1 was 6.37 sparks/(10000*μm^2). The predicted Ca^{2+} spark frequency at y-shaped structures of Exp1 was 8.30 sparks/(10000*μm^2). **h**, Violin plots of Ca^{2+} spark area over y-shaped structures/Ca^{2+} spark area for the three experiments. **i**, Box plots and violin plots of the ratios Ca^{2+} spark area over myosin filaments/Ca^{2+} spark area for the three experiments and the training data set.*

Movie 1

xyt volume shown in Fig. 4c. Box volume 34.88 µm, 34.88 µm, 122.18 s.

Movie 2

xyt volume shown in Fig. 4d. Box volume 34.88 µm, 34.88 µm, 40.18 s.