A positive feedback loop between ZEB2 and ACSL4 regulates lipid metabolism to promote breast cancer metastasis

Jiamin Lin1†, Pingping Zhang1†, Wei Liu2†, Guorong Liu1, Juan Zhang1, Min Yan3, Yuyou Duan4,5*, Na Yang1*

1 Department of Laboratory Medicine, Guangzhou First People's Hospital, South China University of Technology, Guangzhou, Guangdong, 510180, China.

2 Department of Breast Surgery, Guangzhou Red Cross Hospital, Guangzhou Red Cross Hospital of Jinan University, Guangzhou, Guangdong, 510220, China.

3 Sun Yat-sen University Cancer Center, State Key Laboratory of Oncology in South China, Collaborative Innovation Center of Cancer Medicine, Guangzhou, 510060, China.

4 Laboratory of Stem Cells and Translational Medicine, Institutes for Life Sciences and School of Medicine, South China University of Technology, Guangzhou, 510006, China.

5 Institute for Clinical Medicine, Guangzhou First People's Hospital, South China University of Technology, Guangzhou 510180, China.

†These authors contributed equally to this work.

*Correspondence to:

Na Yang, M.D., Ph.D.

Department of Laboratory Medicine

Guangzhou First People's Hospital

South China University of Technology

1 Panfu Road, Guangzhou 510180, P.R. China

E-mail: eyyangna@scut.edu.cn

Yuyou Duan, Ph.D.

Laboratory of Stem Cells and Translational Medicine

Institutes for Life Sciences, School of Medicine

South China University of Technology

No.382 Waihuan East Road, Suite 406

Higher Education Mega Center

Guangzhou 510006, P.R. China

E-mail: yuyouduan@scut.edu.cn

Min Yan, Ph.D

40   State Key Laboratory of Oncology in South China

41   Sun Yat-sen University Cancer Center

42   651 Dongfeng Road East, Guangzhou, 510060, P.R. China

43   Email: yanmin@sysucc.org.cn

Highlights

46   ZEB2 activates FAO through transcription of ACSL4

47   ACSL4 regulates lipid metabolism through FAO, promoting breast cancer metastasis

48   Targeting ZEB2-ACSL4 signaling axis inhibits breast cancer metastasis

Significance

51   We describe a novel positive feedback loop between ZEB2 and ACSL4 that results in

52   enhanced cellular lipid storage and FA oxidation to drive breast cancer metastasis.

54   **Running Title:** A novel positive feedback loop between ZEB2 and ACSL4

55   **Key Words:** Lipid metabolism; Lipid droplet; cancer metastasis; EMT

Abstract

Lipid metabolism plays a critical role in cancer metastasis. However, the mechanisms through which metastatic genes regulate lipid metabolism remain unclear. Here, we describe a new oncogenic-metabolic feedback loop between the epithelial-mesenchymal transition (EMT) transcription factor ZEB2 and the key lipid enzyme ACSL4 (long-chain acyl-CoA synthetase 4), resulting in enhanced cellular lipid storage and fatty acid oxidation to drive breast cancer metastasis. Functionally, Depletion of ZEB2 or ACSL4 significantly reduced lipid droplets (LD) abundance and cell migration. ACSL4 overexpression rescued the invasive capabilities of the ZEB2 knockdown cells, suggesting that ACSL4 is crucial for ZEB2-mediated metastasis. Mechanistically, ZEB2 activated ACSL4 expression by directly binding to the ACSL4 promoter. ACSL4 binds to and stabilizes ZEB2 by reducing ZEB2 ubiquitination. Notably, ACSL4 not only promotes the intracellular lipogenesis and lipid droplet accumulation but also enhances fatty acid oxidation (FAO) and ATP production by upregulating the FAO rate-limiting enzyme CPT1A (carnitine palmitoyltransferase 1 isoform A). Finally, we demonstrated that ACSL4 knockdown significantly reduced metastatic lung nodes in vivo. In conclusion, we reveal a novel positive regulatory loop between ZEB2 and ACSL4, which promotes LD storage to meet the energy needs of breast cancer metastasis, and identify the ZEB2-ACSL4 signaling axis as an attractive therapeutic target for overcoming breast cancer metastasis.

Introduction

More than 90% of breast cancer-related deaths are due to metastasis(Fahad Ullah, 2019; Park et al., 2022). Current treatments, including endocrine therapy, chemotherapy, and radiation therapy, are ineffective in preventing breast cancer

metastasis and remain the greatest clinical challenge for breast cancer
treatment(Barzaman et al., 2020). Cancer metastasis is associated with a process
called epithelial-mesenchymal transition (EMT)(Luond et al., 2021; Mittal, 2018). EMT
is a pre-metastatic state in which epithelial cells lose their tight junctions and convert
to migratory mesenchymal cells(Dongre and Weinberg, 2019). It has become
increasingly recognized that metabolic reprogramming during the EMT process
contributes to metastasis and tumorigenesis(Kang et al., 2019; Sciacovelli and Frezza,
2017). Metastatic invasion is a highly energy-intensive process(Mosier et al., 2021).
Emerging evidence suggests that lipid metabolic reprogramming plays a critical role in
meeting the energy requirements of metastatic invasion(Bian et al., 2021; Cheng et al.,
2018; Li et al., 2020; Snaebjornsson et al., 2020). Elucidating the mechanism by
which reprogrammed lipid metabolism helps us exploit novel and attractive targets for
metastatic therapeutic interventions.
95
Lipid metabolism includes a complex network of pathways that regulate fatty acid (FA)
synthesis, storage, and degradation(Maan et al., 2018). For lipid anabolism, FA are
stored in a dynamic organelle called a lipid droplet, which is composed of a monolayer
of phospholipids that covers a hydrophobic core containing neutral lipids, such as
triacylglycerol (TAG) and cholesteryl esters (CE)(Cruz et al., 2020). Lipid droplets (LD)
accumulation is associated with aggressiveness in many cancer types(Antunes et al.,
2022), including breast(Zembroski et al., 2021), brain(Taib et al., 2019), liver(Niu et al.,
2021), lung(Jin and Yuan, 2020), and prostate(Huang et al., 2012; Roman et al.,
2020). Indeed, the aggregation of FA in LD is considered a priming state to prepare for
metastasis(Petan, 2020; Rozeveld et al., 2020; Senga et al., 2018). In the case of
need, FA can be released and oxidized for energy support. For example, it has been
reported that fatty acids stored in lipid droplets were a crucial resource in fueling the
metastatic process in pancreatic cancer (Rozeveld et al., 2020). Additionally, previous
studies have suggested that metastatic triple-negative breast cancer (TNBC) depends
on fatty acid oxidation (FAO) to produce high ATP levels (Camarda et al., 2016; Wright
et al., 2017). Although lipid metabolism is crucial for cancer metastasis, the signalling
pathway that regulates lipid metabolic reprogramming during breast cancer
metastasis remains unclear.
114
Dysregulation of lipid metabolic enzymes has been documented in cancer
metastasis(Luo et al., 2017). Long-chain fatty acyl synthetase (ACSL) 4 belongs to
the long-chain acyl-CoA synthetase ligase enzyme family (Quan et al., 2021; Rossi
Sebastiano and Konstantinidou, 2019). ACSL4 catalyses the conversion of long-chain
FAs to acyl-CoAs, a necessary step for free long-chain FAs to enter the next
metabolic pathway(Kuwata and Hara, 2019). The increased expression and activity of
ACSL4 have been observed in many cancer types and are well-known biomarkers of
ferroptosis(Doll et al., 2017). Although it has been reported that ACSL4 is a tumor
suppressor that activates ferroptosis, many studies have suggested that ACSL4 is an
oncogene that contributes to tumor progression. For example, ACSL4 promotes
Hepatocellular carcinoma (HCC) cell proliferation and metastasis via lipogenesis and

lipid droplet accumulation(Chen et al., 2020; Niu et al., 2021). In prostate cancer,
ACSL4 promotes cell growth, invasion, and hormonal resistance(Wu et al., 2015). The
function of ACSL4 in breast cancer has been implicated in hormone therapy
resistance involving the regulation of energy-dependent transporter
expression(Orlando et al., 2019). However, regulation of lipid metabolism by ACSL4
during breast cancer invasion remains unclear.

In the present study, we demonstrate a novel positive feedback loop between the
EMT transcription factor ZEB2 and the essential lipid metabolic enzyme ACSL4,
resulting in enhanced cellular lipid droplet accumulation and fatty acid oxidation to
drive breast cancer metastasis. Mechanistically, ZEB2 activates ACSL4 expression by
directly binding to the ACSL4 promoter. ACSL4 stabilizes and upregulates ZEB2 via
transcriptional and post-transcriptional mechanisms. In addition, we also provide
evidence that overexpression of ZEB2 or ACSL4 is associated with worse prognosis
in advanced breast cancer. Our findings provide insights into lipid metabolic
mechanisms during the EMT process and reveal a novel oncogenic-metabolic
pathway critical for breast cancer EMT and metastasis.

Materials and Methods
Breast cancer cell lines and clinical specimens
Breast cancer cell lines MCF-7, MDA-MB-231, and BT549 were purchased from the
American Type Culture Collection (ATCC, Beijing Zhongyuan Ltd., Beijing, China).
Paclitaxel-resistant and epirubicin-resistant cell lines have been reported previously
(Yang et al., 2017; Zheng et al., 2014). All cell lines were maintained in Dulbecco's
modified Eagle's medium (GIBCO) supplemented with 10% (v/v) FBS (BI, Biological
Industries) and 1% (v/v) Pen/Strep (GIBCO) and incubated in a humidified
atmosphere of 5% CO_2 at 37 °C.

Primary breast cancer and adjacent normal tissues were collected from the volunteers
at Guangzhou First People's Hospital, Guangdong, China. This study was approved
by the Ethics Committee of the Guangzhou First People's Hospital (approval no.
K2021-201-01). Fourteen patients were enrolled in this study. The specimens were
then subjected to western blotting and immunohistochemistry (IHC) assays.

Plasmid construction and cell transfection
The PLKO.1-TRC-LUC lentivirus vector containing shRNA (shACSL4-1 and
shACSL4-2) and the control vector were purchased from WZ Biosciences. The virus
package plasmids psPAX2 and pMD2.G were purchased from Addgene. To produce
the lentivirus, the shRNA vector and package plasmids were co-transfected into 293T
cells using polyethyleneimine (Polysciene). After 48 hours or 72 hours, supernatants
were harvested and passed through 0.45 um filters to collect the virus and stored at
-80 °C until further use. MDA-MB-231 cells were transfected with a viral solution
containing 8ug/ml polybrene (Hanbio Biotechnology). 48 hours after transfection, the

virus was removed from the culture and fresh medium was added. The cells were
subsequently selected using 2 mg/mL puromycin to obtain a stable cell line. MCF-7
cells were transfected with HBLV-h-ACSL4-3xflag-ZsGreen-PURO or control lentiviral
plasmid. The plasmids were purchased from Hanbio Biotechnology.

For the siRNA assay, 5 nM negative control siRNA or ACSL4 and ZEB2 siRNA
(GenePharma) were transfected into MDA-MB-231 or paclitaxel-resistant cells using
Lipofectamine 3000 (Invitrogen) according to the manufacturer's instructions. Six
hours after transfection, the medium was replaced with fresh growth medium and the
cells were harvested after 24-48 hours for qPCR or western blot analysis. The siRNA
sequences are listed in Supplementary Table S1.

Western blot analysis
For western blotting, cells were lysed in WIP buffer (Beyotime) containing protease
and phosphatase inhibitors. Protein levels were quantified by BCA assay (Beyotime),
and equal amounts were separated using 10% SDS-PAGE electrophoresis and
transferred onto a 0.22 μm PVDF membrane for probing. After blocking for 1 hour in 5%
non-fat milk diluted in TBST, the membrane was incubated with various primary
antibodies at 4°C overnight. The following primary antibodies were used: ACSL4
(ab155282, Abcam), ZEB2 (sc-271984, Santa Cruz), E-cadherin (3195S, CST),
N-cadherin (610921, BD Biosciences), vimentin (D21H3, 5741T, CST), CPT1A
(ab220789, Abcam), GAPDH (10494-1-AP, Protein Tech), Myc-Tag (71D10, 2278T,
CST), and HA-Tag (C29F4, 3724T, CST). Next, secondary antibodies were added and
the proteins were detected using an ECL kit (Millipore). Immunoreactive signals were
visualized using the GE Amersham Imager 600 chemiluminescence system.

Migration and invasion assay
A wound healing assay was performed to examine the migratory ability of the cells.
Briefly, cells were seeded in 6-well plates at a density of 4×10^5 cells per well in a
complete medium at 37 °C. Until cells reached 80-90% density, a sterile plastic tip
was used to create a wound line across the surface of the plates. The suspended
cells were discarded. After The cells were cultured in reduced serum DMEM medium
in a 5% CO_2 incubator at 37 °C for 48 hours. Images were obtained using a
phase-contrast microscope. Each assay was performed in triplicates.

The cell invasion capacity was measured using Matrigel-coated Transwell chambers
(8.0 μm; Corning Inc.). Briefly, cells were suspended in 200 μL serum-free medium
and seeded in the upper chamber of a Transwell. The lower chamber of the Transwell
was filled with medium containing 10% FBS-containing medium. The Transwell
chambers were then incubated at 37 °C. After culturing for 24 hours, the Transwell
holes were penetrated with 4% paraformaldehyde for 20 minutes and then stained
with 0.1% crystal violet solution for 15 minutes. Invading cells were imaged and

211   counted in five random fields.

Immunofluorescence (IF) and Immunohistochemistry (IHC) assays

For the immunofluorescence assay, cells were seeded on confocal dishes, washed in
PBS three times, and fixed in 4% paraformaldehyde for 15 minutes at room
temperature. The cells were blocked with 10% goat serum (AR0009, BOSTER) for 1 h
at room temperature and washed three times. The cells were then incubated with the
primary antibodies for 1 hour at room temperature. Subsequently, the cells were
rinsed thrice with PBS and incubated with secondary antibodies for 1 hour at room
temperature.

For BODIPY 493/503 staining, cells were fixed in 4% PFA for 15 minutes and washed
twice with PBS. The cells were then incubated with BODIPY 493/503 (1:2500 in PBS,
GLPBIO) for 15 minutes at room temperature. DAPI was used to stain the nuclei.
Images were acquired by confocal microscopy (Ni-E-A1, Nikon, 40x) and analyzed
using NIS Elements Viewer software and ImageJ.

For immunohistochemistry (IHC) assay, paraffin-embedded tissue slides were
dewaxed with xylene and rehydrated using a graded series of alcohols. This was
followed by antigen retrieval and blocking with 5% BSA for 60 minutes. After that,
Tissue slices were incubated with primary antibodies against ACSL4, ZEB2, Vimentin
or ERα at 4 °C overnight in a humidified container and then detected with the SP
Rabbit&Mouse HRP Kit (CWBIO). Images were acquired using a digital pathology
scanning system (Aperio CS2).

Metabolic analysis

For the oxygen consumption rate (OCR) assay, the XF long-chain fatty acid oxidation
(LCFA) stress test Kit (103672-100) and Seahorse XF96 Analyzer were used to
investigate the long-chain fatty acid OCR and extracellular acidification rate (ECAR)
according to the manufacturer's protocol.

ATP levels were determined using an ATP Assay Kit (S0026, Beyotime) according to
the manufacturer's protocol. ATP levels were calculated using luminescence signals
and were normalized to protein concentrations. Glycolytic activity was determined
using a Glycolysis Cell-Based Assay Kit (600450, Cayman Chemical), according to
the manufacturer's protocol.

Untargeted lipidomics and proteomics

Cell Lipids were extracted in a chloroform-methanol mixed solution (2:1, -20 °C). The
extracted cell lysates were immersed in liquid nitrogen and frozen for 5 minutes. The
extracted cell lysates were placed in a 2 ml adapter, and the above steps were

repeated twice. Samples were then centrifuged to pellet the proteins (5 minutes,
12000 rpm), and the supernatant was stored for analysis in a vacuum centrifugal
concentrator. The sample was dissolved in 200 μl isopropanol, filtered through a 0.22
μm membrane, and detected by Liquid Chromatography-Mass Spectrometry (LC-MS).
The lipidomics data were analyzed using LipidSearch software (version 4.0). The
software identified intact lipid molecules based on their molecular weight and
fragmentation pattern using an internal library of predicted fragment ions per lipid
class. The spectra were then aligned based on retention time and the MS1 peak
areas were quantified across the sample conditions. Excel 2010 was used to calculate
intensity, and the R program (version 3.2.5) was used for data manipulation and
statistical analyses, including unsupervised hierarchical clustering and heatmap
visualization.

Quantitative Real-time PCR

266 Total RNA was isolated using the TRIzol reagent (Invitrogen), and cDNAs was
267 synthesized from total RNA using the PrimeScript™ RT Master Mix (Perfect Real
268 Time, Takara). Quantitative real-time PCR was performed in triplicate using PowerUp
269 SYBR Green (Thermo Fisher Scientific). The relative gene expression was measured
270 using the $2^{-\Delta\Delta Ct}$ method. All primers used are listed in Supplementary Table S2.

Luciferase reporter assays

For Luciferase assays, the 2000bp ACSL4 promoter vector containing luciferase was
purchased from WZ Biosciences Inc. The -287bp, -965bp, -1038bp, and -1116bp
regions of the ACSL4 promoter were cloned by PCR amplification using primers
containing restriction sites MIII and Hind III. The nucleotide sequences of primers
used are listed in Supplementary Table S3. 293T cells were transfected with 1ug of
the-287bp, -965bp, -1038bp, -1116bp or 2000bp ACSL4 promoter luciferase reporter
and an empty vector for 24 hours. After transfection, the cells were harvested and
analyzed using Bright-Glo reagent (Promega) according to the manufacturer's
instructions.

Chromatin immunoprecipitation（ChIP）assay

ChIP assays were performed using a ChIP assay kit (Cat.p2078, Beyotime
Biotechnology). Briefly, MDA-MB-231 cells were grown to 90% confluence and
crosslinking was performed with 1% formaldehyde for 10 minutes. Mouse anti-ZEB2
antibody or mouse IgG was used to immunoprecipitate the DNA-containing
complexes. After the DNA purification (Cat. D0033; Beyotime Biotechnology), PCR,
and Q-PCR were performed to detect the ZEB2-binding site in the ACSL4 promoter
region. The primer sequences are listed in Supplementary Table S4.

Co-immunoprecipitation (Co-IP) assay and Ubiquitination assay

For co-IP assays, HEK293T cells were co-transfected with ACSL4-FLAG and
ZEB2-MYC plasmids. 48 hours after transfection, cells were lysed in WIP buffer
(Beyotime) containing a protease inhibitor and phosphatase inhibitors for 30 minutes
at 4 °C followed by centrifugation. The supernatants were immunoprecipitated with
the indicated antibodies overnight at 4 °C, followed by incubation with protein A/G
beads for 1 hour at 4 °C. After incubation, the beads were washed with WIP buffer and
boiled in a 2×loading buffer. Protein samples were analyzed by western blotting.

For the ubiquitination assay, HEK293T cells were transfected with HA-Ubi plasmid,
ACSL4-FLAG plasmid, and ZEB2-MYC or vector plasmid. 48 hours after transfection,
the cells were treated with 20 µM MG-132 for 6 hours to block the proteasomal
degradation of ZEB2 before being lysed with WIP lysis buffer (Beyotime). Equal
amounts of protein lysates were immunoprecipitated with anti-MYC beads and
subjected to SDS-PAGE, followed by blotting with anti-HA (ubiquitin) to visualize
polyubiquitinated ZEB2 protein bends.

In vivo experiments

The Ethics Committee approved the animal experiments for Animal Experiments of
the South China University of Technology. All NSG mice were purchased from the
Medical Laboratory Animal Center of Guangdong Province, China. Six female NSG
mice aged 6-8 weeks were used in each group for the primary tumor growth and
spontaneous lung metastasis experiments. A total of 2×10^6 vector control or
shACSL4 cells were mixed 1:1 by volume with Matrigel (BD Biosciences). Each
mouse was injected orthotopically into both the flanks. Xenograft tumor growth was
measured and tumor volume was calculated as follows: volume=(length×width^2)/2. At
the experimental endpoint, mice were intraperitoneally injected with 150 mg/kg
D-luciferin and imaged for 2 minutes using a live imager. The xenograft tumor and
lung samples were removed and fixed to count the metastatic lung nodules.

Statistical analysis

The results are reported as the mean ± standard deviation (SD) or mean ± standard
error of the mean (SEM), as indicated in the figure legend. Statistical comparisons
were performed using an unpaired two-sided Student's t-test and considered
significant if * P<0.05, ** P <0.01, *** P <0.001, and **** P <0.0001. Statistical
comparisons for the LM2 lung metastasis assay were performed using the Mann–
Whitney U test. All experiments with representative images, including western blotting
and immunofluorescence, were repeated at least twice, and representative images
are shown.

Results

ZEB2 and ACSL4 are overexpressed and correlated in highly invasive breast cancer cells

To explore the molecular mechanism of highly invasive breast cancer, we performed RNA sequencing analysis of wild-type and two drug-resistant luminal breast cancer cell lines. Significant changes in 6155 genes ($P < 0.05$). Notably, EMT and stemness genes such as ZEB2, SNAIL, TWIST, TRAIL, Gli2, WNT, and AKT3, which are overexpressed in basal-like breast cancer (BLBC), were significantly upregulated in drug-resistant cells (Supplementary Fig. S1A). In contrast, differentiated genes such as FOXA1, ERα, E-cadherin, and GATA3, which are highly expressed in the luminal subtype, were significantly downregulated (Supplementary Fig. S1B), suggesting that drug-resistant cells underwent EMT and became stem-like cells. We noticed that the rate-limiting enzymes of fatty acid metabolism, long-chain fatty acyl synthetase 4 (ACSL4), and EMT transcription factor ZEB2 were among the top 200 upregulated (>1.5-fold) genes (Fig. 1A). To verify these findings in the clinical sample, we analyzed ZEB2 and ACSL4 expression in TCGA database and found that ACSL4 expression was positively correlated with ZEB2 expression (Fig. 1B, r=0.7657, p<0.001) and inversely correlated with ERα expression (Fig. 1C, r=-0.3312, p<0.001). Using the Kaplan–Meier plotter database, we compared the post-progression survival (PPS) between ACSL4 or ZEB2 high- and low-expression breast cancer patients. We found that patients with higher ACSL4 or ZEB2 expression had worse prognosis than patients with advanced breast cancer (Fig. 1D and 1E). It is worth noticing that there is no difference in the overall survival between ACSL4 or ZEB2 high and low-expression breast cancer patients (supplementary Fig. S2).

To confirm this correlation, we performed western blot analysis and found that Basal-like and Taxol-resistant cell lines, which lack ERα expression, showed high expression of ZEB2 and ACSL4, whereas low or no ZEB2 and ACSL4 expression was detected in luminal subtype cell lines (Fig. 1F). Consistently, tissue samples from nine breast cancer patients had similar expression patterns, with ZEB2 and ACSL4 being relatively highly expressed in ER-negative patient samples (patients 6, 7, 8, and 9) (Fig. 1G and 1H). The immunohistochemistry staining assay, shown in Fig. 1I, also confirmed that ACSL4 and ZEB2 expression was elevated in ERα-negative breast carcinomas compared with ERα-positive breast carcinomas. Taken together, these results indicate that ACSL4 and ZEB2 are correlated and overexpressed in highly invasive breast cancers.

Overexpression of ACSL4 contributes to ZEB2-mediated breast cancer invasion.

Because ACSL4 is overexpressed in highly invasive breast cancer cells, we

hypothesized that ACSL4 is essential for driving breast cancer migration and invasion.
We then established a stable ACSL4 overexpressed MCF7 cell line, which was less
aggressive than the basal-like breast cancer cells. The overexpression of ACSL4
significantly enhanced the metastatic and invasive capacities of MCF7 cells (Fig. 2A
and 2B). Conversely, ACSL4 knockdown by shRNA significantly reduced the
metastatic and invasive capacity of MDA-MB-231 cells compared to that of the control
cells (Fig. 2C and 2D). We also observed that the metastatic and invasive abilities of
MDA-MB-231 cells were significantly reduced in ZEB2 knockdown cells
(Supplementary Fig. S3). To investigate the downstream genes regulated by ACSL4,
we performed RNA sequencing analysis and identified that the tight junction and focal
adhesion pathways were among the most upregulated pathways in ACSL4
knockdown cells compared to control cells (Fig. 2E). To verify the RNA-seq results,
we examined the expression of adhesion-related genes by immunoblotting and found
that after silencing of ACSL4 and ZEB2, the luminal epithelial marker E-cadherin was
increased, whereas the mesenchymal markers vimentin and N-cadherin were
decreased (Fig. 2F). These results confirm the essential role of ACSL4 in breast
cancer invasion and migration.
389
We hypothesized that the ZEB2-ACSL4 axis played a crucial role in metastatic ability.
To examine whether ACSL4 was required for ZEB2-mediated breast cancer invasion
and migration, ACSL4 was overexpressed in ZEB2-silencing cells (Fig. 2G). As
expected, overexpression of ACSL4 significantly restored the invasive and metastatic
capacities of ZEB2 knockdown cells by 35.1% and 32.4%, respectively (Fig. 2H and
2I), indicating that ACSL4 is essential for ZEB2-mediated breast cancer invasion and
migration.
397

ACSL4 and ZEB2 promote Lipid droplet production and lipogenesis
ACSL4 is a member of the long-chain acyl-CoA synthetase family, which catalyzes the
conversion of long-chain fatty acids into their active forms. However, the mechanism
by which ACSL4 regulates lipid metabolism in breast cancer remains unclear. A
previous study revealed that ACSL4 promotes lipid droplets (LD) accumulation in
hepatocellular carcinoma (HCC)(Niu et al., 2021). Thus, we measured the basal LD
content of breast cancer cells, including MCF-7, MDA-MB-231, and Taxol-resistant
MCF-7 cells. The basal number and size of lipid droplets were significantly larger in
MDA-MB-231 and Taxol-resistant MCF-7 cells, both of which are highly invasive
breast cancer cell lines (Fig. 3A). Fluorescence microscopy revealed that ACSL4
co-localized with LD in MDA-MB-231 cells (Supplementary Fig. S4). Notably, ACSL4
knockdown reduced cytoplasmic LD abundance and LD-containing cells in
MDA-MB-231 cells (Fig. 3B). Consistently, cytoplasmic LD abundance and
LD-containing cells was significantly reduced in ZEB2-depleted cells (Fig. 3C).
412
413   Lipid droplets are phospholipid monolayers containing a hydrophobic core comprising
414   triacylglycerols (TG) and cholesterol esters (CE). As ACSL4 promotes intracellular LD

accumulation in breast cancer cells, we investigated the effect of ACSL4 on the lipid composition of breast cancer cells. Untargeted Lipidomic analyses revealed that ACSL4 knockdown breast cancer cells had a reduced ability to incorporate long-strain monounsaturated (18:1, 17:1, 22:1, 24:1, 26:1) and saturated FA (16:0, 18:0, 24:0, 26:0) into triacylglycerol and phospholipids (PL), indicating that ACSL4 promotes the incorporation of long-chain monounsaturated and saturated fatty acids into triacylglycerol and phospholipids in these cells (Fig. 3D and 3E). Moreover, all these lipids, including triacylglycerol, phospholipids, and cholesterol esters, decreased the incorporation of several polyunsaturated FA, such as 22:6, 20:4, 22:4, and 22:5, in ACSL4 depleted cells as reported previously (Fig. 3D and Supplementary Fig. S5A)(Doll et al., 2017). Consistently, the total amount of different lipid species, as shown in Supplementary Fig. S5B, were significantly decreased after ACSL4 knockdown. All together, these results suggest that ACSL4 directs the long train of free fatty acids into lipid anabolism to form different lipid species in the LD or other cellular membranes.

Exogenous fatty acid promotes LD accumulation and fuel cell invasion

A previous study reported that some tumors tended to increase their intake of extracellular fatty acids to promote migration(Rozeveld et al., 2020). As we observed that depletion of ACSL4 greatly reduced cytoplasmic LD abundance and invasive potential in basal-like breast cancer cells, we hypothesized that LD accumulation is an important step prior to breast cancer invasion. To verify our hypothesis, we examined whether exogenous fatty acids contributed to LD accumulation and the invasive ability of breast cancer cells. Exogenous oleic acid (OA) was added to the cell culture medium. Treatment with oleic acid dramatically enhanced LD abundance in the cells (Fig. 4A and 4B), indicating that breast cancer cells tend to store lipids in LD for energy reserves. Next, we assessed the effects of exogenous oleic acid treatment on cell migration. Transwell and Wound healing assays revealed that oleic acid-treated cells exhibited significantly enhanced invasive and metastatic capacities compared with control cells (Fig. 4C and 4D).

Previous study reported that LD undergo lipolysis during the process of migration in pancreatic cancer(Rozeveld et al., 2020). We reasoned that breast cancer cells utilize stored lipids during migration to fuel metastasis. Lipid droplet content was analyzed using fluorescence microscopy after cell migration. Interestingly, we observed significantly reduced LD in cells on the lower side of the transwell chamber, suggesting that lipids stored in LD were utilized and degraded during cell migration (Fig. 4E). Taken together, these results suggest that lipids stored in lipid droplets are a crucial resource to fuel the process of breast cancer invasion and migration.

ACSL4 and ZEB2 stimulate long-chain fatty acid oxidation and ATP generation in BLBC Cells

To explore the mechanisms by which ACSL4 regulates lipid metabolism, we
performed RNA sequencing to investigate downstream pathways and genes
regulated by ACSL4. KEGG enrichment analysis revealed that the FAO pathway was
among the top 20 regulated pathways (Fig. 5A). ACSL4 knockdown reduced the
expression of genes involved in the FAO pathways (Fig. 5B). Importantly, we
observed that the FAO rate-limiting enzyme CPT1A was significantly reduced in
ACSL4 knockdown cells (Fig. 5B). Reverse transcription PCR confirmed that deletion
of ACSL4 significantly reduced the expression of CPT1A, but did not affect the
expression of CPT1B and CPT1C (Fig. 5C). Consistently, ACSL4 silencing reduced
CPT1A protein expression (supplementary Fig. S6A). In addition, levels of other lipid
metabolic enzymes, such as ATGL, FASN, and SREBP2, were significantly
decreased after ACSL4 knockdown (Supplementary Fig. S6B). As ACSL4 knockdown
decreased CPT1A expression, we reasoned that ACSL4 might stimulate fatty acid
oxidation in highly invasive breast cancer. Oxygen consumption rate (OCR) was
calculated in ACSL4 or ZEB2 silencing and control cells. We observed that the OCR
rate derived from long-chain fatty acids was significantly reduced in ACSL4 or ZEB2
silencing cells (Fig. 5D and 5E), suggesting the metabolic advantage of increased
long-chain fatty acid oxidation and oxidative phosphorylation (OXPHOS).

Since OXPHOS is accompanied by increased ATP generation, we measured ATP
levels and found that ACSL4 or ZEB2 knockout cells had significantly reduced ATP
generation compared to control cells (Fig. 5F and 5G). To exclude OXPHOS derived
from the aerobic glycolysis pathway, we measured lactate production and found no
significant difference in lactate production between ACSL4 or ZEB2 knockdown and
control cells, suggesting that glucose metabolism is not involved in ACSL4 or ZEB2
mediated metabolic process (Fig. 5H and 5I). These results indicated that ACSL4
upregulates CPT1A to stimulate fatty acid oxidation and ATP generation in BLBC
Cells.

ZEB2 transcriptionally activates the expression of ACSL4
ZEB2 is a crucial transcription factor involved in EMT. Because ZEB2 and ACSL4
were highly correlated in the clinical samples in TCGA database, we investigated
whether ACSL4 is a direct downstream target of ZEB2. Silencing of ZEB2 by siRNA in
MDA-MB-231 cells significantly reduced ACSL4 mRNA levels and protein expression
in MDA-MB-231 cells (Fig. 6A and 6 B). We found that the ACSL4 promoter contained
four canonical ZEB2-binding E-boxes (CAGGT/CACCT) located at -287, -965, -1038,
and -1116 of the ACSL4 promoter, respectively (Fig. 6C). Therefore, we cloned five
segments of the ACSL4 promoter and a control segment to generate
promoter-luciferase constructs, based on the location of these E-boxes (Fig. 6C).
ZEB2 overexpression significantly enhanced the luciferase activity of all five
E-box-containing segments of the ACSL4 promoter, whereas no significant change
was observed in the control segment (Fig. 6D). To examine whether ZEB2 directly
binds to the ACSL4 promoter, we performed chromatin immunoprecipitation (ChIP)

using four sets of primers (Fig. 6E). Primer set 1, which covered segment 1, consistently exhibited apparent ZEB2 binding (Fig. 6F and 6G). However, no binding was detectable using the other three sets (Fig. 6F), indicating that ZEB2 binds to the E-Box located at nucleotides -184 to -295 of the ACSL4 promoter.

Since the -287 E-box exhibited apparent ZEB2 binding, we generated mutants of the -287 E-box promoter-luciferase construct. The Luciferase Reporter Assay revealed that ACSL4 promoter activity was almost entirely abolished by the mutations (Fig. 6H), suggesting that this promoter region is essential for ZEB2-mediated ACSL4 promoter activation. Taken together, these data suggest that ZEB2 directly binds to the ACSL4 promoter to activate its mRNA expression.

ACSL4 reduces ZEB2 mRNA expression and protein stabilization

We hypothesized that ACSL4 regulates the expression of ZEB2. We then performed quantitative PCR and immunoblotting and observed that both ZEB2 mRNA and protein levels were reduced after the depletion of ACSL4 in the two BCSC cell lines (Fig. 7A and 7B). A previous study has reported that ACSL4 regulates c-Myc protein stability in HCC. We reasoned that ACSL4 might in turn regulate the stability of ZEB2. Therefore, we performed a ubiquitination assay to investigate whether ACSL4 regulates ZEB2 protein stability via ubiquitination. HEK293T cells were co-transfected with HA-ubiquitin and myc-ZEB2 expression vectors along with either an empty vector or an ACSL4 overexpression vector. As shown in Fig. 7C, the expression of ACSL4 caused a significant decrease in the ubiquitination of ZEB2, suggesting that ACSL4 attenuated the ubiquitin proteolysis of ZEB2. Notably, co-IP assays revealed a specific interaction between ACSL4 and ZEB2 proteins, supporting the notion that ZEB2 and ACSL4 are present in a protein complex (Fig. 7D). To further confirm the role of ACSL4 in the regulation of ZEB2 proteolysis, cycloheximide was added to the cell medium to block the synthesis of new proteins. Interestingly, endogenous ZEB2 protein levels were almost completely suppressed in ACSL4 knockdown cells from 0h to 8h time points (Fig. 7E and 7F). This phenomenon could be explained by the fact that ZEB2 mRNA levels were significantly suppressed by the ACSL4 knockdown. Conversely, ZEB2 protein maintained a relatively steady level at 6 hours in ACSL4 overexpressing cells compared with control cells that undergone obvious ZEB2 proteolysis at 4 hours after adding CHX, indicating an increased ZEB2 protein stability in ACSL4 overexpressing cells (Fig. 7G and 7H). Taken together, these results indicate that ACSL4 not only stabilizes the ZEB2 protein by attenuating its ubiquitination but also upregulates ZEB2 mRNA expression. Therefore, ACSL4 regulates ZEB2 through both transcriptional and post-transcriptional mechanisms.

ACSL4 knockdown inhibits lung metastasis of BLBC in the animal model

To further validate our in vitro findings, we examined the effect of ACSL4 on lung metastasis of breast cancer cells in an animal model. Depletion of ACSL4 resulted in a significant reduction in tumor growth and lung colonization of highly metastatic

MDA-MB-231 cells (Fig. 8A). In contrast, the control group showed significantly more
lung metastatic nodules (Fig. 8B) and rapid tumor growth (Fig. 8C and 8D).
Immunohistochemistry confirmed a striking downregulation of ACSL4, accompanied
by ZEB2 and vimentin, compared to those in the control group (Fig. 8E).

To further validate the role of LD in ACSL4 high-expression cells, we examined LD
content in frozen sections of the two groups. Consistent with the in vitro results, there
was an obvious reduction in LD accumulation in the ACSL4 depletion tissue
compared to that in control tissue (Fig. 8F). Taken together, these data suggest that
ACSL4 promotes the lung metastasis of breast cancer cells in vivo.

Discussion
The rewiring of metabolic pathways during EMT has only recently been recognized. In
addition to glycolysis, dysregulated lipid metabolism has been shown to contribute to
cancer invasion and metastasis(Corbet et al., 2020; Soukupova et al., 2021). Many
EMT-driving factors have been found to reprogram the lipid metabolic pathways by
regulating metabolic enzymes. For example, TGF-β, a major driver of EMT, activates
fatty acid synthase (FASN) and forms a FASN-TGFβ1-FASN positive loop in
cisplatin-resistant cells, resulting in EMT induction(Hua et al., 2020; Jiang et al., 2015;
Yang et al., 2016). However, the direct regulation of lipid enzymes by EMT factors
remains to be elucidated. Herein, we demonstrated that the lipid rating-enzyme
ACSL4 is a direct downstream target of the EMT transcription factor ZEB2 in
controlling lipid metabolism. Mechanistically, ZEB2 activates ACSL4 mRNA
expression by directly binding to its promoter, which contains four ZEB2 consensus
sequences. Importantly, we observed a strong correlation between ZEB2 and ACSL4
expression levels in clinical breast cancer samples. ACSL4 re-expression rescues the
migration ability of ZEB2-depleted cells. These observations indicated that ACSL4 is
crucial for ZEB2-mediated metastasis.

We also demonstrated that ACSL4 directly binds to and stabilizes ZEB2 by reducing
its ubiquitination. Our results are consistent with a recent report that ACSL4 stabilizes
the oncoprotein c-Myc via the ubiquitin-proteasome system(Chen et al., 2020). We
propose that as an LD enzyme, ACSL4 may participate in the protein degradation
system of LD. This is likely to be a novel function of ACSL4. Interestingly, a recent
study reported that lipid droplets recruit Numb through the AP2A/ACSL3 complex to
promote Numb degradation(Liu et al., 2022). Thus, it is likely that lipid droplets act as
a platform for protein degradation. Additionally, ACSL4 promoted ZEB2 expression as
the mRNA level of ZEB2 was significantly reduced after ACSL4 knockdown.
Therefore, ACSL4 regulates ZEB2 not only via a post-transcriptional mechanism but
also via a transcriptional mechanism. Notably, the relationship between ZEB2 and
ACSL4 could form a positive feedback loop: ZEB2 transcriptionally activates ACSL4,
and conversely, ACSL4 stabilizes the ZEB2 protein by reducing its ubiquitination.

Amplification of either gene might therefore lock this loop in an active state, resulting
in the enhanced invasive and metastatic capabilities of breast cancer cells (Fig. 8G).
587

Previous studies have reported that ACSL4 could act as a tumor suppressor or
oncogene, depending on the specific cancer type and tissue environment(Cheng et
al., 2020; Grube et al., 2022; Lei et al., 2020; Ma et al., 2021; Zhang et al., 2021).
Indeed, ACSL4 could either be located at the cytomembrane or at the lipid droplets
and endoplasmic reticulum membrane, indicating the different functions of
ACSL4(Quan et al., 2021). The cytomembrane ACSL4 is likely responsible for
dictating ferroptosis sensitivity by shaping the plasma membrane lipidome, whereas
ACSL4 localized to lipid droplets has other functions. We provide evidence that
ACSL4, which is located in lipid droplets, is a pro-metastatic factor that promotes
invasion and migration. Notably, we demonstrated that ACSL4 depletion significantly
suppressed the invasion and migration of breast cancer cells in vitro and in vivo.
Furthermore, ACSL4 and ZEB2 were found to be preferentially expressed in
basal-like breast cancer cells and clinical samples that lacked ERα expression,
indicating that they are promising targets for highly invasive breast cancer. Survival
analysis revealed that high expression of ACSL4 or ZEB2 is associated with worse
post-progression survival than low expression in patients with breast cancer. It is
worth noting that there was no difference in the overall survival between ACSL4 or
ZEB2 high- and low-expression breast cancer patients (Supplementary Fig. S2),
suggesting that ACSL4 and ZEB2 could specifically be used as prognostic markers
for advanced breast cancer, and these axes hold promise as new metabolic
therapeutic targets for advanced breast cancer.
609

Although ACSL4 has been shown to play an essential role in metastasis in many
types of cancer (Maloberti et al., 2010; Sanchez-Martinez et al., 2017; Wu et al.,
2015), its mechanism of ACSL4-mediated metastasis is not fully understood. ACSL4
is a member of the long-chain fatty acetyl-CoA synthetase enzyme family that
catalyzes fatty acids to their active form, acyl-CoA, which can be directed to
anabolism or catabolism, depending on the cellular background. We observed that
the knockdown of ACSL4 significantly reduced the number and size of LD, indicating
that fatty acids were directed to anabolism to form LD by ACSL4. Indeed, it has been
shown that fatty acids are stored in LD before entering fatty acid oxidation, and the
formation of LD is required for fatty acids oxidation (Li et al., 2020; Petan, 2020; Welte
and Gould, 2017). This is in line with our observation that exogenous fatty acids (oleic
acid) significantly enhance lipid droplet accumulation and migration. Importantly,
lipidomic analysis revealed that the knockdown of ACSL4 significantly decreased the
incorporation of both saturated and unsaturated fatty acids into different species of
lipids, including triacylglycerols (TAG), phospholipids (PE, PC), and cholesteryl esters
(CE). Consistently, Frozen tissue sections from a metastatic animal model confirmed
that high ACSL4 expression was accompanied by apparent LD accumulation. These
results suggest that ACSL4 is a crucial regulator that promotes lipid storage during
breast cancer metastasis. Interestingly, knockdown of ACSL4 restored the expression

of E-cadherin, an essential adhesion molecule, indicating that ACSL4 promotes tumor
invasion through multiple mechanisms.

Reprogramming of lipid metabolism is an essential step in metastasis. Lipid droplets
are highly dynamic monolayer membrane-bound organelles involved in energy
utilization, signal transduction, and cancer invasion. The number and size of LD are
associated with cancer aggressiveness(Cruz et al., 2020). We observed that highly
invasive breast cancer cells were enriched with LD. Using fluorescence microscopy,
we noticed that the number of LD was significantly reduced after cell migration. Our
findings are in line with a recent study showing that LD undergo lipolysis during the
process of migration in pancreatic cancer(Rozeveld et al., 2020). It is likely that fatty
acids are released from lipid droplets and form acyl-CoA, which enters OXPHOS to
support the energy needed for metastasis. Therefore, these data support the notion
that increased LD accumulation is a priming state prior to metastasis, and that LD are
crucial resources to fuel the process of metastasis.

In addition, we observed that ACSL4 also participates in FA catabolism, as the
long-chain fatty acid-derived OCR rate and ATP production were significantly reduced
in ACSL4-depleted cells, indicating that ACSL4 is essential for FAO stimulation. Our
results are in line with a recent study that reported that hexokinase 2 enhances
tumorigenicity by activating the ACSL4-mediated fatty acid β-oxidation pathway (Li et
al., 2022). Notably, the expression of CPT1, a rate-limiting enzyme of FAO, was
significantly reduced in the ACSL4-depleted cells. Among the three isoforms of CPT1,
CPT1A is the only isoform regulated by ACSL4. We proposed that ACSL4 promotes
FAO and ATP production by upregulating CPT1A, thereby providing energy support
for breast cancer metastasis. Our results reveal the mechanism of previous findings
that FAO is a critical energy pathway in triple-negative breast cancer
(TNBC)(Camarda et al., 2016). In addition to the finding that ACSL4 promotes FAO to
fuel metastasis, we observed that lactate levels did not change after ACSL4
knockdown, suggesting that glycolysis is not involved in ACSL4-mediated energy
metabolism. Therefore, we propose a model in which cell migration is a multistep
process accompanied by dynamic cellular lipid metabolic changes. At the
pre-metastatic stage, ACSL4 enhances LD accumulation by promoting lipogenesis.
During metastasis, ACSL4 stimulates fatty acid oxidation to generate adenosine
triphosphate (ATP).

In conclusion, we provide insights into the mechanistic links between EMT and lipid
metabolism and identify the ZEB2/ACSL4 axis as a novel metastatic metabolic
pathway that stimulates both lipogenesis and fatty acid oxidation, resulting in
enhanced breast cancer invasion and metastasis. Importantly, our results
demonstrate that ACSL4 is a direct downstream target of ZEB2 in controlling lipid
storage and LD accumulation, which are important steps and energy pools for
metastasis. Clinically, our findings identified ZEB2-ACSL4 signalling as an attractive
therapeutic target for overcoming breast cancer metastasis. Elevated ACSL4 levels

can be used as an effective marker for predicting cancer progression in patients with
advanced breast cancer.

Additional information

Funding

Funder	Grant reference number	Author
Guangzhou Science and Technology Project	202201020240	Na Yang
Medical Research Foundation of Guangdong Province	A2021112	Na Yang
National Natural Science Foundation of China	81602434	Na Yang
Research Agreement between South China University of Technology and Guangzhou First People's Hospital	D9194290	Yuyou Duan
National Natural Science Foundation of China	81972594	Min Yan
The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication.		

Acknowledgements:

We acknowledge Professors Quentin Liu for the paclitaxel-resistant and
epirubicin-resistant MCF-7 cell lines and Dr. Jue Wang and Dr. Peilin Liao for their
technical support. We thank all members of Professor YY. Duan's lab for helpful
insights and valuable advice.

Authors' Contributions:

N. Yang and J. Lin designed the study. J. Lin, P. Zhang, J. Zhang, and W. Liu
performed experiments. W. Liu and G. Liu provided the clinical samples. N. Yang, J.
Lin, and P. Zhang analyzed the data. N. Yang, J. Lin, M. Yan, and Y. Duan wrote and
revised the manuscript. N. Yang, M. Yan and Y. Duan provided financial support.

Disclosure of Potential Conflicts of Interest

The authors declare no potential conflicts of interest.

Reference

Antunes, P., Cruz, A., Barbosa, J., Bonifacio, V.D.B., and Pinto, S.N. 2022. Lipid Droplets in
Cancer: From Composition and Role to Imaging and Therapeutics. Molecules *27*. DOI:
10.3390/molecules27030991, PMID:35164256
Barzaman, K., Karami, J., Zarei, Z., Hosseinzadeh, A., Kazemi, M.H., Moradi-Kalbolandi, S.,
Safari, E., and Farahmand, L. 2020. Breast cancer: Biology, biomarkers, and treatments. Int
Immunopharmacol *84*, 106535. DOI: 10.1016/j.intimp.2020.106535, PMID:32361569

Bian, X., Liu, R., Meng, Y., Xing, D., Xu, D., and Lu, Z. 2021. Lipid metabolism and cancer. J
Exp Med *218*. DOI: 10.1084/jem.20201606, PMID:33601415

Camarda, R., Zhou, A.Y., Kohnz, R.A., Balakrishnan, S., Mahieu, C., Anderton, B., Eyob, H.,
Kajimura, S., Tward, A., Krings, G.*, et al.* 2016. Inhibition of fatty acid oxidation as a therapy for
MYC-overexpressing triple-negative breast cancer. Nat Med *22*, 427-432. DOI:
10.1038/nm.4055, PMID:26950360

Chen, J., Ding, C., Chen, Y., Hu, W., Lu, Y., Wu, W., Zhang, Y., Yang, B., Wu, H., Peng, C.*, et*
al. 2020. ACSL4 promotes hepatocellular carcinoma progression via c-Myc stability mediated
by ERK/FBW7/c-Myc axis. Oncogenesis *9*, 42. DOI: 10.1038/s41389-020-0226-z,
PMID:32350243

Cheng, C., Geng, F., Cheng, X., and Guo, D. 2018. Lipid metabolism reprogramming and its
potential targets in cancer. Cancer Commun (Lond) *38*, 27. DOI:
10.1186/s40880-018-0301-4, PMID:29784041

Cheng, J., Fan, Y.Q., Liu, B.H., Zhou, H., Wang, J.M., and Chen, Q.X. 2020. ACSL4
suppresses glioma cells proliferation via activating ferroptosis. Oncol Rep *43*, 147-158. DOI:
10.3892/or.2019.7419, PMID:31789401

Corbet, C., Bastien, E., Santiago de Jesus, J.P., Dierge, E., Martherus, R., Vander Linden, C.,
Doix, B., Degavre, C., Guilbaud, C., Petit, L.*, et al.* 2020. TGFbeta2-induced formation of lipid
droplets supports acidosis-driven EMT and the metastatic spreading of cancer cells. Nat
Commun *11*, 454. DOI: 10.1038/s41467-019-14262-3, PMID:31974393

Cruz, A.L.S., Barreto, E.A., Fazolini, N.P.B., Viola, J.P.B., and Bozza, P.T. 2020. Lipid droplets:
platforms with multiple functions in cancer hallmarks. Cell Death Dis *11*, 105. DOI:
10.1038/s41419-020-2297-3, PMID:32029741

Doll, S., Proneth, B., Tyurina, Y.Y., Panzilius, E., Kobayashi, S., Ingold, I., Irmler, M., Beckers,
J., Aichler, M., Walch, A.*, et al.* 2017. ACSL4 dictates ferroptosis sensitivity by shaping cellular
lipid composition. Nat Chem Biol *13*, 91-98. DOI: 10.1038/nchembio.2239, PMID:27842070

Dongre, A., and Weinberg, R.A. 2019. New insights into the mechanisms of
epithelial-mesenchymal transition and implications for cancer. Nat Rev Mol Cell Biol *20*, 69-84.
DOI: 10.1038/s41580-018-0080-4, PMID:30459476

Fahad Ullah, M. 2019. Breast Cancer: Current Perspectives on the Disease Status. Adv Exp
Med Biol *1152*, 51-64. DOI: 10.1007/978-3-030-20301-6_4, PMID:31456179

Grube, J., Woitok, M.M., Mohs, A., Erschfeld, S., Lynen, C., Trautwein, C., and Otto, T. 2022.
ACSL4-dependent ferroptosis does not represent a tumor-suppressive mechanism but ACSL4
rather promotes liver cancer progression. Cell Death Dis *13*, 704. DOI:
10.1038/s41419-022-05137-5, PMID:35963845

Hua, W., Ten Dijke, P., Kostidis, S., Giera, M., and Hornsveld, M. 2020. TGFbeta-induced
metabolic reprogramming during epithelial-to-mesenchymal transition in cancer. Cell Mol Life
Sci *77*, 2103-2123. DOI: 10.1007/s00018-019-03398-6, PMID:31822964

Huang, W.C., Li, X., Liu, J., Lin, J., and Chung, L.W. 2012. Activation of androgen receptor,
lipogenesis, and oxidative stress converged by SREBP-1 is responsible for regulating growth
and progression of prostate cancer cells. Mol Cancer Res *10*, 133-142. DOI:
10.1158/1541-7786.MCR-11-0206, PMID:22064655

Jiang, L., Xiao, L., Sugiura, H., Huang, X., Ali, A., Kuro-o, M., Deberardinis, R.J., and
Boothman, D.A. 2015. Metabolic reprogramming during TGFbeta1-induced

epithelial-to-mesenchymal transition. Oncogene *34*, 3908-3916.   DOI: 10.1038/onc.2014.321,
PMID:25284588

Jin, C., and Yuan, P. 2020. Implications of lipid droplets in lung cancer: Associations with drug
resistance. Oncol Lett *20*, 2091-2104.   DOI: 10.3892/ol.2020.11769, PMID:32782526

Kang, H., Kim, H., Lee, S., Youn, H., and Youn, B. 2019. Role of Metabolic Reprogramming in
Epithelial(-)Mesenchymal Transition (EMT). Int J Mol Sci *20*.   DOI: 10.3390/ijms20082042,
PMID:31027222

Kuwata, H., and Hara, S. 2019. Role of acyl-CoA synthetase ACSL4 in arachidonic acid
metabolism.   Prostaglandins   Other   Lipid   Mediat   *144*,   106363.   DOI:
10.1016/j.prostaglandins.2019.106363, PMID:31306767

Lei, G., Zhang, Y., Koppula, P., Liu, X., Zhang, J., Lin, S.H., Ajani, J.A., Xiao, Q., Liao, Z.,
Wang, H.*, et al.* 2020. The role of ferroptosis in ionizing radiation-induced cell death and tumor
suppression. Cell Res *30*, 146-162.   DOI: 10.1038/s41422-019-0263-3, PMID:31949285

Li, H., Song, J., He, Y., Liu, Y., Liu, Z., Sun, W., Hu, W., Lei, Q.Y., Hu, X., Chen, Z.*, et al.* 2022.
CRISPR/Cas9 Screens Reveal that Hexokinase 2 Enhances Cancer Stemness and
Tumorigenicity by Activating the ACSL4-Fatty Acid beta-Oxidation Pathway. Adv Sci (Weinh) *9*,
e2105126.   DOI: 10.1002/advs.202105126, PMID:35603967

Li, Z., Liu, H., and Luo, X. 2020. Lipid droplet and its implication in cancer progression. Am J
Cancer Res *10*, 4112-4122.   DOI: PMID:33414989

Liu, H., Du, J., Chao, S., Li, S., Cai, H., Zhang, H., Chen, G., Liu, P., and Bu, P. 2022.
Fusobacterium nucleatum Promotes Colorectal Cancer Cell to Acquire Stem Cell-Like
Features by Manipulating Lipid Droplet-Mediated Numb Degradation. Adv Sci (Weinh) *9*,
e2105222.   DOI: 10.1002/advs.202105222, PMID:35170250

Luo, X., Cheng, C., Tan, Z., Li, N., Tang, M., Yang, L., and Cao, Y. 2017. Emerging roles of lipid
metabolism in cancer metastasis. Mol Cancer *16*, 76.   DOI: 10.1186/s12943-017-0646-3,
PMID:28399876

Luond, F., Sugiyama, N., Bill, R., Bornes, L., Hager, C., Tang, F., Santacroce, N., Beisel, C.,
Ivanek, R., Burglin, T.*, et al.* 2021. Distinct contributions of partial and full EMT to breast cancer
malignancy.   Dev   Cell   *56*,   3203-3221   e3211.   DOI: 10.1016/j.devcel.2021.11.006,
PMID:34847378

Ma, L.L., Liang, L., Zhou, D., and Wang, S.W. 2021. Tumor suppressor miR-424-5p abrogates
ferroptosis in ovarian cancer through targeting ACSL4. Neoplasma *68*, 165-173.   DOI:
10.4149/neo_2020_200707N705, PMID:33038905

Maan, M., Peters, J.M., Dutta, M., and Patterson, A.D. 2018. Lipid metabolism and lipophagy
in cancer. Biochem Biophys Res Commun *504*, 582-589.   DOI: 10.1016/j.bbrc.2018.02.097,
PMID:29438712

Maloberti, P.M., Duarte, A.B., Orlando, U.D., Pasqualini, M.E., Solano, A.R., Lopez-Otin, C.,
and Podesta, E.J. 2010. Functional interaction between acyl-CoA synthetase 4,
lipooxygenases and cyclooxygenase-2 in the aggressive phenotype of breast cancer cells.
PLoS One *5*, e15540.   DOI: 10.1371/journal.pone.0015540, PMID:21085606

Mittal, V. 2018. Epithelial Mesenchymal Transition in Tumor Metastasis. Annu Rev Pathol *13*,
395-412.   DOI: 10.1146/annurev-pathol-020117-043854, PMID:29414248

Mosier, J.A., Schwager, S.C., Boyajian, D.A., and Reinhart-King, C.A. 2021. Cancer cell
metabolic plasticity in migration and metastasis. Clin Exp Metastasis *38*, 343-359.   DOI:

10.1007/s10585-021-10102-1, PMID:34076787

Niu, Z., Chen, G., Chang, W., Sun, P., Luo, Z., Zhang, H., Zhi, L., Guo, C., Chen, H., Yin, M.*, et

al.* 2021. Chimeric antigen receptor-modified macrophages trigger systemic anti-tumour

immunity. J Pathol *253*, 247-257.   DOI: 10.1002/path.5585, PMID:33140856

Orlando, U.D., Castillo, A.F., Medrano, M.A.R., Solano, A.R., Maloberti, P.M., and Podesta, E.J.

2019. Acyl-CoA synthetase-4 is implicated in drug resistance in breast cancer cell lines

involving the regulation of energy-dependent transporter expression. Biochem Pharmacol *159*,

52-63.   DOI: 10.1016/j.bcp.2018.11.005, PMID:30414939

Park, M., Kim, D., Ko, S., Kim, A., Mo, K., and Yoon, H. 2022. Breast Cancer Metastasis:

Mechanisms and Therapeutic Implications. Int J Mol Sci *23*.   DOI: 10.3390/ijms23126806,

PMID:35743249

Petan, T. 2020. Lipid Droplets in Cancer. Rev Physiol Biochem Pharmacol.   DOI:

10.1007/112_2020_51, PMID:33074407

Quan, J., Bode, A.M., and Luo, X. 2021. ACSL family: The regulatory mechanisms and

therapeutic implications in cancer. Eur J Pharmacol *909*, 174397.   DOI:

10.1016/j.ejphar.2021.174397, PMID:34332918

Roman, M., Wrobel, T.P., Panek, A., Paluszkiewicz, C., and Kwiatek, W.M. 2020. Lipid droplets

in prostate cancer cells and effect of irradiation studied by Raman microspectroscopy. Biochim

Biophys Acta Mol Cell Biol Lipids *1865*, 158753.   DOI: 10.1016/j.bbalip.2020.158753,

PMID:32504818

Rossi Sebastiano, M., and Konstantinidou, G. 2019. Targeting Long Chain Acyl-CoA

Synthetases for Cancer Therapy. Int J Mol Sci *20*.   DOI: 10.3390/ijms20153624,

PMID:31344914

Rozeveld, C.N., Johnson, K.M., Zhang, L., and Razidlo, G.L. 2020. KRAS Controls Pancreatic

Cancer Cell Lipid Metabolism and Invasive Potential through the Lipase HSL. Cancer Res *80*,

4932-4945.   DOI: 10.1158/0008-5472.CAN-20-1255, PMID:32816911

Sanchez-Martinez, R., Cruz-Gil, S., Garcia-Alvarez, M.S., Reglero, G., and Ramirez de Molina,

A. 2017. Complementary ACSL isoforms contribute to a non-Warburg advantageous energetic

status characterizing invasive colon cancer cells. Sci Rep *7*, 11143.   DOI:

10.1038/s41598-017-11612-3, PMID:28894242

Sciacovelli, M., and Frezza, C. 2017. Metabolic reprogramming and epithelial-to-mesenchymal

transition in cancer. FEBS J *284*, 3132-3144.   DOI: 10.1111/febs.14090, PMID:28444969

Senga, S., Kobayashi, N., Kawaguchi, K., Ando, A., and Fujii, H. 2018. Fatty acid-binding

protein 5 (FABP5) promotes lipolysis of lipid droplets, de novo fatty acid (FA) synthesis and

activation of nuclear factor-kappa B (NF-kappaB) signaling in cancer cells. Biochim Biophys

Acta Mol Cell Biol Lipids *1863*, 1057-1067.   DOI: 10.1016/j.bbalip.2018.06.010,

PMID:29906613

Snaebjornsson, M.T., Janaki-Raman, S., and Schulze, A. 2020. Greasing the Wheels of the

Cancer Machine: The Role of Lipid Metabolism in Cancer. Cell Metab *31*, 62-76.   DOI:

10.1016/j.cmet.2019.11.010, PMID:31813823

Soukupova, J., Malfettone, A., Bertran, E., Hernandez-Alvarez, M.I., Penuelas-Haro, I., Dituri,

F., Giannelli, G., Zorzano, A., and Fabregat, I. 2021. Epithelial-Mesenchymal Transition (EMT)

Induced by TGF-beta in Hepatocellular Carcinoma Cells Reprograms Lipid Metabolism. Int J

Mol Sci *22*.   DOI: 10.3390/ijms22115543, PMID:34073989

Taib, B., Aboussalah, A.M., Moniruzzaman, M., Chen, S., Haughey, N.J., Kim, S.F., and Ahima,
R.S. 2019. Lipid accumulation and oxidation in glioblastoma multiforme. Sci Rep *9*, 19593.
DOI: 10.1038/s41598-019-55985-z, PMID:31863022

Welte, M.A., and Gould, A.P. 2017. Lipid droplet functions beyond energy storage. Biochim
Biophys Acta Mol Cell Biol Lipids *1862*, 1260-1272.   DOI: 10.1016/j.bbalip.2017.07.006,
PMID:28735096

Wright, H.J., Hou, J., Xu, B., Cortez, M., Potma, E.O., Tromberg, B.J., and Razorenova, O.V.
2017. CDCP1 drives triple-negative breast cancer metastasis through reduction of lipid-droplet
abundance and stimulation of fatty acid oxidation. Proc Natl Acad Sci U S A *114*, E6556-E6565.
DOI: 10.1073/pnas.1703791114, PMID:28739932

Wu, X., Deng, F., Li, Y., Daniels, G., Du, X., Ren, Q., Wang, J., Wang, L.H., Yang, Y., Zhang, V.*,
et al.* 2015. ACSL4 promotes prostate cancer growth, invasion and hormonal resistance.
Oncotarget *6*, 44849-44863.   DOI: 10.18632/oncotarget.6438, PMID:26636648

Yang, L., Zhang, F., Wang, X., Tsai, Y., Chuang, K.H., Keng, P.C., Lee, S.O., and Chen, Y.
2016. A FASN-TGF-beta1-FASN regulatory loop contributes to high EMT/metastatic potential
of cisplatin-resistant non-small cell lung cancer. Oncotarget *7*, 55543-55554.   DOI:
10.18632/oncotarget.10837, PMID:27765901

Yang, N., Wang, C., Wang, Z., Zona, S., Lin, S.X., Wang, X., Yan, M., Zheng, F.M., Li, S.S., Xu,
B.*, et al.* 2017. FOXM1 recruits nuclear Aurora kinase A to participate in a positive feedback
loop essential for the self-renewal of breast cancer stem cells. Oncogene *36*, 3428-3440.
DOI: 10.1038/onc.2016.490, PMID:28114286

Zembroski, A.S., Andolino, C., Buhman, K.K., and Teegarden, D. 2021. Proteomic
Characterization of Cytoplasmic Lipid Droplets in Human Metastatic Breast Cancer Cells.
Front Oncol *11*, 576326.   DOI: 10.3389/fonc.2021.576326, PMID:34141606

Zhang, Y., Li, S., Li, F., Lv, C., and Yang, Q.K. 2021. High-fat diet impairs ferroptosis and
promotes cancer invasiveness via downregulating tumor suppressor ACSL4 in lung
adenocarcinoma. Biol Direct *16*, 10.   DOI: 10.1186/s13062-021-00294-7, PMID:34053456

Zheng, F.M., Long, Z.J., Hou, Z.J., Luo, Y., Xu, L.Z., Xia, J.L., Lai, X.J., Liu, J.W., Wang, X.,
Kamran, M.*, et al.* 2014. A novel small molecule aurora kinase inhibitor attenuates breast
tumor-initiating cells and overcomes drug resistance. Mol Cancer Ther *13*, 1991-2003.   DOI:
10.1158/1535-7163.MCT-13-1029, PMID:24899685

Figure Legends

Figure 1, The expression and relationship of ZEB2 and ACSL4 in breast cancer. **A,** Heat
maps of the 38 up-regulated EMT-related genes detected by RNA-seq analysis in the
paclitaxel-resistant MCF-7 cell line (TAXOL) and Epirubicin-resistant MCF-7 cell line (EPI)
compared to wild-type MCF-7 cell line. **B,** The correlation between ACSL4 and ZEB2
mRNA expression in the TCGA cohort consisting of 1222 breast cancer patient samples.
Spearman correlation and linear regression analysis were employed. **C,** The correlation
between ACSL4 and ERα mRNA expression in the TCGA cohort consisting of 666 breast
cancer patient samples. Spearman correlation and linear regression analysis were
employed. **D,** PPS (post-progression survival) was examined by Kaplan–Meier analysis to
compare the survival rates in ACSL high and low expression of breast cancer patients. **E,**

PPS（post-progression survival）was examined by Kaplan–Meier analysis to compare the

survival rates in ZEB2 high and low expression of breast cancer patients. **F,** Expression of

ACSL4 and ZEB2 were analyzed by western blot in a panel of 5 breast cancer cell lines,

including two basal-like, two luminal and one taxol-resistant cell. **G,** Expression of ACSL4

and ZEB2 were analyzed by western blot in a panel of nine cases of tumour samples,

including 5 cases of luminal (Patient 1, 2, 3, 4, 5) and 4 cases of triple-negative breast

cancer (Patient 6, 7, 8, 9). **H,** Scatter plot with linear regression analysis from expression

level of G. **I,** HE staining and IHC analysis of ACSL4, ZEB2, ERα expression in

representative basal-like and luminal subtype breast cancer tissues. Representative

pictures were shown. Scale bar, 50 μm.

Figure 2, Over-expression of ACSL4 contributes to ZEB2-mediated breast cancer

invasion. **A,** Cell metastatic capacity was analyzed by wound healing assay in control or

ACSL4 over-expression MCF-7 cells (Left panel). Quantification of relative migration

distance (Right panel). Scale bar, 5mm. **B,** Cell invasive capacity was analyzed by

transwell invasion assay in control or ACSL4 over-expression MCF-7 cells (Left panel).

Quantification of relative migration distance (Right panel). Scale bar, 1mm. **C,** Cell

metastatic capacity was analyzed by wound healing assays in control or ACSL4 silencing

MDA-MD-231 cells (shACSL4) (Left panel). Quantification of relative migration distance

(Right panel). Scale bar, 5mm. **D,** Cell invasive capacity was analyzed by transwell

invasion assay in control or ACSL4 silencing MDA-MD-231 cells (shACSL4) (Left panel).

Quantification of relative migration distance (Right panel). Scale bar, 1mm. **E,** KEGG

pathway enrichment analysis of differentially expressed genes by RNA-sequencing

between control and ACSL4 knockdown MDA-MB-231 cells. The top 10 deferential

pathways were listed. **F,** Expression of three EMT-related genes, E-cadherin, N-cadherin,

and vimentin, was analyzed by western blotting in control, ACSL4, or ZEB2 silencing

MDA-MB-231 cells. **G,** Cell invasive and metastatic capacity were analyzed by transwell

invasion assay and wound healing assays (Scale bar, 5mm) in control and ZEB2

knockdown, or ZEB2 knockdown cells that over-express ACSL4. **H,** Graphs indicated the

Statistical analysis in G analyzed by Student's t-test (mean ± SEM). ** P<0.01, ***

P<0.001, **** P<0.0001. All results are from three or four independent experiments.

Figure 3, ACSL4 promotes lipid droplets accumulation and lipogenesis. **A,** BODIPY

493/503 staining of lipid droplets in MCF-7, Taxol-resistant MCF-7 cells (TAXOL-7), or

MDA-MB-231 cells (Left panel). Quantification of normalized lipid contents (Right panel).

Scale bar, 5μm. **B,** BODIPY 493/503 staining of lipid droplets in control or ACSL4

knockout MDA-MB-231 cells (Upper panel). Quantification of normalized fluorescence

intensity and percentage of LD-containing cell number (Lower panel). Scale bar, 50μm. **C,**

BODIPY 493/503 staining of lipid droplets in control or ZEB2 knockout MDA-MB-231 cells

(Upper panel). Quantification of normalized fluorescence intensity and percentage of

LD-containing cell number (Lower panel). Scale bar, 50μm. **D,** The heatmap of

representative down-regulated lipid species (TG, PE, PC) with hierarchical clustering in

the control cells and ACSL4 Knockdown cells. Each species was normalized to the

corresponding mean value, as determined by two-way ANOVA. **E,** Quantification of
different FA containing TG, PC, and PE species in control or ACSL4 Knockdown
MDA-MD-231 cells. * P<0.05, ** P<0.01, *** P<0.001, **** P<0.0001. Error bars, SD.

Figure 4, Exogenous lipids promote LD accumulation and fuel cell migration. **A,** BODIPY
493/503 staining of lipid droplets in control cells or oleic acid (OA) loaded (200 µM)
MDA-MB-231 cells. Scale bar, 10 µm. **B,** Quantification of normalized lipid contents from
the conditions in (A). **C,** Cell invasive capacity was analyzed by transwell invasion assay
in control or oleic acid (OA) loaded (200 µM) MDA-MB-231 cells. Cells invaded for 16
hours through a Matrigel-coated filter toward high-serum media (Left panel).
Quantification of relative invasive cells (Right panel). Scale bar, 1 mm. **D,** Cell metastatic
capacity was analyzed by wound healing assays in control or oleic acid (OA) loaded (200
µM) MDA-MB-231 cells. Scale bar, 5 mm. **E,** Cells were seeded in a transwell chamber.
After 24 hours, cells migrated to the lower side of the chamber, and the fluorescence
intensity per cell was calculated (Left panel). Quantitation of total LD area per cell before
and after cell migration (Right panel). Scale bar, 10 µm (small) and 50 µm (big). ***
P<0.001, **** P<0.0001.

Figure 5, ACSL4 and ZEB2 increase fatty acid oxygen consumption and promote ATP
generation in BLBC Cells. **A,** Reactome pathway analysis of differentially expressed
genes by RNA-sequencing between control and ACSL4 knockdown MDA-MB-231 cells.
The representative pathways were shown. **B,** Heatmap showing the differentially
expressed genes between control cells and ACSL4 knockdown cells. The representative
FAO-related genes were shown. **C,** The mRNA levels of CPT1A, CPT1B, and CPT1C
were analyzed by quantitative PCR in control and ACSL4 knockdown cells. **D,**
Quantitation of the normalized oxygen consumption rate (OCR) for long-chain fatty acids
was monitored by Agilent XF Substrate Oxidation Stress Test in control or ACSL4
knockdown cells. Specific inhibitors were added as indicated. **E,** Quantitation of the
normalized oxygen consumption rate (OCR) for long-chain fatty acids was monitored by
Agilent XF Substrate Oxidation Stress Test in control or ZEB2 knockdown cells. Specific
inhibitors were added as indicated. **F and G,** ATP production was quantified in control or
ACSL4 knockdown cells (Left) or ZEB2 knockdown cells (Right). **H and I,** Lactate
production was examined in control and ACSL4 knockdown (shACSL4) or ZEB2
knockdown (shZEB2) cells. Data are represented as mean SEM of three independent
experiments, analyzed by Student's t-test, * P<0.05, **** P<0.0001.

Figure 6, ZEB2 activates ACSL4 expression by directly binding to its promoter. **A,**
Relative mRNA levels of ZEB2 and ACSL4 in control or ZEB2 knockdown MDA-MB-231
cells. **B,** Protein levels of ZEB2 and ACSL4 in control or ZEB2 knockdown MDA-MB-231
cells. **C and D,** Truncated ACSL4 promoter segment activity analyzed by Luciferase
reporter assay in control or ZEB2 over-expressed 293T cells. **E,** The specific primers
designed for ACSL4 promoter according to the E-box position shown in C (set1 to -287bp,
set2 to -995, set3 to -1038, set4 to -1116). **F,** ChIP assay analysis of the occupation of
ZEB2 on ACSL4 promoter by using four primers as indicated in E. **G,** Quantitative PCR

analysis of ZEB2-binding abundance of specific ACSL4 promoter region by using set1
primer indicated in F. Genomic DNA was purified after chromatin immunoprecipitation and
analyzed by Quantitative PCR. **H,** Luciferase reporter analysis of the activity of wild-type
ACSL4 promoter or its mutants in control or ZEB2 over-expression 293T cells. Data are
represented as mean SD of three independent experiments, analyzed by Student's t-test,
** $P<0.01$, *** $P<0.001$, **** $P<0.0001$.

Figure 7, ACSL4 regulates ZEB2 mRNA expression and protein stability. **A,** Relative
mRNA levels of ZEB2 and ACSL4 in control or ACSL4 silencing MDA-MB-231 cells. **B,**
Protein levels of ZEB2 and ACSL4 in control or ACSL4 silencing MDA-MB-231 cells as
indicated. **C,** Ubiquitylation of ZEB2 was examined by in vitro ubiquitin assay. 293T cells
were co-transfected with indicated constructs. Cells were treated with MG132 for 6 hours
before IP. Anti-MYC was used to pull down the ZEB2 protein. The polyubiquitinated ZEB2
protein was detected by an anti-HA antibody. **D,** The interaction between ACSL4 and
ZEB2 was detected by Co-IP assay. 293T cells were co-transfected with ZEB2 and
ACSL4 expressing construct. Anti-HA antibody was used to pull down the protein. **E**, The
stability of ZEB2 protein was detected by CHX treatment assay in control or ACSL4
silencing MDA-MB-231 cells. Cells were treated with 100 μg/mL cycloheximide (CHX) and
were harvested at the indicated times after the addition of CHX. GAPDH was used as the
internal loading control. **F,** Quantification of stability assays shown in E. **G,** The stability of
ZEB2 protein was detected by CHX treatment assay in control or ACSL4 over-expressed
MDA-MB-231 cells. GAPDH was used as the internal loading control. **H,** Quantification of
stability assays shown in G. *** $P<0.001$.

Figure 8, ACSL4 knockdown attenuates lung metastasis of breast cancer. **A,** Cell
metastatic capacity was determined by xenograft experiment in vivo. Control cells and
ACSL4 knockdown MDA-MB-231 cells were mixed with Matrigel in a 1:1 ratio and injected
into NSG mice. Lung metastatic burden in mice was quantified by bioluminescence
imaging at the experimental endpoint (Left panel). Quantification of normalized gray area
(Right panel). **B,** Representative xenograft tumor images and HE staining pictures of
metastatic nodules in lungs (pointed by black arrows) were shown. The right panel shows
the quantification of metastatic lung nodules in two groups, as indicated. **C,** The image of
xenograft tumors developed from the control and ACSL4 knockdown cells. **D,** The tumor
volumes were measured and calculated at the indicated time. **E,** Representative IHC
images of lung metastatic lesions (Left panel) and quantification is shown (Right panel).
Scale. Bar, 50μm. **F,** The representative images of fluorescence assay for ACSL4 and lipid
droplets (Left panel). The fluorescence intensity of lipid droplets and ACSL4 were
calculated in the right panel. Scale. Bar, 100μm. **G,** A proposed mechanism to illustrate
the positive feedback loop of ZEB2 and ACSL4 regulates lipid metabolism, which results
in enhanced breast cancer metastasis. Data are represented as mean SEM of three
independent experiments and analyzed by Student's t-test, * $P<0.05$, *** $P<0.001$, ****
$P<0.0001$.

1004

1005 **Supplementary Figure Legends**

Supplementary Figure S1: The representative differential genes between wild-type MCF-7 cells, paclitaxel-resistant MCF-7 cells (TAXOL), and Epirubicin-resistant MCF-7 cells (EPI).

A, The heat map of the up-regulated genes in drug-resistant cells was shown. **B,** The heat map of the down-regulated genes in the drug-resistant cell was shown.

Supplementary Figure S2: Overall survival of breast cancers was examined by Kaplan–Meier analysis to compare the survival rates between ACSL high- or ZEB2 high- and low-expression breast cancer patients.

Supplementary Figure S3: The metastatic and invasive capacities were reduced in ZEB2 knockdown MDA-MD-231 cells (shZEB2).

A, Wound healing assays were performed in control cells or ZEB2 knockdown (shZEB2) MDA-MD-231 cells. Scale bar, 5 mm. **B,** Transwell invasive assay was performed in control cells or ZEB2 knockdown (shZEB2) MDA-MD-231 cells. Scale bar, 1 mm. *** $P<0.001$, **** $P<0.0001$.

Supplementary Figure S4: Fluorescence staining of lipid droplets and ACSL4 in MDA-MB-231 cells. Scale bar, 10 μm.

Supplementary Figure S5: ACSL4 regulates lipid composition in BLBC cells.

A, The heat map of the down-regulated lipid species in ACSL4 knockdown cells (shACSL4) compared with those in control cells. **B,** Quantification of different species of lipids in two groups as indicated. * $P<0.05$, ** $P<0.01$.

Supplementary Figure S6: ACSL4 regulates the expression of lipid metabolic genes.

A. Western blot analysis of CPT1A expression in control and ACSL4 knockdown, or ZEB2 knockdown MDA-MD-231 cells. **B,** Relative mRNA levels of ATGL, FASN and SREBP2 in control or ACSL4 knockdown MDA-MB-231 cells (shACSL4). ** $P<0.01$, *** $P<0.001$.

Figure1

Figure2

Figure 3

Figure4

Figure 5

Figure 6

Figure 7

Figure 8