Title: **Experimentally induced active and quiet sleep engage non-overlapping transcriptional programs in *Drosophila***

Short title: **Active and quiet sleep in *Drosophila***

Authors: Niki Anthoney[1], Lucy A.L. Tainton-Heap[1], Hang Luong[2], Eleni Notaras[1], Amber B. Kewin[1], Qiongyi Zhao[1], Trent Perry[2], Philip Batterham[2], Paul J. Shaw[3] & Bruno van Swinderen[1*]

Author Affiliation:

[1] Queensland Brain Institute, The University of Queensland, Brisbane, QLD 4072 Australia

[2] School of BioSciences, The University of Melbourne, Melbourne, VIC 3052 Australia

[3] Department of Neuroscience, Washington University School of Medicine, St. Louis, MO USA

*Corresponding Author : b.vanswinderen@uq.edu.au

Abstract

Sleep in mammals can be broadly classified into two different physiological categories: rapid eye movement (REM) sleep and slow wave sleep (SWS), and accordingly REM and SWS are thought to achieve a different set of functions. The fruit fly *Drosophila melanogaster* is increasingly being used as a model to understand sleep functions, although it remains unclear if the fly brain also engages in different kinds of sleep as well. Here, we compare two commonly used approaches for studying sleep experimentally in *Drosophila*: optogenetic activation of sleep-promoting neurons and provision of a sleep-promoting drug, Gaboxadol. We find that these different sleep-induction methods have similar effects on increasing sleep duration, but divergent effects on brain activity. Transcriptomic analysis reveals that drug-induced deep sleep ('quiet' sleep) mostly downregulates metabolism genes, whereas optogenetic 'active' sleep upregulates a wide range of genes relevant to normal waking functions. This suggests that optogenetics and pharmacological induction of sleep in *Drosophila* promote different features of sleep, which engage different sets of genes to achieve their respective functions.

Introduction

There is increasing evidence that sleep is a complex phenomenon in most animals, comprising of distinct stages that are characterized by dramatically different physiological processes and brain activity signatures [1, 2]. This suggests that different sleep stages, such as rapid eye movement (REM) and slow-wave sleep (SWS) in humans and other mammals [3] are accomplishing distinct functions that are nevertheless collectively important for adaptive behavior and survival [4]. While REM and SWS appear to be restricted to a subset of vertebrates (e.g., mammals, birds, and possibly some reptiles [5-7] a broader range of animals, including invertebrates, demonstrate evidence of 'active' versus 'quiet' sleep [1, 2, 8]. Whether active and quiet sleep represent evolutionary antecedents of REM and SWS, respectively, remains speculative [1, 2]. However, during active sleep, although animals are less responsive, brain recordings reveal a level of neural activity that is similar to wakefulness, in contrast to quiet sleep, which is characterized by significantly decreased neural activity in invertebrates [9, 10] as well as certain fish [11], mollusks [12], and reptiles [6].

Although it is likely that even insects such as fruit flies and honeybees sleep in distinct stages [13, 14], sleep studies using the genetic model *Drosophila melanogaster* still mostly measure sleep as a single phenomenon, defined by 5 minutes (or more) of inactivity [15, 16]. As sleep studies increasingly employ *Drosophila* to investigate molecular and cellular processes underpinning potential sleep functions, this simplified approach to measuring sleep in flies carries the risk of overlooking different functions accomplished by distinct kinds of sleep. Sleep physiology and functions are increasingly being addressed in the fly model by imposing experimentally controlled sleep regimes, either pharmacologically or via transient control of sleep-promoting circuits by using opto- or thermogenetic tools [17]. Yet, there is little knowledge available on whether these different approaches are producing qualitatively similar sleep. For example, sleep can be induced genetically in flies by activating sleep-

promoting neurons in the central complex (CX) – a part of the insect brain that has been found to be involved in multimodal sensory processing [18]. In particular, the dorsal fan-shaped body (dFB) of the CX has been proposed to serve as a discharge circuit for the insect's sleep homeostat, whereby increased sleep pressure (e.g., due to extended wakefulness) alters the physiological properties of dFB neurons causing them to fire more readily and thereby promote decreased behavioral responsiveness [19] and thus sleep [20-22]. Crucially, activation of sleep-promoting circuits including the dFB have been shown to be sleep-restorative [10, 23], but confusingly, brain recordings during optogenetically-induced sleep, via electrophysiology as well as whole-brain calcium imaging techniques, reveal wake-like levels of brain activity [9, 10]. This suggests that some approaches to optogenetically-induced sleep might be promoting a form of sleep akin to the 'active' sleep stage detected during spontaneous sleep [9, 10].

An alternate way to induce sleep in *Drosophila* is by feeding flies drugs that have been designed to treat insomnia in humans, such as the GABA-agonist 4,5,6,7-tetrahyrdoisoxazolopyridin-3-ol, (THIP), also known as Gaboxadol [24]. Several studies have shown that THIP-induced sleep in flies is also restorative and achieves key functions ranging from memory consolidation to cellular repair and waste clearance [23, 25-27]. This pharmacological approach centered on GABA function has a solid foundation based on better-understood sleep processes: in mammals, many sleep-inducing drugs also target GABA receptors, and this class of drugs tends to promote SWS [28]. In contrast, there are no obvious drugs that promote REM sleep, although local infusion of cholinergic agonists (e.g., carbachol) to the brainstem has been shown to induce REM-like states in cats [29].

In this study, we compare THIP-induced sleep with optogenetically-induced sleep in *Drosophila*, using behavior, brain activity, and transcriptomics. To ensure the validity of our comparisons, we performed all of our experiments in the same genetic background, employing a canonical Gal4 strain that expresses a transgenic cation channel in a sleep-promoting circuit:

R23E10-Gal4 > UAS-Chrimson [30, 31]. When these flies are fed all-trans-retinal (ATR) and

then exposed to red light, they are put to sleep optogenetically. When these flies are instead fed

THIP, they are put to sleep pharmacologically. By using the same genetic background, we were

thus able to contrast the effects of either kind of sleep at the level of behavior, brain activity,

and gene expression (**Figure 1**). We questioned how similar either form of induced sleep was.

Materials and Methods

Animals

Drosophila melanogaster flies were reared in vials (groups of 20 flies / vial) on standard yeast-

sugar-agar based media (1.0-1.5-0.5 gram ratio) under a 12:12 light/dark (8 AM:8 PM) cycle

and maintained at 25°C with 50% humidity. Adult, 3-5 day-old female, flies were used for all

experiments and randomly assigned to experimental groups. Fly lines used for behavioral and

RNA-sequencing experiments include R23E10-Gal4 (attp2; Bloomington 49032;

Bloomington Drosophila Stock Center, Bloomington, Indiana) and UAS-CsChrimson-mVenus

(attp18; Bloomington 55134; Provided by Janelia Research Campus, Ashburn, Virginia)[31].

For all 2-photon experiments, flies with the genotype 10XUAS-Chrimson88-tdTomato

(attp18) / +: LexAop-nlsGCaMP6f (VIE-260b; kindly provided by Barry J. Dickson) / +: Nsyb-

LexA (attP2) [32], LexAop-PAGFP (VK00005) / R23E10-Gal4 were used. Optogenetically-

manipulated fly lines were maintained on food containing 0.5mg/ml all-trans retinal (ATR;

Merck, Darmstadt, Germany) for 24 hours prior to assay to allow for sufficient consumption.

Pharmacologically-manipulated flies were maintained on food with 0.1 mg/ml of Gaboxadol

(4,5,6,7-tetrahyrdoisoxazolopyridin-3-ol, THIP) for the duration of behavioral experiment

[23].

2-photon imaging

2-photon imaging was performed as described previously [10] using a ThorLabs Bergamo

series 2 multiphoton microscope. Fluorescence was detected with a High Sensitivity GaAsP

photomultiplier tube (ThorLabs, PMT2000). GCaMP fluorescence was filtered through the

microscope with a 594 dichroic beam splitter and a 525/25nm band pass filter.

For imaging experiments, flies were secured to a custom-built holder [10]. Extracellular fluid

(ECF) containing 103 NaCl, 10.5 trehalose, 10 glucose, 26 NaHCO3, 5 C6H15NO6S, 5 MgCl2

(hexa-hydrate), 2 Sucrose, 3 KCl, 1.5 CaCl (dihydrate), and 1 NaH2PO4 (in mM) at room

temperature was used to fill a chamber over the head of the fly. The brain was accessed by

removing the cuticle of the fly with forceps, and the perineural sheath was removed with a

microlance. Flies were allowed to recover from this for one hour before commencement of

experiments. Imaging was performed across 18 z-slices, separated by 6µm, with two additional

flyback frames. The entire nlsGCaMP6f signal was located within a 256 x 256px area,

corresponding to 667 x 667µm. Fly behavior was recorded with a Firefly MV 0.3MP camera

(FMVU-03MTM-CS, FLIR Systems), which was mounted to a 75mm optical lens and an

infrared filter. Camera illumination was provided by a custom-built infrared array consisting

of 24 3mm infrared diodes. Behavioral data was collected for the duration of all experiments.

For THIP experiments, an initial five minutes of baseline activity was captured, followed by

perfusion of 0.2mg/ml THIP in ECF onto the brain at a rate of 1.25ml/minute for five minutes.

An additional twenty minutes of both brain and behavioral activity were recorded to allow

visualization of the fly falling asleep on the ball as a result of THIP exposure. All flies were

removed from imaging after an experiment and confirmed to have awoken by visual inpection.

A subset of THIP-exposed flies (n=3) remained in the imaging setup to measure brain activity

upon recovery.

Behavioral responsiveness probing

For probing behavioral responsiveness in the brain imaging preparation, flies walking on an air-supported ball were subjected to a 50ms long, 10psi air puff stimulus, which was generated using a custom-built apparatus and delivered through a 3mm-diameter tube onto the front of the fly. Flies were subjected to 10 pre-THIP stimuli at a rate of one puff/minute, to characterize the baseline response rate. Flies were then perfused with 0.2mg/ml THIP in ECF for five minutes, followed by continuous ECF perfusion for the remaining experimental time. Five minutes after the fly had fallen asleep on the ball, a further 20 air puff stimuli were delivered, at a rate of one puff/minute. Behavioral responses to the air puff were noted as a 'yes' (1) or 'no' (0), which were characterized as the fly rapidly walking on the ball immediately following the air puff. For statistical analysis, the pre-THIP condition was compared to either the first or last 10 minutes of the post-THIP condition.

Imaging analysis

Preprocessing of images was carried out using custom written Matlab scripts and ImageJ. Motion artifacts of the images were corrected as described previously [10]. Image registration was achieved using efficient sub-pixel image registration by cross-correlation. Each z-slice in a volume (18 z-slices and 2 flyback slices) is acquired at a slightly different time point compared to the rest of the slices. Hence to perform volume (x,y,z) analysis of images, all the slices within a volume need to be adjusted for timing differences. This was achieved by using the 9^{th} z-slice as the reference slice and temporal interpolation was performed for all the other z-slices using 'sinc' interpolation. The timing correction approach implemented here is conceptually similar to the methods using in fMRI for slice timing correction.

For each individual z-slice, a standard deviation projection of the entire time series was used for watershed segmentation with the 'Morphological segmentation' ImageJ plugin [33]. Using

a custom-written MatLab (Mathworks) code, the mean fluorescent value of all pixels within a

given ROI were extracted for the entire time series, resulting in a n x t array for each slice of

each experiment, where n refers to the number of neurons in each Z-slice, and t refers to the

length of the experiment in time frames. These greyscale values were z-scored for each neuron,

and the z-scored data was transformed into a binary matrix where a value of > 3 standard

deviations of the mean was allocated a '1', and every value < 3 standard deviations was

allocated a '0'. To determine whether a neuron fired during the entire time series, a rolling sum

of the binary matrix was performed, where ten consecutive time frames were summed together.

If the value of any of these summing events was greater or equal to seven (indicating a

fluorescent change of > 3 standard deviations in 7/10 time frames), a neuron was deemed to be

active. For THIP sleep experiments, the five minutes of inactivity occurring after an initial 30

seconds of behavioral inactivity were used. After identifying firing neurons for each condition

(wake vs sleep), the percentage of active neurons was calculated in each slice by taking the

number of active neurons and dividing it by the total number of neurons.

Traces of active neurons were used to calculate the number of firing events. This was done

using the 'findpeaks' matlab function on the zscored fluorescent traces, with the parameters

'minpeakheight' of 3, and 'minpeakdistance' of 30. Data resulting from this was crosschecked

by taking the binary matrices of the time traces and finding the number of times each neuron

met the activity threshold described above. Graph-theory analyses of neural connectivity were

performed as described previously [10] .

Behavioral sleep analysis

Behavioral data for flies in imaging experiments was analyzed as previously [10] using a

custom-written MatLab code that measured the pixel change occurring over the legs of the fly

on the ball over the entire time series. Data was analyzed and graphed using Graphpad Prism.

All data was checked for Gaussian distribution using a D'Agostino-Pearson normality test prior to statistical testing. Data from THIP experiments was analyzed using a non-parametric Mann-Whitney test.

Sleep behavior in freely-walking flies was analyzed with the Drosophila ARousal Tracking system (DART) as previously described [34]. Prior to analysis, 3-5 day-old females were collected and loaded individually into 65 mm glass tubes (Trikinetics) that were plugged at one end with our standard fly food (see above), containing either 0.1 mg/ml THIP or 0.5 mg/ml all-trans-retinal (ATR). Controls were placed onto normal food and housed under identical conditions as the experimental groups. The tubes were placed onto platforms (6 total platforms, 17 tubes per platform, up to 102 flies total) for filming. THIP-fed flies were monitored for 3 days/nights, and sleep data were averaged as no significant differences were detected across successive days. Optogenetically-activated were tracked for baseline day without red light, after which they were exposed to ultra-bright red LED (617 nm Luxeon Rebel LED, Luxeon Star LEDs, Ontario, Canada) which produce 0.1-0.2mW/mm2 at a distance of 4-5 cm with the aid of 723 concentrator optics (Polymer Optics 6° 15 mm Circular Beam Optic, Luxeon Star LEDs) for the duration of the experiment for optogenetic activation. Significance was determined by ANOVA with Tukey's multiple comparisons test (GraphPad Prism). Sleep analysis in nAchRα knockout animals was performed using Trikinetics beam-crossing devices, with regular (>5min) and short sleep (1-5min) calculated as previously [10].

Sleep deprivation

Flies were sleep deprived (SD) with the use of the previously described Sleep Nullifying Apparatus (SNAP), an automated sleep deprivation apparatus that has been found to keep flies awake without nonspecifically activating stress responses [35]. Vials containing no greater than 20 flies, which contained either standard food medium or medium containing 0.1mg/ml THIP

were placed on the SNAP apparatus for continuous sleep deprivation. The SNAP apparatus

was programmed to snap the flies once every 20 seconds for the duration of the sleep

deprivation protocol.

RNA-Sequencing

Flies collected for RNA-sequencing analysis were first housed in vials containing either

0.5mg/ml all-trans retinal (ATR) or 0.1mg/ml THIP for sleep induction, along with their

genetically identical controls on standard food medium. Flies undergoing sleep induction by

optogenetic activation with ATR and their controls were placed under constant red-light from

8AM until 6PM to coincide with normal 12:12 light/dark cycles. Flies were collected after 1

hour (ZT 1) and 10 hours (ZT 10) post induction for immediate brain dissection and RNA

extraction. For analysis of pharmacological sleep induction, flies were placed on THIP or

normal food medium at 8AM (ZT 0) and collected for dissection at 6PM (ZT 10).

Whole fly brains were dissected in ice cold RNA*later (Sigma-Aldrich)* with 0.1% PBST as per

previously published protocol [36]. The dissected brains were immediately pooled into five

1.5-mL Eppendorf tubes containing 5 brains (n = 25) each. Total RNA was immediately

purified using TRIzol according to the manufacturer's protocols (Sigma-Aldrich) and stored at

-80°C until commencement of RNA-sequencing.

cDNA libraries were prepared using the Illumina TruSeq stranded mRNA library prep kit.

Image processing and sequence data extraction were performed using the standard Illumina

Genome Analyzer software and CASAVA (version 1.8.2) software. Cutadapt (version 1.8.1)

was used to cut the adaptor sequences as well as low quality nucleotides at both ends. When a

processed read is shorter than 36 bp, the read was discarded by cutadapt, with the parameter

setting of "-q 20,20 --minimum-length=36". Processed reads were aligned to the *Drosophila*

melanogaster reference genome (dm6) using HISAT2 (version 2.0.5) [37], with the parameter

setting of "--no-unal --fr --rna-strandness RF --known-splicesite-infile dm6_splicesites.txt".

This setting is to i) suppress SAM records for reads that failed to align ("--no-unal"), ii) specify

the Illumina's paired-end sequencing assay and the strand-specific information ("--fr --rna-

strandness RF") and iii) provide a list of known splice sites in *Drosophila melanogaster* ("--

known-splicesite-infile dm6_splicesites.txt"). Samtools (version 1.3) [38] was then used to

convert "SAM" files to "BAM" files, sort and index the "BAM" files. The "htseq-count"

module in the HTSeq package (v0.7.1) was used to quantitate the gene expression level by

generating a raw count table for each sample (i.e. counting reads in gene features for each

sample). Based on these raw count tables, edgeR (version 3.16.5) [39] was adopted to perform

the differential expression analysis between treatment groups and controls. EdgeR used a

trimmed mean of M-values to compute scale factors for library size normalization [40]. It used

the Cox-Reid profile-adjusted likelihood method to estimate dispersions [41] and the quasi-

likelihood F-test to determine differential expression [42]. Lowly expressed genes in both

groups (the mean CPM < 5 in both groups) were removed. Differentially expressed genes were

identified using the following criteria: i) FDR < 0.05 and ii) fold changes > 1.5 (or logfc >0.58).

Gene ontology enrichment analysis for differentially expressed genes was performed using the

functional annotation tool in DAVID Bioinformatics Resources (version 6.8) [43, 44].

Gene expression

<u>RNA and cDNA Synthesis</u>

A quantitative reverse transcriptase PCR assay was used to confirm expression of genes

enriched during THIP sleep induction. Nineteen candidate genes were selected (eight

negatively and eleven positively) for the gaboxadol (THIP) sleep analysis and six genes (four

negatively and two positively) for the dFSB activation experiments. Total RNA was isolated

using the Directzol RNA kit (ZymoResearch) from twenty adult brains per condition and each

condition was collected in triplicate. RNA quality was confirmed using a microvolume

spectrophotometer NanoDrop 2000 (Thermo, USA) with only those resulting samples meeting

optimal density ratios between 1.8 and 2.1 used. Up to 1 μg of total RNA was reverse

transcribed using a High-Capacity cDNA Reverse Transcription Kit (Themo, USA) as per the

manufacturer's protocols. The synthesis of cDNA and subsequent amplification was performed

in max volumes of 20 μL per reaction using the T100 Thermal Cycler (Bio-Rad, USA).

Thermocycle conditions were as such; 25 °C for 10 min, 37 °C for 120 min, 85 °C for 5 min,

and held at 4 °C . All cDNA was subsequently stored at − 20 °C until used. Target genes for

THIP experiments included Pxt (CG7660, FBgn0261987), RpS5b (CG7014, FBgn0038277),

Dhd (CG4193, FBgn0011761), CG9377 (CG9377, FBgn0032507), aKHr (CG11325,

FBgn0025595), Acox57D-d (CG9709, FBgn0034629), FASN1 (CG3523, FBgn0283427), Pen

(CG4799, FBgn0287720), CG10513 (CG10513, FBgn0039311), Gasp (CG10287,

FBgn0026077), Act57B (CG10067, FBgn0000044), Bin1 (CG6046, FBgn0024491), verm

(CG8756, FBgn0261341), CG16885 (CG16885, FBgn0032538), CG16884 (CG16884,

FBgn0028544), CG5999 (CG5999, FBgn0038083), Fbp1 (CG17285, FBgn0000639), CG5724

(CG5724, FBgn0038082), Eh (CG5400, FBgn0000564). Target genes for dFSB experiments

included Vmat (CG33528, FBgn0260964), Dop1R1 (CG9652, FBgn0011582), Salt (CG2196,

FBgn0039872), Dysb (CG6856, FBgn0036819), Irk3 (CG10369, FBgn0032706), Blos1

(CG30077, FBgn0050077). Housekeeping genes included Rpl32 (CG7939, FBgn0002626),

Gapdh2 (CG8893, FBgn0001092), Actin 5C (CG4027, FBgn0000042). Primer sequences can

be found in Supplementary Table 8.

<u>Quantitative real-time PCR</u>

Quantitative (q) RT-PCR was carried out using the Luna Universal qPCR Master Mix (NEB)

in the CFX384 Real-Time system (Bio-Rad, USA). Cycling conditions were: 1. 95°C for 60 s,

2. 95°C for 15s, 3. 60°C for 60s with 39 cycles of steps two and three. Melt curve analysis was

then performed with the following conditions 1. 95°C for 15s, 2. 60°C for 60s, 3. 95°C for 15s.

Three biological replicates for each condition as well as three technical replicates per biological

sample were loaded. Each experiment was then repeated on three separate occasions. Cq values

and standard curves were generated using Bio Rad CFX Manager Software to ensure

amplification specificity. Results were normalized to the above housekeeping genes and gene

expression was calculated following the $2^{-\Delta\Delta Cq}$ method (Livak and Schmittgen 2001).

Gene knockouts and knockdowns

Dα1KO harboured an ends-out mediated deletion of Dα1 in a w[1118] background with the X

chromosome replaced with one from the wild type line DGRP line 59 [45]. For Dα2KO, 7kb

of the Da2 locus was removed using the endogenously expressing Actin-Cas9 strain (BDSC

5490) crossed to a strain expressing two Dα2 gene locus specific sgRNAs (Dα_sgRNA_start_1

5'CATGTTTAGCGCTGCAATGC, Dα2_sgRNA_end_1 5'TTACAAGCCATCTGCCTAG).

Transgenic flies were generated using the pCFD4-U6:1_U6:3tandem gRNAs construct. This

was a gift from Simon Bullock. For Dα3KO (*Dα3Δ1020*), Dα4KO (*Dα4ΔBA*), Dα6KO,

Dα7KO (*Dα7ΔD6*), two sgRNAs were designed to target the start and the end of the coding

sequence and cloned into either pU6-BbsI-gRNA or pCFD4 plasmids. These plasmids were

then microinjected into *Drosophila* embryos to generate transgenic strains stably expressing

sgRNAs. These strains were crossed to another strain expressing Cas9 under Actin promoter

(ActinCas9). Their offspring were screened for deletion events with PCR and crossed to

appropriate balancer strains to isolate and generate homozygous knockout strains. Full

deletions were identified for all these subunit genes except for Dα3 which has two partial

deletions at the 3' and 5' ends; these were verified to be true knockouts by the lack of RNA

expression [46]. ActinCas9 strain was used as genetic control for Dα3KO and Dα7KO, while

this same strain with the X chromosome replaced with one from w^{1118} (w^{1118}ActinCas9) was

used as genetic control for Dα2KO, Dα4KO, and Dα6KO. The RNAi strain for the adipokinetic

hormone receptor gene knockdown experiments (UAS-*AkhR*-RNAi) was obtained from the

VDRC (KK109300). This RNAi construct was expressed in neurons across the fly brain by

crossing the UAS flies to R57C10-Gal4 [30]. RNAi strains for the nicotinic alpha receptor gene

knockdown experiments (UAS- nAchRα1-7-RNAi) were obtained from the BDRC. These

were: nAChRα1 RNAi – BL# 28688 - y[1] v[1]; P{y[+t7.7] v[+t1.8]=TRiP.JF03103}attP2 ,

nAChRα2 RNAi – BL# 27493 - y[1] v[1]; P{y[+t7.7] v[+t1.8]=TRiP.JF02643}attP2 ,

nAChRα3 RNAi – BL# 27671 – y[1] v[1]; P{y[+t7.7] v[+t1.8]=TRiP.JF02750}attP2 ,

nAChRα4 RNAi – BL# 31985 - y[1] v[1]; P{y[+t7.7] v[+t1.8]=TRiP.JF03419}attP2 ,

nAChRα5 RNAi – BL# 77418 - y[1] sc[*] v[1] sev[21]; P{y[+t7.7]

v[+t1.8]=TRiP.HMC06550}attP2 ,

nAChRα6 RNAi – BL# 52885 - y[1] sc[*] v[1] sev[21]; P{y[+t7.7]

v[+t1.8]=TRiP.HMC03623}attP40 ,

nAChRα7 RNAi – BL# 27251 - y[1] v[1]; P{y[+t7.7] v[+t1.8]=TRiP.JF02570}attP2.

These RNAi constructs were expressed in the sleep-promoting neurons by crossing the UAS

flies to R23E10-Gal4.

Code and data availability

The data and analysis tools underpinning this study is stored in The University of Queensland

Research Data Management (UQRDM) via UQ eSpace, the institutional data storage repository

of the University of Queensland (UQ). Datasets in this study are also available from the Lead

Contact upon request. All Matlab analysis code for brain imaging have been made available in

a previous publication [10], and are also available from the Lead Contact upon request.

Results

Prolonged optogenetic and THIP-induced sleep have near identical effects on sleep duration

We first compared pharmacological and optogenetic sleep (**Figure 1**) by using the traditional behavioral metrics employed by most *Drosophila* sleep researchers: >5 minutes inactivity for flies confined in small glass tubes over multiple days and nights [15, 16]. We found that optogenetic- and THIP-induced sleep yielded almost identical effects on sleep duration, with both significantly increasing total sleep duration for both the day and night, when compared to controls (**Figure 2A-D**; **Supplementary Table 1**). An increase in total sleep duration can be due to either an increase in the number of sleep bouts that are occurring (reflective of more fragmented sleep), or an increase in the average duration of individual sleep bouts, which indicates a more consolidated sleep structure [47-49]. To investigate whether both sleep induction methods also had similar effects on sleep architecture, we plotted bout number as a function of bout duration for optogenetic and THIP-induced sleep, for the day and night [50]. We found that both optogenetic activation and THIP provision produce a similar increase in sleep consolidation during the day (**Figure 2E, F; Figure 3**). During the night, induced sleep effects were also similar, although less clearly different to the spontaneous sleep seen in control flies (**Figure 2G, H; Figure 3**). Interestingly, red light exposure decreased average night bout duration in non-ATR control flies (**Figure 3A-D**), suggesting a light-induced artefact at night. For THIP, we observed an increase in both bout number and duration during the day, and an increase in bout duration during the night (**Figure 3E,F**). Taken together, these results show that prolonged optogenetic activation and THIP provision have similar behavioral effects on induced sleep duration in *Drosophila*, although some differences in night bout architecture were noted (**Figure 3**). Without any further investigations, this might suggest that both sleep induction methods represent similar underlying processes that uniformly increase sleep in flies.

THIP-induced sleep decreases brain activity and connectivity

The brain presents an obvious place to look for any potential differences between sleep

induction methods. In a previous study employing whole-brain calcium imaging in tethered

flies we showed that optogenetic activation of the R23E10 circuit promotes wake-like sleep,

with neither neural activity levels nor connectivity metrics changing significantly even after

15min of optogenetically-induced sleep [10]. We therefore utilized the same fly strain as in

that study (R23E10-Gal4>UAS-Chrimson88-tdTomato;Nsyb-LexA>LexOp-nlsGCaMP6f) to

examine the effect of THIP-induced sleep on brain activity (**Figure 4A,B**). Since we were

interested in comparing acute sleep induction effects on brain activity (as opposed to

prolonged sleep induction effects on behavior, as in Figures 2,3), we adapted our calcium

imaging approach to allow a brief perfusion of THIP directly onto the exposed fly brain

(**Figure 4A**, see Methods). As done previously for examining optogenetically-induced sleep

[10], we examined calcium transients in neural soma scanning across 18 optical slices of the

central fly brain (**Figure 4B,** left) and identified regions of interest (ROIs) corresponding to

neuronal soma in this volume (**Figure 4B**, right, and see Methods). As shown previously

[10], optogenetically activating the R23E10 circuit renders flies asleep (albeit twitchy at

times) without changing the average level of neural activity measured this way (**Figure 4C**).

We have previously shown that even 15min of R23E10 activation fails to produce a 'quiet'

sleep stage that is evident after 5min in spontaneously sleeping flies [10], suggesting that this

manipulation promotes ongoing active sleep until the red light is turned off – although this

has not been investigated for longer epochs, e.g. the chronic activation experiments described

in Figure 2. To determine the effect of THIP on neural activity in the exact same strain, we

transiently perfused onto the fly brain the minimal THIP dosage required to reliably promote

sleep in flies within five minutes (0.2mg/ml) [9]. In contrast to optogenetic-induced sleep, we

observed overall decreased neural activity coincident with the flies falling asleep, and flies

remained asleep well after the drug was washed out (**Figure 4D**). To ensure that we were

actually putting flies (reversibly) to sleep in this preparation, we probed for behavioral

responsiveness by puffing air onto the fly once every minute (50 ms duration, 10 psi) (**Figure**

5A). Since the time when flies fell asleep following five minutes of THIP perfusion could be

variable [9], arousal probing during sleep was only initiated after 5 min of complete

quiescence (**Figure 5A**, behavioral responsiveness testing). We observed decreased

arousability for flies that had been induced to sleep via THIP perfusion (**Figure 5B**). Drug-

induced sleep was however reversible, with flies returning to baseline levels of behavioral

responsiveness to the air puffs ~20-30 min after sleep initiation. This confirmed that the brief

exposure to THIP was indeed putting flies to sleep, with an expected sleep inertia lasting the

length of a typical spontaneous sleep bout [15, 16]. This contrasts with optogenetic sleep

induction using the R23E10-Gal4 circuit, which does not show appear to show any sleep

inertia (**Figure 4C**) [51].

We then examined more closely neural activity in flies that had been put to sleep with THIP.

We found that neural activity decreased rapidly within 5 min after sleep onset (**Figure 5C**,

+THIP, early). Correlation analysis also revealed a decrease in connectivity among the

remaining active neurons (**Figure 5D**, +THIP, early). We also analyzed the next 5 min of

sleep and observed similar results (**Figure 5C,D**, +THIP, mid). All flies eventually woke up

from THIP-induced sleep, and brain activity returned to wake levels in three flies that were

recorded throughout (**Figure 5C**). These observations suggest that acute THIP exposure is

promoting rapid entry into a 'quiet' sleep stage in flies, bypassing the wake-like sleep evident

during the first 5 min of spontaneous sleep onset, and closely resembling spontaneous quiet

sleep typically seen after 5 minutes of inactivity [10]. Importantly, THIP-induced sleep

appears to be dissimilar from optogenetically-induced sleep in this genotype, at the level of

neural activity as well as connectivity [10].

In recent work we showed that rendering flies unresponsive with a general anesthetic,

isoflurane, in surprising contrast to THIP, does not quieten the fly brain [51]. Isoflurane also

decreased neural connectivity, as well as the diversity of neural activity across the fly brain,

whereas optogenetic-induced sleep did not show any differences in neural activity or

ensemble dynamics [51]. Since in our current THIP experiments we were similarly recording

from neural soma that we could track through time, we were able to assess the level of

overlap between the neurons that remained active during THIP-induced sleep and

wakefulness (**Figure 5E, F**). We found that ~30% of active neurons during THIP-induced

sleep were also active during wake (**Figure 5F, G**). We next examined whether the same

neurons remained active across successive 5min epochs during THIP-induced sleep

compared to wake. We found that there is significantly more overlap between successive

5min sleep epochs (41%), compared to the waking average (**Figure 5G**), suggesting less

neural turnover during THIP-induced sleep than during wake. This contrasts with optogenetic

sleep, where the rate of neural turnover was not different from wake [51]. Taken together, our

calcium imaging data confirm that pharmacological sleep induction promotes a different kind

of sleep than optogenetic sleep induction in the same strain. Henceforth, we call this 'quiet'

sleep, in contrast to the 'active' sleep that seems to be engaged by optogenetic activation [2,

10]. Notably, calcium imaging of spontaneous sleep bouts in *Drosophila* also revealed active

and quiet sleep stages [10], suggesting that both of our experimental approaches are

physiologically relevant. Whether drug perfusion to the brain is equivalent to feeding is of

course less clear. When feeding on 0.1mg/ml THIP-laced food, flies were continuously

exposed to the drug over days, with comparatively less reaching the brain. With perfusion,

the brain was directly exposed to 0.2mg/ml THIP for only 5 minutes. Interestingly, in both

cases this induces daytime sleep bouts which average around 25min (**Figure 3F, Figure 5B**),

the average duration of a spontaneous night-time sleep bout (**Figure 3F; Table S1**).

Transcriptional analysis of flies induced to sleep by THIP provision

Our calcium imaging experiments suggest that different biological processes might be

engaged by optogenetic-induced sleep compared to THIP-induced sleep. Additionally, we

observed neural effects encompassing much of the fly brain (**Figure 5E**), as our recording

approach exploited a pan-neural driver. We therefore wondered if either sleep-induction

method might lead to differences in gene expression across the whole brain, and if these

might highlight distinct molecular pathways engaged by either kind of sleep. To address this,

we collected brains from flies that had been induced to sleep by either method, and compared

the resulting transcriptomes with identically-handled control animals that had not been

induced to sleep by these methods.

To control for genetic background, we again used the same R23E10-Gal4 > UAS-Chrimson

flies as in our multi-day behavioral experiments and fed the flies either THIP or ATR, as in

Figure 2. We only examined daytime sleep-induction effects for either method, as this is when

we observed the greatest increase in sleep compared to controls (**Figure 2**), and previous work

has shown that daytime sleep induction using either method achieves sleep functions [10, 23].

We present our THIP results first. Since THIP is a GABA-acting drug that probably affects a

variety of processes in the brain aside from sleep, we also assessed the effect of THIP on flies

that were prevented from sleeping (**Figure 6A**, left panel). Sleep deprivation (SD) was

performed by mechanically arousing flies once every 20 seconds for the duration of the

experiment, on a 'SNAP' apparatus [23, 35]. RNA was extracted from the brains of all groups

of flies (+/- THIP, +/- SD) after 10 hours of daytime (8am-6pm) THIP (or vehicle) provision.

Samples for RNA-sequencing were collected in replicates of 5 to ensure accuracy, and any

significant transcriptional effects were thresholded at a log fold change of 0.58 (see Methods).

Flies allowed to eat food containing 0.1mg/ml THIP *ad lib* over 10 daytime hours led to 129

significant changes in gene expression compared to vehicle-fed controls, with the large

majority (110) being downregulated and only 19 upregulated (**Figure 6B,C,E;**

Supplementary Table 2). In contrast, when THIP-fed flies were prevented from sleeping

this led to mostly upregulated genes (88 upregulated vs 21 downregulated, **Figure 6B,D,F;**

Supplementary Table 3), with many of the upregulated genes showing a high level of fold-

change in expression. Not surprisingly, preventing sleep in THIP-fed flies led to an almost

entirely non-overlapping set of gene expression changes (**Figure 6B**). This suggests that the

large number of down-regulated genes in +THIP / -SD flies pertain to sleep processes rather

than the effect of ingesting THIP, since only a few (9) genes overlapped with the +THIP /

+SD dataset which revealed mostly upregulated genes.

Gene Ontology analysis on genes that were downregulated as a result of THIP-induced sleep

highlighted a significant enrichment of metabolism pathways (**Figure 6E,F; Supplementary**

Figure 1). The top Gene Ontology biological processes included primary, organic substance,

cellular, biosynthetic and nitrogen compound metabolic pathways, as well as ribosomal

processes. Interestingly, these downregulated processes are largely consistent with a recently

published mouse sleep transcriptome study [52]. Among the metabolism pathways uncovered

in this dataset we observed over-representation of expected genes such as *bgm* (bubblegum

CG4501) and *Acer* (Angiotensin-converting enzyme-related CG10593). Both of these genes

are found in the primary metabolic and organic substance metabolic processes as well as within

the Sleep Gene Ontology dataset (GO:0030431). Another downregulated metabolic gene is AkhR (adipokinetic hormone receptor), which has been found to be involved in starvation-induced sleep loss in *Drosophila* [53]. AkhR belongs to the Class A GPCR Neuropeptide and protein hormone receptors which are a gene class involved in storage fat mobilization, analogous to the glucagon receptor found in mammals [54].

Although THIP-induced sleep overwhelmingly led to gene downregulation, a few genes (19) were significantly upregulated. Gene Ontology analysis on these upregulated genes highlighted enrichment in varying groups including developmental processes and multicellular organismal processes (**Figure 6E,F; Supplementary Figure 1**). Some groups were enriched under the organic substance metabolic process pathways; however, there was no overlap when comparing these to the pathways enriched due to downregulation of genes. There were some overlapping enriched pathways when we compared the gene sets from sleep-deprived flies which had also been treated with THIP (**Figure 6F**; **Supplementary Figure 2**). However, the gene sets they involve are upregulated in the SD dataset but downregulated in sleeping flies. Interestingly, the non-sleeping THIP dataset uncovered a significant enrichment of pathways involved in the response to stress. This might be expected for flies exposed to regular mechanical stimuli over 10 hours. None of these pathways featured in the THIP sleep dataset.

To validate these findings, we conducted qRT-PCR analyses on 19 genes from our THIP sleep dataset and compared these results to our original transcriptional data. The genes represented a range of both up – and down-regulated genes, and we found convincing correspondence between the groups (**Figure 5G**). qRT-PCR comparisons with RNAseq data and associated statistics are presented in **Supplementary Table S9**.

Transcriptional analysis of flies induced to sleep by optogenetic activation

We next examined the effect of optogenetically-induced sleep on the whole-brain transcriptome, to compare to our THIP-induced sleep data. Based on our earlier findings that showed that optogenetic activation results in rapidly inducible sleep behavior that consolidates over at least 12 daytime sleep hours (**Figure 2C,E,G**), as well as our previous study showing that 10 daytime hours of optogenetic activation corrects attention defects in sleep-deprived flies [10], we induced sleep in R23E10-Gal4 x UAS-Chrimson flies for 10 daytime hours and collected tissue for whole-brain RNA-sequencing (**Figure 7A**). We selected two time points for collection, for both the sleep-induced flies (+ATR) as well as their genetically identical controls that were not fed ATR (-ATR; **Figure 7A**). Optogenetic activation was matched to the normal day-time light cycle (8 AM – 8 PM). The first collection point was after 1 hour (ZT1, 9 AM) of red-light exposure, to control for effects of ATR provision (when compared to ZT1, -ATR controls) as well as to uncover any potential short-term genetic effects of optogenetic activation. We then collected flies after 10 hours of red-light exposure (ZT 10, 6 PM) to examine longer-term genetic effects of optogenetic sleep induction, and to match exactly our THIP sleep collection timepoint (i.e., 10 hours of induced daytime sleep by either method). The combined collection points also allowed us to compare transcriptional profiles between conditions (e.g., ZT10 +ATR vs. ZT10 -ATR), to identify sleep genes, as well as within conditions (ZT1 vs. ZT10), to account for genetic effects potentially linked to light-dark cycles. As for the THIP sleep data in the same strain, samples for RNA-sequencing were collected in replicates of 5 to ensure accuracy, and any significant transcriptional effects were thresholded at a log fold change of 0.58 (see Methods).

We first examined the effect of 10 hours of daytime optogenetic-induced sleep. Here, we compared ATR-fed R23E10-Gal4 x UAS-Chrimson flies to genetically identical animals that

were also exposed to red light for 10 hours but not provided with ATR in their food (ATR-). The control flies were therefore never induced to sleep by optogenetic activation, although they were still able to sleep spontaneously (see **Figure 2C,E,G**). We found that 10 hours of optogenetic activation led to 278 significant transcriptional changes, comprising mostly of upregulated genes, with 171 upregulated compared to 107 downregulated (**Figure 7B,C,E; Supplementary Table 4**). In contrast to the THIP-induced sleep dataset, transcriptional analysis of 10hr optogenetic sleep induction uncovered a variety of different processes predominantly related to the regulation of biological and cellular processes, rather than metabolism specifically (**Figure 7E; Supplementary Figure 3**). For example, of the genes that were overexpressed there is an enrichment of the Semaphorin-plexin signaling pathway (GO:0071526, GO:1902287, GO:1902285 and GO:2000305) and the ephrin receptor signaling pathway (GO:0046011), both of which are known to be involved in axonal guidance (**Figure 7F**). Interestingly, several upregulated genes code for different subunits of nicotinic acetylcholine receptors (nAchRα1,3,4 &5; **Figure 7C**). Importantly, there was almost no overlap with our sleep deprivation dataset (**Supplementary Table 3**), suggesting that optogenetic activation is not sleep depriving (or stressing) the flies (only one upregulated gene was shared, CG40198). Of the genes that were downregulated there is enrichment of pathways that relate to synaptic vesicle recycling (GO:0036465 and GO:0036466) as well as neurotransmitter metabolic processes (GO:0042133) (**Figure 7F; Supplementary Figure 3**).

In contrast to the 10hr timepoint, 1 hr of optogenetically-induced sleep had far fewer transcriptomic consequences, with only 17 genes upregulated and 10 downregulated (**Figure 7B,D**). This small number of transcriptomic changes (see **Supplementary Table 5**) may reflect the effect of ATR feeding, rather than any genes relevant to optogenetic sleep. That 9

hours of additional optogenetic sleep increased transcriptomic changes by an order of

magnitude lends confidence to the interpretation that relevant genes linked to prolonged

optogenetic activation are being engaged.

To account for potential genetic effects linked to light-dark expression cycles, we compared

transcriptional profiles between 10 hours of optogenetic-induced sleep to 1 hour of induced

sleep. Here, we found 220 differentially regulated genes (119 upregulated and 101

downregulated) when comparing ATR-fed flies at both time points (ZT10 vs ZT1,

Supplementary Table 6). Since the 1-hour group was collected in the morning and the 10-

hour group was collected in the evening, we expected this dataset to expose a number of

circadian-regulatory genes, and this is indeed what we found (**Supplementary Figure 4A,B**).

We then compared these results with a parallel ZT10 vs ZT1 experiment where flies were not

fed ATR. Here we uncovered 503 differentially expressed genes (252 upregulated and 251

downregulated) when comparing flies that had not been fed ATR at both timepoints

(**Supplementary Table 7**). Importantly, there were 98 genes that overlapped between these

independent Z10 vs ZT1 datasets, suggesting commonalities linked to circadian processes.

Indeed, GO Pathway analysis of Biological Processes revealed a number of genes involved in

the regulation of the circadian rhythm among these 98 overlapping genes, including the well-

known circadian genes *period, timeless, clockwork-orange, clock and vrille*. Notably, co-

factors *period* and *timeless* are both upregulated whereas *clk* is downregulated, and this is

replicated in both independent datasets (**Supplementary Figure 4C**). This correspondence

with expectations for zeitgeber or light-dark effects provides a level of confidence that our

respective sleep datasets are highlighting transcriptomic changes and biological pathways

relevant to either sleep induction approach. Notably, there was no overlap at all in gene

expression changes between optogenetic-induced sleep and THIP-induced sleep (**Supplementary Tables 2 & 4**), and the respective GO pathways analyses of biological processes are also largely non-overlapping (**Supplementary Figure 5**).

To validate these findings, we compared our transcriptional results with qRT-PCR on six genes. This included the dopamine receptor Dop1R1, which regulates arousal levels [55] as well as the schizophrenia susceptibility gene dysbindin (*Dysb*), which has been shown to regulate dopaminergic function [56]. We found convincing correspondence between our qRT-PCR data and our transcriptomic data (**Figure 7G**).). qRT-PCR comparisons with RNAseq data and associated statistics are presented in **Supplementary Table S9**.

Nicotinic acetylcholine receptors regulate sleep architecture

While THIP-induced sleep caused a systemic downregulation of metabolism-related genes, the effect of optogenetic-induced sleep on gene expression was not dominated by a single category. This may be consistent with our earlier observation that brain activity looks similar to wake during optogenetic-induced sleep [10], so we could essentially be highlighting biological processes relevant to an awake fly brain, such as dopamine function [57]. However, optogenetic activation of the R23E10 circuit is not like wake, in that flies are rendered highly unresponsive to external stimuli, so perhaps like REM sleep in mammals a different category of molecular processes could be involved. In mammals, acetylcholine generally promotes wakefulness and alertness, but activity of cholinergic neurons is also high during REM sleep [58]. Neurotransmission in the insect brain is largely cholinergic, with 7 different nicotinic 'alpha' receptor subunits expressed in neural tissue [59]. Interestingly, four

of these subunits were significantly upregulated in our optogenetic sleep dataset: nAchRα1,

nAchRα3, nAchRα4, and nAchRα5 (with nAchRα6 and nAchRα7 approaching

significance, **Supplementary Table 4**). For comparison, none of these were upregulated in

our sleep deprivation dataset, suggesting a sleep-relevant role. Previous studies have

demonstrated a role for some of these same receptor subunits in sleep regulation, in particular

nAchRα4 (also called *redeye*) which is upregulated in short-sleeping mutants [60] and

nAchRα3 which has been reported to regulates arousal levels in flies [61]. Together, these

studies suggest processes that might be reconsidered in the context of active sleep, as

highlighted by our gene expression findings. We therefore sought to examine the role of

cholinergic signalling in sleep more closely, by knocking out each nAchRα subunit and

examining effects of each subunit knockout on sleep architecture. Since our transcriptomic

analysis encompassed effects of active sleep on whole-brain gene expression, we first

eliminated each nAchRα subunit across the brain, by testing confirmed genetic deletions [46,

62].

We first examined the effect of each nAchRα subunit deletion on sleep duration, using the 5-

minute criterion for quantifying sleep in *Drosophila* [15]. We found that the nAchRα mutants

fell into two different categories: removal of nAchRα1 and nAchRα2 significantly decreased

sleep, day and night; whereas removal of nAchRα3, nAchRα4, nAchRα6 and nAchRα7

significantly increased sleep, day and night (**Figure 8A**). The nAchRα5 knockout was found

to be homozygous lethal, so was not included. To examine sleep architecture in these

mutants, we quantified sleep bout number and duration and plotted these together as done

previously for our sleep induction experiments (**Figure 2**). Examining the data this way, it is

clear to see how nAchRα1 and nAchRα2 are different: most sleep bouts are very short, day

and night (**Figure 8B**, top 2 rows, left panels, green dots). In contrast, knocking out the other

alpha subunits seems to consolidate sleep, especially at night (**Figure 8B**, bottom 4 rows, left

panels). nAchRα3 knockouts were most striking in this regard, with these flies sleeping

uninterrupted for an average of 156.53 minutes (± 18.06) during the day and 160.76 minutes

(± 17.92) at night. Increased sleep consolidation in these mutants was however not due to

lack of activity. While awake, nAchRα3 animals were just as active as controls (activity per

waking minute = 2.69 ± 0.18 versus 2.5 ± 0.06, respectively). However, some knockouts did

increase waking activity levels; waking activity data for all of the knockout strains and their

genetic controls are presented in **Supplementary Figure 6**.

We next questioned what kind of sleep the nAchRα knockout flies might be getting. In

previous work we have shown that flies can be asleep already after the first minute of

spontaneous inactivity, and that during the first five minutes of sleep the fly brain displays

wake-like levels of neural activity [10]. We have termed this early sleep stage 'active sleep'

to distinguish it from 'quiet sleep' that typically follows after 5-10 minutes [63]. One way of

estimating the amount of 'active sleep' in *Drosophila* flies is to sum all short sleep epochs

lasting between 1-5 minutes and expressing this as a percentage of total sleep [10]. When we

re-examined our nAchRα knockouts in this way, we found that this behavioral readout for

'active sleep' was significantly affected by the loss of select nAchRα subunits. Short sleep

increased significantly during both the day and the night in nAchRα1 and nAchRα2 (**Figure

8B**, top 2 rows, right panel, green dots). In contrast, and consistent with our sleep architecture

analyses (above), nAchRα3 displayed almost no short sleep (**Figure 8B**, row 3, right panel).

Finally, in nAchRα4 and nAchRα6 short sleep was significantly decreased at night, while in

nAchRα7 short sleep was significantly decreased day and night (**Figure 8B**, rows 4-6, right

panel). In conclusion, every one of the nAchRα knockouts we tested affect short sleep in

some way, either increasing (nAchRα1 and nAchRα2) or decreasing it (nAchRα3,

nAchRα4, nAchRα6, nAchRα7). Additionally, tallying the proportion of short sleep bouts

provides valuable insight into altered sleep architecture in mutant strains.

We nevertheless questioned whether these systemic effects of nicotinic receptors on short

sleep were perhaps a trivial consequence of altered >5min sleep duration in these mutants,

especially regarding the striking differences between nAchRα1&2 and the other subunit

knockouts. We therefore returned to our 'quiet' sleep (THIP) dataset to contrast a gene

derived from that study. We had found that several of the THIP-induced sleep genes are

involved in metabolic processes, which are mostly downregulated (**Supplementary Table 2**).

This included the adipokinetic hormone receptor (AkhR), which has previously been

associated with starvation-induced sleep regulation [53]. We employed an RNAi strategy to

downregulate this metabolic gene's expression across the fly brain in AkhR-RNAi / R57C10-

Gal4 flies (see Methods). We found that downregulating AkhR significantly decreased sleep

duration during the day as well as night, compared to genetic control strains (**Figure 9A,B**).

Accordingly, sleep bout duration and number decreased, especially during the day (**Figure

9C**). However, in contrast to knocking out nAchRα1 and nAchRα2, which also significantly

decreased sleep duration day and night, short sleep was not significantly altered in AkhR

knockdown animals compared to genetic controls (**Figure 9D**). This suggests that short (1-

5min) sleep might be under separate regulatory control than >5min sleep.

We then returned to the nAchRα subunits and employed an RNAi strategy to knock them

down in the R23E10 sleep-promoting neurons specifically. While it is unlikely that these

cholinergic receptor subunits are restricted to just these neurons, we were curious if removing

any of them from this specific circuit had similar effect on sleep as in the knockout strains.

We found that nAchRα1 knockdown in the R23E10 neurons significantly decreased total

sleep during both day and night (**Figure 10A,B**), which was also true for the nAchRα1

knockouts (**Figure 8A**). Also consistent with the knockouts, the proportion of short sleep

bouts was increased in this this specific knockdown (**Figure 10C**). Among all of the other

alpha subunits, knockdown of nAchRα5 and nAchRα6 in the R23E10 neurons significantly

increased daytime and night-time sleep, respectively (**Figure 10A,B**), while short sleep was

correspondingly decreased only in nAchRα5 (**Figure 10C**). None of the other alpha subunits

had any effect in these sleep-promoting neurons. This suggests that their sleep-regulatory role

lies outside of this circuit, while nAchRα1, nAchRα5, and nAchRα6 might regulate sleep-

related functions within this circuit.

Discussion

One of the key advantages of studying sleep in *Drosophila* is that this versatile model

provides a variety of reliable approaches for inducing and controlling sleep. By being able to

induce sleep on demand, either genetically or pharmacologically, researchers have been able

to manipulate sleep as an experimental variable, and in this way be better able to assess

causality when probing potential sleep functions. However, this approach has often

sidestepped the question of whether different sleep induction methods are equivalent, or

whether distinct forms of sleep might be engaged by different genetic or pharmacological

treatments. In mammals, GABA agonists typically promote slow-wave sleep (SWS), which

has been associated with cellular homeostasis and repair process in the brain [4, 64]. In

contrast, drugs targeting acetylcholine receptors, such as carbachol, have been found to

promote brain states more reminiscent of REM sleep [65, 66]. Although these drugs all

induce sedative (or dissociative) states, they are clearly producing dissimilar forms of sleep in

mammals, with likely different functions or consequences for the brain. In *Drosophila*,

evidence suggests that the GABA agonist THIP promotes a form of deep or 'quiet' sleep,

which may be functionally analogous to mammalian SWS [9, 27, 67]. In contrast,

optogenetic activation of the dFB as well as other neurons may promote a form of active

sleep, which we have suggested could be achieving some similar sleep functions as REM [2].

THIP-induced sleep in flies has been associated with waste clearance from the brain [27], just

as SWS has been associated with clearance of waste metabolites via the mammalian brain's

glymphatic system [64]. Such functional homology suggests that the transcriptional changes

we uncovered for THIP-induced sleep in *Drosophila* might also be relevant for mammalian

SWS, with these largely centered on reduced metabolic processes and stress regulation [52].

In contrast, except for the upregulation of cholinergic signaling [68], there is little to compare

to test hypotheses potentially linking active sleep in flies with REM sleep in mammals,

except for the potential upregulation of cholinergic signaling. Even this cholinergic

connection seems odd, seeing that the predominant excitatory neurotransmitters are reversed

in the brains of insects and mammals: glutamate in mammals and acetylcholine in insects

[69]. Additionally, only nicotinic receptor subunits were identified in our analyses, whereas

muscarinic receptors have been more commonly associated with REM sleep in mammals [70,

71]. Nevertheless, it is clear from our results that optogenetic-induced active sleep

upregulates the expression of multiple nicotinic acetylcholine subunits, and that knocking

these out individually has profound (and opposing) effects on sleep architecture in flies, and

that a subset might function intrinsically within sleep promoting neurons. This supports other

studies showing a role for nAchRα's in sleep regulation [60, 61]. It will be especially

interesting in future brain imaging studies to see whether a knockout such as nAchRα3 is

eliminating one kind of sleep (e.g., active sleep) as predicted by our behavioral data, and

whether this is associated with any functional consequences. Similarly, it will be telling to

see whether the opposite sleep phenotypes observed in nAchRα1 for example result in a

distinct class of functional consequences. A previous study has shown that nAchRα1

knockout animals have significantly shorter survivorship compared to controls, with flies

dying almost 20 days earlier [45]. One reason could be because of impaired or insufficient

deep sleep functions (e.g., brain waste clearance [27]). The nAchRα knockouts provide an

opportunity to further examine mutant animals potentially lacking either kind of sleep,

although this will have to be confirmed by brain imaging or electrophysiology.

We found little similarity between two different approaches to inducing sleep in flies, at the

level of gene expression as well as acute effects on brain activity. It may however not be

surprising that these entirely different sleep induction methods produce dissimilar

physiological effects. After all, one method requires flies to ingest a drug which then must

make its way to the brain, while the other method directly activates a subset of neurons in the

central brain, as well as in the ventral nerve cord [72]. Yet both methods yield similarly

increased sleep duration profiles and consolidated sleep architecture (**Figure 2**). One

underlying assumption with focusing on sleep duration as the most relevant metric for

understanding sleep function in *Drosophila* is that sleep is a unitary phenomenon in the fly

model, meaning that primarily one set of functions and one form of brain activity are

occurring when flies sleep. There is now substantial evidence that this is unlikely to be true,

and that like other animals flies probably also experience distinct sleep stages that accomplish

different functions [9, 10, 27, 63, 73, 74]. This does not mean that these functions are

mutually exclusive; for example, both THIP provision and neuronal activation of central

brain circuits have been found to promote memory consolidation in *Drosophila* [23]. Indeed,

it seems reasonable to propose that different sleep stages could be synergistic, accomplishing

a variety of homeostatic functions that might be required for adaptive behaviors in an animal.

Our results suggest that THIP provision promotes a 'quiet sleep' stage in flies, which induces

a brain-wide downregulation of metabolism-related genes. This is consistent with studies in

flies showing that metabolic rate is decreased in longer sleep bouts, especially at night, and

that this is recapitulated by THIP-induced sleep [75]. Our findings also align with a previous

study showing that mitochondrial metabolic processes are upregulated in the absence of sleep

[76]. One argument for why metabolism-related genes are downregulated during THIP-

induced sleep might be that flies are starved (because they are sleeping more, and thus

perhaps not eating). However, flies induced to sleep by optogenetic activation are also

sleeping more (indeed, their average sleep bouts are longer), yet these did not reveal a similar

downregulation of metabolic processes. Another view might be that our sampling was done

after flies had achieved 10 hours of induced sleep, so sleep functions might have already been

achieved by that time. Thus, we might not be uncovering genes required for achieving 'quiet'

sleep functions as much as identifying exactly the opposite: genetic pathways that have been

satisfied by 10 hours of induced quiet sleep. Other studies using THIP to induce sleep have

examined longer timeframes (e.g., 2 days [23]), so it remains unclear whether changes in

gene regulation relate to sleep functions that have been achieved or that are still being

engaged. For our transcriptomic study, both sleep induction protocols were conducted during

the day (thus in presumably well-rested flies) to deliberately avoid any sleep pressure

confounds. Since the dFB has been proposed as a regulator of sleep homeostasis [20-22], one

could question why its prolonged activation did not incur a deep sleep debt – which might

have been detectable at a transcriptomic level by some overlap with our sleep deprivation

controls. That we did not see this suggests induced daytime active sleep might not accrue any

sleep debt. We have shown previously that daytime sleep is qualitatively different than night-

time sleep in flies, which is 'deeper' [63]. It is for example possible that most quiet sleep

happens at night in flies, so a daytime regime of induced active or quiet sleep might not incur

any sleep debt for either. Following this logic, our RNA analyses simply revealed the

transcriptomic changes resulting from either manipulation in well-rested flies, with the key

result being that there are zero shared genes after 10 daytime hours of either manipulation.

In contrast to THIP, our optogenetic manipulation induced a brain-wide upregulation of a

variety of neural mechanisms unrelated to metabolic processes, suggesting a different kind of

sleep was engaged. Although many studies have shown that the R23E10-Gal4 circuit is sleep

promoting (e.g.[19, 21, 74]), and we have shown that acute activation of these neurons

produces a form of active sleep [10], it seems unlikely that active sleep regulation is limited

to the dFB neurons alone [72]. Other circuits in the fly central brain are also sleep-promoting,

including in the ellipsoid body [77] and the ventral fan-shaped body (vFB) [78], although it

remains unknown if activation of these other circuits also promotes an active sleep stage, or

whether a similar transcriptome might be engaged by these alternate approaches to

optogenetic sleep induction in flies. This again highlights a variant of the same problem we

have uncovered in the current study comparing pharmacology with optogenetics: different

circuit-based approaches could all be increasing sleep duration but achieving entirely

different functions by engaging distinct transcriptomes and thus different sleep functions.

How many different kinds of functions are engaged by sleep remains unclear: is it roughly

two functional categories linked to quiet and active sleep, or is it a broader range of sub-

categories that are not so tightly linked to these obviously different brain activity states?

A compelling argument could nevertheless be made for two kinds of sleep in most animals,

with two distinct sets of functions [1, 67]. Most animals have been shown to require a form of

'quiet' sleep to ensure survival, suggesting that these might encompass an evolutionarily

conserved set of cellular processes that promote neural health and development [79], and that

operate best during periods of behavioral quiescence. Nematode worms thus experience a

form of quiet sleep when they pause to molt ('lethargus') into a different life stages during

their development [80], or when cellular repair processes are needed following environmental

stress [81]. In flies, quiet sleep seems to be similarly required for neuronal repair [26] or

waste clearance [27], and there is evidence that glia might play a key role in these cellular

homeostatic processes in flies [26] as well as other animals [82]. Thus, SWS in mammals and

birds might present a narrow neocortical view of a more ancient set of sleep functions

centered on quiescence and decreased metabolic rate. Indeed, neural quiescence is also a

feature of SWS, both at the level of pulsed inhibition (down-states), as well as in other parts

of the brain beyond the cortex [1]. Similar to findings in flies that are induced into a quiet

sleep stage with THIP [75], metabolic rate also decreases during SWS in mammals [83]. In

contrast, metabolic rate is similar to waking during REM sleep in mammals [84], suggesting

an alternate set of functions not linked to cellular homeostasis. What might these functions

be, and could some of these be conserved between active sleep in invertebrates and REM

sleep in mammals? A REM-like sleep stage has now been identified in a variety of

invertebrate species, including cephalopods [12] and jumping spiders [85]. In humans, REM

sleep has been implicated in emotion regulation [86], and cognitive disorders where emotions

are dysregulated, such as depression, are often associated with REM sleep dysfunction [87].

While it is not evident how to study emotions in insects (but see [88]), it could be argued that

arousal systems more generally are employed to detect prediction errors and thereby promote

learning [89]. Thus, we and others have suggested that REM sleep might be crucial for

optimizing prediction, and attention, and learning [2, 67, 90], and this may involve different

kinds of homeostatic mechanisms centered on brain circuits rather than cells. Whether active

sleep in flies plays a similar homeostatic role remains to be determined. Our finding that

optogenetically -induced sleep in *Drosophila* upregulates different nAchRα subunits is

consistent with new findings that these subunits regulate appetitive memories in flies [91] and

that cholinergic systems more generally underpin learning and memory in this animal [92].

Yet learning and memory in flies clearly also benefits from quiet sleep, as evidenced by

multiple studies using THIP as a sleep-promoting agent [23, 25, 93]. One view consistent

with our findings and previous studies is that both kinds of sleep are crucial for optimal

behavior: quiet sleep for cellular homeostasis and active sleep for circuit-level homeostasis.

Manipulating these separately, alongside the non-overlapping pathways engaged by either

kind of sleep, should help further disambiguate the functions potentially associated with these

distinct sleep stages.

Author contributions: Designed experiments, collected and analyzed data: NA, LALT-H,

HL, EN, AK. Designed experiments and analyzed data: QZ. Designed experiments: TP, PB,

PJS. Designed experiments, analyzed data, and wrote the paper: BvS.

Acknowledgements: This work was supported by National Health and Medical Research

Council grant GNT1164499 to BvS and NIH R01 grant NS076980 to PJS and BvS.

Declaration of Interests: The authors declare no conflicts of interest.

References

1.   Jaggard, J.B., G.X. Wang, and P. Mourrain, *Non-REM and REM/paradoxical sleep*

dynamics across phylogeny. Curr Opin Neurobiol, 2021. **71**: p. 44-51.

2.   Van De Poll, M.N. and B. van Swinderen, *Balancing Prediction and Surprise: A Role*

for Active Sleep at the Dawn of Consciousness? Front Syst Neurosci, 2021. **15**: p.

768762.

3.   Dijk, D.J., D.P. Brunner, and A.A. Borbely, *Time course of EEG power density*

during long sleep in humans. Am J Physiol, 1990. **258**(3 Pt 2): p. R650-61.

4.   Tononi, G. and C. Cirelli, *Sleep and the price of plasticity: from synaptic and cellular*

homeostasis to memory consolidation and integration. Neuron, 2014. **81**(1): p. 12-34.

5.   Siegel, J.M., *REM sleep: a biological and psychological paradox.* Sleep Med Rev,

2011. **15**(3): p. 139-42.

6.   Shein-Idelson, M., et al., *Slow waves, sharp waves, ripples, and REM in sleeping*

dragons. Science, 2016. **352**(6285): p. 590-5.

7.   Rattenborg, N.C., et al., *Local Aspects of Avian Non-REM and REM Sleep.* Front

Neurosci, 2019. **13**: p. 567.

8.   Libourel, P.A. and A. Herrel, *Sleep in amphibians and reptiles: a review and a*

preliminary analysis of evolutionary patterns. Biol Rev Camb Philos Soc, 2016.

91(3): p. 833-66.

9.   Yap, M.H.W., et al., *Oscillatory brain activity in spontaneous and induced sleep*

stages in flies. Nat Commun, 2017. **8**(1): p. 1815.

10.   Tainton-Heap, L.A.L., et al., *A Paradoxical Kind of Sleep in Drosophila*

melanogaster. Curr Biol, 2021. **31**(3): p. 578-590 e6.

11.   Leung, L.C., et al., *Neural signatures of sleep in zebrafish.* Nature, 2019. **571**(7764):

p. 198-204.

12.   Iglesias, T.L., et al., *Cyclic nature of the REM sleep-like state in the cuttlefish Sepia*

officinalis. J Exp Biol, 2019. **222**(Pt 1).

13.   Sauer, S., et al., *The dynamics of sleep-like behaviour in honey bees.* J Comp Physiol

A Neuroethol Sens Neural Behav Physiol, 2003. **189**(8): p. 599-607.

14.   Calcagno, B., et al., *Transient activation of dopaminergic neurons during*

development modulates visual responsiveness, locomotion and brain activity in a

dopamine ontogeny model of schizophrenia. Transl Psychiatry, 2013. **2**: p. e2026.

15.   Shaw, P.J., et al., *Correlates of sleep and waking in Drosophila melanogaster.*

Science, 2000. **287**(5459): p. 1834-7.

16.   Hendricks, J.C., et al., *Rest in Drosophila is a sleep-like state.* Neuron, 2000. **25**(1): p.

129-38.

17.   Shafer, O.T. and A.C. Keene, *The Regulation of Drosophila Sleep.* Curr Biol, 2021.

31(1): p. R38-R49.

18.   Wolff, T. and G.M. Rubin, *Neuroarchitecture of the Drosophila central complex: A*

catalog of nodulus and asymmetrical body neurons and a revision of the

protocerebral bridge catalog. J Comp Neurol, 2018. **526**(16): p. 2585-2611.

19.   Troup, M., et al., *Acute control of the sleep switch in Drosophila reveals a role for*

gap junctions in regulating behavioral responsiveness. Elife, 2018. **7**.

20.   Donlea, J.M., et al., *Inducing sleep by remote control facilitates memory*

consolidation in Drosophila. Science, 2011. **332**(6037): p. 1571-6.

21.   Donlea, J.M., D. Pimentel, and G. Miesenbock, *Neuronal machinery of sleep*

homeostasis in Drosophila. Neuron, 2014. **81**(4): p. 860-72.

22.   Donlea, J.M., et al., *Recurrent Circuitry for Balancing Sleep Need and Sleep.* Neuron,

2018. **97**(2): p. 378-389 e4.

23.   Dissel, S., et al., *Sleep restores behavioral plasticity to Drosophila mutants.* Curr

Biol, 2015. **25**(10): p. 1270-81.

24.   Harrison, N.L., *Mechanisms of sleep induction by GABA(A) receptor agonists.* J Clin

Psychiatry, 2007. **68 Suppl 5**: p. 6-12.

25.   Berry, J.A., et al., *Sleep Facilitates Memory by Blocking Dopamine Neuron-Mediated*

Forgetting. Cell, 2015. **161**(7): p. 1656-67.

26.   Stanhope, B.A., et al., *Sleep Regulates Glial Plasticity and Expression of the*

Engulfment Receptor Draper Following Neural Injury. Curr Biol, 2020. **30**(6): p.

1092-1101 e3.

27.   van Alphen, B., et al., *A deep sleep stage in Drosophila with a functional role in*

waste clearance. Sci Adv, 2021. **7**(4).

28.   Lundahl, J., et al., *EEG spectral power density profiles during NREM sleep for*

gaboxadol and zolpidem in patients with primary insomnia. J Psychopharmacol,

2012. **26**(8): p. 1081-7.

29.   Mitler, M.M. and W.C. Dement, *Cataplectic-like behavior in cats after micro-*

injections of carbachol in pontine reticular formation. Brain Res, 1974. **68**(2): p. 335-

43.

30.   Jenett, A., et al., *A GAL4-driver line resource for Drosophila neurobiology.* Cell Rep,

2012. **2**(4): p. 991-1001.

31.   Klapoetke, N.C., et al., *Independent optical excitation of distinct neural populations.*

Nat Methods, 2014. **11**(3): p. 338-46.

32.   Pfeiffer, B.D., J.W. Truman, and G.M. Rubin, *Using translational enhancers to*

increase transgene expression in Drosophila. Proc Natl Acad Sci U S A, 2012.

109(17): p. 6626-31.

33.   Legland, D., I. Arganda-Carreras, and P. Andrey, *MorphoLibJ: integrated library and*

915   *plugins for mathematical morphology with ImageJ.* Bioinformatics, 2016. **32**(22): p.

916   3532-3534.

917   34.   Faville, R., et al., *How deeply does your mutant sleep? Probing arousal to better*

918   *understand sleep defects in Drosophila.* Scientific Reports, 2015. **5**: p. 8454.

919   35.   Seugnet, L., et al., *Persistent short-term memory defects following sleep deprivation*

920   *in a drosophila model of Parkinson disease.* Sleep, 2009. **32**(8): p. 984-92.

921   36.   Chen, C., et al., *Drosophila Ionotropic Receptor 25a mediates circadian clock*

922   *resetting by temperature.* Nature, 2015. **527**(7579): p. 516-20.

923   37.   Kim, D., B. Langmead, and S.L. Salzberg, *HISAT: a fast spliced aligner with low*

924   *memory requirements.* Nat Methods, 2015. **12**(4): p. 357-60.

925   38.   Li, H., et al., *The Sequence Alignment/Map format and SAMtools.* Bioinformatics,

926   2009. **25**(16): p. 2078-9.

927   39.   Robinson, M.D., D.J. McCarthy, and G.K. Smyth, *edgeR: a Bioconductor package*

928   *for differential expression analysis of digital gene expression data.* Bioinformatics,

929   2010. **26**(1): p. 139-40.

930   40.   Robinson, M.D. and A. Oshlack, *A scaling normalization method for differential*

931   *expression analysis of RNA-seq data.* Genome Biol, 2010. **11**(3): p. R25.

932   41.   McCarthy, D.J., Y. Chen, and G.K. Smyth, *Differential expression analysis of*

933   *multifactor RNA-Seq experiments with respect to biological variation.* Nucleic Acids

934   Res, 2012. **40**(10): p. 4288-97.

935   42.   Lun, A.T., Y. Chen, and G.K. Smyth, *It's DE-licious: A Recipe for Differential*

936   *Expression Analyses of RNA-seq Experiments Using Quasi-Likelihood Methods in*

937   *edgeR.* Methods Mol Biol, 2016. **1418**: p. 391-416.

43.   Huang da, W., B.T. Sherman, and R.A. Lempicki, *Bioinformatics enrichment tools:*

paths toward the comprehensive functional analysis of large gene lists. Nucleic Acids

Res, 2009. **37**(1): p. 1-13.

44.   Huang da, W., B.T. Sherman, and R.A. Lempicki, *Systematic and integrative analysis*

of large gene lists using DAVID bioinformatics resources. Nat Protoc, 2009. **4**(1): p.

44-57.

45.   Somers, J., et al., *Pleiotropic Effects of Loss of the Dalpha1 Subunit in Drosophila*

melanogaster: Implications for Insecticide Resistance. Genetics, 2017. **205**(1): p. 263-

271.

46.   Perry, T., et al., *Role of nicotinic acetylcholine receptor subunits in the mode of action*

of neonicotinoid, sulfoximine and spinosyn insecticides in Drosophila melanogaster.

Insect Biochem Mol Biol, 2021. **131**: p. 103547.

47.   Zimmerman, J.E., et al., *A video method to study Drosophila sleep.* Sleep, 2008.

31(11): p. 1587-98.

48.   Huber, R., et al., *Sleep homeostasis in Drosophila melanogaster.* Sleep, 2004. **27**(4):

p. 628-39.

49.   Andretic, R. and P.J. Shaw, *Essentials of sleep recordings in Drosophila: moving*

beyond sleep time. Methods Enzymol, 2005. **393**: p. 759-72.

50.   Kirszenblat, L., R. Yaun, and B. van Swinderen, *Visual experience drives sleep need*

in Drosophila. Sleep, 2019. **42**(7).

51.   Troup, M.J., L.A.L. Tainton-Heap, and B. van Swinderen, *Neural ensemble*

fragmentation in the anesthetized Drosophila brain. J Neurosci, 2023.

52.   Muheim, C.M., et al., *Dynamic- and Frequency-Specific Regulation of Sleep*

Oscillations by Cortical Potassium Channels. Curr Biol, 2019. **29**(18): p. 2983-2992

e3.

53. He, Q., et al., *AKH-FOXO pathway regulates starvation-induced sleep loss through*

remodeling of the small ventral lateral neuron dorsal projections. PLoS Genet, 2020.

16(10): p. e1009181.

54. Bharucha, K.N., P. Tarr, and S.L. Zipursky, *A glucagon-like endocrine pathway in*

Drosophila modulates both lipid and carbohydrate homeostasis. J Exp Biol, 2008.

211(Pt 19): p. 3103-10.

55. Ferguson, L., et al., *Transient Dysregulation of Dopamine Signaling in a Developing*

Drosophila Arousal Circuit Permanently Impairs Behavioral Responsiveness in

Adults. Front Psychiatry, 2017. **8**: p. 22.

56. Shao, L., et al., *Schizophrenia susceptibility gene dysbindin regulates glutamatergic*

and dopaminergic functions via distinctive mechanisms in Drosophila. Proc Natl

Acad Sci U S A, 2011. **108**(46): p. 18831-6.

57. Van Swinderen, B. and R. Andretic, *Dopamine in Drosophila: setting arousal*

thresholds in a miniature brain. Proc Biol Sci, 2011. **278**(1707): p. 906-13.

58. Watson, C.J., H.A. Baghdoyan, and R. Lydic, *Neuropharmacology of Sleep and*

Wakefulness. Sleep Med Clin, 2010. **5**(4): p. 513-528.

59. Rosenthal, J.S. and Q. Yuan, *Constructing and Tuning Excitatory Cholinergic*

Synapses: The Multifaceted Functions of Nicotinic Acetylcholine Receptors in

Drosophila Neural Development and Physiology. Front Cell Neurosci, 2021. **15**: p.

720560.

60. Shi, M., et al., *Identification of Redeye, a new sleep-regulating protein whose*

expression is modulated by sleep amount. Elife, 2014. **3**: p. e01473.

61. Dai, X., et al., *Molecular resolution of a behavioral paradox: sleep and arousal are*

regulated by distinct acetylcholine receptors in different neuronal types in

Drosophila. Sleep, 2021. **44**(7).

62.   Chen, W., et al., *Dual nicotinic acetylcholine receptor subunit gene knockouts reveal*

989   *limits to functional redundancy.* Pestic Biochem Physiol, 2022. **184**: p. 105118.

990   63.   van Alphen, B., et al., *A dynamic deep sleep stage in Drosophila.* J Neurosci, 2013.

991   **33**(16): p. 6917-27.

992   64.   Xie, L., et al., *Sleep drives metabolite clearance from the adult brain.* Science, 2013.

993   **342**(6156): p. 373-7.

994   65.   Kubin, L., *Carbachol models of REM sleep: recent developments and new directions.*

995   Arch Ital Biol, 2001. **139**(1-2): p. 147-68.

996   66.   Torterolo, P., et al., *Neocortical 40 Hz oscillations during carbachol-induced rapid*

997   *eye movement sleep and cataplexy.* Eur J Neurosci, 2016. **43**(4): p. 580-9.

998   67.   Kirszenblat, L. and B. van Swinderen, *The Yin and Yang of Sleep and Attention.*

999   Trends Neurosci, 2015. **38**(12): p. 776-86.

1000   68.   Vazquez, J. and H.A. Baghdoyan, *Basal forebrain acetylcholine release during REM*

1001   *sleep is significantly greater than during waking.* Am J Physiol Regul Integr Comp

1002   Physiol, 2001. **280**(2): p. R598-601.

1003   69.   Smarandache-Wellmann, C.R., *Arthropod neurons and nervous system.* Curr Biol,

1004   2016. **26**(20): p. R960-R965.

1005   70.   Bourgin, P., et al., *Induction of rapid eye movement sleep by carbachol infusion into*

1006   *the pontine reticular formation in the rat.* Neuroreport, 1995. **6**(3): p. 532-6.

1007   71.   Kashiwagi, M. and Y. Hayashi, *Life Without Dreams: Muscarinic Receptors Are*

1008   *Required to Regulate REM Sleep in Mice.* Cell Rep, 2018. **24**(9): p. 2211-2212.

1009   72.   Jones, J.D., et al., *Regulation of sleep by cholinergic neurons located outside the*

1010   *central brain in Drosophila.* PLoS Biol, 2023. **21**(3): p. e3002012.

73.   Wiggin, T.D., et al., *Covert sleep-related biological processes are revealed by probabilistic analysis in Drosophila.* Proc Natl Acad Sci U S A, 2020. **117**(18): p. 10024-10034.

74.   Gong, N.N., et al., *Intrinsic maturation of sleep output neurons regulates sleep ontogeny in Drosophila.* Curr Biol, 2022. **32**(18): p. 4025-4039 e3.

75.   Stahl, B.A., et al., *Sleep-Dependent Modulation of Metabolic Rate in Drosophila.* Sleep, 2017. **40**(8).

76.   Kempf, A., et al., *A potassium channel beta-subunit couples mitochondrial electron transport to sleep.* Nature, 2019. **568**(7751): p. 230-234.

77.   Liu, S., et al., *Sleep Drive Is Encoded by Neural Plastic Changes in a Dedicated Circuit.* Cell, 2016. **165**(6): p. 1347-1360.

78.   Lei, Z., K. Henderson, and K. Keleman, *A neural circuit linking learning and sleep in Drosophila long-term memory.* Nat Commun, 2022. **13**(1): p. 609.

79.   Zimmerman, J.E., et al., *Conservation of sleep: insights from non-mammalian model systems.* Trends Neurosci, 2008. **31**(7): p. 371-6.

80.   Raizen, D.M., et al., *Lethargus is a Caenorhabditis elegans sleep-like state.* Nature, 2008. **451**(7178): p. 569-72.

81.   Hill, A.J., et al., *Cellular stress induces a protective sleep-like state in C. elegans.* Curr Biol, 2014. **24**(20): p. 2399-405.

82.   Artiushin, G. and A. Sehgal, *The Glial Perspective on Sleep and Circadian Rhythms.* Annu Rev Neurosci, 2020. **43**: p. 119-140.

83.   Caron, A.M. and R. Stephenson, *Energy expenditure is affected by rate of accumulation of sleep deficit in rats.* Sleep, 2010. **33**(9): p. 1226-35.

84.   Maquet, P., *Sleep function(s) and cerebral metabolism.* Behav Brain Res, 1995. **69**(1-2): p. 75-83.

85. Rossler, D.C., et al., *Regularly occurring bouts of retinal movements suggest an REM sleep-like state in jumping spiders.* Proc Natl Acad Sci U S A, 2022. **119**(33): p. e2204754119.

86. Hutchison, I.C. and S. Rathore, *The role of REM sleep theta activity in emotional memory.* Front Psychol, 2015. **6**: p. 1439.

87. Harrington, M.O., et al., *The influence of REM sleep and SWS on emotional memory consolidation in participants reporting depressive symptoms.* Cortex, 2018. **99**: p. 281-295.

88. Anderson, D.J. and R. Adolphs, *A framework for studying emotions across species.* Cell, 2014. **157**(1): p. 187-200.

89. Seth, A.K. and K.J. Friston, *Active interoceptive inference and the emotional brain.* Philos Trans R Soc Lond B Biol Sci, 2016. **371**(1708).

90. Hobson, J.A., *REM sleep and dreaming: towards a theory of protoconsciousness.* Nat Rev Neurosci, 2009. **10**(11): p. 803-13.

91. Pribbenow, C., et al., *Postsynaptic plasticity of cholinergic synapses underlies the induction and expression of appetitive and familiarity memories in Drosophila.* Elife, 2022. **11**.

92. Barnstedt, O., et al., *Memory-Relevant Mushroom Body Output Synapses Are Cholinergic.* Neuron, 2016. **89**(6): p. 1237-1247.

93. Melnattur, K., et al., *A conserved role for sleep in supporting Spatial Learning in Drosophila.* Sleep, 2020.

Figure Legends

Figure 1. Study rationale and design. The same genetic background strain (R23E10;UAS-Chrimson) was used for optogenetic or pharmacologically-induced sleep. Flies were fed either all-trans retinal (ATR) or 4,5,6,7-tetrahyrdoisoxazolopyridin-3-ol (THIP) to promote either kind of sleep, which was assessed in three different ways: behavioral analysis, whole brain imaging, and gene expression changes. The comparisons made for each level of analysis are labelled A-E.

Figure 2. Optogenetic- and THIP-induced sleep have similar effects on sleep duration and consolidation. A) Experimental regime for observing the effects of optogenetic activation and THIP provision **(B)**. **C)** Sleep profile across 24 hours in the baseline condition (grey) and optogenetic activation condition (green). **D)** 3-day average of the 24-hour sleep profile of control (grey) and THIP fed (blue) flies. **E)** Daytime sleep consolidation scatterplot for optogenetic baseline and THIP control flies. **F)** Daytime sleep consolidation scatterplot for optogenetic- and THIP-induced sleep. **G)** Night-time sleep consolidation scatterplot for optogenetic baseline and THIP control flies. **H)** Night-time sleep consolidation scatterplot for optogenetic- and THIP-induced sleep. n = 87 for optogenetic activation across three replicates; n = 88 for –THIP, n = 85 for +THIP, across three replicates. Maximum bout duration possible is 720 minutes, or 12 hours. See Supplementary Figure 1 for summary histograms and Supplementary Table 1 for statistics.

Figure 3. Sleep architecture in optogenetic and THIP induced sleep.

A-B. Average number of sleep bouts in control (grey) and optogenetic activation (green) conditions in the day and night for both +ATR **(A)** and –ATR **(B)** fed flies. optogenetic-induced sleep results in an increase in the number of sleep bouts both during the day and the

night, whereas red light alone has no effect. **C-D.** optogenetic activation (green) increases

the average sleep duration during the day, but not the night when compared to controls (grey)

in +ATR flies **(C)**. **D.** –ATR flies show no difference in mean sleep bout duration during the

day, but show a decrease in average bout duration during the night. THIP (blue) increases

both the average number of sleep bouts **(E)** and the average duration of sleep bouts **(F)**

during the day, but not the night, when compared to controls (grey). Analysis for a and b =

Kruskal-Wallis test with Dunn's multiple comparison correction. * = $p<0.05$, *** = $p <$

0.001, **** = $p<0.0001$. For e and f, analysis = Ordinary one-way ANOVA with Tukey

correction for multiple comparisons. *** = $p<0.001$, **** = $p <0.0001$.

Figure 4. Brain imaging during optogenetic and THIP-induced sleep

A) Flies were mounted onto a custom-built holder that allowed a coronal visualization of the

brain through the posterior side of the head. Perfusion of extracellular fluid (ECF) occurred

throughout all experiments. A 617nm LED was delivered to the brain through the imaging

objective during optogenetic experiments. During THIP experiments, 4% THIP in ECF was

perfused onto the brain through a custom perfusion system. Behavior was recorded as the

movement of flies on an air suspended ball. **B)** Left: Imaging was carried out across 18 z-slices,

with a z-step of 6μm. Each z-plane spanned 667μm x 667μm, which was captured across 256

x 256 pixels. Right: A collapsed mask from one fly of neurons found to be active (green) in C

alongside all identified regions of interest (ROIs, gray). **C)** Neural activity in an example fly

brain, represented across cells (top) and as the population mean (middle) did not change

following optogenetic-induced sleep (bottom). **D)** Neural activity in an example fly brain,

represented across cells (top) and as the population mean (middle) showed an initial high level

of activity in the baseline condition, which decreased when the fly fell asleep (bottom)

1107   following THIP exposure. The Y axis scale is standard deviation of the experiment mean, so

1108   the baseline is not absolute but rather reflects any difference with the overall experiment mean.

1109

Figure 5. Brain activity and connectivity decreases during THIP-induced sleep. A)

Experimental protocol for behavioral responsiveness and brain imaging experiments. 5 mins

of baseline condition were recorded, during which the exposed brain was perfused with

extracellular fluid (ECF), followed by 5 mins of THIP perfusion. Following sleep induction,

an additional 10 minutes of calcium activity was recorded, which was separated into 'Early'

and 'Mid' sleep for analysis. Air puff stimuli were delivered to test for behavioral

responsiveness. **B)** Mean behavioral response rate (% ± sem) to air puff stimuli over the

course of an experiment (n = 6). Air puff delivery times are indicated by the solid dots. **C)**

Percent neurons active (± sem) in non ATR-fed UAS:Chrimson / X ; Nsyb:LexA/+ ;

LexOp:nlsGCaMP6f / R23E10:Gal4 flies during wake, THIP-induced sleep, and recovery (n

= 9; 3 flies were recorded post-waking). **D)** Correlation analysis (mean degree ± sem) of

active neurons in (C). **E)** Collapsed mask of neurons active during wake, and both early and

mid THIP sleep. **F)** Overlap in neural identities between wake and THIP-induced sleep in

two example flies. Number indicates active neurons within each condition; same color code

as in (E). **G)** Quantification of neural overlap data. Red dots indicate the flies shown in (F). n

= 9 flies. All tests are one-way ANOVA with Dunnett's multiple comparison test. ns = not

significant, * = $p<0.05$, ** = $p<0.01$, **** = $p < 0.001$.

1127

1128   **Figure 6. Metabolic processes are downregulated during THIP-induced sleep.**

1129   **A)** Schematic representation of the experimental set-up and samples processed using RNA-

1130   Sequencing. **B)** Venn diagram showing the gene expression overlap between flies that had

1131   been treated with THIP versus their control (shaded blue) and flies that had been treated with

THIP in a sleep deprived background versus their control (shaded blue bars). The number of

significant differentially-expressed genes in each category is indicated. **C)** Volcano plot

representing the distribution of differentially expressed genes in the presence or absence of

THIP. Genes that are significantly up/down regulated meeting a Log2Fold change of 0.58

and FDRq value of 0.05 are shown in red. Genes meeting the threshold for FDRq value only

are shown in blue. Fold change only is shown in green. Those genes not meeting any

predetermined criteria are shown in grey. **D)** Volcano plot representing the distribution of

differentially expressed genes in the presence or absence of THIP in a sleep deprived

background. Criteria as above (C). **E)** Schematic representation of Gene Ontology (GO)

enrichment of biological process results. Colour coded to indicate parent and child terms for

comparisons between groups highlighted above (C) - Left and (D) - Right). **F)** Bar chart

representation of a subset of interesting significant GO pathway terms originating from the

organic substance and primary metabolic processes for the dataset shown in (C). **G)**

Comparison between significant gene hits obtained via RNA-Sequencing (Blue) and qRT-

PCR (Grey) in response to THIP, represented by Log2Fold change values. See Supplemental

Tables 2&3 and Supplementary Figures 1&2.

Figure 7. A variety of biological processes including axon guidance are upregulated during optogenetic-induced sleep.

A) Schematic representation of the experimental set-up and samples processed using RNA-

Sequencing. **B)** Venn diagram showing the gene expression overlap between flies that

experienced 10 hours of optogenetic-induced sleep (ZT10) compared to -ATR controls

(ZT10) and those flies where optogenetic activation was restricted to 1 hour (ZT1) and

compared to -ATR controls (ZT1). **C)** Volcano plot representing the distribution of

differentially expressed genes resulting from optogenetic optogenetic activation for 10 hours

versus control flies which were allowed to sleep spontaneously for 10 hours. Genes that are

significantly up/down regulated meeting a Log2Fold change of 0.58 and FDRq value of 0.05

are shown in red. Genes meeting the threshold for FDRq value only are shown in blue. Fold

change only in green. Those genes not meeting any predetermined criteria are shown in grey.

D) Volcano plot representing the distribution of differentially expressed genes resulting from

optogenetic activation for 1 hour versus control flies which were allowed to sleep

spontaneously for 1 hour. Criteria as above (C). **E)** Schematic representation of Gene

Ontology (GO) enrichment of biological process results. Color coded to indicate parent and

child terms comparing flies that had been activated optogenetically for 10 hours versus flies

which had been allowed to spontaneously sleep for the same duration. **F)** Bar chart

representation of a subset of interesting significant GO pathway terms originating from the

regulation of cellular processes and signalling biological processes. **G)** Comparison between

significant gene hits obtained via RNA-Sequencing (Green) and qRT-PCR (Grey) in response

to optogenetic sleep, represented by Log2Fold change values. See Supplemental Tables 4-7

and Supplementary Figures 3&4.

1172

Figure 8. nAchRα subunit knockouts bidirectionally regulate >5min sleep as well as

short sleep. A. Average total day and night sleep duration (minutes±95% confidence

intervals) in nAchRα knockout mutants, expressed as difference to their respective

background controls (see Methods). α1, N=91; control ($X^{59}w^{1118}$) = 93; α2, N=70; control

(w^{1118}ActinCas9) = 65; α3, N=43; (ActinCas9) =9; α4, N=87; (w^{1118}ActinCas9) =98; α6,

N=91; (w^{1118}ActinCas9) =91; α7, N=94; (ActinCas9) =95. *$P<0.05$, ***$P<0.001$,

****$P<0.0001$ by t-test adjusted for multiple comparisons. **B.** Left two panels: sleep

architecture for the same six knockout strains as in A (green), shown against their respective

controls (black). Each datapoint is a fly. Right panels: cumulative short sleep (1-5min)

expressed as a percentage of total sleep duration. Data are the from the same experiment as in A&B. Each datapoint is a fly. **$P<0.01$, ***$P<0.001$, ****$P<0.0001$ Man-Whitney U Test. All data were collected over three days and three nights and averaged.

Figure 9. AkhR knockdown decreases >5min sleep but not short sleep. A,B. Total sleep (>5min) in UAS-AkhR:RNAi / R57C10-Gal4 flies (blue, N=126) compared to genetic controls (light grey: UAS-AkhR:RNAi / + , N= 124; dark grey: R57C10-Gal4 / + , N= 120). **C.** Sleep architecture (average bout duration versus bout number per fly) in data from A,B. **D.** Cumulative short sleep (1-5min, expressed as a % of total sleep) in UAS-AkhR:RNAi / R57C10-Gal4 flies (blue) compared to genetic controls (light grey: UAS-AkhR:RNAi / + ; dark grey: R57C10-Gal4 / +). Wild-type background (+) is Canton-S(w[1118]). Each datapoint is a fly. ***$P<0.001$, ****$P<0.0001$ Man-Whitney U Test. ns, not significant. All data were collected over two days and two nights and averaged.

Figure 10. nAchRα subunit knockdowns in sleep-promoting neurons. A. Average total day and night sleep duration (minutes±95% confidence intervals) in UAS-nAchRα RNAi / R23E10-Gal4 flies, expressed as difference to RNAi / + controls (all were also compared Gal4 / + controls, not shown). α1 N=60, Gal4 N=45, RNAi N=37; α2 N=21, Gal4 N=24, RNAi N=50; α3 N=33, Gal4 N=50, RNAi N=59; α4 N=92, Gal4 N=90, RNAi N=80; α5 N=94, Gal4 N=47, RNAi N=74; α6 N=44, Gal4 N=47, RNAi N=43; α7 N=32, Gal4 N=57, RNAi N=44. **$P<0.01$, ***$P<0.001$ by t-test adjusted for multiple comparisons. **B.** Total >5min sleep duration data for significant knoockdowns in (A). **C.** Cumulative short sleep (1-5min) expressed as a percentage of total sleep duration. Data are the from the same experiment as in A&B. Each datapoint is a fly. ***$P<0.001$, ****$P<0.0001$ Man-Whitney U Test. All data were collected over two days and two nights and averaged.

Supplementary Figure Legends

Figure S1. Gene Ontology (GO) enrichment analysis for THIP-induced sleep (related to Figure 6). Significantly downregulated and upregulated GO categories for THIP-sleep (Table S2), listed from most enriched at the top. Broad GO categories are identified below.

Figure S2. Gene Ontology (GO) enrichment analysis for THIP-provisioned flies that were sleep deprived (related to Figure 6). Significantly downregulated and upregulated GO categories for sleep deprived flies (Table S3), listed from most enriched at the top. Broad GO categories are identified below.

Figure S3. Gene Ontology (GO) enrichment analysis for optogenetic-induced sleep (related to Figure 7). Significantly downregulated and upregulated GO categories for optogenetic-sleep (Table S4), listed from most enriched at the top. Broad GO categories are identified below.

Figure S4. Circadian-related genes uncovered in optogenetic-sleep dataset (related to Figure 7). A. Zeitgeber (ZT) 10 timepoint was compared with ZT in to uncover potential circadian-regulated genes, in two separate datasets (-ATR and +ATR). 98 genes were shared between these datasets. **B.** Of the 98 shared genes, circadian-related processes were highly enriched. **C.** Expression levels of 7 circadian genes drawn from the two different datasets in A.

Figure S5. Summary of different Gene Ontogeny pathways engaged by optogenetic-induced sleep and THIP-induced sleep. A. Either sleep induction method produces

different levels of activity in the fly brain. We term optogenetic-induced sleep 'active sleep' because brain activity levels are not different than during wake. We term THIP-induced sleep 'quiet sleep' because neural activity is significant decreased already in the first 5 minutes. Both of these induced forms of sleep resemble sleep stages seen during spontaneous sleep in flies. **B.** Number of GO pathways engaged by either induced active or quiet sleep, separated by upregulated versus downregulated biological processes.

Figure S6. Waking activity levels of nAchRα knockout mutants. Top: mutants are compared to their genetic background strain. Da1-7 = nAchRα1-7. Bottom: statistical tests for waking activity levels in each knockout compared to its genetic control, during bith day and night.

Supplementary Tables

Table S1, related to Figure 2. A comparison of sleep duration profiles (min/hr) during optogenetic and THIP induced sleep. Tested with 2way ANOVA with Tukey's multiple comparison test.

Table S2, related to Figure 6. List of significant THIP-sleep genes. Raw data for all genes.

Table S3, related to Figure 6. List of significant sleep-deprivation genes in THIP-fed flies. Raw data for all genes.

Table S4, related to Figure 7. List of significant optogenetic-sleep genes after 10 hours activation. Raw data for all genes.

Table S5, related to Figure 7. List of significant optogenetic-sleep genes after 1 hour of activation. Raw data for all genes.

Table S6, related to Figure 7. List of significant ZT10 vs ZT1 genes in ATR+ dataset. Raw data for all genes.

Table S7, related to Figure 7. List of significant ZT10 vs ZT1 genes in ATR- dataset. Raw data for all genes.

Table S8, related to Figures 6 & 7. Primer list for RT qPCR validation experiments.

Table S9, related to Figures 6&7. Raw data and statistics for RT qPCR experiments.

Background

R23e10Gal4 > UAS-Chrimson

Sleep Manipulation

Optogenetic   **Pharmacological**

Treatment

-ATR	+ATR	+THIP	-THIP
Control	Sleep	Sleep	Control

Sleep Readout   **Comparisons**

1. Behaviour

 A

2. Brain Activity

 B

3. Transcriptome

C   **D**

E

A **Optognetically induced sleep**

B **Drug (THIP) induced Sleep**

Sleep duration

C

D

Sleep architecture **Day**   **Night**

E

G

F

H

Figure 6

A

B

C ZT10 +ATR vs -ATR

D ZT1 +ATR vs -ATR

E

Upregulated

Downregulated

Regulation of macromolecular metabolic process
Regulation of cellular process
Regulation of Signalling
Regulation of development process
Regulation of localisation
Establishment of localisation
Regulation of protein localisation

Biosynthetic process
Nitrogen compound metabolic process
Cellular metabolic process
Regulation of biological process
Regulation of biological quality
Reproductive behaviour

F

GO Pathway Analysis

synaptic vesicle recycling via endosome

synaptic vesicle recycling

positive regulation of posttranscriptional gene silencing

neurotransmitter metabolic process

histamine metabolic process

semaphorin−plexin signaling pathway: reg of photoreceptor cell axon guidance

semaphorin−plexin signaling pathway: neuron projection guidance

semaphorin−plexin signaling pathway: axon guidance

semaphorin−plexin signaling pathway

ephrin receptor signaling pathway

Enrichment Value

G

qRT-PCR validation compared to RNA-Seq

Figure 8

Figure 9

Figure 10

	GO Term	Pvalue	Enrichment value
Downregulated	GO:0046460 neutral lipid biosynthetic process	0.0006	52.92
	GO:0046463 acylglycerol biosynthetic process	0.0006	52.92
	GO:0002181 cytoplasmic translation	0.0000	33.41
	GO:0009059 macromolecule biosynthetic process	0.0000	6.44
	GO:0009058 biosynthetic process	0.0000	4.53
	GO:0009064 glutamine family amino acid metabolic process	0.0008	16.54
	GO:0006412 translation	0.0000	11.7
	GO:0043043 peptide biosynthetic process	0.0000	11.54
	GO:0043604 amide biosynthetic process	0.0000	10.48
	GO:0006518 peptide metabolic process	0.0000	8.89
	GO:1901566 organonitrogen compound biosynthetic process	0.0000	6.38
	GO:0044271 cellular nitrogen compound biosynthetic process	0.0000	5.07
	GO:0034641 cellular nitrogen compound metabolic process	0.0000	2.31
	GO:1901564 organonitrogen compound metabolic process	0.0000	2.06
	GO:0006807 nitrogen compound metabolic process	0.0002	1.6
	GO:1901576 organic substance biosynthetic process	0.0000	4.53
	GO:0019752 carboxylic acid metabolic process	0.0008	3.36
	GO:0043436 oxoacid metabolic process	0.0010	3.26
	GO:0006082 organic acid metabolic process	0.0010	3.25
	GO:0019538 protein metabolic process	0.0000	2.27
	GO:0071704 organic substance metabolic process	0.0000	1.9
	GO:0043170 macromolecule metabolic process	0.0001	1.72
	GO:0006591 ornithine metabolic process	0.0006	52.92
	GO:0006525 arginine metabolic process	0.0008	44.1
	GO:1901605 alpha-amino acid metabolic process	0.0003	6.73
	GO:0008152 metabolic process	0.0000	1.83
	GO:0044238 primary metabolic process	0.0000	1.89
	GO:0019432 triglyceride biosynthetic process	0.0003	66.15
	GO:0006414 translational elongation	0.0000	34.51
	GO:0034645 cellular macromolecule biosynthetic process	0.0000	7.55
	GO:0043603 cellular amide metabolic process	0.0000	7.7
	GO:0044249 cellular biosynthetic process	0.0000	4.69
	GO:0044267 cellular protein metabolic process	0.0000	2.58
	GO:0044260 cellular macromolecule metabolic process	0.0000	2.02
	GO:0044237 cellular metabolic process	0.0000	1.67
	GO:0007548 sex differentiation	0.0003	22.05
Upregulated	GO:0006030 chitin metabolic process	0.0003	21.44
	GO:1901071 glucosamine-containing compound metabolic process	0.0004	19.78
	GO:0006040 amino sugar metabolic process	0.0004	19.33
	GO:0006022 aminoglycan metabolic process	0.0006	17.48
	GO:0018990 ecdysis, chitin-based cuticle	0.0002	85.05
	GO:0022404 molting cycle process	0.0004	65.42
	GO:0040003 chitin-based cuticle development	0.0000	32.89
	GO:0042335 cuticle development	0.0000	31.33
	GO:0048856 anatomical structure development	0.0004	4.39

Metabolic process

Biosynthetic metabolic process
Nitrogen compound metabolic process
organic substance metabolic process
primary metabolic process
Cellular metabolic process

Developmental Process

Developmental process involved in reproduction
Anatomical structure development

Mutlicellular organismal process

Molting cycle process

	GO Term	Pvalue	Enrichment value
Downregulated	GO:0104004 cellular response to environmental stimulus	0.0005	19.04
	GO:0050896 response to stimulus	0.0000	3.79
	GO:0034644 cellular response to UV	0.0000	85.05
	GO:0009411 response to UV	0.0001	41.15
	GO:0071482 cellular response to light stimulus	0.0001	32.71
	GO:0071478 cellular response to radiation	0.0003	22.78
	GO:0071214 cellular response to abiotic stimulus	0.0005	19.04
	GO:0009266 response to temperature stimulus	0.0000	16.88
	GO: 0009617 response to bacterium	0.0000	21.07
	GO: 0051707 response to other organism	0.0000	15.39
	GO:0043207 response to external biotic stimulus	0.0000	15.14
	GO: 0009607 response to biotic stimulus	0.0000	15.09
	GO:0050830 defense response to Gram-positive bacterium	0.0000	36.98
	GO:0009605 response to external stimulus	0.0000	9.31
	GO:0034605 cellular response to heat	0.0000	60.75
	GO:0009408 response to heat	0.0000	26.58
	GO:0006979 response to oxidative stress	0.0002	13.83
	GO:0042742 defense response to bacterium	0.0000	12.89
	GO:0098542 defense response to other organism	0.0000	11.63
	GO:0006952 defense response	0.0000	10.12
	GO:0006950 response to stress	0.0000	5.79
	GO:0042381 hemolymph coagulation	0.0001	141.75
	GO:0006955 immune response	0.0007	9.83
	GO:0002376 immune system process	0.0002	9.01
	GO:0007599 hemostasis	0.0001	141.75
	GO:0050878 regulation of body fluid levels	0.0005	56.7
	GO:0051704 multi-organism process	0.0000	12.34
	GO:0050817 coagulation	0.0001	141.75
Upregulated	GO:0008152 metabolic process	0.0000	1.86
	GO:0006032 chitin catabolic process	0.0001	36.6
	GO:1901072 glucosamine-containing compound catabolic process	0.0001	32.53
	GO:0006022 aminoglycan metabolic process	0.0000	13.37
	GO:0006807 nitrogen compound metabolic process	0.0000	1.94
	GO:0046348 amino sugar catabolic process	0.0001	32.53
	GO:0006026 aminoglycan catabolic process	0.0003	21.69
	GO:0006030 chitin metabolic process	0.0000	16.4
	GO:1901071 glucosamine-containing compound metabolic process	0.0000	15.13
	GO:0006040 amino sugar metabolic process	0.0000	14.79
	GO:0006508 proteolysis	0.0000	5.13
	GO:1901135 carbohydrate derivative metabolic process	0.0002	3.94
	GO:1901564 organonitrogen compound metabolic process	0.0000	2.31
	GO:0043170 macromolecule metabolic process	0.0000	2.1
	GO:0071704 organic substance metabolic process	0.0000	1.92
	GO:0019538 protein metabolic process	0.0008	2.01
	GO:0044238 primary metabolic process	0.0004	1.67
	GO:0017144 drug metabolic process	0.0000	6.24

Response to stimulus
Cellular response to stimulus
Response to abiotic stimulus
Response to biotic stimulus
Response to external stimulus
Response to stress
Mutli-organism process
Mutliorganism cellular process
MutliCellular organismal process
Coagulation

Immune system process
Immune response
Biological regulation
Regulation of biological quality
Metabolic process
Metabolic process
Nitrogen compound metabolic process
Organic substance
Primary Metabolic process
Other

	GO Term	Pvalue	Enrichment value
Downregulated	GO:0060148 positive regulation of posttranscriptional gene silencing	0.0005	53.51
	GO:0002181 cytoplasmic translation	0	9.46
	GO:0052803 imidazole-containing compound metabolic process	0.0002	89.19
	GO:0001692 histamine metabolic process	0.0002	89.19
	GO:0042133 neurotransmitter metabolic process	0.0007	16.72
	GO:1900368 regulation of RNA interference	0.0001	133.79
	GO:1900370 positive regulation of RNA interference	0.0001	133.79
	GO:0036466 synaptic vesicle recycling via endosome	0.0003	66.89
	GO:0036465 synaptic vesicle recycling	0.0008	44.6
Upregulated	GO:2000766 negative regulation of cytoplasmic translation	0.0003	63.5
	GO:0017148 negative regulation of translation	0.0002	9.16
	GO:0034249 negative regulation of cellular amide metabolic process	0.0004	7.81
	GO:0006417 regulation of translation	0.0001	5.69
	GO:0034248 regulation of cellular amide metabolic process	0.0001	4.68
	GO:0010608 posttranscriptional regulation of gene expression	0.0001	4.61
	GO:0010468 regulation of gene expression	0	2.21
	GO:0031326 regulation of cellular biosynthetic process	0.0001	2.04
	GO:0009889 regulation of biosynthetic process	0.0001	2.04
	GO:0010556 regulation of macromolecule biosynthetic process	0.0004	1.98
	GO:0060255 regulation of macromolecule metabolic process	0.0001	1.86
	GO:0019222 regulation of metabolic process	0.0001	1.81
	GO:0051171 regulation of nitrogen compound metabolic process	0.0005	1.76
	GO:0080090 regulation of primary metabolic process	0.0006	1.74
	GO:0031323 regulation of cellular metabolic process	0.0007	1.71
	GO:0046011 regulation of oskar mRNA translation	0	29.31
	GO:1902287 semaphorin-plexin signaling pathway involved in axon guidance	0.0001	95.26
	GO:1902285 semaphorin-plexin signaling pathway involved in neuron projection guidance	0.0001	95.26
	GO:0045876 positive regulation of sister chromatid cohesion	0.0001	95.26
	GO:2000305 semaphorin-plexin signaling pathway involved in reg of photoreceptor cell axon guidance	0.0001	95.26
	GO:0048013 ephrin receptor signaling pathway	0.0003	63.5
	GO:0071526 semaphorin-plexin signaling pathway	0	47.63
	GO:0007162 negative regulation of cell adhesion	0.0006	17.86
	GO:0051128 regulation of cellular component organization	0.0002	2.4
	GO:0048522 positive regulation of cellular process	0.0007	1.88
	GO:0099177 regulation of trans-synaptic signaling	0.0001	5.82
	GO:0050804 modulation of chemical synaptic transmission	0.0001	5.82
	GO:0016319 mushroom body development	0.001	6.52
	GO:2000026 regulation of multicellular organismal development	0.0007	2.57
	GO:0032502 developmental process	0.0001	1.81
	GO:0032879 regulation of localization	0.0006	2.72
	GO:2000112 regulation of cellular macromolecule biosynthetic process	0.0004	1.98
	GO:0050794 regulation of cellular process	0	1.84
	GO:0048518 positive regulation of biological process	0.0008	1.8
	GO:0050789 regulation of biological process	0	1.7
	GO:0065007 biological regulation	0	1.63
	GO:0042391 regulation of membrane potential	0.0008	6.9
	GO:0065008 regulation of biological quality	0.0002	2.12
	GO:0120187 positive regulation of protein localization to chromatin	0.0001	95.26
	GO:0008049 male courtship behavior	0.0003	8.36
	GO:0060179 male mating behavior	0.0005	7.56

Legend:

Regulation of biological process
Regulation of metabolic process
Regulation of macromolecular metabolic process
Regulation of cellular process
Regulation of Signalling
Regulation of development process
Regulation of localisation

Metabolic process
Biosynthetic process
Nitrogen compound metabolic process
cellular metabolic process

Biological Regulation
Regulation of biological process
Regulation of biological quality

Localisation
Establishment of localisation
Regulation of protein localisation

Behaviour
Reproductive behaviour

A: Sleep induction method

B: Induced Sleep Transcriptome: GO Pathways of Biological Processes

hour	gene	estimate	estimate1	estimate2	.y.	group1	group2	n1	n2	statistic	p	df	conf.low	conf.high	method	alternative	p.adj	p.adj.signif
Day	Da1	0.747011947979923	1.81085006902361	2.55786201700353	activity_index	control	knockout	91	93	-7.68674248159984	1.63e-12	154.732513840803	0.938986358845125	0.555037537114721	T-test	two.sided	1.63e-11	****
Night	Da1	0.621402556761431	2.20459173404308	2.82599429080452	activity_index	control	knockout	91	93	-6.52394430852639	1.39e-09	129.928135878636	0.809843317756889	0.432961795765974	T-test	two.sided	9.73e-09	****
Day	Da2	0.802741622150127	2.43199942296601	3.23474104511614	activity_index	control	knockout	70	65	-7.4427805261935	2.03e-11	113.734263925721	1.01640704197762	0.589076202322632	T-test	two.sided	1.827e-10	****
Night	Da2	1.23901539806697	2.26258176444676	3.50159716251373	activity_index	control	knockout	70	65	-10.1194970663198	8.52e-17	95.6867888875377	1.48206392653957	0.995966869594365	T-test	two.sided	1.1076e-15	****
Day	Da3	0.415534948767109	2.16037312206754	2.57590807083465	activity_index	control	knockout	91	43	-1.34323807941607	0.186	45.4132347049576	1.03844775713935	-0.207377859605138	T-test	two.sided	0.744	ns
Night	Da3	-0.0227331901857819	2.83159939569465	2.80886620550887	activity_index	control	knockout	91	43	0.0990781987694288	0.921	55.2553143772094	0.43704101471215	-0.482507395083714	T-test	two.sided	1	ns
Day	Da4	-0.61837129538045	2.42331822907521	1.80494693369476	activity_index	control	knockout	98	87	8.66914228246151	2.56e-15	178.965595681112	-0.47761492132397	-0.75912766943693	T-test	two.sided	2.816e-14	****
Night	Da4	-0.110300614309692	2.26586151382992	2.15556089952023	activity_index	control	knockout	98	87	0.835733550651872	0.405	115.736113589999	0.151109873134961	-0.371711101754344	T-test	two.sided	1	ns
Day	Da6	-0.623668327284424	2.88534347323253	2.26167514594811	activity_index	control	knockout	91	91	6.64405236041315	5.29e-10	150.02407366578	-0.43819297313673	-0.809143681432117	T-test	two.sided	4.232e-09	****
Night	Da6	0.0488610995058534	2.54149490606688	2.59035600557274	activity_index	control	knockout	91	91	-0.47619750862764	0.635	153.272240702089	0.251567217217754	-0.153845018206047	T-test	two.sided	1	ns
Day	Da7	0.326181724510013	2.22741317915403	2.55359490366404	activity_index	control	knockout	95	94	-3.12088657353367	0.00211	176.4794317341	0.532443223747717	0.119920225272308	T-test	two.sided	0.01266	*
Night	Da7	-0.236825775153197	2.70854101277	2.4717152376168	activity_index	control	knockout	95	94	1.93009604618506	0.0552	181.674501771687	0.00527760128503382	-0.478929151591428	T-test	two.sided	0.276	ns