Differences in vertical and horizontal transmission dynamics shape plasmid

distribution in clinical enterobacteria

Aida Alonso-del Valle[#,1,*], Laura Toribio-Celestino[#,1], Anna Quirant[2], Carles Tardio Pi[3,4],

Javier DelaFuente[1], Rafael Canton[5,6], Eduardo Rocha[7], Carles Ubeda[2,8], Rafael Peña-

Miller[3], Alvaro San Millan[1,8,*].

Affiliations

1. Centro Nacional de Biotecnología (CNB), CSIC. Madrid, Spain.

2. Fundación para el Fomento de la Investigación Sanitaria y Biomédica de la Comunitat

Valenciana - FISABIO, Valencia, Spain.

3. Centro de Ciencias Genómicas, Universidad Nacional Autónoma de México, Cuernavaca,

Mexico.

4. Instituto de Investigaciones en Matemáticas Aplicadas y en Sistemas, Unidad Académica

Yucatán, Universidad Nacional Autónoma de México, Yucatán, México.

5. Servicio de Microbiología. Hospital Universitario Ramón y Cajal-IRYCIS. Madrid, Spain.

6. Centro de Investigación Biológica en Red de Enfermedades Infecciosas (CIBERINFEC),

Instituto de Salud Carlos III, Madrid, Spain.

7. Institut Pasteur, Université de Paris Cité, CNRS UMR3525, Microbial Evolutionary Genomics,

Paris, France.

8. Centro de Investigación Biológica en Red de Epidemiología y Salud Pública (CIBERESP),

Instituto de Salud Carlos III, Madrid, Spain.

# Contributed equally

* Correspondence and request for materials should be addressed to Aida Alonso-del Valle

(aalonso@cnb.csic.es) & Alvaro San Millan (asanmillan@cnb.csic.es).

Abstract

Conjugative plasmids can transfer both vertically and horizontally in bacterial communities, playing a key role in the dissemination of antimicrobial resistance (AMR) genes across bacterial pathogens. AMR plasmids are widespread in clinical settings, but their distribution is not random, and certain associations between plasmids and bacterial clones are particularly successful. However, knowledge remains limited about the contribution made by vertical and horizontal transmission dynamics to plasmid distribution and maintenance in clinically relevant bacterial communities. In this study, we used a collection of wild type enterobacterial strains isolated from hospitalized patients to perform a comprehensive analysis of the transmission dynamics of the globally spread carbapenem resistance plasmid pOXA-48. We combined *in vitro* and *in vivo* experimental approaches to quantify key traits responsible for vertical (the level of AMR) and horizontal (conjugation frequency) plasmid transmission. Our results reveal significant variability in these traits across different bacterial hosts, with *Klebsiella* spp. strains showing higher pOXA-48-mediated AMR and conjugation frequencies than *Escherichia coli* strains. Using experimentally determined parameters, we developed a simple mathematical model to interrogate the contribution of vertical and horizontal transmission to plasmid distribution in bacterial communities. These simulations revealed that a small subset of clones, combining high vertical and horizontal plasmid transmission ability, play a critical role in stabilizing the plasmid in different polyclonal microbial communities. Our results indicate that strain-specific differences in plasmid transmission dynamics dictate successful associations between plasmids and bacterial clones, shaping AMR evolution.

Significance statement

Conjugative plasmids are the main vehicle for the dissemination of AMR genes across many bacterial pathogens, contributing to one of the most concerning public health problems facing modern societies. Understanding the rules governing plasmid dynamics is therefore crucial to controlling the global AMR crisis. In this study, we show that the plasmid-associated traits responsible for vertical and horizontal plasmid transmission in bacterial communities vary across different bacterial hosts. This information can be used to predict which specific plasmid-bacteria associations are more likely to spread in bacterial communities, thus enabling health care authorities to predict, and potentially control, the evolution of AMR in clinical settings.

Introduction

Plasmids shape bacterial ecology and evolution by spreading accessory genes across populations. Plasmids can disseminate both vertically and horizontally. Vertical transmission is coupled to the division of the bacterial host, whereas horizontal transmission is mediated mainly by conjugation (the direct cell-to-cell transfer of plasmids through a bridge-like connection) (1). Vertical plasmid transmission is favored in the presence of environmental stresses that select for plasmid-encoded genes, but in the absence of plasmid-specific selection, vertical transmission is usually hindered by the fitness costs that plasmids produce in their bacterial hosts (2, 3). Therefore, plasmids' interests can align or conflict with those of their bacterial hosts depending on the environmental conditions, leading to complex eco-evolutionary dynamics (4-6). Horizontal plasmid transmission can in principle make up for a deficit in vertical transmission, although conjugation rates vary considerably depending on the plasmid type (7, 8). The fate of plasmids in bacterial populations is thus determined by the interplay between horizontal and vertical transmission dynamics (and their evolution) (9-12). Multiple theoretical and experimental studies have investigated plasmid dynamics in clonal bacterial populations (7, 8, 13-17). However, information is very limited about how vertical and horizontal plasmid transmission dynamics vary across the diversity of wild type bacteria that plasmids encounter in natural communities and how this variability may affect plasmid distribution in complex microbiota (18-21).

A dramatic example of the ability of plasmids to fuel bacterial evolution is the central role they play in the spread of antimicrobial resistance (AMR) in clinical pathogens, which is one of the most urgent public health threats facing humanity (22, 23). One particularly concerning group of drug resistant pathogens is carbapenemase-producing enterobacteria (order *Enterobacterales*), which appear on the WHO "priority pathogens" list (24, 25). Carbapenemases are β-lactamase enzymes able to degrade carbapenem antibiotics and are mainly acquired through conjugative plasmids (22). pOXA-48-like plasmids (from here on pOXA-48) constitute one of the most clinically important groups of carbapenemase-producing plasmids (26). pOXA-48 is a broad-host-range conjugative plasmid from the plasmid taxonomic unit L (27) that encodes the OXA-48 carbapenemase and is disseminated across enterobacteria worldwide (26). Like most AMR plasmids, pOXA-48 is not randomly distributed across bacterial hosts, and, although it has a broad potential host range, it is strongly associated with *Klebsiella pneumoniae* and is especially

prevalent in high-risk clones of specific sequence-types (ST), such as ST11 or ST101 (28, 29). In

a previous study, we proposed that this bias in host distribution could be explained in part by

pOXA-48-associated fitness costs in the absence of antibiotics (30). However, an analysis of the

distribution of pOXA-48 fitness effects across a collection of *Escherichia coli* and *Klebsiella* spp.

clinical strains revealed that, although pOXA-48 produced a wide distribution of fitness effects

across these hosts, these effects could not explain overall plasmid distribution, especially at the

species level (30).

In the present study, we performed a comprehensive analysis of vertical and horizontal pOXA-48

transmission dynamics across clinical enterobacteria, using multidrug resistant *E. coli* and

Klebsiella spp. strains isolated from the gut microbiota of hospitalized patients at a large hospital

in Madrid, Spain (Ramon y Cajal University Hospital). Through a combination of *in vitro* and *in*

vivo experimental approaches, we demonstrate that (i) pOXA-48 confers a higher level of AMR

in *Klebsiella* spp., and that (ii) *Klebsiella* spp. strains are, on average, more permissive than *E.*

coli strains to pOXA-48 acquisition through conjugation. Integration of these new experimentally

determined parameters with our previous data in a simple population dynamics model allowed us

to interrogate how plasmid transmission dynamics determine the distribution of plasmid pOXA-48

across clinical enterobacteria.

Results

Experimental system

In this study, we used a collection of multidrug-resistant clinical enterobacterial strains isolated

from the gut microbiota of patients at a large hospital in Madrid. We focused on the two most

prevalent species associated with pOXA-48 in this hospital, *K. pneumoniae* and E. *coli* (28). Since

K. quasipneumoniae and *K. variicola* (two species previously misidentified as *K. pneumoniae*) are

also associated with pOXA-48 in the hospital (28), we included strains from these species in the

study. From now on we use *Klebsiella* spp. to refer to the strains in our collection belonging to

these three *Klebsiella* species.

To analyze pOXA-48 transmission dynamics it is necessary to work with pOXA-48-free strains,

which are able to receive the plasmid by conjugation. Therefore, we selected 25 *Escherichia coli*

and 25 *Klebsiella* spp. pOXA-48-free strains representative of the phylogenetic diversity of these

species in our hospital, and sequenced their genomes (Fig. S1 and Table S1, collection partially

overlapping with the one in ref. (30), see Methods). Although these strains were pOXA-48-naive,

they were identified as ecologically compatible with pOXA-48 because they were obtained from

patients on hospital wards where pOXA-48-carrying enterobacteria were commonly isolated from

other patients during the same period (28).

AMR level conferred by pOXA-48 in wild type enterobacteria

To characterize pOXA-48 vertical transmission dynamics in the presence of antibiotics, we

determined the impact of plasmid acquisition on the level of AMR. pOXA-48 was introduced by

conjugation into the pOXA-48-naive enterobacteria collection. Four *E. coli* strains and one *K.*

pneumoniae strain were unable to acquire the plasmid, and whole genome sequencing of the

remaining transconjugants revealed pOXA-48 mutations in six *E. coli* and six *Klebsiella* spp.

transconjugants. After excluding these strains, we used the remaining 15 *E. coli* and 18 *Klebsiella*

spp. isogenic strains pairs with or without pOXA-48 plasmid for this analysis (Fig. S1). For each

of the transconjugant (TC) and plasmid-free (PF) clones, we determined the minimal inhibitory

concentration (MIC) of 4 clinically important β-lactam antibiotics: amoxicillin in combination with

the β-lactamase inhibitor clavulanic acid (AMC) and the carbapenems ertapenem (ERT),

imipenem (IMP), and meropenem (MER) (Fig. 1A). Compared with *E. coli*, *Klebsiella* spp. strains

showed higher constitutive resistance to all four antibiotics (Wilcoxon rank sum test: AMC, $W =$

47.5, $P = 0.001$; ERT, $W = 14.5$, $P = 8.68 \times 10^{-6}$; IMP, $W = 15$, $P = 5.16 \times 10^{-6}$; MER, $W = 56$, $P =$

0.002), as well as higher resistance level after plasmid acquisition (Wilcoxon rank sum test: AMC,

$W = 27$, $P = 8.94 \times 10^{-5}$; ERT, $W = 82$, $P = 0.031$; IMP, $W = 29.5$, $P = 4.79 \times 10^{-5}$; MER, $W = 54$,

$P = 0.002$).

The diverse genomic background and AMR gene content of the strains under study made it

difficult to elucidate a genetic origin for the higher AMR levels reached by pOXA-48-carrying

Klebsiella spp. (Fig. S2). However, a prominent influence on plasmid-mediated AMR level is

plasmid copy number (PCN). AMR level commonly escalates with PCN for AMR mechanisms

showing strong gene dosage effects, such as carbapenemases (31, 32). We used the genome

sequencing data to estimate pOXA-48 PCN in the strains under study (Fig. 1B). The results

showed that PCN was higher in *Klebsiella* spp. (mean = 2.21, SD = 1.53) than in *E. coli* strains

(mean = 1.41, SD = 0.55) (Wilcoxon rank-sum test, $W = 76$, $P = 0.03$). We then compared pOXA-

48 PCN and AMR levels for the antibiotic to which pOXA-48 confers the highest level of resistance: AMC (average fold-change in MIC associated to pOXA-48 presence = 244.7). The final AMC resistance level correlated positively with PCN in the strains under study (Spearman's correlation: R = 0.5, P = 0.003; Fig. S3), supporting the idea that the higher PCN is at least partly responsible for the elevated AMR level in *Klebsiella*. spp. Finally, to investigate if the differences in PCN were also present in strains naturally carrying pOXA-48, we used the same genomic approach to analyze PCN in a recently characterized large collection of 200 pOXA-48-carrying clinical isolates from the same hospital. In line with the results from the transconjugant strains, PCN was higher in *Klebsiella* spp. (mean = 2.35, SD = 0.88) than in *E. coli* (mean = 1.46, SD = 1.33) (Fig. 1C; Wilcoxon rank-sum test, W = 1271, P = 1.32 x 10^{-14}).

Fig. 1. Resistance levels and copy number of pOXA-48 in wild type enterobacteria. pOXA-48 confers higher resistance levels and has a higher PCN in *Klebsiella* spp. than in *E. coli.* (A) MIC (mg/L) of amoxicillin/clavulanic acid, ertapenem, imipenem and meropenem for the plasmid-free/plasmid-carrying strain pairs (median of three biological replicates). (B) pOXA-48 plasmid

copy number (PCN) of the *E. coli* (n = 15) and *Klebsiella* spp. (n = 18) transconjugant strains.

PCN was estimated from sequencing data as the ratio of plasmid/chromosome median coverage

(see Methods). Horizontal lines inside boxes mark median values, the upper and lower hinges

correspond to the 25th and 75th percentiles, and whiskers extend to 1.5 times the interquartile

range. Dots indicate individual PCN values. To aid visualization, the y axis is represented in $\log_{10}$

scale with no value transformation. (C) pOXA-48 PCN in *E. coli* (n = 59) and *Klebsiella* spp. (n =

140) pOXA-48-carrying wild-type clinical strains. Boxplots are structured as in B. *P < 0.05, **P

< 0.01, ***P < 0.001.

Conjugation frequency varies across clinical strains

To analyze pOXA-48 horizontal transmission dynamics, we determined the ability of clinical

strains to acquire the plasmid by conjugation ("conjugation permissiveness"). For recipients, we

used a subset of 10 *Klebsiella spp.* and 10 *E. coli* strains representative of the genetic diversity

of the collection (Fig. S1). For donors, we used three pOXA-48-carrying strains. These were a

laboratory-adapted *E. coli* strain [β3914, a diaminopimelic acid (DAP) auxotrophic laboratory

mutant of *E. coli* K-12], and two wild-type pOXA-48-carrying clinical strains recovered from

hospitalized patients and belonging to STs responsible for between-patient plasmid dissemination

in the same hospital (28): *K. pneumoniae* ST11, and *E. coli* ST10 (strains K93 and C165

described in references 28 and 32).

We first performed classical conjugation assays (one donor and one recipient strain in equal

proportions) for all donor–recipient combinations, and measured conjugation frequencies as the

ratio between transconjugants (TC) and total recipient cells (see Methods for details). Note that

the donor *K. pneumoniae* ST11 could not be distinguished from three recipients of the same

species, either by the phenotype in the differential medium or by antibiotic selection, and therefore

conjugations could not be performed for these donor–recipient pairings. This analysis revealed a

wide diversity of conjugation frequencies across recipient strains. Overall, *Klebsiella spp.* strains

showed a significantly higher conjugation permissiveness than *E. coli* for all three donors (Fig.

2A, Kruskal-Wallis rank sum test: *E. coli* β3914, X^2 = 45.8, P = 1.9 x 10^{-11}; *E. coli* ST10, X^2 = 34.1,

P = 5.16 x 10^{-9}; *K. pneumoniae* ST11, X^2 = 6.98, P = 8.24 x 10^{-3}). Moreover, three *E. coli* strains

produced no TCs with any of the donors, whereas all *Klebsiella spp.* strains produced TCs in at

least one of the conditions and were also the recipients with highest conjugation permissiveness

for every donor.

The gut microbiota is a dense bacterial community that engages in complex ecological

interactions. To analyze if community complexity could affect conjugation permissiveness in our

clinical strains, we repeated the conjugation assays, but this time instead of using a single clone

as the receptor, we used ten pairs of the *E. coli*/*Klebsiella* spp. strains (Fig. 2B and Table S1).

This analysis once again revealed great variability in reception frequencies and a higher overall

conjugation permissiveness in *Klebsiella* spp. strains (Fig. 2B, paired Wilcoxon signed-rank exact

test: *E. coli* β3914, V = 235, P = 0.017; *E. coli* ST10, V = 205, P = 0.01; *K. pneumoniae* ST11, V

= 45, P = 1.2 x 10^{-4}). In general, the conjugation frequencies observed agreed with those from

the single conjugation assays, suggesting that ecological interactions did not markedly affect

conjugation dynamics. However, some of the *E. coli* strains that produced no TCs as single

receptors were able to acquire the plasmid, which could be due to secondary conjugations from

the *Klebsiella* spp. TC (33). Finally, to increase the complexity of the recipient community even

further, we performed conjugation assays using a pool of all 20 strains as recipients (Fig. 2C). In

this case, we could only distinguish between TCs at the species level in the differential medium.

Once again, the results revealed *Klebsiella spp.* to be more conjugation-permissive than *E. coli*,

regardless of the donor (Fig. 2C, Kruskal-Wallis rank sum test: *E. coli* β3914, X^2 = 9.28, P = 2.32

x 10^{-3}; *E. coli* ST10, X^2 = 9.31, P = 2.35 x 10^{-3}; *K. pneumoniae* ST11, X^2 = 12.8, P = 3.49 x 10^{-4}).

Fig. 2. *In vitro* **pOXA-48 conjugation dynamics in wild type enterobacteria**. Conjugation frequencies (transconjugants/recipient) of a subset of 20 isolates from our clinical enterobacteria collection (10 *E. coli* and 10 *Klebsiella* spp., x-axis), obtained from three pOXA-48 donors (*E. coli* β3914, left; *K. pneumoniae* ST11, center; and *E. coli* ST10, right). (A) Conjugation frequencies obtained in conjugation assays with one donor and one recipient in equal proportions. Color represents the recipient species (red, *E. coli*; blue, *Klebsiella* spp.), symbols represent biological replicates (n = 4), and asterisks symbolize conjugation frequencies below the detection limit (dotted line). *Klebsiella spp.* exhibits higher conjugation frequencies than *E. coli* for the three donors. (Kruskal-Wallis rank sum test: *P* < 0.01). (B) Conjugation frequencies using one donor and pairs of *E. coli* and *Klebsiella* spp. as recipients. *Klebsiella spp.* is more conjugation-permissive than *E. coli*, regardless of the donor used (Wilcoxon signed-rank exact test: *P* < 0.02). (C) Conjugation frequencies obtained with a pool of recipients (one donor, 20 recipients). The results give a conjugation frequency value per species, and symbols represent biological replicates (n=9) whose values would correspond to the most successful strains of each species

in plasmid uptake. *Klebsiella spp.* has higher conjugation frequencies than *E. coli*, regardless of

the donor (Kruskal-Wallis rank sum test: $P < 0.01$).

Conjugation dynamics in a mouse model of gut colonization

To confirm our *in vitro* results in a more biologically relevant environment, we studied pOXA-48

conjugation dynamics in a mouse model of gut colonization (C57BL/6J). As donor, we used *K.*

pneumoniae ST11 strain K93, since this clonal group is frequently responsible for gut microbiota

colonization of patients in our hospital (28). As recipients, we used 6 strains representative of the

variability in conjugation frequency observed *in vitro* (*E. coli* PF_EC10, PF_EC12, and PF_EC21;

and *Klebsiella spp*. FP_KQ01, PF_KPN11, and PF_KPN12). Mice (n = 21) were treated orally

with ampicillin (0.5 g/L), vancomycin (0.5 g/L), and neomycin (1 g/L) for one week to reduce

colonization resistance, as we previously described (34), and the antibiotic treatment was stopped

one day before inoculation. A total of 18 mice were inoculated orally with the donor and co-

inoculated 2 hours later with the recipients (3 mice per recipient), and 3 mice were not inoculated

with any bacteria, as controls. Fresh fecal samples were collected every 24 hours during 2 days.

At the end of the second day, mice were sacrificed, and cecum samples were collected. Samples

were processed and plated with antibiotic selection to determine colonization levels and

conjugation frequencies, as described in Methods. Two independent assays were performed, two

months apart, and the results of both assays are compiled below (Fig. 3 and source data).

Colonization levels were high in all inoculated mice at both 24 and 48 hours for donor and recipient

strains (Fig. 3A), while no colonization was observed in the control mice. At 24 hours post-

inoculation, conjugation frequencies were generally low, and no differences in conjugation

frequencies were detected between *E. coli* and *Klebsiella* spp. (Fig. 3B Kruskal-Wallis rank sum

test H = 1.4, P = 0.265). At 48 hours post-inoculation, the levels of donors were similar in mice

co-inoculated with *E. coli* or *Klebsiella* spp. recipient strains (Wilcoxon rank sum test: in feces, W

= 2.71, P = 0.1; in cecum, W = 0.1, P = 0.752). However, despite the fact that the levels of

Klebsiella spp. recipients were slightly lower than those of *E. coli* recipients (Wilcoxon rank sum

test: in feces, W = 245, P = 0.008; in cecum, W = 4.9, P = 0.026), *Klebsiella spp.* showed

significantly higher conjugation frequencies (Fig. 3B, Kruskal-Wallis rank sum test: in feces, $H =$

24.6, $P = 6.89 \times 10^{-7}$; in cecum, $H = 10.2$, $P = 1.4 \times 10^{-3}$). In summary, the *in vivo* results correlated

qualitatively with those from the *in vitro* conjugation assays (Fig. 2A and 3B), and *Klebsiella spp.* strains showed higher conjugation permissiveness than *E. coli* strains.

Fig. 3. pOXA-48 conjugation dynamics in a mouse model of gut colonization. pOXA-48 conjugation frequencies in mouse feces and cecum at 24 and 48 hours post colonization. (A) Colonization levels in the donor (*K. pneumoniae* ST11), recipients (3 *E. coli* and 3 *Klebsiella spp.* strains), and transconjugants, in feces at 24 and 48 hours post-inoculation (top and center) and in cecum content at 48 h (bottom). Species are indicated by colors (red, *E. coli*; blue, *Klebsiella spp.*) and biological replicates (n=6) by symbols. Black lines indicate the median values for the biological replicates. Donor and recipient colonization levels were high in both feces and cecum. (B) Conjugation frequencies (transconjugants/recipients) in feces at 24 and 48 hours (top and center) and in cecum (bottom). Conjugation frequencies were significantly higher in *Klebsiella* spp. than in *E. coli* in both feces and cecum at 48 hours (Kruskal-Wallis rank sum test in feces H = 24.6, P = 6.89 x 10^{-7}; in cecum H = 10.2, P = 1.4 x 10^{-3}).

Klebsiella*-derived capsules are associated with plasmid carriage in *E. coli

The conjugation experiments showed a higher permissiveness for pOXA-48 acquisition in

Klebsiella spp. than in *E. coli*, as well as high variability between strains of the same species. We

sought to investigate the genetic traits that could explain these differences. We analyzed the

genomes of the 20 recipient strains for an array of traits known to affect conjugation efficiency:

CRISPR arrays, restriction-modification (RM) systems and other phage defense systems that

could target pOXA-48, plasmid incompatibilities, and type VI secretion systems (T6SS, known to

mediate bacterial competition, which may affect conjugation frequencies). However, none of

these traits accounted for the observed differences in pOXA-48 acquisition (Fig. S4, Table S2).

Capsules have been reported to hinder DNA transfer and possibly constitute a barrier to plasmid

acquisition (35). We used two software tools to search for capsular systems: Kaptive, to detect

and type *Klebsiella* capsular loci, and CapsuleFinder, to identify other capsular systems. With the

single exception of the non-capsulated PF_KPN14 strain, each recipient strain encoded a specific

set or type of complete capsular loci (Table S2). To assess whether similar sets of capsular loci

were associated with higher or lower conjugation permissiveness, we calculated a weighted gene

repertoire relatedness (wGRR) score. Although we found no clear pattern for the *Klebsiella* spp.

recipients, the *E. coli* strain with the highest pOXA-48 reception rate, PF_EC25, encoded a

Klebsiella capsular locus (KL35, Fig. 4A, Table S2).

To determine if *Klebsiella*-derived capsules better facilitate pOXA-48 acquisition than other

capsular systems, we analyzed their association in the complete genomes of 1,585 *E. coli* and

715 *K. pneumoniae* strains (RefSeq), of which 6 *E. coli* strains and 46 *K. pneumoniae* strains

carry pOXA-48 (Fig. 4B, Table S3). We used phylogenetic logistic regression (*phyloglm*) to

account for phylogenetic dependency of the significant associations ($P < 0.05$, Fisher's exact test;

Table S5). The presence of *Klebsiella*-derived capsules was associated with pOXA-48 carriage

only when *E. coli* and *K. pneumoniae* were analyzed together (Table S5, *phyloglm* $P = 0.0016$).

We next investigated the relationship between the presence of *Klebsiella*-derived capsules and

plasmid carriage more generally, finding significant associations when analyzing *E. coli* and *K.*

pneumoniae together (Table S5, *phyloglm* $P < 2 \times 10^{-16}$) and when analyzing *E. coli* separately

(Table S5, *phyloglm* $P = 2.2 \times 10^{-5}$). Moreover, *E. coli* strains encoding other capsular systems

(not derived from *Klebsiella*) were more likely to be plasmid-free or to carry only one plasmid,

whereas *E. coli* encoding *Klebsiella*-derived capsules had an increased likelihood of carrying

multiple plasmids (Fig. 4C). These results suggest that although capsules generally obstruct

conjugation (35), certain types of *Klebsiella*-derived capsules could be more permissive than

other capsule types to plasmid acquisition by *E. coli*.

Fig. 4. Associations between capsular systems and plasmid carriage. Analysis of capsular

systems and plasmid prevalence in our recipient strains and in *E. coli* and *K. pneumoniae*

genomes obtained from RefSeq. (A) Heatmap of weighted gene repertoire relatedness (wGRR)

scores between capsular sets (*Klebsiella*-derived capsules and/or other capsular systems)

encoded by the recipient strains used in conjugation experiments. A wGRR score of 1 means that

all proteins from the capsular set encoding the fewest number of proteins have a homolog in the

capsular set it is compared with. A score of 0 indicates there are no homologs between the

capsular sets of the two compared strains. Recipients are ordered by mean pOXA-48 conjugation

frequency across replicates and experimental conditions. The PF_EC25 strain, highlighted in

bold, has the highest mean conjugation frequency of any *E. coli* strain and is the only *E. coli* strain

encoding a *Klebsiella*-derived capsule. Strains from the same sequence type (ST) are marked

with a symbol. (B) Number of *E. coli* (Ec) and *K. pneumoniae* (Kpn) strains from the RefSeq

database analyzed for the association between the presence of *Klebsiella*-derived capsules

(Kcaps) and carriage of pOXA-48 (left) or plasmids in general (including pOXA-48; right). In total,

the RefSeq database includes 1,585 Ec and 715 Kpn strains. (C) Association between the

presence of a genome-encoded *Klebsiella*-derived capsule and the number of plasmids carried

in the *E. coli* (Ec) and *K. pneumoniae* (Kpn) RefSeq strains. The chart shows odds ratio (Fisher's

exact test) of plasmid carriage for strains encoding a *Klebsiella*-derived capsule *vs* strains

encoding other capsular systems (not from *Klebsiella*). Values above 1 (horizontal dashed line)

indicate a positive association, and values below 1 a negative association, between encoding

Klebsiella-derived capsules and carrying the indicated number of plasmids. In all cases, the

number of strains carrying each number of plasmids is >100. A significant Fisher's test ($P < 0.05$)

is indicated by a black outline.

Modeling the impact of vertical and horizontal transmission dynamics on plasmid

distribution.

We previously developed a mathematical model to study how pOXA-48 fitness effects impact

plasmid maintenance in complex bacterial communities (32). That model allowed us to predict

the competitive fitness of each strain, with and without pOXA-48, based on growth-kinetic

parameters that were calibrated from experimental growth curves obtained from single clones

(Methods). In the present study, we extended this model to include the new variables informed

by the experimental results: conjugation permissiveness and resistance levels (Fig. 5A). We used

this extended model to numerically simulate different enterobacteria community compositions and

environmental regimes, with the aim of estimating the stability of different plasmid-host

associations. This computational model allowed us to track the distribution of plasmids in

response to fluctuating selection at strain and species resolution.

We first simulated pairwise competition experiments between the 10 *Klebsiella* spp. and 10 *E.*

coli strains in our collection with experimentally determined conjugation frequencies and AMR

levels. Each competition experiment started with a 1:1 ratio, and a 50% plasmid frequency in each

strain (Methods). The total of 1,000 stochastic simulations were performed in environments with

daily changes in drug concentration determined by a Gaussian signal noise, with a probability

density function equal to that of the normal distribution (Fig. S5). In these simulations, most

pairwise associations were able to maintain the plasmid in the absence of selection, with similar

probabilities of finding the plasmid in an *E. coli* strain (35%) or in a *Klebsiella* spp. strain (38%)

(Fig. S6-S7). To assess the role of antibiotics in the distribution of plasmids in the population, we

increased the maximal drug concentration in the environment and repeated all pairwise

competition experiments for a range of selection pressures (Fig. S6). The fraction of pairwise

associations that rendered the plasmid stable peaked at intermediate selection strength (Fig. S7).

Notably, due to the lower resistance levels exhibited by *E. coli* strains, the frequency of *Klebsiella*

spp. strains in the population increased in direct correlation with the selection strength.

We next used the model to reproduce the population dynamics of pOXA-48 in the gut microbiota

of hospitalized patients, in order to analyze the impact of vertical and horizontal transfer dynamics

in plasmid distribution. Our previous epidemiological studies revealed that patients are commonly

colonized by pOXA-48-carrying nosocomial enterobacterial clones (30). During colonization, the

plasmid can spread horizontally to resident enterobacteria in the gut community, and different

clones can subsequently maintain the plasmid in the community, producing long term pOXA-48

gut carriage (30). To reproduce plasmid invasion and dissemination in the gut microbiota

community, we used the mathematical model to perform a multistrain computer simulation in

which only one member of the community carried pOXA-48 at the onset of the simulation (starting

at 0.1% frequency in the community; Fig. S8). Similar to the pairwise comparison, computer

simulations of polymicrobial communities also resulted in a reduction in the frequency of *E. coli*

strains as the drug concentration increased (Fig. 5B). Simulation of the ecological dynamics of

bacterial communities of increasing complexity (increasing number of individuals) revealed that,

in drug-free environments, pOXA-48 was found both in *E. coli* and *Klebsiella* spp. strains

independently of the number of strains in the community (Fig. 5C). In contrast, in the presence of

antibiotics, not only was the plasmid present mostly in highly-resistant *Klebsiella* spp. strains but,

as the number of strains in the community increased, so did the relative abundance of pOXA-48-

carrying *Klebsiella* spp. strains (Fig. 5D).

To explore the theoretical distribution of pOXA-48 among the different strains in a community, we

retrospectively analyzed parameter values of strains in which the plasmid remained stable during

stochastic simulations. In no-drug environments, simulations indicated the presence of pOXA-48

in strains with high growth rates, characterized by high cell efficiency and specific affinity (Fig.

5E,G). In contrast, in the presence of antibiotics, pOXA-48 was mostly found in highly resistant

strains (Fig. 5F,H). Notably, specific strains in the community played a significant role in stabilizing

the plasmid within the population; in our numerical simulations, the plasmid remained stable in

the population when, for example, strains PF_KQ04 or PF_EC16 were present, independently of

the selection strength and the fitness effects pOXA-48 produced in other members of the

community (Fig. S9). These strains were both highly resistant and highly plasmid permissive,

suggesting that successful plasmid-host associations in the microbiota are critical determinants

of plasmid invasion in complex microbial communities.

Fig. 5: **Theoretical modeling results.** (A) Scheme of the theoretical model used to simulate

plasmid invasion of a polymicrobial community composed of M strains. Plasmid dynamics are

driven by the rates of conjugative transfer (γ) and segregational loss (λ), as well as by the degree

of resistance (κ) and growth rate ($G(R)$), in environments with a limiting resource ($R(t)$) and a

bactericidal antibiotic ($A(t)$). (B) Frequency of plasmid-bearing *E. coli* strains (red) and a

Klebsiella spp. strains (blue) as a function of the maximum selection strength imposed by the

environment (A_{max}). As selection for the plasmid increases, the frequency of plasmid-carrying *E.*

coli strains is reduced, whereas the frequency of *Klebsiella* spp. strains remains constant. (C-D)

Frequency of plasmid-bearing strains as a function of the number of individuals in the community.

Each numerical experiment was performed for 100 random communities exposed to 10 different

stochastic environments, with increasing selective pressures. Top panel shows drug free

environments, $A_{max} = 0$, and bottom panel illustrates intermediate selective pressures, $A_{max} =$

0.1. In all cases, increasing the complexity of the population increases the likelihood of the plasmid

finding a successful association with a *Klebsiella* spp. strain. This effect is particularly pronounced

as the strength of selection increases. (E-F) Bidimensional representation of growth kinetic

parameters. *E. coli* strains are represented in red and *Klebsiella* spp. strains in blue ($A_{max} = 0$;

panel E, $A_{max} = 0.1$; panel F). Empty circles represent plasmid-free strains and solid circles

represent plasmid-carrying strains. The diameter of each circle is proportional to the number of

times each strain was detected at the end of a numerical experiment. Ellipses illustrate standard

deviations of best-fit normal distributions (plasmid-bearing strains in green and plasmid-free cells

in orange). (G-H) Plots illustrating conjugation permissiveness (TC/recipients) and level of AMR

(MIC in mg/L) of each plasmid-carrying strain. Vertical dotted line in the conjugation frequency

axis illustrates the detection limit. In the absence of selection ($A_{max} = 0$; panel G), both *Klebsiella*

spp. and *E. coli* strains with a high specific affinity and cell efficiency were frequently detected. In

the presence of positive selection for the plasmid ($A_{max} = 0.1$; panel H), almost exclusively

Klebsiella spp. strains with high levels of resistance survive. These findings suggest that antibiotic

selection plays a crucial role in shaping plasmid distribution in clinical enterobacteria.

Discussion

In this study, we performed a comprehensive analysis of the vertical and horizontal transmission

dynamics of the carbapenem resistance plasmid pOXA-48 in clinical enterobacteria. An important

conclusion of our study is that these dynamics are far from being constant across different wild

type bacterial hosts. We had previously shown that pOXA-48 produces variable fitness effects in

the absence of antibiotics (30), and in this new study we reveal similar variability in traits important

for vertical transmission in the presence of antibiotics (AMR levels, Fig. 1), and horizontal

transmission (conjugation, Figs. 2 and 3). This variability is especially evident for horizontal

transmission, with conjugation frequencies differing by orders of magnitude across different

recipients. Despite this variability, our results indicate that, compared with *E. coli*, *Klebsiella* spp.

clones generally have a higher AMR level associated with pOXA-48 carriage and a greater ability

to take up pOXA-48 by conjugation. These results may help to explain why pOXA-48 is usually

associated with *K. pneumoniae* clones in clinical settings (28, 29).

A key advantage of experimentally determining plasmid-associated traits in a collection of wild

type clinical enterobacteria is that we can confidently feed these variables into mathematical

models in order to ask general questions about pOXA-48 dynamics in complex bacterial

communities. In this study, we developed a model including three key variables informed by our

experimental results: bacterial growth dynamics with and without pOXA-48, conjugation

permissiveness, and AMR levels. We used this model to investigate plasmid dynamics under

different fluctuating regimes of selection pressure and community complexity, aiming to simulate

pOXA-48 invasion of a gut microbiota (Fig. 5). Our results reveal that vertical transmission is

probably the most critical factor modulating the distribution of plasmids in polymicrobial

communities. Specifically, in the absence of antibiotics, the growth dynamics of plasmid-carrying

clones determine which strains maintain the plasmid (Fig. 5E), whereas in the presence of

antibiotics, plasmid distribution is primarily determined by AMR level (Fig 5H). However, as

expected, horizontal transmission also plays a significant role in plasmid distribution in

communities, and strains showing low-to-zero conjugation permissiveness very rarely maintain

pOXA-48 (conjugation frequency $< 10^{-4}$, Fig. 5G,H).

Arguably the most important result of our simulations is that a single successful pOXA-48-

bacterium association is sufficient to allow for plasmid invasion and stable maintenance in a

complex population, regardless of plasmid effects on the remaining community members. These

jackpot associations, combining high growth rates, high AMR levels, and high conjugation

permissiveness, can act as pOXA-48 super-sinks in the gut microbiota of patients, maintaining

and further disseminating the plasmid. More generally, our results indicate that plasmid

transmission dynamics can help predict successful associations between high-risk bacterial

clones and AMR plasmids of epidemiological relevance. For example, we showed that the two *K.*

quasipneumoniae strains tested in this study (PF_KQ01 and PF_KQ04) produced successful

associations with pOXA-48 (Fig. S9), suggesting that this species, which has been previously

associated with pOXA-48 in our hospitals and others (28, 29 36-38), may play an important role

in pOXA-48 epidemiology. Our results could also have implications for the design of interventions

aimed at reducing the spread of antibiotic resistance. For example, new strategies that specifically

target these successful associations, using biotechnological tools such as those based on

CRISPR or toxin-inteins (39, 40), would dramatically increase our ability to control AMR

dissemination in clinical settings.

Given the importance of vertical and horizontal plasmid transmission in the evolution of plasmid-

mediated AMR, a critical research direction in the field will be to characterize the molecular basis

of the observed high variability in transmission-associated traits. Understanding the plasmid and

host factors that determine this variability will help us not only to predict, but also hopefully to

counteract successful associations between AMR plasmids and high-risk bacterial clones. In this

study, we showed that the presence of *Klebsiella*-derived capsules, compared to other capsular

systems, is associated with a higher plasmid frequency in *E. coli* clones (Figure 4) and that pOXA-

48 PCN is higher in *Klebsiella* spp. than in *E. coli* clones (Figure 1). Further work will be needed

to characterize the molecular basis and significance of these and other specific interactions.

Methods

pOXA-48_K8 plasmid, strains, and growth conditions. To analyze the pOXA-48-associated

AMR level, we selected 33 isogenic strains pairs (15 *E. coli* and 18 *Klebsiella* spp.) with (TC) or

without pOXA-48 plasmid (PF) from our collection of ecologically compatible enterobacterial

isolates from the human gut microbiota (Table S1, partially overlapping the collection in (30)).

Plasmid-bearing strains carry the most common pOXA-48-like plasmid variant from our hospital,

pOXA-48_K8 (26). For the conjugation frequencies determination, we selected, as recipients, 10

E. coli and 10 *Klebsiella spp.* pOXA-48-free isolates that cover the phylogenetic diversity of the

collection (Fig. S1) (30) and showed similar β-lactam resistance levels. As plasmid donors we

selected three pOXA-48_K8-carrying strains: *E. coli* β3914 (30, 40), *K. pneumoniae* K93 (32) and

E. coli C165 (32). Bacterial strains were cultured in LB at 37 °C with continuous shaking (250

r.p.m), and on LB or HiCrome™ UTI (Himedia Laboratories, India) agar plates at 37 °C.

Determination of minimum inhibitory concentration (MIC). We determined the MIC of AMC

(Normon, Spain), ERT (Merck Sharp & Dohme B.V., Netherlands), IMP (Fresenius Kabi,

Germany) and MER (SunPharma, India) in all plasmid-carrying and plasmid-free clones selected

for this study following a modified version of the agar dilution protocol (42, 43). This method allows

us to test a large number of isolates simultaneously under identical conditions and to test a wide

range of AMC concentrations, since this β-lactam antibiotic degrades fast in liquid medium (44).

We prepared pre-cultures of plasmid-free and plasmid-carrying strains by inoculating single

independent colonies into LB broth in 96-well plates and overnight incubation at 37 ºC with

continuous shaking (250 r.p.m.). We spotted 10 µl of each overnight culture onto LB agar plates

with increasing concentrations of AMC (from 4 mg/L to 32,768 mg/L), ERT (from 0,25 mg/L to 512

mg/L), IMP (from 0.25 mg/L to 1,024 mg/L) or MER (from 0.25 mg/L to 32 mg/L), and incubated

them overnight at 37 ºC. We determined the MIC of each strain as the lower antimicrobial

concentration able to inhibit the growth of each strain. We performed three biological replicates

per MIC determination in three independent days, and we used the median value of the three

replicates as the MIC.

***In vitro* determination of conjugation frequencies.** To determine the *in vitro* conjugation

frequencies, we selected, as recipient strains, 10 *E. coli* and 10 *Klebsiella spp.* pOXA-48-free

isolates, from our collection of ecologically compatible enterobacterial isolates (Fig. S1). As

plasmid donors we used three pOXA-48_K8-carrying strains: *E. coli* β3914, a counter-selectable

diaminopimelic acid auxotrophic laboratory strain (30); *K. pneumoniae* K93 (322), a pOXA-48-

carrying ST11 strain; and *E. coli* C165 (32), a pOXA-48-carrying ST10 strain. To counter-select

the two natural donors, we transformed the recipients with pBGC plasmid, a non-mobilizable *gfp-*

carrying small plasmid that encodes a chloramphenicol resistance gene (30). It was not possible

to introduce pBGC into three *Klebsiella spp.* isolates (PF_KPN09, PF_KPN14 y PF_KPN16), due

to their level of constitutive resistance to chloramphenicol. We streaked donor and recipient

strains from freezer stocks onto LB agar plates with or without ertapenem 0.5 mg/L –and 0.3 mM

DAP for *E. coli* β3914/pOXA-48_K8–, respectively, and incubated at 37 °C overnight. We

prepared pre-cultures of donors and recipients by inoculating single independent colonies in 15-

ml culture tubes containing 2 ml of LB broth and overnight incubation at 37 ºC with continuous

shaking (250 r.p.m.). To perform classic conjugation assays, we mixed one donor and one

recipient at equal proportions (each donor with each recipient, independently, with a total of 4

recipient biological replicates by donor), plated the mixture as a drop onto LB agar plates –

supplemented with DAP 0.3mM for *E. coli* β3914/pOXA-48_K8 donor– and incubated them at 37

ºC during only 4h (to avoid secondary conjugation events and the impact of potential differences

in donor, recipient, and transconjugant growth rates on conjugation frequency determination).

After incubation, we collected the biomass, resuspended it in 1 ml of NaCl 0.9 % and performed

1/10 serial dilutions up to 10^{-6}. We estimated the final densities of transconjugants and recipients

by plating 10 µl of each dilution on HiCrome™ UTI agar plates (HIMEDIA laboratories, India) with

or without AMC 256 mg/L, respectively, and with chloramphenicol 50 mg/L. We plated the

dilutions as drops, and each drop was subsequently allowed to drain down the plate, this way we

can separate the colonies so they are easily distinguishable and quantifiable. We performed the

donor counter-selection by the absence of DAP in the medium for *E. coli* β3914/pOXA-48_K8, or

by selection with chloramphenicol for clinical *K. pneumoniae* ST11 and *E. coli* ST10. Finally, we

calculated the conjugation frequencies per recipient following the formula below:

516
$$Conjugation\ frequency_R = \frac{CFU_{transconjugants}}{CFU_{recipient}}$$

where *CFU$_{transconjugants}$* are the transconjugant colony-forming units obtained from the number of

colonies in the plate (corrected by the dilution factor); and *CFU$_{recipient}$* are the number of recipient

colony-forming units minus transconjugants. Note that in the cases where no TC colonies were

detected in the plate, we calculated a threshold conjugation frequency assuming the presence of

a single colony in the total cell suspension, thus establishing a detection limit of 10^{-6}.

To perform conjugation assays with recipient pairs we randomly grouped the recipients in pairs

so that each pair comprised one *E. coli* and one *Klebsiella spp.* recipient. We used a modified

version of the above protocol, in which we mixed recipient pre-cultures of each pair at equal

proportions and we subsequently mixed these recipient mixtures with the pre-culture of the donor

at the same proportion, forming the final conjugation reaction (each recipient pair with one donor

per time, with 4 biological replicates of recipient pairs). We determined the recipients and

transconjugants densities by antibiotic selection with AMC 256 mg/L, (to select transconjugants

vs. recipients) and chloramphenicol 50 mg/L (to counter-select donors), and distinguished

between recipients using the differential chromogenic medium HiCrome™ UTI agar. We

calculated the conjugation frequency per each receptor using the formula above.

For conjugation assays with a pool of all the recipients, we mixed the same pre-culture volume of

all recipient strains with one donor per experiment, following the same protocol previously

described. Nine biological replicates were performed.

***In vivo* determination of conjugation frequencies.** For the determination of conjugation

frequencies of clinical isolates *in vivo*, we performed conjugation assays using a mouse model of

gut colonization. We selected a subset of 6 pBGC-carrying (30) recipient strains from the *in vitro*

conjugation assays: 3 *E. coli* (PF_EC10, PF_EC12 and PF_EC21) and 3 *Klebsiella spp.*

(PF_KQ01, PF_KPN11 and PF_KPN12) strains. As plasmid donor we selected the natural pOXA-

48_K8-carrying isolate, *K. pneumoniae* ST11, K93 (32). We carried out the experiment described

below twice, 2 months apart and under identical conditions.

Mouse model and housing conditions. We used 6-week-old C57BL/6J female mice (n =

21) purchased from Charles River Laboratories and housed them with autoclave-sterilized food

(a 1:1 mixture of 2014S Teklad Global diet and 2019S Teklad Global Extrused 19% Protein

Rodent Diet from Envigo) and autoclave-sterilized water. Temperature was kept at 21 ± 2 °C and

humidity was maintained at 60–70%, in 12 h light/dark cycles. To allow intestinal colonization of

the donor and recipient strains, we disrupted the intestinal microbiota by administering ampicillin

(500 mg/L), vancomycin (500 mg/L) and neomycin (1 g/L) in the drinking water for one week, as

previously described (34). Water with antibiotics was changed every 3-4 days to avoid reduction

of antibiotic activity. During antibiotic treatment, mice were co-housed (3-6 mice per cage). After

1 week of treatment, antibiotics were removed from the drinking water and mice were individually

housed to avoid bacterial transfer between mice due to coprophagia during the experiment.

Mice inoculation with donor and recipient strains. 1 day after antibiotic treatment removal,

18 out of 21 mice were inoculated by oral gavage with 10^7 CFU of the donor strain, *K. pneumoniae*

ST11. Two hours later, each donor-inoculated mouse was co-inoculated by oral gavage with 10^7

CFU of a recipient strain (3 mice per recipient). This way, from the initial 21 mice, 18 of them were

inoculated with donor and recipient bacteria (6 different recipients, 3 mice per recipient) and 3

mice remained uninoculated as control.

Sample collection and bacterial levels determination. After antibiotic treatment and prior

to inoculation, we collected fresh fecal samples from each mouse as to control for the absence of

bacteria able to grow on the selective medium used for our bacteria of interest. We weighted the

feces and resuspended them in 1 ml PBS until a homogeneous mixture was obtained. Then, we

diluted mixtures 1:100,000 and plated 100 µl on HiCrome_{TM} UTI agar plates with ampicillin (400

mg/L) and vancomycin (1 mg/L). We incubated the plates overnight at 37 ºC. After incubation,

bacterial growth was found to be absent. During the assay, we collected fresh fecal samples from

each mouse 24 and 48 h post inoculation and processed samples as described before. To

determine the donor, recipients and transconjugants densities, we plated 100 µl of different

dilutions onto HiCrome~TM~ UTI agar plates with: i) vancomycin (1 mg/L) and AMC (256 mg/L) to

detect the donor; ii) vancomycin (1 mg/L) and chloramphenicol (50 mg/L) to detect recipients; and

iii) vancomycin (1 mg/L), AMC (256 mg/L) and chloramphenicol (50 mg/L) to determine the

presence of transconjugants. We calculated the colonization levels of the members of each

conjugation assay with the following formula:

573
$$UFC/mg_{donor,recipient} = \frac{UFC_{donor,recipient} - UFC_{transconjugants}}{mg_{feces}}$$

From the levels of recipients and transconjugants we calculated the conjugation frequencies per

recipient as described before in the *in vitro* protocol. At 48 h post inoculation, after fresh fecal

samples collection, we sacrificed mice and obtained samples of cecum content by manual

extrusion. We performed sample processing, bacterial levels determination and conjugation

frequencies determination as described above.

DNA extraction and genome sequencing. We isolated the genomic DNA of the strains used in

this work using the Wizard genomic DNA purification kit (Promega, WI, USA) and quantified using

the Qubit 4 fluorometer (ThermoFisher Scientific, MA, USA). Whole genome sequencing was

performed at the Wellcome Trust Centre for Human Genetics (Oxford, UK) using the Illumina

HiSeq4000 or NovaSeq6000 platforms with 125 or 150 base pair (bp) paired-end reads,

respectively. Read data is available under the BioProject PRJNA803387.

Processing of sequencing data. All Illumina reads were trimmed with a quality threshold of 20

and reads shorter than 50 bp and adapters were removed using Trim Galore v0.6.6

(https://github.com/FelixKrueger/TrimGalore). PF and TC genomes were *de novo* assembled with

SPAdes v3.15.2 (45) in --isolate mode and with the --cov-cutoff flag set to auto. Assembly quality

was assessed with QUAST v5.0.2 (46). TC were confirmed to correspond to their PF strain by

checking mutations and plasmid content with Snippy v4.6.0 (https://github.com/tseemann/snippy)

and ABRicate v1.0.1 (PlasmidFinder database, https://github.com/tseemann/abricate),

respectively. Snippy was also used to confirm isogenicity of the acquired pOXA-48 plasmids.

Multilocus sequence typing was performed with MLST v2.21.0

(https://github.com/tseemann/mlst).

Construction of phylogenetic trees. Mash distance phylogenies were constructed for the 25 *E.*

coli and 25 *Klebsiella* spp. PF strains from their whole-genome assemblies using mashtree v1.2.0

(47) with a bootstrap of 100 (Fig. S1).

Plasmid copy number (PCN) estimation. PCN of pOXA-48 was determined for the 33 TC

strains and the 200 wild-type *E. coli* and *Klebsiella* spp. R-GNOSIS strains (BioProject

PRJNA626430) carrying pOXA-48 variants sharing 72% of the pOXA-48_K8 core genome (32).

PCN was estimated from read coverage data as the ratio of plasmid median coverage by

chromosome median coverage (45, 46). Since the genome assemblies are fragmented, the

median coverage of the chromosome was calculated from the first three, largest contigs (total

size sum of first three contigs 0.3-4.0 Mb), which were confirmed to correspond to chromosomal

sequences by using BLASTn against the NCBI nr nucleotide database. The median coverage of

pOXA-48 was calculated from the contig containing the IncL replicon, as identified with ABRicate

v1.0.1 with the PlasmidFinder database, which generally corresponds to the largest pOXA-48

contig (size 11.0-59.8 Kb). For computing the PCN, the Illumina trimmed reads were first mapped

to their respective genome assembly using BWA MEM v0.7.17 (50). Strain K25 from the wild-type

pOXA-48-carrying collection was removed from the analysis due to truncated reads. Samtools

depth v1.12 (51) with the -a flag was used to obtain read depths at each genomic position and

the median read coverage for pOXA-48 and chromosome was computed with GNU datamash

v1.4 (gnu.org/software/datamash).

Comparative genomic analysis of recipient strains. We analyzed the genomes of the recipient

strains to find traits that could explain the observed differences in pOXA-48 acquisition. First, the

draft genomes of the 20 recipients and three donors (*E. coli* ST10 C165 and *K. pneumoniae* ST11

K93 from Bioproject PRJNA626430; for *E. coli* β3914, the sequence of the ancestral K-12 strain,

NC_000913.3) were annotated with Prokka v1.14.6 (52) with default settings. A summary of the

results can be found in Table S2.

Restriction-modification (RM) systems were searched using two approaches. First, the protein

sequences of the recipient and donor strains were blasted against the protein "gold standard"

REBASE database (downloaded February 2022, (53)) using BLASTp v2.9.0 (54). The results

were filtered with a criteria based on (55). Briefly, hits with type I and IIG RM systems were kept

if the percentage of identity and alignment was 100%, since it was observed that recognition sites

varied between enzymes with only a few mismatches. For type II systems, a threshold of 70%

identity and 90% alignment was used, as it was previously reported that these systems with

identity over 55% generally share the same target motifs. For type III and type IV this threshold

was higher, as reported in the same study, and was set at 90% identity and 90% alignment. After

filtering, some proteins had hits with more than one enzyme. For these cases, the best hit or the

hit with the enzyme of the corresponding organism (*E. coli* or *Klebsiella* spp.) was kept. The

DefenseFinder webserver (accessed February 2022, (56, 57)) was also used to search for RM

systems in the proteomes of recipients and donors. Then, the REBASE hits and the RM systems

detected by DefenseFinder were unified by protein ID. Enzymes or RM systems present in all

strains were discarded, since they would not explain differences in conjugation frequencies

between isolates. Finally, only complete RM systems that were not present in the donor strains

were retained for each recipient-donor combination, since the donor would be conferring

protection to pOXA-48 against equivalent systems of the recipient. Complete type I systems

comprise a restriction enzyme (REase), a methylase (MTase) and a specificity (S) subunit. Type

II systems include at least a REase and a MTase. No type III systems were detected, and type IV

systems are normally composed of one or two proteins. Donor and recipient systems were

regarded as similar if they shared the same recognition sequence or if a BLASTp alignment of

the enzymes followed the previous criteria of identity and alignment percentage per type of

system. The same method was applied for comparing the final systems between recipient strains,

and similar systems within each type were given the same letter code identifying the RM system

subtype (Fig. S4, Table S2).

Clustered regularly interspaced short palindromic repeats (CRISPRs) were identified in the

nucleotide sequences of the recipient strains with the DefenseFinder webserver. The spacer

sequences were aligned to pOXA-48_K8 (MT441554) using the blastn-short function of BLASTn

v2.9.0 (54) and hits were then filtered by percentage of alignment. Other defense systems were

also detected with the DefenseFinder webserver and PADLOC v1.1.0 (database v1.3.0, (58)).

Plasmid incompatibility was investigated by identifying plasmid replicons using ABRicate v1.0.1

with the PlasmidFinder database.

Secretion systems were searched using the TXSScan models of MacSyFinder v1.0.5 (56, 59),

selecting the option "ordered_replicon" and linear topology. Only complete systems were

considered.

Capsular systems were identified using two software. First, *Klebsiella* capsules, which belong to

the Wzx/Wzy-dependent group I capsular systems, were identified with Kaptive v2.0.0 (60), using

the K locus primary reference and default parameters. The presence of *Klebsiella* capsules was

assigned when the confidence level of the matches was above "good", as recommended by the

authors. Other capsular systems from groups I (Wzx/Wzy-dependent), II and III (ABC-dependent),

IV, synthase-dependent (cps3-like and hyaluronic acid) and PGA (Poly-γ-d-glutamate) were

searched using MacSyFinder v1.0.5 with the CapsuleFinder for diderms models, indicating

"ordered_replicon" as database type and linear topology (56, 61). Only systems reported as

complete were considered. In the cases where *Klebsiella* capsular loci were also identified by

CapsuleFinder as group I capsules, analyses were performed with the Kaptive output. To assess

the similarity of the different sets of capsular loci between strains, a weighted gene repertoire

relatedness (wGRR) score was calculated as in (62). For this, significant sequence similarities (e-

value <10^{-4}, identity ≥35%, coverage ≥50%) among all pairs of proteins in the capsular loci were

searched using MMSeqs2 (Nature Biotechnology release, August 2017, (63)) with the sensitivity

parameter set to 7.5. The best bi-directional hits between pairs of capsule loci sets were then

used to compute the wGRR as:

$$wGRR_{A,B} = \Sigma_i \frac{id(A_i, B_i)}{min(A,B)}$$

where *id(A$_i$, B$_j$)* is the sequence identity between each pair *i* of homologous proteins in *A* and *B*,

and *min(A, B)* is the number of proteins of the capsule loci set encoding the fewest proteins (*A* or

B). The wGRR takes values between 0 (no homologous proteins between capsule loci sets) and

1 (all genes of the smallest capsule loci set have an identical homologous in the loci set of the

other strain), representing the fraction of homologs in the smallest of the two capsule loci set

weighted by their sequence similarity. Protein sequences of the *Klebsiella* capsular loci were

obtained from Prokka annotations of the operon nucleotide sequences output by Kaptive. All

proteins between the borders of CapsuleFinder complete systems were included.

Analysis of capsular systems in a database. The database comprised the genomes of 730 *K.*

pneumoniae and 1,585 *E. coli* downloaded from the NCBI RefSeq database of high quality

complete non-redundant genomes (retrieved from ftp://ftp.ncbi.nlm.nih.gov/genomes/refseq/ on

March 2021; Table S3) (64). pOXA-48-carrying strains were identified in the RefSeq database

when the best, largest BLASTn hit between the sequence of pOXA-48_K8 (MT441554) and the

target plasmid had >95% identity and >10 Kb alignment, and when the plasmid sequence

contained the pOXA-48 IncL replicon and the *bla*$_{OXA-48}$ gene as detected by ABRicate with the

PlasmidFinder and ResFinder databases. The plasmids of 6 *E. coli* and 46 *K. pneumoniae* strains

met this criteria, which had lengths between 50.6-74.2 Kb (average 64.0 Kb; Table S3). Kaptive

and CapsuleFinder were used as previously described to search for *Klebsiella*-derived capsules

and capsular systems, respectively, and to discard non-capsulated strains (15 *K. pneumoniae*

strains).

Associations between the presence of *Klebsiella*-derived capsules and pOXA-48- or plasmid-

carriage were analyzed building 2x2 contingency tables. Significant interactions ($P < 0.05$, Fisher

exact test) were then corrected for phylogenetic dependency, i.e. the tendency of closely related

strains to share the same traits. For this, a set of 128 HMM profiles (Pfam release 35, Table S4)

of conserved bacterial single copy genes, described in (65) and curated in (66), were identified in

the   RefSeq   strains   and   an   *Enterobacter   cloacae*   outgroup   (accession   number

GCF_003204095.1) using HMMER v3.3 (option --cut_ga) (67). Hits were filtered by score using

the cutoffs reported in (65), resulting in 127 gene markers present in >90% of strains. Protein

sequences of each family were aligned with MAFFT v7.453 (option --auto) (68) and alignments

were   trimmed   with   trimAl   v1.4.rev15   (69).   IQ-TREE   v1.6.12   (70)   was   used   to   infer   two

phylogenetic trees from the concatenated alignments (the first including all capsulated strains,

and the second including only *E. coli* strains, both with the *E. cloacae* outgroup) with best

evolutionary model selection and 1000 ultrafast bootstrap. Trees were rooted and rescaled to a

total height of 1. Phylogenetic logistic regression for each significant association was performed

with the function *phyloglm* from the *phylolm* v2.6.4 R package (71), fitting a logistic MPLE model

with 100 independent bootstrap replicates.

Mathematical model. The plasmid population dynamics model consists of a set of differential equations describing changes in the abundance of bacterial subpopulations competing for a single limiting resource in a homogeneous environment. If the time-dependant concentration of the limiting resource is denoted by $R(t)$, then growth of each strain can be modeled as a Monod term with the following growth kinetic parameters: a maximum uptake rate (V_{max}), a half-saturation constant (K_m), and a resource conversion coefficient (ρ). Parameter values of each strain in our collection were previously obtained using a Markov chain Monte Carlo algorithm with a Metropolis-Hastings sampler (30). Convergence of the Markov chains after a burn-in period and a thinning of 100 iterations was verified by obtaining Gelman–Rubin $\hat{R}$ statistical values near one for all strains. Robustness to different priors was verified by obtaining similar maximum likelihood estimates for uniform, lognormal, gamma, and beta prior distributions. The identifiability of growth kinetic parameters was confirmed using a data cloning algorithm (72).

To model the action of a bactericidal antibiotic in each subpopulation, we explicitly considered the environmental concentration of an antibiotic with a time-dependant variable $A(t)$. Parameter α denotes the rate at which antibiotic molecules are inactivated by binding to their cellular targets, and κ the corresponding killing rate. Resistance to the antibiotic is a consequence of a reduced killing rate, resulting from encoding a resistance gene in a mobile genetic element. Therefore the model considers that each strain can be either plasmid-bearing or plasmid-free, with densities of each subpopulation denoted by B_p and $B_{\oslash}$ respectively (we also use the same subindex p or $\oslash$ to denote parameter values corresponding to plasmid-bearing or plasmid-free cells). We assume that plasmids are randomly segregated between daughter cells, so plasmid-free cells are produced from the plasmid-bearing subpopulation at a rate λ. To model horizontal gene transfer, we consider that conjugation is proportional to the densities of both donor and recipient cells. Therefore the rate of conjugation can be represented with γ, a parameter that is determined by the plasmid permissiveness of the receiving cell to pOXA-48. Then the system of differential equations that describes the population dynamics of plasmid-free and plasmid-bearing subpopulations can be written as

$$737 \qquad \frac{dR}{dt} = -u_p(R)B_p - u_\oslash(R)B_\oslash - \delta R,$$

$$738 \qquad \frac{dA}{dt} = -\alpha A(B_p + B_\oslash) - \delta A,$$

$$739 \qquad \frac{dB_p}{dt} = \rho_p u_p(R)(1 - \lambda - \kappa_p A)B_p + \gamma B_p B_\oslash - \delta B_p,$$

$$740 \qquad \frac{dB_\oslash}{dt} = \rho_\oslash u_\oslash(R)(1 - \kappa_\oslash A)B_\oslash + \lambda(\rho_p u_p(R)B_p) - \gamma B_p B_\oslash - \delta B_\oslash,$$

where δ denotes a constant dilution rate and $u(R) = \frac{V_{max}R}{K_m+R}$ denotes a resource uptake function.

Stochastic simulations of polymicrobial communities. Computer experiments were

performed by numerically solving the population dynamics model for different environmental

regimes and population structures. Random environments were produced in Python using a

dedicated package that generates realizations of different stochastic processes, categorized

under continuous, discrete, diffusion, and noise methods. To model random environments with

different strengths of selection, we multiplied this stochastic time series by an amplitude

parameter, $0 \le A_{max} \le 1$, representing the maximum selective pressure imposed by the

environment. In particular, numerical experiments presented were performed with a daily drug

concentration obtained with a probability density function equal to that of a Normal distribution.

We repeated the experiments for environments characterized by other sources of noise and

obtained qualitatively the same distribution of plasmids in the population.

Polyclonal communities of M individuals were assembled by randomly sampling M<N individuals

from our collection of N=20 strains (10 Ec and 10 Kpn). If we denote with B_p^i and $B_\oslash^i$ the densities

of plasmid-bearing and plasmid-free subpopulations of strain i, then the bacterial community can

be represented with vector $B(t) = (B_p^1(t), B_p^2(t), \ldots, B_p^M(t), B_\oslash^1(t), B_\oslash^2(t), \ldots, B_\oslash^M(t))$. As all cells

consume the limiting resource at a rate determined by their corresponding growth kinetic

parameters, then the equation that describes the rate of change of resource R can be written as

$$760 \qquad \frac{dR}{dt} = -\sum_{i=1}^{M} \left(u_p^i(R(t)) \cdot B_p^i(t) + u_\oslash^i(R(t)) \cdot B_\oslash^i(t) \right) - \delta R.$$

761   We assume that antibiotic molecules are inactivated at a rate that is proportional to the total

762   bacterial population and the environmental drug concentration,

763
$$\frac{dA}{dt} = -\alpha A(t) \sum_{i=1}^{M} (B_p^i(t) + B_\oslash^i(t)) - \delta A.$$

764   As the model considers that plasmid conjugation is restricted by the receiving cell, then we denote

765   with γ^i the permissiveness of strain i towards pOXA-48. Therefore bacterial growth of the plasmid-

766   bearing subpopulation of strain i depends on the contribution of all plasmid-bearing populations,

767
$$\frac{dB_p^i}{dt} = \rho_p^i u_p^i(R)(1 - \lambda - \kappa_p^i A) \cdot B_p^i + \gamma^i B_\oslash^i \cdot \sum_{j=1}^{M} B_p^j - \delta B_p^i .$$

768   Similarly, the population dynamics of plasmid-free strain i can be written as

769
$$\frac{dB_\oslash^i}{dt} = \rho_\oslash u_\oslash(R)(1 - \kappa_\oslash A)B_\oslash^i + \lambda(\rho_p u_p(R)B_p^i) - \gamma^i B_\oslash^i \sum_{j=1}^{M} B_p^j - d\delta.$$

Initial conditions of competition experiments assumed that all strains in the community, both with

and without plasmid, were present at equal initial densities at time $t = 0$. For plasmid invasion

experiments, we considered a community of plasmid-free cells, with 0.1% of cells of a random

strain carrying plasmids. Computer simulations were implemented in Matlab using standard

numerical solvers and scientific libraries. All code and data used is available at

https://github.com/ccg-esb/EvK.

Statistical analyses. All statistical analyses were performed with R v4.2.2 (www.R-project.org).

Packages *ggplot2* v3.3.6, *ggpubr* v0.4.0 *pheatmap* v1.0.12, *RColorBrewer* v1.1-3, *ggsignif* v0.6.3

and *tidyverse* v1.3.1 were used for data manipulation and representation. R base packages were

used for statistical tests. Normality of the data was assessed with the Shapiro-Wilk test. To

analyze the differences in initial (PF) and final (TC) MIC of antibiotics and PCN between *E. coli*

and *Klebsiella* spp. a Wilcoxon sum-rank test was used. To analyze the differences between the

members of each recipient couple in the pairwise conjugation experiments (Shapiro-Wilk

normality test, $P < 0.01$ in all cases), the non-parametric Wilcoxon paired signed-rank test was

used. To determine differences between receptor species in the classical and pooled recipient

conjugation experiments, as well as in *in vivo* conjugation assays (Shapiro-Wilk normality test, P

< 0.01 in all cases), the non-parametric Kruskal-Wallis test was performed. Correlations were

performed with the Spearman's rank test. Phylogenetic analyses were performed with *ape* v5.6-

2, *phytools* v1.0-3 and *phylolm* v2.6.4 packages.

Data and code availability. Sequencing data generated in this study is available in the Sequence

Read Archive (SRA) under the BioProject PRJNA803387. The R-GNOSIS sequences were

generated in (28) and a subset of strains were selected as in (32) (see Methods). Source data is

provided. The code generated during the study along with detailed bioinformatic methods is

available at https://github.com/LaboraTORIbio/super-sinks.

Acknowledgements

We thank J. Penadés for constructive comments. This work was supported by the European

Research Council (ERC) under the European Union's Horizon 2020 research and innovation

programme (ERC grant no. 757440-PLASREVOLUTION) and by the Instituto de Salud Carlos III

(PI19/00749) cofunded by the European Development Regional Fund 'A way to achieve Europe'.

The R-GNOSIS project received financial support from the European Commission (grant no. R-

GNOSIS-F P7-HEALTH-F3-2011-282512). C.U. was supported by a grant from MICINN

(PID2020-120292RB-I00). R.P.M. and C.T.P. received funding from CONACYT Ciencia Basica

(grant A1-S-321649). C.T.P: also received funding from CONACYT Fronteras (grant 2019-

217367).

Bibliography

1.   J. P. Gogarten, J. P. Townsend, Horizontal gene transfer, genome innovation and evolution.
Nat Rev Microbiol 3, 679–687 (2005).
2.   A. San Millan, R. C. MacLean, Fitness Costs of Plasmids: a Limit to Plasmid Transmission.
Microbiol. Spectr. 5 (2017).
3.   D. A. Baltrus, Exploring the costs of horizontal gene transfer. Trends Ecol Evol 28, 489–495
(2013).
4.   K. Z. Coyte, et al., Horizontal gene transfer and ecological interactions jointly control
microbiome stability. PLOS Biol. 20, e3001847 (2022).
5.   M. A. Brockhurst, E. Harrison, Ecological and evolutionary solutions to the plasmid paradox.
Trends Microbiol. (2021) https:/doi.org/10.1016/j.tim.2021.11.001.
6.   M. J. Bottery, Ecological dynamics of plasmid transfer and persistence in microbial
communities. Curr. Opin. Microbiol. 68, 102152 (2022).
7.   F. M. Stewart, B. R. Levin, The Population Biology of Bacterial Plasmids: A PRIORI
Conditions for the Existence of Conjugationally Transmitted Factors. Genetics 87, 209–228
(1977).
8.   A. J. Lopatkin, et al., Persistence and reversal of plasmid-mediated antibiotic resistance. Nat.
Commun. 8, 1689 (2017).
9.   P. E. Turner, V. S. Cooper, R. E. Lenski, TRADEOFF BETWEEN HORIZONTAL AND
VERTICAL MODES OF TRANSMISSION IN BACTERIAL PLASMIDS. Evol. Int. J. Org. Evol.
52, 315–329 (1998).
10.   J. H. Bethke, et al., Vertical and horizontal gene transfer tradeoffs direct plasmid fitness. Mol.
Syst. Biol., e11300 (2022).
11.   F. Benz, A. R. Hall, "Host-specific plasmid evolution explains the variable spread of clinical
antibiotic-resistance plasmids" (Evolutionary Biology, 2022)
https:/doi.org/10.1101/2022.07.06.498992 (January 3, 2023).
12.   Jordt, H., et al. Coevolution of host–plasmid pairs facilitates the emergence of novel
multidrug resistance. Nat. Ecol. Evol. 4, 863–869 (2020)
13.   C. T. Bergstrom, M. Lipsitch, B. R. Levin, Natural selection, infectious transfer and the
existence conditions for bacterial plasmids. Genetics 155, 1505–1519 (2000).
14.   L. N. Lili, N. F. Britton, E. J. Feil, The persistence of parasitic plasmids. Genetics 177, 399–
405 (2007).
15.   J. M. Ponciano, L. De Gelder, E. M. Top, P. Joyce, The population biology of bacterial
plasmids: a hidden Markov model approach. Genetics 176, 957–968 (2007).
16.   J. C. R. Hernández-Beltrán, A. San Millán, A. Fuentes-Hernández, R. Peña-Miller,
Mathematical Models of Plasmid Population Dynamics. Front. Microbiol. 12, 3389 (2021).
17.   A. San Millan, et al., Positive selection and compensatory adaptation interact to stabilize
non-transmissible plasmids. Nat Commun 5, 5208 (2014).
18.   J. P. J. Hall, A. J. Wood, E. Harrison, M. A. Brockhurst, Source–sink plasmid transfer
dynamics maintain gene mobility in soil bacterial communities. Proc. Natl. Acad. Sci. 113,
8260–8265 (2016).
19.   L. Li, et al., Plasmids persist in a microbial community by providing fitness benefit to multiple
phylotypes. ISME J. 14, 1170–1181 (2020).
20.   T. Dimitriu, L. Marchant, A. Buckling, B. Raymond, Bacteria from natural populations transfer
plasmids mostly towards their kin. Proc. R. Soc. B 286, 20191110 (2019).
21.   J. H. Bethke, et al., Environmental and genetic determinants of plasmid mobility in
pathogenic *Escherichia coli*. Science Advances. 6, 4 (2020).
22.   S. R. Partridge, S. M. Kwong, N. Firth, S. O. Jensen, Mobile genetic elements associated
with antimicrobial resistance. Clin. Microbiol. Rev. 31 (2018).
23.   C. J. L. Murray, et al., Global burden of bacterial antimicrobial resistance in 2019: a
systematic analysis. The Lancet (2022) https:/doi.org/10.1016/S0140-6736(21)02724-0.
24.   WHO, List of bacteria for which new antibiotics are urgently needed

https://www.who.int/news/item/27-02-2017-who-publishes-list-of-bacteria-for-which-new-
antibiotics-are-urgently-needed (January 3, 2023).

25. A. Cassini, et al., Attributable deaths and disability-adjusted life-years caused by infections
860 with antibiotic-resistant bacteria in the EU and the European Economic Area in 2015: a
861 population-level modelling analysis. Lancet Infect. Dis. 19, 56–66 (2019).

26. J. D. D. Pitout, G. Peirano, M. M. Kock, K.-A. Strydom, Y. Matsumura, The Global
863 Ascendency of OXA-48-Type Carbapenemases. Clin. Microbiol. Rev. 33, e00102-19 (2019).

27. S. Redondo-Salvo, et al., Pathways for horizontal gene transfer in bacteria revealed by a
865 global map of their plasmids. Nat. Commun. 11, 3602 (2020).

28. R. León-Sampedro, et al., Pervasive transmission of a carbapenem resistance plasmid in the
867 gut microbiota of hospitalized patients. Nat. Microbiol. 6, 606–616 (2021).

29. S. David, et al., Epidemic of carbapenem-resistant Klebsiella pneumoniae in Europe is
869 driven by nosocomial spread. Nat. Microbiol. 4, 1919–1929 (2019).

30. A. Alonso-del Valle, et al., Variability of plasmid fitness effects contributes to plasmid
871 persistence in bacterial communities. Nat. Commun. 12, 2653 (2021).

31. A. San Millan, J. A. Escudero, D. R. Gifford, D. Mazel, R. C. MacLean, Multicopy plasmids
873 potentiate the evolution of antibiotic resistance in bacteria. Nat. Ecol. Evol. 1, 10 (2016).

32. J. DelaFuente, et al., Within-patient evolution of plasmid-mediated antimicrobial resistance.
875 Nat. Ecol. Evol. 6, 1980–1991 (2022).

33. R. Fernandez-Lopez, I. del Campo, C. Revilla, A. Cuevas, F. de la Cruz, Negative Feedback
877 and Transcriptional Overshooting in a Regulatory Network for Horizontal Gene Transfer.
878 PLoS Genet. 10, e1004171 (2014).

34 A. Djukovic, et al., Lactobacillus supports Clostridiales to restrict gut colonization by
880 multidrug- resistant Enterobacteriaceae. (2022). Nat. Commun. 13, 5617 (2022).

35. M. Haudiquet, A. Buffet, O. Rendueles, E. P. C. Rocha, Interplay between the cell envelope
882 and mobile genetic elements shapes gene flow in populations of the nosocomial pathogen
883 Klebsiella pneumoniae. PLoS Biol. 19, e3001276 (2021).

36. C. Brehony, et al., An MLST approach to support tracking of plasmids carrying OXA-48-like
885 carbapenemase. J. Antimicrob. Chemother. 74, 1856–1862 (2019).

37. A. M. Hammerum, et al., Investigation of possible clonal transmission of carbapenemase-
887 producing Klebsiella pneumoniae complex member isolates in Denmark using core genome
888 MLST and National Patient Registry Data. Int. J. Antimicrob. Agents 55, 105931 (2020).

38. H. Arabaghian, et al., Molecular Characterization of Carbapenem Resistant Klebsiella
890 pneumoniae and Klebsiella quasipneumoniae Isolated from Lebanon. Sci. Rep. 9, 531
891 (2019).

39. A. K. Brödel, et al., In situ targeted mutagenesis of gut bacteria. bioRxiv, 2022.09.30.509847
893 (2022).

40. R. López-Igual, J. Bernal-Bayard, A. Rodríguez-Patón, J.-M. Ghigo, D. Mazel, Engineered
895 toxin-intein antimicrobials can selectively target and kill antibiotic-resistant bacteria in mixed
896 populations. Nat. Biotechnol. 37, 755–760 (2019).

41. F. Le Roux, J. Binesse, D. Saulnier, D. Mazel, Construction of a Vibrio splendidus Mutant
898 Lacking the Metalloprotease Gene vsm by Use of a Novel Counterselectable Suicide Vector.
899 Appl. Environ. Microbiol. 73, 777–784 (2007).

42. I. Wiegand, K. Hilpert, R. E. W. Hancock, Agar and broth dilution methods to determine the
901 minimal inhibitory concentration (MIC) of antimicrobial substances. Nat. Protoc. 3, 163–175
902 (2008).

43. M. Agarwal, R. S. Rathore, A. Chauhan, A Rapid and High Throughput MIC Determination
904 Method to Screen Uranium Resistant Microorganisms. Methods Protoc. 3, 21 (2020).

44. M. Carlier, A. G. Verstraete, J. J. De Waele, V. Stove, Stability of amoxicillin and
906 amoxicillin/clavulanic acid reconstituted in isotonic saline. J. Chemother. 29, 54–56 (2017).

45. A. Bankevich, et al., SPAdes: a new genome assembly algorithm and its applications to
908 single-cell sequencing. J. Comput. Biol. J. Comput. Mol. Cell Biol. 19, 455–477 (2012).

46. A. Gurevich, V. Saveliev, N. Vyahhi, G. Tesler, QUAST: quality assessment tool for genome

assemblies. Bioinforma. Oxf. Engl. 29, 1072–1075 (2013).

47. L. S. Katz, et al., Mashtree: a rapid comparison of whole genome sequence files. J. Open
Source Softw. 4, 1762 (2019).

48. H. Wang, et al., Increased plasmid copy number is essential for Yersinia T3SS function and
virulence. Science 353, 492–495 (2016).

49. P. K. Jangir, et al., Pre-existing chromosomal polymorphisms in pathogenic E. coli potentiate
the evolution of resistance to a last-resort antibiotic. eLife 11, e78834 (2022).

50. H. Li, Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM
(2013) https:/doi.org/10.48550/arXiv.1303.3997 (January 18, 2023).

51. P. Danecek, et al., Twelve years of SAMtools and BCFtools. GigaScience 10, giab008
(2021).

52. T. Seemann, Prokka: rapid prokaryotic genome annotation. Bioinformatics 30, 2068–2069
(2014).

53. R. J. Roberts, T. Vincze, J. Posfai, D. Macelis, REBASE—a database for DNA restriction
and modification: enzymes, genes and genomes. Nucleic Acids Res. 38, D234–D236 (2010).

54. S. F. Altschul, W. Gish, W. Miller, E. W. Myers, D. J. Lipman, Basic local alignment search
tool. J. Mol. Biol. 215, 403–410 (1990).

55. P. H. Oliveira, M. Touchon, E. P. C. Rocha, Regulation of genetic flux between bacteria by
restriction–modification systems. Proc. Natl. Acad. Sci. 113, 5658–5663 (2016).

56. S. S. Abby, B. Néron, H. Ménager, M. Touchon, E. P. C. Rocha, MacSyFinder: a program to
mine genomes for molecular systems with an application to CRISPR-Cas systems. PLOS
ONE 9, e110726 (2014).

57. F. Tesson, et al., Systematic and quantitative view of the antiviral arsenal of prokaryotes.
Nat. Commun. 13, 2561 (2022).

58. L. J. Payne, et al., Identification and classification of antiviral defence systems in bacteria
and archaea with PADLOC reveals new system types. Nucleic Acids Res. 49, 10868–10878
(2021).

59. S. S. Abby, et al., Identification of protein secretion systems in bacterial genomes. Sci. Rep.
6, 23080 (2016).

60. M. M. C. Lam, R. R. Wick, L. M. Judd, K. E. Holt, K. L. Wyres, Kaptive 2.0: updated capsule
and lipopolysaccharide locus typing for the Klebsiella pneumoniae species complex. Microb.
Genomics 8, 000800 (2022).

61. O. Rendueles, M. Garcia-Garcerà, B. Néron, M. Touchon, E. P. C. Rocha, Abundance and
co-occurrence of extracellular capsules increase environmental breadth: Implications for the
emergence of pathogens. PLOS Pathog. 13, e1006525 (2017).

62. J. A. M. de Sousa, A. Buffet, M. Haudiquet, E. P. C. Rocha, O. Rendueles, Modular
prophage interactions driven by capsule serotype select for capsule loss under phage
predation. ISME J. 14, 2980–2996 (2020).

63. M. Steinegger, J. Söding, MMseqs2 enables sensitive protein sequence searching for the
analysis of massive data sets. Nat. Biotechnol. 35, 1026–1028 (2017).

64. D. H. Haft, et al., RefSeq: an update on prokaryotic genome annotation and curation. Nucleic
Acids Res. 46, D851–D860 (2018).

65. C. Rinke, et al., Insights into the phylogeny and coding potential of microbial dark matter.
Nature 499, 431–437 (2013).

66. C. d'Humières, et al., A simple, reproducible and cost-effective procedure to analyse gut
phageome: from phage isolation to bioinformatic approach. Sci. Rep. 9, 11331 (2019).

67. S. R. Eddy, Accelerated Profile HMM Searches. PLOS Comput. Biol. 7, e1002195 (2011).

68. K. Katoh, D. M. Standley, MAFFT Multiple Sequence Alignment Software Version 7:
Improvements in Performance and Usability. Mol. Biol. Evol. 30, 772–780 (2013).

69. S. Capella-Gutiérrez, J. M. Silla-Martínez, T. Gabaldón, trimAl: a tool for automated
alignment trimming in large-scale phylogenetic analyses. Bioinforma. Oxf. Engl. 25, 1972–
1973 (2009).

70. L.-T. Nguyen, H. A. Schmidt, A. von Haeseler, B. Q. Minh, IQ-TREE: A Fast and Effective

Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. Mol. Biol. Evol. 32,
268–274 (2015).

71. L. si Tung Ho, C. Ané, A linear-time algorithm for Gaussian and non-Gaussian trait evolution
models. Syst. Biol. 63, 397–408 (2014).

72. S. R. Lele, B. Dennis, F. Lutscher, Data cloning: easy maximum likelihood estimation for
complex ecological models using Bayesian Markov chain Monte Carlo methods. Ecol Lett
10, 551–563 (2007).

970
971

972   **Supplementary Figures**

974
Fig. S1. Unrooted phylogeny of *E. coli* (left, n=25) and *Klebsiella* spp. (right, n=25) plasmid-free

(PF) strains. Branch lengths represent mash distances between the whole-genome assemblies

of the strains. Groups of strains belonging to the same multilocus sequence type (ST) are shaded

(note that *E. coli* ST6217 is part of the ST10 group). Filled circles indicate strains for which

transconjugants (TC) with an isogenic pOXA-48 could be obtained and were therefore used to

determine minimum inhibitory concentrations (MIC) to antibiotics and pOXA-48 copy number (*E.*

coli n=15, *Klebsiella* spp. n=18). Empty circles indicate that TC could be obtained but harbored a

mutated pOXA-48 plasmid. PF strains used as recipients in conjugation experiments are marked

with a star.

984
985
Fig. S2. Antimicrobial resistance (AMR) genes encoded by the plasmid-free (PF) strains used to

determine minimum inhibitory concentrations (MICs) to antibiotics. The heatmap shows presence

(colored by percentage of identity to the BLAST hit in the ResFinder database; see Methods) and

absence (values of 0) of the AMR genes. In each of the four panels, the PF strains are ordered

by fold change (FC) in MIC to the corresponding antibiotic when acquiring pOXA-48 (top left,

amoxicillin/clavulanic acid, AMC; top right, ertapenem, ERT; bottom left, imipenem, IMP; bottom

right, meropenem, MER).

993
994

995   **Fig. S3.** Spearman's correlation between pOXA-48 copy number and the minimum inhibitory

996   concentration (MIC) of amoxicillin/clavulanic acid (AMC) in the pOXA-48-carrying clones. pOXA-

997   48 copy number was estimated from sequencing data as the ratio of plasmid/chromosome median

998   coverage (see Methods).

Fig. S4. Analysis of genomic traits in recipient strains that could explain differences in pOXA-48 acquisition by conjugation. (A) Heatmap of presence/absence of restriction-modification (RM) systems. In the top panel, all RM systems subtypes, labeled with different letters, are shown. In the bottom panel, RM systems are collapsed by type. (B) Heatmap of presence/absence of other phage defense systems. (C) Heatmap of presence/absence of plasmid replicons. The intensity of the color scale in A bottom and B indicate the number of systems of a specific type a strain is encoding, while in C indicates the percentage of identity of the replicon to the BLAST hit of the PlasmidFinder database (see Methods). In A-C, recipient strains are ordered by mean pOXA-48 conjugation rates across replicates and experimental conditions. Strains belonging to the same sequence type (ST) are marked with a symbol.

1013

1014 **Fig. S5.** Stochastic environments used in numerical experiments. (A) Each row depicts a unique

1015 environmental condition that changes daily, with the drug concentration based on a Gaussian

1016 probability density function. Drug concentration is indicated using shades of gray. (B) Each row

1017 displays a stochastic environmental condition with increasing levels of selection strength

1018 $(0 \leq A_{max} \leq 1)$.

1019

1020
1021
Fig. S6. Pairwise competition experiments between *E. coli* and *Klebsiella* spp. strains. (A) In a

drug-free environment ($A_{max} = 0$). The diameter of each circle is proportional to the total bacterial

density observed at the end of the experiment. Plasmid-free and plasmid-bearing subpopulations

are denoted with light and dark colors, respectively (*Klebsiella* spp. in blue and *E. coli* in red). (B)

In the presence of antibiotics ($A_{max} = 0.1$), plasmid-bearing *Klebsiella* spp. strains outcompete

most *E. coli* strains. (C) At high drug concentrations ($A_{max} = 0.5$), only highly resistant *Klebsiella*

spp. strains are present at the end of the experiment.

1029
1030

1031
1032
Fig. S7. Percentage of pairwise *E. coli* / *Klebsiella* spp. associations where the plasmid remains

stable over time. The red bars indicate the cases where the plasmid was present in the *E. coli*

strains, and the blue bars indicate those cases where the plasmid was maintained in the *Klebsiella*

spp. strain. Note that the presence of plasmids is highest at intermediate strengths of selection,

which corresponds to drug concentrations that select for drug-resistant *Klebsiella* spp. and *E. coli*

strains.

1039
1040

Fig. S8. Example numerical simulations for plasmid invasion in a community composed of six

strains. (A) In a drug-free environment ($A_{max} = 0$), the plasmid-bearing population is outcompeted

by plasmid-free cells and thus the plasmid is cleared from the population. Top plot illustrates the

bacterial density at the end of each day, with dashed lines representing plasmid-free cells and

solid lines representing plasmid-bearing cells. Bottom plot illustrates the density (in log scale) of

each strain in the community. (B) The plasmid is stably maintained in the population in the

presence of positive selection ($A_{max} = 0.1$). The drug concentration used each day is represented

in the ribbon between the panels (drug concentration is shown in tones of gray). (C) Density of

different strains in a stochastic environment with high levels of selection ($A_{max} = 0.5$).

1050

1051
1052
Fig. S9. Strain-level results obtained after performing 1,000 simulations of a plasmid invasion

experiment in a community composed of six individuals. Each column represents a different

bacterial strain (*Klebsiella* spp. in blue and *E. coli* in red). In light colors, the number of times each

strain was randomly selected for the stochastic simulations. The number of experiments where

the corresponding plasmid-bearing strain was detected at the end of the numerical experiment is

represented with a dark bar. Strains are ordered based on the frequency of plasmid-bearing in

decreasing order. (A) In drug-free environments ($A_{max} = 0$), plasmid-bearing strains can be found

both in *E. coli* and *Klebsiella* spp. strains. (B) In the presence of antibiotics ($A_{max} = 0.1$), most

plasmid-bearing strains are *Klebsiella* spp. strains. (C) At high drug concentrations, $A_{max} = 0.5$,

all *E. coli* strains are driven to extinction and the plasmid is only present in *Klebsiella* spp. strains.