1

Mast fruiting in a large tropical African legume tree provides

evidence for the nutrient resource limitation hypothesis

David M. Newbery[1*], Sarah Schwan[1], George B. Chuyong[2], Godlove A. Neba[2],

Culbertson E. Etta[3], Julian M. Norghauer[1,4], and Martin Worbes[5]

[1] Section for Vegetation Ecology, Institute of Plant Sciences, University of Bern,

Altenbergrain 21, CH-3013 Bern, Switzerland

[2] Department of Plant Science, University of Buea, POB 63, Buea, South-West Region,

Cameroon

[3] Research and Development Section, PAMOL Plantations Plc, Lobe Oil Palm Estate, PMB

3, Ekondo Titi, South-West Region, Cameroon.

[4] Present address: Statistical Scientific Editing, Montréal, Québec, Canada

5 Tropical Plant Production and Agricultural Systems Modelling, Georg-August-University,

Grisebachstraße 6, DE-37077 Göttingen, Germany

* Corresponding author david.newbery@ips.unibe.ch

19   Abstract

Background and aims – The large grove-forming tropical tree *Microberlinia bisulcata*

(Fabaceae subfamily Detarioideae) in southern Korup, SW Cameroon, displays strong mast

fruiting. The soil is poor in phosphorus and potassium: the climate is strongly seasonal.

Evidence is presented that masting is probably determined primarily by mineral nutrients and

secondarily by radiation.

Method and materials – Field data on phenology and climate variation (1989 to 2017), and

soil moisture, nutrients in litter and fruits, insect herbivory and tree growth are integrated.

Some data come from previous studies at Korup, and they are combined with new data to

enable an extended analysis.

Key results – To test the nutrient resource limitation hypothesis, phenological recordings

were matched with climate variables and modelled using logistic and time-series regressions.

A strong predictor was average daily rainfall in the dry season, low in the current year of

masting and high in the year prior. Less strong was the increase in dry season radiation

between prior and mast years. Masting events showed no relationship to annual stem

increment. Masting was reduced after two caterpillar attacks. Fallen-leaf phosphorus

concentration increased in the inter-mast interval and that for potassium peaked mid-interval.

These elements were likely accumulating and being stored, triggering masting when internal

thresholds were crossed. A drier dry season prior to masting may have increased tree carbon,

the wetter one increased nutrient acquisition and uptake via roots and mycorrhizas. A

rooting–fruiting trade-off in carbon is proposed. Storage of phosphorus may be mostly in

bark of branches; that for potassium on soil organic colloids.

Conclusion – The long-term driver appears to be stochastic dry-season rainfall. Synchrony

across masting trees might be enhanced by equilibration of nutrients across the mycorrhizal-

root network. Life history strategy determined by mineral nutrient resource dynamics

provides the best explanation and an advanced model for the observed pattern of masting

events.

<u>Keywords</u> Detarioideae, dry season intensity, ectomycorrhizas, lowland rain forest, mast

fruiting, phosphorus limitation, times-series analysis.

INTRODUCTION

By fruiting and seeding intensively at intervals as opposed to continuously over the years,

seed and seedling survival may be enhanced for plants species with a masting iteroparous

type of life-history as an evolutionary stable strategy (Maynard Smith 1972; Charlesworth

1980; Stearns 1992). Trees provide by far the majority of the documented cases. To explain

how a species' fitness can be maintained or increased, a so-called 'economy of scale' concept

has often been invoked (Norton and Kelly 1988). Three general processes have been

proposed to achieve this type of economy, usually thought of, teleonomically, as the ultimate

causes of mast fruiting and seeding: (1) the efficacy of pollination may be increased by mass

flowering, (2) seed predators may be more readily satiated by mass seeding, and (3) the

chances of successful dispersal and establishment may be improved by larger reproductive

output (Kelly 1994; Kelly and Sork 2002; Crone and Rapp 2014; Pearse et al. 2016). Much of

the reported research into masting, however, has been in the ecological and evolutionary

direction with a strong temperate-biome bias (see reviews aforementioned). Surprisingly,

there have been comparatively few studies on masting in the tropics (e.g., Henkel et al. 2005;

Ichie et al. 2005a; Newbery et al. 2006a; Norden et al. 2007).

A much more critical issue for understanding masting in trees is the present lack of

evidence for, and hypotheses about, the physiological mechanisms which *enable* mast

fruiting and seeding, for both temperate and tropical ecosystems. These mechanisms are

commonly viewed by ecologists as being proximate causes. Recent discussions about

whether a strict distinction exists between ultimate and proximate causes (Laland et al. 2011;

Corning 2019) leave it more open that factors, such as nutrient supply, directly affecting

physiological processes might also qualify as ultimate causes of mast fruiting (Fernandez-

Martinez et al. 2019; Fernandez-Martinez et al. 2020), although it has been contested whether

they can constitute an 'economy of scale' (Kelly 2020).

For flower, fruit and seed production to be allocated efficiently over the life-time of

the adult tree as masting years, a set of physiological resource-mechanisms, probably in serial

form, must be in place (Lloyd 1980). Resource budget models, initiated by Isagi et al. (1997),

have simulated how a resource, usually carbon (C), accumulates in storage to an internal

plant threshold which then triggers flowering. Almost all such models have involved the

coupled wind pollination of neighbouring individuals (Yamauchi 1996; Satake and Bjornstad

2008; Abe et al. 2016; Venner et al. 2016; Ye and Sakai 2016; Vacchiano et al. 2018). They

have often involved the Moran effect whereby masting is synchronized over large distances

by driving climate factors. The triggering is thought to be brought about by a resource level

being crossed in the plant which affects plant hormone concentrations that then cue a leafing

axil to change to a flowering one, and finally enable fruit maturation (see e.g., Marshall and

Grace 1992). In this respect, an individual tree can be modeled as an integrated and dynamic

system which controls the timing and intensity of reproductive allocation to masting. Details

of the involved physiological mechanisms can be expected to differ across tree species and

sites, each achieving a stable yet flexible masting strategy in a different way.

The evidence for C as the main resource directly involved in masting is, however,

quite weak. This conclusion comes from direct measurements on whole trees and

observations of allocation and experimental manipulation of branches; and from overall C

budgets and studies on the dynamics of non-structural carbohydrates (NSCs) (Hoch et al. 2003; Hoch 2005; Wurth et al. 2005; Millard et al. 2007; Mund et al. 2010; Bustan et al. 2011; Sanchez-Humanes et al. 2011; Ichie et al. 2013; Miyazaki 2013; Richardson et al. 2013; Igarashi et al. 2019; Mund et al. 2020). A few studies have suggested that mineral nutrients may be the key resources underlying budget mechanisms, but the evidence remains scarce because the topic has been so little researched (Bogdziewicz et al. 2024). So far, Han et al. (2014) and Han and Kabeya (2017) have shown that nitrogen (N) storage and mobilization is important for triggering flowering in *Fagus*. Fernandez-Martinez et al. (2017b) and Le Roncé et al. (2023) have indicated that phosphorus (P) and zinc (Zn), and N and Zn, respectively are likely involved in mast fruiting of temperate trees more widely. Carbon should not be laid aside completely as a co-resource though. It is needed with N, P, potassium (K), and other macronutrients, not just to build fruits and seeds, but also indirectly to support growth and maintenance of organs such as fine roots and mycorrhizas which are essential for nutrient acquisition.

If macronutrients do underpin the mechanisms of mast fruiting, it could be expected that the element most important in controlling reproduction might be the one which is at the lowest level of availability at the site relative to the others, that is the one that is in most critical supply overall for the trees (Allen et al. 2017; Fernandez-Martinez et al. 2019). This element will need to be accumulated to support successful reproduction, because seed filling depends on minimally attaining the necessary stoichiometric ratios for all elements required and doing so without undue costs to other tree growth and maintenance demands in the short to medium term. Further, extra-stoichiometric requirements may need to be met where the same element's availability is very low in the soil: a rise in elemental concentration in the seeds would likely be advantageous for seedling establishment.

In this way natural selection would be expected to evolve a cueing-threshold-

triggering system using the critical nutrient in the form of a cell storage product. Other

elements in not quite so low supply as the first will also be important, which leads to the idea

of nested systems of stores and thresholds. The search, then, is for those key limiting

elements that most determine tree growth rates and nutrient cycling, which can also be shown

to determine frequency and intensity of reproduction. Whilst the elements involved are

probably different for different species, habitats and ecosystems, in principle, the same

fundamental process overall can be postulated to operate.

Artificial selection of trees for fruiting under controlled nursery conditions has

provided a major insight into tree physiology with regard to fruiting and what may lie as the

basis to masting (Davis 1957; Monselise and Goldschmidt 1982; Crawley and Long 1995;

Garcia et al. 2021). Cultivated trees often follow an 'alternating bearing' fruiting schedule,

with years of 'on' and 'off' harvests over a regular 2-year period. Several temperate tree

species in the Fagaceae show such an alternating pattern in natural forest, which leads to the

idea that, theoretically, the minimal-intrinsic model across all masting species is perhaps

alternating years of reproduction (Newbery et al. 2006a), but in naturally non-fertilized

situations the interval between fruiting becomes extended beyond 1 year to allow time to

accumulate critical resources and thus allow masting to effect maximize fitness. ('Interval' is

here defined as the years of non-masting between masting events.) The fundamental question

then becomes, how is the tree system coordinated to synchronize mast fruiting? Nutrient

cycles have been strongly implicated in several fruit tree studies (e.g. Rosecrance et al. 1996,

1998).

The main hypothesis of the present paper is that limiting mineral nutrients can

determine mast fruiting at the mechanistic, tree-physiological and proximate/ultimate level,

particularly for those species that are ectomycorrhizal. This limitation is expected to be

strongest where soil mineral nutrients at a site are in critically low supply and

ectomycorrhizas are essential to nutrient uptake. Both the tropical Fabaceae subfamily

Detarioideae and the Dipterocarpaceae are often ectomycorrhizal (Alexander 1989a;

Alexander 1989b; Brearley 2012) and both commonly do show mast fruiting. Among the

temperate families (Fagaceae and Pinaceae especially) there is also a strong association

between masting and having this symbiotic mode of nutrient uptake (Newbery 2005). Apart

from the review of Corrales et al. (2018), the potential role of ectomycorrhizas in masting

appears to have largely gone uncommented (e.g. Pearse et al. 2016).

149   Two associated processes could potentially underpin how mast fruiting might be

controlled by mineral nutrient limitation. They are not fully exclusive to one another. (i)

Ectomycorrhizas might achieve a form of economy of scale due to their putatively extensive

shared hyphal networks between the roots of large adult trees (Simard et al. 1997), possibly

also via root grafts in dense stands (Graham and Bormann 1966), which could equilibrate

levels of the limiting nutrient resource (Newbery et al. 1998), and thereby lead to trees

reaching similar internal thresholds simultaneously. (ii) Year-to-year stochasticity in climate

(Kerkhoff 2004; Crone and Rapp 2014), especially seasonally-related rainfall, might cause

soil nutrient availability and uptake to increase in the interval years between masting events,

thus enabling stores to increase across all trees synchronously. For that no economy of scale

per se is necessary. Furthermore, these two processes might interact positively, entraining

common years of masting because they both depend on ectomycorrhizas and rooting activity.

161   An ecosystem well suited for testing the nutrient resource limitation hypothesis would

be high-stature primary forest, for which C is in abundant supply, on soils very low in one or

two elements that are macronutrients. Such an opportunity was provided by our long-term

work at Korup National Park (SW Cameroon) since the 1970's. In testing the main mineral

nutrient hypothesis, it is also necessary to simultaneously test the most likely alternative,

which is that C limitation, driven by radiation levels, determines mast fruiting. This latter aim

is supported by an analysis of fruit production in a nearby oil-palm plantation, and a new

analysis of previously collected tree ring growth data in relation to masting. Supporting

knowledge of tree growth, forest dynamics, nutrient cycling, soils and climate variation at

Korup are laid out in the following Background section. This information is essential for fully

understanding the framework of the hypotheses being tested and the importance of the two

associated processes.

The present paper builds on the study of mast fruiting in *Microberlinia bisulcata* A.

Chev. between 1989 and 2004 (Newbery et al. 2006a) and extends it to 2017 (29 years). For

the parts concerning nutrients, some previously published values (as cited) are brought in

with the newly reported ones to allow a complete five-element masting interval analysis. In

Methods and Materials, and in Results, old and new values are clearly distinguished from one

another. In the Discussion section, a synthesis of the new updated results, an improved main

hypothesis and schematic model are developed and integrated with a review of the literature.

These form a strong basis for new predictions, tests and analyses of mast fruiting in the

African detarioids.

183   BACKGROUND

Forest ecosystem

Korup National Park, in the South-West Region of Cameroon, is covered by primary

rain forest of the Atlantic Coastal type (Letouzey 1968), and lies within the Guinea-

Congolean refugium (Gartlan 1992). Local and regional maps are found in Gartlan et al.

(1986) and Newbery et al. (1988): further details of the site are given in Newbery and Gartlan

(1996), Newbery et al. (2006a) and Newbery et al. (2013).

Microberlinia bisulcata dominates large patches or 'groves', 1–3 km across, in the

primary rain forest of southern Korup. Trees $\geq$ 50 cm stem diameter composed 17.8% of the

overall density and 32.6% of total basal area in 2005, and together with co-dominants

Tetraberlinia bisulcata and *T. korupensis* correspondingly 46.5% and 61.3% (Newbery et al.

2004; Newbery et al. 2013). All three species are strongly ectomycorrhizal as are several

other co-occurring tree species of the subfamily Detarioideae in the Fabaceae, (formerly

Caesalpiniaceae-Leguminosae, Newbery et al. 1988). The taxonomy follows the revision of

LPWG (2017) and de la Estrella et al. (2018). They are not N-fixing trees. *Microberlinia*

bisulcata trees attain considerable stature in the main canopy, characterized by wide dome-

shaped crowns, and very extensive laterally spreading buttress systems (Newbery et al. 2009;

Newbery et al. 2013). The species is a leaf-exchanger and therefore semi-deciduous: new

leaves at the start of the dry season (December to February) push off the old leaves, flowering

follows within a month (Newbery et al. 1998; Newbery et al. 2006a). Flowers are outcrossed

mainly by small bees and pollen seems unlikely to be limiting: bees are highly abundant in

the canopy during the dry season (Newbery et al. 1998). It remains untested, however,

whether insect pollination could form an economy-of-scale factor: in this connection

presumably the density of trees plays a crucial role. Into the wet season (March to October)

pods mature from soft green to hard fibrous brown structures. Seeds are dispersed

ballistically in August-September (Norghauer and Newbery 2015).

The Korup-Ndian area along with Douala at the coast is climatologically special for

western Central Africa because it has one clear dry season each year as opposed to the two

less-distinct ones found elsewhere in the region. Mean annual rainfall for 1984–2016 was

5159 mm (range 4023 to 6531 mm). The typical seasonal course in rainfall is shown in

Newbery et al. (Newbery et al. 1997), with almost none in January and February but reaching

900–1000 mm in August or September. Site elevation of the site in southern Korup National

Park (KNP) is ~120 m a.s.l. (range of 40 to 200 m from south to north inland), slightly

varying between swampy stream-fed lower areas up to local slopes, in places strewn with

quartzite rocks. The soils are very sandy and free-draining, acidic and very low in available P

and K (Gartlan et al. 1986), meaning that here high-stature tropical forest is growing on one

of the most nutrient-poor sites in tropical Africa. Within groves at Korup, soil organic matter

is largely confined to the top 2 cm of the profile, forming a surface mat that is densely

occupied by fine roots and ectomycorrhizal hyphae (Newbery et al. 1988; Newbery et al.

1997). Larger vertical roots can tap deeper water reserves. Nutrient cycling in the groves

shows a 'fast-forward' modus: leaf litter decay is fast, there is no permanent leaf litter layer,

and most cycling via the very efficient mat happens in the early wet season (Chuyong et al.

2000, 2002). At this time throughfall nutrients help to prime decomposition and uptake

(Chuyong et al. 2004). For the rest of the year, heavy rains leach the forest, and this results in

a tight closed ecosystem in which ectomycorrhizas play a central role in P acquisition and

uptake (Newbery et al. 1997). In addition, during each year's dry season, Harmattan dust

brings some P to the forest via aerial deposition. A critical field experiment showed that

growth and recruitment of *M. bisulcata* is not P-limited (Newbery et al. 2002), possibly

because of its highly evolved adaptations to the low P site or another element like K was

limiting.

Recruitment of *M. bisulcata* over at least the last six decades has been poor due to

very low seedling and sapling survival (Newbery et al. 1998; Newbery et al. 2000; Newbery

et al. 2006b; Newbery et al. 2010; Norghauer and Newbery 2010). Seed production in mast

years is copious (Newbery et al. 2006a) but this species' seedlings are shade-intolerant

(Green and Newbery 2001a, 2001b). The groves demonstrate transient dominance: the stands

now are predominantly made up of large to very large trees, with extremely few small ones

present (Newbery et al. 2013). The dynamics is thought to be one of patches arising and then

decaying, being replaced, *in situ* or *ex situ*, within a mosaic of species-richer forest (Newbery

and Gartlan 1996; Newbery et al. 2013). The species is classified as a long-lived light-

demanding tree (see Newbery et al. 2010). Part of the expected life-history strategy of long-

lived and light responsive species such as *M. bisulcata* would be to delay size at the onset of

maturity (SOM) in order that greater seed output later offsets the relatively low probability of

persistence in the forest understorey. Two further coevolved traits are the ectomycorrhizal

habit mentioned and mast fruiting. Ectomycorrhizas not only provide advantages in resource

acquisition for juvenile and young adult trees but are also thought to be a means to store

resources which enable mast fruiting in the older adult ones (Newbery et al. 1998; Newbery

et al. 2006a). A key sub-hypothesis has been that these links between adults synchronize the

timing of masts by equilibrating nutrient sinks across groves (Newbery et al. 1997; Newbery

et al. 1998). There is mounting evidence from other forests that ectomycorrhizas build

intricate below-ground hyphal networks between individual trees (e.g. Simard and Durall

2004; Simard et al. 2012), although the definite evidence for these networks providing a

means of C or mineral nutrient transfer is inconsistent, at least for adults to seedlings (Karst

et al. 2023). However, in the context of mast fruiting of adults, the hypothesized transfers are

between very large trees occurring in dense stands (the groves at Korup).

Mast fruiting

Mast fruiting in *M. bisulcata* was first realized from early phenological recording in

1988 to 1995, with prominent seed crops in 1989, 1992 and 1995 yet barely any seed fall in

the intervening years (Newbery et al. 1998). Flowering occurs to some extent in most years (a

few fail completely); though not always does a heavy seed crop depend on heavy flowering

(Newbery et al. 2006a). The two *Tetraberlinia* species tend to mast in the same years as *M.*

bisulcata but not both together (Newbery et al. 1998, and unpublished data). After a

propitious start with a potential 3-year cycle for masting (i.e. a 2-year interval between mast

years), this cycle turned out to be shorter between 1999 and 2005. Because masting was

associated with peaks in mean daily radiation in the dry season prior to fruiting this suggested

that C was at a premium, with some threshold needed to be crossed before masting occurred.

Detailed nutrient cycling studies in plots dominated by *M. bisulcata* showed, for the first

time, that leaf litter falling after masting in 1989 was low in P concentration compared with

the dry season of the years before and after (1988, 1990) , and this matched a decline in

available P in the soil, especially in the surface organic layer (Newbery et al. 1997). This last

result suggested that P was being shunted into the seed crop and less was remaining to be

shed in litter and, given the low soil P at the site and the presence of ectomycorrhizas, P was

also likely limiting mast fruiting. Tree C and P levels might be operating a double-threshold

control system and this led to the phosphorus and climate proposed ectomycorrhizal response

(PACER) hypothesis (Newbery et al. 1997).

The climate at Korup varies stochastically from year to year, particularly regarding

the start, duration and intensity of the dry season. The dates defining the actual season each

year are based on 30-day running rainfall total (*rft*) falling < 100 mm (Newbery et al. 2006a).

This season plus the first 6 weeks of the wet season very largely determine tree physiological

processes for the whole year. Fine-tuned sensing of the environment to optimize resource

gain and allocation within a life history strategy is central to survival. Under- or over-

allocation at the wrong time could lead to large fitness losses, hence the presumed operation

of thresholds that safely trigger phenological change. That C and P might be dual constraints

within a broader nutrient-limitation hypothesis as an explanation of mast fruiting for many

tree species requires a detailed budget to determine how much C and P (also N and K) are

allocated to fruits and seeds in a main masting event. Censuses of seed and pod counts, along

with their masses and nutrient amounts, were obtained for the 1995 masting (Green and

Newbery 2002), showing that *M. bisulcata* produced on average 26K seeds per tree or 83.5K

seeds/ha. Seeds are small (0.64 g each), and with an average of 2.0 seeds per pod, total pod

mass was 25 g (see images in the Suppl. material of Norghauer and Newbery (2015); for the

2007 and 2010 events, 2.5 seeds per pod were estimated. Masting resulted in an investment of

1034 kg /ha dry mass that year, and compared with leaf litter production in previous non-mast

years (Chuyong et al. 2000), was 52% of it by comparison, or in a mast year 74%, i.e. a

substantial allocation of resources (Green and Newbery 2002). Seeds and pods together had

N and P amounts 13 and 21% of that in annual leaf litter production, indicating an important

use of P during masting. Unfortunately, C, P, N and K investments in flowers are unknown

but if included they would increase the reproductive allocation estimates slightly. Flowers of

Microberlinia are very small in mass compared with seeds and fruits. Masting in *M. bisulcata*

is highly synchronous, and it is unusual for a spatially isolated tree to fruit outside of an event

(Newbery et al. 1998; Newbery et al. 2006a).

Using a detailed 150-tree phenological study in the main P-plot at Korup of 1995–

2000, and incorporating circumstantial reliable observations in the field from previous and

later years up to 2004, enabled a better picture of how dry season variability might control

mast fruiting in *M. bisulcata* (Newbery et al. 2006a). From daily rainfall and radiation data

(recorded locally at Bulu, Ndian; 1984–2017), a clearer model was developed. Masting was

tending to occur when the prior dry season was *drier* than average (more intense, low mean

daily rainfall), plus the dry season in the year prior was *wetter* than average (less intense, high

mean daily rainfall). After 1993, radiation did not show the same trend as in 1989–1992,

mainly because after 1995 it climbed steeply as part of an inferred 11-year solar cycle

(Newbery et al. 2006a). The relative change in radiation between successive years was

important, and not the absolute levels. The notion that time series of relevant environmental

variables at Korup were or ever had been stationary is hard to sustain.

The argument developed was that the current year's dry season dryness reflected also

to some degree increased radiation, but the wetter dry season the year prior was possibly

indicating better conditions promoting tree nutrient uptake at the most effective time. The

detailed phenology data showed that masting was strongest on trees on upper slopes (i.e. sites

with relatively drier surface soils in years with a 'dry' dry season), and that in the moderately

'wet' dry season of 2000, those trees that masted were also on the drier sites. This tentatively

pointed to surface soil moisture in the dry season being a key determinant to what was

happening. The intensity of the dry season is unlikely to have particularly affected the water

balance of adult trees themselves because they would have had access to deeper water

sources: after all, trees flushed, leaves and branches grew, and flowering took place, in the

dry season. Taken together, the preliminary evidence pointed to processes in the organic mat

regarding fine roots and ectomycorrhizas, coupled to nutrient transfers within the trees and

ectomycorrhizal network (Newbery 2005; Newbery et al. 2006a). With the masting time

series extended to 29 years –almost twice its previous length, plus information collated from

several nutrient studies at the site, further tree phenology censuses, a more detailed statistical

modelling of fruiting on rainfall and radiation, and the effects of an unexpected herbivore

attack – this paper assesses the extent to which C, P and K, might be determining the masting

cycle in *M. bisulcata* and, in extension, presents a more refined hypothesis and model of the

phenomenon.

334   METHODS AND MATERIALS

Climate records

Climate data continued to be available from the PAMOL Bulu Station, Ndian (Etta et

al. 2022). The station lies 12 km to the SE of the study site. Daily radiation, rainfall and

temperature were recorded since 1 Jan. 1988 until 27 May 2017. Radiation was estimated

using a Gunn-Bellani (BG) radiometer, with volume evaporated per day (V, ml) convertible

into radiation (R, W m^{-2}). Two periods, however, lacked radiation data because of instrument

failure in March –May 1993 and April 2009–May 2011. Improved new radiometer

calibration equations were developed using several shorter series of pyrometer measurements

and these applied to the complete time series to obtain mean monthly radiation (Suppl.

material 1). The short break in radiometer readings had been previously linearly interpolated,

but the radiometer data for the much longer break 2009–2011 required more refined

statistical methods. These are outlined in the analysis in Suppl. material 1. Since radiometers

are rarely still in use today, this step in methodology is original. It also meant that the

standard BG instrument calibration curve (used in Newbery et al. 2006a) could be replaced

with the now much better site-specific parameters. The equations were applied to the whole

series (1989–2017), with differences to the first half of the time series referenced in Suppl.

material 1. World Climate data bases unfortunately proved quite inadequate for estimating

these missing values due to the sparseness of regional stations in western Central Africa and

the special topography around Mt. Cameroon (SSE of Korup), with the few data records

available for the last decades being incomplete (see Suppl. material 1).

A dry season occurred when the 30-day running rainfall total (*rft*) at Ndian (Korup)

was < 100 mm, and this gave a defined start date and duration (the 'drought-defined dry

season' of Newbery et al. (2006a). This dry season typically moved one month back or

forward around the modal months of December to February, coming earlier or later in

different years, and often being shorter or longer. Hence, in what follows, 'dry season 1991'

for example means December 1990 to February 1991 (modally). For a more general, less

strict, annual matching of phenology to climate, a 12-month year starting on 1 December one

year and ending on 30 November the next was adopted, with quarter 1 being December to

February, and so on. Part-years were taken as November to March (5 months, dry) versus

April to October (7 months, wet) to encompass the weeks before and after the dry season

proper. Since its use in Newbery et al. (2006a), the definition of drought dry season was

slightly relaxed here, in that within the outer defining limits a very few days where *rft* was at

< 103 mm were included as drought days provided that the total returned to below 100 mm

for the rest of the season. This affected four of the 16 years slightly in their dry season

durations and mean daily rainfall values (Table 1).

Phenological recording

The main 'P-plot' is located in the southern part of the Park (5°10' N, 8°70' E; global

coordinates 0476438/ 0553538) at an elevation of 96 m a.s.l.: refer to Newbery et al. (2004)

and Newbery et al. (2013) for further details. Mast fruiting from 1989 to 2004 was reported

and analysed in Newbery et al. (2006a). Recordings were then continued from 2005 to 2017

for a subset of trees, and it is this full time series of 29 yr that is presented, with an extended

analysis, in the paper here. Following pod maturity to completion was not possible in 2017

because of security restrictions preventing further access to KNP. The situation remains

unresolved as of early 2025.

Evidence for masting rested on a combination of intensive phenological scoring of

marked trees in two periods, quantitative trap-based estimates of seeding in two specific

studies, from which masting fruiting is inferred; circumstantial records linked to seedling

demography and seedling experiments in several years, and general yet objective assessments

of the whole forest in other years during the course of large-plot census work, dendroband

recording times, and seedling herbivore exclusion and nutrient lysimeter studies. (Suppl.

material 2, table 1). Barely a year passed without adequate observation and reporting because

of the continuing project workflow and frequent field presence of the Stirling-Bern research

group since 1988.

The two periods of intensive recording were achieved in the following manner. To

increase the frequency of recording per annum between 1995 and 2000, 150 *M. bisulcata*

trees in the main 82.5 ha 'P-plot' (out of 294 in total with stem diameter ≥ 50 cm; Newbery et

al. 2013) had been scored on 61 occasions (mostly at monthly intervals), for intensity of leaf

fall, leaf flush, flowering and immature and mature pods: scale 0, none; 1, ~25; 2, ~50; and 3,

~75 and up to 100% (Newbery et al. 2006a). Later, within an 8-ha sub-area of the 82.5-ha P-

plot (1000–1200 m E, and 100–500 N, with respect to SW origin; see Fig. 2a of Newbery et

al. (2013), 61 of the 65 trees surviving since the last enumeration in 2005 (stem diameter,

mean ± SE = 112 ± 4 cm, range = 54–174 cm), were scored between 2009 and 2013 in a

similar manner as the first period. In 2014, fallen pod density on the ground was taken as a

proxy for mature pod density (Suppl. material 2, table 1).

Masting was categorized as 'high' (labelled 'M') when all, or almost all (> ~85%), of

the recorded trees had many pods (scale = c. 3), and 'moderate' ('m') when fewer trees (~45–

85%) had many pods, or numbers of pods on relatively more trees were significantly lower.

These masting events were clearly differentiated from those with 'no' or 'very low' pod

production, and from those 'low' ones with just a few to several (~ < 20%) trees masting

(Suppl. material 2, table 1: the Evidence Table). The masting scale was therefore semi-

parametric: except for the last 2 years the scale was quasi-binomial. In the Results section,

three categories of event are use: 'M' and 'm' as masting; and the set "none, very low and

low" as non-masting. Direct estimates of seed output, as a quantitative measure of mast

fruiting, or even qualitative crown scoring, for all trees in all years was simply not feasible at

this remote and difficult field location. The reduced and simplified scale for masting provides

the best lowest denominator allowing all years of recording to be combined with very little

bias.

Two canopy caterpillar outbreaks in 2011-12 and 2014 offered the opportunity to

assess how loss of foliage to herbivory, and hence photosynthetic activity, could have

interfered with tree reproduction.

Nutrient sampling

Previous litter samples providing inter-mast nutrient data

In an earlier-conducted litterfall study between May 1990 and June 1992, monthly

bulked samples from twenty 40-cm x 40-cm traps in each of 10 subplots had been divided

into two halves: as set A, for separation into fractions (leaves, small wood, etc.), and set B,

for separation of just leaves into the commonest 26 species (Chuyong et al. 2000). From this

set B, leaves of *M. bisulcata* were selected and later – as for the other species – they are

analyzed for N, P, Ca, K and Mg concentrations. Leaflets and rachises were kept together. As

this species is a leaf-exchanger, it has a very well-defined period of leaf litterfall each year,

starting with onset of the dry season (Newbery et al. 2006a). Five subplots were in low

(LEM) and five in high (HEM) ectomycorrhizal stands (each 40 x 80 m) along the transect P

in Korup (Gartlan et al. 1986; Newbery et al. 1997). Since *M. bisulcata* occurred in only the

HEM subplots, the number of replicate nutrient values per month was five. The 26 months

had been divided into seasons according to the median rainfall pattern at Korup (as defined

above), with those in the dry season being generally December 1990–February 1991 and

December 1991–February 1992, and the other months the longer wet seasons. Only dry

seasons are considered here, and nutrient concentrations for them are now averaged across

the three constituent months. A mast fruiting occurred in (August-September) 1989, so dry

season 1990–1991 is the second dry season after the event, and dry season 1991–1992 the

third one. The next masting was in the wet season of 1992, so seen from this perspective the

two dry seasons selected were ~ 6 and 18 months prior to the second event.

In a later unpublished study of litterfall by Schwan (2003), paired litter traps (same as

those used by Chuyong et al. 2000) were placed at random below the crowns of 20 *M.*

bisulcata trees in the main P-plot. The bulk of the litter (~90%) fell in the 3-week period of 9-

30 Dec. 2002. This 2002–2003 dry season came immediately after a masting in 2002, and

next one was in 2004 (Newbery et al. 2006a). In this study by Schwan (2003) only leaflets

were analyzed for N, P, K and Mg.

New litter samples included in the inter-mast data set

In the dry season 1997–1998 four traps of galvanized metal (same as used in 1990–

1992) were compared with four similar ones but lined with hessian sacking (used in the

1989–1991 study of Newbery et al. (1997), each set being placed at random in each of the

two quarter plots of six HEM half-plots. Litter was collected over four consecutive fortnights

between 21 Dec. 1997 to 15 Feb. 1998. *Microberlinia bisulcata* leaflets and rachises were

again sampled separately, dried and analyzed for N and P concentrations. Not all traps had

sufficient material and the replication per subplot/trap type/date ranged from 2 to 4. The N

and P values were successively averaged for trap type, then subplot, and their weighted 4:1

leaflet-rachis averages found for the 24 date x plot combinations. Trap type had no effect on

these two elements' litter concentrations (initially to test for Zn contamination). This litterfall

was collected midway between a moderate double masting in wet seasons 1997 and 1998.

Chemical analysis followed Chuyong et al. (2000).

An opportunity to sample *M. bisulcata* leaf litter immediately after a later mast

fruiting came in the 2010–2011 dry season. A large system of over 400 seed traps had been

used to follow seed production and dispersal in the August-September 2010 masting

(Norghauer and Newbery 2015). These traps remained in place until early 2011, collecting

leaf litter. The dry season started on 26 Dec. 2010, and samples were taken on 29 Jan. 2011.

This interval of ~1 month would have covered a very large proportion of the leaf fall of this

species that year, although the thin tail into February was not included. Thirty traps, from 68

small clusters of the traps within 25 ha of the 82.5-ha permanent plot Newbery et al. (2013),

had sufficient quantities of almost-only *M. bisulcata* leaves, lying in the traps as a well-

defined readily collectable upper layer. Leaflets and rachises were separated on this occasion

and analyzed for the same five elements as in 1990–1992, later recombining at the whole leaf

level. The mean dry mass ratio of leaflets to rachis in *M. bisulcata* leaves was 0.793:0.207,

very close to a 4-to-1 ratio.

Nutrients in seeds and pods: extending the earlier data set

The 1998 masting event's seeds and pods, previously analyzed and reported in (Green

and Newbery 2002) for N and P, were analyzed further for K, Mg and Ca using standard acid

digest methods and spectrometric determination (see e.g., Newbery and Neba 2019). To find

the nutrient masses allocated to reproduction (fruits) and to compare them to the same used in

foliage, leaf litterfall values from another similarly strong (1992) mast year were taken from

Green and Newbery (2002). In these new calculations, mature leaf concentrations (for N and

P) were used from a retranslocation study in Chuyong et al. (2000), which involved three

dates: viz. December 1991, and April and July 1992. Litter fall dry mass very slightly

underestimated mature leaf mass because of element retranslocation during senescence.

Essentially, these leaf concentrations were from trees sampled on average 5 months before

seed dispersal, and therefore concurrent with pod maturity that year.

Soil moisture measurement

Soil moisture was recorded across two dry seasons to demonstrate that its variation

could underpin soil nutrient processes which were affecting mast fruiting. *Microberlinia*

bisulcata trees spread their extensive lateral fine root systems and ectomycorrhizas mainly in

that drying soil surface layer (Newbery et al. 1997; Newbery et al. 2009). An additional aim

was to determine by how much soil moisture content was reduced by rainfall deficit.

On 2 Dec. 2003, 12 soil moisture content sensors (Theta Probes ML2, Delta-T

Devices, Cambridge, UK) were installed in a stratified random manner across the eastern 30

ha of the P-plot, one per block of 100 m x 250 m. Each probe was inserted horizontally (at 40

cm depth, excepting three at 20 cm due to access problems) into a hole excavated at the side

of a small pit, the latter backfilled and packed down to the original density. A subsample of

soil from the excavated side hole was retained to measure its fresh, water-saturated, and later

oven-dry weight (105 °C). With a hand-held meter, soil moisture content (SMC, %) was

recorded fortnightly until 26 Apr. 2005 (38 times). Readings were repeated between 15 Nov.

2006 and 15 Mar. 2007 (nine times), with one last recording on 29 Jan. 2011. In the years

2004 and 2007 *M. bisulcata* had major mast fruiting, in 2005 and 2011 it had none.

Statistical analysis

The fitting of masting, as a 0/1 binomial response variable, to the climate variables

was first limited to the 24-year period 1989–2012 because all the events were strong and

pronounced ('M' in Table 1; Suppl. material 2, table 1). Logistic regressions were run with

dispersion (φ) either fixed at 1.0 or estimated (Agresti 2007). Whilst the latter gave

marginally more significant fits, the former was adopted because the sample size was small

and the estimation of φ would be unreliable (McCullagh and Nelder 1989). The binomial

error distribution appeared to be satisfactory: the complementary log-log one did not improve

fitting. Simple logistic regression does not account for temporal autocorrelation, however.

With records for only ~60 trees in common across all events, the sample size was too small to

apply Monte Carlo subsampling in order to achieve temporal independence. Because of the

non-stationarity, randomization tests will not provide valid inferences (Cox 1981).

Extending the analysis to the full 29 years (1989–2017) required the ordinal, or

proportional-odds, logistic model (Agresti 2007; Fox 2008; Hosmer et al. 2013), where 'm'

(moderate masting) was scored as 0.5 (see Suppl. material 2, table 1). Binomial and ordinal

logistic models were run with the 'glm' and 'polr' commands in R (Venables and Ripley

2002; Fox and Weisberg 2011), using the package 'MASS' (R_Core_Team 2022), and in

GenStat for comparison where also φ could be estimated (VSN_International 2022). Given

the modest length of the time series and data set, different models were fitted mostly with

single independent terms; and when with two then the same variable with a 1-year lag, and

only additively. No attempt was made to search for a best-fit subset model with interactions;

if fact, most such model fits failed to converge when attempted. Because mean daily dry

season rainfall and radiation were not correlated, these variables were fitted in separate

models (see Results).

Times-series logistic regression – in full, generalized logistic autoregressive moving-

average regression, with the command 'glarma' (R package glarma, Dunsmuir and Scott

2015) – accounts for temporal autocorrelations in binomial data (Cox 1958, 1981; McKenzie

1985; Cox and Snell 1989; McKenzie 2003; Dunsmuir 2016) and it was applied to the

masting data.

RESULTS

Mast fruiting pattern

The extended time series shows that after 1995 there was a double year 1997–1998

with high seed crops, 1998 somewhat weaker than 1997, and then the interval dropped to 1

year (from a 3- to a 2-year cycle), with a masting sequence of 2000, 2002, and 2004,

returning to strong 'M' masts in 2007 and 2010 with a 2-year interval, after which the control

seemingly weakened to result in two pairs of consecutive years, 2013–2014 and 2016–2017

which had moderate 'm' mast fruiting (Fig. 1a). The time series as it extended began to show

increasing non-stationarity (i.e. change in temporal pattern over time), the 3-year cycling of

first period of 1989–1995 returned only once in 2004–2010: the regular cycle was seemingly

temporarily disturbed. Nevertheless, 'on average' over the 29 years, and taking as markers

the first of those pairs of years with partial masting (2013, 2016), there appears to be good

evidence for an intrinsic 3-year cycle. Synchrony across trees within masting events was very

high.

Climate variation in time

Of the four basic climate variables, mean rainfall and radiation per day in the dry

season along with the start and length of the season (current year of masting and year prior),

only the first required ln-transformation to remove positive skew along the independent

variable axis. Dry season rainfall and radiation were not significantly correlated ($r = 0.110$, P

$= 0.57$, df $= 27$) and could be treated as being independent.

Mean daily dry-season rainfall in years of full masting was lower than in those years'

prior with no masting (Fig. 2a) except for 2000. Further, if the averages of the rainfall values

for the two double partial masting years are taken these are on a par with the masting ones.

The time series for rainfall showed weak temporal autocorrelation (Suppl. material 3, fig. 3)

and likewise, albeit a little more strongly, for ln($rain$) at lag 1 (–0.293 vs. –0.221). (Note that

the first correlation differs slightly from the rain versus $rain_{-1}$ one based on data which

included the lagged 1988-value.) Radiation over time showed a distinctive pattern with a

clear central peak between 1996 and 2005, then a rise again, apart from 2009, to those higher

levels (Fig. 2b): accordingly, the series evidenced non-stationarity, and as a consequence

autocorrelation was positive and significant for lags of 1 and 2 years, then significantly

negative ($P < 0.05$) for lags of 6–8 years (Suppl. material 3, fig. 3). The partial

autocorrelation remained significant for a lag of 1 year only, however.

Start of the dry season varied around the third to fourth week of December with no

clear patterns or differences between masting and non-masting years (Fig. 2c), but it was

negatively autocorrelated with a lag of 1 year, i.e. early starts tended to be followed by late

ones, then early again, in successive years (Suppl. material 3, fig. 3). Excepting the two last

years, duration of the dry season revealed a general decline (an approximate halving), those

years with masting being often longer, and conversely of the six shortest-season years there

was no masting, apart from 2014 which was partial (Fig. 2d). Interestingly, autocorrelation of

duration was almost significant ($P < 0.05$) at a lag of 3 years (Suppl. material 3, fig. 3).

Radiation intensity showed a very similar pattern to duration (since radiation varied much

less than duration) (Fig. 2e; and Suppl. material 3, fig. 3). Intensity of radiation in the dry

season was expressed as mean daily radiation x duration. Lastly, difference in radiation

($radi_d$) between successive years (one approach to removing trend and non-stationarity) was

mixed regarding masting response, although again a weakly significant negative

autocorrelation for a lag of 1 year was evident (Fig. 2f; and Suppl. material 3, fig. 3).

Modelling masting and climate

Binomial logistic regressions for 1989–2012 showed a significant negative effect of

mean daily rainfall in the current year's dry season ($P < 0.05$), and a positive effect of mean

daily rainfall in the dry season prior ($P < 0.05$), on masting. Using ln($rain$) and ln($rain_{-1}$)

improved the regressions fits with $P < 0.015$ (Suppl. material 4, table 1). Mean daily rainfall

was only weakly correlated with rainfall the year prior: untransformed values ($r = -0.190$, P

$= 0.374$), transformed ($r = -0.272$, $P = 0.15$). This last result is important because a strong

negative temporal autocorrelation, i.e. a year-to-year increase-decrease reversal, was absent.

Regressions with both rainfall terms led to reduced fitting, because these coefficients were

now partial ones adjusting for the correlation ($P = 0.093$ and 0.055). The same regressions

with ln-transformed variables also changed the significance levels very little ($P < 0.10$;

Suppl. material 4, table 1). Ordinal logistic regressions improved the fitting further with the

full 1989–2017 data. Concentrating on the models using ln-transformed variables, separate

regressions were now more, and highly, significant ($P < 0.01$), having coefficients with the

same signs as for the binomial logistic ones (Suppl. material 4, table 1) and, whilst

significance decreased again using two-term models, the fits were still relatively strongly

significant overall ($P < 0.025$).

The mean difference in ln-transformed dry-season mean daily rainfall, ln($rain$),

between the current masting (t_0) and year prior (t_{-1}) was highly significant ($P < 0.005$), but

much weaker for the same and penultimate year (t_{-2}) ($P = 0.15$). Back-transformed means

were 1.37, 2.49 and 1.69 mm/day respectively (see Suppl. material 4, table 2). Comparing

rain at t_0 with means of $rain$ at t_{-1} and t_{-2} – applying equal or unequal weighting – led to very

similar results as those at t_{-1} alone ($P < 0.005$). As a robustness test of the strong vs moderate

('M'/'m')-grading of mast fruiting, and how that might have affected the relationship with

mean daily rainfall in the dry season where there was some uncertainty in the final grading,

the 1989–2017 ordinal logistic regressions were rerun with 'M' for 1998 and 2000 replaced

by 'm', i.e. 1.0 by 0.5. The dependence of masting on ln($rain$) in the current and prior seasons

became slightly stronger ($t = -3.209$, $P = 0.0036$; $t = 2.913$, $P = 0.0074$, respectively).

Mast fruiting showed no significant dependence on mean daily radiation in the

drought-defined dry season, for either the 1989–2012, or 1989–2017, time series (using

binomial and ordinal regression, respectively), for the current ($radi$; $P = 0.72$ and 0.80) or

prior ($radi_{-1}$; $P = 0.55$ and 0.62) seasons. Correspondingly, start date was also a poor

predictor of masting for the current ($start$; $P = 0.34$ and 0.29) or prior ($start_{-1}$; $P = 0.18$ and

0.27) seasons. Duration, although only weakly significant, was better for the current (dur; $P =$

0.080 and 0.095) or prior (dur_{-1}; $P = 0.071$ and 0.138) seasons. It is important to note here

that radiation values for the 2010 and 2011 dry seasons had been interpolated (Suppl.

material 1), so model predictions for the 2010 masting event come with an additional level of

uncertainty. Masting was only marginally or non-significantly dependent on masting intensity

in the current and previous dry seasons (positively and negatively respectively; binomial

logistic, $P = 0.083$ and 0.097; ordinal logistic, $P = 0.096$ and 0.165). Hence there was some

indication that high intensity reduces the likelihood of masting in the following year.

Differences in mean daily radiation (current-minus-previous, $radi_d$), on the drought-defined

dry-season basis, had no clear effect on masting (binomial logistic, $P = 0.31$; ordinal logistic

$P = 0.84$).

Radiation did show a long-term trend from the start of recording until 2017 (Suppl.

material 3, fig. 2), with evidence of a 11-year solar cycle. This was not related to the change

in instrumentation at Bulu (Suppl. material 1). Using a LOWESS regression fitting model in

R ('lowess' command – a set of local smoothing splines), this trend could be removed and the

differences between empirical measurements and the detrending curve then used as adjusted

local year-to-year changes (Suppl. material 3, fig. 2[a]). The best result was with a span

parameter of $f = 0.25$, which meant that radiation values > 1 year prior or ahead had some

weight in fitting values for any one year, which simple differences between successive years

did not. Binomial (1989–2012) and ordinal (1989–2017) logistic regression fits of masting to

drought-defined dry-season *radi* and *radi*$_{-1}$ were, however, all clearly insignificant ($P = 0.34$

to 1.00).

Lastly, to check whether heavy showers early in the dry season may have damaged

flowers or interfered with pollination success, mean daily rainfall was plotted across each

year's season. Of the 14 years with strong ('M') or moderate ('m') mast fruiting, ten had no

showers of ≥ 15 mm/day in the first 30 days, whilst in the 15 years without mast fruiting

eight had no showers ($\chi^2 = 3.55$, df $= 1$, $P = 0.060$), suggesting only a marginal tendency for

mast years not to have experienced potentially destructive rainfall events.

Taken together, these regression model fits suggested that only mean daily rainfall in

the current and prior dry seasons was strongly affecting the odds of masting. The exponent of

the coefficient (e^β) estimates the fold-change in odds of masting per unit increase in mean

daily rainfall. Since the range in mean daily rainfall over the years was approximately 1 to 3

mm, this translates on the ln-scale to *c*. 1 unit, meaning that in the transformed-variable

models the change in odds covers the range in rainfall. This outcome provided a sound basis

for predicting future masting events.

Predicting masting from models

Using binomial logistic regressions of masting 1989–2004 on ln(*rain*) in the current

and previous years, fitted models were used to predict masting in 2005–2017 (Fig. 1), on the

original response scale of 0 to 1 (π; Agresti 2007). The series started at 1989, not 1987 as

used in Newbery et al. (2006a); and autocorrelation was not incorporated (see 'Time series

analysis' later though). The model fit for ln(*rain*) was good for the mast years 1989, 1992,

1995 and 1997 (marginal for 1998), and good for 2002 and 2004, but it wrongly indicated a

mast in 1993 and missed the mast in 2000 (Fig. 1b; fitted intercept [α] $= 3.61 \pm 1.80$ and

slope [β] $= -5.77 \pm 2.62$, $t = -2.206$, $P = 0.027$). Predictions of masting in 2007 and 2010

were very high (π = 0.97 and 0.83 respectively), and also that it did not occur in the

intervening years. After 2012 though, one each of the pairs of moderate ('m') masts was

predicted as full ('M', π = 0.98 and 0.97). The predicted masting in 2005 (π = 0.75) was

incorrect, however.

With the model fit for ln($rain_{-1}$), 1992, 1995, 1997, 2000 [cf. with ln($rain$)], 2002 and

2004 were all well-fitting as mast years, but 1989 and 1998 were not: 2001 was fitted as a

mast year wrongly (Fig. 1c; fitted α = –2.47 ± 1.40 and β = 3.96 ± 2.00, t = –1.98, P = 0.027).

Again, the model exactly predicted the mast years 2007 and 2010 masts (π = 0.76 and 0.97

respectively), and it managed three of the four 'm' (moderate) years afterwards as being

moderate masts reasonably correctly. Therefore, similar to the outcome for cycles and

intervals in mast fruiting, the model fits were fair-to-good up to 2012 (the caterpillar-outbreak

year; see later subsection) but after that disorder and unpredictability ensued. Binomial

logistic regression fits made with the 1989–2012 data gave predictions (not shown) that were

very similar to those for 1989–2004, in the main because the masting in 2007 and 2010

matched so well the time series to 2004: yearly π-values (fitted and predicted) from the two

runs were in very good agreement.

The median effective level of ln($rain$), that is the level when π (x) = 0.5 (defined as x

= –α/β; Agresti 2007) was 0.624, which back-transformed to 1.869 mm/day – a value very

close to the estimate for the ln(rain$_{-1}$) estimate of 1.867 mm/day. This median level, which

might be interpreted as a form of threshold for non-masting and masting, applied well to the

events and their rainfall values in most years up to 2011 (Table 1, Fig. 2a), although ln($rain$)

and ln($rain_{-1}$) differed in a few years as being better or worse fitting and predicted. Years

2012, 2013 and 2014 had moderately high and quite similar mean rainfall in their dry seasons

(1.90–2.07 mm/day; Table 1), so aside from the caterpillar intervention, the rain threshold

theory would not have been expected to work too well – a form of counterfactual evidence.

Afterwards, in 2014–2017, the return to low–high alternations in rainfall should, *ceteris*

paribus, have led to 'M' (strong) masting in 2014 and 2017 (as predicted by the binomial

logistic regression for ln(*rain*) (Fig. 1b), but it still did not achieve that outcome.

Time series approach

Application here supported the binomial logistic regression results for 1987–2012

even more strongly ($P < 0.01$ and $P < 0.05$ for ln(*rain*) and ln(*rain$_{-1}$*) respectively, for both

lags p = 1 and 2, (Suppl. material 4, table 1). Regressions using the time-series logistic

regression model for *radi*, *radi$_{-1}$*, *start* and *start$_{-1}$* either failed to converge or the fits were

insignificant ($P = 0.17$ to 0.73). Whilst there was a weak positive effect on masting for

duration in the current year ($\beta = 0.041 \pm 0.025$, $z = 1.64$, $P = 0.10$), an even stronger negative

effect of duration was evidenced for the year prior (duration$_{-1}$; $\beta = -0.099 \pm 0.038$, $z = -2.62$,

$P = 0.0087$) (Suppl. material 4, fig. 2). The last relationship analyzed before with a binomial

logistic model was only marginally significant ($P = 0.073$), so the evidence for the effect

improved.

The dependence of mast fruiting on dry season rainfall appears to be limited to the

current and previous (t$_{-1}$) years, because repeat runs replacing *rain$_{-1}$* with *rain$_{-2}$*, the mean

daily rainfall in the second year prior to the current one (t$_{-2}$), were all clearly insignificant (P

= 0.39 and 0.40) for 1989–2012 binomial logistic regressions, rainfall ln-transformed, single

and two-term models respectively; and they were correspondingly insignificant ($P = 0.35$ and

0.76) for the 1989–2017 series with ordinal logistic regression. Moreover, dry-season rainfall

at t$_{-2}$ was poorly correlated with the same in the current year ($r = 0.027$ and 0.110), but

slightly better with that at t$_{-1}$ ($r = -0.187$ and -0.257). Time-series logistic regression fits did

not improve fitting to intensity of radiation (current or prior), differences in dry season

radiation in successive years, or smoothed radiation regression residuals (current or prior) (P

= 0.094, 0.062, 0.25, 0.74 and 0.26; respectively). Finally, masting was unrelated to

minimum temperature in the current year's dry season (1989–2012, Table 1; binomial logistic

$P = 0.18$; and 1989–2017, ordinal logistic $P = 0.24$).

Extending models for radiation

Whilst a rainfall deficit delimitation will define the drought period in terms of limited

water availability, it may not necessarily correspond precisely to times of heightened

radiation. Returning to the modal definition of dry season, namely the calendar months

December through February (DJF), or first quarter of the phenologically-referenced year,

mean daily radiation can likewise be found (Suppl. material 3, fig. 1). Masting was not

strongly related using binomial, ordinal or time-series logistic regressions, to this quarter's

radiation however, in either the current or prior year ($P \geq 0.18$; Suppl. material 4, table 3[a]),

and also not for the other three quarters (Suppl. material 3, fig. 1; $P > 0.4$; model fits not

shown). By contrast, and surprisingly, masting was strongly dependent on the *difference* in

radiation, both in absolute ($radi_d$) and percent change ($radi_{d\%}$) terms, from prior to the current

year, and with all three regression models ($P = 0.007$ to 0.024; Suppl. material 4, table 3[a]

and fig. 3; and Suppl. material 3, fig. 1). Using again LOWESS smoothing regressions to

remove trend in radiation (Suppl. material 3, fig. 2[b]), with binomial logistic regression

(1989–2012) masting weakly positively depended on differences in *radi* ($P = 0.051$) and

negatively, though more marginally, for $radi_{-1}$ ($P < 0.1$) (Suppl. material 4, fig. 3 and table

3[b]). Time-series logistic regressions improved the fits especially for *radi* ($P < 0.02$),

although only slightly for $radi_{-1}$ ($P = 0.08$). Fits to detrended differences (1989–2017) with

ordinal logistic regression were intermediate in significance to those with binomial and time-

series logistic ones (Suppl. material 4, table 3[b]). Overall, there emerged moderate to strong

evidence that the larger the increase in radiation between successive years, or the more

positive the difference from the trend, the more likely a mast fruiting. This difference effect

was not obvious in the earlier analyses of Newbery et al. (2006a).

An interesting feature of the LOWESS fits (Suppl. material 3, fig. 2), when taking

radiation according to the two definitions of dry season, is how the outliers changed in

relation to the smoothed curves. In the drought-defined case, 2001 and 2003 (both 1 year

before masting) were very high, and 2006 and 2009 (also both 1 year before masting) were

very low. Changing to the DJF-defined case brought 2001 and 2009 radiation much closer to

the curve, and the binomial and time-series regression fits were improvements. Further, the

autocorrelation was notably stronger, positive and significant, at lags of 1 and 2 years for this

DJF-defined dry season than the drought-defined one, shifting to a significant negative

correlation at an 8-year lag (Suppl. material 3, fig. 4). The partial correlation, again, remained

significant for just the lag of 1 year. In the third (JJA) and fourth (SON) quarters a similar but

slightly weaker pattern for the function occurred, the negative lag now at 10–12 years (Suppl.

material 3, fig. 4). More interesting, however, was the almost unstructured weak pattern of

the function for the second (MAM) quarter, which is no temporal autocorrelation over the

early wet seasons (Suppl. material 3, fig. 4). Difference in radiation was significant for a lag

of 3 years, and remained almost so for the partial correlation, hinting at the 3-year cycle.

Start and duration of the dry season

To shed some light on the apparent paradox of the different definitions of dry season

determining the relationship between masting and radiation, regression models were rerun

with an extended dry season of the months of November through to March. All binomial

logistic model fits, 1989–2012, using radiation in the current and prior dry seasons on this

new basis were clearly insignificant ($P = 0.27$ and 0.69). However, dependence on masting

on $radi_d$ was marginally significant at $P = 0.071$ ($radi_{d\%}$, $P = 0.052$), and correspondingly so

was it for the time-series logistic model with p = 2 (P = 0.056 and 0.043). Compared with the

fits for the models using DJF as dry season (Suppl. material 4, table 2[a]), these results for the

5-month season were weaker, suggesting that the influence of radiation was being diluted by

inclusion of the additional 2 months; and this was reflecting the early wet season

autocorrelation. Comparing radiation defined by DJF and that defined by the drought dry

season graphically, both for differences between successive years and as residuals from the

smoother curve over time (Suppl. material 4, fig. 4), indicated that a majority of the years

lying above the 1:1 reference line were ones with masting and most of those below were non-

masting. This confirmed that raised radiation differences and deviations were associated with

mast fruiting.

The placing of start and end dates of the dry season was evidently critical to the

analysis outcome because the degree of overlap between the two types of dry season were

very variable from year to year (Table 1, Fig. 1a). Mean, standard deviations and ranges in

absolute differences were actually larger and wider for the drought- than DJF-defined dry

periods (1.41 ± 11.14 [–21.0 - 21.1] and 0.72 ± 9.62 [–15.0 - 16.2] respectively). Even so, the

reasons for the considerable overlap could be various because start and end dates of the

drought-defined dry season were often very different from those of the DJF one.

When the mean daily radiation of the non-overlapping weeks with DJF increased or

decreased, or vice versa, the $radi_{d\%}$ altered between successive seasons. Since masting fitted

to $radi_d$ much better for the DJF- than drought-defined seasons, this would suggest that (a)

radiation levels outside of the DJF months were most likely important, and (b) DJF-defined

seasons allowed a more consistent year-to-year comparison than the drought-defined ones.

That rain may have been lacking, which defined the duration of the drought-defined season,

did not also imply radiation was changing proportionally. Radiation could have been

sometimes high in the day and rain fell at night: conversely radiation could have been low

when there was cloud cover and yet no rain, as was the case in some years at the transition to

the wet season. Differences in successive years' dry season radiation were more, respectively

less, pronounced in the first and second halves of the time series, for the DJF- than the

drought-defined dry seasons.

Soil moisture content in dry season

On graphing the time series for 2003–2005, three of the 12 sensors had three

anomalous and coinciding high peaks ($\sim$ 40 - <60%) in the 2004 and 2005 early-to-mid wet

seasons suggesting that their locations were prone to flooding. These sensors' data for the

whole period were accordingly omitted. Two of them were actually destroyed later by

hunters. The time series of mean SMC is shown in Fig. 3: it included the two dry seasons of

2004 and 2005, defined as before as when *rft* fell < 100 mm. Mean SMC over the 1.4 years

was $16.80 \pm 0.48\%$. Repeated measures analysis of variance for six times within each of the

two dry seasons (Fig. 3), using all 12 sensors' values, indicated that differences in mean SMC

at the 20-and 40-cm depths were very small and insignificant: 10.98 vs 10.69% in 2003–2004

and 16.20 vs 15.89% in 2004–2005, $P = 0.88$ and 0.89 respectively, df = 1, 10). In a similar-

length period in the late wet season 2004, starting soon after the peaked occurrences that

year, a small difference was evident but again it was clearly not significant (20.20 vs 18.13%,

df = 1, 10; $P = 0.39$). Interactions between depth and time were all insignificant for all three

short periods ($P = 0.19$ to 0.36).

After the strong dry season of 2003–2004, during which SMC reached a minimum of

10.4%, its values remained fairly constant at $18.24 \pm 0.31\%$. The dry season of 2004–2005

had much less influence on SMC (Fig. 3). The lowest SMC-values in 2004 coincided with the

lowest *rft*-values (down to 12 mm then 0 mm over 4 weeks); in 2005 the SMC either did not

go so low or remain low for long, and the match with SMC was weaker. At the start of the

2003–2005 series, the means ($\pm$ SE) of SMC recorded and those found by gravimetric

analysis were 15.09 ± 0.52 and $15.50 \pm 0.82\%$ respectively. The estimates were positively

correlated ($r = 0.677$, df $= 10$, $P = 0.016$). Saturated water holding capacity was on average

$38.15 \pm 1.16\%$, less than the SMC at the peak times, confirming that some locations were not

always free draining.

Between 2 Dec. 2003 and 26 Apr. 2005, SMC was related to 30-day running rainfall

total as follows: SMC $= 14.4 + 0.0070 \cdot$rft ($R^2_{adj} = 40.5$ %, $F_{1,36} = 26.18$, $P < 0.001$, $n = 38$),

inferring 14.4 % at 0 mm, and 21.4% at 1000 mm – the last *rft*-value being close to the June

2004 maximum of 995 mm. The difference in SMC between the very driest and wettest times

of the mast year was therefore only 7% – for those locations assumed free-draining. SMC

before the dry season of 2006–2007 was $20.7 \pm 0.10\%$ and declined sharply and almost

linearly to $14.0 \pm 0.67\%$ ($n = 9$ sensors), a difference of 6.7%. With the dry season that year

lying within the span of SMC recording dates, this last value was likely very close, if not at,

the 2007 SMC minimum. The mean value on 29 January 2011, a date also well into that

year's dry season was $16.2 \pm 0.8\%$ ($n = 8$, one sensor lost because its cable had been

chewed), a value close to that in the dry season of 2005 (Fig. 3).

The 30-day running rainfall total, *rft*, used to define the dry season in terms of drought

when this total goes below 100 mm, does not indicate either how much is the deficit, or how

steep is the decline into, or incline out of, the dry season. Suppl. material 5, fig. 1 shows the

course of *rft* as the daily rainfall events occurred. The intense dry season of 2003–2004 (2004

was a masting year) resulted of *rft* going to zero, or very close to zero, for about 6 weeks

(Suppl. material 5, fig. 1[b]), while in 2004–2005 (2005 was a non-mast year) it was near

zero for perhaps just 2 weeks, and shorter overall (Suppl. material 5, fig. 1[c]). In 2002–2003,

for comparison, the also long dry season *rft* reached zero for a couple of days, and whilst

many days had no rain the season was broken by three high rainfall events keeping *rft* just

below 100 mm (Suppl. material 5, fig. 1[a]). The relatively small decrease in mean daily

rainfall between 2005 and 2004 (Fig. 2) translated into a surprisingly large difference towards

the lowered SMC and suggested a non-linear effect, that is a little less rain led to

proportionally much more soil drying when rainfall was low-to-average.

Caterpillar outbreaks

In the latter half of December 2011, and again in the first half of March 2012, an

outbreak of a black caterpillar was observed that fed on young foliage of *M. bisulcata* trees

recorded for phenology in the P-plot. Although not quantitatively measured, the attacks were

extensive, as evidenced by the substantial amounts of falling frass, numerous caterpillars

hanging below the crowns on threads, and litterfall consisting of soft, green leaf tissues (G.

A. Neba and S. Njibili, pers. obs.). The same caterpillar was also recorded at another *M.*

bisulcata grove 4.2 km due south, in March 2012 (Norghauer et al. 2023). However, no visit

was made to that site in December 2011. Fallen caterpillars were also observed feeding on

young leaves of *Oubanguia alata*, a common understorey tree growing below *M. bisulcata*

crowns at Korup.

Also in the latter half of December, but in 2008, an outbreak of most likely the same

caterpillar species (an *Achaea* sp., probably *A. catocaloides*) had been recorded on *T.*

korupensis trees in the same P-plot (Norghauer et al. 2023). At that time, feeding upon *M.*

bisulcata foliage was limited to low levels on just those crowns closely adjacent to the more

heavily attacked ones of *T. korupensis*. Furthermore, and separately, at the end of January

2014 a morphologically different and larger species of caterpillar, yellow-green and behaving

very differently from the smaller black one, was seen on five of the 65 monitored *M.*

bisulcata trees (Suppl. material 6, text and fig. 1). Most probably it was another *Achaea*

species. Intriguingly, this caterpillar species was feeding on *M. bisulcata* flowers, not young

foliage. How influential the new insect herbivore was for *M. bisulcata* phenology that year is

uncertain because its distribution was localized to a 2–3-ha patch where the *M. bisulcata*

basal area was high.

Tree phenology and masting intensity

Tree phenology of *M. bisulcata* was compared between the heavy mast fruiting year

of 2010, and the year of 2012 in which fruiting was very low (Fig. 4; Suppl. material 2, table

1). Both time series covered the drought-defined dry season of that year and followed

changes in leaf-flush and leaf-fall (leaf exchange), and flowering, pod formation (green) and

pod maturity (brown). Attaining continuous fortnightly scoring was not always feasible and

each series has two gaps. Leaf flush likely began a little before the start of the time series

recorded for 2010, i.e. earlier than December 2009 (Fig. 4a), but was adequately captured for

2012 (Fig. 4b). The bulk of the leaf-flush happened within each year's dry season, flowering

well within it in 2010 but coming at the end in 2012. Even though the sum of the flowering

scores was similar in both years, pollination resulted in substantially more green and brown

pods in 2010 than 2012 (Fig. 4a, b). Caterpillars were not seen in the other years between

2009 and 2013, notably not in 2010. In 2010, green pods were apparent between 18 January

and 3 July, that is for 5.5 months, with a 1-month overlap as they turned brown between 5

June and 28 August, over 2.5 months (Fig. 4a). A similar time course was recorded in 2004,

and likewise (for green plus brown) in 1995 and 2000 (Fig. 4b of Newbery et al. 2006a).

Interestingly the overall period of pod formation was much shorter in the two back-back

years of masting 1997 and 1998.

Trees in 2012 suffered unusually in two definite caterpillar attacks by the same black

Achaea species, both coming just prior to peaks in leaf-flush (marked 'C' in Figs 1a and 4).

The earlier flush in early January 2012 was part of the annual leaf exchange process, but the

later one in early April 2012 not. The first was in part associated with the leaf fall around 15

Jan. 2012: 21 of the 61 trees lost new leaves then. The third leaf flush in late February 2012

(Fig. 4b) had no associated caterpillar attack. What happened in late March was unfortunately

not recorded, but a building up of the re-flush starting then could be surmised if this event

followed the same flushing patterns of the earlier and later ones. The sequence of events

suggests that leaf flush in early January enabled the caterpillar population to initially increase,

and that at the end of February there was a response by the trees to the first attack. The flush

at the end of April is more difficult to explain. It could have been either a further delayed

response to the first attack, or a direct response to the second attack, by a possibly now larger

caterpillar population. The life-cycle dynamics of the herbivore would have allowed,

however, only a second cohort of caterpillars to feed on that third and last flush for the year.

The most flowering accompanied the middle leaf-flush peak in 2012, which speaks against

caterpillars being around then because feeding on leaves might be expected to indirectly

reduce flowering.

The presence of the black caterpillar in 2012 appears to have upset the smooth

phenological stage-replacement of 2010, and led to a much-reduced reproductive output. The

interval from the last year of full masting (1 year) was theoretically suboptimal, so likely the

resource levels of P but not N were reduced and still below a threshold. A high N: P ratio

may have helped the caterpillar outbreak. In addition, the masting pattern changed after 2011

(Fig. 1a), to the two moderate double-year form, possibly a carryover of the disruption of the

feeding in 2012, and a destabilization of the proposed inherent 3-year cycle. In the dry season

2010, mean daily rainfall was 1.42 mm, yet the year prior it was 4.66 mm; correspondingly

the rainfall for 2012 were 1.95 and 2.07 mm: prior-to-current ratios of 3.3 and 1.1

respectively. Interestingly, the lower caterpillar activity noted in the other grove to the south

was associated with average green-pod score of 1.7 in May 2013 ($n = 33$ trees; Suppl.

material 2, table 1), showing that masting was more intense there than in the P-plot that year.

The second herbivore attack in January 2014 of the other unknown caterpillar species

was before the second of the moderate 'm' mast fruiting of the double-years 2013–2014.

Much less is known about its effects particularly because detailed phenological recording did

not continue to end 2014. However, given that this species eats flowers, as well as young

leaves (Suppl. material 6) it most likely reduced the number of trees able to fruit that year,

even if the outbreak was contained to a far smaller known area than the successive outbreaks

of the black caterpillar in late December 2011 and early March 2012.

It is instructive, once again, to look at the course of rainfall events and *rft* during and

immediately after the dry seasons of 2009–2010 and 2011–2012 to find clues for the second,

early March, caterpillar attack. This last came during a second short dry period when there

was that unusual third leaf flush (cf. Fig. 4b with Suppl. material 5, fig. 2[c]). With an

interval of ca. 14 days between phenology census recordings, it is difficult to say which came

first; but the most likely explanation is the further dry period caused flushing and this

promoted renewed feeding, possibly from a next cohort of caterpillars that year. The dry

season 2009–2010 had also several dry days at a similar time but was broken by a large

wetting event (and there was no further flush after the dry season one). For comparison, the

pattern of 2010–2012 shows the shorter, later-starting and not so intense, dry season for a

typical non-masting year (Suppl. material 5, fig. 2[b]).

Tree size-fruiting relationship

Consistency in the intensity of masting over time at the individual tree level was

tested in the following way. Scores for brown pods in 2007 and 2010 were matched with

those for an equally strong masting in 1995 on a tree-by-tree basis. Score categories '0' and

'1' were combined. Fruiting was significantly positively associated in both cases ($\chi^2 = 10.61$,

df = 1, $P = 0.031$ and $\chi^2 = 10.28$, df = 1, $P = 0.036$, respectively). There is good evidence that

high- and low-scored trees tended to remain as such over a 12–15-year period, although

considerable between-trees variability existed.

Ordinal logistic regressions of brown pod scores per tree versus tree stem diameter (at

last census in 2005), in 2010, 2012 and 2013, or as fallen pods in 2014, were weakly and

insignificantly related though ($t = 1.40$, 0.45, –0.68 and 0.42, e.d.f. = 58 to 61, $P = 0.17$ to

0.60). As the phenology score was recording the *relative* proportion of a crown fruiting, this

meant that large trees did not have proportionally more pods than smaller ones.

Oil palm yield comparison

Access to fruit yield data for oil palm (*Elaeis guineensis*) growing in the estates of

PAMOL Plc on the eastern border of the KNP (Bulu-Mundemba, Ndian) allowed a

comparison with *M. bisulcata* mast fruiting under the same local climate conditions. An

analysis of the monthly yields, for 2003 to 2016, showed no year-to-year differences in

means attributable to changes in either rainfall or radiation, although there was a very clear

repeating pattern for each of the three variables within each year (Suppl. material 7, text and

fig. 1). Moreover, mast fruiting of *M. bisulcata* in this period (see Fig. 1a again) did not

correspond to peaks or lows in this proxy measure of productivity, for either the current or

prior year. This suggested that radiation and C fixation were not the primary drivers of mast

fruiting.

The poor relationship with radiation also means that yield cannot be used in an

inverse estimation to confirm the interpolated missing months of radiation at Bulu in 2009–

2011 (Suppl. material 1). The Ndian estates have the same substrate and soil type as southern

Korup. After initial trials in the 1960s through to the late 1970s, phosphate fertilization was

discontinued because little beneficial effect on palm fruit production was found (C. E. Etta,

unpubl. data). It is unlikely that much of the added P was still available in the soil by 2003.

Stem increment growth and masting

Annual radial increments (*incr*, mm) in stem growth of 20 cored *M. bisulcata* trees $\geq$

50 cm diameter in the P-plot from Newbery et al. (2013) were matched to occurrences of

masting over the 15 years of 1989–2003 (Suppl. material 8, fig. 1[a]). The end date was the

last year which had a reliably determined annual band for all trees. Standard deviation

increased linearly with the mean over years (slope [CV] = 1.14, adj. R^2 = 50%, P = 0.002),

indicating a considerable variability in increment between trees in any one year (Suppl.

material 8, fig. 1[b]). No masting was noted for 1988 (Newbery et al. 1997) but that is not

fully certain.

Linear mixed-effects models were run using ln(*incr*) as the response variable, trees as

subjects (or 'blocks') and ± 'M' as the factor. A binomial logistic regression fit of masting

versus increment was not feasible as the power, with only seven events in the period, was

low. There were no significant differences for increment in either the current year (F = 0.79,

df = 1, 279; P = 0.38), or in the year prior (F = 0.12, df = 1, 279, P = 0.72: increments

available for 1988), or in the year after (F = 1.90, df = 1, 259, P = 0.17). Thus, larger

respectively smaller increments were not associated with current mast fruiting, not (by

inference) caused by it beforehand, or conversely masting did not cause a change in

increment afterwards. Graphs of the increment series for each of 20 trees individually (not

shown) had no common correspondences with 'M' years, and the differences in mean

individual tree increment for fruiting type were highly inconsistent (+M: mean 2.49 ± SD

1.97, –M: 2.38 ± 1.98; differences 0.11 ± 0.77, range = –2.02 to 1.67). Mast fruiting appeared

little connected to stem growth in *M. bisulcata*.

Increment was positively correlated, however, with mean daily radiation in the

typically modal dry season period (using all records, i.e. 1984–2003), this 'quarter 1' being

defined by months DJF – ($r = 0.507$, df =18, $P = 0.023$), but not in the early and mid-wet

season quarters 2 and 3 ($P = 0.32$ and 0.41), although again in late wet season 'quarter 4' it

was ($r = 0.479$, $P = 0.033$). Fruiting would have been largely 50% over then, by mid-

September. For the year as a whole the correlation was also reasonably strong ($r = 0.503$, $P =$

0.024). Increment was not strongly correlated with rainfall in any quarter, nor overall ($P =$

0.12 to 0.70), and not with radiation in the year prior to masting ($P = 0.10$ to 0.73).

Binomial logistic regression fits of masting for 1989–2003 to mean annual core

increment with either *radi* or *radi$_{sr}$* (current or prior year for both variables) were all

insignificant for the increment term (current: $P = 0.49$ to 0.63; prior: $P = 0.46$ to 0.95). The

same approach applied using increment of the year prior brought little improvement in fitting

($P = 0.18$ to 0.33; $P = 0.18$ to 0.62, respectively). Thus, whilst increment and dry-season

radiation remained significantly positively correlated ($r = 0.532$, df = 13, $P = 0.041$),

partialling out a radiation effect was of little avail in unravelling a fruiting-stem growth

interaction. These new analyses on the stem increment data provided further important

opposing evidence for C not being the limiting factor for mast fruiting.

Nutrients during the masting cycle

The newly reported nutrient data, coming from the approaches described in the

'Methods' section above, are combined here with other nutrient data already published (see

citations within the footnotes to Tables 2 and 3). To begin with, for 1997 and 1998, and for

2011, the leaflet-to-rachis ratio of N was 1.568 and 1.524, and correspondingly for P it was

1.344 and 1.274 respectively, with means of 1.546 and 1.309. For 2011 only, the L/R ratios

for K and Mg were 0.540 and 1.686, respectively. It was striking how much higher the K

concentration was in the rachises compared to leaflets.

In the 1990–1992 litter sampling the bulked samples each represented 0.320 ha of

forest. In 2011 sampling the trap clusters represented on average 0.368 ha. This means that

each nutrient value is similarly based on ~ 1/3 of a hectare, and the respective samples sizes

of 5-and-5 versus 30 can be retained in statistical comparisons without needing to group the

2011 ones. Likewise, the 1997–1998 study used some of the same half-plots, and accordingly

had the same scale as in 1990–1992. Working with the typical masting cycle of 3 years

shown by *M. bisulcata* at Korup over the period 1989–2010 (Newbery et al. 1998; Newbery

et al. 2006a), the three sampling dates together allow a series of immediately-after (2010–

2011), and 1 (1990–1991) and 2 (1991–1992) years after (and thus also immediately before

the next), masting events to be constructed. The two consecutive years are likely temporally

correlated, but the time difference between them and 2011 is large. The further data from

1997–1998 present an interesting comparison in that masting in 1997 had evidently not

prevented a repeat even the following year. Over the successive four fortnights,

concentrations of N were 18.69, 17.20, 18.58 and 18.63 mg g^{-1}, and those of P were 0.802,

0.799, 0.787, 0.956 mg g^{-1}.

The concentrations across the three weekly litter samples in December 2002 (Schwan

2003), were averaged per trap. To obtain whole-leaf concentrations, the averaged values per

tree were first divided by the 1997–1998/2011 leaflet-to-rachis ratios to have indirect rachis

estimates, and then the 4:1 leaflet-to-rachis mass ratio was applied. This assumed that the

ratios from the 1997–1998 and 2011 studies could be used here for 2002 (which lay between

them). For the complete five-element data sets and direct estimation of leaf concentrations in

Table 2 (viz. 1991–1992 and 2011), years 1-3 after masting, N, P, K and Mg all differed

strongly and significantly (F = 13.4, 12.2, 27.2 and 6.01, all $P < 0.01$) but not Ca ($F = 1.0$, P

> 0.05).

The results show that in the inter-mast period, N was low for the first two dry seasons and

then rose the year before the next masting, P increased from low to high approximately

linearly in the period, Ca barely changed, K and Mg were relatively high for the first two

seasons but fell in the third before masting (Table 2). This suggested that N and especially P

concentrations were depleted by the last mast fruiting and were re-instated before the next

event. Both K and Mg increased earlier in the interval but were presumably deployed

elsewhere before masting. Despite the different sampling approaches, standard errors of

means, across years per element, were quite similar. The concentration within the double-

masting period 1997–1998 intriguingly lay between the 1991 and 1992 ones for N, but were

higher than both for P. That P was accumulating before masting is clearly demonstrated. The

rise in P at the end of the litter fall series, but not of N, in 1998 is also remarkable (Table 2).

The 1998 litter was collected immediately before a masting like the 1992 one, so they could

be considered comparable in this respect even though the previous mast fruiting events were

1 (unusually) and 3 years prior respectively. Averaging where possible values in Table 2 for

2011 and 2002, and separately those for 1992 and 1998, gives composite estimates of leaf

concentration respectively immediately post and prior to masting as: N: 16.73 and18.80

(+12.4%); P: 0.615 and 0.825 (+33.9%); K, 5.00 and 2.44 (–51.1%); Mg: 3.42 and 2.53 (–

26.0%). These inter-mast changes suggested that N and P were depleted by fruiting and

restored for the next event – proportionally three-fold more for P than N, but the opposite

occurred for K and Mg.

Reproductive allocation of N, P, Mg and Ca was in the order of 8% to 16% (Table 3),

but for K it was considerably higher at 37%. This last value can be traced back to the

relatively high concentration of K in *M. bisulcata* pods, slightly higher than in leaves. The

fruit to leaf mass ratio reached almost 0.6 for K compared with < 0.2 for the other elements.

These calculations do not include root allocation, however, because estimates for them are

lacking, in terms of both root growth rate per year and root concentrations. Fruiting in *M.*

bisulcata is much costlier in terms of K than of P. *Microberlinia bisulcata* leaf litterfall rates

in the 12-month July-to-June periods of 1990–1991 (non-masting in 1991) and 1991–92

(masting in 1992) – found by integrating the areas under curves of litter fall versus time in

Fig. 2d of Chuyong et al. (2000) – were practically the same (1710 and 1670 kg ha^{-1} yr^{-1},

respectively). Multiplying these values by their corresponding P concentrations (Table 2), led

to a difference of 0.168 kg P ha^{-1} yr^{-1}, which is very close to 50% of the investment in

masting in 1998 (Table 3), assuming that the mast fruiting that year was similar in intensity to

the one in 1992.

1048

DISCUSSION

Mast fruiting as a non-stationary process

If masting is driven by the availability of resources, and these are partly stochastic and

partly trending over time, then a range of interval lengths might be accommodated for *M.*

bisulcata. In reality this range is limited to 1 to 3 years. The extended time series showed that

between 1989 and 1995 there was a 3-year cycle, but after 1995, as radiation increased, there

came a double year 1997-98 with high fruiting – 1998 somewhat weaker than 1997, and then

the cycle dropped to 2 years with a masting sequence 2000, 2002, and 2004, returning to very

strong masting in 2007 and 2010 on a 3-year cycle, and finally after that the control

seemingly weakened and resulted in two double-years, 2013-14 and 2016-17, which showed

only moderate mast fruiting. Apart from being quite short at 29 years, the time series was

non-stationary. There would be little reason to assume that it would *a priori* be stationary,

especially, given the late transient cycle stage of the stand (Newbery et al. 2013) and the

varying climatic variables on all time scales ranging from years to decades and centuries

(Nicholson et al. 2000; Nicholson et al. 2018). The inconstancy might be best understood in a

novel way with a two- or three-factor threshold model, where C, P and K reach potentially

different thresholds, simultaneously or sequentially. Thresholds themselves may not even be

constant over time but relate to age, climate trends, and other tree-internal demands. When

one of the resources reaches its threshold early, its effectiveness could be limited by the

slower accumulation of another resource. The first resource will need to be stored or

allocated elsewhere in the tree, even to then support the acquisition of the lagging resource.

In time the second resource accumulates to a level that crosses a pre-set threshold, and this

allows the first resource to come into operation.

The allocations of C to mast fruiting in *M. bisulcata* are substantial. In the earlier mast

fruiting event of 1995, 28 out of 30 trees recorded had fruits. The average number of seeds

per tree was ~26K (maximum 92K). Using estimates of mean dry masses of seeds and pods,

and mean numbers of seeds per pod, the total dry mass investment was 17 and 307 kg per tree

respectively, summing to 324 kg per tree (Green and Newbery 2002). These per-tree masses

translate to 53 plus 981 = 1034 kg/ha, ~55% the mass of average annual leaf-fall (Green and

Newbery 2002). Applying the individual seed and pod masses to counts made for the 2007

and 2010 events, for trees that had pods (Norghauer and Newbery 2015), gave 181 kg per tree

in 2007 and 122 kg per tree in 2010. (Excluding the two zero-producers, output values for

1995 were 346 kg/tree.) On this basis the investments per masting tree in 2007 and 2010 were

52% and 35% respectively of that in 1995, illustrating an up to almost two-fold variation in

masting intensity across these full fruiting events. Other more semi-quantitative scorings of

masting across the rest of the time series suggest a similar range in reproductive output.

Flowering is a precondition to fruiting, but the dependence is not exact. Heavy flowering was recorded in 1992, a mast year; but in 1991 there was also especially heavy flowering but no mast fruiting (Chuyong 1994; Newbery et al. 1998). In 1990 there was some low fruiting (flowering unmonitored), this coming though in the year after a main mast fruiting in 1989: further, 1993 had very little flowering and consequently almost no fruiting, and 1994 relatively high flowering but only scattered individuals fruiting (Green and Newbery 2001a, 2002). From the later, more detailed, phenology recordings, 1995, 1997, 1998 and 2000 had strong flowering and were masting years, whilst 1996 and 1999 flowering was practically absent and had no masting (Newbery et al. 2006a). Whilst valvate bracteoles protect flower buds before anthesis, it is possible that the intense occasional shower could damage flowers enough to interfere with pollination. This is the surmise for 1991, two years after the 1989 masting: fruiting was probably checked and masting was delayed to 1992. In 2004, the dry season started 43 days after 1 December 2003, on 12 January 2004 (Newbery et al. 2006a): recall that the dry season for a given year starts in December the year prior. Flowering peaked over the week before and after that date. But in 2012, the season started 21 days in, on 22 December 2011, which was 1 month before flowering started in 2012, and ended another month later. (In both 2004 and 2012 there were no heavy showers that could have disturbed pollination in the first halves of those dry seasons.) These facets indicate that, firstly, occasional external and unpredictable influences can alter the processes behind a masting cycle; and secondly, the timing of flowering (also linked to the timing of leaf-exchange) can change by up to a month with respect to the start of the dry season. If internal tree nutrient levels do, via deterministic physiological mechanisms, trigger the flowering – which leads most often to fruiting at some level of strength (Newbery et al. 2006a), masting remains also subject to climatic stochasticity causing interference to pollination (may be once in 10–15 years) and variability in dry season timing and intensity (every year).

Abortion of small immature green pods falling to the ground was not seen in the 29-

year period. It remains possible that outside of the three more intensively recorded periods

this phenomenon was missed. If in occasional years there was strong flowering, but no mast

fruiting followed, then usually those trees that did develop any pods carried them through to

maturity. The control of masting therefore most likely happens before flowering (see

(Newbery et al. 2006a) for further discussion). After that seed failure is simply a chance

unpredictable cost within the evolved life history schedule.

Nutrient resource interdependencies

Analysis of the seed and pod nutrient concentrations and amounts in relation to those

in mature leaves indicated ratios of between ~ 0.1 and 0.2 for N, P, Mg and Ca, but

substantially higher at 0.6 for K. This difference is very largely due to the low concentrations

of the first four elements, and a much higher one for K in the pods. Pods collected were

recently fallen. Caesalp pods are hard and rather water-repellent for several weeks, so it is

very unlikely they adsorbed much K coming down in canopy throughfall (G. Neba, D. M.

Newbery and G. B. Chuyong, in prep.), or were they leached to any extent. Had that been a

cause of an increase, then Mg should have increased too, yet it did not. Presently it is difficult

though to explain why so much K was being invested in pods. The comparative leaf K values,

coming admittedly from a retranslocation study where replication of bulked samples per date

was small, do closely match those of mature leaves of plantation-grown *M. bisulcata* trees

(averaging ~ 5.8 mg g$^{-1;}$ at ages of 13 to 18 years; D. M. Newbery unpubl. data). Whilst

ectomycorrhizas are strongly linked to P acquisition and uptake in the P-poor soils, K is also

very low in availability at Korup (Gartlan et al. 1986), and the surface-soil root mat (with

dense mycorrhizal networks) is an efficient collector of K (and Mg) coming down in

throughfall (G. A. Neba, D. M. Newbery and G. B. Chuyong, unpubl. data). In plots at the

same site, Chuyong (1994) found N, P, K, Mg and Ca concentrations of 13.8, 2.09, 6.41, 2.22

and 3.50 mg g^{-1} respectively in *M. bisulcata* seeds that fell in 1992. Pods were not separated

in the 1992 general collections but included within 'all reproductive parts' in the HEM plots,

those with high abundance of *M. bisulcata* had a mean K concentration of 6.04 mg g^{-1}.

Although other elements in fruits were albeit a little higher than for *M. bisulcata* alone in

1998, the earlier data confirm the K values of interest well. Investing P in seeds would

presumably be important to the survival of establishing seedlings up until the time they could

become infected with ectomycorrhizas and link into the mycelial network. Could it be that

also K, not P alone – as for some time thought (Newbery et al. 1988; Newbery et al. 2002), is

a second key controlling element of masting in the Korup groves? Several indications suggest

that P and K might act synergistically.

The main pattern explaining the masting time series was the sequence of relatively

wet dry season followed by relatively dry one. There was no evidence of water limitation to

large trees in the dry seasons because of their ability to root deeply (Newbery et al. 1997;

Newbery et al. 2009), so this points to a soil or nutrient factor governed by soil moisture

content in the upper soil layer, where fine roots and ectomycorrhizas form a root-mat and are

most abundant and active in nutrient cycling. What is it about relative dryness and wetness of

the dry months that leads to these relationships? Possibly dryness affected the fine roots and

ectomycorrhizas. The drier the dry season, then likely the more restricted the root activity

under limiting surface soil water conditions; which would possibly signal stored P in the

ectomycorrhizas to be moved to intermediate locations ready for fruiting, and less would be

used in maintaining new root growth. In a relatively wetter dry season this root activity would

continue longer and more effectively, so that any additional P stored in the roots would stay

there and not be signaled away to fruiting. Nevertheless, the main mineralization and uptake

occurs in early weeks of the wet season (Chuyong et al. 2002), and this would be similar each

year once the soil water content was high enough. This dryness is thought to be a proxy for a

mechanism that triggers translocation, but it is one that cannot operate without the previous

year acquiring more P and setting up a form of 'relay'. It is interesting that moderate 'm'

events occur in double-years and twice later, mirroring the two strong 'M' years 1997–1998.

At Korup in the HEM plots, sand, silt and clay proportions were 77%, 9% and 14%

respectively (Newbery et al. 1997). There exist very few published water content vs water

potential curves for sandy loam tropical soils, but at a site in NE Brazil with a very similar

physical composition to Korup, Fisher et al. (2008) found that a change in SMC (at 30 cm

depth) from 20% to 12% (cf. the wet-to-dry-season change in Fig. 3) led to a decrease in soil

water potential (SMP) from –10 to –100 kPa – which is very much higher than the generally

accepted value of –1500 kPa where wilting starts. Such a translation across sites should

however be treated cautiously because any SMC-SMP relationship is finely dependent on the

exact physical composition of the soil (Or and Wraith 2002). Wilting of understorey treelets

and shrubs, even in the strongest dry seasons, was very rarely observed at Korup.

Nevertheless, the drying of the top surface soil with the root mat might still be expected to

affect nutrient uptake capabilities in the dry season. In those dry months the surface leaf litter

is definitely dried out, and in many places the sandy-loam soil just below the organic layer is

visible, leaving the root mat exposed. The implied sensitivity of fine roots and

ectomycorrhizas of *M. bisulcata* to surface soil water conditions might in part result from this

species very pronounced buttressing and an extensive lateral root system lying just below the

surface soil mat (Newbery et al. 2009).

A cycle of 3 years would most likely allow a restocking of P, not only enabling

growth of other tree parts, but also storage of P in wood rays and roots ready for fruiting (as

the model later explains). A 4-year cycle could arise but, *ex hypothesis*, that would need a

sequence of two dry dry-seasons: none was recorded, however. The regression modelling

gave an indication of year-to-year increases in radiation, and positive deviations from the

smoothed trend of radiation with time. Changes were relatively small, radiation varied rather

little from year to year, both for the dry seasons and the years as a whole. The additional

radiation probably did not contribute to much increase in photosynthesis which would cater

completely for the extra demand in years of masting, above the potential competition for C by

leaves, stems and roots. Variation also in start and duration of the dry season, with the fairly

steady daily radiation inputs, meant that there was probably sufficient radiation supplying C

on top of tree storage, to meet the high dry mass investment in fruits. Carbon supply was

therefore likely not the limiting factor to masting. One way to resolve this possible

contradiction is to propose that, internal to the tree, there was strong temporal division in the

allocation of C between rooting (one year, wetter) and fruiting (the next, drier). Further, the

lack of a relationship between proportion of crown fruiting and tree size may be further

evidence that C is not the predominant limiting factor for mast fruiting – contrary to what was

originally suggested by Green and Newbery (2002), if it can be assumed that large trees store

more NSCs than small ones. The lack of year-to-year differences in oil palm yield relating to

radiation is also moderate evidence of C from photosynthesis not being so critical for their

yielding either. Palms can store up to ~20% of their C in stems from year to year (Legros et

al. 2009; Corley and Tinker 2016). The lack of correspondence between oil palm yield and

M. bisulcata fruiting suggests no common climatic driver and points indirectly to nutrients as

the key factor in the forest, though possibly not P alone since palms did not respond to P

fertilizer.

The very few studies on C and nutrients involved in mast fruiting of dipterocarps that

have been made largely support the thesis of this paper. Defoliation and girdling experiments

on branches of *Dryobalanops aromatica* trees by Ichie et al. (2005b) indicated that flowering

drew on C stored in local branches and the filling of fruits came from nearby leaves. This

confirms the many similar conclusions for temperate trees mentioned in the Introduction.

Relative allocation of P to fruits was shown to be higher than of C and N for of another

masting dipterocarp, *Dipterocarpus tempehes* (Ichie et al. 2005a), again emphasizing the

importance of P in reproduction, especially on P-poor tropical sites. During a different

masting event for *D. aromatica*, Ichie and Nakagawa (2013) found that P concentrations in

branch, stem and roots steadily decreased during flowering and fruiting (but not in twigs and

leaves), N fluctuated slightly (in all parts) and, whilst K decreased during flowering, it

increased during fruiting (in all parts). Of the P needed for fruiting, 68% was estimated to

come from branch storage, compared with only 20% for N and very little for K. This was the

first time root-stem-branch stored P was shown to be important for masting, and the result

corroborated the hypotheses of Henkel et al. (2005) and Newbery et al. (2006a). A further

study at the forest stand level provided more support for P and K both being key nutrients in

mast fruiting. Aoyagi et al. (2018) followed nutrient fluxes in soil, stems and fine litter in a

lowland dipterocarp forest during one such event and for 4 years afterwards. Fluxes of P and

K were much higher in total fine litter in the mast year than later years, whilst those of N, Ca

and Mg altered little; the higher P and K was coming from the high allocations to fruits.

Interestingly, P and K concentrations in coarse roots – but not stems – of the most abundant

dipterocarp, *Shorea multiflora*, decreased more in flowering than non-flowering trees. This

might indicate root storage of P and K ahead of stem and branch storage, in preparation for

fruiting. Kitayama et al. (2015) have more generally demonstrated in NE Borneo that across a

P-use efficiency gradient, forest trees allocate proportionally much more P than C or N to

reproduction (in relation to total allocation), even at P-poor sites.

Masting and carbon supply

The internal distribution of C between potentially competing sinks within each tree

has a further interesting aspect. A majority of the *M. bisulcata* trees in the grove are presently

of large to very large ($\geq$ 50 cm stem diameter) sizes which means they likely have

considerable respiration and maintenance costs. Carbon at least would not be so readily

available in an average year to sustain such a heavy fruiting without causing deficits in

growth elsewhere in the tree, perhaps in fine root and ectomycorrhizal production. This

would lead, hypothetically, to an even stronger competition between rooting and fruiting,

once stem growth is partitioned aside. That seedlings and saplings survive very poorly close

to adults, led to the proposal of a reverse drain operating towards the old adults especially in

mast fruiting years, for P as well as for C (Newbery et al. 1998; Newbery et al. 2006b;

Newbery et al. 2010; Norghauer and Newbery 2016).

Previous estimates of leaf litterfall were 1880 and 1380 kg ha^{-1} yr^{-1} in non-masting

1990 and masting 1989 respectively (the first estimate is used in Table 3), and came from

Green and Newbery (Green and Newbery 2002), who in turn based their calculations on

January-to-December, plot-level, leaf litterfall estimates given in Newbery et al. (1997). The

1990–1992 data later indicated 39% of that litter was from *M. bisulcata* trees. This 0.74:1

ratio may have been an overestimate, because it used calendar year estimates which would

have missed linking November-December litterfall with that in January to April prior to

masting, and the newer 1990–1991 species-specific data were more precise. Or it simply

reflects variation in litterfall, viz. 1380, 1880, 1710 and 1670 kg/ha over the four years 1989

through 1992, which suggests that C supply was likely again not to have played an important

role in determining mast fruiting since in around one masting event there was an implied cost

to leafing and around the other none. The fruit-to-leaf amount ratios and reproductive

allocations percentages in Table 3 may well be ~10% lower than the true values, if the 1990–

1992 litterfall estimates did apply. The 1990–1992 data also display a small shoulder in May

1992, possibly the result of the last older leaves being exchanged for the new green pods.

Green pods stay on the trees for more than 5 months and therefore will contribute C to

their structures and to developing seeds. This is likely an important source, and pods and

seeds may indeed be almost autonomous in this respect. Pods are also held up high on the

outer canopy edges where their photosynthesis is maximized (Norghauer and Newbery 2015).

Bennett et al. (2011) reviewed the role of pods in their photosynthetic capacity, and Zhang et

al. (2017) provide a detail analysis of C balance in the pods of *Medicago sativa* to show the

considerable C inputs that are possible. This source of C would make the *M. bisulcata* trees at

masting even more independent of C stores within the rest of the tree. It could explain the

effect of raised radiation in the current fruiting year when compared with the one prior.

That stem increment growth was positively related to current year radiation, but mast

fruiting was not always, or at least not so strongly, suggests that C from photosynthesis was

being used for stem growth and in mast years it was additionally contributing to mast fruiting.

An internal trade-off between root growth plus ectomycorrhizal activity to enable the uptake

of nutrients allowing nutrient uptake, and an allocation to fruits and seeds when cued, may

probably well be the basis to the masting cycle. In the year following a masting, relatively

more C would be allocated to roots because, if fruiting is 'switched off', roots become the

main alternative sink; and then if in the second year there is sufficient P to cross the

threshold, with some increase in radiation and extra C input, a next masting would occur after

2 years. If P accumulation was limited the tree would wait until the next year, forming a 3-

year cycle. Extra C available in that second interval year might then determine the intensity

of the masting ('M' or 'm'). A notable difference between P and K, is the rate of change in

concentration in the leaf litter as the interval between mast fruiting progresses: P increased

slowly over 3 years, but K rose rapidly then fell lower than at the start. This hints at faster

recycling of K than P, with a stronger withdrawal of K than P at senescence later as storage

cells were being refilled prior to masting.

Rainfall and nutrient uptake

The effect of mean daily rainfall in the dry season is dependent on the pattern of rainfall distribution over time, through the dry season and just into the early wet season weeks, which is up until the 100 mm threshold defining the season is exceeded. The first week's rainfall governs root growth and ectomycorrhizal physiology, and the later ones connect to the early wet season and the throughfall which primes the start period of optimal leaf litter decomposition (Chuyong et al. 2002, 2004). A dry season may well limit root activity, and it also would delay decomposition: one which was short would only affect the former. On the other hand, a relative wet dry season will be expected to restrict rooting and ectomycorrhizal activity much less and encourage the earlier start of decomposition, both processes leading to better nutrient uptake into the trees. Ectomycorrhizal activity doubtless plays a role in P uptake, and the regrowth and maintenance of the root mat itself determines the level of retention of K leached from the canopy and its uptake.

An intriguing aspect in connection with the P-supply for fruiting is how P gets taken up and stored in inter-mast years. In the dry season of 1988–1989 and 1990–1991 there were pronounced peaks of P in fallen leaf litter in the HEM but not LEM plots (Newbery et al. 1997). This peak was absent in 1989–1990, directly after the heavy mast fruiting. The HEM litter would have been composed of ~80% *M. bisulcata* leaves (Chuyong et al. 2000). In a relatively dry dry-season with likely more die back of fine roots and ectomycorrhizas than in a wetter one, P might be translocated out of these fine roots and ectomycorrhizas, going into storage for fruiting. Altogether, the integrated aspects of this P nutrient cycle do provide reasonably strong support for P storage and use leading up to, and during, the masting events as proposed by the original PACER hypothesis.

Another result providing important evidence for P-storage is from the comparison between concentrations and amounts in the years across masting intervals which suggests further that the 50% of P used in masting was coming from retranslocation of P out of

senescing leaves a few months before. This P was probably being stored in branches. By

difference it might be deduced that the other 50% was older and coming from local

branch/wood storage built up in the prior 1 or 2 years, still within the interval. That result

would fit with the notion of a temporary store for P and not all of the P needed for masting

could be supplied in the current year alone. Rosell et al. (2021) in a study of a range of

tropical tree species found the inner bark of branches was especially important for NSC

storage and, although not yet tested, this may be where P is also held, ready for fruiting.

Jones et al. (2019) indicate this as a possibility for premontane tropical forests.

If P in leaf litter is relatively low in concentration after allocation to fruiting (Newbery

et al. 1997; and new evidence in this paper), then the fast-forward cycling proposed by

Chuyong et al. (2000) could be temporarily slowed because mineralization by saprophytic

fungi would have a low-litter resource to decompose. Root activity would either need to

become more intense and efficient to take up the now lower soil levels, or at this time the

ectomycorrhizal activity would come more to the fore by not only accessing low molecular-

weight organic P molecules efficiently but also by effectively reaching low-concentration

inorganic P sources further way from roots. This impasse would strongly select for the

several co-adapted root mat and ectomycorrhizal traits seen in the field in order to achieve

restocking of P. The rise in P accumulated just before masting could lead to higher P in litter

falling that year which on one hand would mean fast re-uptake but on the other appears to be

a dissipative risk when the perceived strategy is to save P (in the tree) for immediate use in

fruiting. However, the ectomycorrhizas being part of the whole tree system, and cycling

highly efficient, P is being 'saved dynamically' within the nutrient cycle. Also, N and K

would in parallel complement C in stores – the latter benefitting from the small increase due

to radiation enhanced photosynthesis, all three also needed for seeds and pods. That P was

high in falling litter just prior to masting might furthermore indicate that P stores were quite

full after a 2-year interval, and some of that 'extra' P was also coming from root

retranslocation.

The analysis of mast fruiting in 1989–2004 (Newbery et al. 2006a) used 150 trees

across the whole P-plot, but the later recording 2009–2014 used a subset of 61 trees in the

eastern half, an area 100–500 N by 1000–1200 E, on the higher ground with denser stands

and larger trees. The masting time series could potentially have been confounded by sampling

extent changing over time, because for the earlier half (1989–2004) it was found that masting

was significantly stronger on the relatively higher and drier sites of this eastern part of the P-

plot compared with the rest of the plot (Newbery et al. 2006a). Allaying these doubts, the full

and strong masting in the second half (2007 and 2010) were confirmed by seed trapping

across the eastern 25 ha of the plot (Norghauer and Newbery 2015). The two double years, in

the second half of the time series, were only moderate masting despite being at the more

favorable topographic location.

The very strong masting achieved by *Dicymbe corymbosa* in Guyana, with 3–4 t dry

pod and seed mass per ha, were associated with pronounced El Nino Southern Oscillation

(ENSO) peaks in radiation (Henkel et al. 2005; Henkel and Mayor 2019). However, this

ectomycorrhizal detarioid, although phylogenetically very close to *M. bisulcata*, is unusual in

that it reproduces annually also by sprouting, which must impose a considerable demand on

C and nutrients for growth. For this reason, it might be conjectured that C stores are probably

being continually used up and hence, to achieve a mast fruiting, a dependence on an

additional boost of incoming radiation in those ENSO years would be necessary. Again, in

contrast to most African detarioids (including *M. bisulcata*), *Gilbertiodendron dewevrei* –

which forms monodominant stands in Central Africa – exhibits mast fruiting which is

spatially unsynchronized and at irregular intervals (Hart 1995). Its individual seeds are very

heavy (~30 g), so the mass allocated to reproduction per tree is probably far greater than for

M. bisulcata. Compared with SE Asia and C. and northern S. America, the lack of a strong

ENSO signal in C. Africa is pertinent (Poccard et al. 2000; Camberlin et al. 2001). Masting in

dipterocarps may well be for different suite of factors from those controlling the detarioids,

achieving an evolutionary stable strategy along an alternative channel.

Interference effect of herbivores

The loss of a 2- or 3-year cycle in masting after 2011 suggests that the rhythm of

wet/dry dry-season release of nutrients needed for full masting was disrupted by a

combination of lack of an alternating pattern in rainfall plus heavy caterpillar attacks. This

latter unforeseen intervention, combined with an unusual climate regularity over 3 years,

acted as a 'quasi-treatment' in that C gain and storage were likely temporarily reduced. From

the above argumentation it was not so much the decrease in C to supporting fruiting that was

alone important but also the loss of C input for roots to allow P to be taken up and

accumulated even in the relatively wet dry seasons. With some commitment to flowering

already, leaf grazing would have nullified any C input from the local branches attacked to

fruiting, requiring the tree to draw on stores in the branches. Nevertheless, some fruits

resulted, indicating a means by which tolerance to herbivory could have occurred (Norghauer

et al. 2023). The caterpillar attacks may have interfered with the accumulation of P and

thereby interrupted masting in full, leading to the unusual double 'm' years. The evidence of

an herbivore effect is strongest for fruiting in 2013–2014 because of the detailed phenology

data in 2011–2012. It is tempting to suggest that something similar might have been

happening in 2015 before the very similar masting pattern of 2016–2017: alas, herbivore and

tree phenology records were not possible during those later 3 years. A further unknown

aspect is how herbivory may have induced stress hormones in the trees which, interacting

perhaps with K levels, affected cytokinin concentrations.

The pooled field notes and observations suggest two lepidopteran herbivores occur in the crowns of co-dominant *M. bisulcata* and *T. korupensis* in the natural old-growth forest of Korup. Both species have pinnate leaves – *T. bifoliolata* does not – and similar stem densities at ~ 3.5 per ha for ≥ 50 cm DBH (Newbery et al. 1998). But *M. bisulcata* trees are the greatest in size with a near-zero mortality rate over recent decades (Newbery et al. 2013). If caterpillars indeed fed only on expanding leaves (or flowers), then severe damage could ensue quickly, within 1 to 2 weeks, inducing premature abscission (or lost pollination). For example, within 1 week caterpillars defoliated young leaves in crowns of two monodominant leguminous tree species in rain forests of Brazil and Gabon (Nascimento and Proctor 1994; Maisels 2004). Such phenomena are quite easy to miss though. This could explain why the black caterpillar was not noticed before at Korup, but it is unlikely the same reason for the yellow-caterpillar given the extensive multi-decadal phenological monitoring of *M. bisulcata* (Newbery et al. 1998; Newbery et al. 2006a; Norghauer et al. 2023). The detailed monitoring between 2009 and 2013, like that between 1995 and 2000, did not note any caterpillar attacks. To have an importance they would surely have been seen before from their feeding effects (droppings, noise of chewing, fallen individuals, hanging threads, frass in traps). Many other periods of intensive work back to 1984, focused on the dry months and early wet season when caterpillars would occur, did not record them. Whether these outbreaks are triggered by favourable climate factors, or palatable leaf chemistry or less parasitoid pressure, either perhaps induced by drier conditions (Van Bael et al. 2004) or simply mounting abundance over years (Wong et al. 1990), is unknown. Herbivore attacks to flowers, occurring stochastically, might be viewed as being similar in end effect to intense rainfall events disrupting pollination: both limit reproduction.

Refined updated hypothesis for mast fruiting in caesalps

The mast fruiting cycle in *M. bisulcata* could thus be explained in terms of the

availability of three principal resources which govern the timing of reproductive output: the

limiting resource P, water as a controlling resource, and C as a qualifying resource. If

sufficient P accumulates within 2 years to trigger flowering and there is sufficient C coming

from storage laid down over the last year or supplemented due to a positive radiation

difference between prior to current year, then fruiting is to be expected. But when P is slower

at accumulating, the interval is extended by one year and fruiting happens within 3 years. It is

highly likely that in the latter case C would still be sufficient either from reserves laid down

in the tree in the previous year and/or a small surplus the current year of fruiting. When there

is sufficient P accumulated after one year but not quite enough C for a full masting, fruiting

may follow anyway at a moderate level; and in doing so it is likely that not all of the

accumulated P would have been allocated 'as planned', and the part remaining, plus last

year's continuing uptake, will sustain a second moderate fruiting. According to these resource

requirements, it is difficult to see how the pattern could be 'm-M' or 'M-m', and neither of

these were observed: in the first, there would be insufficient time after a 'm' fruiting to build

up P so rapidly to have an 'M' one, and in the second, the 'M' first would exhaust P levels

down below a threshold and no 'm' would likely be possible either. The pattern M-M did

occur once, however, at the time of an exceptional rise in radiation levels over a 3–4-year

period: it can only be hypothesized that there was unusually even more surplus C for extra

root growth and P uptake in both the year preceding the first 'M' year and during the second

one. The main processes leading to mast fruiting in *M. bisulcata* are schematically shown in

Fig. 5.

An interesting aspect is the feedback control that might operate by way of C going

either into fine roots and ectomycorrhizas, which would enhance nutrient uptake, or into

wood storage (stem, branch and large root), which could sustain a next fruiting. The 2- or 3-

year cycle is simply a consequence of the contingent conditions of these trees growing in

low-P soils aided by ectomycorrhizas. Rainfall and radiation are stochastic processes, to some

extent inversely related, which overlies the intrinsic deterministic cycle: the important

difference between them is that rain, acting primarily on the soil, facilitates P release,

acquisition and uptake (needed to accumulate P), and whilst C is needed also for fruiting, it is

essential for root production and maintenance of ectomycorrhizas that makes this P uptake

possible. That root growth should always precedes fruiting in a masting year, the first in late-

dry into early wet season (March–April), the latter in late wet season (September–October)

decides the sequence unambiguously: C in the dry season will go to the root sink first and

what remains stored, used for stem growth or reserved for fruiting when it happens. The low

P supply is therefore the main determinant: C must be sufficient but does not appear to be

decisive for the interval length. Although N and K will also be involved the evidence more

strongly supports the P thesis. Nitrogen is not limiting at Korup with ample N-fixation and

aerial input (Newbery et al. 1997), and whilst K follows P in being intrinsically quite low in

supply in Korup soils, the root-mat appears to be highly adapted to catching and retaining

canopy-leached K and so enabling its efficient recycling (G. A. Neba, D. M. Newbery and G.

B. Chuyong, unpubl. data). The K allocated to pods could, like N, be accommodated by wood

ray storage.

Carbon and N will be important for masting, even when P and K are limiting and are

the key storage resources, because all elements will be needed in their stoichiometric ratios in

order to fill pods with seeds. As in other plant physiological problems, simplified reductionist

approaches do not recognize the integrated nature of growth and reproduction. Carbon

(sugars) and K are required for transport and storage, while P is bound into C-H-O structures

which act as 'carriers'. It is possible though that a small increase in C from radiation might

have a non-linearly disproportional benefit to P transport and storage. Understanding masting

physiology requires an integrated and dynamic approach, one of shifting sink strengths over time. After a last masting, roots would likely act as the strongest sink, then the stems (and rays and bark) and branches – part growth, part storage – and finally reproductive parts. No abrupt switches in C, in contrast to a threshold switch for P, but a gradual altering of the components over a 2–3-year cycle occurs. For this to operate it is necessary to hypothesize feedback from the C-depleted branches to the more C-depleted roots. With the fruiting sink removed, C should move from stems to roots. Whether this needs additional hormonal control is an interesting question. It remains a conundrum as to why 'in the background' to the rooting-fruiting axis, stem wood growth is positively correlated in a quasi-independent way with radiation. Stems may be working perhaps as the storage stabilizers, being placed centrally between shoots and roots. As they grow, they store and balance the terminal demands. The levelling adjustment of C between the first and second interval years could perhaps explain the lack of clear correlation between fruiting (and inversely rooting) with stem growth increment. The time sequence within the year is also important here. In the second (drier) year of fruiting, C is not, *ex hypothesis* predominantly going to roots in the early wet season but more to stems for storage, and en route there for filling fruits later in the wet season (Fig. 5).

    If radiation were to increase during an increasing part of a cycle or as a longer-term trend, and C were alone the limiting resource for masting, then fruiting would come closer together and this would defeat the putative economy-of-scale selection. In this respect a non-stationary time series offer advantages when matched to an also non-constant climate environment. On this basis it would seem also unlikely that C could be the key resource for *M. bisulcata*. Further, C available per tree will decrease with tree age after the point of maximum crown extension because of the higher respiratory loads incurred in maintenance of the whole tree. This may be connected with the observed crown thinning of the largest trees

(J. M. Norghauer, unpubl. data). When seed production was substantial, in proportion to the

high flowering per crown then there would be a deficit of C to fill pods, or pod production

would decrease, or another cost incurred (e.g. at the roots or wood production). But the

evidence was that in *M. bisulcata* the relative crown-cover with pods remained constant with

size and, by broad inference, age. As a tree grew to have a larger root system with

ectomycorrhizas to maintain, to acquire P and retain K, when absolute pod and seed output

rose then the demand for P and K would have also. As with C and N, we can expect that P

and K is stored in 'quick-release' forms in wood rays and bark of the stem and the main

branches. A phenomenon often remarked is of one part of a large crown flowering and

fruiting much more than the others, which could be explained by a correspondingly more

active root system of the tree, and a matching storage of nutrients in the wood and branches,

on that side: see Newbery et al. (2009) for the strong crown asymmetry in *M. bisulcata*.

The arguments going from nutrient concentrations in fallen litter through to the

possibly the trees building up and storing P and K before masting are tenuous because of the

many uncertainties about the processes involved. If some translocation occurs out of leaves

before they are shed, then why are the P levels so high just before masting? If P was being

actively stored for fruiting, it might be expected that it would be lower in concentration in the

fallen leaves. However, the high concentration reflects that all the trees were well stocked

with P by that time, the stores were full, or even over-full, and there was still plenty of P to

fall. From earlier work, it was established that ectomycorrhizal species, especially *M.*

bisulcata, drop P-richer leaf litter than non-ectomycorrhizal ones (Newbery et al. 1997) partly

because the retranslocation was less (Chuyong et al. 2000), and it was argued then that this

allows the decomposition process to operate without P limitation, the 'fast forward' strategy.

However, the concentration of K (to a lesser extent that of Mg also) reached a peak mid-

interval and then fell just before masting. This could mean that K was being translocated into

storage before fruiting. In general, the export of K and Mg from leaves of trees is usually low

compared with that of N and P. Understanding the differences in dynamics of P and K in this

context is highly challenging, and also explains why N showed no clear mast-interval trends.

Whilst K was likely building up on soil colloids in wetter dry seasons, mineralized from

fallen leaves, dead roots and old pods during decomposition, and that this process was more

limited in the drier dry seasons, suggests that K must also have been transported and actively

stored in the tree between the prior and mast years. To recycle more K via litterfall would

involve a potentially high loss due to leaching when the cation exchange capacity of the soil

has already been reached, compared with the seemingly tighter and 'safer' direct access by

ectomycorrhizas to organic P. This reliability aspect therefore favours P over K as the better

controlling factor in masting (Fig. 5).

Phosphorus storage and control

What could the likely form of storage of P in trees be? Since Bieleski (1973) the most

likely candidate is inositol hexa*kis* phosphate (= phytic acid; $C_6H_{18}O_{24}P_6$). It has been shown

many times to be the main storage form in seeds, and has been detected in stem and roots. It

forms salts with Ca, K, Mg, Mn and Zn, and in soils occurs commonly bound to Fe. It occurs

in 19 hexa-structures which are interconvertible (Raboy 2003). It is viewed as the most

frequent P-storage form when P is in excess, that is in passive storage; the P is readily

retrievable and mobilizable. Other molecules can have seven or eight atoms of P but seem

less common (Raboy 2003). Inositol hexa*kis* phosphate is involved in cell metabolism, in

ATP regeneration, RNA export and the repair of DNA. It is prominent in cell storage

biochemistry, which may make it a candidate for *active* storage. Yet so very little has been

researched in this area, and accounts for trees are very sparse. A notable exception is the

study of Kurita et al. (2017), showing P storage in *Populus alba*, followed by release in

spring, and indicating perhaps a common function in many deciduous tree species. It is

conceivable that the same mechanism could operate in seasonal tropical trees and even be a

basis to P storage for mast fruiting species. How inositol hexa*kis* phosphate is linked to

polyphosphate formation in roots and mycorrhizas and comes under hormonal control

remains unclear.

In an experiment with *Eucalyptus grandis* seedlings, comparing low and high P

media, Mulligan (1988) detected stored P in the form of esters and residual P (mostly

phosphorus proteins), especially in roots, and more so the low P medium. The idea is that

these stores buffer the plant (maybe actively stored), from there P can be mobilized and sent

to other stores in the shoots. In a similar vein, Netzer et al. (2017), comparing mature *Fagus*

sylvatica stands – one on high and one on very-low P available soils – found that in the latter

P built up in roots in late summer, and from there was moved on to stem and branches,

suggestive of at least two storage sites in a series enabling adaptation to the poor site.

However, hardly anything is known about how P stores are internally controlled, or how

hormones signal transfers of P (Rennenberg and Herschbach 2013), which could give even a

lead as to how P might be built up in stores in masting intervals, and once a threshold level is

met how the switch to flowering is signaled.

Much evidence suggests that senescence in leaves, and possibly pod formation, may

also involve cytokinins (Marschner 2012). Chiou and Lin (2011) have proposed different

hypothetical models for inorganic P being regulated by cytokinins and sugar levels. How

these promising models can be extended to storage and release of P has yet to be worked out.

The strongest candidate for the elusive 'florigen' is now widely held to be a flowering locus

T [FT-] protein, which again appears to be controlled by a cytokinin (Turnbull 2011).

Cytokinins act at long distances throughout the plant and are bidirectional in their effects,

both of which might hypothetically implicate them in inositol hexa*kis* phosphate metabolism

and P storage. It remains likely that floral initiation in *M. bisulcata* takes place at the end of

the wet season, before leaf exchange occurs in the dry season (see discussion in Newbery et

al. 2006a). Earlier results on cultivated apple (*Malus*) trees suggested already that a two-way

process of hormonal control could be operating (Davis 1957), and the idea that sugars and

other NSCs destined for the usually stronger root sink (Goldschmidt and Golomb 1982, for

mandarin (*Citrus*)), (Hansen et al. 1982; for prune (*Prunus*) on K-deficient soils) might be

diverted to the flowering meristem sinks under the control of gibberellins (Sachs 1977). How

exactly the 'switch' occurs remains elusive (Monselise and Goldschmidt 1982), and still to

this day. The precise consistency of alternate bearing in pistachio (*Pistacea vera*) orchards

has been questioned though, on statistical grounds (Rosenstock et al. 2010).

Active storage hypothesis

Storage of C as reserves in trees (stem rays, bark and branch wood) was widely seen

in the past as being passive. The C was being made available for regular annual growth

processes in ways that optimized the economy of the tree (Zimmerman and Brown 1971).

Extending this idea to nutrients, Chapin et al. (1990) produced a systematic table for N and P,

reserves of the latter mainly featuring as inorganic phosphate, polyphosphates, or

phospholipids, leading them to conclude that dependence on mineral nutrients is likely much

more pronounced than it is on C for reproduction. Sala et al. (2012a) argued that the build-up

of NSCs in stores in trees might be acting as a buffer against stochastic variation in C inputs

due to environmental stresses (e.g. drought, herbivore attacks), which as a bet-hedging

strategy would insure against temporary collapses in maintenance and growth. The storage

would need to be active, rather than passive, otherwise in the latter case it would be used up

immediately in extra current growth. Extending this idea to mineral nutrients, Dietze et al.

(2014) hypothesized that N, P and K might also be accumulated and stored to guard against

periods of reduced nutrient uptake, stabilizing the internal nutrient cycles. Hormones might

therefore control source-sink processes actively. The empirical evidence for active vs. passive

control of C storage in trees still remains weak and unresolved though (Hartmann and

Trumbore 2016). But where P and K especially are limiting growth, evolution of such an

active mechanism could be of considerable evolutionary advantage. Mast fruiting imposes a

quasi-unpredictable (partially stochastic) demand on mineral nutrients supplies, and therefore

for P and K active storage would serve it well, particularly in sites where uptake patterns

themselves are subject to environmental stochasticity (the intensity of the dry season at

Korup, for example).

Nutrients in alternate bearing of tree crops

Studies of alternate-bearing pistachio trees have provided several valuable insights

into how tree nutrient balance changes between 'on' and 'off' years. Whilst fruiting depresses

root growth in fruiting years, uptake rates from the soil remained similar. If N, P and K

moved from uptake mainly to leaves and shoots in 'off' years, and to fruits in 'on' ones at 3–

6-fold the investment compared with 'off'-year uptake, this implies that stores were being

drawn upon (Rosecrance et al. 1996). Trees stored 7- and 14-fold more N and P respectively

in 'off' than 'on' years, mostly in branches, which also suggests active storage (Rosecrance et

al. 1998). Of what went into fruits in an 'on' year, 70% of N and P came from storage, and

30% from retranslocation. For N and P, storage forms are well known (e.g. phosphoproteins),

but for K they are not; it would need to be intensively taken up in the 'on' years (study cited

here was made on well-fertilized plots, however), despite the slower root growth, transported

and used immediately in the fruits (Brown et al. 1995). Interestingly, K concentrations in

pistachio are inversely related to NSC concentrations, because K is needed for sugar transport

to the apices where fruits are being formed (Marschner 2012). If this last process applied to

M. bisulcata at Korup, then K would be taken up in the mast years and, assuming also it

cannot be stored in roots or branches, the only other feasible (temporary) store would be

outside the plant, this as exchangeable forms held on soil colloids, built up in the wetter non-

masting years (G. A. Neba, D. M. Newbery and G. B. Chuyong, unpubl. data). How far K

might be transported laterally in the mat by mass flow is unknown.

Picchioni et al. (1997) further showed, for pistachio again, that N and P were

translocated out of the mature pericarp into the seeds of the same fruits, although K was not,

which left the pericarp relative rich in K on fruit fall (harvesting). This could have been

happening in *M. bisulcata*, and in this way explains the K-rich pods. Possibly as pods

browned they too moved N and P to the seeds. With K having achieved its role in aiding

sugar transport (under the hypothetical hormonal control suggested), it would then be

redundant as regards reproductive investment *per se*: the fallen pods would be leached much

later by the end wet season rains, even some remaining until next year, and in this way

returning the extra K to the forest floor and soil colloid sites. Returning to the question of

possible CHO storage in 'off' years, direct estimates showed that very little of the stores laid

down in those 'off' years (8%) contributed to the 'on'-year yield (Stevenson and Shackel

1998) – confirming again the general finding mentioned above in the *Introduction*. Overall,

the main evidence is in favour of nutrients plus hormones being the controllers of alternate

bearing and, in extension, masting in forest tree species where nutrients are limiting.

Soil nutrient status

Nitrogen supply has been shown to be important for fruiting in *Nothofagus* (Smaill et

al. 2011). In either N-fertilized stands, or unfertilized ones when rainfall was high (allowing

higher N mineralization), masting was enhanced, subject to a temperature cue also operating.

Likewise, Leeper et al. (2020) showed in N-poor sites, that masting of spruce was less

frequent and less intense than at N-rich sites. Cone production was positively correlated with

soil N in the year prior, not the current one. Both studies found lagged N-effects, implying

that N was being stored. Nevertheless, unravelling the causes is difficult because the better,

wetter, conditions one year for N supply are not conducive to floral initiation, which depends

more on warm drier ones affording higher sugar transport to apices. An inherent inverse

correlation often exists which leads to an unstable alternation: a similar pattern is proposed

for *M. bisulcata* at Korup with the relatively wetter and drier dry seasons.

Evidence for depletion of nutrients during masting, needing then a restocking for the

next event, is unfortunately limited. For white-bark pines, Sala et al. (2012b) showed that P

and N declined in branches after masting – cones having especially high P concentrations;

and this could be linked to the species being ectomycorrhizal. Depletion spread over other

plant stores according to the amount needed for masting. Not only do Sala et al. (2012b)

provide evidence of resource switching (from storage to reproduction), but weather seems to

be cuing for masting by synchronized nutrient uptake being made possible. This supports the

PACER hypothesis and the main thesis of the current paper on *M. bisulcata*.

Climate factors which affect nutrient resource availability driving masting was a

central argument of the review by Allen et al. (2017). This underlying process is how

economies of scale become operational: the predisposition of trees to avoid seed predation or

increasing their flowering efficacy is enhanced by synchronization. Allen et al. (2017) made

strong recognition of the role of ectomycorrhizas, building on Henkel et al. (2005) and

Newbery et al. (2006a). That either N or P (rarely both) are often limiting where

ectomycorrhizas tree species are successfully dominant in forests, must place a high demand

on these elements for seed production. A set of resource acquisition and storage traits that

would benefit growth and defense in these nutrient-poor habitats generally would be expected

to be channel-selected through evolution to support an optimal life history strategy (Allen et

al. 2017). Different storage routes could be postulated to work in temperate vs. tropical

systems where respectively N and P are the more limiting. The Korup work posits that

resource synchronization might, in part, be achieved via a shared mycorrhizal network. It is a

working hypothesis that requires testing in the field at Korup. Given the lack of support for C

storage being important for mast fruiting, clearly the C budget model of Isagi et al. (1997)

needs to be rethought in terms P uptake, storage and usage.

Fernandez-Martinez et al. (2019) recognized that many mast fruiting species grow in

N- or P-poor soils, or at least on sites where the demand for these elements is high. The

authors relied on foliar N and P concentration data. There are two problems with this

approach: firstly, from among the masting species some samples may be from mast years,

others from inter-mast ones pulling the differences between masting and non-masting species

apart; and secondly, many species well adapted physiologically to nutrient poor soils do have

nutrient levels similar to those on richer soils but display different nutrient dynamics. Again,

the obvious association with, and role of, ectomycorrhizas was overlooked. Contrary to the

claim of Kelly (2020), nutrients can be hypothesized to determine masting when both their

availability fluctuates, and their tree nutrient concentrations are synchronized via the

mycorrhizal network. Stronger competition between masting and non-masting neighbours

need not be invoked, as was proposed by Fernandez-Martinez et al. (2020). It follows from

ecophysiological adaption of the masting species to their nutrient-poor conditions. From the

nutrient-supply model the other economy of scale factors can operate; in the case of *M.*

bisulcata this would probably be mainly to satiate seed predators (Norghauer and Newbery

2011).

Janzen (1974) asked why the intervals between mast fruiting in dipterocarps are so

large and irregular. That would not fit with resources like P accumulating to given common

levels. Resources could doubtless be stored like NSCs, but not stored only for fruiting, and

when not so used would be moved to other sinks such as roots. Dipterocarps form diverse

mixtures of species in many forests, with their individuals spaced out among those of other

ectomycorrhizal and non-ectomycorrhizal species. Thus, the functioning of a shared below-

ground network seems less feasible, especially when the fungal species differ across tree

species. However, Janzen (1974) interestingly raised the idea that masting may have

originated in species when they were much more clumped in the evolutionary past than they

are today, growing as patches of near single-species dominating stands. If nutrient resources

were more commonly distributed, shared and equilibrated within such patches in the manner

postulated for extant *M. bisulcata*, and over time the tendency to clump by some species

degenerated as they locally formed mixtures with other tree species, then the nutrient-

threshold mechanism behind masting may have lost its effectiveness.

Tree-tree root connections

Root grafts could potentially provide another means of nutrient transfer between *M.*

bisulcata trees within groves. Soil surface grafts have been observed at Korup although as yet

none anatomically investigated and mapped. Early field experiments on *Pinus strobilus* in

North America by Bormann (1966) point to this as a real possibility. It was suggested that the

process would mostly operate between large dominant trees of similar size to enable an

equilibration of resources across trees. Many temperate and tropical tree species form root

grafts (Larue 1952; Graham and Bormann 1966), often close to stems and among large roots,

but there is also evidence for grafts between roots at greater distances when the species has

long surface roots, e.g. for *Dacryodes excelsa* (Burseraceae) in Puerto Rica (Basnet et al.

1993), or among smaller roots well below the surface, e.g. in oaks (Saunier and Wagle 1965).

The extensively spreading root systems of *M. bisulcata*, connected to large buttresses

(Newbery et al. 2009), make the crossing of roots at 15–20 m away from stems quite feasible.

Because grafts form as a result of physical pressing together of roots, the restricted vertical

space of the root mat would likely also encourage grafts between medium to small roots, and

in this way create a large multi-nodal root network. Apart from possibly providing increased

resistance to wind throw – which would not be important for *M. bisulcata* given its strong

buttressing – and temporarily supporting suppressed, damaged or dying neighbours, other

advantages of root grafts have not been so obvious (Loehle and Jones 1990). However,

synchronizing nutrient supplies between, and storage within, trees across whole clusters

would clearly be beneficial and strengthen masting. Root grafts would moreover provide a

strong 'platform' for the ectomycorrhizal network and faster local transfers. A role for root

grafts enabling masting has recently been suggested by Noble et al. (2018) to partly explain

alternate bearing in pistachio in a year when pollination synchrony (the Moran effect) was

relatively weak. The focus in that plantation study was on C as the putative resource being

transferred.

More directly relevant to *M. bisulcata* was the 1953 work of Yli-Vakkuri, cited by

Bormann (1966), showing that P was moving across root grafts of *Pinus sylvestris*: the author

made no connection to ectomycorrhizal status of pines or association of ectomycorrhizas with

mast fruiting (see Newbery 2005). Conceivably, ectomycorrhizal and grafted root-networks

might operate together within *M. bisulcata* tree clusters, either contemporaneously or in

successive phases around the end of the dry into the early wet season. If hormones are also

transportable across grafts, an added level of synchronizing control is afforded. The role of

tree root interconnections (ectomycorrhizas and grafts) becomes an essential auxiliary to the

main masting hypothesis. Bormann (1966) advanced the thesis that root grafts were

established first by alignment of xylem tissues which rerouted water flow, and connecting

phloem tissues permitted organic and nutrient transfer later. Between the mid- and late-

ontogeny stages, *M. bisulcata* trees growing in dense patches become large adults, as

described by the transient dominance model (Newbery et al. 2013), and root grafting may

have started when water was at a premium in the dry season. It then changed to having a

more organic-flux transfer role once deep roots had secured all-year-round water supplies, the

root mat had fully formed, the litter nutrient cycle was stable, and P (and possibly K) were by

then in greater demand for growth and reproduction. To know whether or not codominants *Tetraberlinia bifoliolata* and *T. korupensis* graft below ground onto *M. bisulcata* would be interesting. That the three species do not usually mast synchronously suggests, though, that they probably do not form interspecific connections as predicted by Graham and Bormann (1966).

Role of environmental stochasticity

Dynamics simulation models involving storage and thresholds, resource-switching in a randomly varying environment, yet with no economy of scale operating, are able to generate masting cycles when either the variation is strong (i.e. very good vs poor years occur) or, when the variation is not so strong, there is an increasing non-linear response to the positive variations (Fernandez-Martinez et al. 2017a). Even then masting can be diluted over successive years and is not quite bimodal. Under weaker variation in the environment an economy of scale is needed to achieve a well-defined masting cycle (Kerkhoff 2004; Lyles et al. 2009). Such large differences between very good and poor years in terms of resources (assuming linear responses) are ecologically rather unrealistic though for medium-length time series covering fairly regular 2- or 3-year masting cycles. This in turn suggests that without an economy of scale a non-linear response is more likely to be involved. An essential requirement of such masting would be to avoid degeneration of the time series into chaos – as shown by the models of Rees et al. (2002).

The systems view would postulate feedback control to avoid such chaos, but it could allow for an unpredictable mix of 1-, 2- and 3-year cycles over time to the same effect in thwarting predator tracking. For a limited resource like P, such a system means regulation of internal fluctuations across the tree parts within a masting cycle, so that some equilibrium in

the P dynamics is maintained medium-term. An economy of scale which itself involves a mechanism for P distribution would facilitate masting when based on storage and thresholds of that same resource. In this respect, ectomycorrhizal networks, which enable P acquisition and uptake (and possibly P storage), could bring about an equilibration of P across the connected roots of neighbouring trees, which would allow a contemporaneous storing and triggering of mast fruiting, as P levels rose in concert across all trees in the population (Fig. 5). In the case of *M. bisulcata* in this study, the key climate variables vary in a somewhat normal way, and it is unlikely they can cue masting in a simple directly linear manner. However, if wetter dry seasons lead to non-linear increases, or pulses, of P-and/or K-uptake, coordinated across the forest, synchrony is also possible. The network idea would also explain why one or both *Tetraberlinia* species often mast fruit along with *M. bisulcata*, because they largely share the same mycorrhizal species (Newbery et al. 1988; Newbery et al. 2000). The non-stationary aspect of the main hypothesis suggests a finely tuned evolved ecophysiological system in *M. bisulcata* allowing it to respond to variation of the climate in order to maximize its fitness.

CONCLUSIONS

The results in this paper confirm and extend the findings based on those for 1989–2004 presented in the Newbery et al. (2006a) with greater statistical detail and confidence. It would have been valuable to know whether the 3-year cycle returned after 2017. Post-dispersal predation is currently important, especially in well-lighted gaps although not in the shaded understorey or below *M. bisulcata* crowns. It could have been much more common in the past when either *M. bisulcata* individuals were scattered across the landscape (Newbery and Gartlan 1996), or when grove dynamics was at the rebuilding stage (Newbery et al. 2013).

Since, seeds and seedlings need P to establish and reach to the network of ectomycorrhizas, it

would be evolutionarily advantageous for an ectomycorrhizal species to mast fruit and

increase its chances of establishment. It is interesting that masting is achieved in the most

clustered caesalps and was not apparent for the more spatially dispersed species in southern

Korup.

Most of the C dynamics models for trees to date (e.g. Cannell and Dewar 1994;

Lacointe 2000; Makela et al. 2008; Franklin et al. 2012) do not cater adequately, if at all, for

environmental stochasticity or non-constant reproductive allocation schedules. Source-sink

models for herbaceous plants, let alone for more complicated trees, have made little

conceptual progress in the last three decades since Farrar (1993), Kozlowski (1971, 1992),

Minchin and Thorpe (1996), and Wardlaw (1990), apart from raising an unresolved debate

over passive versus active storage. They usually assume steady-state equilibrium conditions,

are often teleonomic (e.g. Thornley and Parsons 2014), and are strongly focussed on stem

diameter growth and maximizing tree yield. They are, therefore, still too far away from an

understanding of trees in natural forest ecosystems (Le Roux et al. 2001) to explain even

reproductive schedules like masting. Source-sink relationships, within trees especially, turns

out to be very complicated: a more integrated systems approach is probably required. An

considerable challenge is working out how P, K and N interact with C dynamics over the

complete life-history schedule of a tree. A very interesting hypothesis coming from this

extended phenological analysis is that both K and P levels might synergistically control mast

fruiting of *M. bisulcata*, mediated by the physical structure and functioning of the surface

root mat for nutrient cycling. To test this hypothesis further requires much more detailed

recording of nutrient processes at the tree-level and within the system, replicated over several

mast fruiting events.

The literature review of the likely mechanisms lying behind mast fruiting in *M.*

bisulcata gives valuable pointers to detailed aspects that need investigation. The most critical

data for the nutrient resource limitation hypothesis would be phosphorus concentrations in

roots, branches, leaves and fruits of a large sample of trees over several masting cycles.

Detection of phytic acid in branch wood, prior to fruiting especially, could confirm that P was

being stored. Lack of predicted changes in P over time would falsify the hypothesis. In

addition, to test the related economy-of-scale mechanism for equilibration of P between adult

trees at masting would require showing everywhere a flux of P through the mycelia of the

shared ectomycorrhizal network, i.e. between neighbouring large trees.

The fundamental driving variable behind the deterministic mechanically intricate

system operating to cause mast fruiting in *M. bisulcata*, is the stochastic variation in rainfall

(Fig. 5). The vicissitudes of rainfall in the dry season over successive years could generate

the non-linear differences needed to enable the storage process. Stochasticity becomes the

ultimate 'causal' factor over evolutionary time. This can operate at Korup because P as a

resource is strongly limiting the system, there is seasonality, and rainfall alters access,

availability and uptake of P. The hypothesis argues for a fruiting-rooting trade-off as the basis

to alternate bearing and masting. For a common nutrient threshold to operate trees must be

spatially close and connected via their roots and mycorrhizas. The new evidence and analyses

here, showing that both climate and ectomycorrhizas might operate respectively on and

within the forest nutrient cycle, makes for a major step forward in refining the original

PACER hypothesis (Newbery et al. 1997).

1824

1825   ACKNOWLEDGEMENTS

The Ministries of Forests and Wildlife (MINFOF) and Scientific Research and Innovation

(MINRESI) granted permission to conduct field research, and the Conservator's Office of

Korup National Park approved access to the field site. We thank Sylvanos Ngibili

(phenological recording) and Marlise Zimmermann (chemical analysis of plant and soil

materials), both of late, for their valuable assistance. Xander van der Burgt helped with the

soil moisture readings.

1832

1833   REFERENCES

Abe T, Tachiki Y, Kon H, Nagasaka A, Onodera K, Minamino K, Han QM, Satake A (2016)
Parameterisation and validation of a resource budget model for masting using spatiotemporal
flowering data of individual trees. Ecology Letters 19: 1129-1139. doi:10.1111/ele.12651.

Agresti A (2007) An introduction to categorical data analysis. J. Wiley & Sons, New Jersey, USA, pp.

Alexander I (1989a) Mycorrhizas in tropical forests. In: Proctor J (Ed) Mineral nutrients in tropical
forest and savanna ecosystems. Blackwell Scientific Publications, Oxford, 169-188.

Alexander IJ (1989b) Systematics and ecology of ectomycorrhizal legumes. Monographs in
Systematic Botany of the Missouri Botanical Garden 29: 607-624.

Allen RB, Millard P, Richardson SJ (2017) A resource centric view of climate and mast seeding in
trees. In: Canovas F, Lüttige U, Matyssek R (Eds) Progress in Botany Vol 79. Springer,
Cham, Switzerland, 233-268.

Aoyagi R, Imai N, Hidaka A, Samejima H, Kitayama K (2018) Abrupt increase in phosphorus and
potassium fluxes during a masting event in a Bornean tropical forest. Ecological Research 33:
1193-1205. doi:10.1007/s11284-018-1642-9.

Basnet K, Scatena FN, Likens GE, Lugo AE (1993) Ecological consequences of root grafting in
tabonuco (*Dacryodes excelsa*) trees in the Luquillo Experimental-Forest, Puerto Rico.
Biotropica 25: 28-35. doi:10.2307/2388976.

Bennett EJ, Roberts JA, Wagstaff C (2011) The role of the pod in seed development: strategies for
manipulating yield. New Phytologist 190: 838-853. doi:10.1111/j.1469-8137.2011.03714.x.

Bieleski RL (1973) Phosphate pools, phosphate transport, and phosphate availability. Annual Review
of Plant Physiology and Plant Molecular Biology 24: 225-252.
doi:10.1146/annurev.pp.24.060173.001301.

Bogdziewicz M, Kelly D, Ascoli D, Caignard T, Chianucci F, Crone EE, Fleurot E, Foest JJ, Gratzer
G, Hagiwara T, Han QM, Journé V, Keurinck L, Kondrat K, McClory R, LaMontagne JM,

Mundo IA, Nussbaumer A, Oberklammer I, Ohno M, Pearse IS, Pesendorfer MB, Resente G,

Satake A, Shibata M, Snell RS, Szymkowiak J, Touzot L, Zwolak R, Zywiec M, Hacket-Pain

AJ (2024) Evolutionary ecology of masting: mechanisms, models, and climate change.

Trends in Ecology & Evolution 39: 851-862. doi:10.1016/j.tree.2024.05.006.

Bormann FH (1966) Structure, function, and ecological significance of root grafts in *Pinus strobus* L.

Ecological Monographs 36: 1-26. doi:10.2307/1948486.

Brearley FQ (2012) Ectomycorrhizal associations of the Dipterocarpaceae. Biotropica 44: 637-648.

doi:10.1111/j.1744-7429.2012.00862.x.

Brown PH, Weinbaum SA, Picchioni GA (1995) Alternate bearing influences annual nutrient

consumption and the total nutrient content of mature pistachio trees. Trees-Structure and

Function 9: 158-164. doi:10.1007/bf02418205.

Bustan A, Avni A, Lavee S, Zipori I, Yeselson Y, Schaffer AA, Riov J, Dag A (2011) Role of

carbohydrate reserves in yield production of intensively cultivated oil olive (*Olea europaea*

L.) trees. Tree Physiology 31: 519-530. doi:10.1093/treephys/tpr036.

Camberlin P, Janicot S, Poccard I (2001) Seasonality and atmospheric dynamics of the teleconnection

between African rainfall and tropical sea-surface temperature: Atlantic vs. ENSO.

International Journal of Climatology 21: 973-1005. doi:10.1002/joc.673.

Cannell MGR, Dewar RC (1994) Carbon allocation in trees: a review of concepts for modeling. In:

Advances in Ecological Research, Vol 25. 59-104. doi:10.1016/S0065-2504(08)60213-5.

Chapin FS, Schulze ED, Mooney HA (1990) The ecology and economics of storage in plants. Annual

Review of Ecology and Systematics 21: 423-447. doi:10.1146/annurev.ecolsys.21.1.423.

Charlesworth B (1980) Evolution in age-structured populations. Cambridge University Press,

Cambridge, UK, pp.

Chiou TJ, Lin SI (2011) Signaling network in sensing phosphate availability in plants. Annual Review

of Plant Biology 62: 185-206. doi:10.1146/annurev-arplant-042110-103849.

Chuyong GB (1994) Nutrient cycling in ectomycorrhizal legume-dominated forest in Korup National

Park, Cameroon. PhD thesis. Stirling University, UK, 239 pp.

Chuyong GB, Newbery DM, Songwe NC (2000) Litter nutrients and retranslocation in a central

African rain forest dominated by ectomycorrhizal trees. New Phytologist 148: 493-510.

doi:10.1046/j.1469-8137.2000.00774.x.

Chuyong GB, Newbery DM, Songwe NC (2002) Litter breakdown and mineralization in a central

African rain forest dominated by ectomycorrhizal trees. Biogeochemistry 61: 73-94.

doi:10.1023/a:1020276430119.

Chuyong GB, Newbery DM, Songwe NC (2004) Rainfall input, throughfall and stemflow of nutrients

in a central African rain forest dominated by ectomycorrhizal trees. Biogeochemistry 67: 73-

91. doi:10.1023/B:BIOG.0000015316.90198.cf.

Corley RHV, Tinker PB (2016) The oil palm. J. Wiley, pp.

Corning PA (2019) Teleonomy and the proximate-ultimate distinction revisited. Biological Journal of
the Linnean Society 127: 912-916.

Corrales A, Henkel TW, Smith ME (2018) Ectomycorrhizal associations in the tropics -
biogeography, diversity patterns and ecosystem roles. New Phytologist 220: 1076-1091.
doi:10.1111/nph.15151.

Cox DR (1958) The regression analysis of binary sequences. Journal of the Royal Statistical Society
Series B-Statistical Methodology 20: 215-242. doi:10.1111/j.2517-6161.1958.tb00292.x.

Cox DR (1981) Statistical analysis of time series: some recent developments. Scandinavian Journal of
Statistics 8: 93-115.

Cox DR, Snell EJ (1989) Analysis of binary data. Chapman & Hall/CRC, pp.

Crawley MJ, Long CR (1995) Alternate bearing, predator satiation and seedling recruitment in
Quercus robur L. Journal of Ecology 83: 683-696. doi:10.2307/2261636.

Crone EE, Rapp JM (2014) Resource depletion, pollen coupling, and the ecology of mast seeding.
Annals of the New York Academy of Sciences 1322: 21-34. doi:10.1111/nyas.12465.

Davis LD (1957) Flowering and alternate bearing. American Society for Horticultural Science 70:
545-556.

de la Estrella M, Forest F, Klitgard B, Lewis GP, Mackinder BA, de Queiroz LP, Wieringa JJ,
Bruneau A (2018) A new phylogeny-based tribal classification of subfamily Detarioideae, an
early branching clade of florally diverse tropical arborescent legumes. Scientific Reports 8: 1-
14. doi:10.1038/s41598-018-24687-3.

Dietze MC, Sala A, Carbone MS, Czimczik CI, Mantooth JA, Richardson AD, Vargas R (2014)
Nonstructural Carbon in Woody Plants. Annual Review of Plant Biology 65: 667-687.
doi:10.1146/annurev-arplant-050213-040054.

Dunsmuir WTM (2016) Generalized linear autoregressive moving average models. In: David RA,
Holan SH, Lund R, Ravishanker N (Eds) Handbook of discrete-valued time series. CRC
Press, Boca Raton, USA, 51-76.

Dunsmuir WTM, Scott DJ (2015) The glarma package for observation-driven time series regression
of counts. Journal of Statistical Software 67: 1-36.

Etta CE, Chuyong GB, Newbery DM (2022) Climate records for Bulu, Ndian Division, SW
Cameroon. [Data set]. https://zenodo.org/records/7220319.

Farrar JF (1993) Sink strength: what is it and how do we measure it? A summary. Plant Cell and
Environment 16: 1045-1046. doi:10.1111/j.1365-3040.1996.tb02061.x.

Fernandez-Martinez M, Bogdziewicz M, Espelta JM, Penuelas J (2017a) Nature beyond linearity:
meteorological variability and Jensen's Inequality can explain mast seeding behavior.
Frontiers in Ecology and Evolution 5: 1-8. doi:10.3389/fevo.2017.00134.

Fernandez-Martinez M, Pearse I, Sardans J, Sayol F, Koenig WD, LaMontagne JM, Bogdziewicz M,
Collalti A, Hacket-Pain A, Vacchiano G, Espelta JM, Penuelas J, Janssens IA (2019) Nutrient

scarcity as a selective pressure for mast seeding. Nature Plants 5: 1222-1227. doi:10.1038/s41477-019-0549-y.

Fernandez-Martinez M, Sardans J, Sayol F, LaMontagne JM, Bogdziewicz M, Collalti A, Hacket-Pain A, Vacchiano G, Espelta JM, Penuelas J, Janssens IA (2020) Reply to: Nutrient scarcity cannot cause mast seeding. Nature Plants 6: 763-765. doi:10.1038/s41477-020-0703-6.

Fernandez-Martinez M, Vicca S, Janssens IA, Espelta JM, Penuelas J (2017b) The role of nutrients, productivity and climate in determining tree fruit production in European forests. New Phytologist 213: 669-679. doi:10.1111/nph.14193.

Fisher RA, Williams M, Ruivo MD, de Costa AL, Meira P (2008) Evaluating climatic and soil water controls on evapotranspiration at two Amazonian rainforest sites. Agricultural and Forest Meteorology 148: 850-861. doi:10.1016/j.agrformet.2007.12.001.

Fox J (2008) Applied regression analysis and generalized linear models. Sage Publications, Los Angeles, USA, pp.

Fox J, Weisberg S (2011) An R companion to applied regression. Sage Publications, Los Angeles, CA., pp.

Franklin O, Johansson J, Dewar RC, Dieckmann U, McMurtrie RE, Brannstrom A, Dybzinski R (2012) Modeling carbon allocation in trees: a search for principles. Tree Physiology 32: 648-666. doi:10.1093/treephys/tpr138.

Garcia G, Re B, Orians C, Crone E (2021) By wind or wing: pollination syndromes and alternate bearing in horticultural systems. Philosophical Transactions of the Royal Society B-Biological Sciences 376: 1-9. doi:10.1098/rstb.2020.0371.

Gartlan JS (1992) Cameroon. In: Sayer JA, Harcourt CS, Collins NM (Eds) The Conservation Atlas of Tropical Forests: Africa. Macmillan Publishers, London, UK, 110-118.

Gartlan JS, Newbery DM, Thomas DW, Waterman PG (1986) The influence of topography and soil phosphorus on the vegetation of Korup Forest Reserve, Cameroun. Vegetatio 65: 131-148. doi:10.1007/bf00044814.

Goldschmidt EE, Golomb A (1982) The carbohydrate balance of alternate-bearing citrus trees and the significance of reserves for flowering and fruiting. Journal of the American Society for Horticultural Science 107: 206-208. doi:10.21273/JASHS.107.2.206.

Graham BF, Bormann FH (1966) Natural root grafts. Botanical Review 32: 255-292. doi:10.1007/bf02858662.

Green JJ, Newbery DM (2001a) Light and seed size affect establishment of grove-forming ectomycorrhizal rain forest tree species. New Phytologist 151: 271-289. doi:10.1046/j.1469-8137.2001.00156.x.

Green JJ, Newbery DM (2001b) Shade and leaf loss affect establishment of grove-forming ectomycorrhizal rain forest tree species. New Phytologist 151: 291-309. doi:10.1046/j.1469-8137.2001.00157.x.

Green JJ, Newbery DM (2002) Reproductive investment and seedling survival of the mast-fruiting
rain forest tree, *Microberlinia bisulcata* A. Chev. Plant Ecology 162: 169-183.
doi:10.1023/a:1020304212118.

Han QM, Kabeya D (2017) Recent developments in understanding mast seeding in relation to
dynamics of carbon and nitrogen resources in temperate trees. Ecological Research 32: 771-
778. doi:10.1007/s11284-017-1494-8.

Han QM, Kabeya D, Iio A, Inagaki Y, Kakubari Y (2014) Nitrogen storage dynamics are affected by
masting events in *Fagus crenata*. Oecologia 174: 679-687. doi:10.1007/s00442-013-2824-3.

Hansen P, Ryugo K, Ramos DE, Fitch L (1982) Influence of cropping on Ca, K, Mg, and
carbohydrate status of 'French' prune trees grown on potassium limited soils. Journal of the
American Society for Horticultural Science 107: 511-515. doi:10.21273/JASHS.107.3.511.

Hart TB (1995) Seed, seedling and sub-canopy survival in monodominant and mixed forests of the
Ituri Forest, Africa. Journal of Tropical Ecology 11: 443-459.
doi:10.1017/s0266467400008919.

Hartmann H, Trumbore S (2016) Understanding the roles of nonstructural carbohydrates in forest
trees - from what we can measure to what we want to know. New Phytologist 211: 386-403.
doi:10.1111/nph.13955.

Henkel TW, Mayor JR (2019) Implications of a long-term mast seeding cycle for climatic
entrainment, seedling establishment and persistent monodominance in a Neotropical,
ectomycorrhizal canopy tree. Ecological Research 34: 472-484. doi:10.1111/1440-
1703.12014.

Henkel TW, Mayor JR, Woolley LP (2005) Mast fruiting and seedling survival of the
ectomycorrhizal, monodominant *Dicymbe corymbosa* (Caesalpiniaceae) in Guyana. New
Phytologist 167: 543-556. doi:10.1111/j.1469-8137.2005.01431.x.

Hoch G (2005) Fruit-bearing branchlets are carbon autonomous in mature broad-leaved temperate
forest trees. Plant Cell and Environment 28: 651-659. doi:10.1111/j.1365-3040.2004.01311.x.

Hoch G, Richter A, Körner C (2003) Non-structural carbon compounds in temperate forest trees.
Plant, Cell and Environment 26: 1067-1081. doi:10.1046/j.0016-8025.2003.01032.x.

Hosmer DW, Lemeshow S, Sturdivant RX (2013) Applied logistic regression. J. Wiley & Sons, New
Jersey, USA, pp.

Ichie T, Igarashi S, Yoshida S, Kenzo T, Masaki T, Tayasu I (2013) Are stored carbohydrates
necessary for seed production in temperate deciduous trees? Journal of Ecology 101: 525-531.
doi:10.1111/1365-2745.12038.

Ichie T, Kenta T, Nakagawa M, Sato K, Nakashizuka T (2005a) Resource allocation to reproductive
organs during masting in the tropical emergent tree, *Dipterocarpus tempehes*. Journal of
Tropical Ecology 21: 237-241. doi:10.1017/s0266467404002214.

Ichie T, Kenzo T, Kitahashi Y, Koike T, Nakashizuka T (2005b) How does *Dryobalanops aromatica* supply carbohydrate resources for reproduction in a masting year? Trees-Structure and Function 19: 703-710. doi:10.1007/s00468-005-0434-3.

Ichie T, Nakagawa M (2013) Dynamics of mineral nutrient storage for mast reproduction in the tropical emergent tree *Dryobalanops aromatica*. Ecological Research 28: 151-158. doi:10.1007/s11284-011-0836-1.

Igarashi S, Shibata M, Masaki T, Tayasu I, Ichie T (2019) Mass flowering of *Fagus crenata* does not depend on the amount of stored carbohydrates in trees. Trees-Structure and Function 33: 1399-1408. doi:10.1007/s00468-019-01867-w.

Isagi Y, Sugimura K, Sumida A, Ito H (1997) How does masting happen and synchronize? Journal of Theoretical Biology 187: 231-239. doi:10.1006/jtbi.1997.0442.

Janzen DH (1974) Tropical blackwater rivers, animals and mast fruiting by the Dipterocarpaceae. Biotropica 6: 69-103. doi:10.2307/2989823.

Jones JM, Heineman KD, Dalling JW (2019) Soil and species effects on bark nutrient storage in a premontane tropical forest. Plant and Soil 438: 347-360. doi:10.1007/s11104-019-04026-9.

Karst J, Jones MD, Hoeksema JD (2023) Positive citation bias and overinterpreted results lead to misinformation on common mycorrhizal networks in forests. Nature Ecology & Evolution 7: 501-+. doi:10.1038/s41559-023-01986-1.

Kelly D (1994) The evolutionary ecology of mast seeding. Trends in Ecology & Evolution 9: 465-470. doi:10.1016/0169-5347(94)90310-7.

Kelly D (2020) Nutrient scarcity cannot cause mast seeding. Nature Plants 6: 760-762. doi:10.1038/s41477-020-0702-7.

Kelly D, Sork VL (2002) Mast seeding in perennial plants: Why, how, where? Annual Review of Ecology and Systematics 33: 427-447. doi:10.1146/annurev.ecolsys.33.020602.095433.

Kerkhoff AJ (2004) Expectation, explanation and masting. Evolutionary Ecology Research 6: 1003-1020.

Kitayama K, Tsujii Y, Aoyagi R, Aiba S (2015) Long-term C, N and P allocation to reproduction in Bornean tropical rain forests. Journal of Ecology 103: 606-615. doi:10.1111/1365-2745.12379.

Kozlowski TT (1971) Growth and development of trees. Vol. II. Cambial growth, root growth and reproductive growth. Academic Press, New York, Ch. 5-7-8 pp.

Kozlowski TT (1992) Carbohydrate sources and sinks in woody plants. Botanical Review 58: 107-222. doi:10.1007/bf02858600.

Kurita Y, Baba K, Ohnishi M, Matsubara R, Kosuge K, Anegawa A, Shichijo C, Ishizaki K, Kaneko Y, Hayashi M, Suzaki T, Fukaki H, Mimura T (2017) Inositol hexakis phosphate is the seasonal phosphorus reservoir in the deciduous woody plant *Populus alba* L. Plant and Cell Physiology 58: 1477-1485. doi:10.1093/pcp/pcx106.

2042 Lacointe A (2000) Carbon allocation among tree organs: A review of basic processes and
2043   representation in functional-structural tree models. Annals of Forest Science 57: 521-533.
2044   doi:10.1051/forest:2000139.

2045 Laland KN, Sterelny K, Odling-Smee J, Hoppitt W, Uller T (2011) Cause and effect in biology
2046   revisited: Is Mayr's proximate-ultimate dichotomy still useful? Science 334: 1512-1516.
2047   doi:10.1126/science.1210879.

2048 Larue CD (1952) Root-grafting in tropical trees. Science 115: 296-296.
2049   doi:10.1126/science.115.2985.296.

2050 Le Roncé I, Dardevet E, Venner S, Schönbeck L, Gessler A, Chuine I, Limousin JM (2023)
2051   Reproduction alternation in trees: testing the resource depletion hypothesis using
2052   experimental fruit removal in *Quercus ilex*. Tree Physiology 43: 952-964.
2053   doi:10.1093/treephys/tpad025.

2054 Le Roux X, Lacointe A, Escobar-Gutierrez A, Le Dizes S (2001) Carbon-based models of individual
2055   tree growth: A critical appraisal. Annals of Forest Science 58: 469-506.

2056 Leeper AC, Lawrence BA, LaMontagne JM (2020) Plant-available soil nutrients have a limited
2057   influence on cone production patterns of individual white spruce trees. Oecologia 194: 101-
2058   111. doi:10.1007/s00442-020-04759-w.

2059 Legros S, Mialet-Serra I, Clement-Vidal A, Caliman JP, Siregar FA, Fabre D, Dingkuhn M (2009)
2060   Role of transitory carbon reserves during adjustment to climate variability and source-sink
2061   imbalances in oil palm (*Elaeis guineensis*). Tree Physiology 29: 1199-1211.
2062   doi:10.1093/treephys/tpp057.

2063 Letouzey R (1968) Etude Phytogéographique du Cameroun. P. LeChevalier, Paris, pp.

2064 Lloyd DG (1980) Sexual strategies in plants. I. An hypothesis of serial adjustment of maternal
2065   investment during one reproductive season. New Phytologist 86: 69-79. doi:10.1111/j.1469-
2066   8137.1980.tb00780.x.

2067 Loehle C, Jones RH (1990) Adaptive significance of root grafting in trees. Functional Ecology 4: 268-
2068   271.

2069 LPWG (2017) A new subfamily classification of the Leguminosae based on a taxonomically
2070   comprehensive phylogeny. Taxon 66: 44-77. doi:10.12705/661.3.

2071 Lyles D, Rosenstock TS, Hastings A, Brown PH (2009) The role of large environmental noise in
2072   masting: General model and example from pistachio trees. Journal of Theoretical Biology
2073   259: 701-713. doi:10.1016/j.jtbi.2009.04.015.

2074 Maisels F (2004) Defoliation of a monodominant rain-forest tree by a noctuid moth in Gabon. Journal
2075   of Tropical Ecology 20: 239-241. doi:10.1017/s0266467403001044.

2076 Makela A, Valentine HT, Helmisaari HS (2008) Optimal co-allocation of carbon and nitrogen in a
2077   forest stand at steady state. New Phytologist 180: 114-123. doi:10.1111/j.1469-
2078   8137.2008.02558.x.

Marschner P (Ed) (2012) Marschner's mineral nutrition of higher plants. Vol. Elsevier, Amsterdam,
pp.

Marshall C, Grace J (Eds) (1992) Fruit and seed production: aspects of development, environmental
physiology and ecology. Vol. Cambridge University Press, Cambridge, UK, pp.

Maynard Smith J (1972) On evolution. Edinburgh University Press, Edinburgh, UK, pp.

McCullagh P, Nelder JA (1989) Generalized linear models. Chapman & Hall, London, UK, pp.

McKenzie E (1985) Some simple models for discrete variate time series. Water Resources Bulletin
21: 645-650. doi:10.1111/j.1752-1688.1985.tb05379.x.

McKenzie E (2003) Discrete variate time series. In: Rao CR, Shanbhag DN (Eds) Handbook of
Statistics. Elsevier, 573-606.

Millard P, Sommerkorn M, Grelet GA (2007) Environmental change and carbon limitation in trees: a
biochemical, ecophysiological and ecosystem appraisal. New Phytologist 175: 11-28.
doi:10.1111/j.1469-8137.2007.02079.x.

Minchin PEH, Thorpe MR (1996) What determines carbon partitioning between competing sinks?
Journal of Experimental Botany 47: 1293-1296. doi:10.1093/jxb/47.Special_Issue.1293.

Miyazaki Y (2013) Dynamics of internal carbon resources during masting behavior in trees.
Ecological Research 28: 143-150. doi:10.1007/s11284-011-0892-6.

Monselise SP, Goldschmidt EE (1982) Alternate bearing in fruit trees. Horticultural Research 4: 128-
173. doi:10.1002/9781118060773.ch5.

Mulligan DR (1988) Phosphorus concentrations and chemical fractions in *Eucalyptus* seedlings grown
for a prolonged period under nutrient- deficient conditions. New Phytologist 110: 479-486.
doi:10.1111/j.1469-8137.1988.tb00286.x.

Mund M, Herbst M, Knohl A, Matthäus B, Schumacher J, Schall P, Siebicke L, Tamrakar R, Ammer
C (2020) It is not just a 'trade-off': indications for sink- and source-limitation to vegetative
and regenerative growth in an old-growth beech forest. New Phytologist 226: 111-125.
doi:10.1111/nph.16408.

Mund M, Kutsch WL, Wirth C, Kahl T, Knohl A, Skomarkova MV, Schulze ED (2010) The
influence of climate and fructification on the inter-annual variability of stem growth and net
primary productivity in an old-growth, mixed beech forest. Tree Physiology 30: 689-704.
doi:10.1093/treephys/tpq027.

Nascimento MT, Proctor J (1994) Insect defoliation of a monodominant Amazonian rainforest.
Journal of Tropical Ecology 10: 633-636. doi:10.1017/s0266467400008312.

Netzer F, Schmid C, Herschbach C, Rennenberg H (2017) Phosphorus-nutrition of European beech
(*Fagus sylvatica* L.) during annual growth depends on tree age and P-availability in the soil.
Environmental and Experimental Botany 137: 194-207.
doi:10.1016/j.envexpbot.2017.02.009.

Newbery DM (2005) Ectomycorrhizas and mast fruiting in trees: linked by climate-driven tree
resources? New Phytologist 167: 324-326. doi:10.1111/j.1469-8137.2005.01468.x.
Newbery DM, Alexander IJ, Rother JA (1997) Phosphorus dynamics in a lowland African rain forest:
The influence of ectomycorrhizal trees. Ecological Monographs 67: 367-409.
doi:10.1890/0012-9615(1997)067[0367:Pdiala]2.0.Co;2.
Newbery DM, Alexander IJ, Rother JA (2000) Does proximity to conspecific adults influence the
establishment of ectomycorrhizal trees in rain forest? New Phytologist 147: 401-409.
doi:10.1046/j.1469-8137.2000.00698.x.
Newbery DM, Alexander IJ, Thomas DW, Gartlan JS (1988) Ectomycorrhizal rain-forest legumes and
soil phosphorus in Korup National Park, Cameroon. New Phytologist 109: 433-450.
doi:10.1111/j.1469-8137.1988.tb03719.x.
Newbery DM, Chuyong GB, Green JJ, Songwe NC, Tchuenteu F, Zimmermann L (2002) Does low
phosphorus supply limit seedling establishment and tree growth in groves of ectomycorrhizal
trees in a central African rainforest? New Phytologist 156: 297-311. doi:10.1046/j.1469-
8137.2002.00505.x.
Newbery DM, Chuyong GB, Zimmermann L (2006a) Mast fruiting of large ectomycorrhizal African
rain forest trees: importance of dry season intensity, and the resource-limitation hypothesis.
New Phytologist 170: 561-579. doi:10.1111/j.1469-8137.2006.01691.x.
Newbery DM, Chuyong GB, Zimmermann L, Praz C (2006b) Seedling survival and growth of three
ectomycorrhizal caesalpiniaceous tree species in a Central African rain forest. Journal of
Tropical Ecology 22: 499-511. doi:10.1017/s0266467406003427.
Newbery DM, Gartlan JS (1996) Structural analysis of the rain forest at Korup and Douala Edea,
Cameroon. Proceedings of the Royal Society of Edinburgh B 104: 177-224.
doi:10.1017/S0269727000006138.
Newbery DM, Neba GA (2019) Micronutrients may influence the efficacy of ectomycorrhizas to
support tree seedlings in a lowland African rain forest. Ecosphere 10: 1-28.
doi:10.1002/ecs2.2686.
Newbery DM, Praz CJ, van der Burgt XM, Norghauer JM, Chuyong GB (2010) Recruitment
dynamics of the grove-dominant tree *Microberlinia bisulcata* in African rain forest: extending
the light response versus adult longevity trade-off concept. Plant Ecology 206: 151-172.
doi:10.1007/s11258-009-9631-2.
Newbery DM, Schwan S, Chuyong GB, van der Burgt XM (2009) Buttress form of the central
African rain forest tree *Microberlinia bisulcata*, and its possible role in nutrient acquisition.
Trees-Structure and Function 23: 219-234. doi:10.1007/s00468-008-0270-3.
Newbery DM, Songwe NS, Chuyong GB (1998) Phenology and dynamics of an African rainforest at
Korup, Cameroon. In: Newbery DM, Prins HHT, Brown ND (Eds) Dynamics of tropical
communities. Blackwell Science, 267-308.

Newbery DM, van der Burgt XM, Moravie MA (2004) Structure and inferred dynamics of a large
grove of *Microberlinia bisulcata* trees in central African rain forest: the possible role of
periods of multiple disturbance events. Journal of Tropical Ecology 20: 131-143.
doi:10.1017/s0266467403001111.

Newbery DM, van der Burgt XM, Worbes M, Chuyong GB (2013) Transient dominance in a central
African rain forest. Ecological Monographs 83: 339-382. doi:10.1890/12-1699.1.

Nicholson SE, Funk C, Fink AH (2018) Rainfall over the African continent from the 19th through the
21st century. Global and Planetary Change 165: 114-127.
doi:10.1016/j.gloplacha.2017.12.014.

Nicholson SE, Some B, Kone B (2000) An analysis of recent rainfall conditions in West Africa,
including the rainy seasons of the 1997 El Nino and the 1998 La Nina years. Journal of
Climate 13: 2628-2640. doi:10.1175/1520-0442(2000)013<2628:Aaorrc>2.0.Co;2.

Noble AE, Rosenstock TS, Brown PH, Machta J, Hastings A (2018) Spatial patterns of tree yield
explained by endogenous forces through a correspondence between the Ising model and
ecology. Proceedings of the National Academy of Sciences of the United States of America
115: 1825-1830. doi:10.1073/pnas.1618887115.

Norden N, Chave J, Belbenoit P, Caubère A, Châtelet P, Forget P-M, Thébaud C (2007) Mast fruiting
is a frequent strategy in woody species of eastern South America. PlosOne 10: 1-9 (e1079).
doi:10.1371/journal.pone.0001079.

Norghauer JM, Newbery DM (2010) Recruitment limitation after mast-seeding in two African rain
forest trees. Ecology 91: 2303-2312. doi:10.1890/09-0071.1.

Norghauer JM, Newbery DM (2011) Seed fate and seedling dynamics after masting in two African
rain forest trees. Ecological Monographs 81: 443-468. doi:10.1890/10-2268.1.

Norghauer JM, Newbery DM (2015) Tree size and fecundity influence ballistic seed dispersal of two
dominant mast-fruiting species in a tropical rain forest. Forest Ecology and Management 338:
100-113. doi:10.1016/j.foreco.2014.11.005.

Norghauer JM, Newbery DM (2016) Density-dependent dynamics of a dominant rain forest tree
change with juvenile stage and time of masting. Oecologia 181: 207-223.
doi:10.1007/s00442-015-3534-9.

Norghauer JM, Newbery DM, Neba GA (2023) Indications of an *Achaea* sp. caterpillar outbreak
disrupting fruiting of an ectomycorrhizal tropical tree in Central African rainforest. Plant
Ecology and Evolution 156: 46-58. doi:10.5091/plecevo.96572.

Norton DA, Kelly D (1988) Mast seeding over 33 years by *Dacrydium cupressinum* Lamb. (rimu)
(Podocarpaceae) in New Zealand: the importance of economies of scale. Functional Ecology
2: 399-408. doi:DOI10.2307/2389413.

Or D, Wraith JM (2002) Soil water content and water potential relationships. In: Warrick AW (Ed)
Soil physics companion. CRC Press, Boca Raton, FL, USA, 49-84.

Pearse IS, Koenig WD, Kelly D (2016) Mechanisms of mast seeding: resources, weather, cues, and selection. New Phytologist 212: 546-562. doi:10.1111/nph.14114.

Picchioni GA, Brown PH, Weinbaum SA, Muraoka TT (1997) Macronutrient allocation to leaves and fruit of mature, alternate-bearing pistachio trees: Magnitude and seasonal patterns at the whole-canopy level. Journal of the American Society for Horticultural Science 122: 267-274. doi:10.21273/jashs.122.2.267.

Poccard I, Janicot S, Camberlin P (2000) Comparison of rainfall structures between NCEP/NCAR reanalyses and observed data over tropical Africa. Climate Dynamics 16: 897-915. doi:10.1007/s003820000087.

R_Core_Team (2022) R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/. pp.

Raboy V (2003) Myo-Inositol-1,2,3,4,5,6-hexakisphosphate. Phytochemistry 64: 1033-1043. doi:10.1016/s0031-9422(03)00446-1.

Rees M, Kelly D, Bjornstad ON (2002) Snow tussocks, chaos, and the evolution of mast seeding. American Naturalist 160: 44-59. doi:10.1086/340603.

Rennenberg H, Herschbach C (2013) Phosphorus nutrition of woody plants: many questions - few answers. Plant Biology 15: 785-788. doi:10.1111/plb.12078.

Richardson AD, Carbone MS, Keenan TF, Czimczik CI, Hollinger DY, Murakami P, Schaberg PG, Xu XM (2013) Seasonal dynamics and age of stemwood nonstructural carbohydrates in temperate forest trees. New Phytologist 197: 850-861. doi:10.1111/nph.12042.

Rosecrance RC, Weinbaum SA, Brown PH (1996) Assessment of nitrogen, phosphorus, and potassium uptake capacity and root growth in mature alternate-bearing pistachio (*Pistacia vera*) trees. Tree Physiology 16: 949-956. doi:10.1093/treephys/16.11-12.949.

Rosecrance RC, Weinbaum SA, Brown PH (1998) Alternate bearing affects nitrogen, phosphorus, potassium and starch storage pools in mature pistachio trees. Annals of Botany 82: 463-470. doi:10.1006/anbo.1998.0696.

Rosell JA, Piper FI, Jimenez-Vera C, Vergilio PCB, Marcati CR, Castorena M, Olson ME (2021) Inner bark as a crucial tissue for non-structural carbohydrate storage across three tropical woody plant communities. Plant Cell and Environment 44: 156-170. doi:10.1111/pce.13903.

Rosenstock TS, Rosa UA, Plant RE, Brown PH (2010) A reevaluation of alternate bearing in pistachio. Scientia Horticulturae 124: 149-152. doi:10.1016/j.scienta.2009.12.007.

Sachs RM (1977) Nutrient diversion: an hypothesis to explain the chemical control of flowering. HortScience 12: 220-222. doi:10.21273/HORTSCI.12.3.220.

Sala A, Hopping K, McIntire EJB, Delzon S, Crone EE (2012a) Masting in whitebark pine (*Pinus albicaulis*) depletes stored nutrients. New Phytologist 196: 189-199. doi:10.1111/j.1469-8137.2012.04257.x.

Sala A, Woodruff DR, Meinzer FC (2012b) Carbon dynamics in trees: feast or famine? Tree Physiology 32: 764-775. doi:10.1093/treephys/tpr143.

Sanchez-Humanes B, Sork VL, Espelta JM (2011) Trade-offs between vegetative growth and acorn production in *Quercus lobata* during a mast year: the relevance of crop size and hierarchical level within the canopy. Oecologia 166: 101-110. doi:10.1007/s00442-010-1819-6.

Satake A, Bjornstad ON (2008) A resource budget model to explain intraspecific variation in mast reproductive dynamics. Ecological Research 23: 3-10. doi:10.1007/s11284-007-0397-5.

Saunier RE, Wagle RF (1965) Root grafting in *Quercus turbinella* Greene. Ecology 46: 749-750. doi:10.2307/1935020.

Schwan S (2003) Phenology, resource conservation and tree architecture of large ectomycorrhizal trees in a lowland African rain forest at Korup, Cameroon. Diploma (MSc) thesis. University of Bern, Switzerland, 77 pp.

Simard SW, Beiler KJ, Bingham MA, Deslippe JR, Philip LJ, Teste FP (2012) Mycorrhizal networks: mechanisms, ecology and modelling. Fungal Biology Reviews 26: 39-60.

Simard SW, Durall DM (2004) Mycorrhizal networks: a review of their extent, function, and importance. Canadian Journal of Botany 82: 1140-1165. doi:10.1139/b04-116.

Simard SW, Perry DA, Jones MD, Myrold DD, Durall DM, Molina R (1997) Net transfer of carbon between ectomycorrhizal tree species in the field. Nature 388: 579-582.

Smaill SJ, Clinton PW, Allen RB, Davis MR (2011) Climate cues and resources interact to determine seed production by a masting species. Journal of Ecology 99: 870-877. doi:10.1111/j.1365-2745.2011.01803.x.

Stearns SC (1992) The evolution of life-history strategies. Oxford University Press, Oxford, UK, pp.

Stevenson MT, Shackel KA (1998) Alternate bearing in pistachio as a masting phenomenon: Construction cost of reproduction versus vegetative growth and storage. Journal of the American Society for Horticultural Science 123: 1069-1075. doi:10.21273/jashs.123.6.1069.

Thornley JHM, Parsons AJ (2014) Allocation of new growth between shoot, root and mycorrhiza in relation to carbon, nitrogen and phosphate supply: Teleonomy with maximum growth rate. Journal of Theoretical Biology 342: 1-14. doi:10.1016/j.jtbi.2013.10.003.

Turnbull C (2011) Long-distance regulation of flowering time. Journal of Experimental Botany 62: 4399-4413. doi:10.1093/jxb/err191.

Vacchiano G, Ascoli D, Berzaghi F, Lucas-Borja ME, Caignard T, Collalti A, Mairota P, Palaghianu C, Reyer CPO, Sanders TGM, Schermer E, Wohlgemuth T, Hacket-Pain A (2018) Reproducing reproduction: How to simulate mast seeding in forest models. Ecological Modelling 376: 40-53. doi:10.1016/j.ecolmodel.2018.03.004.

Van Bael SA, Aiello A, Valderrama A, Medianero E, Samaniego M, Wright SJ (2004) General herbivore outbreak following an El Nino-related drought in a lowland Panamanian forest. Journal of Tropical Ecology 20: 625-633. doi:10.1017/s0266467404001725.

Venables WN, Ripley BD (2002) Modern applied statistics with S. Springer, Berlin, pp.

Venner S, Siberchicot A, Pelisson PF, Schermer E, Bel-Venner MC, Nicolas M, Debias F, Miele V,
Sauzet S, Boulanger V, Delzon S (2016) Fruiting strategies of perennial plants: a resource
budget model to couple mast seeding to pollination efficiency and resource allocation
strategies. American Naturalist 188: 66-75. doi:10.1086/686684.

VSN_International (2022) The guide to the Genstat Command Language (Release 22), Part 1 Syntax.
Part 2 Statistics. VSN International, Hemel Hempstead, UK., pp.

Wardlaw IF (1990) The control of carbon partitioning in plants. New Phytologist 116: 341-381.
doi:10.1111/j.1469-8137.1990.tb00524.x.

Wong M, Wright SJ, Hubbell SP, Foster RB (1990) The spatial pattern and reproductive
consequences of outbreak defoliation in *Quararibea asterolepis*, a tropical tree. Journal of
Ecology 78: 579-588. doi:10.2307/2260885.

Wurth MKR, Pelaez-Riedl S, Wright SJ, Korner C (2005) Non-structural carbohydrate pools in a
tropical forest. Oecologia 143: 11-24. doi:10.1007/s00442-004-1773-2.

Yamauchi A (1996) Theory of mast reproduction in plants: Storage-size dependent strategy.
Evolution 50: 1795-1807. doi:10.2307/2410737.

Ye XJ, Sakai K (2016) A new modified resource budget model for nonlinear dynamics in citrus
production. Chaos Solitons & Fractals 87: 51-60. doi:10.1016/j.chaos.2016.03.016.

Zhang WX, Mao PS, Li Y, Wang MY, Xia FS, Wang H (2017) Assessing of the contributions of pod
photosynthesis to carbon acquisition of seed in alfalfa (*Medicago sativa* L.). Scientific
Reports 7: 1-7 (e42026). doi:10.1038/srep42026.

Zimmerman MH, Brown CL (1971) Trees: structure and function. Springer, New York, pp.

2284

2285

Supporting information

Suppl. material 1. Radiometry at Bulu, Ndian: calibration and interpretation.

Suppl. material 2. Evidence table for the levels of mast fruiting per year.

Suppl. material 3. Time series analysis of dry season climate variables.

Suppl. material 4. Regression model fits of masting on rainfall and radiation.

Suppl. material 5. Rainfall and 30-day rainfall totals defining dry seasons.

Suppl. material 6. Description and feeding of the yellow-green caterpillar.

Suppl. material 7. Analysis of Ndian PAMOL oil-palm plantation yields.

Suppl. material 8. Stem increments from tree stem cores.

Year/dry season	Masting event	Start (days from 1.12)	Duration (days)	Mean daily rainfall (mm)	Mean daily radiation (W m^{-2})	Minimum temperature (°C)
2004–05		20	65	1.55	148	18
2005–06		39	38	2.50	157	22
2006–07	M	12	85	1.01	152	19
2007–08[a]		47	50	1.71	158	19
2008–09		34	21	4.66	137	22
2009–10	M	4	58	1.42	152	22
2010–11		26	43	2.07	158	19
2011–12[b1]		21	48	1.95	163	21
2012–13	m	26	54	1.90	165	20
2013–14 [b2]	m	39	21	0.95	165	21
2014–15		32	47	2.22	173	20
2015–16	m	13	92	1.02	154	21
2016–17	m	26	83	2.19	167	22

9   **Table 2.** Mean (± se) concentrations (mg g^{-1}) of the macro-elements in leaf litterfall of *Microberlinia bisulcata.* Collections were made in the dry

10   season immediately following the mast fruiting of 2010, and in the second and third seasons following the earlier masting of 1989 (t). NA = no

11   data available. Values from collection years with small superscripted letters (see footnotes) are taken from the earlier studies cited, otherwise the

12   values are newly reported (see Methods).

$t_{>mast}$ (n)	year	N	P	Ca	K	Mg
1 (30)	2011	16.06 ± 0.25	0.597 ± 0.018	14.71 ± 0.58	4.41 ± 0.11	3.40 ± 0.13
1 (20)	2002[a]	17.39 ± 0.33	0.632 ± 0.026	NA	5.58 ± 0.45	3.43 ± 0.17
2 (5)	1991[b]	15.72 ± 0.66	0.692 ± 0.025	13.99 ± 0.82	4.67 ± 0.23	3.89 ± 0.17
3 (5)	1992[b]	19.33 ± 0.55	0.809 ± 0.047	12.70 ± 0.90	2.44 ± 0.07	2.53 ± 0.09
<1 (6)	1998	18.27 ± 0.64	0.836 ± 0.057	NA	NA	NA

14   [a], Schwan (2003); [b], Chuyong (1994) and Chuyong et al. (2000).

16 **Table 3.** Concentrations of the five macro-elements in fallen mature seeds and pods (= fruits) of *M. bisulcata*. Collections were made during the

17 1998 masting event, together with concentrations of the same in mature leaves of this species collected in the same year of a previous (1992)

18 event. Prominent values are shown in boldface. Values with small superscripted letters a – c (see footnotes) are from the earlier studies cited,

19 otherwise the values are newly reported (see Methods).

	Concentration (mg g^{-1})			Amount (kg ha^{-1} yr^{-1})^c				Ratio	Reproductive
	Seed	Pod	Leaf^b	Seed	Pod	Fruit	Leaf	[fruit: leaf]	allocation (%)^d
N	14.5^a	3.80^a	22.3	0.770	3.728	4.498	41.924	0.107	9.69
P	1.80^a	0.24^a	1.18	0.096	0.235	0.331	2.218	0.149	12.99
K	7.40	6.13	5.91	0.393	6.014	6.407	11.111	**0.577**	**36.57**
Mg	3.01	1.17	3.77	0.160	1.148	1.308	7.088	0.185	15.58
Ca	3.77	2.17	13.2	0.200	2.129	2.329	24.816	0.094	8.58

^{a} as reported in Table 1 of Green and Newbery (2002); ^{b} mature, green leaves, from Chuyong et al. (2000); ^{c} using estimates of oven-dry mass per

ha from Table 1 of Green and Newbery (2002): seed, pods and fallen litter: 53.1, 981.1, 1880 (kg ha^{-1}) respectively; ^{d} RA = fruit density/(fruit +

leaf densities).

Figure Legends

Fig. 1. The time series of mast fruiting of *Microberlinia bisulcata* at Korup, 1989-2017. (a) The

observed data. Symbol colours: brown, full; orange, partial; and green, no; masting event

recorded. 'C' marks the year 2012, in the dry season of which the strong caterpillar attack

occurred, and 'c' the one in 2014 which was a smaller more localized attack. (b, c) The

probability of masting as fitted for 1989-2004 (squares), and predicted from 2005-2017

(triangles), using binomial logistic regressions of masting on the logarithm of rainfall in

respectively the current and prior year's dry season. Symbol colours: brown, full; orange, partial;

and green, no; masting event recorded.

Fig. 2. Times series of the dry season climatic variables associated with the masting series. (a)

mean daily rainfall; (b) mean daily radiation; (c) start date (from 1 December of year prior) and

(d) duration of the season, i.e. that which preceded the masting event; (e) radiation intensity (=

radiation x duration·10^{-3}); and (f) the difference in radiation ($radi_d$) between successive years

(current minus prior). Symbol colours: brown, full; orange, partial; and green, no; masting event.

Fig. 3. The course of soil moisture content (SMC, %), shown as means and SEs, at (mostly) 40

cm depth, for 2003-2005 at Korup. The horizontal red lines indicate the span of the dry season

for each year (see also Suppl. material 5, fig. 1).

Fig. 4. Changes in the mean scores of five phenological stages of *M. bisulcata* trees at Korup in

(a) 2009-2010 and (b) 2011-2012. Stages: leaf fall (blue), leaf flush (purple), flowering (green);

immature (yellow) and mature (red) pods. The horizontal red lines indicate the span of the dry

season for each year (see also Suppl. material 5, fig. 2). Dates of caterpillar attacks indicated

with 'C'.

Fig. 5. A schematic flow diagram of the interrelated hypothetical processes behind mast fruiting (M) in the grove-forming population of *Microberlinia bisulcata* at Korup. The boxes with light blue shading are either empirically established in the present study, or at least can be strongly inferred from the literature. They involve the deterministic factors soil phosphorus (P) and potassium (K), and the stochastic effects of dry-season (DS) rainfall; SMC means soil moisture content, and ECM stands for ectomycorrhizal. Those in grey concern radiation and carbon (C)-supply, dynamics and allocation. The box in pale brown-orange concerns the likely important role of ectomycorrhizal networks and nutrient fluxes within the root mat, which are hypothetical and require empirical confirmation. The central double-headed arrow represents the interplay between within-tree and between-tree processes. Green shading and lines indicate the proposed threshold triggering flowering and fruiting, and repeated annual uptake and storage over successive years within the masting cycle of typically 2-3 years. Hormonal control of masting, shown in yellow, is strongly suspected: it also requires critical testing. The small polygon in lilac, centre-left, is the primary condition for mast fruiting to occur, as shown by statistical model fitting in this paper (t is time in years). It is the process-link between the start and end of the diagram. Changes in the degree with which flowering results in fruiting, and in the intensity of fruiting itself (e.g. when lowered to 'm'), are explained in the main text.

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5