Targeting the TCA cycle can ameliorate widespread axonal energy deficiency in neuroinflammatory lesions

Yi-Heng Tai[1,2,3], Daniel Engels[1,2], Giuseppe Locatelli[1,2], Caroline Fecher[3,4,+], Delphine Theodorou[1,2], Stephan A. Müller[4,5], Simon Licht-Mayer[6], Mario Kreutzfeldt[7], Ingrid Wagner[7], Ioanna Emmanouilidis[1,2,3], Sofia-Natsouko Gkotzamani[1,2], Laura Trovo[3], Arek Kendirli[1,2], Almir Aljovic[1,2], Michael O. Breckwoldt[1,2,3,++], Ronald Naumann[8], Florence M. Bareyre[1,2,9], Don Mahad[6], Doron Merkler[7], Stefan F. Lichtenthaler[4,5,9], Martin Kerschensteiner[1,2,9]*# and Thomas Misgeld[3,4,9]*#

[1] Institute of Clinical Neuroimmunology, University Hospital, Ludwig-Maximilians Universität (LMU) München, Munich, Germany

[2] Biomedical Center (BMC), Faculty of Medicine, Ludwig-Maximilians Universität München, Martinsried, Germany

[3] Institute of Neuronal Cell Biology, Technical University of Munich, Germany

[4] German Center for Neurodegenerative Diseases (DZNE), Munich, Germany

[5] Neuroproteomics, School of Medicine, Klinikum rechts der Isar, Technical University of Munich, Germany

[6] Centre for Clinical Brain Sciences, University of Edinburgh, Edinburgh, UK

[7] Department of Pathology and Immunology, Division of Clinical Pathology, University & University Hospitals of Geneva, Switzerland

[8] Transgenic Core Facility, Max Planck Institute of Molecular Cell Biology and Genetics, Dresden, Germany

[9] Munich Cluster for Systems Neurology (SyNergy), Munich, Germany

+Present address: Washington University in St. Louis, Department of Cell Biology and Physiology, St. Louis, USA

++ present address: Department of Neuroradiology, University Hospital Heidelberg, Im Neuenheimer Feld 400, 69120 Heidelberg, Germany

#Co-senior authors

*Correspondence to:

        M.K. (martin.kerschensteiner@med.uni-muenchen.de)

        T.M. (thomas.misgeld@tum.de)

Words: 12,302 words (excluding abstract, methods, fig legends, refs)
Figures: 6 (+ 5 Extended Data Figures + 1 Extended Data Table)
References: 52

ABSTRACT

Inflammation in the central nervous system (CNS) can impair the function of neuronal mitochondria and contributes to axon degeneration in the common neuroinflammatory disease multiple sclerosis (MS). Here we combine cell type-specific mitochondrial proteomics with in vivo biosensor imaging to dissect how inflammation alters the molecular composition and functional capacity of neuronal mitochondria. We show that neuroinflammatory lesions in the mouse spinal cord cause widespread and persisting axonal ATP deficiency, which precedes mitochondrial oxidation and calcium overload. This axonal energy deficiency is associated with impaired electron transport chain function, but also an upstream imbalance of tricarboxylic acid (TCA) cycle enzymes. Among these, isocitrate dehydrogenase 3 (Idh3), a rate limiting enzyme of the TCA cycle, is depleted in neuronal mitochondria in experimental models and in MS lesions. Notably, viral overexpression of an Idh3 subunit ameliorates the axonal energy deficits in neuroinflammatory lesions, suggesting that TCA cycle dysfunction in MS may be amendable to therapy.

KEYWORDS

INTRODUCTION

Multiple sclerosis (MS) is a common neurological disease in which CNS inflammation results in myelin loss and progressive neurodegeneration. While the importance of such neurodegenerative processes for long-term disability of MS patients is well established[1-3], the mechanistic links between CNS inflammation and neurodegeneration remain to be resolved. Emerging evidence from MS patients and models indicates that neuronal mitochondria could be critical hubs that render inflammatory signals into neurodegenerative consequences. This concept is not only supported by the essential function of mitochondria in neuronal energy homeostasis[4], but also by previous studies showing: (i) that damaged mitochondria accumulate in axons located in experimental and human neuroinflammatory lesions[5-7], (ii) that neuronal mitochondria acquire DNA deletions over the course of the disease[8], and (iii) that such inflammation-induced mitochondrial damage impairs electron transport chain (ETC) function[2,9].

Mitochondrial damage could thus be initiated already in the highly inflamed lesions of early MS and then further amplify over the course of the disease. This process would provide a link not only between inflammation and neurodegeneration, but also between initial relapsing-remitting and later progressive pathology[2,9]. As a result of these insights, mitochondria have emerged as a promising therapy target to prevent neurodegeneration throughout the disease course of MS. Unfortunately, so far, such strategies have failed to provide robust clinical benefits, at least in large clinical trials[10]. One of the reasons underlying this failure is that our understanding of how the molecular machinery and functional capability of mitochondria is impaired in neuroinflammatory lesions is still incomplete. Moreover, most studies so far have focused on ETC impairments[8,11], which might be hard to target therapeutically given the complex structure of the underlying macromolecular complexes and their genetics.

Here we apply novel in vivo imaging strategies combined with selective ex vivo proteomic

analysis of neuronal mitochondria in a murine MS model (experimental autoimmune encephalomyelitis, EAE) to investigate the molecular underpinnings and functional consequences of inflammation-induced mitochondrial pathology. We reveal that widespread axonal ATP deficiency is initiated in acute neuroinflammatory lesions and persists in the chronic disease stage. These bioenergetic deficits precede mitochondrial redox or calcium dyshomeostasis. Selective *MitoTag*-based proteomic analysis of neuronal mitochondria isolated from acute and chronic EAE spinal cords instead revealed a imbalance of critical tricarboxylic acid (TCA) cycle enzymes, with a prominent loss of a rate-limiting enzyme, isocitrate dehydrogenase 3 (Idh3). Notably, Idh3 depletion in neuronal mitochondria is also apparent in human MS lesions. Finally, we show that viral gene therapy, overexpressing the catalytic subunit of Idh3, can counteract axonal ATP deficits in neuroinflammatory lesions. Our study thus provides a refined understanding of how the molecular composition of neuronal mitochondria changes in response to neuroinflammation. We identify the depletion of Idh3 as a critical mediator of the axonal energy crisis that occurs in neuroinflammatory lesions, thus defining a potential target for therapeutic intervention.

RESULTS

Pervasive and persistent axonal ATP deficits emerge early in neuroinflammatory lesions

As ATP production is a central function of mitochondria, we established an in vivo imaging approach to record the ATP/ADP ratio of individual axons in neuroinflammatory lesions[12]. We generated *Thy1*-PercevalHR mice, in which neurons express PercevalHR, a genetically-encoded excitation ratiometric biosensor that monitors the cytoplasmic ATP/ADP ratio[13]. Histological characterization confirmed the targeting of the sensor to a broad range of neuronal populations in the brain and spinal cord (**Extended Data Figure 1a**). To assess whether the sensor was indeed capable of recording ATP deficits in spinal axons, we pharmacologically interfered with ATP production and recorded the ATP/ADP ratio by measuring fluorescence emission after excitation at 950 and 840 nm using in vivo 2-photon microscopy. Application of either the glycolysis inhibitor IAA (100 μM) or the mitochondrial uncoupling agent CCCP (100 μM) to the exposed dorsal spinal cord resulted in swift reduction of the ATP/ADP ratio in spinal axons (**Extended Data 1b,c**).

To assess the emergence of axonal energy deficits in neuroinflammatory lesions, we next induced EAE, a widely used model of MS, by immunizing *Thy1*-PercevalHR mice with myelin-oligodendrocyte glycoprotein (MOG). In vivo 2-photon imaging then allowed us to record the ATP/ADP ratio in axons that cross through a neuroinflammatory lesion and at the same time to stage axonal morphology, a predictor of axonal fate as we established previously[6]. We found widespread axonal ATP deficits that not only affected axons that were swollen (stage 1 axons) or fragmented (stage 2) and had hence already entered the focal axonal degeneration (FAD) process[6,14], but also morphologically normal axons (stage 0; **Figure 1a-e**). Axon were not only showing a reduced ATP/ADP ratio in acute lesions (analyzed 2 days after EAE symptom onset), but ATP deficits persisted in axons of all FAD stages in chronic lesions (analyzed an additional 20 days later; **Figure 1d,e**). To explore, whether neurons in

their entirety lacked ATP, or whether this was a local axonal deficit, we traced individual sensory axons from the lesion back into the dorsal roots, where neuroinflammation is typically not present in EAE. Comparing the ATP/ADP ratios in the same axons close to and far from the intra-spinal lesion area revealed that ATP deficits were most pronounced at the sites of CNS inflammation (**Figure 1f-g**). This finding is in line with the localized accumulation and damage of mitochondria that we have previously described in spinal cord lesions of the same MS model[6,14,15].

Axonal ATP deficits precede mitochondrial redox and calcium dyshomeostasis

Oxidative stress to axons had previously been proposed as a mediator of mitochondrial damage in neuroinflammatory lesions[1-3,9,16]. Thus, we set out to determine if and when an altered redox status of axonal mitochondria would emerge in relation to the onset of neuroinflammation-induced ATP deficits. We used *Thy1*-mitoGrx-roGFP mice, which allow recording the glutathione redox potential in neuronal mitochondria, as we previously established[17]. After crossing these redox reporter mice to *Thy1*-OFP mice[18] to allow staging of axonal morphology, we induced EAE and performed ratiometric in vivo confocal imaging using single photon excitation at 405 and 488 nm at the peak of acute disease symptoms. An oxidative shift in mitochondrial redox state was only apparent in fragmented (stage 2) axons, while mitochondria in normal-appearing (stage 0) and swollen (stage 1) axons were unaltered compared to control axons in a healthy spinal cord (**Figure 2a-d**). This is notable, as the analysis of individual mitochondria confirmed that damage-related mitochondria shape changes, such as rounding up (reduced shape factor), are already present at the onset of FAD (**Figure 2e**)[6,15]. Decompensated mitochondrial redox stress thus seems to be a late event during inflammatory axon degeneration and affects only a small proportion of non-fragmented axons in neuroinflammatory lesions. Oxidative mitochondrial damage per se is hence unlikely to underlie the widespread axonal ATP deficits in EAE.

In addition to oxidative damage, calcium overload was proposed to mediate mitochondrial dysfunction[1,19]. To monitor calcium handling of axonal mitochondria in vivo, we generated a transgenic reporter mouse line that expresses the ratiometric calcium sensor Twitch2b[20] selectively in neuronal mitochondria. Histological characterization of these *Thy1*-mitoTwitch2b mice confirmed widespread sensor expression in CNS neurons. Moreover, induction of localized laser lesions in the spinal cord demonstrated that the sensor can record relevant pathological calcium signals in vivo (**Extended Data Figure 2**). We then crossed these reporter mice to *Thy1*-OFP mice to relate axonal morphology to mitochondrial calcium levels and shape in acute EAE lesions. Similar to our analysis of mitochondrial redox state, we found that mitochondrial calcium levels are primarily increased in fragmented (stage 2) axons, i.e., during the end stage of FAD, while normal appearing (stage 0) and swollen (stage 1) axons showed no pronounced calcium dyshomeostasis, despite clear morphological signs of organelle damage (**Figure 2f-j**).

Taken together, these experiments indicate that overt dysregulation of the mitochondrial redox state and calcium handling are rather late events during inflammatory axon degeneration. Hence, a molecularly distinct dysregulation of mitochondrial function and molecular composition must explain the prodromal and pervasive ATP deficits in EAE axons.

Proteomics of neuronal mitochondria reveals ETC and TCA cycle enzyme depletion

To obtain an unbiased characterization of the neuroinflammatory changes to the molecular make-up of neuronal mitochondria, we adapted a cell type-specific proteomics approach that we recently established[21]. For this, we performed intrathecal injections of a rAAV.hSyn:Cre virus into the ventricles of *MitoTag* neonates, which resulted in the tagging of the outer membrane of neuronal mitochondria with GFP. This enabled selective isolation and subsequent mass spectrometry analysis of these mitochondria from the spinal cords of healthy mice, as well as from different stages of EAE (**Figure 3a,b** and **Extended Data Figure 3a-c**),

which revealed pronounced changes to the mitochondrial proteome of neurons. These changes could not be predicted by protein lifetime, location inside the mitochondria or whether a protein was encoded in the nucleus vs. the mitochondrial DNA (**Extended Data Figure 3d-f**). Gene set analysis of the major dysregulated pathways converged on the ETC and the TCA cycle, which are abundant in healthy neuronal mitochondria, but depleted in acute neuroinflammatory lesions (**Figure 3c,d** and **Extended Data Figure 3g,h**). The ETC complexes were uniformly affected, typically more strongly in acute than in chronic EAE lesions (**Figure 3d** and **Extended Data Figure 3g,h**). This corresponded with reduced axon complex IV (COX IV) activity as measured in situ by a histochemical assay[22] (**Extended Data Figure 4b,c**). In addition, TCA cycle enzymes showed pronounced dysregulation (**Figure 3d,e**), prominently affecting the subunits of isocitrate dehydrogenase 3 (Idh3), and to a lesser extent malate dehydrogenase 2 (Mdh2). Idh3 is of particular interest as a rate-limiting enzyme of the TCA cycle, which mediates the irreversible oxidative decarboxylation of isocitrate generating NADH, which is subsequently used for ATP generation by the ETC. Notably, Idh2, which catalyzes the reversible NADP-dependent conversion of isocitrate to α-ketoglutarate, appears to be differentially affected, as its abundance is unchanged in acute and increased in chronic neuroinflammatory lesions (**Figure 3d,e**). Expression changes of key TCA cycle related enzymes, including Idh3 and Idh2, were also present on the transcriptional level as revealed by analysis of published RiboTag translatomes of spinal motoneurons in EAE[23] (**Figure 3f**). Hence, neuroinflammation caused a marked alteration in the expression of TCA cycle enzymes, including the rate-limiting enzyme Idh3, which likely hinders neuronal mitochondria from generating sufficient amounts of ATP.

Idh3 is depleted in neuronal mitochondria not only in EAE but also in MS lesions

We next aimed to confirm the proteomics results by immunofluorescence analysis of neuronal mitochondria in EAE as well as in MS lesions. First, we quantified the expression

levels of a subset of TCA cycle enzymes, for which the neuronal *MitoTag* analysis predicted dysregulation in EAE. We performed immunostainings on spinal cord sections from *Thy1-mitoRFP* mice[17], where analysis can be restricted to red fluorescent protein (RFP)-tagged neuronal mitochondria. Indeed, the levels of Idh3a (the catalytic subunit of Idh3) as well as Mdh2 were decreased by more than 50 % in EAE lesions compared to controls (**Figure 4a-e**). As predicted by our proteomics analysis, the expression of Idh2 was slightly increased (**Figure 4f,g**). Second, to explore whether similar expression changes of TCA cycle enzymes were also present in MS lesions, we performed immunofluorescence analysis of Idh3a in brain biopsy and autopsy sections from 7 MS patients (**Extended Data Table 1**). As fixation protocols are typically variable in patient-derived material, we internally normalized the expression of a given TCA cycle enzyme between the lesion area and the adjacent normal-appearing white matter (NAWM) on the same section (**Figure 5a**). This analysis revealed a reduction of Idh3a in MS lesions (**Figure 5b,c**), which also affected chronic active MS lesions indicating that Idh3 depletion persists long-term. Taken together, our findings show that TCA cycle disruption is a persistent neuronal change during MS lesion formation and progression.

Restoring Idh3a expression rectifies axonal ATP deficits in neuroinflammatory lesions

Considering the pronounced depletion of Idh3a in EAE and MS, as well as its critical role in the TCA cycle, we explored whether overexpression of this catalytic subunit, could restore axonal ATP levels in neuroinflammatory lesions. For this, we expressed Idh3a and a fluorescent marker (tdTomato) under a pan-neuronal promoter (human Synapsin) via systemic injection of a rAAV.PHP.eB virus[24]. Confocal analysis confirmed that a substantial fraction of CNS neurons was transduced with the rAAV.hSyn:Idh3a-tdTomato virus as indicated by tdTomato expression. Moreover, tdTomato-labelled neurons in EAE spinal cords showed a marked increase in mitochondrial Idh3a expression (**Extended Data Figure 5**). We then directly assessed the effects of Idh3a restoration on axonal energy deficits in neuroinflammatory lesions,

by inducing EAE in *Thy1*-PercevalHR mice injected with either the rAAV.hSyn:Idh3a-tdTomato virus or a control rAAV.hSyn:cre-tdTomato virus, carrying a similarly-sized, but in our setting inert transgene. In vivo imaging of the surgically exposed spinal cord in EAE-induced animals again confirmed strong tdTomato-expression in a fraction of axons, while other PercevalHR-expressing axons were tdTomato-negative and provided an internal control (**Figure 6a,b**). Compared to this control axon population (which rules out general effects on lesion activity by the virus), transduction with a rAAV.Syn:Idh3a-tdTomato virus significantly increased the ATP/ADP ratio and thus partially restored the axonal energy deficiency in neuroinflammatory lesions (**Figure 6c**). No such rescue was observed in axons transduced with the control rAAV.hSyn:cre-tdTomato (**Figure 6d**). Notably, ATP/ADP ratios rose in axons of all damage stages, suggesting that correcting Idh3a levels could also counteract ATP deficiencies in more advanced stages of FAD. While we could only achieve partial restoration of ATP levels in these experiments, the finding that a widespread improvement of axonal energy homoeostasis could be achieved by solely supplementing the catalytic subunit of a rate-limiting enzyme reveals the potential of targeting the TCA cycle to rectify the axonal energy crisis in neuroinflammatory conditions.

DISCUSSION

That axons in neuroinflammatory lesions exist in a state of 'energy failure' linked to mitochondrial damage has long been inferred based on (1) abnormalities in mitochondrial density, dynamics and morphology; (2) mitochondrial dysfunction as revealed by in situ assays of respiratory complex function or mitochondrial polarization; and (3) the accumulation of mitochondrial DNA mutations in inflamed CNS tissue[1,2,16,20]. Furthermore, there is molecular evidence that axons in MS lesions might exist in a hypoxic state, which would further restrict mitochondrial respiration[25]. These data have directed most attention to the ETC, which covers most of the uniquely high energy demands of neurons and determines the mitochondrial potential and other aspects of respiration that can be assayed in situ, e.g. using enzymatic assays. Thus, the focus on the ETC is both well justified by its importance, but also biased by experimental accessibility. At the same time, the ETC is not necessarily a good therapeutic target, given its complex molecular make-up at the protein, as well as at the genome level, the extremely low turnover of some of its components and its Janus-faced nature as the major source not only of ATP, but also of oxygen radicals. Finally, the ETC needs to be fed with a steady and sufficient flow of redox equivalents and other substrates, most of which derive from the TCA cycle. Hence the TCA cycle – as the 'upstream' bioenergetic hub of metabolism – also deserves attention in the context of the presumed 'energy failure' of axons in neuroinflammation[26]. Especially so, as the TCA cycle is increasingly emerging as a metabolic hub relevant to neurodegeneration[27] and a 'druggable' target, for example in cancer[28,29].

Against this backdrop, the present study makes the following contributions. (1) We demonstrate directly by *in vivo* biosensor measurements that most axons in neuroinflammatory lesion exist in a state of depressed cytoplasmic ATP availability, already during early stages of the axon degeneration pathway. Moreover, this change precedes overt signs of mitochondrial redox or calcium dyshomeostasis. (2) We show by molecular profiling of neuronal

mitochondria not only a disruption of the ETC, but a similarly profound alteration of the 'up-stream' TCA cycle, where amongst others the pace-making enzyme Idh3 is depleted in neuronal mitochondria in EAE and MS lesions. (3) Finally, we demonstrate that over-expressing the catalytic Idh3a subunit suffices to boost axonal ATP provision suggesting that the full extent of the axonal energy crisis is due to a dual deficiency, both of redox substrate provision from the TCA cycle to the ETC, as well of oxidative phosphorylation itself.

Together, these results thus expand our understanding of the molecular pathogenesis of immune-mediated mitochondrial damage and propose new avenues for therapeutic intervention. They also raise the question where the dysregulation of the TCA cycle originates. Overall, the pattern of mitochondrial proteome changes does not relate to any simple pattern that we could discern. While the overall mitochondrial mass in neurons cannot be estimated easily by this method, the data suggest that within the mitochondrial proteome of neurons, specific dysregulation happens, as e.g. some biosynthetic pathways (such as mitochondrial translation) are upregulated, while most TCA cycle and ETC components are suppressed. Indeed, the parallel and progressive upregulation of Idh2, while Idh3 is downregulated, suggests a specific metabolic 'rewiring' response[30], rather than just a global loss of mitochondrial biogenesis or increase in mitophagy. Most likely, this response involves biosynthetic, as well as proteostatic mechanisms, as the observed dysregulation of mitochondrial proteins does neither correspond to whether a protein is encoded in the mitochondrial DNA (where mutations are known to accumulate in MS)[2,8] vs. in the nucleus, nor with the estimated lifetime[31]. Indeed, the correlation of published neuronal translatome data from a similar model[23] with our neuronal proteomes provides direct support for an element of 'anterograde regulation' from the nucleus, while the long life-times of some down-regulated enzymes (including subunits of Idh3[32]) argue for a parallel degradation, given the swift development of the proteomic phenotype in our acute EAE model (which is present in acute lesions two days after disease onset).

Irrespective of the origin of the reduced ATP/ADP ratio that our biosensor measurements reveal, the further question arises what this implies for axonal survival, and hence – given the central role of axon degeneration in MS disease course[33] – for possible lasting functional consequences. While limited ATP availability is often cited as a sufficient argument to assume axonal demise, and profound and lasting ATP deficiency can clearly cause axon degeneration[34], our observations rather argue for a chronic 'dystrophy' than for an acute bioenergetic collapse of axons in neuroinflammation. Indeed, we find reduced ATP/ADP ratios in the majority of axons, while we know from previous work that only a subset of these axons is at immediate risk of degeneration, and some even recover morphological and mitochondrial integrity[6]. Notably, a similarly pervasive pattern of dysfunction has been apparent in our previous studies of axonal transport[15]. As axonal transport is highly ATP dependent, and in converse, mitochondria are a major axonal transport cargo[4], a vicious cycle could result in the long-term break-down of the ATP supply chain – which might not cause immediate axon degeneration, but rather subacute dysfunction. Furthermore, previous reports of in situ ATP measurements in mouse white matter tracts have shown that neuronal activity puts an immediate strain on energy supply[35], which could sensitize specific activity-related patterns of degeneration.

So how to counterbalance the axonal energy deficiency? Indeed, 'brain energy rescue' has been hailed as a possible therapeutic strategy across neurological disorders[36]. The most direct approach would be the provision of energy substrates or key co-factors for the function of bioenergetic enzymes. Indeed, first trials along these lines – e.g. providing biotin – have been conducted in MS, but failed[37]. Notably, biotin is required for the activity of enzymes that feed into the TCA cycle upstream of Idh3 and Mdh2, which are the most dysregulated TCA enzymes in EAE. Our data on TCA cycle disruptions – combined with the previous insights on ETC dysfunction – would argue that simply feeding a substrate into a broken supply chain will indeed not be efficient. Therefore, key bottlenecks in the ATP-providing metabolic pathways need to be unblocked in addition. Our data now argue that Idh3 – a key 'pace-maker' enzyme of the

TCA cycle[38] – might be such a bottleneck and that a rather focussed intervention, such as the genetic or pharmacological upregulation of its catalytic subunit, could result in marked changes to the neuronal energy state. Indeed, the fact that Idh3 activity should be directly allosterically stimulated by a low ATP/ADP-ratio[39] and activated by calcium[40], two changes that are induced in axons in response to an inflammatory challenge, could argue in favour of Idh3a overexpression having an unblocking effect on the TCA cycle.

Interestingly, the Idh family of enzymes is the focus of substantial translational interest[41-43], given the key role of Idh1 and Idh2 in cancer biology. Notably, also Idh3 appears to be dysregulated in some neoplasms, including glioblastomas[44]. This has resulted in the development of Idh targeting drugs, e.g. for certain leukaemia and brain tumor subtypes[42-44]. In cancer, the detrimental consequence of Idh mutations appears to be metabolic rerouting that enhances tumor cell proliferation and immune cell dysregulation[45]. Indeed, also in neuroinflammation some aspects of axonal dysfunction resulting from TCA cycle disruptions could be due to metabolite imbalances rather than mere lack of efficient fuelling of the ETC and hence ATP deficiency. In this context, the concomitant upregulation of Idh2 in chronic stages of our model is notable and raises important questions for future exploration. For instance, could the available Idh blockers – most of which are deliberately not targeting Idh3 – still be useful in forcing metabolic flow back into the TCA cycle? Similarly, could approaches of increasing mitochondrial mass (such as overexpression of PGC1α, which is beneficial in inflammatory demyelination and regulates Idh3a expression as part of its broad transcriptional effects[11,43]) be synergistic with targeting the TCA cycle to unblock energy provision in axons and resolve their dystrophy in conditions like MS? The identification of the TCA cycle and some of its key enzymes as a disease contributor and a potential therapy target indeed opens the possibility for such a comprehensive evaluation of strategies to mitigate the neuronal energy crisis in the inflamed CNS.

ACKNOWLEDGEMENTS

We would like to thank A. Schmalz, J. Schmitt, K. Plesniar, B. Fiedler, Y. Hufnagel and A. Berghofer for excellent technical assistance, D. Matzek, B. Stahr, M. Korica, N. and M. Budak for animal husbandry. We also acknowledge N. de Mello and F. Perocchi (Helmholtz Center Munich/TUM) for advice on bioenergetics measurements; K. Dyar (Helmholtz Center Munich) for advice on metabolomics, O. Griesbeck (Max Planck Institute of Neurobiology) for constructs, J. Lichtman (Harvard University) for *Thy1*-OFP mice; and C. de la Rosa (BMC/LMU) support for PHP.eB virus establishment.

This project was supported by the Deutsche Forschungsgemeinschaft (DFG) via TRR 274/1 (Projects B03, C02, C05, Z01, Z02 – ID 408885537). Work in M.K.'s laboratory is further financed through grants from the DFG (TRR128, Project B10 and B13), the European Research Council (ERC) under the European Union's Seventh Framework Program (FP/2007-2013; ERC Grant Agreement n. 310932), the German Federal Ministry of Research and Education (BMBF; Competence Network Multiple Sclerosis) and the "Verein Therapieforschung für MS-Kranke e.V.". Work in T.M.'s lab is supported by the DFG (CRC870 A11-ID 118803580, Mi 694/8-1, Mi 694/9-1 A03-ID 428663564, FOR Immunostroke) and the ERC under the European Union's Seventh Framework Program (FP/2007-2013; ERC Grant Agreement n. 616791). T.M. and S.L. are members of and supported by the German Center for Neurodegenerative Diseases (DZNE). M.K. and T.M. were further supported by the DFG through a common grant (Ke 774/5-1/Mi 694/7-1) and – together with S.L. and F.B. – receive support from the Munich Center for Systems Neurology (SyNergy EXC 2145; Project ID 390857198). D.M. is supported by the Swiss National Science Foundation (SNSF; 310030B_201271 & 310030_185321) and the ERC (865026). M.O.B. was recipient of a doctoral fellowship from the Gertrud Reemtsma Foundation and supported by the Emmy Noether Program of the DFG (BR 6153/1-1). G.L. and L.T. were supported by EMBO Fellowships, G.L. further received a SNSF fellowship. Y.T.

AUTHOR CONTRIBUTIONS

M.K. and T.M. conceived and designed the experiments. Y.T. performed in vivo imaging and disease modeling experiments, with contributions from G.L. and D.T. for mitochondrial calcium and redox imaging. C.F., G.L, D.T., R.N., and M.B. established and characterized mouse lines for in vivo imaging, L.T., C.F. and Y.T. established, supported and performed the MitoTag analysis, for which S.M. and S.L. did the proteomic analysis. D.E. did the bioinformatics analysis with input from S.M. and S.L. M.Kr., I.W. and D. Me. did histological analysis of MS, while Y.T., I.E. and S.G. did mouse histology and S.M.-L. and D. Ma. contributed the COX IV assay in EAE. A.A., A.K., Y.T. and F.B. designed and generated the PHP.eB viruses. Y.T. coordinated and contributed data analysis across all experimental approaches and designed the final data representation with input from M.K. and T.M.. Y.T., M.K. and T.M. wrote the paper with input from all authors.

ONLINE METHODS

Animals

All experiments were performed on either postnatal day 3 pups or adult mice according to the protocols on a C57BL/6 (strain designation C57BL/6J, Jackson Laboratories) background. All animals were bred and housed under standard conditions with a 12-hours light/12-hours dark cycle. Food and water for mice were provided ad libitum. Female and male were equally allocated into control and experimental groups if not explicitly mentioned otherwise. All experimental procedures were conducted in accordance with regulations of the relevant animal welfare acts and protocols approved by the responsible regulatory office.

To measure neuronal mitochondrial redox changes in spinal axons, we used the *Thy1*-mitoGrx-roGFP mouse line, which selectively expresses the redox sensor Grx-roGFP2 in neuronal mitochondria[17]. To visualize axonal morphology, these mice were crossed to *Thy1*-OFP mice[18], in which neurons are labelled by cytoplasmic expression of an orange fluorescent protein (OFP). To investigate the proteomic profile of neuronal mitochondria in the context of neuroinflammation, we used the *Rosa26* MitoTag knock-in mouse line that allows a Cre-dependent expression of GFP targeted to the outer mitochondrial membrane[21]. *Thy1*-mitoRFP mice, in which tagRFP is localized to the matrix of neuronal mitochondria[17] were used to ascertain the localization of immunofluorescence signals to neuronal mitochondria.

Generation of reporter mouse lines

To measure the neuronal ATP-to-ADP ratio, *Thy1*-PercevalHR mouse lines were generated using a blunt-end cloning strategy with Perceval GW1-HR plasmid (Addgene, #49082) purchased from Addgene[13]. In brief, the GW1-Perceval-HR plasmid was digested with Xba1 and EcoRI enzymes, blunted, and inserted into the Thy1-vector cut beforehand with XhoI, blunted, and dephosphorylated. After ligation and electroporation, minipreps of ampicillin-

selected E. coli cultures were performed, followed by verifying the correct plasmid orientation and correct enzyme digestions. The construct was transfected in human embryonic kidney 293T (HEK) cells, and the cytoplasmic expression was confirmed by confocal microscopy. Maxipreps of the adequate cloning candidates were carried out using Qiagen kits. The Thy1-Perceval-HR plasmid was further linearized using PvuI and EcoRI restrictions enzymes.

To measure the calcium level in neuronal mitochondria, *Thy1*-mitoTwitch2b mouse lines were generated. The Twitch2b-pcDNA3 plasmid[20] was kindly provided by O. Griesbeck (Max Planck Institute of Neurobiology, Martinsried, Germany). The Twitch2b sensor was extracted using NotI restriction enzyme and inserted into the pCMV:myc-mito plasmid (Addgene, #71542) cut beforehand with NotI restriction enzyme and dephosphorylated. After ligation and electroporation, minipreps of ampicillin-selected E. coli cultures were performed, followed by verifying the correct plasmid orientation and correct enzyme digestions. The construct was transfected in HEK cells and verified by live imaging. After retesting the construct and its functional responsiveness by chemical application in HEK cell culture, maxipreps of the adequate cloning candidates were carried out using Qiagen kits. The pCMV-myc-mito-Twitch2b sequence was further extracted using PmI and XbaI, blunted and inserted into the *Thy1*-vector cut beforehand with XhoI, blunted and dephosphorylated. The *Thy1*-mitoTwitch2b plasmid was linearized using ZraI and AflIII restrictions enzymes. Sequencing of *Thy1*-mitoTwitch2b and *Thy1*-Perceval-HR constructs were performed by Eurofins. The sample purity of the linearized DNA was determined using the absorbance ratio at 260 nm vs 280 nm. The concentration of the linearized DNA used for pronuclear injections was 45.2 ng/µl in 60 µl for Thy1-Perceval-HR (with a ratio A260/280 at 1.83) and 36.5 ng/µl in 120 µl (with a ratio A260/280 at 1.83) for *Thy1*-mitoTwitch2b. The generation of the *Thy1*-PercevalHR and *Thy1*-mito-Twitch2b was conducted at the Transgenic Core Facility of the Max Planck Institute for Molecular Cell Biology and Genetics in Dresden by standard pronuclear injections into pseudo-

pregnant host mice. For analysis of mitochondrial calcium level in spinal axons *Thy1*-mitoTwitch2b mice were crossed to *Thy1*-OFP mice.

AAV vector generation and virus production

The AAV.hSyn: Cre plasmid, which encodes for Cre expression under the control of the human Synapsin promoter, was generated by replacing the DIO-GFP sequence (Addgene, #50457) to generate the AAV.hSyn backbone plasmid. In brief, the AAV.hSyn:DIO-GFP plasmid was initially digested with SalI/XhoI, followed by incubating with BglII to remove DIO.GFP and ligated with GFP.ires.Cre, which was excised from AAV.CMV:GFP.ires.Cre with BglII using Quick Ligase according to the manufacturer's instruction. Further from AAV.hSyn: GFP.ires.Cre, GFP.ires was excised with BmgBI, and the backbone was ligated to acquire AAV.hSyn:Cre plasmid. The AAV.PHP.eB-hSyn:Idh3a.P2A.tdTomato plasmid, which encodes for Idh3a.P2A.tdTomato expression under the control of the human Synapsin promotor was generated as follows. In brief, the AAV.hSyn:DIO.FGF.P2A.EGFP was initially digested with SalI/HindIII to remove DIO.FGF.P2A.EGFP, ligated with amplified Idh3a fragment (Sino biological, MG53630-UT) and amplified P2A.tdTomato sequence, which was excised from CAG.Cre.P2A.tdTomato with KpnI/EcoRI by using Gibson assembly master mix (New England BioLabs, E2611S).

The AAV.hSyn:Cre.P2A.tdTomato plasmid was purchased from Addgene (Addgene, #107738). The construct with the correctly oriented insert was introduced to Stellar Competent cells (Clontech, 636763), and the plasmid purification was performed with a Qiagen Plasmid Maxi Kit according to the manufacturer's protocol.

AAV vector packaging was performed using HEK 293T (ATCC, crl-3216) and produced as described before68. In short, HEK 293T cells were transfected with pAD-helper, AAV-capsid 9, and AAV-construct (molar ratio 1:1:1) using a RPMI: PEI incubation protocol. AAV vectors

were harvested from the supernatant with polyethylene glycol (PEG, Sigma-Aldrich, no. 25322-68-3) solution and the cell pellets. Freeze-thaw cycles were performed to lyse the cells, and the residual DNA from the packaging was further degraded with benzonase (Sigma-Aldrich, E1014). Following the purification procedure using iodixanol gradient ultracentrifugation, the virus was concentrated by subsequent centrifugation and incubation with formulation buffer (Pluronic-F68 0.001% in saline PBS). The product was then further collected (with a genomic titer of $\sim 10^{12}$ to 10^{14}) and stored in small aliquots at -80°C.

Neonatal intraventricular injection of rAAV

For the proteomic profiling of mitochondria in neuroinflammation, AAV.hSyn:Cre was injected into neonatal *MitoTag* pups (genotype: WT/MitoTag) according to a previously published protocol[46]. In short, P3 pups were anaesthetized with isoflurane (Abbott) and injected with 3 µl ($\sim 1 \times 10^{12}$ viral particles per animal) AAV9.hSyn:Cre with 0.05% (wt/vol) trypan blue to the right lateral ventricle at a rate of 30 nl/s using a fine glass pipette (Drummond; 3.5", #3-000-203-G/X) attached to a nanoliter injector (World Precision Instruments; Micro4 MicroSyringe Pump Controller connected with Nanoliter 2000) held by the stereotaxic manipulator. All surgeries were conducted under ultrasound guidance (VisualSonics; Vevo1100 Imaging System, with a Microscan MS550D transducer).

Intravenous injection of rAAV

To achieve overexpression of Idh3a or Cre in spinal axons, AAV.PHP.eB.hSyn: Idh3a.P2A.tdTomato or AAV.PHP.eB.hSyn:Cre.P2A.tdTomato was intravenously injected 14 days prior to the EAE induction. In brief, the animal was immobilized, and the tail was locally warmed to dilate the vessels facilitating the injection. The mice were injected with the virus (~1

x 10^{11} viral particles per animal) at a final volume of 120 µl, followed by disinfection of the operation area with 80% ethanol.

Induction of experimental autoimmune encephalitis (EAE)

Mice were immunized subcutaneously with 250 µl of an emulsion containing 200 µg of purified recombinant MOG (N1-125) and complete Freund's adjuvant (Sigma-Aldrich, F5506) supplemented with 10 mg/ml Mycobacterium tuberculosis (H37RA, BD Difco, 231191). The mice received intraperitoneal (i.p.) injections with 200 ng pertussis toxin at day 0 and 2. After immunization, mice were weighed daily, and neurological deficits were evaluated according to the following EAE score: 0, no clinical signs; 0.5, partial tail weakness; 1, tail paralysis; 1.5, gait instability or impaired righting ability; 2, hind limb paresis; 2.5, hind limb paresis with dragging of one foot; 3, total hind limb paralysis; 3.5, hind limb paralysis and forelimb paresis; 4, hind limb and forelimb paralysis; 5, death. The onset of disease was defined by the first day of neurological symptoms; the acute stage as 2–3 days following clinical onset and the chronic stage as 22 days following the onset of EAE in all chronic experiments.

Spinal cord laminectomy

Mice were imaged at the acute stage of EAE, two days after onset of disease (only animals with an EAE score ≥ 2.5 were included) and at onset plus 22 days for the chronic stage. The mice were anesthetized by intraperitoneal injection of medetomidine (0.5 mg/kg), midazolam (5.0 mg/kg), and fentanyl (0.05 mg/kg) and tracheotomized and intubated to minimize breathing if needed. The dorsal spinal cord was surgically exposed as previously described[6]. In short, the skin was disinfected and incised along the spinal column. The dorsal spinal cord was surgically exposed by performing a laminectomy, and the surgical field was constantly superfused with artificial cerebrospinal fluid (aCSF) to stay moisturized. The vertebral column was then

position-fixed on a spinal clamping device (Narishige STS-a), allowing controlled movement in x, y, z directions during the imaging session. A 3.5% agarose well was built up around the spinal opening and filled with aCSF. During the imaging sessions, mice were kept under constant anaesthesia, and their breathing and reflexes were controlled every 30 min. Animals showing signs of traumatic damage after laminectomy were excluded from the analysis.

In vivo multiphoton and confocal imaging

Imaging of Thy1-PercevalHR mice: to detect the axonal ATP/ADP ratio, the genetically encoded ATP/ADP indicator PercevalHR was imaged using a set of wavelengths of 950 nm and 840 nm to excite the cpmVenus fluorescent protein sequentially. The signals were collected simultaneously in cyan and yellow channels using emission barrier filter pairs (bandwidth of 455–490 nm and 526–557 nm) on the Olympus MPE-resonant scanner system. Iodoacetic acid (IAA), 10 mM, or Carbonyl cyanide m-chlorophenyl hydrazone (CCCP), 100 µM, at final concentration were added to the imaging solution to obtain the baseline signal of the PercevalHR sensor. For imaging of the effects of Idh3a-tdTomato overexpression on ATP/ADP ratio, we used the same setting as described above. In addition, tdTomato signals were recorded using a wavelength of 1040 and collected in the red channel with a barrier filter pair (bandwidth 655-725nm).

Imaging of Thy1-mitoTwitch2b × Thy1-OFP mice: to measure mitochondrial calcium levels in axons, the genetically encoded calcium indicator Twitch-2b was excited using a wavelength of 840 nm to excite mCerulean3 and cpVenusCD fluorescent proteins simultaneously[20]. The signals were collected in cyan and yellow channels. To reveal the axonal morphology, the orange fluorescent signal was excited with a wavelength of 750 nm. The customized emission barrier filter pairs with bandwidths of 457-487 nm, 500-540 nm (GaAsP detectors), and 560-600 nm (RXD2 detector) were used on the Olympus MPE-FV1200 system. Images (12 bit)

were acquired with a 25x/1.05 dipping cone water-immersion objective, a pixel size of 0.26 mm pixel^{-1} or smaller, a dwell time of 2.0 μs pixel^{-1}, and a laser power of 30 mW measured in the back focal plane. Volume stacks penetrating ~50 μm into the dorsal spinal cord from the surface were acquired with a Z-spacing of 1 μm.

Imaging of Thy1-mitoGrx-roGFP × Thy1-OFP mice: to detect the mitochondrial redox state in spinal axons, the genetically encoded redox indicator Grx-roGFP was sequentially excited at 405 and 488 nm wavelengths as previously described[17]. The signals were collected with emission barrier filter pairs (bandwidth 492–592 nm) in separate channels using a 50/50 beam splitter on an Olympus FV1000 confocal system. Smaller stacks were acquired with a Z-spacing of 1 μm at a frame rate of 0.1–0.2 Hz.

Image processing and analysis

Images were analyzed with the open-source image analysis software ImageJ (Fiji, http://fiji.sc/Fiji) and Photoshop (Adobe). To determine PercevalHR signals in axons, background intensities (non-axonal areas) were measured and subtracted for every axon in every experiment to create a background mask, and pixel-by-pixel ratios were calculated from the mean over three regions of the same axon and further normalized to the mean value of the control mice. For representative images, maximum intensity projections of image stacks were gamma-adjusted and processed with a despeckle filter using Photoshop software (Adobe). To measure the redox state of a single mitochondrion or clusters of mitochondria signals in *Thy1*-mitoGrx-roGFP × *Thy1*-OFP, the mean intensity values were measured in the 405 and 488 nm channels as previously described[17]. The values for the two channels were divided (405/488 nm) to obtain a ratio related to the redox state of the sensor. This ratio was normalized to the mean value of the control mice. Mitochondrial morphology is quantified as the mitochondrial shape factor calculated by dividing the length by the width of a single mitochondrion as previously

described[6,17]. To measure the calcium levels of a single mitochondrion or clusters of mitochondria in *Thy1*-mitoTwitch2b × *Thy1*-OFP, the mean intensity values were measured in the cyan and yellow channels. The FRET signal (YFP channel) was corrected by subtracting the measured crosstalk-fraction of the CFP signal and determined as cFRET. The mitochondrial calcium levels were then expressed as the background-corrected cFRET/CFP ratio normalized to the mean value of healthy mice as previously established for FRET-based calcium sensors[14].

Mitochondrial isolation

Mitochondrial samples for mass spectrometry were prepared from adult male mice described in Fecher at al.[21] In brief, virus-injected control and EAE Mito*Tag* mice were anesthetized with isoflurane and transcardially perfused with 1× PBS/heparin. The lumbar spinal cord was dissected, weighed, and homogenized with a Dounce glass homogenizer using three complete up and down cycles with an A-type pestle in isolation buffer (IB). The sample was then transferred to a cell disruption vessel and processed with nitrogen cavitation at 800 psi and under stirring at 60 rpm for 10 min. After pressure release, a protease inhibitor (cOmplete™, EDTA-free Protease Inhibitor Cocktail, Sigma-Aldrich, 5056489001) was added to the resulting tissue fraction, and subcellular sediments were removed through two times centrifugation at 600g for 10 min. The resulting post-nuclear tissue fraction was filtered through a 30 μm cell strainer. For immunopurification (IP), the post-nuclear tissue fraction was diluted to a maximal concentration of 2 mg tissue/ mL in immunopurification buffer (IPB) and 50μl microbeads coated with mouse IgG$_1$ subtype antibody (Miltenyi Biotec, 130-091-125) against GFP was added to the sample and incubated for 1 hr at 4°C on the shaker. Magnetic-activated cell sorting (MACS) was applied to separate the microbead-coated mitochondria. The LS columns (Miltenyi Biotec, 130-042-401) were placed in a magnetic QuadroMACS™ separator (Miltenyi Biotec, 130-090-976) followed by the equilibration step with 3 mL IPB. The samples were applied to the column in repeating 3ml steps and washed three times with IPB. The

columns were then removed from the magnetic separator, and the microbead-coated mitochondria were gently flushed out with the plunger. Mitochondria were pelleted by centrifugation at 12,000g for 3 min at 4°C and washed twice with IB (without BSA and EDTA), and the pellets were immediately stored at -20°C for further experiments. Subsequently, the protein amount was determined using the BCA assay (Pierce BCA Protein Assay Kit; Thermo Fisher Scientific, 23227) according to the manufacturer's instructions. BSA was used as the standard, and sample buffer was used to correct the measurement alterations caused by detergent or BSA.

Sample preparation for mass spectrometry

Mitochondria were immunocaptured from the spinal cord according to the above-described protocol. Samples were lysed in a modified RIPA buffer (1% Triton X-100, 0.5% sodium deoxycholate, 0.1% SDS, 150 mM NaCl, 5 mM EDTA, 50 mM Tris–HCl, pH 8). Cell debris and undissolved material were removed by centrifugation at 16,000g for 10 min at 4 °C. Protein concentrations were assessed using the BCA assay. A protein amount of 20 μg was further diluted with distilled water at a 1:2 ratio, and 50 mM $MgCl_2$ was added to a final concentration of 10 mM, and DNA/RNA was digested using 12.5U Benzonase. Disulfide bonds were reduced by adding dithiothreitol (DTT) to a final concentration of 10 mM and incubation for 30 min at 37°C. Cysteine alkylation was performed by adding iodoacetamide (IAA) to a final concentration of 40 mM and incubation for 30 min at 24 °C in the dark. The alkylation step was quenched by adding 4 μL of 200 mM DTT.

Protein digestion was performed using the single-pot, solid phase, sample preparation (SP3) protocol[47]. Briefly, 10 μL of a 4 μg/μL bead slurry of Sera-Mag SpeedBeads A and B (GE Healthcare) were added to 20 μg of alkylated protein lysate. Protein binding to the magnetic beads was achieved by adding acetonitrile (ACN) to a final volume of 70% (v/v) and mixing at 1200 rpm at 24 °C for 30 min in a Thermomixer (Eppendorf). Magnetic beads were retained in

a DynaMag-2 magnetic rack (Thermo Fisher Scientific), and the supernatant was discarded. Detergents were removed using four washing steps with 200 µL 80% (v/v) ethanol. Proteins were digested with 0.25 µg LysC (Promega, V1671) at 37°C for 3 hrs followed by a second digestion step with 0.25 µg trypsin (Promega, V5111) for 16 hr at room temperature. Tubes were placed in a magnetic rack, and peptides were transferred to 0.22 µm Costar Spin-X filter tubes (Corning) to remove the remaining magnetic beads. Samples were dried by vacuum centrifugation. Finally, peptides were dissolved in 20 µL 0.1% (v/v) formic acid (FA). Furthermore, the peptide concentration was estimated using the Qubit protein assay (Thermo Fisher, Q33211).

Liquid chromatography-tandem mass spectrometry data acquisition

Samples were analyzed on an Easy nLC-1200 nano UHPLC (Thermo Fisher Scientific) coupled online via a Nanospray Flex electrospray ion source (Thermo Fisher Scientific) equipped with a column oven (Sonation) to a Q-Exactive HF mass spectrometer (Thermo Fisher Scientific). An amount of 1.3 µg of peptides was separated on self-packed C18 columns (300mm × 75 µm, ReproSilPur 120 C18-AQ, 1.9 µm; Dr. Maisch) using a binary gradient of water (A) and acetonitrile (B) supplemented with 0.1% formic acid (gradient: 0 min., 2.4% B; 2 min., 4.8% B; 92 min., 24% B; 112 min., 35.2% B; 121 min., 60% B). Full mass spectrometry spectra were acquired at a resolution of 120,000 (automatic gain control (AGC) target: 3E+6). The 15 most intense peptide ions were chosen for fragmentation by higher-energy collisional dissociation (resolution: 15,000, isolation width: 1.6 m/z, AGC target: 1E+5, normalized collision energy (NCE): 26%). A dynamic exclusion of 120 s was applied for fragment ion spectra acquisition. For cEAE samples, 2 technical replicates were measured per sample.

Liquid chromatography-tandem mass spectrometry data analysis

The LC-MS/MS raw data was analysed with the software Maxquant (Version 1.6.3.3 or 1.6.10.43)[48]. The data was searched against a one gene per protein canonical database of Mus musculus downloaded from UniProt (download date: 2019-03-14, 22294 entries). Carbamidomethylation of cyteines was defined as fixed modification, whereas oxidation of methionines and acetylation of protein N-termini were set as variable modifications. Peptide mass recalibration using the first search option with 20 ppm mass tolerance was enabled. For the main search, peptide and peptide fragment mass tolerances were set to 7 and 20 ppm, respectively. Peptide spectrum match and protein false discovery rates (FDR) were set to 1%, and FDRs were controlled using a forward and reverese concatenated database search approach. The option match between runs was enabled with a matching time window of 1.5 s. Protein label-free quantification was performed on the basis of at least two ratio counts of unique peptides per protein. For cEAE, LFQ intensities of technical replicates were averaged. If a protein was detected in < 50 % of the samples within one experimental group (EAE or control), all values were imputed with values drawn from a standard distribution with a mean value which was 2 standard deviations below the mean LFQ value of the entire dataset and a 0.3-fold standard deviation of the overall standard deviation of the data set (to control for zero inflation). The remaining missing values (of proteins detected in > 50 % of the samples within one experimental group) were imputed by multiple imputations by chained equations (MICE, R package 'mice'). Label-free quantification (LFQ) values were $\log_2$-transformed. For subsequent analysis, LFQ values were treated as interval scale values. To test the similarity of samples within the respective experimental group and to identify outliers, principal component analysis (PCA) was performed. Subsequently, outliers were removed from further analyses. The first two principal components (PC) with their explained variance of each sample were visualized. To test if a protein was differentially expressed between the EAE and control group, we

calculated a Student's t-test and a fold change of the LFQ values between the experimental groups. To control type I error inflation, p values were corrected according to Bonferroni.

MitoCarta 2.0[49] was used to annotate mitochondrial proteins as well as their sub-mitochondrial location. Gene set enrichment analysis was performed with the Python package nezzworker (https://github.com/engelsdaniel/nezzworker) with REACTOME (version 7.4) as a reference gene set library. Proteins which are members of gene sets that showed the most extreme normalized enrichment scores (NES) were further analyzed. To test if mitochondrial, non-mitochondrial proteins and proteins from different mitochondrial respiratory chain complexes were equally abundant in EAE and control samples, one-way analysis of variance was calculated to test for statistically significant differences between the mean LFQ values. To assess transcriptional changes of motor neurons in EAE, we re-analyzed previously published data from Schattling et al.[23], using a custom R script with the DESeq2 library. To assess the correlation of protein lifetime and the alterations of neuronal mitochondrial proteome, we extracted the protein half-lives dataset from Fornasiero *et al*[31]. All analysis was done in a R or Python environment (https://github.com/engelsdaniel/mitoproteomics).

Histology and immunofluorescence staining

Immunofluorescence staining on mouse tissues was performed as described in Fecher et al.[21]. Animals were anesthetized with isoflurane and perfused transcardially with 5000 U/mL heparin (Sigma-Aldrich, H3149) in PBS followed by 4% PFA (Morphisto, 11762.01000) in phosphate buffer (PB) or 4% formalin (Sigma-Aldrich, HT501128). Tissues were kept in 4% PFA or 4% formalin overnight at 4 °C and subsequently cut using a vibratome (Leica). The tissue sections were permeabilized with 0.3% Triton-X100 (Sigma-Aldrich, X100) and blocked using 2% fish gelatin (Sigma-Aldrich, G7765), 2% FBS (Thermo Fisher Scientific, 10500064), and 2% BSA (Sigma, A7030) in PBS for 1 hr. Antigen retrieval procedures were applied to unmask antigens

and epitopes. For this purpose, sections were treated with either 10mM citrate buffer pH 6.0, 10 mM EDTA buffer pH 8.0, or citrate-EDTA (10 mM citric acid, 2 mM EDTA and 0.05% Tween20) pH 6.2 for 10 min at RT followed by 30 min at 90°C and 30 min at 4°C. The sections were further rinsed with 0.05% Tween20 (Sigma-Aldrich, P9416) before the blocking procedure. Primary antibodies were incubated overnight at 4°C at a dilution of 1:400 (Mdh2, Novus Biologicals, NBP1-32259; Idh3a, Novus Biologicals, NBP1-32396; Idh2, Thermo Scientific, 702713) or 1:1000 (RFP, Novus Biologicals, NBP1-97371 and GFP, Abcam, ab13970). The sections were incubated with secondary antibodies at a dilution of 1:1000 for 2 hrs and mounted using Vectashield mounting medium (Vector Laboratories, H-1000) on the following day.

Heat-induced antigen retrieval was performed on PFA fixed tissue following deparaffination for human tissue sections. After 10% FCS/PBS unspecific binding blockade, sections were incubated with primary antibodies in Dako Diluent (Dako, 52022) overnight at 4°C. After washing with Wash Buffer (Dako, 53006) and autofluorescence removal treatment (Merck, 2160), secondary antibodies and DAPI (Life, D3571) at a dilution of 1:200 were incubated at RT for 1 hr and further mounted with Fluoromount Mounting Medium (Sigma, F4680). The use of human samples followed institutional ethical guidelines and was approved by the ethics committee of the University of Geneva (Switzerland). Written informed consent to use autopsy samples for research purposes was obtained for all samples, with the exceptions of autopsies that were performed more than 20 years ago. In all cases, no samples were used from patients who refused involvement in research projects. Patient information is provided in Extended Data Table 1. Data from the human cortex are representative of two experiments performed on 7 different cases.

Confocal imaging and image processing

Sections stained by immunofluorescence labelling were imaged with an upright Olympus FV1000 confocal system equipped with x10/0.4 air, x20/0.85, and x60/1.42 oil immersion objectives. Images were obtained using standard filter sets and processed with Fiji. For representative figure panels, different channels of image series were pseudo-color-coded in Fiji or Adobe Photoshop; schematics were created using BioRender.com. Contrast and brightness were equally adjusted across the entire image. For the panels displayed in **Figure 4**, immunofluorescence images of both control and EAE *Thy1*-mitoRFP tissues were acquired with the same settings and were adjusted with the same processing parameters. In panels that do not primarily illustrate quantitative differences, gamma was adjusted nonlinearly to enhance the visibility of low-intensity objects. Figures were assembled in Adobe Illustrator. Human sections were scanned using a whole slide scanner (Pannoramic P250 II, 3DHistech). Regions of interest were manually selected using SlideViewer software (v2.3, 3DHistech) and exported as individual images for further processing using a custom rule-based script in Definiens Developer XD (v2.7, Definiens AG). In short, specific signals for each marker were detected based on their respective intensity and ratio to tissue background intensity. NF70 (neurofilament) was detected first, and Idh3a detection was restricted to the inside of NF70 positive structures. The total area was quantified for each marker. Additionally, mean intensities were exported for each object individually.

Sequential complex IV histochemistry and immunofluorescence histochemistry

Complex IV histochemistry was combined with immunofluorescence histochemistry with a primary antibody against the orange fluorescent protein (OFP) followed by a directly conjugated secondary antibody to assess complex IV activity in single axons. This sequential technique was performed in the same tissue section and has already been described and validated in the previous studies[5,22,50]. Complex IV media consisted of 100µM cytochrome c, 4mM diaminobenzidine tetrahydrochloride, and 20µg/ml catalase in 0.1M phosphate buffer

pH7.0. Cryosections were incubated at 37 °C for 30 min and washed in PBS. The cryosections were processed through the immunofluorescent histochemistry steps as mentioned above. The sequentially stained sections were mounted in glycerol with Hoechst nuclear stain and stored at -20 °C until required for imaging by the Zeiss Imager Z1 Apotome 2 microscope.

To assess complex IV activity in axons, a mask of the individual axon identified by the fluorescent labelling of OFP was generated in Image J and superimposed onto the brightfield image of complex IV histochemistry. The total area occupied by complex IV active elements within a single axon was determined as a percentage of the axonal area. Twenty axons (at least 25μm in length) per region were randomly selected from each animal for quantitation. Assessors were blinded by coding the axons in ascending numerical order.

Statistics

Statistical analysis was performed using Microsoft excel software and GraphPad Prism (GraphPad Software, Version 7.0, La Jolla, USA). Sample sizes were chosen according to previous *in vivo* imaging studies of spinal axons[6,14]. Normality of distribution was assessed with the Shapiro-Wilk test. When normal distribution was confirmed, a Student's t-test was applied for two groups comparison, and one-way ANOVA followed by Turkey's post hoc comparison was used for more than two groups. The Kruskal–Wallis test followed by Dunn's multiple comparisons test or Mann–Whitney U test was used where normal distribution could not be confirmed. Obtained p-values were stated as significance levels in the figure legends (**** $P<0.0001$, *** $P<0.001$; **$P<0.01$; *$P<0.05$).

SUPPLEMENTARY INFORMATION

Extended Data Table 1: Information on human tissue samples used in this study.

Case	Age	Sex	Histological specimen	State
MS1	41	female	Frontal lobe, subcortical lesion	Chronic active
MS3	45	female	Frontal lobe, subcortical lesion	Chronic active
MS4	51	female	Right frontal-parietal -lobe	Chronic active
MS5	55	male	Left frontal lobe	Chronic active
MS6	58	male	Frontal lobe, subcortical lesion	Chronic
MS8	25	female	Frontal lobe, subcortical lesion	Acute active
MS10	77	female	Not indicated	Acute active

REFERENCES

Friese, M. A., Schattling, B. & Fugger, L. Mechanisms of neurodegeneration and axonal dysfunction in multiple sclerosis. *Nat Rev Neurol* **10**, 225-238, doi:10.1038/nrneurol.2014.37 (2014).

Mahad, D. H., Trapp, B. D. & Lassmann, H. Pathological mechanisms in progressive multiple sclerosis. *Lancet Neurol* **14**, 183-193, doi:10.1016/S1474-4422(14)70256-X (2015).

Absinta, M., Lassmann, H. & Trapp, B. D. Mechanisms underlying progression in multiple sclerosis. *Curr Opin Neurol* **33**, 277-285, doi:10.1097/WCO.0000000000000818 (2020).

Misgeld, T. & Schwarz, T. L. Mitostasis in Neurons: Maintaining Mitochondria in an Extended Cellular Architecture. *Neuron* **96**, 651-666, doi:10.1016/j.neuron.2017.09.055 (2017).

Mahad, D. J. *et al.* Mitochondrial changes within axons in multiple sclerosis. *Brain* **132**, 1161-1174, doi:10.1093/brain/awp046 (2009).

Nikic, I. *et al.* A reversible form of axon damage in experimental autoimmune encephalomyelitis and multiple sclerosis. *Nat Med* **17**, 495-499, doi:10.1038/nm.2324 (2011).

Sadeghian, M. *et al.* Mitochondrial dysfunction is an important cause of neurological deficits in an inflammatory model of multiple sclerosis. *Sci Rep* **6**, 33249, doi:10.1038/srep33249 (2016).

Campbell, G. R. *et al.* Mitochondrial DNA deletions and neurodegeneration in multiple sclerosis. *Ann Neurol* **69**, 481-492, doi:10.1002/ana.22109 (2011).

Witte, M. E., Mahad, D. J., Lassmann, H. & van Horssen, J. Mitochondrial dysfunction contributes to neurodegeneration in multiple sclerosis. *Trends Mol Med* **20**, 179-187, doi:10.1016/j.molmed.2013.11.007 (2014).

Cree, B. A. C. *et al.* Safety and efficacy of MD1003 (high-dose biotin) in patients with progressive multiple sclerosis (SPI2): a randomised, double-blind, placebo-controlled, phase 3 trial. *Lancet Neurol* **19**, 988-997, doi:10.1016/S1474-4422(20)30347-1 (2020).

Rosenkranz, S. C. *et al.* Enhancing mitochondrial activity in neurons protects against neurodegeneration in a mouse model of multiple sclerosis. *Elife* **10**, doi:10.7554/eLife.61798 (2021).

Pathak, D., Berthet, A. & Nakamura, K. Energy failure: does it contribute to neurodegeneration? *Ann Neurol* **74**, 506-516, doi:10.1002/ana.24014 (2013).

Tantama, M., Martinez-Francois, J. R., Mongeon, R. & Yellen, G. Imaging energy status in live cells with a fluorescent biosensor of the intracellular ATP-to-ADP ratio. *Nat Commun* **4**, 2550, doi:10.1038/ncomms3550 (2013).

Witte, M. E. *et al.* Calcium Influx through Plasma-Membrane Nanoruptures Drives Axon Degeneration in a Model of Multiple Sclerosis. *Neuron* **101**, 615-624 e615, doi:10.1016/j.neuron.2018.12.023 (2019).

Sorbara, C. D. *et al.* Pervasive axonal transport deficits in multiple sclerosis models. *Neuron* **84**, 1183-1190, doi:10.1016/j.neuron.2014.11.006 (2014).

Kawachi, I. & Lassmann, H. Neurodegeneration in multiple sclerosis and neuromyelitis optica. *J Neurol Neurosurg Psychiatry* **88**, 137-145, doi:10.1136/jnnp-2016-313300 (2017).

Breckwoldt, M. O. *et al.* Multiparametric optical analysis of mitochondrial redox signals during neuronal physiology and pathology in vivo. *Nat Med* **20**, 555-560, doi:10.1038/nm.3520 (2014).

Brill, M. S., Lichtman, J. W., Thompson, W., Zuo, Y. & Misgeld, T. Spatial constraints dictate glial territories at murine neuromuscular junctions. *J Cell Biol* **195**, 293-305, doi:10.1083/jcb.201108005 (2011).

Giorgi, C., Marchi, S. & Pinton, P. The machineries, regulation and cellular functions of mitochondrial calcium. *Nat Rev Mol Cell Biol* **19**, 713-730, doi:10.1038/s41580-018-0052-8 (2018).

Thestrup, T. *et al.* Optimized ratiometric calcium sensors for functional in vivo imaging of neurons and T lymphocytes. *Nat Methods* **11**, 175-182, doi:10.1038/nmeth.2773 (2014).

Fecher, C. *et al.* Cell-type-specific profiling of brain mitochondria reveals functional and molecular diversity. *Nat Neurosci* **22**, 1731-1742, doi:10.1038/s41593-019-0479-z (2019).

Zambonin, J. *et al.* Identification and investigation of mitochondria lacking cytochrome c oxidase activity in axons. *J Neurosci Methods* **192**, 115-120, doi:10.1016/j.jneumeth.2010.07.022 (2010).

Schattling, B. *et al.* Bassoon proteinopathy drives neurodegeneration in multiple sclerosis. *Nat Neurosci* **22**, 887-896, doi:10.1038/s41593-019-0385-4 (2019).

Chan, K. Y. *et al.* Engineered AAVs for efficient noninvasive gene delivery to the central and peripheral nervous systems. *Nat Neurosci* **20**, 1172-1179, doi:10.1038/nn.4593 (2017).

Trapp, B. D. & Stys, P. K. Virtual hypoxia and chronic necrosis of demyelinated axons in multiple sclerosis. *Lancet Neurol* **8**, 280-291, doi:10.1016/S1474-4422(09)70043-2 (2009).

Nijland, P. G. *et al.* Differential expression of glucose-metabolizing enzymes in multiple sclerosis lesions. *Acta Neuropathol Commun* **3**, 79, doi:10.1186/s40478-015-0261-8 (2015).

Martinez-Reyes, I. & Chandel, N. S. Mitochondrial TCA cycle metabolites control physiology and disease. *Nat Commun* **11**, 102, doi:10.1038/s41467-019-13668-3 (2020).

Anderson, N. M., Mucka, P., Kern, J. G. & Feng, H. The emerging role and targetability of the TCA cycle in cancer metabolism. *Protein Cell* **9**, 216-237, doi:10.1007/s13238-017-0451-1 (2018).

Vasan, K., Werner, M. & Chandel, N. S. Mitochondrial Metabolism as a Target for Cancer Therapy. *Cell Metab* **32**, 341-352, doi:10.1016/j.cmet.2020.06.019 (2020).

Motori, E. *et al.* Neuronal metabolic rewiring promotes resilience to neurodegeneration caused by mitochondrial dysfunction. *Sci Adv* **6**, eaba8271, doi:10.1126/sciadv.aba8271 (2020).

Fornasiero, E. F. *et al.* Precisely measured protein lifetimes in the mouse brain reveal differences across tissues and subcellular fractions. *Nat Commun* **9**, 4230, doi:10.1038/s41467-018-06519-0 (2018).

Bomba-Warczak, E., Edassery, S. L., Hark, T. J. & Savas, J. N. Long-lived mitochondrial cristae proteins in mouse heart and brain. *J Cell Biol* **220**, doi:10.1083/jcb.202005193 (2021).

Trapp, B. D. & Nave, K. A. Multiple sclerosis: an immune or neurodegenerative disorder? *Annu Rev Neurosci* **31**, 247-269, doi:10.1146/annurev.neuro.30.051606.094313 (2008).

Sheng, Z. H. The Interplay of Axonal Energy Homeostasis and Mitochondrial Trafficking and Anchoring. *Trends Cell Biol* **27**, 403-416, doi:10.1016/j.tcb.2017.01.005 (2017).

Trevisiol, A. *et al.* Monitoring ATP dynamics in electrically active white matter tracts. *Elife* **6**, doi:10.7554/eLife.24241 (2017).

Cunnane, S. C. *et al.* Brain energy rescue: an emerging therapeutic concept for neurodegenerative disorders of ageing. *Nat Rev Drug Discov* **19**, 609-633, doi:10.1038/s41573-020-0072-x (2020).

Motte, J. & Gold, R. High-dose biotin in multiple sclerosis: the end of the road. *Lancet Neurol* **19**, 965-966, doi:10.1016/S1474-4422(20)30353-7 (2020).

Gabriel, J. L., Zervos, P. R. & Plaut, G. W. Activity of purified NAD-specific isocitrate dehydrogenase at modulator and substrate concentrations approximating conditions in mitochondria. *Metabolism* **35**, 661-667, doi:10.1016/0026-0495(86)90175-7 (1986).

Sun, P., Bai, T., Ma, T. & Ding, J. Molecular mechanism of the dual regulatory roles of ATP on the alphagamma heterodimer of human NAD-dependent isocitrate dehydrogenase. *Sci Rep* **10**, 6225, doi:10.1038/s41598-020-63425-6 (2020).

Denton, R. M. Regulation of mitochondrial dehydrogenases by calcium ions. *Biochim Biophys Acta* **1787**, 1309-1316, doi:10.1016/j.bbabio.2009.01.005 (2009).

Upadhyay, V. A., Brunner, A. M. & Fathi, A. T. Isocitrate dehydrogenase (IDH) inhibition as treatment of myeloid malignancies: Progress and future directions. *Pharmacol Ther* **177**, 123-128, doi:10.1016/j.pharmthera.2017.03.003 (2017).

Sharma, H. Development of Novel Therapeutics Targeting Isocitrate Dehydrogenase Mutations in Cancer. *Curr Top Med Chem* **18**, 505-524, doi:10.2174/1568026618666180518091144 (2018).

Licht-Mayer, S. *et al.* Enhanced axonal response of mitochondria to demyelination offers neuroprotection: implications for multiple sclerosis. *Acta Neuropathol* **140**, 143-167, doi:10.1007/s00401-020-02179-x (2020).

May, J. L. *et al.* IDH3alpha regulates one-carbon metabolism in glioblastoma. *Sci Adv* **5**, eaat0456, doi:10.1126/sciadv.aat0456 (2019).

Liu, S., Cadoux-Hudson, T. & Schofield, C. J. Isocitrate dehydrogenase variants in cancer - Cellular consequences and therapeutic opportunities. *Curr Opin Chem Biol* **57**, 122-134, doi:10.1016/j.cbpa.2020.06.012 (2020).

Wang, M., Misgeld, T. & Brill, M. S. Neural labeling and manipulation by neonatal intraventricular viral injection in mice. *STAR Protoc* **3**, 101081, doi:10.1016/j.xpro.2021.101081 (2022).

Hughes, C. S. *et al.* Single-pot, solid-phase-enhanced sample preparation for proteomics experiments. *Nat Protoc* **14**, 68-85, doi:10.1038/s41596-018-0082-x (2019).

Cox, J. *et al.* Accurate proteome-wide label-free quantification by delayed normalization and maximal peptide ratio extraction, termed MaxLFQ. *Mol Cell Proteomics* **13**, 2513-2526, doi:10.1074/mcp.M113.031591 (2014).

Calvo, S. E., Clauser, K. R. & Mootha, V. K. MitoCarta2.0: an updated inventory of mammalian mitochondrial proteins. *Nucleic Acids Res* **44**, D1251-1257, doi:10.1093/nar/gkv1003 (2016).

Mahad, D. J. *et al.* Detection of cytochrome c oxidase activity and mitochondrial proteins in single cells. *J Neurosci Methods* **184**, 310-319, doi:10.1016/j.jneumeth.2009.08.020 (2009).

Jassal, B. *et al.* The reactome pathway knowledgebase. *Nucleic Acids Res* **48**, D498-D503, doi:10.1093/nar/gkz1031 (2020).

Gillespie, M. *et al.* The reactome pathway knowledgebase 2022. *Nucleic Acids Res* **50**, D687-D692, doi:10.1093/nar/gkab1028 (2022).

FIGURES

Figure 1: Early and pervasive axonal ATP deficits in EAE lesions.

(**a**) Experimental design for axonal ATP/ADP ratio ([ATP/ADP]$_{axon}$) measurement in experimental autoimmune encephalomyelitis (EAE); FAD: focal axonal degeneration; on: onset. (**b**) Maximum intensity projections of *in vivo* multi-photon image stacks of spinal cord axons of control (left) and acute EAE (right) in *Thy1*-PercevalHR mice. **Top**: Grayscale look-up table (LUT; λ_{ex} 950 nm). **Bottom**: Ratiometric [ATP/ADP]$_{axon}$ LUT (λ_{ex} ratio 950 nm/840 nm). (**c**) Details from b. **Top to bottom**: [ATP/ADP]$_{axon}$ images of control axon in healthy spinal cord, and normal-appearing (stage 0), swollen (1), and fragmented (2) axons in acute EAE. (**d**) Spinal cord axons during chronic EAE shown as in b. (**e**) [ATP/ADP]$_{axon}$ of single axons in healthy and EAE mice normalized to mean of controls (magenta: axons after CCCP, 100 μM; mean ± s.e.m .; n > 800 axons, 6 control and 4 EAE, acute and 6 EAE, chronic compared by Kruskal-Wallis and Dunn's multiple comparison test). (**f**) [ATP/ADP]$_{axon}$ gradient in a dorsal root axon traced through a lesion using *in vivo* multiphoton imaging. **Left**: Traced axon pseudo-colored yellow, running from root at top left to the lesion center at lower right - superimposed on full volume projection, grayscale LUT, λ_{ex} 950 nm. Boxes: Locations of high-resolution stacks. **Right**: Details showing the [ATP/ADP]$_{axon}$ gradient between locations far from (1) vs. close to the lesion (3). LUTs as in b. (**g**) Paired analysis of [ATP/ADP]$_{axon}$ far vs. close to the lesion (n = 22 axons from 3 EAE mice; paired t-test; normalized to control). Scale bars: 25 μm in b, also applied to d; 10 μm in c and f (right); 100 μm in f (left). ***, $p < 0.005$; ****, $p < 0.001$.

Figure 2: Late changes in axonal ROS and calcium homeostasis in EAE lesions.

(**a**) Experimental design to measure mitochondrial ROS ($[ROS]_{mito}$) in EAE. (**b**) Maximum intensity projections of *in vivo* confocal image stacks of spinal cord axons of control (top) and acute EAE (bottom) in *Thy1*-OFP $\times$ *Thy1*-mitoGrx-roGFP mice. **Left**: OFP channel shown with grayscale LUT. **Right**: Ratiometric LUT of $[ROS]_{mito}$ (λ_{ex} ratio 405 nm/488 nm). (**c**) Details from b. **Top to bottom**: mitochondria morphologies and $[ROS]_{mito}$ in normal-appearing (stage 0), swollen (1), and fragmented (2) axons in acute EAE. (**d**) Average $[ROS]_{mito}$ of single axons in control and acute EAE mice normalized to mean of controls (mean $\pm$ s.e.m.; n > 200 axons, 6 control and 5 EAE compared by one-way ANOVA and Tukey's multiple comparison test). (**e**) Single organelle correlation analysis of mitochondrial shape factor (length/width ratio) and $[ROS]_{mito}$ of axons plotted in d. Percentages indicate fraction of mitochondria with $[ROS]_{mito}$ > control mean + 3SD (yellow line) in each axon stage. (**f**) Experimental design to measure mitochondrial Ca^{2+} level ($[Ca^{2+}]_{mito}$) in EAE. (**g**) Maximum intensity projections of *in vivo* multiphoton images of spinal cord axons of control (top) and acute EAE (bottom) in *Thy1*-OFP $\times$ *Thy1*-mitoTwitch2b mice. **Top**: OFP channel shown with grayscale look-up table. **Bottom**: Ratiometric LUT of $[Ca^{2+}]_{mito}$ (YFP/CFP emission ratio). (**h**) Details from g. **Top to bottom**: mitochondria morphologies and $[Ca^{2+}]_{mito}$ represented as in c; grayscale LUT of YFP channel above ratiometric $[Ca^{2+}]_{mito}$ image. (**i**) Average $[Ca^{2+}]_{mito}$ of single axons in control and acute EAE mice normalized to mean of controls. (mean $\pm$ s.e.m.; n > 300 axons, 9 control and 11 EAE mice compared by one-way ANOVA and Tukey's multiple comparison test). (**j**) Single organelle correlation analysis of mitochondrial shape factor (length/width ratio) and $[Ca^{2+}]_{mito}$ of axons plotted in i. Percentages indicate fraction of mitochondria with $[Ca^{2+}]_{mito}$ > control mean + 3SD (yellow line) in each axon stage. Arrow heads indicate axons with different FAD stages. Scale bars: 25 µm in b and g; 10 µm in c and h. ****, p < 0.001

Figure 3: ETC depletion and TCA cycle imbalance in neurons during EAE.

(**a**) Experimental design for combined AAV/*MitoTag*-based proteomic analysis of neuronal mitochondria in EAE. (**b**) Confocal image of outer membrane GFP expression (green) in rAAV9.hSyn:Cre transduced neurons (NeuN, red) in a control and EAE *MitoTag* mouse spinal cord. (**c**) Annotations of the most down-regulated pathways (Reactome[51,52], version 7.4) in *MitoTag* proteomes of neuronal mitochondria in acute EAE. Dot size indicates min-max normalized mean protein expression level ranged from 0 to 1. (**d**) Relative abundance of the TCA and ETC components in neuronal mitochondria. Average shown as color-coded $\log_2$(EAE/Control) for acute and chronic EAE compared to respective controls. (**e**) Rank of TCA (orange) and ETC (green) proteins according to $\log_2$FC expression in acute (top) and chronic (bottom) EAE. (**g**) Correlation between neuronal transcript and protein levels of TCA and ETC proteins in EAE. Transcriptomic data was re-analyzed from Schattling et al.[23] Scale bars: 25 μm in b.

Figure 4: TCA cycle enzymes are dysregulated in neurons in EAE lesions.

(**a**) Experimental design to validate dysregulated TCA enzymes identified by *MitoTag* proteomics in EAE. (**b-g**) Immunofluorescence analysis of Idh3a (**b, c**), Mdh2 (**d, e**) and Idh2 (**f, g**) in spinal cord neurons of control (left panels) or acute EAE (right panels) using *Thy1-mitoRFP* mice (TCA enzymes, green, RFP, red). Graphs (**c, e, g,**) show mean fluorescence intensities in EAE neurons normalized to the mean of control (mean ± s.e.m.; n = 4 mice for Idh3a and Mdh2, 5 for Idh2, compared per animal by two-tailed, unpaired Student's t-test. Scale bar: 25 μm in f, applies also to b, d, and their details. *, p < 0.05; ***, p < 0.005.

Figure 5: TCA cycle enzymes are dysregulated in neurons in MS.

(**a**) Overview of an MS lesion in human cortex, with areas of normal-appearing white matter (NAWM) and the lesion site marked and magnified in insets. Myelin (Luxol fast blue, LFB, blue; periodic acid–Schiff, PAS, purple) and macrophages (CD68, black) are labelled. (**b, c**) Immunofluorescence analysis of Idh3a in cortical axons in NAWM (left panels) or lesion areas (right; Idh3a, green; neurofilament, NF, red). Graphs (**c**) show area occupancy, mean fluorescence intensities and their product - integrated density - in pairs of NAWM and lesion areas from n = 7 cases, as listed in Extended Data Table 1 and compared by paired t-test. Scale bars: 1000 μm in a; 25 μm in b. *, p < 0.05; **, p<0.01.

Figure 6: Idh3 overexpression ameliorates axonal ATP deficits in EAE lesions.

(**a**) Experimental design for [ATP/ADP]$_{axon}$ in acute EAE in *Thy1*- PercevalHR mice that virally overexpressed Idh3a or a control protein (Cre recombinase) together with tdTomato in a subset of axons. (**b**) Maximum intensity projections of *in vivo* multi-photon image stacks of spinal cord axons in Idh3a-overexpressing *Thy1*-PercevalHR mice. **Left**: Grayscale LUT of tdTomato. **Right**: Ratiometric [ATP/ADP]$_{axon}$ LUT (λ_{ex} ratio 950 nm/840 nm). Details show image pairs of tdTomato-negative (left) and –positive (right) normal-appearing (stage 0), swollen (1), and fragmented (2) axons in acute EAE. (**c**) Comparison of [ATP/ADP]$_{axon}$ in tdTomato-positive and -negative axons (plotted as λ_{ex} ratio 950 nm/840 nm, normalized to control axon mean indicated as dash line). **Left**: [ATP/ADP]$_{axon}$ of single tdTomato negative (gray) and positive (orange) axons in Idh3a-overexpressing EAE mice. **Right**: Lesion-specific paired analysis of mean [ATP/ADP]$_{axon}$ in tdTomato-negative (gray) and -positive (orange) axon populations of the three morphological stages. (**d**) Same analysis as c, but in a mouse cohort overexpressing a control protein (Cre recombinase). Mean ± s.e.m. Comparison of n > 300 axons in > 9 lesions from 4 mice in c; n > 300 axons in > 8 lesions from 3 mice in d using a two-tailed, unpaired Student's t-test (c, d, left graphs) and a paired t-test (c, d, right graphs). Scale bar: 25 μm in b. **, $p < 0.01$; ***, $p < 0.005$; ****, $p < 0.001$.