An orphan protein drove the ecological expansion of nitrogen fixation

Bruno Cuevas-Zuviría[1,2,*], Amanda K. Garcia[1,*], Brooke M. Carruthers[1], Betül Kaçar[1,**]

[1]Department of Bacteriology, University of Wisconsin-Madison, Madison, WI, USA

[2]Centro de Biotecnología y Genómica de Plantas, Universidad Politécnica de Madrid, Madrid, Spain

*These authors contributed equally to this work

**To whom correspondence should be addressed: bkacar@wisc.edu

ABSTRACT

Nitrogenases have catalyzed biological nitrogen fixation for billions of years and revolutionized planet Earth by supplying essential nitrogen to the biosphere. How these molecular machines were built and distributed by microbial processes and a shifting geochemical landscape remains an open question. Here, we probe the birth and functional evolution of the G-subunit, an integral, Precambrian component of certain nitrogenases that makes its appearance midway through the history of nitrogen fixation. Our results establish the G-subunit as an unprecedented molecular novelty that was integrated into preexisting interaction pathways for nitrogenase metallocluster incorporation. In the wake of Earth oxygenation, the G-subunit primed nitrogenases and their microbial hosts to exploit newly diversified geochemical environments. The history of the nitrogenase G-subunit showcases the elaborate tango between contingent evolutionary events and environmental conditions that together open an adaptive frontier for molecular innovations with global consequences.

INTRODUCTION

Over billions of years, life has generated an enormous wealth of biomolecular diversity, producing an estimated 10^5 extant protein families[1] and a much larger but inaccessible multitude of extinct proteins. In parallel, biology has become a defining component of the Earth system, which has experienced tremendous revolutions in climate and biogeochemistry that are largely attributable to the molecular innovations of life itself[2]. A unified understanding of the Earth system requires investigating how the proteins and metabolic processes that power planetary phenomena have been built and distributed.

In many cases, nature has evidently been a tinkerer[3], generating novel functions through the progressive duplication, accretion, and recombination of pre-existing protein elements[4-7]. In rarer cases, complete novelty can arise through the *de novo* birth of genes from ancestral, non-genic DNA[4,8]. This scenario represents a more difficult evolutionary hurdle due to the improbability that formerly non-genic DNA would produce a structurally or functionally coherent peptide[9]. Retracing these different evolutionary events in the histories of biogeochemically critical enzymes is itself a considerable challenge given their antiquity, with many of these enzymes, including those required for essential elemental cycling, likely having evolved more than three billion years ago[2,10,11]. Nevertheless, from an evolutionary perspective, it is of considerable interest to uncover the unique conditions under which the benefits of molecular innovations in these systems have outweighed their inherent costs[4] and expanded the planet's biogeochemical repertoire.

Here, we demonstrate an unusual case of molecular novelty in the evolutionary history of biological nitrogen fixation. For >3 billion years[12,13], nitrogenase metalloenzymes have provided the sole biochemical gateway for essential nitrogen into the Earth biosphere, consequently transforming the planet[10,14-16]. This family is today represented by three homologous isozymes—molybdenum- (Nif), vanadium- (Vnf), and iron- (Anf) nitrogenases—that differ in their metal dependencies but otherwise share a core, two-component oxidoreductase architecture[17-19]. The antiquity and biological significance of this architecture is highlighted by its conservation in distantly related enzymes within the nitrogenase superfamily[20], the origin of which likely dates back to the last universal common ancestor[20,21]. Nevertheless, nitrogenase evolution appears to have since uniquely elaborated upon this structural scheme. A striking distinction in all experimentally characterized Vnf and Anf nitrogenases is an additional, functionally enigmatic protein subunit that is not present in Nif, referred to here as the "G-subunit"[22-25]. However, the

origin and role of the G-subunit in the diversification of Vnf and Anf nitrogenases—and thus in the
ability of biological nitrogen fixation to exploit a different suite of geochemical ingredients—
remains unknown. Given the ancient transitions in redox-sensitive metal availabilities triggered
by Earth surface oxygenation[26,27], elucidating these evolutionary features would elucidate the
roles of environmental conditions versus molecular evolutionary events in the ecological
diversification of biological nitrogen fixation[26,28,29].

In the present study, we replay the sequence of events that gave rise to the nitrogenase G-
subunit. We interrogate the genetic and environmental context of its emergence, recruitment, and
functional significance. Our findings reveal the unprecedented nature of G-subunit sequence and
structure within broader protein diversity, plausibly derived from the serendipitous, *de novo* birth
of an orphan gene. By reconstructing its ancestral interactions, we find that G-subunit recruitment
>1.5 billion years ago likely exploited and improved the specificity of existing pathways that shape
nitrogenase metal usage. The G-subunit thus represents an unusual molecular novelty that, in
the wake of Earth oxygenation and its impact on surface geochemistry, fueled the proliferation of
nitrogen fixation strategies to diverse microorganisms and ecological niches.

RESULTS

The nitrogenase G-subunit is an ancient orphan protein

We began our evolutionary study of the nitrogenase G-subunit by constraining its place within the
broader diversity of protein sequence and structure. The G-subunit protein is a small, structural
element of the larger, two-component nitrogenase complex. The first component is a H-subunit
homodimer that delivers electrons to the second, catalytic component, which, in Nif, is a D- and
K-subunit heterotetramer but, in Vnf/Anf (collectively referred to as "alternative nitrogenases"), is
a D-, G-, and K-subunit heterohexamer[22] (**Figure 1A**). Thus, the G-subunit is a primary distinction
in the otherwise structurally comparable Nif, Vnf, and Anf complexes. To date, the sole
crystallographic structure of the G-subunit is from *A. vinelandii* Vnf[22], a 113-amino-acid, four-helix
protein (**Figure 1B**). In Vnf (and presumably Anf), the G-subunit neighbors the D-subunit that
houses the active-site metallocluster where N_2 is bound and reduced[30]. The metal content of the
metallocluster differs between Nif, Vnf, and Anf nitrogenase isozymes, incorporating Fe and Mo
("FeMo-co"), Fe and V ("FeV-co"), or only Fe ("FeFe-co"), respectively. Finally, though sharing a

common ancestry[28,31,32] and core N_2-reduction mechanism[33], Nif, Vnf, and Anf nitrogenases vary

in their kinetic behavior and reactivities to different substrates[19,33-35].

Figure 1. Structure and genetics of the nitrogenase G-subunit. A) Left, crystallographic

structure of the extant A. vinelandii Vnf nitrogenase complex (VnfH, PDB 6Q93[36]; VnfDGK, PDB

5N6Y[22]) containing the G-subunit (dark blue). The Anf nitrogenase is inferred to be comparable

in its multimeric structure, binding FeFe-co instead of FeV-co. Right, crystallographic structure of

the extant Nif nitrogenase complex (NifH, PDB 1M34[37]; NifDK, PDB 3U7Q[38]) lacking the G-

subunit. B) Secondary structure of the Vnf G-subunit. C) Arrangement of the Nif, Vnf, and Anf

structural genes in A. vinelandii. Vnf/AnfG (dark blue) is located between Vnf/AnfD and Vnf/AnfK

genes.

In addition to surveying the diversity of members of the G-subunit protein family, we sought to

identify distant homologs that might provide insights into G-subunit ancestry and function. Though

genetic studies have established the G-subunit as essential for an active, N_2-reducing Vnf/Anf

nitrogenase complex[24,39], its precise functional contribution remains enigmatic. We implemented

a homology search workflow that included both sequence-based (BLASTP, TBLASTN[40], PSI-

BLAST[41], HMMER[42]) and structure-based (Foldseek[43]) methods appropriate for detection of

distantly related proteins. Our search returned >400 canonical G-subunit proteins, but, notably,

we did not find convincing similarity to any other protein. PSI-BLAST and HMMER searches

together returned <10 hits that we could exclude from the canonical G-subunit protein family (i.e.,

were not encoded by genes within *vnf*/*anf* structural operons) (**Table 1, Figure S1**). In both cases,

only short sequence regions of scattered taxonomic affinities aligned to our query with poor

statistical significance. We were thus unable to rule out spurious matches. Additionally, we used

the structure-based method, FoldSeek, leveraging the greater conservation of protein structure

compared to sequence[44] in homology detection. We identified two uncharacterized, AlphaFold-

predicted structures that are not canonical G-subunit proteins but nevertheless share its four-helix

structural arrangement (**Table 1, Figure S2, S3**). However, sequence-level similarity to G-subunit

proteins is negligible and, given the lack of functional annotations for these hits, their relevance

to early G-subunit evolution is unclear. Finally, we note that we did not detect sequence or

structural similarity to NafY and CowN proteins, both of which have previously been proposed to

resemble the G-subunit in their interactions with the nitrogenase complex[22,45,46].

If the G-subunit shares an ancient ancestry with another extant protein family, no strong vestige

of this evolutionary relationship appears presently detectable. The absence of distant homologs

to the G-subunit family is comparable to the case of "orphan genes," which are individual genes

that either have no resemblance to other sequences or have few, taxonomically restricted

relatives[8,47]. Proposed mechanisms for the emergence of orphan genes include duplication

followed by rapid divergence and *de novo* birth from previously non-genic DNA[4,8]. The latter has

more recently been recognized as a significant source of orphan genes[48-50] and thus would

plausibly explain G-subunit origins. Regardless of the specific originating mechanism, our results

indicate that the G-subunit family is a molecular innovation as unique to nitrogenases as it is to

broader protein diversity.

Table 1. Summary of G-subunit homolog search results. *Template modeling score for structural alignments.

Method	Database	Total hits (including canonical G-subunit proteins)	Distant hits	E-value range of distant hits	TM-score* of distant hits	Annotation of distant hits	Taxonomy of distant hits
BLASTP	NCBI non-redundant proteins (nr)	433	0	n/a	n/a	n/a	n/a
TBLASTN	NCBI nucleotide collection (nr/nt)	205	0	n/a	n/a	n/a	n/a
PSI-BLAST	NCBI non-redundant proteins (nr)	446	7	0.026-0.064	n/a	TIGR04141 family sporadically distributed protein, Uncharacterized protein	*Pseudomonas* spp.
HMMER	UniProt Reference Proteomes	94	1	0.061	n/a	Uncharacterized protein	*Coregonus* sp.
FoldSeek	Protein Data Bank	3	0	n/a	n/a	n/a	n/a
FoldSeek	AlphaFold-Uniprot/Swiss-Prot	39	2	5.2-8.7	0.52-0.54	Uncharacterized protein	*Vibrio vulnificus, Deinococcus grandis*

Invention of the G-subunit coincided with expansion of nitrogenase metal dependence and host ecology

We reconstructed phylogenies of nitrogenase G-subunit proteins and examined their genetic distribution across extant genomes and microbial host environments. We identified G-subunit homologs exclusively from *vnf* and *anf* gene clusters in bacterial and archaeal genomes (similarly, all identified *vnf*/*anf* clusters include the G-subunit), confirming that the G-subunit is a significant genetic distinction between Vnf/Anf and Nif nitrogenases[51]. In all analyzed clusters, *vnf/anfG* genes are located between *vnf/anfD and vnf/anfK* loci (**Figure 1C**). By contrast, *nifD* and *nifK* are universally contiguous with small intergenic distances (median ≈ 25 bp) (**Figure S4**). Finally, all complete genomes harboring *vnf* and/or *anf* nitrogenase genes in our dataset also possess *nif* genes. Therefore, the presence of the G-subunit correlates with the capacity of a host diazotroph to express at least two different nitrogenase isozymes with distinct metal dependencies. We built two maximum-likelihood G-subunit phylogenies: the first from individual G-subunit protein sequences and the second from G-subunit sequences concatenated with other nitrogenase subunit sequences. Both trees show separate clustering of Vnf and Anf sequences, though deep branch support in the concatenated tree is significantly improved (**Figure 2, S5, S6**).

The overall topology of our concatenated nitrogenase tree mirrors previous phylogenetic analyses of nitrogenase proteins reconstructed without G-subunit sequences. Nesting of Vnf and Anf clades within Nif sequences indicates that Vnf/Anf are more recently evolved[28,32], which, by extension, establishes the G-subunit as the most recently evolved structural component of the nitrogenase complex. Minimum age estimates of the Vnf/Anf clade range from ~1.5 to 2.5 Ga, based on the timing of horizontal gene transfer events that pervade nitrogenase evolutionary history[13]. These estimates indicate that the Precambrian origin of the G-subunit roughly coincided with the initial accumulation of oxygen in the Earth surface environment[27], the earliest whiffs of which date back to ~2.5 Ga[52].

Our analyses demonstrate that the origin of the G-subunit was associated both with the gain of novel metal dependence and the ecological expansion of nitrogen fixation following the rise of atmospheric oxygen. Vnf and Anf sequences specifically branch from within the "Group III" nitrogenase lineage[31], which is additionally populated by Nif homologs of either experimentally characterized or computationally predicted Mo-dependence[28,53]. Group III Nif are exclusively hosted by anaerobic taxa (primarily methanogenic Euryarchaeota and Firmicutes), many of which

are probable thermophiles or hyperthermophiles (optimal growth temperatures predicted from genome content, see **Materials and Methods**) (**Figure 2, Supplementary Tables**). By contrast, Vnf/Anf are hosted by comparatively more taxonomically diverse and mesophilic archaea and bacteria that also harbor "Group I and II" Nif nitrogenases[31], including the well-studied diazotrophic models like *A. vinelandii*, *Clostridium pasteurianum,* and *Rhodopseudomonas palustris*. Vnf/Anf are not hosted by organisms that harbor Group III Nif, despite Vnf/Anf emerging from within the Group III lineage. Within Group III, only Vnf and Anf clades contain sequences from aerobic taxa, including cyanobacteria which appear to only host Nif and Vnf nitrogenases. Together, these data indicate that G-subunit containing Vnf and Anf nitrogenases were able to proliferate into ecological niches inaccessible to earlier diverged Group III Nif.

Figure 2. Evolutionary and environmental history of the nitrogenases and the G-subunit. *Maximum-likelihood phylogeny built from concatenated nitrogenase subunit proteins. Oxygen tolerance, predicted optimal growth temperatures (OGT), and taxonomy of represented host microbes, as well as the metal content of the active-site metallocluster, are mapped to the*

 phylogeny (light grey bars for O$_2$ tolerance and predicted OGT indicate missing data). Nif clades
are labeled according to the nomenclature used by Raymond et al.[31] (e.g., Nif-I = Group I Nif,
etc.). The G-subunit origin event is indicated by a black hash mark. Branches that track the
emergence of the G-subunit and its diversification into Vnf and Anf groups are shown in black,
whereas branches lacking the G-subunit are shown in grey. Ancestral nodes (Nif^{Anc}, VnfAnf^{Anc},
Vnf^{Anc}, Anf^{Anc}) targeted for sequence reconstruction and analysis are highlighted by bold outlined,
white circles.

An ancient assembly pathway preconditioned early nitrogenases for G-subunit interaction

To replay the first evolutionary steps in the birth of the G-subunit, we reconstructed and analyzed
ancestral nitrogenase protein sequences. We examined the degree to which ancestral protein
surface features were remodeled to accommodate a new subunit, with the expectation that G-
subunit interactions would have driven the evolution of complementary residues within the
interface. Because the G-subunit is exclusive to Vnf and Anf nitrogenases, the most parsimonious
evolutionary scenario is that it emerged along the branch leading to the Vnf/Anf last common
ancestor, VnfAnf^{Anc} (**Figure 2**). We therefore compared sequences and structures of nitrogenase
ancestors reconstructed both immediately before (Nif^{Anc}) and after (VnfAnf^{Anc}) gain of the G-
subunit.

We mapped the interface between ancestral G- and D-subunit proteins ("G-D interface") by
building its residue interaction network from the predicted VnfAnf^{Anc} structure (**Figure 3A-3D**).
Nearly all well-conserved, G-subunit residues lie within the G-D interface and contribute
significantly to its stability (**Figure S7**). The few exceptions are conserved residues distal from the
interface with side chains oriented towards the interior of the helical bundle, which likely stabilize
its globular arrangement.

We tracked the evolution of D-subunit residues through the emergence of the ancestral G-D
interface. We identified 42 D-subunit sites interacting with the G-subunit following its initial
recruitment in ancestral VnfAnf^{Anc} (**Supplementary Tables**). Of the 42 sites, 26 are well-
conserved among extant Vnf/Anf nitrogenases and form a core interaction patch that was likely
integral to the early G-D interface (**Figure 3E, 3F**). We inspected these 26 sites and found that
nine sites evolved (i.e., were substituted) alongside G-subunit emergence, imposing subtle
changes to the electrostatic surface potential of the VnfAnf^{Anc} interaction patch (**Figure S8, S9**).

Figure 3. Birth of the nitrogenase G-subunit interface. A) Predicted Nif^{Anc} D-subunit structure, prior to G-subunit emergence. Inset shows cross-section at the putative metallocluster insertion site, the entrance of which forms a "cleft" in the D-subunit surface. Superimposed G-subunit

*structure (transparent) highlights relative position of the cleft. **B)** Predicted VnfAnf^{Anc} D- and G-*
subunit structures, following G-subunit emergence, shown in the same view as A). Side chain of
*the G-subunit Leu18 residue is anchored within the D-subunit cleft. **A-B)** Heterometal content for*
*the ancestor metalloclusters is inferred. **C)** Nif^{Anc} G-D proto-interaction network. Purple residues*
are located within an interface patch conserved in extant Vnf/Anf. Residues that are later
substituted in VnfAnf^{Anc} are shown bold and labeled to permit comparison with D). Residues within
*the cleft are circled. **D)** VnfAnf^{Anc} G-D interaction network, shown in the same view as C).*
*Residues newly substituted in VnfAnf^{Anc} are shown in dark purple and labeled. **E)** Surface view of*
*the Nif^{Anc} D-subunit proto-interface patch (purple). **F)** Surface view of the VnfAnf^{Anc} D-subunit*
interface patch, shown in the same view as E). Residues that are maintained between NifAnc and
*VnfAnfAnc are shown in purple, and those that are substituted are shown in dark purple. **G)***
Evolution of residues within the interaction patch through the emergence of the G-subunit. For
substituted sites, the posterior probability of the most likely VnfAnf^{Anc} residue is <.3 in Nif^{Anc} and
>.7 in VnfAnf^{Anc} and, for maintained sites, >.7 in Nif^{Anc} and >.7 in VnfAnf^{Anc}.

Intriguingly, we find that multiple ancestral D-subunit surface features appear to have been
preconditioned for G-subunit binding, collectively forming a "proto-interface" upon which
interactions with the newly evolved G-subunit is built. Out of 26 residues within the core, 16
(~62%) show that the D-subunit interaction patch had already become VnfAnf^{Anc}-like prior to G-
subunit emergence (**Figure 3F,G**). This residue-level conservation translates to maintained
hydrophobicity (median apolar fraction ≈ .2) of the (proto-)interaction patch both before and after
G-subunit recruitment (**Figure S10**). Further, shape complementarity as an integral feature of the
G-D interface appears to have preceded the G-subunit. In *A. vinelandii* Vnf[22], a cleft within the D-
subunit interaction patch is occupied by a strictly conserved G-subunit residue, Leu18. This
residue is among the most well-connected within the G-D interaction network (**Figure 3A-D**), is
critical for interface stability (**Figure S7**), and is among the most allosterically active, capable of
generating long-distance effects that reach across opposite ends of the nitrogenase complex
(**Figure S11**). These effects reflect the importance of the G-D interface in controlling the global
motions of the nitrogenase complex and evoke similar, long-distance structural perturbations that
are induced through the nitrogenase catalytic cycle[54,55]. The cleft itself provides an important
anchor point for G-subunit binding and is present in both ancestral and representative extant
nitrogenase structures (**Figure 3A,B, Figure 4**).

Figure 4. Conservation of the D-subunit cofactor insertion cleft across extant and ancestral nitrogenases. A) *Crystallographic structure of the A. vinelandii apo-NifD protein (PDB 1L5H). Cross-section view (right) reveals the proposed metallocluster insertion site[56], the entrance of which structurally aligns with the G-D interface of VnfD. Aligned, cross-section views are displayed for extant* **B)** *A. vinelandii NifD (PDB 3U7Q),* **C)** *Clostridium pasteurianum NifD (PDB 4WES), and* **D)** *A. vinelandii VnfD (PDB 5N6Y) crystallographic structures as well as predicted, ancestral* **E)** *Nif[Anc] D-subunit and* **F)** *VnfAnf[Anc] D-subunit structures.* **(A-C, E)** *A. vinelandii VnfG (PDB 5N6Y; transparent) is superimposed to enable comparison across nitrogenases with and without native G-subunit proteins.* **(A-F)** *Small black arrow indicates position of the proposed cofactor insertion cleft.*

The functional significance of the D-subunit cleft provides an explanation for why the interface
patch resisted significant remodeling with G-subunit addition, as well as implicates an adaptive
role for the G-subunit. In extant Nif, the cleft aligns with the entrance to a proposed nitrogenase
metallocluster insertion site, which, in the apo-NifD protein, adopts an open conformation to permit
cluster loading[56]. Given the consistent presence of the cleft across Nif, Vnf, and Anf nitrogenases,
a similar mechanism can be predicted for Vnf and Anf and is likely an ancient, universal feature
of nitrogenase maturation that predates the G-subunit and was maintained due to its essentiality.
In fact, we find that 12 interface residues surrounding the insertion site are well-conserved in
extant Group III Nif clades that diverge prior to the origin of the G-subunit, and five of these are
either universally or nearly universally conserved among all nitrogenases (**Figure S12**). Structural
alignments indicate that the G-subunit caps the opening of the insertion site, with the side chain
of the conserved Leu18 residue oriented into the cleft and toward the loaded active site. An
unusual feature of the G-subunit, a carboxyl-terminal tail that terminates in a strictly conserved
tyrosine residue (Tyr113), bridges the presumed opening of the apo-D-subunit protein and
neighbors universally conserved D-subunit residues. Together, these features implicate a role for
the G-subunit in nitrogenase assembly, which is supported by experimental evidence that VnfG
and VFe-co associate *in vitro*, as well as of the requirement of both VnfG and VFe-co to stabilize
an active complex[24]. Conservation of G-subunit residues surrounding the cleft (and thus the
metallocluster insertion site) suggests that this putative role was early-evolved and maintained
through its evolution.

The G-subunit confers specialization between Vnf and Anf nitrogenases

We assessed to what degree sequence variation within the G-subunit is capable of driving the
functional distinctions between Vnf and Anf isozymes, which are each dependent on distinct
metalloclusters and exhibit varied N_2-reduction efficiencies and reactivities to different chemical
substrates[19,33-35]. We identified protein sites that are "specialized" between Vnf and Anf (i.e.,
diverge between Vnf and Anf but are otherwise conserved in each of the two groups), and
therefore most likely drive functional variation. We find the degree of specialization (relative to
protein size) within the G-subunit to be on par with that of the D-subunit but observe comparatively
little specialization within the K-subunit (**Figure 5A, 5B, Supplementary Tables**). Therefore, the
G-subunit contributes an outsized fraction of the total sequence specialization between Vnf and
Anf, despite its remoteness from any nitrogenase metalloclusters. Specialized sites within the G-

subunit are distal from the G-D interface. These include one residue close to the presumed H-subunit binding site, which might contribute to a previously proposed role in mediating H-subunit binding specificity[25].

We next examined how these signatures of sequence-level specialization manifested early in G-subunit evolution, tracking their amino acid content through the divergence of Vnf and Anf clades. We find that the specialized sequence motifs of Vnf[Anc] (Vnf ancestor) and Anf[Anc] (Anf ancestor) most closely resemble those of their respective descendants (**Figure 5C**). The most likely Vnf[Anc] motif across divergent sites is "FIKIDV" (extant Vnf consensus "FIKVDH") whereas the most likely Anf[Anc] motif is "VNVDYD" (extant Anf consensus "VNVDYD"). However, we find that the most likely motif of the common VnfAnf[Anc] ancestor, "VIKIDV," is more like that of Vnf than Anf, even accounting for the statistical uncertainty associated with ancestral reconstruction. This result suggests that the earliest G-subunit proteins were Vnf-like (in agreement with previous inferences of ancestral nitrogenases that indicate Vnf evolved prior to Anf[28]), suggesting that the G-subunit contributed to the specificity required for vanadium dependency early in the birth of a new nitrogenase.

Figure 5. Coevolution of G-subunit specialization and Vnf/Anf metal dependence. A) Specialized sites mapped to the nitrogenase VnfDGK structure (PDB 5N6Y[22]). Scale in A) applies to B). B) Detailed view of specialized sites across the G-subunit. C) Evolution of specialized G-subunit sequence motifs through the divergence of Vnf and Anf. Sequence logos scale either with amino acid frequencies (for extant Vnf or Anf nitrogenases) or posterior probabilities (for ancestral VnfAnf^{Anc}) at each site. VnfAnf^{Anc} dependence on FeV-co is inferred based on comparisons of specialized sequence motifs. D) The bulk marine concentrations of redox-sensitive metals involved in nitrogen fixation (Mo, V, Fe) have changed with progressive oxygenation of the Earth surface environment, with Fe more abundant in the Archean and Mo/V more abundant today. Though quantitative estimates of historical V concentrations are not presently available, oxygenation likely impacted the speciation and solubility of V and Mo similarly over geologic time[57,58]. Metal concentrations from Zerkle et al.[59] and references therein and p_{O2} from Lyons et al.[27]. Age constraints of nitrogenase origins and VnfAnf^{Anc} (i.e., G-subunit emergence) from Stueken et al.[12] and Parsons et al.[13], respectively. GOE: Great Oxidation Event, PAL: present atmospheric level.

DISCUSSION

In the present study, we interrogated the origin, early evolution, and functional significance of a major novelty in the history of nitrogen fixation: the emergence of the G-subunit in Vnf and Anf nitrogenases >1.5 billion years ago. With the birth of the nitrogenase G-subunit, not only did a rare evolutionary event occur with associated fitness costs overcome[4], but a novel structural domain became fully integrated into the nitrogenase complex and was retained in one of the most central metabolic processes to Earth's biosphere.

We demonstrate that there currently exists no candidate, evolutionary predecessor of the G-subunit family, whether another protein, protein fragment, or segment of non-genic DNA. This finding sets the origin of the G-subunit apart from other major evolutionary events in the history of nitrogen fixation that are explainable by gene duplication, including the divergence between other catalytic subunits, between nitrogenases and their maturation protein scaffolds, and between nitrogenase isozymes[32,60]. The addition of such a small and evolutionarily enigmatic protein domain appears to be an anomaly across biogeochemical enzymes.

We propose that the G-subunit was formed *de novo* from previously non-genic DNA, more
recently acknowledged as a viable process for generating genetic novelty[48-50]. Such an event may
have accompanied the genetic duplication and divergence of Nif that produced alternative
nitrogenases[32]. Regardless of its originating mechanism, retainment of the G-subunit gene within
the nitrogenase operon appears to have been improbable. Insertions between nitrogenase D- and
K-subunit genes must either be rare or frequently deleterious, as evidenced by their small
intergenic distance and absence of any other disrupting open reading frames across extant Nif
clusters in our dataset. Thus, the evolution of the G-subunit constitutes an unprecedented gain in
structural and genetic complexity in the evolutionary history of nitrogen fixation.

By analyzing predicted, nitrogenase ancestors, we reveal an evolutionary steppingstone in the
initial recruitment of the G-subunit. Our reconstruction of ancestral, D-subunit surface features
indicates that the G-subunit binding site was built from an essential, preexisting metallocluster
insertion site, which additionally appears preconditioned for G-subunit interactions. How did this
essential surface feature accommodate burial by a new interaction partner? Metallocluster
delivery in extant nitrogenases is thought to be mediated by transient protein-protein
interactions[56,61,62] that would presumably have shaped nitrogenase surface properties, perhaps
explaining the early evolved hydrophobicity of the ancestral, proto-interface. Our evolutionary
analyses, coupled with experimental support of a role in nitrogenase assembly[24], suggest that the
G-subunit was integrated within the assembly pathway, replacing and/or mediating earlier evolved
protein or metallocluster interactions. Thus, the birth of the permanent G-D interface represents
a gradual subsummation of formerly transient interactions that provided an evolutionary "cradle"
for a permanent, novel structural domain.

In its capacity to mediate newly evolved incorporation of V- and Fe-based clusters, the G-subunit
would have enabled the evolutionary burst of both metal dependence and ecological diversity
within Group III nitrogenase lineage from which alternative nitrogenases originated. We find that
G-subunit residues, despite being relatively distal from the active-site and not retaining bound,
reactive metalloclusters like other nitrogenase subunits, have an outsized contribution to the
sequence-level specialization between alternative nitrogenases. Possible functional outcomes
include improved specificity toward metalloclusters or associated protein partners during
nitrogenase assembly (preventing costly mismetallation[63])[24] and mediating crosstalk between
electron transfer and catalytic nitrogenase components[25]. Each of these functions would have
equipped diazotrophs to coordinate multiple nitrogenases systems with unique metal

dependencies within the same cellular environment. The G-subunit therefore primed these

organisms for fluctuations in redox-sensitive metal availabilities following the rise of oxygenic

phototrophs[27,64], one of the most significant ecological revolutions in Earth history.

An ongoing debate in evolutionary biology centers on the degree to which environmental

conditions drive biological innovations. In the evolution of biological nitrogen fixation, an intuitive

narrative is that diversification of nitrogenase metal dependence, intimately tied to the productivity

and ecological distribution of nitrogen fixation, was triggered largely by transitions in bulk,

environmental metal availabilities[26], the historical records of which are primarily built from

geochemical analyses of ancient marine sediments (**Figure 5D**). Our study instead highlights the

importance of chance events shaping evolutionary innovations with global consequences. The

birth of the nitrogenase G-subunit, by providing a crucial component in novel metal incorporation,

expanded the adaptive landscape for biological nitrogen fixation, forever altering the evolutionary

trajectory of the biosphere. This ancient, orphan protein was birthed by seemingly improbable

events but was nevertheless fostered by molecular features that were predisposed for its arrival.

Whether this evolutionary transition in the history of biological nitrogen fixation could have been

crossed in the absence of the G-subunit awaits further studies. Similar, future synergies between

historical studies of molecular novelties and the environmental circumstances that shaped them

can refine our understanding of how the Earth-life planetary system has evolved over billions of

years.

METHODS

Nitrogenase protein sequence curation

A preliminary 412-sequence dataset of nitrogenase Vnf/AnfG proteins was compiled by BLASTP[40]

search (Expect value threshold = 0.1) against the National Center for Biotechnology Information

(NCBI) non-redundant protein sequence database (nr)[65] (accessed January 2022), using the *A.*

vinelandii VnfG query sequence (WP_012698949.1). BLASTP hits were curated by filtering

against partial sequences and homologs whose encoding genes are distantly located from *vnf* or

anf operons (and thus unlikely to be canonical *vnf/anfG* genes). Certain VnfG sequences were

manually extracted from fused VnfDG proteins, using MAFFT v7.490[66] alignments to other D- or

G-subunit sequences as a guide. The preliminary Vnf/AnfG sequence dataset was pruned by CD-

HIT[67] (sequence identity threshold = 90%), yielding a final 188-sequence G-subunit dataset.

G-subunit homolog search

In addition to the BLASTP search described above, three sequence-based search tools–
TBLASTN[40], PSI-BLAST[41], and HMMER[42]–were used to identify distantly related homologs of the
nitrogenase G-subunit (conducted January 2022). TBLASTN and PSI-BLAST searches against
the NCBI nucleotide collection (nr/nt) and nr protein database, respectively, were executed with
the *A. vinelandii* VnfG query (WP_012698949.1), an Expect value threshold = 0.1, and word size
= 2 (all other parameters default). HMMER search (hmmsearch) against the UniProt Reference
Proteome database[68] was performed using an HMM profile generated from our MAFFT-aligned,
188-sequence G-subunit dataset (see above) with an Expect value threshold = 0.1 for both
sequences and hits (all other parameters default).

Structure-based searches of distant G-subunit homologs were performed by the FoldSeek
server[43] using the *A. vinelandii* VnfG crystallographic structure (PDB 5N6Y[22]) query against the
Protein Data Bank and AlphaFold-UniProt/Swiss-Prot databases. We employed both 3Di-AA and
TM-Align modes with an 3Di-AA Expect value threshold = 0.01 or a TM-score = 0.5 (template
modeling score).

Phylogenetic and ancestral sequence reconstruction

The curated, 188-sequence Vnf/AnfG dataset was aligned by MAFFT. No outgroup to G-subunit
sequences was included because the distant homology search described above did not yield
convincing hits (see **Results**). IQ-TREE v1.6.12[69] was used to reconstruct a maximum-likelihood
G-subunit protein phylogeny, using the best-fit evolutionary model, WAG+R6 (selected by the
Bayesian Information Criterion, BIC), tested by ModelFinder[70], with branch support calculated by
ultra-fast bootstrap approximation (UFBoot)[71] and the Shimodaira-Hasegawa-like approximate
likelihood-ratio test (SH-like aLRT)[72].

A phylogeny of concatenated Vnf/AnfHDGK proteins was generated by retrieving syntenous
subunit sequences to the curated VnfG/AnfG dataset. Certain VnfH sequences that recently
diverged from NifH clades were omitted due to having different evolutionary histories than those
of other Vnf/Anf proteins. NifHDK sequences from a prior phylogenetic analysis[73] were included
as an outgroup (n = 67, closely branching to Anf/Vnf) and were used to calculate intergenic *nifD-*

to-*nifK* distances (n = 306). Each subunit sequence dataset was aligned by MAFFT prior to

concatenation. Maximum-likelihood phylogenetic reconstruction was performed by IQ-TREE as

described above (best-fit evolutionary model, BIC: LG+F+R9).

Marginal maximum-likelihood ancestral sequence reconstruction of concatenated Vnf/AnfHDGK

sequences was performed by PAML v4.9j[74] (LG+F+G4 model). Mean sitewise posterior

probabilities D- and G-subunit ancestors analyzed in this study range from .75 to .96 (**Table S1**).

Sequence gaps were reconstructed as described by Aadland et al.[75]. Briefly, the input protein

sequence alignment was recoded as a binary amino-acid presence/absence matrix. Ancestral

presence/absence states were reconstructed by PAML using a binary character model and state

frequencies inferred by maximum likelihood.

Logo representations of extant multiple sequence alignments were generated with the Python

library LogoMaker[76] using the probability transformation of the count matrix. Representations of

ancestral sequences were generated from the posterior probability matrices produced by PAML.

Sequence   and   phylogenetic   tree   datasets   are   available   at

https://github.com/kacarlab/nitrogenase-g-subunit.

Optimal growth temperature prediction

For taxa represented in the nitrogenase G-subunit sequence dataset, optimal growth

temperatures (OGT) were predicted from genomic content using the method described by Sauer

and Wang[77]. Due to genome incompleteness and the absence of 16S rRNA sequences in

fragmented genomes for certain representative taxa, the selected "Superkingdom Bacteria" and

"Superkingdom Archaea" regression models for OGT prediction excluded genome size and 16S

rRNA.

Protein conservation and divergence analysis

493   Sequence conservation for D- and G-subunit proteins was determined by the ConSurf server[78].

494   Sites were considered well-conserved if their ConSurf grade were >7. Sequence specialization

495   was determined by the TwinCons package[79]. TwinCons was run separately on individual

496   alignments of extant H, D, G, and K subunit protein sequences with user-defined Vnf and Anf

groups, using default parameters and the LG substitution matrix. TwinCons specialization scores were normalized to a 0-to-1 scale per analyzed subunit.

Structural Modeling

Ancestral and extant NifHDGK protein structures were predicted using ColabFold[80], which combines the AlphaFold2 method for structure prediction from multiple-sequence alignments[18] with the MMSeq2 method to obtain evolutionary information[81]. Specifically, the colabfold_batch v1.4.0 implementation (https://github.com/YoshitakaMo/localcolabfold) was used with the AlphaFold-v2.0-multimer prediction model and standard options (3 recycles and AMBER all-atom optimization in GPU).Protein structures were visualized by ChimeraX[82].

Structure Analysis

Residue interaction networks[83] were generated using ProLif[84], which finds the different possible interactions (hydrophobic, electrostatic, hydrogen bonds, etc) between two groups of atoms (here defined as G- and D-subunit groups) within a structure snapshot (e.g. single structure, trajectory frame). Interactions were compiled into networks by NetworkX[85], which were later visualized in Gephi[86]. Scripts used to generate the networks are available at https://github.com/kacarlab/nitrogenase-g-subunit.

Surface electrostatic maps were computed using the Poisson Boltzmann Solvent-Accessible model implemented in APBS[87] through the PyMOL plugin (Schrödinger, Inc), using standard PDB2PQR and APBS protocols at pH 7.

Sitewise contribution to D-G interface stability was computed using *in-silico* alanine scanning, which substitutes residues with a minimal methyl moiety (alanine)[88]. The changes on the binding free energy values upon substitution were obtained using the machine-learning enabled mCSM2 method[89] with default parameters.

Allosteric effects across the nitrogenase complex was modeled using Perturbation Response Scanning[90], implemented in ProDy[90] with default parameters and provided with PDB 5N6Y[22] alpha carbon coordinates. Scripts used to model allosteric effects are available at https://github.com/kacarlab/nitrogenase-g-subunit.

ACKNOWLEDGEMENTS

This research was supported by the National Aeronautics and Space Administration (NASA)
Interdisciplinary Consortium for Astrobiology Research: Metal Utilization and Selection Across
Eons, MUSE (19-ICAR19_2–0007), the NASA Arizona Space Grant (B.M.C), the John Templeton
Foundation (B.K.; 61926), the National Science Foundation (B.K.; 2228495), the NASA Early
Career Faculty Award (B.K.), the Hypothesis Fund Award (B.K.), and the Universidad Politécnica
de Madrid (UPM) Margarita Salas Fellowship, founded by the Unión Europea - Next Generation
EU (B.C.-Z.; UP2021-035). We thank Morgan Sobol for assistance with optimal growth
temperature prediction, as well as Alex Rivier, Holly Rucker, Lance Seefeldt, Derek Harris,
Joanna Masel and Steven Russell for valuable discussions and feedback.

REFERENCES

1   Choi, I.-G. & Kim, S.-H. Evolution of protein structural classes and protein sequence
families. *Proc Natl Acad Sci USA* **103**, 14056-14061, doi:10.1073/pnas.0606239103
(2006).
2   Knoll, A. H. The geological consequences of evolution. *Geobiology* **1**, 3-14,
doi:10.1046/j.1472-4669.2003.00002.x (2003).
3   Jacob, F. Evolution and Tinkering. *Science* **196**, 1161-1166, doi:10.1126/science.860134
(1977).
4   Andersson, D. I., Jerlstrom-Hultqvist, J. & Nasvall, J. Evolution of new functions de novo
and from preexisting genes. *Cold Spring Harb Perspect Biol* **7**,
doi:10.1101/cshperspect.a017996 (2015).
5   Eck, R. V. & Dayhoff, M. O. Evolution of the Structure of Ferredoxin Based on Living Relics
of Primitive Amino Acid Sequences. *Science* **152**, 363-366,
doi:10.1126/science.152.3720.363 (1966).
6   Copley, S. D. Evolution of new enzymes by gene duplication and divergence. *FEBS J* **287**,
1262-1283, doi:10.1111/febs.15299 (2020).
7   Alva, V., Soding, J. & Lupas, A. N. A vocabulary of ancient peptides at the origin of folded
proteins. *Elife* **4**, e09410, doi:10.7554/eLife.09410 (2015).
8   Van Oss, S. B. & Carvunis, A. R. De novo gene birth. *PLoS Genet* **15**, e1008160,
doi:10.1371/journal.pgen.1008160 (2019).

9   Beasley, J. R. & Hecht, M. H. Protein design: the choice of de novo sequences. *J Biol*
Chem **272**, 2031-2034, doi:10.1074/jbc.272.4.2031 (1997).

10   Falkowski, P. G., Fenchel, T. & Delong, E. F. The microbial engines that drive Earth's
biogeochemical cycles. *Science* **320**, 1034-1039, doi:10.1126/science.1153213 (2008).

11   Moore, E. K., Jelen, B. I., Giovannelli, D., Raanan, H. & Falkowski, P. G. Metal availability
and the expanding network of microbial metabolisms in the Archaean eon. *Nat Geosci* **10**,
629-636, doi:10.1038/ngeo3006 (2017).

12   Stueken, E. E., Buick, R., Guy, B. M. & Koehler, M. C. Isotopic evidence for biological
nitrogen fixation by molybdenum-nitrogenase from 3.2 Gyr. *Nature* **520**, 666-669,
doi:10.1038/nature14180 (2015).

13   Parsons, C., Stüeken, E. E., Rosen, C. J., Mateos, K. & Anderson, R. E. Radiation of
nitrogen-metabolizing enzymes across the tree of life tracks environmental transitions in
Earth history. *Geobiology* **19**, 18-34, doi:10.1111/gbi.12419 (2020).

14   Falkowski, P. G. Evolution of the nitrogen cycle and its influence on the biological
sequestration of CO_2 in the ocean. *Nature* **387**, 272-275, doi:10.1038/387272a0 (1997).

15   Allen, J. F., Thake, B. & Martin, W. F. Nitrogenase Inhibition Limited Oxygenation of
Earth's Proterozoic Atmosphere. *Trends Plant Sci* **24**, 1022-1031,
doi:10.1016/j.tplants.2019.07.007 (2019).

16   Sanchez-Baracaldo, P., Ridgwell, A. & Raven, J. A. A neoproterozoic transition in the
marine nitrogen cycle. *Curr Biol* **24**, 652-657, doi:10.1016/j.cub.2014.01.041 (2014).

17   Einsle, O. & Rees, D. C. Structural Enzymology of Nitrogenase Enzymes. *Chem Rev* **120**,
4969-5004, doi:10.1021/acs.chemrev.0c00067 (2020).

18   Mus, F., Alleman, A. B., Pence, N., Seefeldt, L. C. & Peters, J. W. Exploring the
alternatives of biological nitrogen fixation. *Metallomics* **10**, 523-538,
doi:10.1039/c8mt00038g (2018).

19   Eady, R. R. Structure-function relationships of alternative nitrogenases. *Chemical*
Reviews **96**, 3013-3030, doi:10.1021/cr950057h (1996).

20   Ghebreamlak, S. M. & Mansoorabadi, S. O. Divergent Members of the Nitrogenase
Superfamily: Tetrapyrrole Biosynthesis and Beyond. *ChemBioChem* **21**, 1723-1728,
doi:10.1002/cbic.201900782 (2020).

21   Weiss, M. C. *et al.* The physiology and habitat of the last universal common ancestor. *Nat*
Microbiol **1** (2016). <https://www.ncbi.nlm.nih.gov/pubmed/27562259>.

22   Sippel, D. & Einsle, O. The structure of vanadium nitrogenase reveals an unusual bridging
ligand. *Nat Chem Biol* **13**, 956-960, doi:10.1038/nchembio.2428 (2017).

23   Eady, R. R., Robson, R. L., Richardson, T. H., Miller, R. W. & Hawkins, M. The vanadium
600   nitrogenase of Azotobacter chroococcum. Purification and properties of the VFe protein.
601   *Biochemical Journal* **244**, 197-207, doi:10.1042/bj2440197 (1987).

602   24   Chatterjee, R., Ludden, P. W. & Shah, V. K. Characterization of VNFG, the delta subunit
603   of the vnf-encoded apodinitrogenase from Azotobacter vinelandii. Implications for its role
604   in the formation of functional dinitrogenase 2. *J Biol Chem* **272**, 3758-3765,
605   doi:10.1074/jbc.272.6.3758 (1997).

606   25   Pence, N., Lewis, N., Alleman, A. B., Seefeldt, L. C. & Peters, J. W. Revealing a role for
607   the G subunit in mediating interactions between the nitrogenase component proteins. *J*
608   *Inorg Biochem* **214**, 111273, doi:10.1016/j.jinorgbio.2020.111273 (2021).

609   26   Anbar, A. D. & Knoll, A. H. Proterozoic ocean chemistry and evolution: a bioinorganic
610   bridge? *Science* **297**, 1137-1142, doi:10.1126/science.1069651 (2002).

611   27   Lyons, T. W., Reinhard, C. T. & Planavsky, N. J. The rise of oxygen in Earth's early ocean
612   and atmosphere. *Nature* **506**, 307-315, doi:10.1038/nature13068 (2014).

613   28   Garcia, A. K., McShea, H., Kolaczkowski, B. & Kacar, B. Reconstructing the evolutionary
614   history of nitrogenases: Evidence for ancestral molybdenum-cofactor utilization.
615   *Geobiology* **18**, 394-411, doi:10.1111/gbi.12381 (2020).

616   29   Glass, J. B., Wolfe-Simon, F. & Anbar, A. D. Coevolution of metal availability and nitrogen
617   assimilation in cyanobacteria and algae. *Geobiology* **7**, 100-123, doi:10.1111/j.1472-
618   4669.2009.00190.x (2009).

619   30   Hoffman, B. M., Lukoyanov, D., Yang, Z. Y., Dean, D. R. & Seefeldt, L. C. Mechanism of
620   nitrogen fixation by nitrogenase: the next stage. *Chem Rev* **114**, 4041-4062,
621   doi:10.1021/cr400641x (2014).

622   31   Raymond, J., Siefert, J. L., Staples, C. R. & Blankenship, R. E. The natural history of
623   nitrogen fixation. *Mol Biol Evol* **21**, 541-554, doi:10.1093/molbev/msh047 (2004).

624   32   Boyd, E. S. & Peters, J. W. New insights into the evolutionary history of biological nitrogen
625   fixation. *Front Microbiol* **4**, 201, doi:10.3389/fmicb.2013.00201 (2013).

626   33   Harris, D. F. *et al.* Mo-, V-, and Fe-Nitrogenases Use a Universal Eight-Electron
627   Reductive-Elimination Mechanism To Achieve N_2 Reduction. *Biochemistry* **58**, 3293-3301,
628   doi:10.1021/acs.biochem.9b00468 (2019).

629   34   Zheng, Y. *et al.* A pathway for biological methane production using bacterial iron-only
630   nitrogenase. *Nat Microbiol* **3**, 281-286, doi:10.1038/s41564-017-0091-5 (2018).

35 Hu, Y., Lee, C. C. & Ribbe, M. W. Extending the Carbon Chain: Hydrocarbon Formation
Catalyzed by Vanadium/Molybdenum Nitrogenases. *Science* **333**, 753-755,
doi:10.1126/science.1206883 (2011).

36 Rohde, M., Trncik, C., Sippel, D., Gerhardt, S. & Einsle, O. Crystal structure of VnfH, the
iron protein component of vanadium nitrogenase. *J Biol Inorg Chem* **23**, 1049-1056,
doi:10.1007/s00775-018-1602-4 (2018).

37 Schmid, B. *et al.* Biochemical and structural characterization of the cross-linked complex
of nitrogenase: comparison to the ADP-AlF4(-)-stabilized structure. *Biochemistry* **41**,
15557-15565 (2002).

38 Spatzal, T. *et al.* Evidence for interstitial carbon in nitrogenase FeMo cofactor. *Science*
334, 940, doi:10.1126/science.1214025 (2011).

39 Waugh, S. I. *et al.* The genes encoding the delta subunits of dinitrogenases 2 and 3 are
required for mo-independent diazotrophic growth by Azotobacter vinelandii. *Journal of*
Bacteriology **177**, 1505-1510, doi:10.1128/jb.177.6.1505-1510.1995 (1995).

40 Altschul, S. F., Gish, W., Miller, W., Myers, E. W. & Lipman, D. J. Basic local alignment
search tool. *Journal of Molecular Biology* **215**, 403-410, doi:10.1016/s0022-
2836(05)80360-2 (1990).

41 Altschul, S. F. *et al.* Gapped BLAST and PSI-BLAST: a new generation of protein
database search programs. *Nucleic Acids Research* **25**, 3389-3402,
doi:10.1093/nar/25.17.3389 (1997).

42 Finn, R. D., Clements, J. & Eddy, S. R. HMMER web server: interactive sequence
similarity searching. *Nucleic Acids Res* **39**, W29-37, doi:10.1093/nar/gkr367 (2011).

43 van Kempen, M. *et al.* Foldseek: fast and accurate protein structure search.
doi:10.1101/2022.02.07.479398 (2022).

44 Illergard, K., Ardell, D. H. & Elofsson, A. Structure is three to ten times more conserved
than sequence--a study of structural response in protein cores. *Proteins* **77**, 499-508,
doi:10.1002/prot.22458 (2009).

45 Medina, M. S. *et al.* CowN sustains nitrogenase turnover in the presence of the inhibitor
carbon monoxide. *J Biol Chem* **296**, 100501, doi:10.1016/j.jbc.2021.100501 (2021).

46 Rubio, L. M. & Ludden, P. W. Maturation of nitrogenase: a biochemical puzzle. *J Bacteriol*
187, 405-414, doi:10.1128/JB.187.2.405-414.2005 (2005).

47 Tautz, D. & Domazet-Loso, T. The evolutionary origin of orphan genes. *Nat Rev Genet*
12, 692-702, doi:10.1038/nrg3053 (2011).

48   Heames, B., Schmitz, J. & Bornberg-Bauer, E. A Continuum of Evolving De Novo Genes
Drives Protein-Coding Novelty in Drosophila. *J Mol Evol* **88**, 382-398, doi:10.1007/s00239-
020-09939-z (2020).

49   Vakirlis, N. *et al.* De novo emergence of adaptive membrane proteins from thymine-rich
genomic sequences. *Nat Commun* **11**, 781, doi:10.1038/s41467-020-14500-z (2020).

50   Vakirlis, N., Carvunis, A. R. & McLysaght, A. Synteny-based analyses indicate that
sequence divergence is not the main source of orphan genes. *Elife* **9**,
doi:10.7554/eLife.53500 (2020).

51   Dos Santos, P. C., Fang, Z., Mason, S. W., Setubal, J. C. & Dixon, R. Distribution of
nitrogen fixation and nitrogenase-like sequences amongst microbial genomes. *BMC
Genomics* **13**, 162, doi:10.1186/1471-2164-13-162 (2012).

52   Anbar, A. D. *et al.* A whiff of oxygen before the great oxidation event? *Science* **317**, 1903-
1906, doi:10.1126/science.1140325 (2007).

53   Kessler, P. S., McLarnan, J. & Leigh, J. A. Nitrogenase phylogeny and the molybdenum
dependence of nitrogen fixation in Methanococcus maripaludis. *J Bacteriol* **179**, 541-543
(1997).

54   Rutledge, H. L., Cook, B. D., Nguyen, H. P. M., Herzik, M. A. & Tezcan, F. A. Structures
of the nitrogenase complex prepared under catalytic turnover conditions. *Science* **377**,
865-869, doi:10.1126/science.abq7641 (2022).

55   Huang, Q. *et al.* Mechanical coupling in the nitrogenase complex. *PLoS Comput Biol* **17**,
e1008719, doi:10.1371/journal.pcbi.1008719 (2021).

56   Schmid, B. *et al.* Structure of a cofactor-deficient nitrogenase MoFe protein. *Science* **296**,
352-356, doi:10.1126/science.1070010 (2002).

57   Moore, E. K., Hao, J., Spielman, S. J. & Yee, N. The evolving redox chemistry and
bioavailability of vanadium in deep time. *Geobiology* **18**, 127-138, doi:10.1111/gbi.12375
(2020).

58   Emerson, S. R. & Huested, S. S. Ocean anoxia and the concentrations of molybdenum
and vanadium in seawater. *Marine Chemistry* **34**, 177-196, doi:10.1016/0304-
4203(91)90002-e (1991).

59   Zerkle, A. L., House, C. H. & Brantley, S. L. Biogeochemical signatures through time as
inferred from whole microbial genomes. *Am J Sci* **305**, 467-502, doi:DOI
10.2475/ajs.305.6-8.467 (2005).

60   Garcia, A. K., Kolaczkowski, B. & Kacar, B. Reconstruction of Nitrogenase Predecessors
697   Suggests Origin from Maturase-Like Proteins. *Genome Biol Evol* **14**,
698   doi:10.1093/gbe/evac031 (2022).

699   61   Ruttimann-Johnson, C., Rubio, L. M., Dean, D. R. & Ludden, P. W. VnfY is required for
700   full activity of the vanadium-containing dinitrogenase in Azotobacter vinelandii. *J Bacteriol*
701   **185**, 2383-2386, doi:10.1128/JB.185.7.2383-2386.2003 (2003).

702   62   Perez-Gonzalez, A. *et al.* Specificity of NifEN and VnfEN for the Assembly of Nitrogenase
703   Active Site Cofactors in Azotobacter vinelandii. *mBio* **12**, e0156821,
704   doi:10.1128/mBio.01568-21 (2021).

705   63   Rebelein, J. G., Lee, C. C., Newcomb, M., Hu, Y. & Ribbe, M. W. Characterization of an
706   M-Cluster-Substituted Nitrogenase VFe Protein. *MBio* **9**, doi:10.1128/mBio.00310-18
707   (2018).

708   64   Hazen, R. M. *et al.* Mineral evolution. *American Mineralogist* **93**, 1693-1720,
709   doi:10.2138/am.2008.2955 (2008).

710   65   Sayers, E. W. *et al.* Database resources of the national center for biotechnology
711   information. *Nucleic Acids Res* **50**, D20-D26, doi:10.1093/nar/gkab1112 (2022).

712   66   Katoh, K. & Standley, D. M. MAFFT multiple sequence alignment software version 7:
713   improvements in performance and usability. *Mol Biol Evol* **30**, 772-780,
714   doi:10.1093/molbev/mst010 (2013).

715   67   Fu, L., Niu, B., Zhu, Z., Wu, S. & Li, W. CD-HIT: accelerated for clustering the next-
716   generation sequencing data. *Bioinformatics* **28**, 3150-3152,
717   doi:10.1093/bioinformatics/bts565 (2012).

718   68   The Uniprot Consortium. UniProt: the Universal Protein Knowledgebase in 2023. *Nucleic*
719   *Acids Res* **51**, D523-D531, doi:10.1093/nar/gkac1052 (2023).

720   69   Nguyen, L. T., Schmidt, H. A., von Haeseler, A. & Minh, B. Q. IQ-TREE: a fast and effective
721   stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol Biol Evol* **32**,
722   268-274, doi:10.1093/molbev/msu300 (2015).

723   70   Kalyaanamoorthy, S., Minh, B. Q., Wong, T. K. F., von Haeseler, A. & Jermiin, L. S.
724   ModelFinder: fast model selection for accurate phylogenetic estimates. *Nature Methods*
725   **14**, 587-589, doi:10.1038/nmeth.4285 (2017).

726   71   Hoang, D. T., Chernomor, O., von Haeseler, A., Minh, B. Q. & Vinh, L. S. UFBoot2:
727   Improving the Ultrafast Bootstrap Approximation. *Mol Biol Evol* **35**, 518-522,
728   doi:10.1093/molbev/msx281 (2018).

72 Guindon, S. *et al.* New algorithms and methods to estimate maximum-likelihood
phylogenies: assessing the performance of PhyML 3.0. *Syst Biol* **59**, 307-321,
doi:10.1093/sysbio/syq010 (2010).

73 Garcia, A. K. *et al.* Nitrogenase resurrection and the evolution of a singular enzymatic
mechanism. *Elife* **12**, doi:10.7554/eLife.85003 (2023).

74 Yang, Z. PAML 4: phylogenetic analysis by maximum likelihood. *Mol Biol Evol* **24**, 1586-
1591, doi:10.1093/molbev/msm088 (2007).

75 Aadland, K., Pugh, C. & Kolaczkowski, B. High-Throughput Reconstruction of Ancestral
Protein Sequence, Structure, and Molecular Function. *Methods Mol Biol* **1851**, 135-170,
doi:10.1007/978-1-4939-8736-8_8 (2019).

76 Tareen, A. & Kinney, J. B. Logomaker: beautiful sequence logos in Python. *Bioinformatics*
36, 2272-2274, doi:10.1093/bioinformatics/btz921 (2020).

77 Sauer, D. B. & Wang, D. N. Predicting the optimal growth temperatures of prokaryotes
using only genome derived features. *Bioinformatics* **35**, 3224-3231,
doi:10.1093/bioinformatics/btz059 (2019).

78 Ashkenazy, H. *et al.* ConSurf 2016: an improved methodology to estimate and visualize
evolutionary conservation in macromolecules. *Nucleic Acids Res* **44**, W344-350,
doi:10.1093/nar/gkw408 (2016).

79 Penev, P. I., Alvarez-Carreno, C., Smith, E., Petrov, A. S. & Williams, L. D. TwinCons:
Conservation score for uncovering deep sequence similarity and divergence. *PLoS*
Comput Biol **17**, e1009541, doi:10.1371/journal.pcbi.1009541 (2021).

80 Mirdita, M. *et al.* ColabFold: making protein folding accessible to all. *Nature Methods* **19**,
679-682, doi:10.1038/s41592-022-01488-1 (2022).

81 Steinegger, M. & Söding, J. MMseqs2 enables sensitive protein sequence searching for
the analysis of massive data sets. *Nature Biotechnology* **35**, 1026-1028,
doi:10.1038/nbt.3988 (2017).

82 Pettersen, E. F. *et al.* UCSF ChimeraX: Structure visualization for researchers, educators,
and developers. *Protein Sci* **30**, 70-82, doi:10.1002/pro.3943 (2021).

83 Csermely, P. Creative elements: network-based predictions of active centres in proteins
and cellular and social networks. *Trends Biochem Sci* **33**, 569-576,
doi:10.1016/j.tibs.2008.09.006 (2008).

84 Bouysset, C. & Fiorucci, S. ProLIF: a library to encode molecular interactions as
fingerprints. *Journal of Cheminformatics* **13**, doi:10.1186/s13321-021-00548-6 (2021).

85   Hagberg, A. A., Schult, D. A. & Swart, P. J. in *Proceedings of the 7th Python in Science*
Conference. (eds Gäel Varoquaux, Travis Vaught, & Jarrod Millman) 11-15.

86   Bastian, M., Heymann, S. & Jacomy, M. Gephi: An Open Source Software for Exploring
and Manipulating Networks. *Proceedings of the International AAAI Conference on Web*
and Social Media **3**, 361-362, doi:10.1609/icwsm.v3i1.13937 (2009).

87   Jurrus, E. *et al.* Improvements to the APBS biomolecular solvation software suite. *Protein*
Science **27**, 112-128, doi:10.1002/pro.3280 (2017).

88   Kortemme, T., Kim, D. E. & Baker, D. Computational Alanine Scanning of Protein-Protein
Interfaces. *Science's STKE* **2004**, doi:10.1126/stke.2192004pl2 (2004).

89   Rodrigues, C. H. M., Myung, Y., Pires, D. E. V. & Ascher, D. B. mCSM-PPI2: predicting
the effects of mutations on protein–protein interactions. *Nucleic Acids Research* **47**,
W338-W344, doi:10.1093/nar/gkz383 (2019).

90   Gerek, Z. N. & Ozkan, S. B. Change in allosteric network affects binding affinities of PDZ
domains: analysis through perturbation response scanning. *PLoS Comput Biol* **7**,
e1002154, doi:10.1371/journal.pcbi.1002154 (2011).