1 **The nitrogenase G-subunit is an ancient orphan protein that drove the**

2 **ecological expansion of nitrogen fixation**

Bruno Cuevas-Zuviría[1,2,*], Amanda K. Garcia[1,*], Brooke M. Carruthers[1], Betül Kaçar[1,**]

[1]Department of Bacteriology, University of Wisconsin-Madison, Madison, WI, USA

[2]Centro de Biotecnología y Genómica de Plantas, Universidad Politécnica de Madrid, Madrid,

Spain

*These authors contributed equally to this work

**To whom correspondence should be addressed: bkacar@wisc.edu

13 **ABSTRACT**

Nitrogenase metalloenzymes have catalyzed biological nitrogen fixation for billions of years and

revolutionized planet Earth by supplying essential nitrogen to the biosphere. How these

enzymes were built and distributed by microbial and evolutionary processes in a shifting

geochemical landscape remains an open question. Here, we probe the birth and evolution of the

G-subunit protein, an integral, Precambrian-age structural component of certain nitrogenase

isozymes that makes its appearance midway through nitrogenase evolutionary history. We

establish that the G-subunit is an orphan protein, with no homologs detected across wider

protein diversity. We find that G-subunit emergence accompanied both the diversification of

nitrogenase metal usage and an ecological expansion of nitrogen-fixing microbes during the

transition in enviromental metal availabilities triggered by Earth surface oxygenation ~2.5 billion

years ago. Further, analyses of ancestral nitrogenase structures implicate a role for the G-

subunit in novel metal incorporation, which would have primed nitrogenases and their hosts to

exploit these newly diversified geochemical environments. However, permanent recruitment of

the G-subunit into the nitrogenase complex was likely only enabled by tuning preexisting,

protein interaction features that were selected prior to Earth oxygenation. Our results showcase

how contingent evolutionary novelties shape microbial ecological responses and their global

consequences.

INTRODUCTION

Over billions of years, life generated an enormous wealth of biomolecular diversity, producing an estimated 10^5 extant protein families[1] and a much larger multitude of extinct proteins. Through this process, biology became a defining component of the Earth system, which experienced tremendous revolutions in climate and biogeochemistry that are largely attributable to the molecular innovations of life itself[2]. A unified understanding of the Earth-life system requires investigating how the proteins and biogeochemical processes that power planetary phenomena emerged, evolved and proliferated.

In many cases, nature was a tinkerer[3], generating novel functions through the progressive duplication, accretion, and recombination of pre-existing protein elements[4-7]. In rarer cases, complete novelty arose through the *de novo* birth of genes[4,8]. This scenario represents a more difficult evolutionary hurdle due to the improbability that formerly noncoding DNA would produce a structurally or functionally coherent peptide[9]. Retracing these different evolutionary events in the histories of biogeochemically critical enzymes is a considerable challenge given their antiquity, since many of these enzymes, including those required for essential elemental cycling, likely evolved more than three billion years ago[2,10,11].

Here, we identify a case of molecular novelty in the evolutionary history of biological nitrogen fixation. Nitrogenase metalloenzymes provide the sole biochemical gateway for essential nitrogen into Earth's biosphere, consequently transforming the planet[10,12-14]. They have done so for more than 3 billion years[12,13]. This metalloenzyme family is today represented by three homologous isozymes—molybdenum- (Nif), vanadium- (Vnf), and iron-only (Anf) nitrogenases, the latter two which evolved more recently from Nif[15]. Each are named for the differing compositions of their active-site metalloclusters but otherwise share a core, two-component oxidoreductase architecture[16-18]. The antiquity and biological significance of this architecture is highlighted by its conservation in distantly related enzymes within the nitrogenase superfamily[19], the origin of which likely dates back to the last universal common ancestor[19,20].

A striking distinction in all known Vnf and Anf nitrogenases is a specific protein subunit called the "G-subunit"[21-24]. G-subunit does not exist in any known Nif nitrogenases, and it is an essential component for nitrogen fixation in modern Vnf and Anf nitrogenases[23,25]. The origin and evolutionary role of the G-subunit is not known. We know that the adaptive landscape of

67 nitrogenases shifted with transitions in environmental metal availabilities brought on by Earth
68 surface oxygenation ~2.5 billion years ago[26,27] which may have impacted nitrogen fixation
69 diversification[15,26,28]. To what degree was a novelty such as the G-subunit necessary for the
70 expansion of metal usage in nitrogen fixation?
71
In the present study, we elucidate the evolutionary history of the nitrogenase G-subunit and
address the specific interplay between molecular evolutionary events and environmental
triggers in the ecological diversification of biological nitrogen for the first time. We infer the
ancestral sequences and structures of nitrogenase enzymes through the emergence of the G-
subunit, replaying the sequence of events that gave rise to this nitrogenase protein component
>1.5 billion years ago. We further interrogate the genetic and environmental context of its
emergence, recruitment, and functional significane within the context of an ancient ecosystem.
79
80 **RESULTS**
81
82 **The nitrogenase G-subunit is an ancient orphan protein**
83
In modern nitrogen-fixing microbes, the nitrogenase G-subunit protein is a small, structural
element of the larger, two-component enzyme complex. The first component is a H-subunit
homodimer that delivers electrons to the second, catalytic component, which, in Nif, is a D- and
K-subunit heterotetramer but, in Vnf/Anf (collectively referred to as "alternative nitrogenases"), is
a D-, G-, and K-subunit heterohexamer[21] (**Figure 1A**). Thus, the G-subunit is a primary
distinction in the otherwise structurally comparable Nif, Vnf, and Anf complexes. To date, the
sole, publicly available crystallographic structure of the G-subunit is from *A. vinelandii* Vnf[21], a
113-amino-acid, four-helix protein (**Figure 1B;** the *A. vinelandii* Anf crystallographic structure
has only recently been described[29]). In both Vnf and Anf, the G-subunit neighbors the D-subunit
that houses the active-site metallocluster where N_2 is bound and reduced[30]. The metal content
of the metallocluster differs between Nif, Vnf, and Anf nitrogenase isozymes, incorporating Fe
and Mo ("FeMo-co"), Fe and V ("FeV-co"), or only Fe ("FeFe-co"), respectively. Finally, though
sharing a common ancestry[15,31,32] and core N_2-reduction mechanism[33], Nif, Vnf, and Anf
nitrogenases vary in their kinetic behavior and reactivities to different substrates[18,33-35].
98

Figure 1. Structure and genetics of the nitrogenase G-subunit. A) *Left, crystallographic structure of the extant A. vinelandii Vnf nitrogenase complex (VnfH, PDB 6Q93[36]; VnfDGK, PDB 5N6Y[21]) containing the G-subunit (dark blue). The Anf nitrogenase is inferred to be comparable in its multimeric structure, binding FeFe-co instead of FeV-co. Right, crystallographic structure of the extant Nif nitrogenase complex (NifH, PDB 1M34[37]; NifDK, PDB 3U7Q[38]) lacking the G-subunit.* **B)** *Secondary structure of the Vnf G-subunit.* **C)** *Arrangement of the Nif, Vnf, and Anf structural genes in A. vinelandii. Vnf/AnfG (dark blue) is located between Vnf/AnfD and Vnf/AnfK genes.*

We began our evolutionary study of the nitrogenase G-subunit by constraining its place within the broader diversity of protein sequence and structure. Specifically, we sought to identify distant homologs that might provide insights into G-subunit ancestry and function. Though genetic studies have established the G-subunit as essential for an active, N_2-reducing Vnf/Anf nitrogenase complex[23,25], its precise functional contribution remains enigmatic. We implemented a homology search workflow that included both sequence-based (BLASTP, TBLASTN[39], PSI-BLAST[40], HMMER[41]) and structure-based (Foldseek[42]) methods appropriate for detection of distantly related proteins. Our search returned >400 canonical G-subunit proteins, but, notably, we did not find convincing similarity to any other protein. PSI-BLAST and HMMER searches together returned <10 hits that we could exclude from the canonical G-subunit protein family

(i.e., were not encoded by genes within *vnf/anf* structural operons) (**Table 1, Figure S1**). In both

cases, only short sequence regions of scattered taxonomic affinities aligned to our query with

poor statistical significance. We were thus unable to rule out spurious matches. Additionally, we

used the structure-based method, FoldSeek, leveraging the greater conservation of protein

structure compared to sequence[43] in homology detection. We identified two uncharacterized,

AlphaFold-predicted structures that are not canonical G-subunit proteins but nevertheless share

its four-helix structural arrangement (**Table 1, Figure S2, S3**). However, sequence-level

similarity to G-subunit proteins is negligible and, given the lack of functional annotations for

these hits, their relevance to early G-subunit evolution is unclear. Finally, we note that we did

not detect sequence or structural similarity to NafY and CowN proteins, both of which have

previously been proposed to resemble the G-subunit in their interactions with the nitrogenase

complex[21,44,45].

If the G-subunit shares an ancient ancestry with another extant protein family, no strong vestige

of this evolutionary relationship appears presently detectable. The absence of distant homologs

to the G-subunit family is comparable to the case of "orphan genes," which are individual genes

that either have no resemblance to other sequences or have few, taxonomically restricted

relatives[8,46]. Proposed mechanisms for the emergence of orphan genes include duplication

followed by rapid divergence and *de novo* birth from previously non-genic DNA[4,8]. The latter has

more recently been recognized as a significant source of orphan genes[47-49] and thus would

plausibly explain G-subunit origins. Regardless of the specific originating mechanism, our

results indicate that the G-subunit family is a molecular innovation as unique to nitrogenases as

it is to broader protein diversity.

Table 1. Summary of G-subunit homolog search results. *Template modeling score for structural alignments.

Method	Database	Total hits (including canonical G-subunit proteins)	Distant hits	E-value range of distant hits	TM-score* of distant hits	Annotation of distant hits	Taxonomy of distant hits
BLASTP	NCBI non-redundant proteins (nr)	433	0	n/a	n/a	n/a	n/a
TBLASTN	NCBI nucleotide collection (nr/nt)	205	0	n/a	n/a	n/a	n/a
PSI-BLAST	NCBI non-redundant proteins (nr)	446	7	0.026-0.064	n/a	TIGR04141 family sporadically distributed protein, Uncharacterized protein	*Pseudomonas* spp.
HMMER	UniProt Reference Proteomes	94	1	0.061	n/a	Uncharacterized protein	*Coregonus* sp.
FoldSeek	Protein Data Bank	3	0	n/a	n/a	n/a	n/a
FoldSeek	AlphaFold-Uniprot/Swiss-Prot	39	2	5.2-8.7	0.52-0.54	Uncharacterized protein	*Vibrio vulnificus, Deinococcus grandis*

Invention of the G-subunit coincided with diversification of nitrogenase metal usage and host microbial ecology

We reconstructed phylogenies of nitrogenase G-subunit proteins and examined their genetic distribution across extant genomes and microbial host environments. We identified G-subunit homologs exclusively from *vnf* and *anf* gene clusters in bacterial and archaeal genomes (similarly, all identified *vnf*/*anf* clusters include the G-subunit), confirming that the G-subunit is a significant molecular distinction between Vnf/Anf and Nif nitrogenases[50]. In all analyzed clusters, *vnf/anfG* genes are located between *vnf/anfD and vnf/anfK* loci (**Figure 1C**). By contrast, *nifD* and *nifK* genes in our dataset are universally contiguous with small intergenic distances (median ≈ 25 bp) (**Figure S4**). Finally, all complete genomes harboring *vnf* and/or *anf* nitrogenase genes in our dataset also possess *nif* genes. Therefore, the presence of the G-subunit correlates with the capacity of a host diazotroph to express at least two different nitrogenase isozymes with distinct metal dependencies. We built two maximum-likelihood G-subunit phylogenies: the first from individual G-subunit protein sequences and the second from G-subunit sequences concatenated with other nitrogenase subunit sequences. Both trees show separate clustering of Vnf and Anf sequences, though deep branch support in the concatenated tree is significantly improved (**Figure 2, S5, S6**).

The overall topology of our concatenated nitrogenase tree mirrors previous phylogenetic analyses of nitrogenase proteins reconstructed without G-subunit sequences. Nesting of Vnf and Anf clades within Nif sequences indicates that Vnf/Anf are more recently evolved[15,32], which, by extension, establishes the G-subunit as the most recently evolved structural component of the nitrogenase complex. Minimum age estimates of the Vnf/Anf clade range from ~1.5 to 2.5 Ga, based on the timing of horizontal gene transfer events that pervade nitrogenase evolutionary history[51]. These estimates indicate that the Precambrian origin of the G-subunit roughly coincided with the initial accumulation of oxygen in the Earth surface environment[27], the earliest whiffs of which have been identified from geochemical signatures in ~2.5 to 3.0 Ga sediments[52,53] and did not reach modern levels until ~0.5 Ga[54].

Our analyses demonstrate that the origin of the G-subunit was associated both with the gain of novel metal dependence and the ecological expansion of nitrogen fixation following the rise of atmospheric oxygen. Vnf and Anf sequences specifically branch from within the "Group III" nitrogenase lineage[31], which is additionally populated by Nif homologs of either experimentally

characterized or computationally predicted Mo-dependence[15,55]. Group III Nif are exclusively
hosted by anaerobic taxa (primarily methanogenic Euryarchaeota and Firmicutes), many of
which are probable thermophiles or hyperthermophiles (optimal growth temperatures predicted
from genome content, see **Materials and Methods**) (**Figure 2, Supplementary Tables**). By
contrast, Vnf/Anf are hosted by comparatively more taxonomically diverse and mesophilic
archaea and bacteria that also harbor "Group I and II" Nif nitrogenases[31], including the well-
studied   diazotrophic   models   like   *A. vinelandii*,   *Clostridium   pasteurianum,*   and
Rhodopseudomonas palustris. Vnf/Anf are not hosted by organisms that harbor Group III Nif,
despite Vnf/Anf emerging from within the Group III lineage. Within Group III, only Vnf and Anf
clades contain sequences from aerobic taxa, including cyanobacteria which appear to only host
Nif and Vnf nitrogenases. Together, these data indicate that G-subunit-containing Vnf and Anf
nitrogenases were able to proliferate into ecological niches inaccessible to earlier diverged
Group III Nif.

Figure 2. Evolutionary and environmental history of the nitrogenases and the G-subunit.
Maximum-likelihood phylogeny built from concatenated nitrogenase subunit proteins. Oxygen
tolerance, predicted optimal growth temperatures (OGT), and taxonomy of represented host
microbes, as well as the metal content of the active-site metallocluster, are mapped to the
phylogeny (light grey bars for O_2 tolerance and predicted OGT indicate missing data). Nif clades
are labeled according to the nomenclature used by Raymond et al.[31] (e.g., Nif-I = Group I Nif,
etc.). The G-subunit origin event is indicated by a black hash mark. Branches that track the
emergence of the G-subunit and its diversification into Vnf and Anf groups are shown in black,
whereas branches lacking the G-subunit are shown in grey. Ancestral nodes (Nif^{Anc}, VnfAnf^{Anc},
Vnf^{Anc}, Anf^{Anc}) targeted for sequence reconstruction and analysis are highlighted by bold
outlined, white circles.

An ancient assembly pathway preconditioned early nitrogenases for G-subunit
interaction

To replay the first evolutionary steps in the birth of the G-subunit, we reconstructed and
analyzed ancestral nitrogenase protein sequences. We examined the degree to which ancestral
protein surface features were remodeled to accommodate a new subunit, with the expectation
that G-subunit interactions would have driven the evolution of complementary residues within
the interface. Because the G-subunit is exclusive to Vnf and Anf nitrogenases, the most
parsimonious evolutionary scenario is that it emerged along the branch leading to the Vnf/Anf
last common ancestor, VnfAnf^{Anc} (**Figure 2**). We therefore compared sequences and structures
of nitrogenase ancestors reconstructed both immediately before (Nif^{Anc}) and after (VnfAnf^{Anc})
gain of the G-subunit.

We mapped the interface between ancestral G- and D-subunit proteins ("G-D interface") by
building its residue interaction network from the predicted VnfAnf^{Anc} structure (**Figure 3A-3D**).
Nearly all well-conserved, G-subunit residues lie within the G-D interface and contribute
significantly to its stability (**Figure S7**). The few exceptions are conserved residues distal from
the interface with side chains oriented towards the interior of the helical bundle, which likely
stabilize its globular arrangement.

We tracked the evolution of D-subunit residues through the emergence of the ancestral G-D
interface. We identified 42 D-subunit sites interacting with the G-subunit following its initial

227 recruitment in ancestral VnfAnf^{Anc} (**Supplementary Tables**). Of the 42 sites, 26 are well-

228 conserved among extant Vnf/Anf nitrogenases and form a core interaction patch that was likely

229 integral to the early G-D interface (**Figure 3E, 3F**). We inspected these 26 sites and found that

230 nine sites evolved (i.e., were substituted) alongside G-subunit emergence, imposing subtle

231 changes to the electrostatic surface potential of the VnfAnf^{Anc} interaction patch (**Figure S8, S9**).

Figure 3. Birth of the nitrogenase G-subunit interface. A) *Predicted Nif^Anc D-subunit*
structure, prior to G-subunit emergence. Inset shows cross-section at the putative metallocluster
insertion site, the entrance of which forms a "cleft" in the D-subunit surface. Superimposed G-
subunit structure (transparent) highlights relative position of the cleft. **B)** *Predicted VnfAnf^Anc D-*

*and G-subunit structures, following G-subunit emergence, shown in the same view as A). Side chain of the G-subunit Leu18 residue is anchored within the D-subunit cleft. **A-B)** Heterometal content for the ancestor metalloclusters is inferred. **C)** Nif^Anc G-D proto-interaction network. Purple residues are located within an interface patch conserved in extant Vnf/Anf. Residues that are later substituted in VnfAnf^Anc are shown bold and labeled to permit comparison with D). Residues within the cleft are circled. **D)** VnfAnf^Anc G-D interaction network, shown in the same view as C). Residues newly substituted in VnfAnf^Anc are shown in dark purple and labeled. **E)** Surface view of the Nif^Anc D-subunit proto-interface patch (purple). **F)** Surface view of the VnfAnf^Anc D-subunit interface patch, shown in the same view as E). Residues that are maintained between NifAnc and VnfAnfAnc are shown in purple, and those that are substituted are shown in dark purple. **G)** Evolution of residues within the interaction patch through the emergence of the G-subunit. For substituted sites, the posterior probability of the most likely VnfAnf^Anc residue is <.3 in Nif^Anc and >.7 in VnfAnf^Anc and, for maintained sites, >.7 in Nif^Anc and >.7 in VnfAnf^Anc.*

Intriguingly, we find that multiple ancestral D-subunit surface features appear to have been preconditioned for G-subunit binding, collectively forming a "proto-interface" upon which interactions with the newly evolved G-subunit was built. Out of 26 residues within the core, 16 (~62%) show that the D-subunit interaction patch had already become VnfAnf^Anc-like prior to G-subunit emergence (**Figure 3F,G**). This residue-level conservation translates to maintained hydrophobicity (median apolar fraction ≈ .2) of the (proto-)interaction patch both before and after G-subunit recruitment (**Figure S10**). Further, shape complementarity as an integral feature of the G-D interface appears to have preceded the G-subunit. In *A. vinelandii* Vnf[21], a cleft within the D-subunit interaction patch is occupied by a strictly conserved G-subunit residue, Leu18. This residue is among the most well-connected within the G-D interaction network (**Figure 3A-D**), is critical for interface stability (**Figure S7**), and is among the most allosterically active, capable of generating long-distance effects that reach across opposite ends of the nitrogenase complex (**Figure S11**). These effects reflect the importance of the G-D interface in controlling the global motions of the nitrogenase complex and evoke similar, long-distance structural perturbations that are induced through the nitrogenase catalytic cycle[56,57]. The cleft itself provides an important anchor point for G-subunit binding and is present in both ancestral and representative extant nitrogenase structures (**Figure 3A,B, Figure 4**).

Figure 4. Conservation of the D-subunit cofactor insertion cleft across extant and ancestral nitrogenases. *Small black arrow indicates position of the proposed cofactor insertion cleft.* **A)** *Crystallographic structure of the A. vinelandii apo-NifD protein (PDB 1L5H). Cross-section view (right) reveals the proposed metallocluster insertion site[58], the entrance of which structurally aligns with the G-D interface of VnfD. Aligned, cross-section views are displayed for extant* **B)** *A. vinelandii NifD (PDB 3U7Q),* **C)** *Clostridium pasteurianum NifD (PDB 4WES), and* **D)** *A. vinelandii VnfD (PDB 5N6Y) crystallographic structures as well as predicted, ancestral* **E)** *Nif^{Anc} D-subunit and* **F)** *VnfAnf^{Anc} D-subunit structures.* **(A-C, E)** *A. vinelandii VnfG (PDB 5N6Y; transparent) is superimposed to enable comparison across nitrogenases with and without native G-subunit proteins.* **(A-F)**

The functional significance of the D-subunit cleft provides an explanation for why the interface
patch resisted significant remodeling with G-subunit addition, as well as implicates an adaptive
role for the G-subunit. In extant Nif, the cleft aligns with the entrance to a proposed nitrogenase
metallocluster insertion site, which, in the apo-NifD protein, adopts an open conformation to
permit cluster loading[58]. Given the consistent presence of the cleft across Nif, Vnf, and Anf
nitrogenases, a similar mechanism can be predicted for Vnf and Anf and is likely an ancient,
universal feature of nitrogenase assembly that predates the G-subunit and was maintained due
to its essentiality. In fact, we find that 12 interface residues surrounding the insertion site are
well-conserved in extant Group III Nif clades that diverge prior to the origin of the G-subunit, and
five of these are either universally or nearly universally conserved among all nitrogenases
(**Figure S12**). Structural alignments indicate that the G-subunit caps the opening of the insertion
site, with the side chain of the conserved Leu18 residue oriented into the cleft and toward the
loaded active site. An unusual feature of the G-subunit, a carboxyl-terminal tail that terminates
in a strictly conserved tyrosine residue (Tyr113), bridges the presumed opening of the apo-D-
subunit protein and neighbors universally conserved D-subunit residues. The identified
interactions between the bound G-subunit and the proposed metallocluster insertion site in the
D-subunit implicate a role for the G-subunit in nitrogenase assembly, which is supported by
experimental evidence that VnfG and VFe-co of *A. vinelandii* associate *in vitro*, as well as of the
requirement of both VnfG and VFe-co to stabilize an active complex[23]. Conservation of G-
subunit residues surrounding the cleft (and thus the metallocluster insertion site) across our
phylogenetic dataset suggests that this putative role was early-evolved and maintained through
its evolution.

The G-subunit confers specialization between Vnf and Anf nitrogenases

We assessed to what degree sequence variation within the G-subunit is capable of driving the
functional distinctions between Vnf and Anf isozymes, which are each dependent on distinct
metalloclusters and exhibit varied N_2-reduction efficiencies and reactivities to different chemical
substrates[18,33-35]. We identified protein sites that are "specialized" between Vnf and Anf (i.e.,
diverge between Vnf and Anf but are otherwise conserved in each of the two groups), and
therefore most likely drive functional variation. We find the degree of specialization (relative to
protein size) within the G-subunit to be on par with that of the D-subunit but observe
comparatively little specialization within the K-subunit (**Figure 5A, 5B, Supplementary Tables**).

Therefore, the G-subunit contributes an outsized fraction of the total sequence specialization
between Vnf and Anf, despite its remoteness from any nitrogenase metalloclusters. Specialized
sites within the G-subunit are distal from the G-D interface. These include one residue close to
the presumed H-subunit binding site, which might contribute to a previously proposed role in
mediating H-subunit binding specificity[24].
323

We next examined how these signatures of sequence-level specialization manifested early in G-
subunit evolution, tracking their amino acid content through the divergence of Vnf and Anf
clades. We find that the specialized sequence motifs of Vnf[Anc] (Vnf ancestor) and Anf[Anc] (Anf
ancestor) most closely resemble those of their respective descendants (**Figure 5C**). The most
likely Vnf[Anc] motif across divergent sites is "FIKIDV" (extant Vnf consensus "FIKVDH") whereas
the most likely Anf[Anc] motif is "VNVDYD" (extant Anf consensus "VNVDYD"). However, we find
that the most likely motif of the common VnfAnf[Anc] ancestor, "VIKIDV," is more like that of Vnf
than Anf, even accounting for the statistical uncertainty associated with ancestral reconstruction.
This result suggests that the earliest G-subunit proteins were Vnf-like (in agreement with
previous inferences of ancestral nitrogenases that indicate Vnf evolved prior to Anf[15]), and that
the G-subunit contributed to specificity required for vanadium dependency early in the birth of a
new nitrogenase isozyme.
336

Figure 5. Coevolution of G-subunit specialization and Vnf/Anf metal dependence*. A)*
Specialized sites mapped to the nitrogenase VnfDGK structure (PDB 5N6Y[21]). Scale in A)
applies to B). ***B)*** *Detailed view of specialized sites across the G-subunit.* ***C)*** *Evolution of*
specialized G-subunit sequence motifs through the divergence of Vnf and Anf. Sequence logos
scale either with amino acid frequencies (for extant Vnf or Anf nitrogenases) or posterior
probabilities (for ancestral VnfAnf[Anc]) at each site. VnfAnf[Anc] dependence on FeV-co is inferred
based on comparisons of specialized sequence motifs. ***D)*** *The bulk marine concentrations of*
redox-sensitive metals involved in nitrogen fixation (Mo, V, Fe) have changed with progressive
oxygenation of the Earth surface environment, with Fe more abundant in the Archean and Mo/V
more abundant today. Though quantitative estimates of historical V concentrations are not
presently available, oxygenation likely impacted the speciation and solubility of V and Mo
similarly over geologic time[59,60]. Metal concentrations from Zerkle et al.[61] and references therein
and p$_{O2}$ from Lyons et al.[27]. Age constraints of nitrogenase origins and VnfAnf[Anc] (i.e., G-subunit
emergence) from Stueken et al.[62] and Parsons et al.[51], respectively. GOE: Great Oxidation
Event, PAL: present atmospheric level.

DISCUSSION

In the present study, we interrogated the origin, early evolution, and functional significance of a

major novelty in the history of nitrogen fixation: the emergence of the G-subunit in Vnf and Anf

nitrogenases >1.5 billion years ago[51]. With the birth of the nitrogenase G-subunit, not only did a

rare evolutionary event occur with associated fitness costs overcome[4], but a novel structural

domain became fully integrated into the nitrogenase complex and was retained in one of the

most central metabolic processes to Earth's biosphere.

We demonstrate that there currently exists no candidate, evolutionary predecessor of the G-

subunit family, whether another protein, protein fragment, or segment of non-genic DNA. This

finding sets the origin of the G-subunit apart from other major evolutionary events in the history

of nitrogen fixation that are explainable by gene duplication, including the divergence between

other catalytic subunits, between nitrogenases and their maturation protein scaffolds, and

between nitrogenase isozymes[32,63]. The addition of such a small and evolutionarily enigmatic

protein domain appears to be an anomaly across biogeochemical enzymes.

We propose that the G-subunit was formed *de novo* from previously non-genic DNA, more

recently acknowledged as a viable process for generating genetic novelty[47-49]. Such an event

may have accompanied the genetic duplication and divergence of Nif that produced alternative

nitrogenases[32]. Regardless of its originating mechanism, retainment of the G-subunit gene

within the nitrogenase operon appears to have been improbable. Insertions between

nitrogenase D- and K-subunit genes must either be rare or frequently deleterious, as evidenced

by their small intergenic distance and absence of any other disrupting open reading frames

across extant Nif clusters in our dataset. Thus, the evolution of the G-subunit constitutes an

unprecedented gain in structural and genetic complexity in the evolutionary history of nitrogen

fixation.

By analyzing predicted, nitrogenase ancestors, we reveal an evolutionary steppingstone in the

initial recruitment of the G-subunit. Our reconstruction of ancestral, D-subunit surface features

indicates that the G-subunit binding site was built from an essential, preexisting metallocluster

insertion site, which additionally appears preconditioned for G-subunit interactions. Given that

all known extant nitrogenases share this particular assembly scheme, its presence likely dates

as far back as the oldest geochemical evidence of biological nitrogen fixation at ~3.2 Ga[62],

hundreds of millions of years prior to the earliest whiffs of Earth oxygenation[52,53]. How then did

this essential surface feature accommodate burial by a new interaction partner? Metallocluster
delivery in extant nitrogenases is thought to be mediated by transient protein-protein
interactions[58,64,65] that would presumably have shaped nitrogenase surface properties, perhaps
explaining the early evolved hydrophobicity of the ancestral, proto-interface. Our evolutionary
analyses, coupled with experimental support of a role in nitrogenase assembly[23], suggest that
the G-subunit was integrated within the assembly pathway, replacing and/or mediating earlier
evolved protein or metallocluster interactions. Thus, the birth of the permanent G-D interface
represents a gradual subsummation of formerly transient interactions that provided an
evolutionary "cradle" for a permanent, novel structural domain.

In its capacity to mediate newly evolved incorporation of V- and Fe-based clusters, the G-
subunit would have enabled the evolutionary burst of both metal dependence and ecological
diversity within the Group III nitrogenase lineage from which alternative nitrogenases originated.
We find that G-subunit residues, despite being relatively distal from the active-site and not
permanently retaining bound, reactive metalloclusters like other nitrogenase subunits, have an
outsized contribution to the sequence-level specialization between alternative nitrogenases.
Possible functional outcomes include improved specificity toward metalloclusters or associated
protein partners during nitrogenase assembly (preventing costly mismetallation[66])[23] and
mediating crosstalk between electron transfer and catalytic nitrogenase components[24]. Each of
these functions would have equipped diazotrophs to coordinate multiple nitrogenases systems
with unique metal dependencies within the same cellular environment. This ability is
advantageous in ancient or modern environments that experience transient molybdenum
limitation, with alternative nitrogenases providing a "fail-safe" strategy under conditions when
molybdenum-dependent nitrogen fixation cannot be supported[67]. The G-subunit therefore
primed these organisms for new, molybdenum-limited ecological niches, as well as for
fluctuations in redox-sensitive metal availabilities following the rise of oxygenic phototrophs[27,68],
one of the most significant ecological revolutions in Earth history.

An ongoing debate in evolutionary biology centers on the degree to which environmental
conditions drive biological innovations. In the evolution of biological nitrogen fixation, an intuitive
narrative is that the diversification of nitrogenase metal dependence, intimately tied to the
productivity and ecological distribution of nitrogen fixation, was triggered largely by transitions in
bulk, environmental metal availabilities beginning ~2.5 to 3.0 Ga[26,52,53], the historical records of
which are primarily built from geochemical analyses of ancient marine sediments (**Figure 5D**).

Our study highlights the additional importance of molecular evolutionary events emerging from
specific historical trajectories that predate, and are therefore removed from, candidate
environmental triggers. The birth of the nitrogenase G-subunit, by providing a crucial component
in novel metal incorporation, expanded the adaptive landscape for biological nitrogen fixation.
This ancient, orphan protein was birthed by seemingly improbable events but its recruitment
was nevertheless enabled by a conserved, nitrogenase assembly pathway that arose hundreds
of millions of years prior to the rise of Earth surface oxygen and the accompanying shifts in
metal availabilities. Whether this evolutionary transition in the history of biological nitrogen
fixation could have been crossed in the absence of the G-subunit awaits further studies. Similar,
future synergies between historical studies of molecular novelties and the environmental
circumstances that shaped them can refine our understanding of how the Earth-life planetary
system has evolved over billions of years.
435
436   **METHODS**
437
438   **Nitrogenase protein sequence curation**
439
A preliminary 412-sequence dataset of nitrogenase Vnf/AnfG proteins was compiled by
BLASTP[39] search (Expect value threshold = 0.1) against the National Center for Biotechnology
Information (NCBI) non-redundant protein sequence database (nr)[69] (accessed January 2022),
using the *A. vinelandii* VnfG query sequence (WP_012698949.1). BLASTP hits were curated by
filtering against partial sequences and homologs whose encoding genes are distantly located
from *vnf* or *anf* operons (and thus unlikely to be canonical *vnf/anfG* genes). Certain VnfG
sequences were manually extracted from fused VnfDG proteins, using MAFFT v7.490[70]
alignments to other D- or G-subunit sequences as a guide. The preliminary Vnf/AnfG sequence
dataset was pruned by CD-HIT[71] (sequence identity threshold = 90%), yielding a final 188-
sequence G-subunit dataset.
450
451   **G-subunit homolog search**
452
In addition to the BLASTP search described above, three sequence-based search tools–
TBLASTN[39], PSI-BLAST[40], and HMMER[41]–were used to identify distantly related homologs of
the nitrogenase G-subunit (conducted January 2022). TBLASTN and PSI-BLAST searches
against the NCBI nucleotide collection (nr/nt) and nr protein database, respectively, were

executed with the *A. vinelandii* VnfG query (WP_012698949.1), an Expect value threshold =
0.1, and word size = 2 (all other parameters default). HMMER search (hmmsearch) against the
UniProt Reference Proteome database[72] was performed using an HMM profile generated from
our MAFFT-aligned, 188-sequence G-subunit dataset (see above) with an Expect value
threshold = 0.1 for both sequences and hits (all other parameters default).
462
463   Structure-based searches of distant G-subunit homologs were performed by the FoldSeek
464   server[42] using the *A. vinelandii* VnfG crystallographic structure (PDB 5N6Y[21]) query against the
465   Protein Data Bank and AlphaFold-UniProt/Swiss-Prot databases. We employed both 3Di-AA
466   and TM-Align modes with an 3Di-AA Expect value threshold = 0.01 or a TM-score = 0.5
467   (template modeling score).
468
469   **Phylogenetic and ancestral sequence reconstruction**
470
The curated, 188-sequence Vnf/AnfG dataset was aligned by MAFFT. No outgroup to G-subunit
sequences was included because the distant homology search described above did not yield
convincing hits (see **Results**). IQ-TREE v1.6.12[73] was used to reconstruct a maximum-
likelihood G-subunit protein phylogeny, using the best-fit evolutionary model, WAG+R6
(selected by the Bayesian Information Criterion, BIC), tested by ModelFinder[74], with branch
support calculated by ultra-fast bootstrap approximation (UFBoot)[75] and the Shimodaira-
Hasegawa-like approximate likelihood-ratio test (SH-like aLRT)[76].
478
A phylogeny of concatenated Vnf/AnfHDGK proteins was generated by retrieving syntenous
subunit sequences to the curated VnfG/AnfG dataset. Certain VnfH sequences that recently
diverged from NifH clades were omitted due to having different evolutionary histories than those
of other Vnf/Anf proteins. NifHDK sequences from a prior phylogenetic analysis[77] were included
as an outgroup (n = 67, closely branching to Anf/Vnf) and were used to calculate intergenic
nifD-to-*nifK* distances (n = 306). Each subunit sequence dataset was aligned by MAFFT prior to
concatenation. Maximum-likelihood phylogenetic reconstruction was performed by IQ-TREE as
described above (best-fit evolutionary model, BIC: LG+F+R9).
487
Marginal maximum-likelihood ancestral sequence reconstruction of concatenated Vnf/AnfHDGK
sequences was performed by PAML v4.9j[78] (LG+F+G4 model). Mean sitewise posterior
probabilities D- and G-subunit ancestors analyzed in this study range from .75 to .96 (**Table**

S1). Sequence gaps were reconstructed as described by Aadland et al.[79]. Briefly, the input

protein sequence alignment was recoded as a binary amino-acid presence/absence matrix.

Ancestral presence/absence states were reconstructed by PAML using a binary character

model and state frequencies inferred by maximum likelihood.

Logo representations of extant multiple sequence alignments were generated with the Python

library LogoMaker[80] using the probability transformation of the count matrix. Representations of

ancestral sequences were generated from the posterior probability matrices produced by PAML.

Sequence and phylogenetic tree datasets are available at

https://github.com/kacarlab/nitrogenase-g-subunit.

Optimal growth temperature prediction

For taxa represented in the nitrogenase G-subunit sequence dataset, optimal growth

temperatures (OGT) were predicted from genomic content using the method described by

Sauer and Wang[81]. Due to genome incompleteness and the absence of 16S rRNA sequences

in fragmented genomes for certain representative taxa, the selected "Superkingdom Bacteria"

and "Superkingdom Archaea" regression models for OGT prediction excluded genome size and

16S rRNA.

Protein conservation and divergence analysis

Sequence conservation for D- and G-subunit proteins was determined by the ConSurf server[82].

Sites were considered well-conserved if their ConSurf grade were >7. Sequence specialization

was determined by the TwinCons package[83]. TwinCons was run separately on individual

alignments of extant H, D, G, and K subunit protein sequences with user-defined Vnf and Anf

groups, using default parameters and the LG substitution matrix. TwinCons specialization

scores were normalized to a 0-to-1 scale per analyzed subunit.

Structural Modeling

Ancestral and extant NifHDGK protein structures were predicted using ColabFold[84], which

combines the AlphaFold2 method for structure prediction from multiple-sequence alignments[18]

with the MMSeq2 method to obtain evolutionary information[85]. Specifically, the colabfold_batch v1.4.0 implementation (https://github.com/YoshitakaMo/localcolabfold) was used with the AlphaFold-v2.0-multimer prediction model and standard options (3 recycles and AMBER all-atom optimization in GPU).Protein structures were visualized by ChimeraX[86].

Structure Analysis

Residue interaction networks[87] were generated using ProLif[88], which finds the different possible interactions (hydrophobic, electrostatic, hydrogen bonds, etc) between two groups of atoms (here defined as G- and D-subunit groups) within a structure snapshot (e.g. single structure, trajectory frame). Interactions were compiled into networks by NetworkX[89], which were later visualized in Gephi[90]. Scripts used to generate the networks are available at https://github.com/kacarlab/nitrogenase-g-subunit.

Surface electrostatic maps were computed using the Poisson Boltzmann Solvent-Accessible model implemented in APBS[91] through the PyMOL plugin (Schrödinger, Inc), using standard PDB2PQR and APBS protocols at pH 7.

Sitewise contribution to D-G interface stability was computed using *in-silico* alanine scanning, which substitutes residues with a minimal methyl moiety (alanine)[92]. The changes on the binding free energy values upon substitution were obtained using the machine-learning enabled mCSM2 method[93] with default parameters.

Allosteric effects across the nitrogenase complex was modeled using Perturbation Response Scanning[94], implemented in ProDy[94] with default parameters and provided with PDB 5N6Y[21] alpha carbon coordinates. Scripts used to model allosteric effects are available at https://github.com/kacarlab/nitrogenase-g-subunit.

ACKNOWLEDGEMENTS

This research was supported by the National Aeronautics and Space Administration (NASA) Interdisciplinary Consortium for Astrobiology Research: Metal Utilization and Selection Across Eons, MUSE (19-ICAR19_2–0007), the NASA Arizona Space Grant (B.M.C), the John Templeton Foundation (B.K.; 61926), the National Science Foundation (B.K.; 2228495), the

NASA Early Career Faculty Award (B.K.), the Hypothesis Fund Award (B.K.), and the Universidad Politécnica de Madrid (UPM) Margarita Salas Fellowship, founded by the Unión Europea - Next Generation EU (B.C.-Z.; UP2021-035). We thank Morgan Sobol for assistance with optimal growth temperature prediction, as well as Ariel Anbar, Alex Rivier, Derek Harris, Holly Rucker, Joanna Masel, Lance Seefeldt and Steven Russell for valuable discussions and feedback.

REFERENCES

Choi, I.-G. & Kim, S.-H. Evolution of protein structural classes and protein sequence families. *Proc Natl Acad Sci USA* **103**, 14056-14061, doi:10.1073/pnas.0606239103 (2006).

Knoll, A. H. The geological consequences of evolution. *Geobiology* **1**, 3-14, doi:10.1046/j.1472-4669.2003.00002.x (2003).

Jacob, F. Evolution and Tinkering. *Science* **196**, 1161-1166, doi:10.1126/science.860134 (1977).

Andersson, D. I., Jerlstrom-Hultqvist, J. & Nasvall, J. Evolution of new functions de novo and from preexisting genes. *Cold Spring Harb Perspect Biol* **7**, doi:10.1101/cshperspect.a017996 (2015).

Eck, R. V. & Dayhoff, M. O. Evolution of the Structure of Ferredoxin Based on Living Relics of Primitive Amino Acid Sequences. *Science* **152**, 363-366, doi:10.1126/science.152.3720.363 (1966).

Copley, S. D. Evolution of new enzymes by gene duplication and divergence. *FEBS J* **287**, 1262-1283, doi:10.1111/febs.15299 (2020).

Alva, V., Soding, J. & Lupas, A. N. A vocabulary of ancient peptides at the origin of folded proteins. *Elife* **4**, e09410, doi:10.7554/eLife.09410 (2015).

Van Oss, S. B. & Carvunis, A. R. De novo gene birth. *PLoS Genet* **15**, e1008160, doi:10.1371/journal.pgen.1008160 (2019).

Beasley, J. R. & Hecht, M. H. Protein design: the choice of de novo sequences. *J Biol Chem* **272**, 2031-2034, doi:10.1074/jbc.272.4.2031 (1997).

Falkowski, P. G., Fenchel, T. & Delong, E. F. The microbial engines that drive Earth's biogeochemical cycles. *Science* **320**, 1034-1039, doi:10.1126/science.1153213 (2008).

11   Moore, E. K., Jelen, B. I., Giovannelli, D., Raanan, H. & Falkowski, P. G. Metal
availability and the expanding network of microbial metabolisms in the Archaean eon.
Nat Geosci **10**, 629-636, doi:10.1038/ngeo3006 (2017).

12   Falkowski, P. G. Evolution of the nitrogen cycle and its influence on the biological
sequestration of CO_2 in the ocean. *Nature* **387**, 272-275, doi:10.1038/387272a0 (1997).

13   Allen, J. F., Thake, B. & Martin, W. F. Nitrogenase Inhibition Limited Oxygenation of
Earth's   Proterozoic   Atmosphere.   *Trends   Plant   Sci*   **24**,   1022-1031,
doi:10.1016/j.tplants.2019.07.007 (2019).

14   Sanchez-Baracaldo, P., Ridgwell, A. & Raven, J. A. A neoproterozoic transition in the
marine nitrogen cycle. *Curr Biol* **24**, 652-657, doi:10.1016/j.cub.2014.01.041 (2014).

15   Garcia, A. K., McShea, H., Kolaczkowski, B. & Kacar, B. Reconstructing the evolutionary
history of nitrogenases: Evidence for ancestral molybdenum-cofactor utilization.
Geobiology **18**, 394-411, doi:10.1111/gbi.12381 (2020).

16   Einsle, O. & Rees, D. C. Structural Enzymology of Nitrogenase Enzymes. *Chem Rev*
120, 4969-5004, doi:10.1021/acs.chemrev.0c00067 (2020).

17   Mus, F., Alleman, A. B., Pence, N., Seefeldt, L. C. & Peters, J. W. Exploring the
alternatives   of   biological   nitrogen   fixation.   *Metallomics*   **10**,   523-538,
doi:10.1039/c8mt00038g (2018).

18   Eady, R. R. Structure-function relationships of alternative nitrogenases. *Chemical
Reviews* **96**, 3013-3030, doi:10.1021/cr950057h (1996).

19   Ghebreamlak, S. M. & Mansoorabadi, S. O. Divergent Members of the Nitrogenase
Superfamily: Tetrapyrrole Biosynthesis and Beyond. *ChemBioChem* **21**, 1723-1728,
doi:10.1002/cbic.201900782 (2020).

20   Weiss, M. C. *et al.* The physiology and habitat of the last universal common ancestor.
Nat Microbiol **1** (2016). <https://www.ncbi.nlm.nih.gov/pubmed/27562259>.

21   Sippel, D. & Einsle, O. The structure of vanadium nitrogenase reveals an unusual
bridging ligand. *Nat Chem Biol* **13**, 956-960, doi:10.1038/nchembio.2428 (2017).

22   Eady, R. R., Robson, R. L., Richardson, T. H., Miller, R. W. & Hawkins, M. The
vanadium nitrogenase of Azotobacter chroococcum. Purification and properties of the
VFe protein. *Biochemical Journal* **244**, 197-207, doi:10.1042/bj2440197 (1987).

23   Chatterjee, R., Ludden, P. W. & Shah, V. K. Characterization of VNFG, the delta subunit
of the vnf-encoded apodinitrogenase from Azotobacter vinelandii. Implications for its role
in the formation of functional dinitrogenase 2. *J Biol Chem* **272**, 3758-3765,
doi:10.1074/jbc.272.6.3758 (1997).

24   Pence, N., Lewis, N., Alleman, A. B., Seefeldt, L. C. & Peters, J. W. Revealing a role for
the G subunit in mediating interactions between the nitrogenase component proteins. *J*
Inorg Biochem **214**, 111273, doi:10.1016/j.jinorgbio.2020.111273 (2021).

25   Waugh, S. I. *et al.* The genes encoding the delta subunits of dinitrogenases 2 and 3 are
required for mo-independent diazotrophic growth by Azotobacter vinelandii. *Journal of*
Bacteriology **177**, 1505-1510, doi:10.1128/jb.177.6.1505-1510.1995 (1995).

26   Anbar, A. D. & Knoll, A. H. Proterozoic ocean chemistry and evolution: a bioinorganic
bridge? *Science* **297**, 1137-1142, doi:10.1126/science.1069651 (2002).

27   Lyons, T. W., Reinhard, C. T. & Planavsky, N. J. The rise of oxygen in Earth's early
ocean and atmosphere. *Nature* **506**, 307-315, doi:10.1038/nature13068 (2014).

28   Glass, J. B., Wolfe-Simon, F. & Anbar, A. D. Coevolution of metal availability and
nitrogen assimilation in cyanobacteria and algae. *Geobiology* **7**, 100-123,
doi:10.1111/j.1472-4669.2009.00190.x (2009).

29   Trncik, C., Detemple, F. & Einsle, O. Iron-only Fe-nitrogenase underscores common
catalytic principles in biological nitrogen fixation. *Nature Catalysis* **6**, 415-424,
doi:10.1038/s41929-023-00952-1 (2023).

30   Hoffman, B. M., Lukoyanov, D., Yang, Z. Y., Dean, D. R. & Seefeldt, L. C. Mechanism of
nitrogen fixation by nitrogenase: the next stage. *Chem Rev* **114**, 4041-4062,
doi:10.1021/cr400641x (2014).

31   Raymond, J., Siefert, J. L., Staples, C. R. & Blankenship, R. E. The natural history of
nitrogen fixation. *Mol Biol Evol* **21**, 541-554, doi:10.1093/molbev/msh047 (2004).

32   Boyd, E. S. & Peters, J. W. New insights into the evolutionary history of biological
nitrogen fixation. *Front Microbiol* **4**, 201, doi:10.3389/fmicb.2013.00201 (2013).

33   Harris, D. F. *et al.* Mo-, V-, and Fe-Nitrogenases Use a Universal Eight-Electron
Reductive-Elimination Mechanism To Achieve N_2 Reduction. *Biochemistry* **58**, 3293-
3301, doi:10.1021/acs.biochem.9b00468 (2019).

34   Zheng, Y. *et al.* A pathway for biological methane production using bacterial iron-only
nitrogenase. *Nat Microbiol* **3**, 281-286, doi:10.1038/s41564-017-0091-5 (2018).

35   Hu, Y., Lee, C. C. & Ribbe, M. W. Extending the Carbon Chain: Hydrocarbon Formation
Catalyzed by Vanadium/Molybdenum Nitrogenases. *Science* **333**, 753-755,
doi:10.1126/science.1206883 (2011).

36   Rohde, M., Trncik, C., Sippel, D., Gerhardt, S. & Einsle, O. Crystal structure of VnfH, the
iron protein component of vanadium nitrogenase. *J Biol Inorg Chem* **23**, 1049-1056,
doi:10.1007/s00775-018-1602-4 (2018).

37 Schmid, B. *et al.* Biochemical and structural characterization of the cross-linked complex
of nitrogenase: comparison to the ADP-AlF4(-)-stabilized structure. *Biochemistry* **41**,
15557-15565 (2002).
38 Spatzal, T. *et al.* Evidence for interstitial carbon in nitrogenase FeMo cofactor. *Science*
334, 940, doi:10.1126/science.1214025 (2011).
39 Altschul, S. F., Gish, W., Miller, W., Myers, E. W. & Lipman, D. J. Basic local alignment
search tool. *Journal of Molecular Biology* **215**, 403-410, doi:10.1016/s0022-
2836(05)80360-2 (1990).
40 Altschul, S. F. *et al.* Gapped BLAST and PSI-BLAST: a new generation of protein
database search programs. *Nucleic Acids Research* **25**, 3389-3402,
doi:10.1093/nar/25.17.3389 (1997).
41 Finn, R. D., Clements, J. & Eddy, S. R. HMMER web server: interactive sequence
similarity searching. *Nucleic Acids Res* **39**, W29-37, doi:10.1093/nar/gkr367 (2011).
42 van Kempen, M. *et al.* Foldseek: fast and accurate protein structure search.
doi:10.1101/2022.02.07.479398 (2022).
43 Illergard, K., Ardell, D. H. & Elofsson, A. Structure is three to ten times more conserved
than sequence--a study of structural response in protein cores. *Proteins* **77**, 499-508,
doi:10.1002/prot.22458 (2009).
44 Medina, M. S. *et al.* CowN sustains nitrogenase turnover in the presence of the inhibitor
carbon monoxide. *J Biol Chem* **296**, 100501, doi:10.1016/j.jbc.2021.100501 (2021).
45 Rubio, L. M. & Ludden, P. W. Maturation of nitrogenase: a biochemical puzzle. *J
Bacteriol* **187**, 405-414, doi:10.1128/JB.187.2.405-414.2005 (2005).
46 Tautz, D. & Domazet-Loso, T. The evolutionary origin of orphan genes. *Nat Rev Genet*
12, 692-702, doi:10.1038/nrg3053 (2011).
47 Heames, B., Schmitz, J. & Bornberg-Bauer, E. A Continuum of Evolving De Novo Genes
Drives Protein-Coding Novelty in Drosophila. *J Mol Evol* **88**, 382-398,
doi:10.1007/s00239-020-09939-z (2020).
48 Vakirlis, N. *et al.* De novo emergence of adaptive membrane proteins from thymine-rich
genomic sequences. *Nat Commun* **11**, 781, doi:10.1038/s41467-020-14500-z (2020).
49 Vakirlis, N., Carvunis, A. R. & McLysaght, A. Synteny-based analyses indicate that
sequence divergence is not the main source of orphan genes. *Elife* **9**,
doi:10.7554/eLife.53500 (2020).

50 Dos Santos, P. C., Fang, Z., Mason, S. W., Setubal, J. C. & Dixon, R. Distribution of
nitrogen fixation and nitrogenase-like sequences amongst microbial genomes. *BMC*
Genomics **13**, 162, doi:10.1186/1471-2164-13-162 (2012).
51 Parsons, C., Stüeken, E. E., Rosen, C. J., Mateos, K. & Anderson, R. E. Radiation of
nitrogen-metabolizing enzymes across the tree of life tracks environmental transitions in
Earth history. *Geobiology* **19**, 18-34, doi:10.1111/gbi.12419 (2020).
52 Anbar, A. D. *et al.* A whiff of oxygen before the great oxidation event? *Science* **317**,
1903-1906, doi:10.1126/science.1140325 (2007).
53 Ostrander, C. M., Johnson, A. C. & Anbar, A. D. Earth's First Redox Revolution. *Annu*
Rev Earth Pl Sc **49**, 337-366, doi:10.1146/annurev-earth-072020-055249 (2021).
54 Cole, D. B. *et al.* On the co-evolution of surface oxygen levels and animals. *Geobiology*
18, 260-281, doi:10.1111/gbi.12382 (2020).
55 Kessler, P. S., McLarnan, J. & Leigh, J. A. Nitrogenase phylogeny and the molybdenum
dependence of nitrogen fixation in Methanococcus maripaludis. *J Bacteriol* **179**, 541-543
(1997).
56 Rutledge, H. L., Cook, B. D., Nguyen, H. P. M., Herzik, M. A. & Tezcan, F. A. Structures
of the nitrogenase complex prepared under catalytic turnover conditions. *Science* **377**,
865-869, doi:10.1126/science.abq7641 (2022).
57 Huang, Q. *et al.* Mechanical coupling in the nitrogenase complex. *PLoS Comput Biol* **17**,
e1008719, doi:10.1371/journal.pcbi.1008719 (2021).
58 Schmid, B. *et al.* Structure of a cofactor-deficient nitrogenase MoFe protein. *Science*
296, 352-356, doi:10.1126/science.1070010 (2002).
59 Moore, E. K., Hao, J., Spielman, S. J. & Yee, N. The evolving redox chemistry and
bioavailability of vanadium in deep time. *Geobiology* **18**, 127-138, doi:10.1111/gbi.12375
(2020).
60 Emerson, S. R. & Huested, S. S. Ocean anoxia and the concentrations of molybdenum
and vanadium in seawater. *Marine Chemistry* **34**, 177-196, doi:10.1016/0304-
4203(91)90002-e (1991).
61 Zerkle, A. L., House, C. H. & Brantley, S. L. Biogeochemical signatures through time as
inferred from whole microbial genomes. *Am J Sci* **305**, 467-502, doi:DOI
10.2475/ajs.305.6-8.467 (2005).
62 Stueken, E. E., Buick, R., Guy, B. M. & Koehler, M. C. Isotopic evidence for biological
nitrogen fixation by molybdenum-nitrogenase from 3.2 Gyr. *Nature* **520**, 666-669,
doi:10.1038/nature14180 (2015).

63 Garcia, A. K., Kolaczkowski, B. & Kacar, B. Reconstruction of Nitrogenase Predecessors
Suggests Origin from Maturase-Like Proteins. *Genome Biol Evol* **14**,
doi:10.1093/gbe/evac031 (2022).

64 Ruttimann-Johnson, C., Rubio, L. M., Dean, D. R. & Ludden, P. W. VnfY is required for
full activity of the vanadium-containing dinitrogenase in Azotobacter vinelandii. *J*
Bacteriol **185**, 2383-2386, doi:10.1128/JB.185.7.2383-2386.2003 (2003).

65 Perez-Gonzalez, A. *et al.* Specificity of NifEN and VnfEN for the Assembly of
Nitrogenase Active Site Cofactors in Azotobacter vinelandii. *mBio* **12**, e0156821,
doi:10.1128/mBio.01568-21 (2021).

66 Rebelein, J. G., Lee, C. C., Newcomb, M., Hu, Y. & Ribbe, M. W. Characterization of an
M-Cluster-Substituted Nitrogenase VFe Protein. *MBio* **9**, doi:10.1128/mBio.00310-18
(2018).

67 Harwood, C. S. Iron-Only and Vanadium Nitrogenases: Fail-Safe Enzymes or Something
More? *Annu Rev Microbiol* **74**, 247-266, doi:10.1146/annurev-micro-022620-014338
(2020).

68 Hazen, R. M. *et al.* Mineral evolution. *American Mineralogist* **93**, 1693-1720,
doi:10.2138/am.2008.2955 (2008).

69 Sayers, E. W. *et al.* Database resources of the national center for biotechnology
information. *Nucleic Acids Res* **50**, D20-D26, doi:10.1093/nar/gkab1112 (2022).

70 Katoh, K. & Standley, D. M. MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. *Mol Biol Evol* **30**, 772-780,
doi:10.1093/molbev/mst010 (2013).

71 Fu, L., Niu, B., Zhu, Z., Wu, S. & Li, W. CD-HIT: accelerated for clustering the next-
generation sequencing data. *Bioinformatics* **28**, 3150-3152,
doi:10.1093/bioinformatics/bts565 (2012).

72 The Uniprot Consortium. UniProt: the Universal Protein Knowledgebase in 2023. *Nucleic*
Acids Res **51**, D523-D531, doi:10.1093/nar/gkac1052 (2023).

73 Nguyen, L. T., Schmidt, H. A., von Haeseler, A. & Minh, B. Q. IQ-TREE: a fast and
effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol Biol*
Evol **32**, 268-274, doi:10.1093/molbev/msu300 (2015).

74 Kalyaanamoorthy, S., Minh, B. Q., Wong, T. K. F., von Haeseler, A. & Jermiin, L. S.
ModelFinder: fast model selection for accurate phylogenetic estimates. *Nature Methods*
14, 587-589, doi:10.1038/nmeth.4285 (2017).

75 Hoang, D. T., Chernomor, O., von Haeseler, A., Minh, B. Q. & Vinh, L. S. UFBoot2:
Improving the Ultrafast Bootstrap Approximation. *Mol Biol Evol* **35**, 518-522,
doi:10.1093/molbev/msx281 (2018).

76 Guindon, S. *et al.* New algorithms and methods to estimate maximum-likelihood
phylogenies: assessing the performance of PhyML 3.0. *Syst Biol* **59**, 307-321,
doi:10.1093/sysbio/syq010 (2010).

77 Garcia, A. K. *et al.* Nitrogenase resurrection and the evolution of a singular enzymatic
mechanism. *eLife* **12**, doi:10.7554/eLife.85003 (2023).

78 Yang, Z. PAML 4: phylogenetic analysis by maximum likelihood. *Mol Biol Evol* **24**, 1586-
1591, doi:10.1093/molbev/msm088 (2007).

79 Aadland, K., Pugh, C. & Kolaczkowski, B. High-Throughput Reconstruction of Ancestral
Protein Sequence, Structure, and Molecular Function. *Methods Mol Biol* **1851**, 135-170,
doi:10.1007/978-1-4939-8736-8_8 (2019).

80 Tareen, A. & Kinney, J. B. Logomaker: beautiful sequence logos in Python.
Bioinformatics **36**, 2272-2274, doi:10.1093/bioinformatics/btz921 (2020).

81 Sauer, D. B. & Wang, D. N. Predicting the optimal growth temperatures of prokaryotes
using only genome derived features. *Bioinformatics* **35**, 3224-3231,
doi:10.1093/bioinformatics/btz059 (2019).

82 Ashkenazy, H. *et al.* ConSurf 2016: an improved methodology to estimate and visualize
evolutionary conservation in macromolecules. *Nucleic Acids Res* **44**, W344-350,
doi:10.1093/nar/gkw408 (2016).

83 Penev, P. I., Alvarez-Carreno, C., Smith, E., Petrov, A. S. & Williams, L. D. TwinCons:
Conservation score for uncovering deep sequence similarity and divergence. *PLoS*
Comput Biol **17**, e1009541, doi:10.1371/journal.pcbi.1009541 (2021).

84 Mirdita, M. *et al.* ColabFold: making protein folding accessible to all. *Nature Methods* **19**,
679-682, doi:10.1038/s41592-022-01488-1 (2022).

85 Steinegger, M. & Söding, J. MMseqs2 enables sensitive protein sequence searching for
the analysis of massive data sets. *Nature Biotechnology* **35**, 1026-1028,
doi:10.1038/nbt.3988 (2017).

86 Pettersen, E. F. *et al.* UCSF ChimeraX: Structure visualization for researchers,
educators, and developers. *Protein Sci* **30**, 70-82, doi:10.1002/pro.3943 (2021).

87 Csermely, P. Creative elements: network-based predictions of active centres in proteins
and cellular and social networks. *Trends Biochem Sci* **33**, 569-576,
doi:10.1016/j.tibs.2008.09.006 (2008).

88   Bouysset, C. & Fiorucci, S. ProLIF: a library to encode molecular interactions as
793   fingerprints. *Journal of Cheminformatics* **13**, doi:10.1186/s13321-021-00548-6 (2021).
794   89   Hagberg, A. A., Schult, D. A. & Swart, P. J. in *Proceedings of the 7th Python in Science*
795   *Conference.* (eds Gäel Varoquaux, Travis Vaught, & Jarrod Millman) 11-15.
796   90   Bastian, M., Heymann, S. & Jacomy, M. Gephi: An Open Source Software for Exploring
797   and Manipulating Networks. *Proceedings of the International AAAI Conference on Web*
798   *and Social Media* **3**, 361-362, doi:10.1609/icwsm.v3i1.13937 (2009).
799   91   Jurrus, E. *et al.* Improvements to the APBS biomolecular solvation software suite.
800   *Protein Science* **27**, 112-128, doi:10.1002/pro.3280 (2017).
801   92   Kortemme, T., Kim, D. E. & Baker, D. Computational Alanine Scanning of Protein-
802   Protein Interfaces. *Science's STKE* **2004**, doi:10.1126/stke.2192004pl2 (2004).
803   93   Rodrigues, C. H. M., Myung, Y., Pires, D. E. V. & Ascher, D. B. mCSM-PPI2: predicting
804   the effects of mutations on protein–protein interactions. *Nucleic Acids Research* **47**,
805   W338-W344, doi:10.1093/nar/gkz383 (2019).
806   94   Gerek, Z. N. & Ozkan, S. B. Change in allosteric network affects binding affinities of PDZ
807   domains: analysis through perturbation response scanning. *PLoS Comput Biol* **7**,
808   e1002154, doi:10.1371/journal.pcbi.1002154 (2011).